

*Pacific
Journal of
Mathematics*

A SHARP DIVISION ESTIMATE FOR
ULTRADIFFERENTIABLE GERMS

VINCENT THILLIEZ

Volume 205 No. 1

July 2002

A SHARP DIVISION ESTIMATE FOR ULTRADIFFERENTIABLE GERMS

VINCENT THILLIEZ

Let f (resp. φ) be a C^∞ (resp. real-analytic) function germ near the origin in \mathbf{R}^n . Assume that f is divisible by φ in C^∞ , and that it belongs to a sufficiently regular ultradifferentiable class $\{\ell! M_\ell\}$ of Carleman type (for example, one of the Gevrey rings familiar in the theory of differential equations). What can then be said about the regularity of the quotient f/φ ? In this paper, we obtain first a complete solution of this problem in the case $n = 2$. Namely, it is shown that f/φ belongs to the Carleman class $\{\ell! M_\ell^{d(\varphi)}\}$, where $d(\varphi)$ is a suitable Lojasiewicz exponent for the regular separation between the space \mathbf{R}^2 and certain components of the complex zero set Z_φ of φ . This number can be explicitly computed by means of Puiseux expansions. We prove moreover that the division result is sharp for any φ and M . Finally, we apply it to get a characterization of closed principal ideals generated by real-analytic functions in Carleman classes of two variables, improving a result which was known previously only in the case of generators with isolated real zeros.

Introduction.

Many studies in differential analysis involve a more or less explicit version of the following division problem: Let φ be a given real-analytic function germ at the origin in \mathbf{R}^n and let f be another germ enjoying some regularity properties. What kind of control on the quotient f/φ is it then possible to derive from the knowledge of geometric and differential properties of φ and f ? As typical examples of questions in this spirit, we refer the reader to the Lojasiewicz division theorem [10], [11], the work of Mostow-Shnider on continuity of C^∞ quotients [12], theorems of Izumi [8] and Wang [17] on Chevalley estimates, and so on. Of course, the kind of regularity property which is dealt with (order of vanishing, smoothness, estimates on derivatives...) has a strong influence on the nature of the corresponding result.

The present article studies division estimates in the setting of very smooth germs, namely ultradifferentiable classes. Denote by $C^\infty(\mathbf{R}^n, 0)$ the ring of infinitely differentiable function-germs at the origin in \mathbf{R}^n , and recall

that the Carleman class of germs $C_M(\mathbf{R}^n, 0)$ is defined as the subclass of $C^\infty(\mathbf{R}^n, 0)$ containing all those germs f whose derivatives at any order ℓ are majorized by $C_f^{\ell+1} \ell! M_\ell$ in some fixed neighborhood of 0, where C_f is a constant depending on f , and $(M_\ell)_{\ell \geq 0}$ is a given sequence of real numbers satisfying suitable regularity assumptions. In some sense, the role of this sequence consists in putting a bound on the defect of analyticity of the elements of $C_M(\mathbf{R}^n, 0)$. Thus, such classes provide scales of regularity between the analytic and the C^∞ ones. A well-known and much studied example is given by the scale of Gevrey classes ($M_\ell = \ell!^\alpha$ with $\alpha > 0$). In Carleman classes, the general division problem described above can be precisely stated as follows: Being given a germ φ of real-analytic function at the origin in \mathbf{R}^n and a Carleman class $C_M(\mathbf{R}^n, 0)$, investigate the smallest possible class $C_{M^+}(\mathbf{R}^n, 0)$ containing $C_M(\mathbf{R}^n, 0)$ and enjoying the following property: If f is an element of $C_M(\mathbf{R}^n, 0)$ which can be written as φg for some g belonging to $C^\infty(\mathbf{R}^n, 0)$, then the quotient $g = f/\varphi$ belongs to $C_{M^+}(\mathbf{R}^n, 0)$. Of course, the sequence M^+ should depend only on M and on geometric and differential properties of φ . The reason why we consider different classes C_M and C_{M^+} lies in the fact that some loss of regularity turns out to be generally unavoidable in the division process; hence the class of the quotient has to be (possibly strictly) wider than C_M . The only exception arises for $n = 1$, where obvious arguments show that $M^+ = M$.

In this paper, we obtain a complete solution of the problem for $n = 2$, that is, for germs in two variables. As a starting point, we mention that in the particular case where the real zero set of φ reduces to the single point $\{0\}$, it is possible to see, as a corollary of the results in [15], that the solution is given by $M_\ell^+ = M_\ell^{\tau(\varphi)}$, where $\tau(\varphi)$ is the Lojasiewicz exponent for the regular separation between \mathbf{R}^2 and the complex zero set of φ . Moreover, this choice of M^+ is sharp. However, after [15], the previous division problem remained unsolved for general zero sets, even for two variables. In this setting, we shall show here (Theorem 2.1) that the previous solution is still valid, provided we extend the definition of M^+ by putting

$$M_\ell^+ = M_\ell^{d(\varphi)},$$

where $d(\varphi)$ is, roughly speaking, the greatest number among all Lojasiewicz exponents for the regular separation between \mathbf{R}^2 and one of those branches of the complex zero set of φ whose intersection with a half-plane in \mathbf{R}^2 reduces to $\{0\}$ (see Section 1 for details). Of course, we have $d(\varphi) = \tau(\varphi)$ when the intersection of the zero set itself with \mathbf{R}^2 reduces to $\{0\}$.

Our method of proof relies exclusively on explicit estimates, striving to stay as accurate as possible at each step. The underlying reason why dimension 2 plays a special role lies in the fact that, as in [15], it allows to relate $d(\varphi)$ to some numbers occurring explicitly in Puiseux expansions of

the roots of φ . However, some particular results can be extended to the higher dimensional case; see the details in Subsection 4.3 below. We owe these extensions to Edward Bierstone and we take here the opportunity to thank him for his insightful comments on the subject.

Finally, we mention that another strategy to study such division problems in certain classes of smooth functions has been developed by Bos-Milman [4] and Chaumat-Chollet [7]: Using resolution of singularities, these authors are led to a quite different kind of results, as explained in Remark 4.4 below.

The article is organized as follows: Section 1 recalls definitions and various prerequisites; Section 2 states and partly proves the main result (Theorem 2.1). Section 3 is devoted to establishing a technical estimate, thus completing the proof of Theorem 2.1; this is the most delicate part of the paper. In Section 4, we discuss the sharpness of the main result (see Theorem 4.1); we apply it to a characterization of closed ideals (Theorem 4.2) and give some comments.

1. Preliminaries.

1.1. Notations. Throughout the paper, for any multi-index $L = (\ell_1, \dots, \ell_n)$ in \mathbf{N}^n , we shall denote by the corresponding lower case letter ℓ the length $\ell_1 + \dots + \ell_n$ of L . We also put $L! = \ell_1! \dots \ell_n!$ and denote by D^L the operator $\partial_1^{\ell_1} \dots \partial_n^{\ell_n}$, with $\partial_j = \partial/\partial x_j$ for $j = 1, \dots, n$. We say that a C^∞ germ f is flat at a point a if f and all its derivatives vanish at a . For any subset E of \mathbf{R}^n or \mathbf{C}^n , we shall sometimes write $(E, 0)$ to denote $E \cap U$ where U is some neighborhood of 0. When we write that a property $\mathcal{P}(x, y)$, depending on $x \in \mathbf{R}^n$ and on some family of other parameters y , holds for x in $(E, 0)$, it is assumed that the neighborhood U does not depend on y . We denote by $d(x, E)$ the euclidean distance from a point x to E . At last, the following classical notation will be useful: If $A(t)$ and $B(t)$ are two positive functions of a variable t belonging to a set T , one writes $A(t) \lesssim B(t)$ (or equivalently $B(t) \gtrsim A(t)$) to say that there exists a real C , not depending on t , such that the inequality $A(t) \leq CB(t)$ holds for all $t \in T$. The notation $A(t) \approx B(t)$ means that one has $A(t) \lesssim B(t)$ and $B(t) \lesssim A(t)$ simultaneously.

1.2. On ultradifferentiable germs. Let $M = (M_\ell)_{\ell \geq 0}$ be an increasing sequence of real numbers with $M_0 = 1$. The sequence M is said to be *strongly regular* if there exists a positive constant A such that the following assumptions are satisfied:

- (1) M is logarithmically convex,
- (2) $M_{j+k} \leq A^{j+k} M_j M_k$ for any $(j, k) \in \mathbf{N}^2$,
- (3) $\sum_{j \geq \ell} \frac{M_j}{(j+1)M_{j+1}} \leq A \frac{M_\ell}{M_{\ell+1}}$ for any $\ell \in \mathbf{N}$.

Condition (1), which amounts to saying that $M_{\ell+1}/M_\ell$ increases with ℓ , implies

$$(4) \quad M_j M_k \leq M_{j+k} \text{ for any } (j, k) \in \mathbf{N}^2.$$

Thus (2), which is in some sense converse to (4), means that the growth of M is not too wild. The role of condition (3) will be recalled later, after the definition of ultradifferentiable germs. To each strongly regular sequence M , one associates classically the function h_M defined by $h_M(t) = \inf_{\ell \geq 0} t^\ell M_\ell$ for any $t \in \mathbf{R}_+$. This function, which is a sort of Legendre transform, is continuous, non-decreasing; we have $h_M(0) = 0$ and $h_M(t) = 1$ for $t \geq 1$. In virtue of (1), the knowledge of h_M fully determines the sequence M since we have

$$(5) \quad M_\ell = \sup_{t > 0} t^{-\ell} h_M(t) \text{ for any } \ell \in \mathbf{N}.$$

For the typical example of Gevrey sequences $M_\ell = \ell!^\alpha$ ($\alpha > 0$), we have $h_M(t) = \exp(-\lambda(t))$ with $\lambda(t) \approx t^{-1/\alpha}$.

Now let s be a real number with $s \geq 1$. Obviously, the sequence $M^s = (M_\ell^s)_{\ell \geq 0}$ is also strongly regular, and we have $h_{M^s}(t^s) = (h_M(t))^s$. We recall also that there exists a constant $\rho(s) \geq 1$, depending only on s and M , such that

$$(6) \quad h_M(t) \leq (h_M(\rho(s)t))^s \text{ for any } t \in \mathbf{R}_+.$$

From (2), (5) and (6), it is straightforward to deduce that there exists a positive constant $A(s)$, depending only on s and M , such that

$$(7) \quad t^{-s\ell} h_M(t) \leq A(s)^\ell M_\ell^s h_M(A(s)t) \text{ for any } t \in \mathbf{R}_+^* \text{ and } \ell \in \mathbf{N}.$$

As in [15], for any strongly regular sequence M , we define now $C_M(\mathbf{R}^n, 0)$ as the class of germs f in $C^\infty(\mathbf{R}^n, 0)$ for which there exist a neighborhood U of 0 in \mathbf{R}^n and a constant $C > 0$ such that the inequality

$$|D^L f(x)| \leq C^{\ell+1} \ell! M_\ell$$

holds for any x in U and any multi-index L in \mathbf{N}^n . The class $C_M(\mathbf{R}^n, 0)$ is a local algebra, stable under derivation. As remarked in the introduction, the sequence M majorizes, in some sense, the defect of analyticity of its elements. Condition (3) is the so-called *strong non-quasianalyticity* condition ensuring the existence of “good” cutoff functions with C_M regularity [5]. For a more detailed interpretation of (1), (2), (3) in terms of ultradifferentiable classes, we refer the reader to [5], [9], [13]. Beware that the notations in [5], [9], [13] are different, since M_ℓ plays there the role played by $\ell! M_\ell$ in the present paper. The latter notation, viewing M as a defect of analyticity, is more convenient here, since it seems, as a general rule, that any loss of regularity involved by the various operations of differential analysis (division, etc.) in the setting of ultradifferentiable functions, acts only on the defect of analyticity.

1.3. On analytic germs. Denote by $\mathcal{O}(\mathbf{R}^n, 0)$ (resp. $\mathcal{O}(\mathbf{C}^n, 0)$) the ring of real-analytic (resp. holomorphic) function germs at the origin in \mathbf{R}^n (resp. \mathbf{C}^n). By means of the power series expansion, $\mathcal{O}(\mathbf{R}^n, 0)$ can obviously be viewed as the subset of elements of $\mathcal{O}(\mathbf{C}^n, 0)$ taking real values on \mathbf{R}^n . Thus, to each real-analytic germ γ in $\mathcal{O}(\mathbf{R}^n, 0)$, one can associate the set of its complex zeros $Z_\gamma = \gamma^{-1}(0)$ as a germ of complex analytic set.

From now on, we shall denote by $\omega(v)$ the order of any element v of $\mathcal{O}(\mathbf{C}, 0)$, that is the least degree of monomials occurring in the power series expansion of v , with the convention $\omega(0) = \infty$.

Let φ be a given element of $\mathcal{O}(\mathbf{R}^2, 0)$. After a linear change of coordinates with real matrix, one can clearly assume, without changing the fact that φ is real, that the z_1 -axis is transversal to the zero set Z_φ at 0. Using Puiseux’s theorem as in [15], we can write, for some integer m in \mathbf{N}^* ,

$$\varphi(z_1, z_2^m) = u(z_1, z_2^m) \prod_{p=1}^{p_1} (z_1 - U_p(z_2))^{n_p},$$

where n_1, \dots, n_{p_1} are positive integers, u is a unit in $\mathcal{O}(\mathbf{C}^2, 0)$ and U_1, \dots, U_{p_1} is a suitable family of germs in $\mathcal{O}(\mathbf{C}, 0)$, satisfying $\omega(U_p) \geq m$ for $p = 1, \dots, p_1$. Put $U_p = R_p + iS_p$ with $R_p \in \mathcal{O}(\mathbf{R}, 0)$ and $S_p \in \mathcal{O}(\mathbf{R}, 0)$. Using the fact that φ is real, it is not difficult to see that for x_2 in \mathbf{R}^* , the roots $U_p(x_2)$ for which $S_p(x_2)$ is non-identically zero are pairwise conjugate. After reindexation, we can thus write, for some suitable integer p_0 satisfying $2p_0 \leq p_1$,

$$U_p = \begin{cases} R_p + iS_p & \text{with } m \leq \omega(S_p) < \infty \text{ for } p = 1, \dots, p_0, \\ R_{p-p_0} - iS_{p-p_0} & \text{for } p = p_0 + 1, \dots, 2p_0, \\ R_p & \text{with } \omega(R_p) \geq m \text{ for } p = 2p_0 + 1, \dots, p_1. \end{cases}$$

Note that we can write

$$\varphi(z_1, z_2^m) = u(z_1, z_2^m) \prod_{p=1}^{p_0} (\psi_p(z_1, z_2))^{n_p} \prod_{p=2p_0+1}^{p_1} (z_1 - R_p(z_2))^{n_p}$$

with $\psi_p(z_1, z_2) = (z_1 - R_p(z_2))^2 + (S_p(z_2))^2$; and we have also $u \in \mathcal{O}(\mathbf{R}^2, 0)$ since φ is real. The special case $p_1 = 2p_0$, occurring when $Z_\varphi \cap (\mathbf{R} \times \mathbf{R}_+) = \{0\}$, has been studied in detail in [15]. In the same spirit, we define here

$$\begin{aligned} \mu_p &= \omega(S_p) \text{ and } \tau_p^+ = \mu_p/m \text{ for } p = 1, \dots, p_0, \\ d^+(\varphi) &= \max\{\tau_p^+; p = 1, \dots, p_0\} \text{ if } p_0 \geq 1 \text{ and } d^+(\varphi) = 1 \text{ if } p_0 = 0, \\ d^-(\varphi) &= d^+(\tilde{\varphi}) \text{ with } \tilde{\varphi}(z_1, z_2) = \varphi(z_1, -z_2), \\ d(\varphi) &= \max(d^+(\varphi), d^-(\varphi)). \end{aligned}$$

We denote by X_φ the set $Z_\varphi \cap \mathbf{R}^2$ of real zeros of φ . For technical reasons, we also define, for $p = 1, \dots, p_1$,

$$Z_p = \{z; \operatorname{Re} z_2 \geq 0 \text{ and } z_1 = U_p(z_2^{1/m})\}, \quad X_p = Z_p \cap \mathbf{R}^2,$$

$$Z_\varphi^\pm = \{z \in Z_\varphi; \operatorname{Re} z_2 \in \mathbf{R}_\pm\}, \quad X_\varphi^\pm = Z_\varphi^\pm \cap \mathbf{R}^2,$$

where $z_2^{1/m}$ is the principal determination of the m^{th} -root of z_2 .

1.4. On Łojasiewicz exponents. Let \mathcal{X} and \mathcal{Y} be two germs of closed subsets at the origin in \mathbf{R}^n , with $0 \in \mathcal{X} \cap \mathcal{Y}$; and let \mathcal{T} be the set of real numbers $t \geq 1$ for which there exists a constant $c_t > 0$ such that one has

$$d(x, \mathcal{Y}) \geq c_t d(x, \mathcal{X} \cap \mathcal{Y})^t \text{ for any } x \in (\mathcal{X}, 0).$$

Recall that \mathcal{X} and \mathcal{Y} are said to be regularly separated if \mathcal{T} is not empty (it is classical that this definition is symmetric with respect to \mathcal{X} and \mathcal{Y} , see e.g., [11]). The Łojasiewicz exponent for the regular separation between \mathcal{X} and \mathcal{Y} is then defined as the infimum of \mathcal{T} . As a fundamental fact of subanalytic geometry, it is known that the subanalyticity of \mathcal{X} and \mathcal{Y} implies regular separation, and that the Łojasiewicz exponent is rational and attained (in other words, it belongs to \mathcal{T}), see [3]. If it is equal to 1, the sets \mathcal{X} and \mathcal{Y} are said to intersect transversally.

Here, according to [15], the number $d^+(\varphi)$ defined in the previous subsection is precisely the Łojasiewicz exponent for the regular separation between \mathbf{R}^2 and the union $\bigcup_{1 \leq p \leq 2p_0} Z_p$ of all those “branches” Z_p in Z_φ^+ for which $X_p = \{0\}$ (for the other ones, X_p is a C^1 real arc originating in $\{0\}$, and Z_p intersects \mathbf{R}^2 transversally, as shown in Lemma 2.3 below). Notice also that when we have $Z_\varphi \cap \mathbf{R}^2 \neq \{0\}$, the number $d(\varphi)$, which has been defined as $\max(d^+(\varphi), d^-(\varphi))$, generally does not coincide anymore with the Łojasiewicz exponent for the regular separation between \mathbf{R}^2 and Z_φ . Indeed, it is possible to check that the latter exponent $\tau(\varphi)$ always satisfies $\tau(\varphi) \leq d(\varphi)$; but this inequality can be strict, see the following Example (iv).

1.5. Examples.

- (i) $\varphi(x) = x_1^2 - x_2^{2k}$ ($k \in \mathbf{N}^*$). This the simplest kind of example: We can write $\varphi(x) = \check{\varphi}(x) = (x_1 - x_2^k)(x_1 + x_2^k)$, hence $m = 1, p_1 = 2, p_0 = 0$ for both φ and $\check{\varphi}$. We derive $d^+(\varphi) = d^-(\varphi) = d(\varphi) = 1$.
- (ii) $\varphi(x) = x_1^2 - x_2^{2k+1}$ ($k \in \mathbf{N}^*$), for which X_φ is a cusp. We can write $\varphi(x_1, x_2) = (x_1 - x_2^{2k+1})(x_1 + x_2^{2k+1})$, hence $m = 2, p_1 = 2, p_0 = 0$ and $d^+(\varphi) = 1$. On the other hand, we have $\check{\varphi}(x) = x_1^2 + x_2^{2k+1}$, thus $\check{\varphi}(x_1, x_2) = (x_1 - ix_2^{2k+1})(x_1 + ix_2^{2k+1})$: For this one, we get $m = 2, p_1 = 2, p_0 = 1, \mu_1 = 2k + 1$, hence $d^-(\varphi) = d^+(\check{\varphi}) = k + \frac{1}{2}$. We derive finally $d(\varphi) = k + \frac{1}{2}$.

(iii) $\varphi(x) = (x_1^2 - x_2^3)(x_1^3 - x_2^4)$. We have

$$\varphi(x_1, x_2^6) = \left(x_1 + \frac{1}{2}x_2^8 + i\frac{\sqrt{3}}{2}x_2^8\right) \left(x_1 + \frac{1}{2}x_2^8 - i\frac{\sqrt{3}}{2}x_2^8\right) (x_1 - x_2^8)(x_1 - x_2^9)(x_1 + x_2^9).$$

This yields $m = 6, p_1 = 5, p_0 = 1, \mu_1 = 8$ and consequently $d^+(\varphi) = 4/3$. We have also $\check{\varphi}(x) = (x_1^2 + x_2^3)(x_1^3 - x_2^4)$, hence

$$\check{\varphi}(x_1, x_2^6) = (x_1 - ix_2^9)(x_1 + ix_2^9) \left(x_1 + \frac{1}{2}x_2^8 + i\frac{\sqrt{3}}{2}x_2^8\right) \left(x_1 + \frac{1}{2}x_2^8 - i\frac{\sqrt{3}}{2}x_2^8\right) (x_1 - x_2^8).$$

For $\check{\varphi}$, we can thus take $m = 6, p_1 = 5, p_0 = 2, \mu_1 = 9, \mu_2 = 8$ and $d^-(\varphi) = d^+(\check{\varphi}) = 3/2$. Finally, we get $d(\varphi) = 3/2$.

(iv) $\varphi(x) = x_1((x_1 - x_2^2)^2 + x_2^8)$. We can write

$$\varphi(x) = \check{\varphi}(x) = x_1(x_1 - x_2^2 + ix_2^4)(x_1 - x_2^2 - ix_2^4),$$

hence $m = 1, p_1 = 3, p_0 = 1, \mu_1 = 4$ and $d(\varphi) = 4$. Now we compute the Lojasiewicz exponent $\tau(\varphi)$ for the regular separation between \mathbf{R}^2 and Z_φ . In \mathbf{C}^3 , the set Z_φ has 3 smooth irreducible components Z', Z''_+, Z''_- respectively given by $Z' = \{z_1 = 0\}$ and $Z''_\pm = \{z_1 - z_2^2 \pm iz_2^4 = 0\}$. Pick $x \in (\mathbf{R}^2, 0)$. Elementary arguments show that $d(x, Z') = |x_1|$ and $d(x, Z''_\pm) \approx |x_1 - x_2^2 \pm ix_2^4| \approx |x_1 - x_2^2| + x_2^4$, hence $d(x, Z_\varphi) \approx \min(|x_1|, |x_1 - x_2^2| + x_2^4)$. For $|x_1| \leq 2x_2^2$, we have $|x_1 - x_2^2| + x_2^4 \geq x_2^4 \gtrsim x_1^2$ and otherwise we have $|x_1 - x_2^2| + x_2^4 \geq |x_1 - x_2^2| \gtrsim |x_1|$. We derive thus the separation inequality $d(x, Z_\varphi) \gtrsim x_1^2 = d(x, X_\varphi)^2$. On the other hand, the point $x(\epsilon) = (\epsilon^2, \epsilon)$ satisfies $d(x(\epsilon), X_\varphi) = \epsilon^2$ and $d(x(\epsilon), Z_\varphi) \leq \epsilon^4$, since $x(\epsilon) + (i\epsilon^4, 0)$ belongs to Z_φ . Therefore, the exponent 2 can not be lowered in the separation inequality, and we have $\tau(\varphi) = 2$ whereas $d(\varphi) = 4$.

(v) Note finally that if φ is a homogeneous polynomial, we have obviously $d(\varphi) = \tau(\varphi) = 1$.

2. A division theorem.

In the sequel, φ denotes a given germ in $\mathcal{O}(\mathbf{R}^2, 0)$. Using the notations of Section 1, the main result of this paper reads as follows:

Theorem 2.1. *Let M be a strongly regular sequence and let g be a germ in $C^\infty(\mathbf{R}^2, 0)$ such that φg belongs to $C_M(\mathbf{R}^2, 0)$. Then g belongs to $C_{M^{d(\varphi)}}(\mathbf{R}^2, 0)$.*

This theorem reveals that in some sense, the real arcs of Z_φ do not contribute to the loss of regularity involved in the division process. We give now a few geometric lemmas to prepare the proof.

Lemma 2.2. *For $1 \leq p \leq 2p_0$ and $x \in (\mathbf{R} \times \mathbf{R}_+, 0)$, one has $d(x, Z_p) \gtrsim |x|^{r_p^+}$.*

Proof. Use the same argument as in [15, Lemma 3.2]. □

Lemma 2.3. *For $2p_0 + 1 \leq p \leq p_1$ and $x \in (\mathbf{R} \times \mathbf{R}_+, 0)$, one has $d(x, Z_p) \gtrsim d(x, X_p)$.*

Proof. Put $y = (R_p(x_2^{1/m}), x_2)$; then y belongs to X_p and one has thus $d(x, X_p) \leq |x - y| = |x_1 - R_p(x_2^{1/m})|$. Besides this, as in [15, Lemma 3.2], the function given by $z \mapsto |z_1 - R_p(z_2^{1/m})|$ for $\text{Re } z_2 \geq 0$ can be extended to a C^1 -function T_p in a neighborhood of 0 in \mathbf{C}^2 ; and $T_p^{-1}(0)$ is a 2-dimensional real submanifold of \mathbf{C}^2 containing Z_p . Hence we derive $d(x, Z_p) \geq d(x, T_p^{-1}(0)) \approx |T_p(x)| = |x_1 - R_p(x_2^{1/m})|$, which ends the proof. □

Lemma 2.4. *For $1 \leq p \leq p_1$ and $x \in (\mathbf{R} \times \mathbf{R}_-, 0)$, one has $d(x, Z_p) \gtrsim |x|$.*

Proof. Use the same argument as in [15, Lemma 3.3]. □

Proposition 2.5. *For $x \in (\mathbf{R}^2, 0)$, one has $d(x, Z_\varphi) \gtrsim \min(|x|^{d(\varphi)}, d(x, X_\varphi))$.*

Proof. We remark first that we have

$$(8) \quad d(x, Z_\varphi^+) \gtrsim \min(|x|^{d^+(\varphi)}, d(x, X_\varphi^+)).$$

Indeed, for $x \in (\mathbf{R} \times \mathbf{R}_+, 0)$, the estimate (8) is a direct consequence of Lemmas 2.2 and 2.3. For $x \in (\mathbf{R} \times \mathbf{R}_-, 0)$, it is a consequence of Lemma 2.4. Now, applying (8) to $\check{\varphi}(z) = \varphi(z_1, -z_2)$, it is easy to obtain symmetrically

$$(9) \quad d(x, Z_\varphi^-) \gtrsim \min(|x|^{d^-(\varphi)}, d(x, X_\varphi^-)).$$

The estimates (8) and (9) yield the proposition. □

The previous geometric proposition, which extends [15, Proposition 3.1], is one of the two key ingredients in the proof of Theorem 2.1. The other key ingredient is of a more analytic nature; it can be stated as follows:

Proposition 2.6. *Under the assumptions of Theorem 2.1, one can find constants B_1 and B_2 such that one has, for any bi-index L and any point a of the real zero set X_φ ,*

$$|D^L g(a)| \leq B_1 B_2^\ell \ell! M_\ell^{d(\varphi)}.$$

The proof of Proposition 2.6 is quite technical and all of Section 3 will be devoted to it. Taking the result for granted, we shall end Section 2 by the proof of Theorem 2.1. Before this, the following important first step has to be carried out:

2.7. Reduction to the case of flat germs at the origin. Under the assumptions of Theorem 2.1, we claim that for any point a in X_φ , there exist constants $B_{1,a}$ and $B_{2,a}$, depending on a , such that we have

$$(10) \quad |D^L g(a)| \leq B_{1,a} B_{2,a}^\ell \ell! M_\ell \text{ for any } L \in \mathbf{N}^2.$$

It is indeed an easy consequence of results of Chaumat-Chollet [6, Theorem 9 and Remark 17], but it can be proved by elementary arguments as well [16]. Such a statement means that division estimates hold pointwise (that is: On formal power series) without loss of regularity. However, the constants in (10) are by no means uniformly bounded with respect to a , whereas Proposition 2.6 gives a uniform estimate, at the cost of some loss of regularity. Nevertheless, the above estimate (10) will be useful, in the following way: Taking into account (10) for $a = 0$ and Borel’s theorem for Carleman classes [13], we get some g_0 in $C_M(\mathbf{R}^2, 0)$ such that $g - g_0$ is flat at 0. Clearly, replacing g by $g - g_0$ does not change the assumption nor the conclusion of Theorem 2.1. Thus we see that it is always possible to make the extra assumption that φg is flat at 0, without disturbing the generality of the theorem. In the sequel, we shall use systematically this property, which, by standard arguments (see e.g., the estimate (2.17) in [15]) yields the majorization

$$(11) \quad |D^K(\varphi g)(x)| \leq B_3^{k+1} k! M_k h_M(B_4|x|)$$

for any bi-index K and any point x in a suitable neighborhood of 0 in \mathbf{R}^2 , where B_3 and B_4 are constants depending only on g and φ .

2.8. Proof of Theorem 2.1. From now on, put $f = \varphi g$. Taking Proposition 2.6 into account, we still have to estimate the derivatives of g at each point of $\mathbf{R}^2 \setminus X_\varphi$. By the Leibniz formula, we have

$$(12) \quad D^L g(x) = \sum_{J,K; J+K=L} \frac{L!}{J!K!} (D^J(1/\varphi)(x)) D^K f(x).$$

The Łojasiewicz inequality asserts that there exist constants B_5 and ν , with $B_5 > 0$ and $\nu \geq 1$, such that the estimate $|\varphi(z)| \geq B_5 d(z, Z_\varphi)^\nu$ holds for any z in $(\mathbf{C}^2, 0)$. Applying the Cauchy formula to $1/\varphi$ on the bi-disk of radius $\frac{1}{4}d(x, Z_\varphi)$ centered at x , we derive, in virtue of Proposition 2.5 and of the previous inequality, that one has

$$(13) \quad |D^J(1/\varphi)(x)| \leq B_6^{j+1} j! \left(\min \left(|x|^{d(\varphi)}, d(x, X_\varphi) \right) \right)^{-(j+\nu)}.$$

Now there are two distinct situations:

First case: $\min(|x|^{d(\varphi)}, d(x, X_\varphi)) = |x|^{d(\varphi)}$. Then we proceed exactly as in [15, Theorem 2.3] to deduce from (7), (11), (12), (13) that we have $|D^L g(x)| \leq B_7^{\ell+1} \ell! M_\ell^{d(\varphi)}$ for some suitable constant B_7 .

Second case: $\min(|x|^{d(\varphi)}, d(x, X_\varphi)) = d(x, X_\varphi)$. Using Proposition 2.6 together with Petzsche’s continuous version of the Borel theorem for Carleman classes [13], we see that for any point a in X_φ , there exists a C^∞ function g_a satisfying, for any bi-index L ,

$$(14) \quad D^L g_a(a) = D^L g(a),$$

$$(15) \quad |D^L g_a(y)| \leq B_1 B_8 (B_2 B_9)^\ell \ell! M_\ell^{d(\varphi)} \quad \text{for any } y \text{ in } \mathbf{R}^2,$$

with constants B_8 and B_9 depending only on M ; in particular, uniform with respect to a . Put $f_a = \varphi g_a$. From (14) and (15) we derive that $f - f_a$ belongs to $C_{M^{d(\varphi)}}(\mathbf{R}^2, 0)$, is flat at a and we have, in the same way as for (11),

$$(16) \quad |D^K (f - f_a)(x)| \leq B_{10}^{k+1} k! M_k^{d(\varphi)} h_{M^{d(\varphi)}}(B_{11}|x - a|)$$

for any bi-index K , with constants B_{10} and B_{11} not depending on a . We have also $h_{M^{d(\varphi)}}(B_{11}|x - a|) \leq (B_{11}|x - a|)^{j+[\nu]+1} M_{j+[\nu]+1}^{d(\varphi)}$ and, using (4) and (2), $M_{j+[\nu]+1}^{d(\varphi)} M_k^{d(\varphi)} \leq B_{12}^{\ell+1} M_\ell^{d(\varphi)}$ for $j + k = \ell$. Pick a point a in X_φ satisfying $|x - a| = d(x, X_\varphi)$. Then, applying (12) with $g - g_a$ instead of g and $f - f_a$ instead of f , one gets, in virtue of (7), (13), (16) and the previous inequalities, $|D^L (g - g_a)(x)| \leq B_{13}^{\ell+1} \ell! M_\ell^{d(\varphi)}$. Taking (15) into account, we derive $|D^L g(x)| \leq B_{14}^{\ell+1} \ell! M_\ell^{d(\varphi)}$ for some suitable B_{14} , thus ending the proof. \square

We turn now to the proof of Proposition 2.6.

3. Estimates along the zero set.

We shall use the analytic vector field $\mathbf{Y} = m^{-1} x_2^{1-m} \partial_2$ on $\mathbf{R} \times \mathbf{R}^*$, in such a way that we have, for any h in $C^\infty(\mathbf{R}^2, 0)$ and any x in $(\mathbf{R} \times \mathbf{R}^*, 0)$,

$$(\partial_2 h)(x_1, x_2^m) = \mathbf{Y}(h(x_1, x_2^m)).$$

The following proposition will play a crucial role.

Proposition 3.1. *Let R be an element of $\mathcal{O}(\mathbf{R}, 0)$ satisfying $\omega(R) \geq m$ and let N be an integer with $N \geq 1$. Consider a germ g in $C^\infty(\mathbf{R}^2, 0)$ and assume that for every real $\beta > 0$, there exist constants C_1 and C_2 , depending on β , such that for any point x in $(\mathbf{R}^2, 0)$ satisfying $|x_1| < \beta|x_2|^m$, the estimate*

$$\left| \partial_1^k \mathbf{Y}^\ell ((x_1 - R(x_2))^N G(x)) \right| \leq C_1^{k+\ell+1} (k + \ell)! M_{k+\ell} h_M(C_2|x_2|^m)$$

holds for every bi-index (k, ℓ) in \mathbf{N}^2 . Then, for every real $\beta > 0$, there exist constants C_3 and C_4 , depending only on C_1, C_2, R, N, m, β and M , such

that for any point x in $(\mathbf{R}^2, 0)$ satisfying $|x_1| < \beta|x_2|^m$, the estimate

$$\left| \partial_1^k \mathbf{Y}^\ell G(x) \right| \leq C_3^{k+\ell+1} (k + \ell)! M_{k+\ell} h_M (C_4 |x_2|^m)$$

holds for every bi-index (k, ℓ) in \mathbf{N}^2 .

Proof. The proof is quite technical and it will be cut into several lemmas. We define, for $0 \leq t \leq 1$ and $x_2 \in \mathbf{R}^*$,

$$Q(x_2, t) = (1 - t)m^{-1}x_2^{1-m}R'(x_2).$$

We also put, for $0 \leq t \leq 1$ and $x \in (\mathbf{R}^2, 0)$,

$$y = y(x, t) = (tx_1 + (1 - t)R(x_2), x_2).$$

When x satisfies $|x_1| < \beta|x_2|^m$, one gets, for some constant C_5 depending only on R ,

$$(17) \quad |y_1| < \beta|x_2|^m + |R(x_2)| < (\beta + C_5)|y_2|^m$$

since $y_2 = x_2$ and $\omega(R) \geq m$.

Lemma 3.2. *For any x_2 in $(\mathbf{R}^*, 0)$, any p in \mathbf{N} and any t in $[0, 1]$, one has*

$$|\mathbf{Y}^p Q(x_2, t)| \leq C_6 C_7^p p! |x_2|^{-mp},$$

where the constants C_6 and C_7 depend only on R and m .

Proof. The assumption $\omega(R) \geq m$ ensures that Q is analytic for x_2 in a neighborhood of 0. We have therefore

$$(18) \quad |\partial_2^j Q(x_2, t)| \leq C_8 C_9^j j!$$

for any $j \in \mathbf{N}$, where the constants C_8 and C_9 depend only on R and m . Besides this, an induction on the integer p gives the expansion

$$(19) \quad \mathbf{Y}^p = \sum_{j=1}^p \mathbf{Y}_j^{(p)}(x_2) \partial_2^j$$

where the coefficients $\mathbf{Y}_j^{(p)}(x_2)$ satisfy the estimate

$$(20) \quad |\mathbf{Y}_j^{(p)}(x_2)| \leq C_{10}^{p+1} (p - j)! |x_2|^{j-mp} \quad \text{for } j = 1, \dots, p.$$

This estimate can be obtained as a very special case of [14, Proposition 2.5]. Finally, Lemma 3.2 is easy to deduce from (18), (19), (20). \square

Lemma 3.3. *For any germ h in $C^\infty(\mathbf{R}^2, 0)$, any point x in $(\mathbf{R} \times \mathbf{R}^*, 0)$, any integer $\ell \geq 1$ and any real t in $[0, 1]$, one has*

$$(21) \quad \mathbf{Y}^\ell (h(y(x, t))) = \sum_{(i,j) \in \mathbf{N}^2; 1 \leq i+j \leq \ell} \Gamma_{i,j}^{(\ell)}(x_2, t) (\partial_1^i \mathbf{Y}^j h)(y(x, t)),$$

where the coefficients $\Gamma_{i,j}^{(\ell)}(x_2, t)$ satisfy, for every integer $k \geq 0$, the estimate

$$(22) \quad \left| \mathbf{Y}^k \Gamma_{i,j}^{(\ell)}(x_2, t) \right| \leq (4C_{11})^\ell (2C_7)^{k+\ell-i-j} (k+\ell-i-j)! |x_2|^{-m(k+\ell-i-j)}$$

with $C_{11} = \max(1, C_6)$.

Proof. We proceed by induction on ℓ . A simple computation shows that

$$\mathbf{Y}(h(y(x, t))) = Q(x_2, t)(\partial_1 h)(y(x, t)) + (\mathbf{Y}h)(y(x, t)),$$

which proves (21) for $\ell = 1$, with $\Gamma_{1,0}^{(1)}(x_2, t) = Q(x_2, t)$ and $\Gamma_{0,1}^{(1)}(x_2, t) = 1$. In this case, the estimate (22) is a direct consequence of Lemma 3.2. Now, an integer ℓ being given, the computation of $\mathbf{Y}(\mathbf{Y}^\ell(h(y(x, t))))$ starting from the expansion (21) gives

$$(23) \quad \Gamma_{i,j}^{(\ell+1)}(x_2, t) = \mathbf{Y}\Gamma_{i,j}^{(\ell)}(x_2, t) + \Gamma_{i,j-1}^{(\ell)}(x_2, t) + Q(x_2, t)\Gamma_{i-1,j}^{(\ell)}(x_2, t),$$

where we have put $\Gamma_{i,j}^{(\ell)} = 0$ for $i < 0$ or $j < 0$, for $i = j = 0$, and for $i+j > \ell$. In virtue of the Leibniz formula, (23) yields

$$(24) \quad \begin{aligned} \mathbf{Y}^k \Gamma_{i,j}^{(\ell+1)}(x_2, t) &= \mathbf{Y}^{k+1} \Gamma_{i,j}^{(\ell)}(x_2, t) + \mathbf{Y}^k \Gamma_{i,j-1}^{(\ell)}(x_2, t) \\ &\quad + \sum_{p+q=k} \frac{k!}{p!q!} (\mathbf{Y}^p Q(x_2, t)) \left(\mathbf{Y}^q \Gamma_{i-1,j}^{(\ell)}(x_2, t) \right). \end{aligned}$$

By induction assumption, both terms $\mathbf{Y}^{k+1} \Gamma_{i,j}^{(\ell)}(x_2, t)$ and $\mathbf{Y}^k \Gamma_{i,j-1}^{(\ell)}(x_2, t)$ are majorized by $(4C_{11})^\ell (2C_7)^{k+\ell+1-i-j} (k+\ell+1-i-j)! |x_2|^{-m(k+\ell+1-i-j)}$. For each term occurring with $p+q=k$ in the sum at the right-hand side of (24), we have also, in virtue of Lemma 3.2,

$$\begin{aligned} \left| (\mathbf{Y}^p Q(x_2, t)) \left(\mathbf{Y}^q \Gamma_{i-1,j}^{(\ell)}(x_2, t) \right) \right| &\leq C_6 C_7^p p! |x_2|^{-mp} (4C_{11})^\ell (2C_7)^{q+\ell+1-i-j} \\ &\quad \cdot (q+\ell+1-i-j)! |x_2|^{-m(q+\ell+1-i-j)}. \end{aligned}$$

Using the elementary fact $p!(q+\ell+1-i-j)! \leq (k+\ell+1-i-j)!$, we deduce, from these estimates and from the definition of C_{11} , that we have

$$(25) \quad \left| \mathbf{Y}^k \Gamma_{i,j}^{(\ell+1)}(x_2, t) \right| \leq (4C_{11})^\ell (2C_7)^{k+\ell+1-i-j} (k+\ell+1-i-j)! |x_2|^{-m(k+\ell+1-i-j)} \mathcal{S},$$

with $\mathcal{S} = 2 + C_{11} \sum_{p+q=k} \frac{k!(q+\ell+1-i-j)!}{q!(k+\ell+1-i-j)!} 2^{-p}$. It is easy to check that one has

$$\frac{k!(q+\ell+1-i-j)!}{q!(k+\ell+1-i-j)!} \leq 1 \quad \text{for } 0 \leq q \leq k \text{ and } i+j \leq \ell+1.$$

We have thus $\mathcal{S} \leq 2 + C_{11} \sum_{0 \leq p \leq k} 2^{-p} \leq 2 + 2C_{11} \leq 4C_{11}$. Therefore the estimate (25) completes the induction and Lemma 3.3 is proved. \square

3.4. End of the proof of Proposition 3.1. Put

$$(26) \quad F(x) = (x_1 - R(x_2))^N G(x)$$

and consider the real analytic change of coordinates near the origin of \mathbf{R}^2 given by $u_1 = x_1 - R(x_2)$, $u_2 = x_2$. In this system of coordinates, (26) becomes

$$F \circ \psi(u) = u_1^N (G \circ \psi)(u) \quad \text{with} \quad \psi(u) = (u_1 + R(u_2), u_2).$$

This yields easily, for $x = \psi(u)$,

$$\begin{aligned} G \circ \psi(u) &= \int_0^1 \frac{(1-t)^{N-1}}{(N-1)!} \frac{\partial^N (F \circ \psi)}{\partial u_1^N} (tu_1, u_2) dt \\ &= \int_0^1 \frac{(1-t)^{N-1}}{(N-1)!} (\partial_1^N F)(y(x, t)) dt. \end{aligned}$$

Taking into account Lemma 3.3 and the fact that ∂_1 and \mathbf{Y} commute, we get thus

$$(27) \quad \partial_1^k \mathbf{Y}^\ell G(x) = \sum_{1 \leq i+j \leq \ell} \int_0^1 \frac{t^k (1-t)^{N-1}}{(N-1)!} \Gamma_{i,j}^{(\ell)}(x_2, t) \left(\partial_1^{i+k+N} \mathbf{Y}^j F \right) (y(x, t)) dt.$$

Now, the assumptions on F in Proposition 3.1, together with the estimate (17) on $y(x, t)$, ensure that one has

$$(28) \quad \left| \left(\partial_1^{i+k+N} \mathbf{Y}^j F \right) (y(x, t)) \right| \leq C_{12}^{i+j+k+N+1} (i+j+k+N+1)! \cdot M_{i+j+k+N+1} h_M(C_{13}|x_2|^m)$$

for any t in $[0, 1]$ and any x in $(\mathbf{R}^2, 0)$ satisfying $|x_1| < \beta|x_2|^m$, where C_{12} and C_{13} are suitable constants depending on β . Besides this, the estimate (22) in Lemma 3.3 gives

$$(29) \quad \left| \Gamma_{i,j}^{(\ell)}(x_2, t) \right| \leq C_{14}^{\ell+1} (\ell-i-j)! |x_2|^{-m(\ell-i-j)}$$

with $C_{14} = \max(4C_{11}, 2C_7)$. We have also, in virtue of (6), the inequality

$$(30) \quad h_M(C_{13}|x_2|^m) \leq (C_{15}|x_2|^m)^{\ell-i-j} M_{\ell-i-j} h_M(C_{15}|x_2|^m)$$

with $C_{15} = \rho(2)C_{13}$. Finally, we have the elementary estimates $(i+j+k+N+1)!(\ell-i-j)! \leq (k+\ell+N+1)! \leq 2^{k+\ell+N+1}(k+\ell)!(N+1)!$ as well as $M_{i+j+k+N+1}M_{\ell-i-j} \leq M_{k+\ell+N+1} \leq A^{k+\ell+N+1}M_{k+\ell}M_{N+1}$ by (2) and (4). Thus, gathering (28), (29) and (30) in (27), we obtain the desired estimate of Proposition 3.1. □

Now, going back to Section 2, we have to obtain bounds for the derivatives of g on the real zero set X_φ . In this purpose, we shall carry out the estimates on X_p for any fixed p with $2p_0 + 1 \leq p \leq p_1$. This will yield a global estimate, first on X_φ^+ , then on the whole of X_φ by symmetry. Of course, if

$2p_0 = p_1$ (which means $X_\varphi^+ = \{0\}$), we have nothing to do. Otherwise, it is clearly enough (since $X_p \cap (\mathbf{R} \times \{0\}) = \{0\}$) to prove a uniform bound on $X_p \cap (\mathbf{R} \times \mathbf{R}_+^*)$. In the sequel, a denotes a point on this set; in other words, we have

$$(31) \quad a_2 > 0 \quad \text{and} \quad a_1 = R_p(a_2^{1/m}).$$

Let H be the half-plane $\{w \in \mathbf{C}; \operatorname{Re} w > 0\}$. For any z in $(\mathbf{C} \times H, 0)$, put

$$(32) \quad \tilde{\varphi}(z) = u(z) \prod_{q=1}^{2p_0} \left(z_1 - U_q(z_2^{1/m}) \right)^{n_q}.$$

We also define, for any x in $(\mathbf{R} \times \mathbf{R}_+^*, 0)$,

$$(33) \quad \tilde{g}(x) = \tilde{\varphi}(x)g(x),$$

in such a way that we have

$$\tilde{g}(x) \prod_{q=2p_0+1}^{p_1} \left(x_1 - R_q(x_2^{1/m}) \right)^{n_q} = (\varphi g)(x).$$

Lemma 3.5. *There exist constants C_{16} and C_{17} such that one has, for any bi-index L ,*

$$|D^L \tilde{g}(a)| \leq C_{16}^{\ell+1} \ell! M_\ell h_M(C_{17}|a|).$$

Proof. For $j = 2p_0 + 1, \dots, p_1 + 1$, put

$$G_j(x) = \tilde{g}(x_1, x_2^m) \prod_{j \leq q \leq p_1} (x_1 - R_q(x_2))^{n_q}.$$

Using (32) and (33), the functions G_j defined above for $x \in (\mathbf{R} \times \mathbf{R}_+^*, 0)$ are easily seen to be, in fact, of class C^∞ in a whole neighborhood of 0 in \mathbf{R}^2 . Moreover, one has

$$(34) \quad (x_1 - R_j(x_2))^{n_j} G_{j+1}(x) = G_j(x),$$

$$(35) \quad G_{2p_0+1}(x) = (\varphi g)(x_1, x_2^m).$$

In particular, we have $\partial_1^k \mathbf{Y}^\ell G_{2p_0+1}(x) = (\partial_1^k \partial_2^\ell (\varphi g))(x_1, x_2^m)$. In virtue of the estimate (11) and the induction relations (34) and (35), it is possible to apply Proposition 3.1 repeatedly with $j = 2p_0 + 1, \dots, p_1$. We get thus

$$(36) \quad \left| \partial_1^k \mathbf{Y}^\ell G_{p_1+1}(x) \right| \leq C_{18}^{k+\ell+1} (k + \ell)! M_{k+\ell} h_M(C_{19}|x_2|^m)$$

for any (k, ℓ) in \mathbf{N}^2 and any x in $(\mathbf{R}^2, 0)$ satisfying

$$(37) \quad |x_1| < \beta |x_2|^m,$$

with suitable constants C_{18} and C_{19} depending on β . Finally, the point $x = (a_1, a_2^{1/m})$ satisfies (37) for some $\beta > 0$ depending only on φ , in virtue of (31) and since $\omega(R_p) \geq m$. Therefore, applying (36) for this choice of x completes the proof of the lemma, since we have $G_{p_1+1}(x) = \tilde{g}(x_1, x_2^m)$. \square

In order to estimate the derivatives of $g = \tilde{g}/\tilde{\varphi}$ at the point a , we finally need some bounds on $1/\tilde{\varphi}$. Such bounds are obtained in the following lemma. Put first

$$\tilde{Z} = \bigcup_{1 \leq q \leq 2p_0} Z_q.$$

Lemma 3.6. *One has*

$$(38) \quad |\tilde{\varphi}(z)| \geq C_{20}d(z, \tilde{Z})^{\tilde{\nu}} \text{ for any } z \in (\mathbf{C} \times H, 0),$$

$$(39) \quad d(a, \tilde{Z}) \geq C_{21}|a|^{d^+(\varphi)},$$

with $\tilde{\nu} = \sum_{1 \leq q \leq 2p_0} n_q$ and constants C_{20}, C_{21} depending only on φ .

Proof. For $z \in \mathbf{C} \times H$, the point $z(q) = (U_q(z_2^{1/m}), z_2)$ belongs to Z_q , hence $d(z, Z_q) \leq |z - z(q)| = |z_1 - U_q(z_2^{1/m})|$. One has also $d(z, \tilde{Z}) \leq d(z, Z_q)$ for $q = 1, \dots, 2p_0$. Taking $\tilde{\nu}$ as indicated, we get thus

$$d(z, \tilde{Z})^{\tilde{\nu}} \leq \prod_{1 \leq q \leq 2p_0} d(z, Z_q)^{n_q} \leq \prod_{1 \leq q \leq 2p_0} |z_1 - U_q(z_2^{1/m})|^{n_q},$$

which, by definition of $\tilde{\varphi}$, proves (38). We have also, using (31) and the same sort of arguments as in Lemma 2.3, $d(a, Z_q) \gtrsim |a_1 - U_q(a_2^{1/m})| = |R_p(a_2^{1/m}) - U_q(a_2^{1/m})| \geq |S_q(a_2^{1/m})| \gtrsim a_2^{\mu_q/m}$ for $q = 1, \dots, 2p_0$. The estimate (39) follows, since (31) implies $|a| \lesssim a_2$. □

Lemma 3.7. *For any bi-index J , one has the estimate*

$$|D^J (1/\tilde{\varphi})(a)| \leq C_{22}^{j+1} j! |a|^{-d^+(\varphi)(j+\tilde{\nu})},$$

where the constant C_{22} depends only on φ .

Proof. Since we have $|a| \lesssim a_2$, we derive $d(a, \tilde{Z}) \lesssim a_2$. Therefore, for a sufficiently small $\epsilon > 0$, the bi-disk of radius $\epsilon d(a, \tilde{Z})$ centered at a lies in $\mathbf{C} \times H$ and we can apply the Cauchy formula, together with the estimate (38) of Lemma 3.6, to obtain

$$|D^J (1/\tilde{\varphi})(a)| \leq C_{23}^{j+1} j! d(a, \tilde{Z})^{-(j+\tilde{\nu})}$$

for some suitable C_{23} . Then the estimate (39) of Lemma 3.6 yields the conclusion. □

3.8. End of the proof of Proposition 2.6. It is now enough to estimate $D^L(\tilde{g}/\tilde{\varphi})(a)$ by means of the Leibniz formula, with the help of Lemmas 3.5 and 3.7. By arguments similar to those in Subsection 2.8, we obtain

$$|D^L g(a)| \leq C_{24}^{\ell+1} \ell! M_\ell^{d^+(\varphi)}$$

for any L in \mathbf{N}^2 and any a in X_p , with $2p_0 + 1 \leq p \leq p_1$, hence any a belonging to X_φ^+ . We have symmetrically

$$|D^L g(a)| \leq C_{25}^{\ell+1} \ell! M_\ell^{d^-(\varphi)}$$

when a belongs to X_φ^- . Clearly, Proposition 2.6 follows. □

4. Further results and comments.

We shall discuss here first the sharpness of Theorem 2.1. In the particular situation $Z_\varphi \cap \mathbf{R}^2 = \{0\}$, it can be obtained as a consequence of [15, Theorem 2.6]. The result is in fact optimal in full generality, as shown by the following theorem.

Theorem 4.1. *For any germ φ in $\mathcal{O}(\mathbf{R}^2, 0)$ and any strongly regular sequence M , one can always find a germ g in $C_{M^{d(\varphi)}}(\mathbf{R}^2, 0)$, flat at the origin, such that φg belongs to $C_M(\mathbf{R}^2, 0)$ and g does not belong to any Carleman class strictly smaller than $C_{M^{d(\varphi)}}(\mathbf{R}^2, 0)$.*

Proof. It goes partly along the same lines as [15, Subsection 3.3], so we shall skip some details. Without losing generality, we can assume $d(\varphi) = d^+(\varphi)$. Put

$$(40) \quad g(x) = \frac{\eta(x_2)}{\tilde{\varphi}(x_1, x_2)} \text{ for } x_2 > 0 \quad \text{and} \quad g(x) = 0 \text{ for } x_2 \leq 0,$$

where η is the special flat germ in $C_M(\mathbf{R}, 0)$ given by [15, Lemma 3.6]; and $\tilde{\varphi}$ has been defined by (32). Proceeding as in [15], it is readily seen that g belongs to $C_{M^{d(\varphi)}}(\mathbf{R}^2, 0)$ but not to any smaller class. Now, the only thing which remains to be checked is that φg belongs to $C_M(\mathbf{R}^2, 0)$. By definition of g in (40), it is enough to obtain estimates in $\mathbf{R} \times \mathbf{R}_+$. For $x_2 \geq 0$, we have

$$(41) \quad (\varphi g)(x_1, x_2^m) = \eta(x_2^m) \prod_{q=2p_0+1}^{p_1} (x_1 - R_q(x_2))^{n_q}.$$

Now, using the flatness of η and the definition of \mathbf{Y} at the beginning of Section 3, we have

$$(42) \quad \left| \mathbf{Y}^\ell (\eta(x_2^m)) \right| \leq D_1^{\ell+1} \ell! M_\ell h_M(D_2 |x_2|^m)$$

for any positive integer ℓ , where D_1 and D_2 are suitable constants. Besides this, in the same way as for Lemma 3.2, we have

$$(43) \quad |\mathbf{Y}^\ell R_q(x_2)| \leq D_3^{\ell+1} \ell! |x_2|^{-m\ell}$$

for $q = 2p_0 + 1, \dots, p_1$. Using (41), (42), (43) and the Leibniz formula, we obtain, by the same sort of arguments as in the end of the proof of Proposition 3.1,

$$(44) \quad \left| \partial_1^k \mathbf{Y}^\ell ((\varphi g)(x_1, x_2^m)) \right| \leq D_4^{k+\ell+1} (k + \ell)! M_{k+\ell} h_M(D_5 |x_2|^m)$$

for any bi-index (k, ℓ) . Applying (44) with $x = (y_1, y_2^{1/m})$, where y is any given point of $\mathbf{R} \times \mathbf{R}_+$, we get thus

$$\left| \partial_1^k \partial_2^\ell (\varphi g)(y) \right| \leq D_4^{k+\ell+1} (k + \ell)! M_{k+\ell} h_M(D_5|y_2|),$$

hence the desired result. □

Next we describe an application of Theorem 2.1 to the study of closed ideals in ultradifferentiable classes. It requires only a few basic facts about topology on Carleman classes, recalled hereafter. For any smoothly bounded neighborhood U of 0 in \mathbf{R}^n and any real $\sigma > 0$, let $C_{M,\sigma}(\overline{U})$ be the set of those functions f belonging to $C^\infty(\overline{U})$ and such that there exists a constant $C_f > 0$ for which we have $|D^L f(x)| \leq C_f \sigma^\ell \ell! M_\ell$ for any x in U and any multi-index L . This set is a Banach space for the norm $\|f\|_{U,\sigma}$ defined as the smallest possible C_f in the previous condition. One can then define the Carleman class $C_M(\overline{U})$ as the inductive limit of all spaces $C_{M,\sigma}(\overline{U})$ for increasing $\sigma > 0$ (we refer the reader to [9] for a detailed study of such spaces). We shall say that the ideal $I_{\varphi,M}$ generated by a real-analytic germ φ in $C_M(\mathbf{R}^n, 0)$ is closed if any germ which, for some suitable neighborhood U of 0, belongs to the closure of the ideal $\varphi C_M(\overline{U})$ generated by φ in $C_M(\overline{U})$, actually belongs to $I_{\varphi,M}$ (that is, it belongs to $\varphi C_M(\overline{V})$ for some neighborhood V of 0, maybe smaller than U). Now, the following result extends [15, Corollary 4.2] to the case of general zero sets. Note that we state it for germs in order to stay in the setting of the paper, but a corresponding theorem for functions can be written as well.

Theorem 4.2. *The ideal $I_{\varphi,M}$ generated by φ in $C_M(\mathbf{R}^2, 0)$ is closed if and only if one has $d(\varphi) = 1$.*

Proof. First, any element f of $C_M(\mathbf{R}^2, 0)$ belonging to the closure of $\varphi C_M(\overline{U})$ in $C_M(\overline{U})$ for some neighborhood U of 0, also belongs to the C^∞ -closure of $\varphi C^\infty(U)$. The Lojasiewicz-Malgrange division theorem [10], [11] ensures that the latter ideal is C^∞ -closed, thus f can be written φg for some g in $C^\infty(\mathbf{R}^2, 0)$. Assuming $d(\varphi) = 1$, Theorem 2.1 implies that g belongs to $C_M(\mathbf{R}^2, 0)$. Therefore the condition is sufficient for closedness. Now, we prove the converse: Let g be the germ constructed in Theorem 4.1. From the argument of Subsection 2.7, we deduce that for any point a in a suitable neighborhood U of 0, there exists some function f_a belonging to the ideal $\varphi C_M(\overline{U})$ and such that $\varphi g - f_a$ is flat at the point a . Applying the ultradifferentiable version of Whitney’s spectral theorem [6] and assuming that $I_{\varphi,M}$ is closed, we derive that φg belongs to $I_{\varphi,M}$, hence g belongs to $C_M(\mathbf{R}^2, 0)$. In virtue of Theorem 4.1, this implies $d(\varphi) = 1$. □

4.3. A discussion of the higher-dimensional case. We shall see here that Theorem 4.1 can be extended to germs in \mathbf{R}^n with $n \geq 3$, provided that,

as in [15], we assume to be in the special situation $Z_\varphi \cap \mathbf{R}^n = \{0\}$, and we replace consequently $d(\varphi)$ by the Łojasiewicz exponent $\tau(\varphi)$ for the regular separation between Z_φ and \mathbf{R}^n . As a consequence, the answer to Problem 2.7 in [15] is affirmative and the assumption $n = 2$ is no longer necessary in [15, Theorem 2.6]. Note that Theorem 4.2 extends as well, since it reduces to [15, Corollary 4.2] in this situation. The scheme of proof, pointed out by Bierstone [1], is as follows:

- First, using the curve-selecting lemma as in [2, Theorem 6.4] or [3], one can find a real-analytic curve $\sigma : [-\varepsilon, \varepsilon] \rightarrow Z_\varphi$, with $\sigma(0) = 0$, which realizes the Łojasiewicz exponent $\tau(\varphi)$; in other words, such that $\tau(\varphi)$ is also the Łojasiewicz exponent for the regular separation between \mathbf{R}^n and $\sigma([-\varepsilon, \varepsilon])$. Now, consider the germ of complex curve Σ defined as the image of $\sigma : (\mathbf{C}, 0) \rightarrow (\mathbf{C}^n, 0)$ and denote by $\tau^+(\varphi)$ the Łojasiewicz exponent for the regular separation between \mathbf{R}^n and $Z_\varphi^+ = Z_\varphi \cap \{z \in \mathbf{C}^n; \operatorname{Re} z_n \geq 0\}$. We have clearly $\sigma([-\varepsilon, \varepsilon]) \subset \Sigma \subset Z_\varphi$, thus $\tau^+(\varphi)$ is also the Łojasiewicz exponent for the regular separation between \mathbf{R}^n and $\Sigma^+ = \Sigma \cap \{z \in \mathbf{C}^n; \operatorname{Re} z_n \geq 0\}$. Write $\sigma(t) = (\sigma_1(t), \dots, \sigma_n(t))$ for $t \in (\mathbf{C}, 0)$. Up to analytic changes of coordinates, we can assume that $\sigma_n(t) = t^m$ for some integer $m \geq 1$, and that $\omega(\sigma_j) \geq m$ for $j = 1, \dots, n - 1$. Put $\sigma_j(t) = X_j(t) + iY_j(t)$, where both X_j, Y_j belong to $\mathcal{O}(\mathbf{R}, 0)$; and put $\mu = \inf \{\omega(Y_j); j = 1, \dots, n - 1\}$ (the integer μ is finite, since we have $\Sigma \cap \mathbf{R}^n = \{0\}$). We claim that

$$(45) \quad \tau^+(\varphi) = \mu/m.$$

Indeed, this can be proved in the same way as [15, Lemma 3.2], reasoning with Σ^+ instead of Z_φ^+ : Consider the map $z \mapsto T(z) = (T_1(z), \dots, T_{n-1}(z))$, where each T_j is a C^1 extension to $(\mathbf{C}^n, 0)$ of the function $z \mapsto z_j - \sigma_j(z_n^{1/m})$, first defined on $\{z \in \mathbf{C}^n; \operatorname{Re} z_n \geq 0\}$. One has then, for $x \in (\mathbf{R}^n, 0)$, the estimate $d(x, T^{-1}(0)) \approx |T_1(x)| + \dots + |T_{n-1}(x)|$ with $|T_j(x)| \approx |x_j - X_j(x_n^{1/m})| + |Y_j(x_n^{1/m})|$, and arguments similar to those in [15] can be applied.

- Now, by the definition of μ , we can write $Y_j(z_n) = z_n^\mu Y_j^*(z_n)$ for $j = 1, \dots, n - 1$, where each Y_j^* belongs to $\mathcal{O}(\mathbf{R}, 0)$. Put $X = (X_1, \dots, X_{n-1})$, $Y^* = (Y_1^*, \dots, Y_{n-1}^*)$ and, for $(t, z_n) \in (\mathbf{C}^2, 0)$,

$$\varphi^*(t, z_n) = \varphi(X(z_n) + tY^*(z_n), z_n).$$

Note that $\varphi^*(iz_n^\mu, z_n^m) = \varphi(\sigma(z_n)) = 0$ for any $z_n \in (\mathbf{C}, 0)$. Moreover, for any real x_n , the roots of the real-analytic germ $t \mapsto \varphi^*(t, x_n^m)$ are non-real, pairwise conjugate. Therefore, this germ is divisible by $(t - ix_n^\mu)(t + ix_n^\mu) = t^2 + x_n^{2\mu}$. Now, as in the proof of Theorem 4.1, we can assume $\tau(\varphi) = \tau^+(\varphi)$ and consider the germ g defined, for

$x \in (\mathbf{R}^n, 0)$, by

$$(46) \quad g(x) = \eta(x_n)/\varphi(x) \text{ for } x_n > 0 \quad \text{and} \quad g(x) = 0 \text{ for } x_n \leq 0.$$

In virtue of [15, Theorem 2.6], g belongs to $C_{M\tau(\varphi)}(\mathbf{R}^n, 0)$ and it is flat at the origin. Conversely, assuming that g belongs to $C_{M^s}(\mathbf{R}^n, 0)$ and replacing x by $(X(x_n) + tY^*(x_n), x_n^m)$ in (46), we get

$$\eta(x_n^m) = (t^2 + x_n^{2\mu})g_1(t, x_n)$$

for some germ $g_1 \in C_{M^s}(\mathbf{R}^2, 0)$. It is now possible to argue just as in [15, Subsection 3.3] to derive the inequality $s \geq \mu/m$, which, in view of (45), yields the desired conclusion.

4.4. Concluding remark. As mentioned in the introduction, the division problem for ultradifferentiable functions has been studied under another angle by Chaumat-Chollet [7]. Roughly speaking, they are able, by means of desingularization, to show that the quotient f/φ enjoys C_{M^+} regularity with $M_\ell^+ = M_\ell^T$ for some $T \geq 1$, not only in the case of two variables, but for higher dimensions as well. In counterpart, they have no control over the number T and, in particular, nothing can be derived about the closedness of the ideal $I_{\varphi, M}$. Therefore, the viewpoint of [7] consists in working afterwards in a special subclass of C_M , say \mathcal{C} , defined as the intersection of all classes C_{M^λ} for $\lambda > 0$. The class \mathcal{C} does not change when M is replaced by M^T ; thus, if f belongs to \mathcal{C} , so does f/φ . In the same spirit, Bos-Milman [4] had previously considered some sort of ‘‘Gevrey estimates’’ under a fairly general, yet less precise, form. In contrast with these results, our approach gives optimal estimates for natural Carleman classes, but does not seem to work in the general higher-dimensional case. Also, we have to assume strong non-quasianalyticity since we use Borel type theorems, whereas [7] does not. It would be very interesting to get rid of both restrictions, especially regarding the dimension.

References

- [1] E. Bierstone, Personal communication (1999).
- [2] E. Bierstone and P.D. Milman, *Semianalytic and subanalytic sets*, Publ. Math. IHES, **67** (1988), 5-42, MR 89k:32011, Zbl 0674.32002.
- [3] J. Bochnak and J.-J. Risler, *Sur les exposants de Lojasiewicz*, Comment. Math. Helvetici, **50** (1975), 493-507, MR 53 #8474, Zbl 0321.32006.
- [4] L. Bos and P.D. Milman, *Sobolev-Gagliardo-Nirenberg and Markov type inequalities on subanalytic domains*, Geom. and Funct. Anal., **5** (1995), 853-923, MR 97e:46038, Zbl 0848.46022.
- [5] J. Bruna, *An extension theorem of Whitney type for non-quasianalytic classes of functions*, J. London Math. Soc., **22** (1980), 495-505, MR 82e:58024, Zbl 0441.26015.

- [6] J. Chaumat and A.-M. Chollet, *Caractérisation des anneaux noëtheriens de séries formelles à croissance contrôlée. Application à la synthèse spectrale*, Publ. Mat., **41** (1997), 545-561, [MR 98k:32010](#), [Zbl 0906.46017](#).
- [7] ———, *Sur la division et la composition dans des classes ultradifférentiables*, Studia Math., **136** (1999), 49-70, [MR 2000k:26033](#), [Zbl 0942.26029](#).
- [8] S. Izumi, *A measure of integrity for local analytic algebras*, Publ. RIMS, Kyoto University, **21** (1985), 719-735, [MR 87i:32014](#), [Zbl 0587.32016](#).
- [9] H. Komatsu, *Ultradistributions, I. Structure theorems and a characterization*, J. Fac. Sci. Univ. Tokyo, Sect. IA, **20** (1973), 25-105, [MR 47 #9277](#), [Zbl 0258.46039](#).
- [10] S. Lojasiewicz, *Sur le problème de la division*, Studia Math., **18** (1959), 87-136, [MR 21 #5893](#), [Zbl 0115.10203](#).
- [11] B. Malgrange, *Ideals of Differentiable Functions*, Oxford University Press (1966), [MR 35 #3446](#), [Zbl 0177.17902](#).
- [12] M.A. Mostow and S. Shnider, *Joint continuity of division of smooth functions I: Uniform Lojasiewicz estimates*, Trans. Amer. Math. Soc., **292** (1985), 573-583, [MR 87f:58018a](#), [Zbl 0603.46034](#).
- [13] H.-J. Petzsche, *On E. Borel's theorem*, Math. Ann., **282** (1988), 299-313, [MR 89m:46076](#), [Zbl 0633.46033](#).
- [14] V. Thilliez, *Sur les fonctions composées ultradifférentiables*, J. Math. Pures et Appl., **76** (1997), 499-524, [MR 99e:58018](#), [Zbl 0878.58008](#).
- [15] ———, *On closed ideals in smooth classes*, Math. Nachr., **227** (2001), 143-157, [MR 2002f:46101](#).
- [16] ———, *Bounds for quotients in rings of formal power series with growth constraints*, Studia Math., **151** (2002), 49-65.
- [17] T. Wang, *Linear Chevalley estimates*, Trans. Amer. Math. Soc., **347** (1995), 4877-4897, [MR 96k:32071](#), [Zbl 0848.32002](#).

Received January 12, 2000

CNRS - UMR 8524

MATHÉMATIQUES - BÂTIMENT M2

UNIVERSITÉ DES SCIENCES ET TECHNOLOGIES DE LILLE

59655 VILLENEUVE D'ASCQ CEDEX

FRANCE

E-mail address: Vincent.Thilliez@univ-lille1.fr