

*Pacific
Journal of
Mathematics*

SPRINGER THEORY IN BRAID GROUPS AND THE
BIRMAN–KO–LEE MONOID

DAVID BESSIS, FRANÇOIS DIGNE, AND JEAN MICHEL

SPRINGER THEORY IN BRAID GROUPS AND THE BIRMAN–KO–LEE MONOID

DAVID BESSIS, FRANÇOIS DIGNE, AND JEAN MICHEL

We state a conjecture about centralizers of certain roots of central elements in braid groups, and check it for braid groups of type A , B , $G(d, 1, r)$ and a couple of other cases. Our proof makes use of results from Birman-Ko-Lee, of which we give a new intrinsic account.

Notations. If G is a group acting on a set X , we denote by X^G the subset of X of elements fixed by all elements of G . If (X, x) is a pointed topological space, we denote by $\Omega(X, x)$ the corresponding loop space, by \sim the homotopy relation on $\Omega(X, x)$ and by $\pi_1(X, x)$ the fundamental group. For all $n \in \mathbf{N}$, we denote by μ_n the set of n -th roots of unity in \mathbf{C} .

0. Introduction.

Springer theory of regular elements (introduced in [Sp]) explains how certain complex reflection groups naturally arise as centralizers in other complex reflection groups of particular elements, the *regular elements*. The construction relies on invariant theory and elementary algebraic geometry, and gives precise information on the relation between the two reflection groups involved (in terms of basic invariants, hyperplane arrangements,...). In their work on Deligne-Lusztig varieties ([BrMi]), Broué and the third author gave a partial interpretation of Springer theory in terms of roots of central elements in braid groups associated to complex reflection groups. Following some of the implicit ideas in [BrMi], we state (and prove in some cases) a conjecture which explains how Springer theory of regular elements can be seen as the image in complex reflection groups of a more general theory holding in braid groups. Of course, Springer's original invariant-theoretical arguments have no analog in braid groups. Instead, we rely on properties of the monoid structure recently discovered by Birman-Ko-Lee ([BiKoLee]) and on a geometric interpretation of this monoid.

Before stating our conjecture, let us recall some properties of complex reflection groups. Let V be a finite dimensional complex vector space and let $W \subset \mathrm{GL}(V)$ be an irreducible finite group generated by complex reflections (that is, elements $s \in \mathrm{GL}(V)$ such that $\ker(s - \mathrm{Id})$ is a hyperplane). We denote by \mathcal{M} the complement in V of the union of all reflecting hyperplanes.

The elements of \mathcal{M} are said to be *regular vectors*. The space \mathcal{M}/W is called the *complement of the discriminant of W* . Choose a basepoint in \mathcal{M} , take its image as basepoint for \mathcal{M}/W . The *braid group \mathbf{B}_W* associated to W is, by definition, the fundamental group of \mathcal{M}/W . The fundamental group of \mathcal{M} is the *pure braid group*, denoted by \mathbf{P}_W . The covering $\mathcal{M} \rightarrow \mathcal{M}/W$ is unramified and one has an exact sequence

$$1 \longrightarrow \mathbf{P}_W \longrightarrow \mathbf{B}_W \longrightarrow W \longrightarrow 1.$$

Let $\zeta \in \mathbf{C}$ be a root of unity of order d . An element $w \in W$ is ζ -*regular*, as defined in [Sp], if and only if it admits a regular eigenvector with eigenvalue ζ or, in other words, if and only if

$$\ker(w - \zeta \text{Id}) \cap \mathcal{M} \neq \emptyset.$$

Assume that there exists a ζ -regular element w (ζ is then said to be a *regular eigenvalue*, its order d is a *regular number*). Let $V(w) := \ker(w - \zeta \text{Id})$ and $\mathcal{M}(w) := \ker(w - \zeta \text{Id}) \cap \mathcal{M}$. The centralizer $C_W(w)$ stabilizes each eigenspace of w . In particular, $C_W(w)$ has a linear representation in $V(w)$. According to the main result in [Sp], this representation is faithful and $C_W(w)$, viewed as a linear group acting on $V(w)$, is a complex reflection group. Moreover, the set of regular vectors for this reflection group is precisely $\mathcal{M}(w)$. Let $\mathbf{B}_{C_W(w)}$ be the braid group associated with $C_W(w)$, i.e., the fundamental group of $\mathcal{M}(w)/C_W(w)$.

Let k be such that $\zeta = e^{2i\pi k/d}$. Assume that the basepoint $x_0 \in \mathcal{M}$ is chosen in $\mathcal{M}(w)$. Let γ be the path:

$$\begin{aligned} \gamma : [0, 1] &\longrightarrow \mathcal{M} \\ t &\longmapsto e^{2i\pi kt/d} x_0. \end{aligned}$$

Let p be the quotient map $\mathcal{M} \rightarrow \mathcal{M}/W$. As noticed in [BrMi] (p. 92), the composed map $p \circ \gamma$ is a loop in \mathcal{M}/W , defining an element $\mathbf{w} \in \mathbf{B}_W$, such that \mathbf{w}^d is central in \mathbf{B}_W , and w is the image of \mathbf{w} by the canonical epimorphism $\mathbf{B}_W \rightarrow W$. In other words, there is a natural way of lifting a regular element of W to a root of a central element in \mathbf{B}_W . Moreover, the natural map $\mathcal{M}(w)/C_W(w) \rightarrow \mathcal{M}/W$ defines a morphism $\mathbf{B}_{C_W(w)} \rightarrow \mathbf{B}_W$, whose image can be easily checked to lie in the centralizer $C_{\mathbf{B}_W}(\mathbf{w})$.

The following conjecture claims that there is an analog of Springer theory in braid groups, where the regular elements have to be replaced by the corresponding roots of central elements:

Conjecture 0.1. *The natural morphism $\mathbf{B}_{C_W(w)} \rightarrow \mathbf{B}_W$ induces an isomorphism between $\mathbf{B}_{C_W(w)}$ and the centralizer $C_{\mathbf{B}_W}(\mathbf{w})$.*

The conjecture can be reformulated in a more intrinsic way:

Let (X, x) be a pointed topological space and G be a group acting on (X, x) (by morphisms in the category of pointed spaces, so $x \in X^G$). The

action of G can be naturally extended to the loop space $\Omega(X, x)$. As clearly $\forall g \in G, \forall \gamma, \gamma' \in \Omega(X, x), \gamma \sim \gamma' \Leftrightarrow g(\gamma) \sim g(\gamma')$, this induces a morphism $G \rightarrow \text{Aut}(\pi_1(X, x))$. For a given G , the construction of $\pi_1(X, x)^G$ from (X, x) is actually functorial, from the category of pointed topological spaces with G -action to the category of groups. Thus the natural injection $(X^G, x) \subseteq (X, x)$ induces a natural morphism from $\pi_1(X^G, x) = \pi_1(X^G, x)^G$ to $\pi_1(X, x)^G$.

If W is as above a finite irreducible complex reflection group in $\text{GL}(V)$, we denote by X the corresponding space \mathcal{M}/W . For all integers m , the inclusion $\mu_m \subset \mathbf{C}^\times$ and the identification of \mathbf{C}^\times with the center of $\text{GL}(V)$ define a natural quotient action of μ_m on X . As noticed in [Be] 1.2, when d is regular and w and \mathbf{w} are as defined above, $\mathcal{M}(w)/C_W(w)$ is homeomorphic to its image X^{μ_d} in X . Let $x \in X^{\mu_d}$. It is an easy calculation to check that the subgroup of $\text{Aut}(\pi_1(X, x))$ generated by the conjugation by \mathbf{w} coincides with the one arising from the action of μ_d on X .

Thus the above conjecture is equivalent to the statement that the natural morphism

$$\pi_1(X^{\mu_d}, x) \rightarrow \pi_1(X, x)^{\mu_d}$$

is an isomorphism.

The fact that the morphism is an isomorphism doesn't depend on the choice of $x \in X^{\mu_d}$.

Our main theorem checks the conjecture for some specific groups (we use the standard notations from [ShTo]):

Theorem 0.2. *Let W be an irreducible complex reflection group. Suppose W is of one of the following types: \mathfrak{S}_n ; $G(p, 1, r)$ with $p > 1$; G_4 ; G_5 ; G_8 ; G_{10} ; G_{16} ; G_{18} ; G_{25} ; G_{26} ; G_{32} . Let X be the complement of the discriminant of W . Let d be a regular number for W , and let $x \in X^{\mu_d}$. The natural morphism*

$$\pi_1(X^{\mu_d}, x) \rightarrow \pi_1(X, x)^{\mu_d}$$

is an isomorphism.

1. The local monoid.

We will deduce our theorem from the particular case where W is the symmetric group \mathfrak{S}_n , and \mathbf{B} the Artin braid group on n strings. In [BiKoLee], Birman, Ko and Lee describe a remarkable monoid for this group. The properties of their monoid will be crucial in our proof. However, contrary to what is done in [BiKoLee], where the Artin braid group is initially given by the Artin presentation, we prefer to use its more intrinsic definition as a

fundamental group. Of course, both viewpoints give (non-canonically) isomorphic groups; however, we believe our reformulation is more natural than the original description.¹

Let X_n be the space of subsets of \mathbf{C} of cardinal n , with its natural topology. In the setting of the introduction, when taking the natural irreducible reflection representation of the symmetric group $W = \mathfrak{S}_n$ on \mathbf{C}^{n-1} , the complement of the discriminant \mathcal{M}/W is homotopy equivalent (in a way compatible with the action of \mathbf{C}^\times) to X_n . We choose the usual direct (= anti-clockwise) orientation on \mathbf{C} .

Let us choose a basepoint $x \in X_n$, and let $B_x = \pi_1(X_n, x)$. We define in this section a monoid M_x , which is a set of group generators for B_x . The structure of M_x depends on the choice of x : When x is taken to be the “usual” basepoint $\{1, \dots, n\}$, the monoid M_x will be isomorphic to the usual Artin monoid; choosing μ_n (the set of n -roots of unity) will yield what we call the Birman-Ko-Lee monoid, which is isomorphic to the one described in [BiKoLee].

If $\gamma \in \Omega(X_n, x)$ and $z \in x$, we denote by γ_z the “string” of γ with origin z . It is a path $[0, 1] \rightarrow \mathbf{C}$, with $\gamma_z(0) = z$ and $\gamma_z(1) \in x$. The element γ is uniquely determined by $(\gamma_z)_{z \in x}$. Conversely, a set of n such strings which do not intersect define an element of $\Omega(X_n, x)$.

Definition 1.1. A pair $\{z, z'\} \subset x$ is said to be *non-crossing* if and only if the closed line segment $[z, z']$ intersects x only at z and z' .

We denote by S_x the set of non-crossing pairs of elements of x .

Examples. The two crucial examples are $x = \{1, \dots, n\}$ and $x = \mu_n$. The corresponding S_x have respectively cardinal $n - 1$ and $n(n - 1)/2$.

For each $\{z, z'\} \in S_x$, we denote by $\delta_{\{z, z'\}} \in B_x$ the generator of the monodromy naturally associated to $[z, z']$, as in the appendix of [Be]. A representative of this element can be defined for instance by the set of strings $\gamma_{z''}(t) = z''$ if $z'' \notin \{z, z'\}$, $\gamma_z(t) = f(z, z', \epsilon)(t)$, $\gamma_{z'}(t) = f(z', z, \epsilon)(t)$ for $\epsilon \in \mathbf{R}_+^*$ small enough (depending only on x) where

$$\forall t, f(z, z', \epsilon)(t) = \frac{z + z'}{2} + \frac{z - z'}{2} \cos(\pi t) + i\epsilon \frac{z - z'}{2} \sin(\pi t)$$

(the corresponding arc is a half-ellipse with great axis $[z, z']$ and small axis of length $\epsilon|z - z'|$. The exact choice of $f(z, z', \epsilon)$ is not important. One could for example replace the half-ellipse by a half-rhomb, or any other variation. However, for later use where defining a loop up to homotopy will not be sufficient, it is convenient for us to define this explicit element of $\Omega(X_n, x)$).

¹After the present article was accepted for publication, the authors were informed that T. Brady had independently obtained a similar description of the Birman-Ko-Lee monoid.

Definition 1.2. The submonoid M_x of B_x generated by

$$\{\delta_{\{z,z'\}} \mid \{z, z'\} \in S_x\}$$

is called the *local monoid* at x .

Clearly, the map $\{z, z'\} \mapsto \delta_{\{z,z'\}}$ is injective, so we may identify S_x with its image in M_x . Let l be the natural length function on B_x (the map $B_x \rightarrow \mathbf{Z}$ induced by the discriminant function $\mathcal{M} \rightarrow \mathbf{C}^\times$). The elements of S_x , being generators of the monodromy, have length 1, so the monoid M_x is \mathbf{N} -graded (only the trivial element being of length 0).

We denote by \prec the left divisibility relation in M_x , i.e.,

$$\forall m, m' \in M_x, m \prec m' \Leftrightarrow \exists m'' \in M_x, mm'' = m'.$$

It results from the \mathbf{N} -grading of M_x that the relation \prec is a partial order on M_x .

We will later study extensively the structure of M_{μ_n} . We complete this section with some first properties which are valid for all x .

Proposition 1.3. *The set S_x is a set of group generators for B_x (i.e., by taking $S_x \cup \{s^{-1} \mid s \in S_x\}$, one has a set of monoid generators for B_x).*

Proof. Distinguish one point $z \in x$. Draw the segments $\{[z, z'] \mid z' \in x\}$. Some of them may be crossing, but by splitting these into smaller ones, one gets a planar graph connecting all points in x and whose edges are non-crossing. The result then follows from the main theorem in [Se] and its reformulation in the appendix of [Be]. □

Notations and conventions. We write $\lambda \vdash x$ to say that λ is a partition of x , in the usual set theoretical sense. If $y \subset x$ and $\lambda \vdash y$, we will also use λ to denote the partition of x obtained by completing λ with parts of cardinal 1. In other words, we will sometimes, for convenience, omit to write the cardinal 1 parts of a partition. By *convex polygon*, we mean either a point (if the number of vertices is 1), a segment (if the number of vertices is 2), or (if the number of vertices is 3 or more) a nondegenerate convex polygon, i.e., such that three vertices never lie on the same line. When $y \subset \mathbf{C}$, we write \bar{y} for the convex hull of y .

We now extend the notion of *non-crossedness* to partitions of x :

Definition 1.4.

- A finite nonempty subset $y \subset \mathbf{C}$ is said to be *convex* if and only if it is the set of vertices of a convex polygon in \mathbf{C} .
- A partition λ of x is said to be *non-crossing* if and only if it satisfies the following two properties:
 - Every part ν in λ is convex.
 - If two parts ν_1 and ν_2 are distinct, their convex hulls $\bar{\nu}_1$ and $\bar{\nu}_2$ do not intersect.

- The set of non-crossing partitions of x is denoted by \mathcal{P}_x . We use the notation $\lambda \models x$ to express that $\lambda \in \mathcal{P}_x$.

We already have a notion of non-crossing pairs. The partition $\{\{z, z'\}\}$ is non-crossing if and only if $\{z, z'\}$ is a non-crossing pair.

Let ν be a part of a non-crossing partition of x . Choose z_1 an arbitrary element of ν . For $i \in \{1, \dots, k\}$, denote by z_i the i -th element of ν for the **clockwise** order on ν starting at z_1 . In the next lemma, when $k = 1$, the product is the empty product, thus the trivial element in M_x .

Lemma 1.5. *The element $\delta_{\{z_1, z_2\}} \delta_{\{z_2, z_3\}} \cdots \delta_{\{z_{k-2}, z_{k-1}\}} \delta_{\{z_{k-1}, z_k\}} \in M_x$ does not depend on the choice of z_1 in ν .*

Proof. This is a consequence of one of the Sergiescu relations (see [Se], 1.1(ii) or [Be], Théorème A.6) namely that

$$\begin{aligned} &\delta_{\{z_1, z_2\}} \delta_{\{z_2, z_3\}} \cdots \delta_{\{z_{k-2}, z_{k-1}\}} \delta_{\{z_{k-1}, z_k\}} \\ &= \delta_{\{z_k, z_1\}} \delta_{\{z_1, z_2\}} \delta_{\{z_2, z_3\}} \cdots \delta_{\{z_{k-2}, z_{k-1}\}}. \end{aligned}$$

The proof of this relation is by induction on k . The case $k = 3$ is checked by a direct computation, and for other k we have

$$\begin{aligned} &\delta_{\{z_1, z_2\}} \delta_{\{z_2, z_3\}} \cdots \delta_{\{z_{k-2}, z_{k-1}\}} \delta_{\{z_{k-1}, z_k\}} \\ &= \delta_{\{z_2, z_3\}} \cdots \delta_{\{z_{k-2}, z_{k-1}\}} \delta_{\{z_{k-1}, z_1\}} \delta_{\{z_{k-1}, z_k\}} \\ &= \delta_{\{z_2, z_3\}} \cdots \delta_{\{z_{k-2}, z_{k-1}\}} \delta_{\{z_{k-1}, z_k\}} \delta_{\{z_k, z_1\}} \end{aligned}$$

where the first equality is by induction and the second by the case $k = 3$. \square

We denote δ_ν the element of 1.5 (when $\{z, z'\}$ is a non-crossing pair, $\delta_{\{\{z, z'\}\}}$ coincides with the element $\delta_{\{z, z'\}}$ defined earlier).

Lemma 1.6. *Let ν and ν' be two finite nonempty subsets of \mathbf{C} . Suppose $\{\nu, \nu'\}$ can be completed to a non-crossing partition of x . Then we have $\delta_\nu \delta_{\nu'} = \delta_{\nu'} \delta_\nu$.*

Proof. The convex hulls of ν and ν' cannot intersect, thus the generators corresponding to their edges commute pairwise. \square

Let $\lambda \models x$. The above lemma makes it natural (and non-ambiguous) to define

$$\delta_\lambda = \prod_{\nu \in \lambda} \delta_\nu.$$

Definition 1.7. Given two partitions λ and λ' of x , we say that λ is *finer* than λ' (or equivalently that λ' is *coarser* than λ), and we write $\lambda \prec \lambda'$, if and only if $\forall \nu \in \lambda, \exists \nu' \in \lambda', \nu \subset \nu'$.

Clearly \prec is a partial order on the set of partitions of x . We intentionally use the same symbol \prec , as for the left divisibility order in M_x . The next proposition justifies this notation.

Let $P_x = \{\delta_\lambda | \lambda \in \mathcal{P}_x\}$. The main result in this section is the following proposition.

Proposition 1.8.

- (i) *The left divisibility order in P_x coincides with the restriction of \prec from M_x to P_x ; in other words,*

$$\forall p, p' \in P_x, (\exists p'' \in P_x, pp'' = p') \Leftrightarrow (\exists m'' \in M_x, pm'' = p').$$

We denote this partial order by \prec .

- (ii) *The map*


$$\begin{aligned} D : \mathcal{P}_x &\longrightarrow P_x \\ \lambda &\longmapsto \delta_\lambda \end{aligned}$$

is a poset isomorphism from (\mathcal{P}_x, \prec) to (P_x, \prec) .

Before proving the proposition, we need some definitions and lemmas.

Definition 1.9. Let ν be a convex subset of \mathbf{C} , and let ν' be a subset of ν . Let z_1, \dots, z_k be a clockwise numbering of the elements of ν , such that $z_{i_1}, \dots, z_{i_{k'}}$ is a clockwise numbering of ν' with $1 = i_1 < i_2 < \dots < i_{k'} \leq k$. Then we denote by $\nu' \setminus \nu$ (ν “cut at” ν') the partition with parts the sets $\{z_{i_j}, z_{i_j+1}, \dots, z_{i_{j+1}-1}\}$ (where for $j = k'$ we make the convention that $i_{k'+1} = k + 1$).

Example. In the picture below, the grey points are the points of ν' , the other points of ν are black, and the parts of $\nu' \setminus \nu$ are enclosed by dashed curves.



Note that $\nu' \setminus \nu$ is well-defined (does not depend on the chosen numbering) and is a non-crossing partition of ν . We will need the following alternative description of $\nu' \setminus \nu$: Its parts of cardinal ≥ 2 are the intersections of ν with each connected component of the complement of $\overline{\nu'}$ in $\overline{\nu}$, to which has been added the element of ν' just before the connected component (in clockwise order).

Denote by ϕ_x the natural epimorphism from B_x to \mathfrak{S}_x . It maps $\delta_{\{z,z'\}}$ to the transposition (z, z') . Thus, when $\lambda \models x$, it is clear by construction that λ is the orbit decomposition for the action in x of $\phi_x(\delta_\lambda)$. Thus if $\lambda \neq \lambda'$, then $\phi_x(\delta_\lambda)$ and $\phi_x(\delta_{\lambda'})$ have different cycle decompositions, and are different. We have proved the:

Lemma 1.10. *The restriction of ϕ_x to P_x is injective.*

Lemma 1.11.

- (i) *Let ν be a convex subset of \mathbf{C} , and let ν' be a subset of ν . We have $\delta_\nu = \delta_{\nu'}\delta_{\nu'\setminus\nu}$.*
- (ii) *Suppose $\lambda, \lambda' \models x$, λ' has only one part of cardinal ≥ 2 , and λ is finer than λ' . Then there exists a unique non-crossing partition of x , which we denote by $\lambda \setminus \lambda'$ such that $\delta_\lambda \delta_{\lambda \setminus \lambda'} = \delta_{\lambda'}$.*
- (iii) *Suppose $\lambda, \lambda' \models x$ and λ is finer than λ' . Then there exists a unique non-crossing partition of x , which we denote by $\lambda \setminus \lambda'$ such that $\delta_\lambda \delta_{\lambda \setminus \lambda'} = \delta_{\lambda'}$.*

Proof. We prove (i) by induction on the cardinality of ν' . With the notations of 1.9, let $\nu_1 = \{z_1, \dots, z_{i_2-1}\}$, $\nu_2 = \nu - \nu_1$ so that $\delta_\nu = \delta_{\nu_1} \delta_{\{z_{i_2-1}, z_{i_2}\}} \delta_{\nu_2}$, and let $\nu'_2 = \nu' - \{z_1\}$, so that $\delta_{\nu'} = \delta_{\{z_1, z_{i_2}\}} \delta_{\nu'_2}$. Then by 1.5 applied to $\nu_1 \cup \{z_{i_2}\}$ we have $\delta_\nu = \delta_{\{z_1, z_{i_2}\}} \delta_{\nu_1} \delta_{\nu_2}$. By the induction hypothesis, we have $\delta_{\nu_2} = \delta_{\nu'_2} \delta_{\nu'_2 \setminus \nu_2}$. As $\nu' \setminus \nu = \{\nu_1\} \cup \nu'_2 \setminus \nu_2$, and as $\delta_{\nu'_2}$ commutes to δ_{ν_1} (since $\overline{\nu'_2} \cap \overline{\nu_1} = \emptyset$), we get the result by induction.

(ii): Denote by ν the only nontrivial part of λ' , and by $\lambda_1, \dots, \lambda_l$ the nontrivial parts of λ . To prove the result, the essential step is to notice that since for $i > 1$ we have $\overline{\lambda_i} \cap \overline{\lambda_1} = \emptyset$, each λ_i lies inside a single connected component of the complement of $\overline{\lambda_1}$ in $\overline{\nu}$, thus each is included in a part of $\lambda_1 \setminus \nu$. Thus (ii) follows by induction on the number of parts of λ from (i). The uniqueness comes from the fact that the identity $\delta_\lambda \delta_{\lambda \setminus \lambda'} = \delta_{\lambda'}$ is valid in $M_x \subset B_x$, and B_x is a group.

Finally, (iii) is easily obtained by applying (ii) to all the nontrivial parts of λ' . □

For $\sigma \in \mathfrak{S}_x$, we denote by $|\sigma|$ the minimum number of transpositions in a decomposition of σ as a product of transpositions; such a decomposition is called *reduced* (note that we allow all transpositions). The map $\sigma \mapsto |\sigma|$ is not a morphism, but we have the relation $|\sigma_1 \sigma_2| \leq |\sigma_1| + |\sigma_2|$.

Lemma 1.12.

- (i) *Let $\lambda \vdash x$ be the orbit decomposition of $\sigma \in \mathfrak{S}_x$. We have*

$$|\sigma| = \sum_{\nu \in \lambda} (|\nu| - 1) = |x| - |\lambda|.$$

- (ii) Let $\sigma \in \mathfrak{S}_x$. Let $t = (z, z')$. If z and z' are in the same orbit of σ , then $|\sigma t| = |\sigma| - 1$. If z and z' are in different orbits of σ , then $|\sigma t| = |\sigma| + 1$.
- (iii) Let $\sigma \in \mathfrak{S}_x$. Let $t_1 t_2 \dots t_{|\sigma|}$ be a reduced decomposition of σ . Then if $t_i = (z, z')$ is one of the transpositions, z and z' belong to the same orbit of σ . Let $\lambda \vdash x$ be the orbit decomposition of σ and, for $i = 1, \dots, |\sigma|$, $\lambda_i \vdash x$ be the orbit decomposition of $t_1 t_2 \dots t_i$. Then $\lambda_i \prec \lambda$.
- (iv) For all $m \in M_x$, we have $l(m) \geq |\phi_x(m)|$.
- (v) If $\lambda \models x$, we have $l(\delta_\lambda) = |\phi_x(\delta_\lambda)|$.

Proof. (i) is easy.

(ii) comes from (i) and the following remark: When z and z' are in the same orbit, multiplying by t splits this orbit into two orbits, thus increasing by 1 the number of orbits; when z and z' are in different orbits, multiplying by t merges their orbits, thus decreasing by 1 the number of orbits.

(iii) is an easy induction from (ii) and its proof: As the decomposition is reduced, the relation $|\sigma| = |t_1 t_2 \dots t_{|\sigma|}|$ can only be achieved if the successive multiplications by the t_i merge orbits.

(iv) and (v) are easy. □

We can now prove Proposition 1.8:

Proof. (i) Let $\lambda, \lambda' \in \mathcal{P}_x$. As the converse implication is trivial, we only have to check that

$$(\exists m'' \in M_x, \delta_\lambda m'' = \delta_{\lambda'}) \Rightarrow (\exists \lambda'' \in \mathcal{P}_x, \delta_\lambda \delta_{\lambda''} = \delta_{\lambda'}).$$

Let $m'' \in M_x$ such that $\delta_\lambda m'' = \delta_{\lambda'}$. Consider the image in \mathfrak{S}_x of this identity: $\phi_x(\delta_\lambda) \phi_x(m'') = \phi_x(\delta_{\lambda'})$. By Lemma 1.12 (v), we have $|\phi_x(\delta_\lambda)| = l(\delta_\lambda)$ and $|\phi_x(\delta_{\lambda'})| = l(\delta_{\lambda'})$, and, using Lemma 1.12 (iv), we have $|\phi_x(\delta_{\lambda'})| = l(\delta_{\lambda'}) = l(\delta_\lambda m'') = l(\delta_\lambda) + l(m'') \geq |\phi_x(\delta_\lambda)| + |\phi_x(m'')|$ and thus $|\phi_x(\delta_{\lambda'})| = |\phi_x(\delta_\lambda)| + |\phi_x(m'')|$. Consequently, when concatenating reduced decompositions for $\phi_x(\delta_\lambda)$ and $\phi_x(m'')$, one gets a reduced decomposition for $\phi_x(\delta_{\lambda'})$. By Lemma 1.12 (iii), this implies that the orbit decomposition for $\phi_x(\delta_\lambda)$ is finer than the one for $\phi_x(\delta_{\lambda'})$, i.e., $\lambda \prec \lambda'$. We conclude by Lemma 1.11 (iii).

(ii) By Lemma 1.11 (iii), the map D is a poset morphism. It is injective: From δ_λ , one recovers λ by considering the orbit decomposition of $\phi_x(\delta_\lambda)$. The fact that the inverse map is a poset morphism, i.e., $\delta_\lambda \prec \delta_{\lambda'} \Rightarrow \lambda \prec \lambda'$, has already been obtained in our proof of (i). □

Remark 1.13. If x is convex, then there is a largest element $\delta_{\{x\}}$ in the poset (P_x, \prec) (which corresponds to the coarsest partition which has just one part equal to x).

2. Pre-Garside structures and Garside monoids.

The existence of nice normal forms in the Birman-Ko-Lee monoid will be of crucial importance in the proof of our main theorem. This property is

proved in [BiKoLee], and we could have just quoted and translated it into our setting. However, it appeared that our geometric interpretation allows us to give a new proof, far less computational, of some of the main results in [BiKoLee]. The Birman-Ko-Lee monoid is a *Garside monoid*, in the sense of [DePa], and this implies (among others) the existence of the normal form. To give a simple proof of this fact, we make use a new criterion of “Garsiditude”. This criterion relies on the notion of *pre-Garside structure*, which can be seen as an axiomatization of the context in which most of the proofs in [Mi] are actually valid.

When A and B are two sets, we mean by “partial map” from A to B a datum consisting of a subset $A' \subset A$ and a map $f : A' \rightarrow B$. It is convenient to refer to A' implicitly, and to use a slightly abusive language, e.g., we will write “ $f(a)$ is defined” instead of “ $a \in A'$ ”.

Definition 2.1. Let P be a set. An atomic partial product on P is a partial map $m : P \times P \rightarrow P$ (we will denote $m(a, b)$ by $a.b$ or ab), satisfying the following axioms:

- (i) (Unit element and associativity.) There exists an element $1 \in P$ such that for all $a \in P$, both $1.a$ and $a.1$ exist and are equal to a . For any $a, b, c \in P$, it is equivalent for ab and $(ab)c$ to be defined or for bc and $a(bc)$ to be defined and then $a(bc) = (ab)c$.
- (ii) (Finite number of atoms.) Let $P^* = P - \{1\}$; the image of $P^* \times P^*$ is in P^* , and the complement $S = P^* - m(P^* \times P^*)$ is finite (the elements of S are the *atoms* of (P, m)).
- (iii) (Grading.) There exists a function $l : P \rightarrow \mathbf{N}$ such that $p \in P^* \Rightarrow l(p) > 0$ and $l(ab) \geq l(a) + l(b)$ whenever ab is defined.

If $a, b \in P$ are such that ab is defined, we say that a (resp. b) is a *left* (resp. *right*) *divisor* of ab . Conversely, ab is a *right* (resp. *left*) *multiple* of a (resp. b).

Definition 2.2. Suppose P is a set together with an atomic partial product. The associated monoid $M(P)$ is the monoid defined by the following presentation:

- As a set of generators we take P .
- For relations we take $ab = c$ whenever $a, b, c \in P$ are such that ab is defined in P and equal to c .

Note that if P is a subset of a monoid M' , and if the partial product on P is a restriction of the monoid law in M' , then there is a natural morphism $M(P) \rightarrow M'$. If P generates M' , the morphism is surjective.

As in [Mi], we note that we can identify $M(P)$ to the set of finite sequences of elements of P , quotiented by the minimal equivalence relation \sim such that

$$(p_1, \dots, p_{i-1}, p_i, p_{i+1}, \dots, p_n) \sim (p_1, \dots, p_{i-1}, a, b, p_{i+1}, \dots, p_n)$$

whenever the product ab is defined and equal to p_i .

Let (x_1, \dots, x_n) be a sequence of elements of P . There are $(2(n-1))!/(n!(n-1)!)$ different ways to put brackets on the product $x_1 \dots x_n$. By an obvious induction from the associativity axiom, if the product is defined in P for one of these bracketings, then it is defined in P for any other bracketing, and the value of this product does not depend on the choice of the bracketing. When this is the case, we write $x_1 \dots x_n$ for this product.

Lemma 2.3. *Let (x_1, \dots, x_n) be a sequence of elements of P equivalent, in $M(P)$, to a single term sequence (y) . Then the product $x_1 \dots x_n$ is defined in P and we have $x_1 \dots x_n = y$.*

Proof. By assumption (x_1, \dots, x_n) can be transformed into (y) by a finite rewriting process

$$l_0 = (x_1, \dots, x_n) \rightarrow l_1 \rightarrow \dots \rightarrow l_k = (y)$$

in which, at each step, the elementary transformation $l_{j-1} \rightarrow l_j$ is

- either of the type

$$(p_1, \dots, p_{i-1}, a, b, p_{i+1}, \dots, p_m) \rightarrow (p_1, \dots, p_{i-1}, p_i, p_{i+1}, \dots, p_m)$$

- or of the type

$$(p_1, \dots, p_{i-1}, p_i, p_{i+1}, \dots, p_m) \rightarrow (p_1, \dots, p_{i-1}, a, b, p_{i+1}, \dots, p_m)$$

where a, b are such that their product is defined in P and equal to p_i . Suppose the length k of the rewriting process is minimal.

Suppose some of the transformations are of the second type, and chose j maximal such that $l_{j-1} \rightarrow l_j$ is of the second type

$$(p_1, \dots, p_{i-1}, p_i, p_{i+1}, \dots, p_m) \rightarrow (p_1, \dots, p_{i-1}, a, b, p_{i+1}, \dots, p_m).$$

As all further steps are of the first type, we have $k = j + m$ and the product $p_1 \dots p_{i-1} a b p_{i+1} \dots p_m$ is defined in P (and is equal to y). Choosing a bracketing starting by $\dots (ab) \dots$, we see that the product $p_1 \dots p_{i-1} p_i p_{i+1} \dots p_m$ must also be defined in P (and equal to y). But this yields a rewriting of length $j + m - 2 < k$ and we have a contradiction.

Thus there are no transformation of the second type in a minimal rewriting. The result follows. □

The following proposition is a straightforward consequence of the lemma:

Proposition 2.4.

- (i) *The natural map from P to $M(P)$ is injective.*
- (ii) *If $x \in M(P)$ divides $a \in P$ then $x \in P$.*

A *right common multiple* of s and t is an element u which is a right multiple of both s and t . A common right multiple u of s and t is a *least right common multiple* if and only if, for all common right multiple v of s

and t, u is a left divisor of v . Similarly, there are notions of (least) left common multiple, of (greatest) right common divisor and of (greatest) left common divisor.

Definition 2.5. A *pre-Garside* structure on a set P is an atomic partial product, with set of atoms S , satisfying the following additional axioms:

- (iv) If two elements of S have a common right multiple in P , they have a least common right multiple. When $s, t \in S$ have a least right common multiple, we write it $\Delta_{s,t}$.
- (iv') If two elements of S have a common left multiple in P , they have a least common left multiple.
- (v) If $s, t \in S$ have a common right multiple in P , and if $a \in P$ is such that $as \in P$ and $at \in P$, then $a\Delta_{s,t} \in P$.
- (vi) For all $m \in M(P)$ and $a, b \in P$, if either $am = bm$ or $ma = mb$, then $a = b$.

In [Mi], S was taken to be the set of usual Artin generators, and P was the set of reduced braids. To handle the Birman-Ko-Lee monoid, we will take for S the set S_{μ_n} and for P the set P_{μ_n} .

Condition (v) is satisfied e.g., if there exists a common right multiple in P of all elements of P , which is the case in the usual braid monoid and also in the Birman-Ko-Lee monoid (see 1.13). Note that such an element is necessarily unique and if it exists, the finiteness of S implies that P is finite. We shall see (2.22) that its existence is equivalent to the fact that all pairs of elements of S have a right lcm.

The following lemmas and propositions are rephrasings of [Mi], 1.4 to 1.9.

Lemma 2.6. *Let X be a finite subset of $M(P)$ such that:*

- *If $x \in X, a \in M(P), a \prec x$ then $a \in X$.*
- *If $a \in M(P), s, t \in S, as, at \in X$, then $\Delta_{s,t}$ exists and $a\Delta_{s,t} \in X$.*

Then there exists $g \in X$ such that X is the set of left divisors of g .

Proof. The statement follows [Mi] 1.4, and the proof is exactly the same (after having extended the length function to $M(P)$ by setting $l(m) := \max\{\sum_{i=1}^n l(p_i) | p_1, \dots, p_n \in P, m = p_1 \dots p_n\}$). □

Proposition 2.7. *Any two elements of P have a left g.c.d. in P .*

Proof. We can follow the proof of [Mi] 1.6, replacing 1.5 in [Mi] by Axiom (iv). □

We shall denote by $a \wedge b$ the left g.c.d. of a and b .

Proposition 2.8. *For a and b in P there exists a unique maximal $c \prec b$ such that $ac \in P$.*

Proof. We apply Lemma 2.6 to the set X of c such that $c \prec b$ and $ac \in P$. To check the assumptions of that lemma, we need that if $s, t \in S$ and if $cs, ct \in X$ then $\Delta_{s,t}$ exists and $ac\Delta_{s,t} \in P$ (then we have $c\Delta_{s,t} \in X$). Since cs and ct divide $b \in P$, by cancellability and 2.4 s and t have a common multiple in P , so by Axiom (iv) $\Delta_{s,t}$ exists and by Axiom (v) $ac\Delta_{s,t} \in P$. \square

Definition 2.9. In the situation of 2.8 we denote by $\alpha_2(a, b)$ the element ac and we denote by $\omega_2(a, b)$ the unique $d \in P$ such that $b = cd$. We thus have $ab = \alpha_2(a, b)\omega_2(a, b)$.

Note that the uniqueness of ω_2 follows from Axiom (vi).

Proposition 2.10. For $a, b, c, ab \in P$ we have $\alpha_2(ab, c) = \alpha_2(a, \alpha_2(b, c))$.

Proof. The statement is [Mi], 1.8 and the proof is the same. \square

Proposition 2.11. For $a, b, c, ab \in P$ we have

$$\omega_2(ab, c) = \omega_2(a, \alpha_2(b, c))\omega_2(b, c).$$

Proof. By Propositions 2.8 and 2.10 the products of both sides with $\alpha_2(ab, c)$ are equal. By Axiom (vi) we will be done if we show that $\omega_2(a, \alpha_2(b, c))\omega_2(b, c)$ is in P . By definition 2.9 there exists $z \in P$ such that $\alpha_2(b, c) = bz$ and $c = z\omega_2(b, c)$. As $ab \prec abz$ and $ab \in P$ we have $\alpha_2(a, bz) = abz_1$ for some $z_1 \prec z$ (by definition of α_2 and cancellability in P). Hence $bz = bz_1\omega_2(a, bz)$ and $z = z_1\omega_2(a, bz)$. So $z_1\omega_2(a, \alpha_2(b, c))\omega_2(b, c) = z_1\omega_2(a, bz)\omega_2(b, c) = z\omega_2(b, c) = c$. The result follows since any divisor of c is in P by 2.4. \square

We now extend the definition of α_2 to $M(P)$, following [Mi], 2.1 to 2.6. All the proofs of [Mi] can be reproduced, replacing Proposition 1.5 in [Mi] by Axiom (iv).

Proposition 2.12. There is a unique function $\alpha : M(P) \rightarrow P$ extending the identity of P and satisfying $\alpha(ab) = \alpha_2(a, b)$ for $a, b \in P$ and $\alpha(gh) = \alpha(g\alpha(h))$ for $g, h \in M(P)$. Moreover $\alpha(g)$ is the unique maximal element in $\{c \in P | c \prec g\}$.

Proposition 2.13. There exists a unique function $\omega : M(P) \rightarrow M(P)$ such that $\omega(ab) = \omega(a, b)$ for $a, b \in P$ (in particular $\omega(a) = 1$ for $a \in P$) and such that $\omega(gh) = \omega(g\alpha(h))\omega(h)$ for $g, h \in M(P)$.

Proposition 2.14. Let $g \in M(P)$; then $\omega(g)$ is the unique $y \in M(P)$ such that $g = \alpha(g)y$.

Proposition 2.15. The monoid $M(P)$ has left and right cancellation property (i.e., in Axiom (vi) we can replace the condition $a, b \in P$ by $a, b \in M(P)$).

Proposition 2.16. If $s, t \in S$ divide $a \in M(P)$ on the left, then $\Delta_{s,t}$ exists and divides a .

Proposition 2.17. *For $a, b \in M(P)$ there exists a unique maximal $c \in M(P)$ (for \prec) such that $c \prec a$ and $c \prec b$.*

We shall still denote this left g.c.d. by $a \wedge b$.

Proposition 2.18. *A family of elements of $M(P)$ which has a right (resp. left) common multiple has a right (resp. left) lcm in $M(P)$.*

Proof. Assume that all elements of the family $\{a_i\}_{i \in I}$ divide c . If we can apply 2.6 to the set X of elements of $M(P)$ which divide all common multiples of the a_i , it will give the result. Let us check the assumption of 2.6. This set X is finite as it is included in the set of divisors of c . The first assumption of 2.6 is clearly satisfied. The second assumption is a consequence of the fact that if $s, t \in S$ are such that xs and xt divide some element xz , then s and t divide $\alpha(z)$, so $\Delta_{s,t}$ exists and divides z , whence $x\Delta_{s,t}$ divides xz . \square

We have a more precise result for elements of P :

Proposition 2.19. *If a family of elements of P has a common right (resp. left) multiple in $M(P)$ then its right (resp. left) lcm exists and is in P .*

Proof. The lcm exists by the preceding proposition. Let m be this lcm. Any divisor of m divides $\alpha(m) \in P$, whence the result. \square

We now get a normal form for any element of $M(P)$ exactly as in [Mi].

Definition 2.20. A decomposition (g_1, \dots, g_n) of an element $g_1 \dots g_n$ of $M(P)$ is said to be its normal form if no g_i is equal to 1 and for any i we have $g_i = \alpha(g_i \dots g_n)$.

The following statement is 4.2 of [Mi] and the same proof applies.

Proposition 2.21. *A decomposition (g_1, \dots, g_i) with $g_i \in P$ is a normal form if and only if (g_i, g_{i+1}) is a normal form for any i . In particular any segment (g_i, \dots, g_j) of a normal form (g_1, \dots, g_n) is a normal form.*

In the same way, statements 4.6 to 4.9 and 5.1 to 5.3 of [Mi] generalize to our setting.

Assume that all elements of P have a right common multiple (which is in P by 2.19). By 2.18, this is the same as assuming that any finite subset of elements of P has a lcm. As already noticed, the existence of a lcm Δ of all elements in P implies the finiteness of P . The converse is not true, but we have:

Proposition 2.22. *The elements of P have a common right multiple if and only if P is finite and any pair of elements of S has a common multiple.*

Proof. Implication “only if” is clear. The converse is an immediate application of 2.6, taking $X = P$. \square

By 2.12, for any $a \in M(P)$ we have $\alpha(a) = a \wedge \Delta$.

Proposition 2.23. *Assume P has a right lcm Δ .*

- *There is an automorphism $x \mapsto \bar{x}$ of $M(P)$ mapping S to itself and such that $x\Delta = \Delta\bar{x}$.*
- *The element Δ is the left lcm of P .*

Proof. It is sufficient to define the automorphism on P . As Δ is a right multiple of all elements of P , for any $a \in P$ there exists a unique $\Delta_a \in P$ such that $\Delta = a\Delta_a$ and there exists a unique $\bar{a} \in P$ such that $\Delta = \Delta_a\bar{a}$, so that $a\Delta = a\Delta_a\bar{a} = \Delta\bar{a}$. The map $a \mapsto \bar{a}$ is injective by cancellability in $M(P)$ and is thus compatible with the product. As P is finite, it is surjective. If $a \in S$, by surjectivity \bar{a} cannot be the product of two nontrivial elements of P , so has to be in S by Axiom (ii).

As $a \mapsto \bar{a}$ is a bijection of P , the above proof shows that Δ is a left multiple of all elements of P . So it is the left lcm of P . □

We can now compare our formalism with the one in [DePa]: If P is a pre-Garside structure with right common multiples, then, it is readily seen that any two elements of $M(P)$ have left and right common multiples: Indeed, if $m_1, m_2 \in M(P)$ are both products of n or less elements of S , then Δ^n is a left and right multiple of both m_1 and m_2 (use Proposition 2.22: The conjugation by Δ maps S into itself). Thus, by Proposition 2.18, pairs of elements in $M(P)$ have both a right lcm and a left lcm. Moreover, by Proposition 2.22, Δ is both the right lcm and the left lcm of P . Conversely, elements of P are left and right divisors of Δ , and, by Proposition 2.4, Δ has no other left or right divisors. We have proved that $M(P)$ is a Garside monoid, as defined in [DePa].

Conversely, if M is a Garside monoid, then the restriction of the monoid product to the set P of simple elements (see [DePa]) is a pre-Garside structure with common multiples.

We have proved the:

Theorem 2.24. *Let P be set, endowed with a pre-Garside structure, such that all atoms have a common right multiple. Then $M(P)$ is a Garside monoid.*

Conversely, when M is a Garside monoid with fundamental element Δ , the set P of divisors of Δ has a pre-Garside structure for the partial product obtained by restriction of the product in M ; all elements of P have a common right multiple and $M \simeq M(P)$.

Note that our approach already gives interesting results (e.g., 2.26 below) even when some elements of P have no common multiple, e.g., in the case of braid groups attached to infinite type Coxeter groups; in this case, $M(P)$ is no longer a Garside monoid. More important is, for the present article,

that pre-Garside structures provide us with a convenient criterion to check that the Birman-Ko-Lee monoid is a Garside monoid.

Let us recall the following proposition about Garside monoids, which is already in [DePa].

Proposition 2.25. *Assume that P has a right lcm. The monoid $M(P)$ injects into the group $G(P)$ having the same presentation and any element of G can be written uniquely as $x^{-1}y$ with $x, y \in M(P)$ and $x \wedge y = 1$.*

Proof. The statement follows [Mi] 3.2 and the same proof applies. \square

We now generalize [Mi] 4.4.

Proposition 2.26. *Let Γ be a group of automorphisms of $M(P)$ stabilizing S . Let Σ be the set of lcms of Γ -orbits in S which exist and are not the product of other such lcms; then P^Γ has a pre-Garside structure with atoms Σ ; the monoid $M(P)^\Gamma$ identifies with $M(P^\Gamma)$. If moreover we assume that P has a right lcm then the group $G(P)^\Gamma$ identifies with $G(P^\Gamma)$ (cf. 2.25).*

Proof. Axiom (i) holds as it holds in P and as the product of two Γ -fixed elements is Γ -fixed. Let $x \in P^\Gamma$, and let $s \in S$ such that $s \prec x$; then x is divisible by all elements in the orbit of s , so is divisible by their lcm (which exists). So P^Γ is generated by the lcms of Γ -orbits in S which exist, thus by Σ , and we have arranged for elements of Σ to be atoms, so Axiom (ii) is satisfied. The length inherited from P is still compatible with the product so we have Axiom (iii) (but note that the elements of Σ may have length greater than 1 even if all elements of S had length 1). The lcm of two elements of P^Γ is Γ -fixed by its uniqueness and is in P by 2.19, whence Axioms (iv) and (iv'). If σ and τ are in Σ and have a lcm $\Delta_{\sigma,\tau}$ and if $a \in P^\Gamma$ is such that $a\sigma$ and $a\tau$ are in P then $a\Delta_{\sigma,\tau}$ is the lcm of $a\sigma$ and $a\tau$ so is in P by 2.19, and is clearly Γ -fixed, whence Axiom (v). The monoid $M(P^\Gamma)$ is by definition a submonoid of $M(P)$, so Axiom (vi) holds.

Let $x \in M(P)^\Gamma$; the uniqueness of its normal form implies that each term is in P^Γ . On the other hand, if $x \in M(P)^\Gamma$, as $\alpha(x)$ is the unique maximal element in P dividing x it is also the unique maximal element in P^Γ dividing x , so, by the definition of normal forms, the normal form of an element of $M(P)^\Gamma$ is a normal form in $M(P^\Gamma)$. This shows that $M(P)^\Gamma$ identifies with $M(P^\Gamma)$. The same argument shows that $G(P)^\Gamma$ identifies with $G(P^\Gamma)$ when Δ exists, as Δ is in P^Γ . \square

3. The Birman-Ko-Lee monoid.

Let $x \in X_n$, and let S_x , P_x and M_x as defined as in Section 1.

Lemma 3.1. *The restriction of the product in M_x defines an atomic partial product in P_x with set of atoms S_x .*

Proof. Axiom (i) is a consequence of the associativity in the monoid M_x and of 1.8(i). Axioms (ii) and (iii) are easy consequences of the existence of the length function $l : M_x \rightarrow \mathbf{N}$ introduced in Section 1. The atoms are clearly, by construction, the elements of S_x . \square

We call the monoid M_{μ_n} the *Birman-Ko-Lee monoid* (whenever x is convex, we have $M_x \simeq M_{\mu_n}$).

Remark 3.2. If we map the base point μ_n to the usual base point $\{1, \dots, n\}$ by choosing $z_1 \in \mu_n$ and numbering clockwise the elements of μ_n starting at z_1 as z_1, z_2, \dots, z_n , then the $\delta_{\{z_i, z_{i+1}\}}$ map to the usual Artin generators σ_i of $\pi_1(X_n, \mu_n)$ and the $\delta_{\{z_i, z_j\}}$ map to the generators a_{ij} considered in [BiKoLee] (note that they always suppose $i > j$ while we don't order the pairs $\{z, z'\}$). Then the elements of P_{μ_n} map to the *canonical factors* defined in [BiKoLee]. In order to compare our definition to that of [BiKoLee] one should note that [BiKoLee] call a_{ij}, a_{kl} an “obstructing pair” exactly when the partition $\{\{z_i, z_j\}, \{z_k, z_l\}\}$ is crossing.

The monoid M_{μ_n} enjoys remarkable properties which are consequences of the discussion in the previous section and of the following theorem.

Theorem 3.3. *The restriction of the product in M_{μ_n} to a partial product in P_{μ_n} is a pre-Garside structure, and P_{μ_n} has a lcm δ .*

Proof. Since μ_n is convex, $\delta = \delta_{\{\mu_n\}}$ is an lcm of P_{μ_n} , as explained in 1.13. This eliminates the need to check Axiom (v).

Let us now prove Axiom (iv). Using the isomorphism in Proposition 1.8, and noticing that any pair of elements of μ_n defines a non-crossing edge, we have to prove: For all $z_1, z'_1, z_2, z'_2 \in \mu_n$, the set

$$E = \{\lambda \mid \mu_n \mid \{\{z_1, z'_1\}\} \prec \lambda \text{ and } \{\{z_2, z'_2\}\} \prec \lambda\}$$

has a minimum element for \prec . We discuss by cases:

- Assume $[z_1, z'_1] \cap [z_2, z'_2] = \emptyset$. Then let $\lambda = \{\{z_1, z'_1\}, \{z_2, z'_2\}\}$. Clearly, $\lambda \in E$, and any partition coarser than $\{\{z_1, z'_1\}\}$ and $\{\{z_2, z'_2\}\}$ must be coarser than λ .
- Assume $[z_1, z'_1] \cap [z_2, z'_2] \neq \emptyset$. Let $\lambda = \{\{z_1, z'_1, z_2, z'_2\}\}$ (the nontrivial part may have three or four elements). Clearly $\lambda \in E$. Now let $\lambda' \in E$. Consider ν_1 the part of λ' in which z_1 lies, ν_2 the part of λ' in which z_2 lies. Because $\{\{z_1, z'_1\}\} \prec \lambda'$ and $\{\{z_2, z'_2\}\} \prec \lambda'$, we have $z'_1 \in \nu_1$ and $z'_2 \in \nu_2$. As λ' is non-crossing, we must have $\nu_1 = \nu_2$ (otherwise we would have $\overline{\nu_1} \cap \overline{\nu_2} = \emptyset$, which contradicts $[z_1, z'_1] \cap [z_2, z'_2] \neq \emptyset$). Thus λ is finer than λ' .

(iv') is proved similarly.

To prove (vi), consider the natural morphism $M(P_{\mu_n}) \rightarrow M_{\mu_n}$ and its composition with the epimorphism $M_{\mu_n} \rightarrow \mathfrak{S}_{\mu_n}$. Let $m \in M(P_{\mu_n})$, $\lambda, \lambda' \mid =$

μ_n such that either $\delta_\lambda m = \delta_{\lambda'} m$ or $m\delta_\lambda = m\delta_{\lambda'}$. Denote by σ the image of m in \mathfrak{S}_{μ_n} . We have $\phi_{\mu_n}(\delta_\lambda)\sigma = \phi_{\mu_n}(\delta_{\lambda'})\sigma$ or $\sigma\phi_{\mu_n}(\delta_\lambda) = \sigma\phi_{\mu_n}(\delta_{\lambda'})$. As \mathfrak{S}_{μ_n} is a group, this implies $\phi_{\mu_n}(\delta_\lambda) = \phi_{\mu_n}(\delta_{\lambda'})$. By Lemma 1.10, the restriction of ϕ_{μ_n} to P_{μ_n} is injective, thus we have as required $\delta_\lambda = \delta_{\lambda'}$. \square

Remark 3.4. According to the isomorphism in Proposition 1.8, the above theorem implies that the poset of non-crossing partitions of μ_n is a lattice. If $\lambda_1, \lambda_2 \models \mu_n$, the set

$$E = \{\lambda \models \mu_n \mid \lambda_1 \prec \lambda \text{ and } \lambda_2 \prec \lambda\}$$

admits a minimum element. We leave to the reader to check that this can be proved directly, using the following arguments: Given two (possibly crossing) partitions, the set of (possibly crossing) partitions coarser than the two partitions admits a minimum element; given a (possibly crossing) partition, the set of non-crossing partitions which are coarser admits a minimum element.

Of course, by construction, M_x embeds in B_x for all $x \in X_n$. What is specific to the Birman-Ko-Lee monoid is the following result, which is the analog in our intrinsic setting of the embedding theorem in [BiKoLee].

Corollary 3.5. *The natural morphism $M(P_{\mu_n}) \rightarrow M_{\mu_n}$ is an isomorphism.*

Proof. Consider the group $G(P_{\mu_n})$ given by the group presentation corresponding to the monoid presentation of $M(P_{\mu_n})$. As the partial product in P_{μ_n} is a restriction of the one in B_{μ_n} , there is a natural morphism

$$G(P_{\mu_n}) \rightarrow B_{\mu_n}.$$

By Proposition 1.3, this morphism is surjective.

Note that it is easy to adapt Sergiescu’s presentations to find a presentation of B_{μ_n} where the generators corresponds to the elements of S_{μ_n} . A set of defining relations is for example given in [BiKoLee], Proposition 2.1:

- $\delta_{\{z_1, z'_1\}}\delta_{\{z_2, z'_2\}} = \delta_{\{z_2, z'_2\}}\delta_{\{z_1, z'_1\}}$ when $\{\{z_1, z'_1\}, \{z_2, z'_2\}\}$ is non-obstructing,
- $\delta_{\{z_1, z_2\}}\delta_{\{z_2, z_3\}} = \delta_{\{z_2, z_3\}}\delta_{\{z_3, z_1\}} = \delta_{\{z_3, z_1\}}\delta_{\{z_1, z_2\}}$ for z_1, z_2, z_3 coming in clockwise order.

These relations are valid in P_{μ_n} , thus in $G(P_{\mu_n})$, and the morphism $G(P_{\mu_n}) \rightarrow B_{\mu_n}$ is an isomorphism.

We conclude using the natural commutative diagram:

$$\begin{array}{ccc} M(P_{\mu_n}) & \twoheadrightarrow & M_{\mu_n} \\ \downarrow & & \downarrow \\ G(P_{\mu_n}) & \xrightarrow{\sim} & B_{\mu_n} \end{array}$$

and the injectivity of the map $M(P_{\mu_n}) \rightarrow G(P_{\mu_n})$ (Proposition 2.25). \square

We identify $M(P_{\mu_n})$ and M_{μ_n} through the natural isomorphism. As M_{μ_n} is a Garside monoid, we have nice normal forms, an algorithm for the word problem, . . . The following property is the one we need for the proof of our main theorem.

Proposition 3.6. *Let $\zeta = e^{\frac{2i\pi}{n}}$.*

- (i) *For $\lambda \models \mu_n$ the automorphism of M_{μ_n} induced by δ maps δ_λ to $\delta_\zeta \lambda$.*
- (ii) *For d dividing n , the centralizer of $\delta^{n/d}$ in $\pi_1(X_n, \mu_n)$ is generated by the elements δ_λ for $\lambda \models \mu_n$ such that $\zeta^{n/d} \lambda = \lambda$.*

Proof. It is an immediate consequence of 1.5 that (i) holds for a generator of the form $\delta_{\{s_i, s_{i+1}\}}$ where s_i and s_{i+1} are two consecutive points in a numbering of μ_n . It follows that it holds for any generator $\delta_{\{s_1, s_a\}}$ by using that $\delta_{\{s_1, s_a\}} = \delta_{\{s_1, s_2\}} \cdots \delta_{\{s_{a-1}, s_a\}} \delta_{\{s_{a-2}, s_{a-1}\}}^{-1} \cdots \delta_{\{s_1, s_2\}}^{-1}$ (which follows also from 1.5), and it follows thus for any element of P . Part (ii) is then a consequence of 2.26. □

Remark. Note that 2.26 gives a “Birman-Ko-Lee” presentation of $C_{\mathbf{B}}(\delta^i)$. Let us work out an example to show that one has to take only a part of the lcms of δ^i -orbits on S to get an atomic set. Take $n = 6$ and $i = 2$. Then the δ^2 -orbits in S are

$$\{\delta_{\{s_1, s_2\}}, \delta_{\{s_3, s_4\}}, \delta_{\{s_5, s_6\}}\}, \{\delta_{\{s_2, s_3\}}, \delta_{\{s_4, s_5\}}, \delta_{\{s_1, s_6\}}\},$$

$$\{\delta_{\{s_1, s_3\}}, \delta_{\{s_3, s_5\}}, \delta_{\{s_1, s_5\}}\} \text{ and } \{\delta_{\{s_1, s_4\}}, \delta_{\{s_2, s_5\}}, \delta_{\{s_3, s_6\}}\}$$

whose respective lcm are

$$\delta_{(\{s_1, s_2\}, \{s_3, s_4\}, \{s_5, s_6\})}, \delta_{(\{s_2, s_3\}, \{s_4, s_5\}, \{s_1, s_6\})}, \delta_{\{s_1, s_3, s_5\}} \text{ and } \delta;$$

but $\delta = \delta_{\{s_1, s_3, s_5\}} \delta_{(\{s_1, s_2\}, \{s_3, s_4\}, \{s_5, s_6\})}$, so must be eliminated.

4. A geometric normal form for canonical factors.

Let n be a positive integer. If K_1, K_2 are non-intersecting compact subsets of \mathbf{C} , we denote by $d(K_1, K_2)$ the positive number

$$\inf_{(z_1, z_2) \in K_1 \times K_2} |z_1 - z_2|.$$

It is clear that

$$\inf_{\lambda \models \mu_n} \inf_{\substack{\nu_1, \nu_2 \in \lambda \\ \nu_1 \neq \nu_2}} d(\overline{\nu_1}, \overline{\nu_2}) > 0.$$

We define

$$\epsilon_n := \frac{1}{3} \inf_{\lambda \models \mu_n} \inf_{\substack{\nu_1, \nu_2 \in \lambda \\ \nu_1 \neq \nu_2}} d(\overline{\nu_1}, \overline{\nu_2}).$$

The exact value is not important, we will only use the fact that ϵ_n is fixed and small enough. In particular, the reader should check for himself that in the

following definition, the strings do not intersect and thus γ_λ is well-defined (note that the function f has been defined in the first section).

Definition 4.1. Let $\lambda \models \mu_n$.

- Let $z \in \mu_n$. We define a path $\gamma_z : [0, 1] \rightarrow \mathbf{C}$ in the following way:
 - If z is in a part of λ with exactly one element then we set

$$\forall t, \gamma_z(t) = z.$$

- If z is in a part of λ with exactly two elements z and z' we set:

$$\forall t, \gamma_z(t) = f(z, z', \epsilon_n)(t).$$

- If z is in a part ν of λ with three or more elements, we denote by z' the element of ν immediately after z in the direct (i.e., anti-clockwise) cyclic order on the vertices of $\bar{\nu}$ and we set:

$$\forall t, \gamma_z(t) = z + t(z' - z).$$

- These strings uniquely determine an element of $\Omega(X_n, \mu_n)$ which we denote by γ_λ .

Lemma 4.2. Let $\lambda \models \mu_n$, and let $\zeta = e^{\frac{2i\pi}{n}}$.

- The loop γ_λ represents δ_λ .
- $\forall z \in \mu_n, \gamma_{z \cdot \lambda} = z \cdot \gamma_\lambda$.
- Let $\lambda' \models \mu_n$. We have $\lambda = \lambda' \Leftrightarrow \delta_\lambda = \delta_{\lambda'} \Leftrightarrow \gamma_\lambda = \gamma_{\lambda'}$.
- Let $k \in \mathbf{N}$. Denote by d the order of the root of unity ζ^k . Then

$$\zeta^k \lambda = \lambda \Leftrightarrow \delta^k \delta_\lambda = \delta_\lambda \delta^k \Leftrightarrow \gamma_\lambda \in \Omega(X_n^{\mu_d}, \mu_n).$$

Proof. The first three statements are obvious consequences of the previous definition.

As $\delta^{-k} \delta_\lambda \delta^k = \delta_{\zeta^k \lambda}$, we have, using (iii),

$$e^{\frac{2ik\pi}{n}} \lambda = \lambda \Leftrightarrow \delta^k \delta_\lambda = \delta_\lambda \delta^k \Leftrightarrow \gamma_{\zeta^k \lambda} = \gamma_\lambda.$$

By (ii), we have

$$\gamma_{\zeta^k \lambda} = \gamma_\lambda \Leftrightarrow \zeta^k \gamma_\lambda = \gamma_\lambda.$$

It is clear that $\zeta^k \gamma_\lambda = \gamma_\lambda$ is equivalent to $\gamma_\lambda \in \Omega(X_n^{\mu_d}, \mu_n)$. We have proved (iv). \square

5. Proof of the theorem.

We start with one particular case.

Proposition 5.1. Let $r, d \in \mathbf{N}$. Let $n = dr$. The inclusion $X_n^{\mu_d} \subset X_n$ induces an isomorphism

$$\pi_1(X_n^{\mu_d}, \mu_n) \xrightarrow{\sim} \pi_1(X_n, \mu_n)^{\mu_d}.$$

Proof. The injectivity part of the proposition has been proved in [Be]. Let us prove the surjectivity. Let $\zeta = e^{\frac{2i\pi}{d}}$. Multiplication by ζ coincides with conjugating by $\delta^{n/d}$, so, by 3.6(ii), we know that $\pi_1(X_n, \mu_n)^{\mu_d}$ is generated by the δ_λ such that $\zeta\lambda = \lambda$. Using (iv) of Lemma 4.2, this means that $\gamma_\lambda \in \Omega(X_n^{\mu_d}, \mu_n)$. Thus any such δ_λ is in the image of $\pi_1(X_n^{\mu_d}, \mu_n)$. So we have proved that the image of $\pi_1(X_n^{\mu_d}, \mu_n)$ is $\pi_1(X_n, \mu_n)^{\mu_d}$. \square

We now have to study the other type of regular numbers for \mathfrak{S}_n . We denote by ν_n the basepoint of X_n defined by $\nu_n := \mu_{n-1} \cup \{0\}$.

Proposition 5.2. *Let $r, d \in \mathbf{N}$. Let $n = dr + 1$. The inclusion $X_n^{\mu_d} \subset X_n$ induces an isomorphism*

$$\pi_1(X_n^{\mu_d}, \nu_n) \xrightarrow{\sim} \pi_1(X_n, \nu_n)^{\mu_d}.$$

Proof. For all integer m , let us denote by X_m^* the space of configurations of m points in \mathbf{C}^* , with its natural topology.

There is a natural inclusion $X_m^* \subset X_m$ and a natural injection $X_m^* \subset X_{m+1}$ defined by $x \mapsto x \cup \{0\}$. The action of μ_d on X_m restricts to an action on X_m^* . Thus we have a commutative diagram of continuous maps:

$$\begin{array}{ccccc} X_n^{\mu_d} & \longleftarrow & X_{dr}^{*\mu_d} & \longrightarrow & X_{dr}^{\mu_d} \\ \downarrow & & \downarrow & & \downarrow \\ X_n & \longleftarrow & X_{dr}^* & \longrightarrow & X_{dr} \end{array}$$

According to [Be], Lemme 3.1., the first line consists of homeomorphisms. Consider the following commutative diagram of group morphisms:

$$\begin{array}{ccccc} \pi_1(X_n^{\mu_d}, \nu_n) & \xleftarrow{\sim} & \pi_1(X_{dr}^{*\mu_d}, \mu_{dr}) & \xrightarrow{\sim} & \pi_1(X_{dr}^{\mu_d}, \mu_{dr}) \\ \downarrow & & \downarrow & & \downarrow \\ \alpha \left(\pi_1(X_n, \nu_n) \right. & \xleftarrow{A} & \pi_1(X_{dr}^*, \mu_{dr}) & \xrightarrow{B} & \left. \pi_1(X_{dr}, \mu_{dr}) \right) \beta \\ \uparrow & & \uparrow & & \uparrow \\ \pi_1(X_n, \nu_n)^{\mu_d} & \xleftarrow{a} & \pi_1(X_{dr}^*, \mu_{dr})^{\mu_d} & \xrightarrow{b} & \pi_1(X_{dr}, \mu_{dr})^{\mu_d} \end{array}$$

where α and β are defined by functoriality, as in the introduction. According to our Proposition 5.1, β is an isomorphism. What we want to prove is that α is also an isomorphism. This will result from the fact that both a and b are isomorphisms.

- The map a is an isomorphism: By an easy (and almost classical) argument, one can see that A is injective and identifies $\pi_1(X_{dr}^*, \mu_{dr})$ with the subgroup of $\pi_1(X_n, \nu_n)$ consisting of elements whose associated

permutations of ν_n fix the point 0. As clearly $\pi_1(X_n, \nu_n)^{\mu_d}$ is included in this subgroup, a is an isomorphism.

- The map b is an isomorphism: As noticed in [Be] (fact (*), used in the proof of Theorem 3.2.(I), page 14), there is an exact sequence


$$1 \longrightarrow F_n \longrightarrow \pi_1(X_{dr}^*, \mu_{dr}) \xrightarrow{B} \pi_1(X_{dr}, \mu_{dr}) \longrightarrow 1$$

where F_n is the free group on n generators and such that the action of μ_d permutes without fixed points the images of the generators of F_n . As a consequence, the intersection of $\ker B$ with $\pi_1(X_{dr}^*, \mu_{dr})^{\mu_d}$ is trivial, and b is injective. The surjectivity of b results from the surjectivity of B and from the μ_d -equivariance of the diagram. □


Together, the last two propositions prove the part of Theorem 0.2 about symmetric groups, where the only regular numbers are the divisors of n and $n - 1$.

Now consider the irreducible reflection group $G(p, 1, n)$, denote by $X_{p,n}$ the complement of its discriminant. As $G(p, 1, n)$ is the centralizer of a p regular element of \mathfrak{S}_{pn} , we can identify $X_{p,n}$ and $X_{pn}^{\mu_p}$ (in a way compatible with the action of μ_p).

Let d be a regular number for $G(p, 1, n)$. Consider the following diagram of inclusions:



where e is the lcm of d and p . By Proposition 5.1, the π_1 -images of downward arrows factorize through isomorphisms with centralizers in $\pi_1(X_{pn}, \mu_{pn})$ (all base points being μ_{pn}):



Using the identification between $X_{p,n}$ and $X_{pn}^{\mu_p}$, the part of the theorem about $G(p, 1, n)$ says that the natural morphism $\pi_1(X_{pn}^{\mu_e}) \rightarrow \pi_1(X_{pn}^{\mu_p})$ induces an isomorphism $\pi_1(X_{pn}^{\mu_e}) \xrightarrow{\sim} \pi_1(X_{pn}^{\mu_p})^{\mu_d}$; this is obvious on the above diagram, which identifies $\pi_1(X_{pn}^{\mu_e})$ with $\pi_1(X_{pn})^{\mu_e}$ and $\pi_1(X_{pn}^{\mu_p})$ with $\pi_1(X_{pn})^{\mu_p}$.

To complete the proof of the theorem, it is enough to notice that the exceptional groups mentioned are Shephard groups (see [OrSo]) and that their discriminants are isomorphic to ones we have already studied.

References

- [Be] D. Bessis, *Groupes des tresses et éléments réguliers*, J. Reine Angew. Math., **518** (2000), 1-40, [MR 2001c:20082](#), [Zbl 0961.20033](#).
- [BiKoLee] J. Birman, K.H. Ko and S.J. Lee, *A new approach to the word and conjugacy problem in the braid groups*, Adv. Math., **139(2)** (1998), 322-353, [MR 99m:20082](#), [Zbl 0937.20016](#).
- [BrMi] M. Broué and J. Michel, *Sur certains éléments réguliers des groupes de Weyl et les variétés de Deligne-Lusztig associées*, Proceedings de la Semaine de Luminy 'Représentations des groupes réductifs finis,' 73-139, Birkhäuser, 1996, [MR 98h:20077](#).
- [DePa] P. Dehornoy and L. Paris, *Gaussian groups and Garside groups, two generalizations of Artin groups*, Proc. of London Math. Soc., **79** (1999), 569-604, [MR 2001f:20061](#).
- [Mi] J. Michel, *A note on words in braid monoids*, J. of Algebra, **215** (1999), 366-377, [MR 2000b:20046](#), [Zbl 0937.20017](#).
- [OrSo] P. Orlik and L. Solomon, *Discriminants in the invariant theory of reflection groups*, Nagoya Math. J., **109** (1988), 23-45, [MR 89i:32025](#), [Zbl 0614.20032](#).
- [Se] V. Sergiescu, *Graphes planaires et présentations des groupes de tresses*, Math. Z., **214(3)** (1993), 477-490, [MR 95j:20034](#), [Zbl 0819.20040](#).
- [ShTo] G.C. Shephard and J.A. Todd, *Finite unitary reflection groups*, Canad. J. Math., **6** (1954), 274-304, [MR 15,600b](#), [Zbl 0055.14305](#).
- [Sp] T.A. Springer, *Regular elements of finite reflection groups*, Invent. Math., **25** (1974), 159-198, [MR 50 #7371](#), [Zbl 0287.20043](#).

Received October 4, 2000 and revised January 2, 2001.

YALE UNIVERSITY
DEPARTMENT OF MATHEMATICS
PO BOX 208 283, NEW HAVEN CT 06520-8283
E-mail address: db278@math.yale.edu

LAMFA, UNIVERSITÉ DE PICARDIE-JULES VERNE
33, RUE SAINT-LEU
80039 AMIENS CEDEX
FRANCE
E-mail address: digne@u-picardie.fr

LAMFA, UNIVERSITÉ DE PICARDIE-JULES VERNE
33, RUE SAINT-LEU
80039 AMIENS CEDEX
FRANCE

INSTITUT DE MATHÉMATIQUES
UNIVERSITÉ PARIS VII
175, RUE DU CHEVALERET
75013 PARIS
FRANCE
E-mail address: jmichel@math.jussieu.fr