

*Pacific  
Journal of  
Mathematics*

MATRIX  $A_p$  WEIGHTS VIA MAXIMAL FUNCTIONS

MICHAEL GOLDBERG

Volume 211 No. 2

October 2003



## MATRIX $A_p$ WEIGHTS VIA MAXIMAL FUNCTIONS

MICHAEL GOLDBERG

The matrix  $A_p$  condition extends several results in weighted norm theory to functions taking values in a finite-dimensional vector space. Here we show that the matrix  $A_p$  condition leads to  $L^p$ -boundedness of a Hardy-Littlewood maximal function, then use this estimate to establish a bound for the weighted  $L^p$  norm of singular integral operators.

### 1. Preliminaries.

Weighted Norm theory forms a basic component of the study of singular integrals. Here one attempts to characterize those measure spaces over which a broad class of singular integral operators remain bounded. For the case of singular integral operators on  $\mathbf{C}$ -valued functions in Euclidean space, the answer is given by the Hunt-Muckenhoupt-Wheeden theorem [10]. It states that the necessary and sufficient condition for boundedness in  $L^p(d\mu)$  is that  $d\mu = W(x) dx$  and the function  $W$  satisfies the  $A_p$  condition, namely:  $\left(\frac{1}{|B|} \int_B W dx\right)^{1/p} \left(\frac{1}{|B|} \int_B W^{-p'/p} dx\right)^{1/p'} \leq C$  for all balls  $B \subset \mathbf{R}^n$ .

The  $A_p$  condition requires considerable interpretation in order to apply it to weighted measures of  $\mathbf{C}^d$ -valued functions. First, the weight  $W(x)$  should take values in the space of positive  $d \times d$  Hermitian forms. This raises concerns about the order in which products are taken, since matrices need not commute, and also what it means for the quantity on the left-hand side to be uniformly bounded. Treil' [21] conjectured that the correct statement of the matrix  $A_2$  condition should be

$$\sup_B \left\| \left( \frac{1}{|B|} \int_B W dx \right)^{1/2} \left( \frac{1}{|B|} \int_B W^{-1} dx \right)^{1/2} \right\| < \infty$$

where exponents  $1/2$  indicate operator powers of a nonnegative matrix. This was subsequently proven in [23] and again in [24].

If  $p$  is different from 2, the matrix  $A_p$  condition cannot be written in terms of averages of operator powers of weight  $W$ . Averages still play a crucial role, however it is more accurate to regard  $W(x)$  as a Banach space norm on  $\mathbf{C}^d$  rather than a matrix. A correct formulation of the matrix  $A_p$  condition, which is also the subject of this note, first appeared in [12] and [25]. Because their statements do not appear similar, it is especially

important to understand what properties matrix  $A_p$  weights share with their scalar counterparts. This is discussed further in the next section.

Boundedness estimates on singular integral operators were originally obtained by way of the Hardy-Littlewood maximal function  $M$ . If a scalar weight  $W$  possesses the  $A_\infty$  property (several equivalent definitions are given in [18]), then the  $L^p$  norm of any singular integral is dominated by the  $L^p$  norm of  $M$  via a distributional argument commonly known as the good- $\lambda$  inequality. The  $A_p$  condition is specifically required to ensure that  $\|Mf\|_{L^p(W)} \leq C\|f\|_{L^p(W)}$ .

Some of these techniques fail to generalize to the case of vector-valued functions with matrix weights. There is no known analogue of the  $A_\infty$  property to create simultaneous estimates for every exponent  $p$ . The weak- $L^p(W)$  spaces used to prove boundedness of the Hardy-Littlewood maximal function are not well defined in this setting. In general, much of the ability to compare objects and dominate one by another is lost when the objects are vectors rather than scalars. The theory of matrix weights has consequently evolved along much different lines. One fundamental technique employed in both [23] and [25] is to choose a good basis (often inspired by Haar functions) in  $L^p(W)$  and consider the integral operator as a matrix acting on the coefficient space. Estimates may then be made separately on the matrix and on the coefficient embedding operator. Even in the scalar case these ideas have yielded new results and new ways of approaching weighted norm problems.

In this note we attempt to tackle the difficulties of extending the classical theory, or else circumvent them. Some arguments may be borrowed nearly word for word, some remain intact only if they are presented in a specific manner. Our hope is to discover which properties of scalar  $A_p$  weights admit some generalization to the case of vector-valued functions and matrix weights, leading to a more complete understanding of the matrix  $A_p$  class.

Let  $T$  be a singular integral operator associated to kernel  $K(x)$  in the sense that  $Tf(x) = \int_{\mathbf{R}^n} K(x-y)f(y)dy$  for almost every  $x$  outside the support of  $f$ . The following regularity hypotheses are to be assumed for  $K$ :

$$(1) \quad |K(x)| \leq C|x|^{-n} \text{ and } |\nabla K(x)| \leq C|x|^{-n-1}$$

and additionally we suppose that for some choice of  $p$ ,  $1 < p < \infty$ , the bound  $\|Tf\|_{L^p} \leq A\|f\|_{L^p}$  holds for all  $f \in L^p$ . One may then apply  $T$  to functions taking values in  $\mathbf{C}^d$  by allowing it to act separately on each coordinate function, that is:  $(Tf)_j = Tf_j$ . This new operator, also denoted by  $T$ , is a singular integral operator whose associated convolution kernel is  $K$  times the identity matrix.

In a similar manner, define the truncated operators  $T_\epsilon$  to be convolution with  $K_\epsilon(x) = \chi_{\{|x|>\epsilon\}}K(x)$  for all  $\epsilon > 0$ . Note that  $T$  and the  $T_\epsilon$  all commute

with pointwise multiplication by any constant matrix  $\Lambda$ , in other words  $\Lambda T f = T(\Lambda f)$ .

A matrix weight  $W$  is a function on  $\mathbf{R}^n$  taking values in  $d \times d$  positive-definite matrices, with weighted norm space  $L^p(W)$  defined by

$$(2) \quad \|f\|_{L^p(W)}^p = \int_{\mathbf{R}^n} |W^{1/p} f|^p dx.$$

One is often concerned with the relationship between a weight and its average over arbitrary balls. The most straightforward notion of an average,  $W_B = \frac{1}{|B|} \int_B W dx$ , turns out to be useful only in the study of  $L^2(W)$ . With any exponent  $p \neq 2$ , this does not properly respect the structure of the underlying  $L^p$ -space. The following definitions are needed instead:

A metric  $\rho = \rho_x(\cdot)$  denotes a family of Banach space norms on  $\mathbf{C}^d$ , indexed by  $x \in \mathbf{R}^n$ . The weighted norm space  $L^p(\rho)$  is given by

$$\|f\|_{L^p(\rho)}^p = \int_{\mathbf{R}^n} [\rho_x(f(x))]^p dx.$$

Note that for any matrix weight  $W$ ,  $L^p(W)$  is isometrically equivalent to  $L^p(\rho)$  with the metric  $\rho_x(\mathbf{e}) = |W^{1/p}(x)\mathbf{e}|$ . Given a ball  $B \subset \mathbf{R}^n$  and an exponent  $p > 1$ , let  $\rho_{p,B}$  be defined by the formula

$$\rho_{p,B}(\mathbf{e}) = \left( \frac{1}{|B|} \int_B [\rho_x(\mathbf{e})]^p dx \right)^{1/p}.$$

This will be our method for averaging the metric  $\rho$  over a ball  $B$ .

The dual metric  $\rho^*$  is defined pointwise in  $x$  to be

$$\rho_x^*(\mathbf{e}) = \sup_{\mathbf{f} \in \mathbf{C}^d} \frac{|(\mathbf{e}, \mathbf{f})|}{\rho_x(\mathbf{f})}.$$

One immediate consequence is that  $(\mathbf{e}, \mathbf{f}) \leq \rho_x^*(\mathbf{e})\rho_x(\mathbf{f})$ .

**Proposition 1.1.** *For any  $\mathbf{e} \in \mathbf{C}^d$  and any ball  $B \subset \mathbf{R}^n$ ,*

$$\rho_{p',B}^*(\mathbf{e}) \geq (\rho_{p,B})^*(\mathbf{e}).$$

*Proof.* Given two vectors  $\mathbf{e}, \mathbf{f} \in \mathbf{C}^d$ ,

$$\begin{aligned} (\mathbf{e}, \mathbf{f}) &\leq \frac{1}{|B|} \int_B \rho_x^*(\mathbf{e})\rho_x(\mathbf{f}) dx \\ &\leq \left( \frac{1}{|B|} \int_B [\rho_x^*(\mathbf{e})]^{p'} dx \right)^{1/p'} \cdot \left( \frac{1}{|B|} \int_B [\rho_x(\mathbf{f})]^p dx \right)^{1/p} \\ &= \rho_{p',B}^*(\mathbf{e})\rho_{p,B}(\mathbf{f}). \end{aligned}$$

In other words,  $\rho_{p',B}^*(\mathbf{e}) \geq \frac{(\mathbf{e}, \mathbf{f})}{\rho_{p,B}(\mathbf{f})}$ . The proof is completed by taking the supremum over all  $\mathbf{f} \in \mathbf{C}^d$ . □

A metric  $\rho$  is called an  $A_p$  metric if there exists some constant  $C < \infty$  so that the opposite statement

$$(3) \quad \rho_{p',B}^*(\mathbf{e}) \leq C(\rho_{p,B})^*(\mathbf{e}) \quad \text{for all balls } B \subset \mathbf{R}^n$$

is also true. Since the averages over cubes and balls in  $\mathbf{R}^n$  differ by no more than a fixed constant,  $A_p$  metrics satisfy an analogous condition for cubes, and vice versa. Stated either way, the  $A_p$  condition characterizes an important class of weighted measures.

**Theorem 1** (Nazarov, Treil' [12], Volberg [25]). *Let  $d < \infty$ . The following statements are equivalent:*

- 1) *The Hilbert Transform is bounded on  $L^p(\rho)$ .*
- 2)  *$\rho$  is an  $A_p$  metric.*

We will prove this theorem again for metrics which are induced by some matrix weight  $W$ . There is no loss of generality because for fixed dimension  $d < \infty$  every metric can be uniformly approximated by matrix weights.

**Proposition 1.2.** *Let  $d < \infty$ . Given a Banach space norm  $\rho_x$  on  $\mathbf{C}^d$ , there exists a positive selfadjoint matrix  $W_x$  such that*

$$(4) \quad \rho_x(\mathbf{e}) \leq |W_x(\mathbf{e})| \leq \sqrt{d} \cdot \rho_x(\mathbf{e}) \quad \text{for all } \mathbf{e} \in \mathbf{C}^d.$$

*Proof.* Let  $O$  represent the unit ball of  $\rho_x$ , and  $E$  the ellipsoid of maximal volume contained in  $O$ . There exists a positive selfadjoint matrix  $W_x$  such that  $W_x(E)$  is the standard unit ball in  $\mathbf{C}^d$ . The image  $W_x(O)$  is a convex balanced set containing the unit ball, and containing no ellipsoid of greater volume.

If there exists a point  $\mathbf{v} \in W_x(O)$  with  $|\mathbf{v}| > \sqrt{d}$ , then by convexity the boundary of  $W_x(O)$  can only be tangent to the unit sphere at points  $\mathbf{w}$  such that

$$(\mathbf{w}, \mathbf{v}) \leq \frac{1}{|\mathbf{v}|} < \frac{1}{\sqrt{d}}.$$

For some  $\delta > 0$  the ellipsoid with major axis length  $e^\delta$  in the direction of  $\mathbf{v}$  and minor axes length  $e^{-\delta/(|\mathbf{v}|^2-1)}$  in every direction perpendicular to  $\mathbf{v}$  is also contained in  $W_x(O)$ . This has strictly greater volume than the unit ball, contradicting the property of  $W_x(O)$  stated above.  $\square$

It is now possible to state the  $A_p$  condition in terms of matrix weights, though some precision is lost in the process. Given a matrix weight  $W$  and a ball  $B \subset \mathbf{R}^n$ , define a Banach space norm  $X_B$  on  $\mathbf{C}^d$  by considering the  $L^p(W)$  norm of characteristic functions on  $B$ .

$$\|\mathbf{v}\|_{X_B} = |B|^{-1/p} \|\chi_B \mathbf{v}\|_{L^p(W)}.$$

By Proposition 1.2 there exists a positive-definite  $d \times d$  matrix  $V_B$  such that  $\|\mathbf{v}\|_{X_B} \leq |V_B \mathbf{v}| \leq d^{1/2} \|\mathbf{v}\|_{X_B}$ . From a heuristic standpoint,  $V_B$  might

be considered an “ $L^p$  average” of  $W^{1/p}$  over ball  $B$ . With  $p' = \frac{p}{p-1}$  the dual exponent to  $p$ , let  $V'_B$  be an  $L^{p'}$  average of  $W^{-1/p}$ . In summary, matrices  $V_B, V'_B$  enjoy the following properties:

$$(5) \quad \begin{aligned} |V_B \mathbf{v}| &\sim |B|^{-1/p} \|\chi_B W^{1/p} \mathbf{v}\|_{L^p} \\ |V'_B \mathbf{v}| &\sim |B|^{-1/p'} \|\chi_B W^{-1/p} \mathbf{v}\|_{L^{p'}}. \end{aligned}$$

**Remark.** The definition of  $V_B$  and  $V'_B$  depends implicitly on the method used to approximate Banach space norms by matrices. For the purposes of our discussion,  $V_B$  and  $V'_B$  may be any two matrices satisfying (5).

The statement about weights taking the place of Proposition 1.1 is

$$|V_B V'_B \mathbf{e}| \geq |\mathbf{e}| \text{ for all vectors } \mathbf{e} \in \mathbf{C}^d \text{ and balls } B \subset \mathbf{R}^n.$$

A matrix weight  $W$  satisfies the *matrix  $A_p$  condition* if  $V_B V'_B$  are uniformly bounded as operators on  $\mathbf{C}^d$ ; that is

$$(6) \quad \|V_B V'_B\| \leq C < \infty \text{ for all balls } B \subset \mathbf{R}^n.$$

The exact value of  $C$  depends on the choice of  $V_B$  and  $V'_B$ , and is therefore determined here only up to a factor of  $d$ .

Our approach to Theorem 1 is styled after Coifman and Fefferman’s proof [5] in the scalar ( $d = 1$ ) case. Two technical problems arise immediately: First that general  $d \times d$  matrices do not commute with one another, and second the matter of defining a maximal operator for vector-valued functions. To choose pointwise a vector with the largest  $\ell^2(\mathbf{C}^d)$  magnitude is clearly wrong because the effect of weight  $W(x)$  may depend strongly on the direction. In the special case where  $W$  is uniformly nonsingular (i.e.,  $\|W(x)\| \cdot \|W^{-1}(x)\| \leq C$  for all  $x$ ) this can be controlled by a constant factor, but we have no such *a priori* assumptions about  $W$ .

For this reason our analysis will take place primarily in unweighted  $L^q$  spaces, following [4]. Rather than deal with  $T$  directly, we consider the action of  $W^{1/p} T W^{-1/p}$  on functions in  $L^q(dx)$ . With the family of truncated operators  $W^{1/p} T_\epsilon W^{-1/p}$  in mind, we define the maximal truncated operator  $(W^{1/p} T)_*$  to be

$$(7) \quad (W^{1/p} T)_* f(x) = \sup_{\epsilon > 0} |W^{1/p} T_\epsilon f(x)|$$

with the convention that  $f = W^{-1/p} g$  and  $g$  is a function in  $L^q(dx)$ . One estimate from the scalar theory that remains wholly intact is the bound

$$(8) \quad |W^{1/p} T W^{-1/p} g|(x) \leq |(W^{1/p} T)_* W^{-1/p} g|(x) + C|g(x)|.$$

The constant  $C$  depends only on our choice of operator  $T$  but not on the function  $g$ . This will allow us to infer the boundedness of  $T$  by controlling the behavior of its truncations. Our primary results are the following four theorems, numbered according to the section in which they appear:

**Four Theorems.**

(3.2) If  $W$  is a matrix  $A_p$  weight, there exists  $\delta > 0$  such that the vector Hardy-Littlewood maximal function  $M_w$  (defined in Section 3) is a bounded operator from  $L^q(\mathbf{R}^n; \mathbf{C}^d)$  to  $L^q(\mathbf{R}; \mathbf{R})$  whenever  $|p - q| < \delta$ .

(4.2): Given a singular integral operator  $T$  as above, and a weight  $W \in A_p$ , there exists  $\delta > 0$  such that  $(W^{1/p}T)_*W^{-1/p}$  is a bounded operator from  $L^q(\mathbf{R}^n; \mathbf{C}^d)$  to  $L^q(\mathbf{R}; \mathbf{R})$  whenever  $|p - q| < \delta$ .

(5.1): Consequently  $W^{1/p}TW^{-1/p}$  is bounded on  $L^q(\mathbf{R}^n; \mathbf{C}^d)$  for this range of exponents  $q$ .

(5.2): In particular,  $T$  is bounded on  $L^p(W)$  if  $W \in A_p$ . With one additional hypothesis on the structure of  $T$ , the converse statement is also true.

**Remark.** The exponent  $W^{1/p}$  is used throughout, even when we are considering functions under an  $L^q$  norm with  $q \neq p$ . This places us squarely in the setting of [25], where the  $A_p$  metric  $W^{1/p}$  is the basic object of study. Theorem 5.1 then asserts that any  $A_p$  metric is also an  $A_q$  metric for all  $q$  in some open interval containing  $p$ .

**2. Properties of  $A_p$  weights.**

We would like first to characterize the matrix  $A_p$  class in a more transparent manner by borrowing a lemma from [12]:

**Proposition 2.1.** *A metric  $\rho_x$  satisfies the  $A_p$  condition if and only if the operators  $f \rightarrow \chi_B \frac{1}{|B|} \int_B f dx$  are uniformly bounded on  $L^p(\rho)$ . In fact, the uniform bound is equal to the  $A_p$  constant of  $\rho$ .*

*Proof.* The  $L^p(\rho)$  norm of  $\chi_B \frac{1}{|B|} \int_B f dx$  is given by

$$\frac{1}{|B|} \left( \int_B \left[ \rho_y \left( \int_B f dx \right) \right]^p dy \right)^{1/p},$$

which in turn is equal to  $|B|^{-1/p'} \rho_{p,B} \left( \int_B f dx \right)$ . Therefore

$$\begin{aligned} \sup_{\|f\|_{L^p(\rho)}=1} \left\| \chi_B \frac{1}{|B|} \int_B f dx \right\|_{L^p(\rho)} &= \sup_f \sup_{\mathbf{e} \in \mathbf{C}^d} |B|^{-1/p'} \frac{\int_B(\mathbf{e}, f(x)) dx}{(\rho_{p,B})^*(\mathbf{e})} \\ &= \sup_{\mathbf{e} \in \mathbf{C}^d} |B|^{-1/p'} \frac{\|\chi_B \mathbf{e}\|_{L^{p'}(\rho^*)}}{(\rho_{p,B})^*(\mathbf{e})} \\ &= \sup_{\mathbf{e} \in \mathbf{C}^d} \frac{\rho_{p',B}^*(\mathbf{e})}{(\rho_{p,B})^*(\mathbf{e})}. \end{aligned}$$

Equality between the first and second lines takes place because  $L^p(\rho)$  is the dual space of  $L^{p'}(\rho^*)$ . □

**Corollary 2.2.** *Let  $\rho$  be an  $A_p$  metric. For any vector  $\mathbf{v} \in \mathbf{C}^d$ ,  $\rho_x(\mathbf{v})^p$  is a scalar  $A_p$  weight with constant less than or equal to that of  $\rho$ .*

*Proof.* Let  $\phi$  be any scalar function and consider  $f = \phi\mathbf{v}$ . The weighted norm of  $f$  is  $\|f\|_{L^p(\rho)} = (\int_B \phi^p [\rho_x(\mathbf{v})]^p dx)^{1/p}$ . Proposition 2.1 applied to  $f$  states that all maps  $\phi \rightarrow \chi_B \frac{1}{|B|} \int_B \phi dx$  are uniformly bounded on the  $L^p$  space with measure  $[\rho_x(\mathbf{v})]^p dx$ , with norms less than the  $A_p$  constant of  $\rho$ . We now apply Proposition 2.1 again, this time in the scalar setting, to conclude that  $[\rho_x(\mathbf{v})]^p$  is a scalar  $A_p$  weight whose constant is also less than the  $A_p$  constant of  $\rho$ .  $\square$

**Corollary 2.3.** *If  $W$  is a matrix  $A_p$  weight, then  $\|W\|$  is a scalar  $A_p$  weight.*

*Proof.* Let  $\mathbf{e}_i$  be the standard unit basis for  $\mathbf{C}^d$ . Since  $W(x)$  is a nonnegative and selfadjoint operator at each point  $x$ ,

$$\begin{aligned}
 (9) \quad \|W(x)\| &= \|W^{2/p}(x)\|^{p/2} \sim [\text{tr}(W^{2/p}(x))]^{p/2} \\
 &= \left( \sum_{i=1}^d |W^{1/p}(x)\mathbf{e}_i|^2 \right)^{p/2} \\
 &\sim \sum_{i=1}^d |W^{1/p}(x)\mathbf{e}_i|^p
 \end{aligned}$$

pointwise in  $x$ . By Corollary 2.2, each individual function  $|W^{1/p}(x)\mathbf{e}_i|^p$  is a scalar  $A_p$  weight, therefore their sum is as well.  $\square$

**Remarks.** Both of these corollaries are proven in [23] for the case  $p = 2$ , and are adapted here with minimal alteration.

From this point forward we will work exclusively in the language of matrix weights. While our primary definition of  $A_p$  weights (6) is decidedly less elegant than that of  $A_p$  metrics (3), the ability to use notation and theorems from linear algebra makes it a worthwhile sacrifice.

One crucial feature in the theory of scalar  $A_p$  weights is the presence of ‘‘Reverse Hölder’’ inequalities estimating the average of  $W^{1+\epsilon}$  in terms of the average of  $W$ . We will employ inequalities of a similar character as the centerpiece of our analysis.

**Proposition 2.4.** *Let  $W$  be an  $A_p$  weight. Then there exist  $\delta > 0$  and constants  $C_q$  such that for all balls  $B \subset \mathbf{R}^n$ ,*

$$(10) \quad \frac{1}{|B|} \int_B \|W^{1/p}(y)V'_B\|^q dy \leq C_q, \text{ all } q < p + \delta$$

$$(11) \quad \frac{1}{|B|} \int_B \|V_B W^{-1/p}(y)\|^q dy \leq C_q, \text{ all } q < p' + \delta.$$

*Proof.* We will verify only the first of these statements. The second one is proven in an identical manner with the starting point that  $W^{-p'/p}$  is an  $A_p$  weight.

By Corollary 2.2, all functions of the form  $|W^{1/p}(y)V'_B\mathbf{e}|^p$  are scalar  $A_p$  weights with  $A_p$  norms bounded uniformly in  $\mathbf{e}$ . It is therefore possible to choose  $q$  and  $C_q$  so that the Reverse Hölder inequality

$$\frac{1}{|B|} \int_B |W^{1/p}(y)V'_B\mathbf{e}|^q dy \leq C_q \left( \frac{1}{|B|} \int_B |W^{1/p}(y)V'_B\mathbf{e}|^p dy \right)^{q/p}$$

is satisfied for all  $\mathbf{e} \in \mathbf{C}^d$ .

Let  $\mathbf{e}_i$  once again be the standard unit basis for  $\mathbf{C}^d$ . It is useful to remember that the norm of any  $d \times d$  matrix  $M$  (not necessarily Hermitian) is controlled by its action on the vectors  $\mathbf{e}_i$  via the formula

$$\|M\| \leq d^{1/2} \sup_i |M\mathbf{e}_i|.$$

We may now estimate the desired integral:

$$\begin{aligned} \frac{1}{|B|} \int_B \|W^{1/p}(y)V'_B\|^q dy &\leq \frac{1}{|B|} \int_B (d^{1/2} \sup_i |W^{1/p}(y)V'_B\mathbf{e}_i|)^q dy \\ &\leq d^{q/2} \sum_{i=1}^d \frac{1}{|B|} \int_B |W^{1/p}(y)V'_B\mathbf{e}_i|^q dy \\ &\leq C_q \sum_{i=1}^d \left( \frac{1}{|B|} \int_B |W^{1/p}(y)V'_B\mathbf{e}_i|^p dy \right)^{q/p} \\ &\sim C_q \sum_{i=1}^d |V_B V'_B \mathbf{e}_i|^q \leq d \cdot C_q \|V_B V'_B\|^q \leq C_q. \end{aligned}$$

□

**Note.** In later sections we will also use the slightly weaker inequality

$$(12) \quad |B|^{-1} \int_B \|W^{1/p}(y)V_B^{-1}\|^q dy \leq C_q, \text{ all } q < p + \delta$$

whose proof follows the above calculations almost word for word.

### 3. The Hardy-Littlewood maximal function.

There is a wide variety of possible maximal functions to choose from, each of which has its own advantages and limitations. In [4] we first considered an auxiliary maximal function  $M'_w$ , given by

$$(13) \quad M'_w g(x) = \sup_{x \in B} \frac{1}{|B|} \int_B |V_B W^{-1/p}(y)g(y)| dy.$$

Although the intuitive meaning of  $M'_w$  is unclear, one may approach it with the classical tools of weak-type inequalities and interpolation. A direct application of the second reverse Hölder inequality (11) proves the following lemma:

**Lemma 3.1.** *Let  $W$  be an  $A_p$  weight. Then there exists  $\delta > 0$  such that*

$$\|M'_w g\|_{L^q} \leq C_q \|g\|_{L^q(\mathbf{R}^n; \mathbf{C}^d)}, \text{ all } g \in L^q, \text{ all } q > p - \delta.$$

*Sketch of Proof.* The operators  $g \mapsto \chi_B \frac{1}{|B|} \int_B |V_B W^{-1/p} g| dy$  are uniformly bounded in  $L^q$  if  $p - \delta < q$ . This is a consequence of (11) together with the inequality

$$\int_B |V_B W^{-1/p}(y)g(y)| dy \leq \left( \int_B \|V_B W^{-1/p}(y)\|^{q'} dy \right)^{1/q'} \|g\|_{L^q}$$

One may use the Vitali Covering Lemma to obtain a weak-type  $(q, q)$  estimate on the associated maximal function  $M'_w$ . The lemma then follows from the Marcinkiewicz Interpolation Theorem.  $\square$

The vector Hardy-Littlewood maximal function  $M_w$  is defined as

$$(14) \quad M_w g(x) = \sup_{x \in B} \frac{1}{|B|} \int_B |W^{1/p}(x)W^{-1/p}(y)g(y)| dy.$$

The following equivalent definition of  $M_w$  is often quite useful:

$$(15) \quad M_w g(x) = M(|W^{1/p}(x)W^{-1/p}(\cdot)g(\cdot)|)(x).$$

Here  $M$  denotes the classical Hardy-Littlewood maximal operator acting on scalar-valued functions. The only difference between  $M_w$  and  $M'_w$  is the presence of a weight  $W^{1/p}(x)$  rather than an average weight  $V_B$  over a ball containing  $x$ . The reverse Hölder inequalities suggest that  $A_p$  weights are often pointwise comparable to their averages, in which case  $\|M_w g\|$  would be controlled by  $\|M'_w g\|$ . For a range of exponents near  $p$ , this line of reasoning can be made precise.

**Theorem 3.2.** *Let  $W$  be an  $A_p$  weight. Then there exists  $\delta > 0$  such that*

$$\|M_w g\|_{L^q} \leq C_q \|g\|_{L^q(\mathbf{R}^n; \mathbf{C}^d)}, \text{ all } g \in L^q, \text{ all } |p - q| < \delta.$$

*Proof.* Let us suppose for a moment that the suprema defining  $M_w g$  and  $M'_w g$  are taken over cubes in some dyadic grid. The entire preceding discussion holds for maximal functions over cubes, so in particular we can still estimate  $\|M'_w g\|$  via Lemma 3.1. For each point  $x$ , choose a (dyadic) cube  $R_x$  such that

$$\begin{aligned} M_w g(x) &\leq 2|R_x|^{-1} \int_{R_x} |W^{1/p}(x)W^{-1/p}(y)g(y)| dy \\ &\leq 2\|W^{1/p}(x)V_{R_x}^{-1}\| \cdot \left( |R_x|^{-1} \int_{R_x} |V_{R_x} W^{-1/p}(y)g(y)| dy \right). \end{aligned}$$

For each integer  $j$ , define  $\{S_j\}$  to be the collection of dyadic cubes  $R = R_x$  that are maximal with respect to the property

$$2^j \leq |R|^{-1} \int_R |V_R W^{-1/p}(y)g(y)|dy < 2^{j+1}.$$

Maximality insures that whenever  $M_w g(x) \neq 0$  the cube  $R_x$  is contained in some  $S_j$  with

$$|R_x|^{-1} \int_{R_x} |V_{R_x} W^{-1/p}(y)g(y)|dy \leq 2|S_j|^{-1} \int_{S_j} |V_{S_j} W^{-1/p}(y)g(y)|dy.$$

When  $j$  is fixed, the disjoint union  $\cup_j S_j$  is contained in the set where  $M'_w g(x) \geq 2^j$ .

Consider the functions  $N_Q(x) = \sup_{x \in RCQ} \|W^{1/p}(x)V_R^{-1}\|$ , defined on their respective cubes  $Q$ . By virtue of the preceding two statements, the inequality  $M_w g(x) \leq 4 \cdot 2^{j+1} N_{S_j}(x)$  must hold for some number  $j$  (this is trivial at the points where  $M_w g(x) = 0$ ). It follows that

$$(16) \quad \|M_w g\|_{L^q}^q \leq C \sum_{j=-\infty}^{\infty} 2^{jq} \sum_{S_j} \int_{S_j} (N_{S_j}(x))^q dx.$$

By Lemma 3.3 below, we can continue the estimate as follows:

$$\begin{aligned} \|M_w g\|_{L^q}^q &\leq C \sum_{j=-\infty}^{\infty} 2^{jq} \sum_{S_j} |S_j| \\ &\leq C \sum_{j=-\infty}^{\infty} 2^{jq} |\{M'_w g \geq 2^j\}| \leq C \|M'_w g\|_{L^q}^q. \end{aligned}$$

The proof is then complete by Lemma 3.1. □

**Lemma 3.3.** *Let  $W$  be a matrix  $A_p$  weight and functions  $N_Q(x)$  be defined as above. Then there exist  $\delta > 0$  and  $C_q < \infty$  such that for all dyadic  $Q$ ,*

$$\int_Q (N_Q(x))^q dx \leq C_q |Q| \text{ for all } q < p + \delta.$$

*Proof.* We present an informal argument here, assuming that  $\int_Q N_Q^q \leq B|Q|$  with some finite  $B$  then deriving an *a priori* bound for  $B$ . This may be readily adapted into a rigorous proof.

Let  $A < \infty$  be a large constant to be specified later. Denote by  $\{R_j\}$  the set of maximal cubes satisfying  $\|V_Q V_{R_j}^{-1}\| > A$ . Outside of  $\cup_j R_j$ ,  $N_Q(x) \leq A \|W^{1/p}(x)V_Q^{-1}\|$ . Thus  $\int_{Q \setminus \cup_j R_j} (N_Q(x))^q dx \leq C|Q|$ , seen by applying reverse Hölder inequality (12).

We claim that  $\sum_j |R_j| < \frac{1}{2}|Q|$  if  $A$  is sufficiently large. Remember first that  $\|V_Q V_{R_j}^{-1}\| = \|V_{R_j}^{-1} V_Q\| \leq C \|V_{R_j}' V_Q\|$ , by Proposition 1.1. It follows that

$$\begin{aligned}
 (17) \quad |R_j| \cdot \|V_Q V_{R_j}^{-1}\|^{p'} &\leq C \sup_{|\mathbf{e}|=1} |R_j| \cdot |V_{R_j}' V_Q \mathbf{e}|^{p'} \\
 &\sim C \sup_{|\mathbf{e}|=1} \int_{R_j} |W^{-1/p}(y) V_Q \mathbf{e}|^{p'} dy \\
 &\leq C \int_{R_j} \|W^{-1/p}(y) V_Q\|^{p'} dy.
 \end{aligned}$$

The cubes  $R_j$  are disjoint from one another, so

$$\begin{aligned}
 A^{p'} \sum_j |R_j| &< C \int_{\cup_j R_j} \|W^{-1/p}(y) V_Q\|^{p'} dy \\
 &\leq C \int_Q \|W^{-1/p}(y) V_Q\|^{p'} \leq C|Q|.
 \end{aligned}$$

This estimate shows that for  $A$  large enough,  $\sum_j |R_j| < \frac{1}{2}|Q|$ , and the value of  $A$  may be chosen independently of  $Q$ .

Inside each cube  $R_j$ , we may assume that  $N_Q(x) = N_{R_j}(x)$ , otherwise the bound  $N_Q(x) \leq A \|W^{1/p}(x) V_Q^{-1}\|$  still holds. Then

$$(18) \quad \int_{\cup_j R_j} (N_Q(x))^q = \sum_j \int_{R_j} (N_{R_j}(x))^q \leq B \sum_j |R_j| < \frac{1}{2} B|Q|.$$

Putting these pieces together, we would discover that  $B \leq C + \frac{1}{2}B$ , where  $C < \infty$  is determined by the constants in the reverse Hölder inequality.  $\square$

This concludes the proof that matrix  $A_p$  weights enjoy  $L^q$ -boundedness of the dyadic Hardy-Littlewood maximal function for a range of exponents  $|q - p| < \delta$ . There is a standard argument employing two incompatible dyadic grids [7] for extending results of this kind to the general setting. Thus the Hardy-Littlewood maximal function as we originally defined it (as a supremum over balls containing  $x$ ) is bounded in  $L^q$  for the same range of exponents  $q$ .

#### 4. A distributional inequality.

**Proposition 4.1.** *Let  $W$  be a matrix  $A_p$  weight and fix  $q < p + \delta$ . Then there exist positive constants  $0 < b < 1$ ,  $c > 0$  depending only on  $q$ , the  $A_p$*

“norm” of  $W$ , and the dimensions  $d, n$  such that

$$(19) \quad \left| \left\{ x \in \mathbf{R}^n : (W^{1/p}T)_*f(x) > \alpha; \right. \right. \\ \left. \left. \max(M'_w(W^{1/p}f)(x), M_w(W^{1/p}f)(x)) < c\alpha \right\} \right| \\ < \frac{1}{2}b^q |\{x \in \mathbf{R}^n : (W^{1/p}T)_*f(x) > b\alpha\}|$$

for all  $f \in C_c^\infty(\mathbf{R}^n; \mathbf{C}^d)$ .

From this point onward we follow as closely as possible in the footsteps of Coifman and Fefferman [5], decomposing the set where  $(W^{1/p}T)_*f > b\alpha$  into a union of cubes and proving the desired inequality on each cube separately. Our decomposition uses a slightly modified version of the Whitney covering lemma, stated below.

**Covering Lemma.** *Given a set  $E \subset \mathbf{R}^n$  of finite (Lebesgue) measure, there exists a collection  $\{Q_j\}$  of pairwise disjoint cubes such that:*

- i)  $E \subset \cup_j Q_j$  up to sets of measure zero,
- ii)  $|Q_j \cap E| \geq \frac{1}{2}|Q_j|$ ,
- iii)  $|3Q_j \cap E^c| \geq C_n|3Q_j|$ .

A simple consequence of Statements i) and ii) is that  $\sum_j |Q_j| \leq 2|E|$ .

*Proof.* Let  $\{Q_j\}$  be the collection of dyadic cubes maximal under the property that  $|Q \cap E| \geq \frac{1}{2}|Q|$ . Then Conditions ii) and iii) hold with constant  $C_n = \frac{1}{2} \cdot (\frac{2}{3})^n$ . The first condition also holds because as  $\epsilon \rightarrow 0$ , the ratio  $|B(x, \epsilon) \cap E|/|B(x, \epsilon)| \rightarrow 1$  at almost every  $x \in E$ . □

*Proof of Proposition 4.1.* Write  $f = W^{-1/p}g$  and let

$$E = \{x \in \mathbf{R}^n : (W^{1/p}T)_*f(x) > b\alpha\}.$$

Apply the covering lemma to obtain cubes  $\{Q_j\}$  with the specified properties. It suffices to verify that in each cube  $Q = Q_j$  there is a distributional inequality

$$(20) \quad \left| \left\{ x \in Q : (W^{1/p}T)_*f(x) > \alpha; \max(M'_w g(x), M_w g(x)) < c\alpha \right\} \right| < \frac{1}{4}b^q|Q|.$$

For this we use a construction similar to the one in [5]. Let  $O$  be the ball with the same center as  $Q_j$  and radius  $5 \text{ diam}(Q)$ . By the covering lemma and inequality (11), there exists a point  $\bar{x} \in 3Q$  such that

$$(W^{1/p}T)_*f(\bar{x}) < b\alpha \text{ and } \|V_O W^{-1/p}(\bar{x})\| < C.$$

Let  $B = B(\bar{x}, 3 \text{ diam}(Q_j))$ . Since  $B \subset O$  and is of comparable size,  $\|V_B V_O^{-1}\|$  is bounded by a constant and hence  $\|V_B W^{-1/p}(\bar{x})\| < C$ .

Assume  $|\{x \in Q : M'_w g(x) < c\alpha\}| \geq \frac{1}{4}b^q|Q|$ , otherwise the proposition is trivially satisfied. Then there exists a point  $\bar{y} \in Q$  such that

$$M'_w g(\bar{y}) < c\alpha \text{ and } \|V_B W^{-1/p}(\bar{y})\| \leq Cb^{-1}.$$

Write  $f_1 = \chi_B f$  and  $f_2 = \chi_{B^c} f$ . By the sublinearity of  $(W^{1/p}T)_*$ , the set where  $(W^{1/p}T)_* f(x) > \alpha$  is contained in the union of sets  $\{(W^{1/p}T)_* f_i(x) > \alpha/2\}$ ,  $i = 1, 2$ .

The operator  $T_*$  is weak-type  $(1, 1)$ . This fact is easily obtained from the scalar case when  $d$  is finite, but is also true in general [17]. Consequently,

$$\left| \left\{ (V_B T)_* f_1(x) > \frac{\alpha}{2R} \right\} \right| \leq \frac{AR}{\alpha} \|V_B f_1\|_{L_1(\mathbf{R}^n; \mathbf{C}^d)}.$$

Here we are using the property that operator  $T_*$  commutes with multiplication by any constant matrix, in this case  $V_B$ . Furthermore,

$$\|V_B f_1\|_{L^1} = \int_B |V_B f(y)| dy \leq |B| M'_w g(\bar{y}) \leq Cc\alpha|Q|$$

with the end result that  $|\{x \in Q : (V_B T)_* f_1(x) > \frac{\alpha}{2R}\}| \leq CcR|Q|$ .

It follows that  $|\{x \in Q : (W^{1/p}T)_* f_1(x) > \frac{\alpha}{2}\}| \leq (CcR + C'R^{-p})|Q|$  for all  $R > 0$ , because the Reverse Hölder inequality (10) guarantees that  $\|W^{1/p}(x)V_B^{-1}\| < R$  except on a set of measure less than  $C'R^{-p}$ . Taking the infimum over  $R$ ,

$$(21) \quad |\{x \in Q : (W^{1/p}T)_* f_1(x) > \alpha/2\}| \leq C_0 c^{p/(p+1)}|Q|.$$

For the second estimate, we begin by noting that the point  $\bar{x}$  is chosen so that  $(W^{1/p}T)_* f(\bar{x}) < b\alpha$  and  $\|V_B W^{-1/p}(\bar{x})\| < C$ . Then  $(V_B T)_* f(\bar{x}) < Cb\alpha$ . Our estimate for  $|\{(W^{1/p}T)_* f_2(x) > \alpha/2\}|$  relies on the following inequality which holds for all  $x \in Q$ :

$$\begin{aligned} (22) \quad (V_B T)_* f_2(x) &\leq (V_B T)_* f(\bar{x}) + C' M(|V_B f|)(\bar{y}) \\ &\leq Cb\alpha + C' \|V_B W^{-1/p}(\bar{y})\| \cdot M(|W^{1/p} f|)(\bar{y}) \\ &\leq Cb\alpha + C' \|V_B W^{-1/p}(\bar{y})\| \cdot M_w g(\bar{y}) \\ &\leq (Cb + C'b^{-1}c)\alpha. \end{aligned}$$

In the preceding expressions  $M(\cdot)$  denotes the scalar Hardy-Littlewood maximal function.

Imitating the method for the  $|(W^{1/p}T)_* f_1|$  estimate, we see that

$$|\{x \in Q : (W^{1/p}T)_* f_2(x) > R(Cb + C'b^{-1}c)\alpha\}| \leq AR^{-r}|Q|$$

where  $r$  may be chosen so that  $q < r < p + \delta$ . Once again (10) has been invoked, this time to guarantee that  $\|W^{1/p}V_B^{-1}\| > R$  only on a set of measure less than  $CR^{-r}|B|$ . Set  $R$  equal to  $(4bC)^{-1}$ . Then

$$(23) \quad |\{x \in Q : (W^{1/p}T)_* f_2(x) > (1/4 + C_1 b^{-2}c)\alpha\}| \leq C_2 b^r |Q|.$$

Statement (20) is then verified by choosing  $b < (8C_2)^{1/q-r}$  and  $c$  sufficiently small. Summing over all cubes  $Q_j$  proves the proposition.  $\square$

**Corollary 4.2.** *With  $c$  as in Proposition (4.1),*

$$\|(W^{1/p}T)_*f\|_{L^q}^q \leq 2c^{-q} \|\max(M'_w(W^{1/p}f), M_w(W^{1/p}f))\|_{L^q}^q$$

for all  $f \in C_c^\infty(\mathbf{R}^n; \mathbf{C}^d)$ .

*Proof.* If both sides of (19) are multiplied by  $q\alpha^{q-1}$  and integrated over the interval  $0 \leq \alpha < \infty$ , the resulting inequality is

$$\begin{aligned} & \int_{\mathbf{R}^n} \left( [(W^{1/p}T)_*f]^q - c^{-q} \max([M'_w(W^{1/p}f)]^q, [M_w(W^{1/p}f)]^q) \right)_+ dx \\ & \leq \frac{1}{2} \int_{\mathbf{R}^n} [(W^{1/p}T)_*f]^q dx \end{aligned}$$

from which it follows that

$$\begin{aligned} & \|(W^{1/p}T)_*f\|_{L^q}^q - \frac{1}{c^q} \|\max(M'_w(W^{1/p}f), M_w(W^{1/p}f))\|_{L^q}^q \\ & \leq \frac{1}{2} \|(W^{1/p}T)_*f\|_{L^q}^q. \end{aligned}$$

The remaining task is to verify that the  $L^q$  norm of  $(W^{1/p}T)_*f$  is finite. A key estimate is the fact that  $T_*f(x) \leq C_f(1+|x|)^{-n}$  for all  $f \in C_c^\infty$ , where  $C_f$  depends on  $f$ . Then

$$(W^{1/p}T)_*f(x) \leq C\|W\|^{1/p}(1+|x|)^{-n}.$$

There are many ways to show that the expression on the right-hand side is in  $L^q$ , all exploiting the fact that  $\|W\|$  is a scalar  $A_p$  weight. One possibility is to choose any nontrivial (scalar) function  $\phi \geq 0 \in C_c^\infty$ . We have shown in Theorem 3.2 that  $\|W\|^{1/p}M(\|W\|^{-1/p}\phi) \in L^q$  whenever  $|p-q| < \delta$ .

On the other hand,  $C(1+|x|)^{-n} \leq M(\|W\|^{-1/p}\phi)$ , which completes the proof.  $\square$

### 5. The main theorem.

**Theorem 5.1.** *Let  $T$  be a linear operator whose associated convolution kernel  $K(x)$  satisfies the hypotheses in (1), and which acts separately on each coordinate function of  $f$  (in other words,  $(Tf)_j = Tf_j$ ). Let  $W$  be a matrix  $A_p$  weight.*

*There exists  $\delta > 0$  such that  $W^{1/p}TW^{-1/p}$  is a bounded operator on  $L^q(\mathbf{R}^n; \mathbf{C}^d)$  whenever  $|q-p| < \delta$ .*

*Proof.* As in the scalar case, the truncated operators  $T_\epsilon$  possess a weak limit  $T_0$ , and  $T = T_0 + A$ , where  $A$  is a bounded pointwise multiplier. In dimensions  $d > 1$ ,  $A = A(x)$  is a matrix-valued function, but the hypothesis

$(Tf)_j = Tf_j$  requires  $A(x)$  to be a scalar  $L^\infty$  function multiplied by the identity matrix.

The function  $W^{1/p}TW^{-1/p}g$  is dominated pointwise by  $g$  and  $(W^{1/p}T)_*(W^{-1/p}g)$ , as in Equation (8):

$$\begin{aligned} |W^{1/p}TW^{-1/p}g(x)| &= |W^{1/p}T_0W^{-1/p}g(x) + A(x)g(x)| \\ &\leq |(W^{1/p}T)_*(W^{-1/p}g)(x)| + C|g(x)|. \end{aligned}$$

The triangle inequality for  $L^q$ -norms immediately yields the result

$$(24) \quad \|W^{1/p}TW^{-1/p}g\|_{L^q} \leq \|(W^{1/p}T)_*W^{-1/p}g\|_{L^q} + C\|g\|_{L^q}.$$

For all  $g$  such that  $W^{-1/p}g \in C_c^\infty$ , the right-hand side is controlled by  $\|g\|_{L^q}$ . Observe that  $W^{q/p}$  is a locally integrable matrix-valued function. Then  $C_c^\infty(\mathbf{R}^n; \mathbf{C}^d)$  is a dense subset of  $L^q(W^{q/p})$ . The map  $f \in L^q(W^{q/p}) \rightarrow g = W^{1/p}f \in L^q(dx)$  is an invertible isometry, so its image  $W^{1/p}(C_c^\infty)$  is dense in  $L^q$ . Thus the boundedness of  $W^{1/p}TW^{-1/p}$  may then be extended to all functions  $g \in L^q(\mathbf{R}^n; \mathbf{C}^d)$ ,  $|p - q| < \delta$ .  $\square$

A converse statement, with some minor modifications, is also true.

**Theorem 5.2.** *Suppose that  $T$  is a convolution operator as above, with the additional nondegeneracy hypothesis that there exists some unit vector  $\mathbf{u} \in \mathbf{R}^n$  such that  $|K(r\mathbf{u})| \geq a|r|^{-n}$ , all  $r \in \mathbf{R} \setminus \{0\}$ . If  $T$  is a bounded operator on  $L^p(W)$ , then  $W$  is an  $A_p$  weight.*

In order to prove this theorem we first need a result about integral operators with bounded and compactly supported kernels:

**Proposition 5.3.** *Let  $S$  be an integral operator  $Sf(x) = \int_{\mathbf{R}^n} S(x, y)f(y)$  whose (scalar) kernel  $S(x, y)$  is supported in  $B \times B$  and satisfies the bound  $|S(x, y)| \leq |B|^{-1}$  for all  $(x, y) \in B \times B$ .*

*The norm of  $S$  as an operator on  $L^p(W)$  is less than  $C_d\|V_B V'_B\|$ , where  $C_d$  is a dimensional constant independent of the particular choice of  $S$ . In the special case  $S_0(x, y) = |B|^{-1}\chi_{B \times B}$ , the operator norm of  $S_0$  is also greater than  $C_d^{-1}\|V_B V'_B\|$ .*

*Proof.* This is a straightforward calculation similar to those found in Section 2. Let  $f$  be any function in  $L^p(W)$ . We first estimate the size of

$W^{1/p}(x)Sf(x)$  pointwise for each  $x$ .

$$\begin{aligned}
|W^{1/p}(x)Sf(x)| &= \left| W^{1/p}(x) \int_B S(x, y)f(y) dy \right| \\
&= \left| \int_B S(x, y)W^{1/p}(x)f(y) dy \right| \\
&\leq |B|^{-1} \int_B |W^{1/p}(x)f(y)| dy \\
&\leq |B|^{-1} \left( \int_B \|W^{1/p}(x)W^{-1/p}(y)\|^{p'} dy \right)^{1/p'} \cdot \|f\|_{L^p(W)}.
\end{aligned}$$

As in Section 2, we now introduce an orthonormal basis of vectors  $\mathbf{e}_i$  spanning  $\mathbf{C}^d$ .

$$\begin{aligned}
&\left( \int_B \|W^{1/p}(x)W^{-1/p}(y)\|^{p'} dy \right)^{1/p'} \\
&\leq \left( \int_B \left( d^{1/2} \sup_i |W^{-1/p}(y)W^{1/p}(x)\mathbf{e}_i| \right)^{p'} dy \right)^{1/p'} \\
&\leq d^{1/2} \left( \sum_{i=1}^d \int_B |W^{-1/p}(y)W^{1/p}(x)\mathbf{e}_i|^{p'} dy \right)^{1/p'} \\
&\leq C_d \left( \sum_{i=1}^d |B| \cdot |V'_B W^{1/p}(x)\mathbf{e}_i|^{p'} \right)^{1/p'} \\
&\leq C_d |B|^{1/p'} \|V'_B W^{1/p}(x)\|
\end{aligned}$$

which leads to the estimate

$$|W^{1/p}(x)Sf(x)| \leq C_d |B|^{-1/p} \|V'_B W^{1/p}(x)\| \cdot \|f\|_{L^p(W)}.$$

Then for all  $\|f\|_{L^p(W)} \leq 1$ , it follows that

$$\begin{aligned}
(25) \quad \|Sf\|_{L^p(W)} &\leq C \left( |B|^{-1} \int_B \|V'_B W^{1/p}(x)\|^p dx \right)^{1/p} \\
&\leq C_d \left( |B|^{-1} \int_B \left( d^{1/2} \sup_i |W^{1/p}(x)V'_B \mathbf{e}_i| \right)^p dx \right)^{1/p} \\
&\leq C_d \left( \sum_i |B|^{-1} \int_B |W^{1/p}(x)V'_B \mathbf{e}_i|^p dx \right)^{1/p} \\
&\sim C_d \left( \sum_i |V_B V'_B \mathbf{e}_i|^p \right)^{1/p} \leq C_d \|V_B V'_B\|.
\end{aligned}$$

The second assertion is a restatement of Proposition 2.1. □

*Proof of Theorem 5.2.* First, let  $\epsilon > 0$  be small enough so that  $2\epsilon + \epsilon^2 < \frac{1}{2}C_d^{-2}$ . There exists a number  $t_0 < \infty$  such that

$$(26) \quad |K(\mathbf{v}) - K(rt_0\mathbf{u})| \leq \epsilon|K(rt_0\mathbf{u})| \text{ whenever } \mathbf{v} \in B(rt_0\mathbf{u}, 2r), \text{ all } r \in \mathbf{R} \setminus \{0\}.$$

This is seen to be true because  $|K(rt_0\mathbf{u})| \geq \frac{a}{t_0^n|r|^n}$  but  $|\nabla K(x)| \leq \frac{C}{t_0^{n+1}r^{n+1}}$  for all  $x \in B(rt_0\mathbf{u}, r)$ . It suffices to choose  $t_0 > \frac{2C}{\epsilon a}$ .

Let  $B$  denote the ball  $B(y, r)$  in  $\mathbf{R}^n$ , and  $B'$  the translated ball  $B' = B(y + rt_0\mathbf{u}, r)$ . We wish to consider the operator  $S_B$  defined by

$$S_B f = \chi_B T(\chi_{B'} T(\chi_B f)).$$

This is an integral operator whose kernel

$$S_B(x, y) = \chi_{B \times B} \int_{B'} K(x - z)K(z - y) dz$$

is supported in  $B \times B$ . If  $T$  acts boundedly on  $L^p(W)$ , so too does  $S_B$  with operator norm less than or equal to  $\|T\|^2$ .

The restrictions  $\{x, y \in B, z \in B'\}$  guarantee that  $z - y \in B(rt_0\mathbf{u}, 2r)$  and  $x - z \in B(-rt_0\mathbf{u}, 2r)$ . Thus the values of  $K(z - y)$  and  $K(x - z)$  do not vary much over the region of integration. Using the bounds established in (26), we rewrite  $S_B(x, y)$  as the sum of a characteristic function and a small remainder:

$$(27) \quad S_B(x, y) = |B|K(rt_0\mathbf{u})K(-rt_0\mathbf{u})\chi_{B \times B} + S_1(x, y),$$

where  $|S_1(x, y)| \leq \frac{1}{2}C_d^{-2}|B| \cdot |K(rt_0\mathbf{u})K(-rt_0\mathbf{u})|$ .

According to Proposition 5.3, the first term corresponds to an operator with norm at least  $C\|V_B V'_B\|$ . In terms of other constants,  $C$  is proportional to  $a^2 t_0^{-2n} C_d^{-1}$ . The operator corresponding to the second term has norm no more than half as great. It follows that  $\|S_B\| \geq \frac{1}{2}C\|V_B V'_B\|$ . Then

$$(28) \quad \|V_B V'_B\| \leq 2C^{-1}\|S_B\| \leq 2C^{-1}\|T\|^2 < \infty$$

for all balls  $B \subset \mathbf{R}^n$ , and  $W$  is an  $A_p$  weight. □

**Corollary 5.4.** *If  $W$  is a matrix  $A_p$  weight, there exists  $\delta > 0$  such that  $W^{q/p}$  is an  $A_q$  weight whenever  $|q - p| < \delta$ . In other words, an  $A_p$  metric is also an  $A_q$  metric for all  $|q - p| < \delta$ .*

**Remarks.** We could have proven this statement directly in Section 2, using the reverse Hölder inequality to show that operators  $f \rightarrow \chi_B \frac{1}{|B|} \int_B f dx$  are uniformly bounded on  $L^q(W^{q/p})$ . To do so would have added another computation without simplifying the subsequent discussion in any way.

Recall that a matrix weight  $W \in A_p$  if and only if the averaging operators  $A_B$  defined by

$$A_B f = \chi_B \frac{1}{|B|} \int_B f \, dx$$

are uniformly bounded on  $L^p(W)$ . An equivalent statement is that the conjugated operators  $W^{1/p} A_B W^{-1/p}$  are uniformly bounded on the unweighted space  $L^p(\mathbf{C}^d)$ . It is trivial to observe that  $A_B$  are uniformly bounded on  $L^\infty(\mathbf{C}^d)$  with norm 1. By interpolation on the analytic family of operators<sup>1</sup>

$$\{W^{(1-z)/p} A_B W^{(z-1)/p}, \quad 0 \leq \operatorname{Re}(z) \leq 1\}$$

we find that  $W^{1/r} A_B W^{-1/r}$  are uniformly bounded on  $L^r(dx)$  for all  $r > p$ , leading to another result well-known in the scalar case:

**Proposition 5.5.** *If  $W$  is a matrix  $A_p$  weight, then  $W$  is also a matrix  $A_r$  weight for all  $r > p$ .*

One crucial difference must be noted. We cannot use the reverse Hölder inequality in this setting to extend the range of exponents to  $r > p - \delta$ . If we could, then by Corollary 5.4 and Proposition 5.5 for each weight  $W \in A_p$  there would exist numbers  $r < q < p$  such that  $W^{q/r} \in A_q \subset A_p$ . Instead, counterexamples are known; in [1] a matrix  $A_2$  weight  $W$  is constructed for which  $W^s \notin A_2$  for any  $s > 1$ .

On a speculative note, perhaps this (suspected) lack of self-improvement is related to the absence of a unifying matrix  $A_\infty$  class whose elements are all contained in some  $A_p$  with  $p$  finite. We do not claim to have proven anything here, nor have we investigated thoroughly the union of the  $A_p$ -weight classes in search of a common  $A_\infty$  property. It has been suggested [25] that the scalar  $A_\infty$  condition generalizes instead to an entire spectrum of  $A_{p,\infty}$  conditions, one for each exponent  $p$ , in the matrix setting.

## 6. The case $d = \infty$ .

Most of the estimates in the preceding discussion fail when the dimension  $d$  is infinite. Banach space norms may not be representable by matrices, and traces (when defined at all) are no longer comparable to operator norms. Most importantly, the main theorem is false. Gillespie et al. [9] have constructed operator  $A_2$  weights  $W$  for which the Hilbert Transform is unbounded on  $L^2(W)$ .

The test function  $f$  in their counterexample is constructed out of Haar functions on different length-scales, with the signs chosen so that each new piece contributes positively to the overall  $L^2(W)$  norm of  $Tf$ . Linearity of  $T$  is needed to ensure that the whole of  $Tf$  will be equal to the sum of the

<sup>1</sup>Following [16], with the slight modification  $F_z(\psi) = |\psi|^{\frac{\alpha(z)}{\alpha}-1}\psi$ .

various parts, and also to bound from below an expectation over choices of signs. When applied to merely sublinear operators such as a maximal function, the argument is less successful. We do not presently know if the Hardy-Littlewood maximal operator  $M_w$  is bounded or not.

## References

- [1] M. Bownik, *Inverse volume inequalities for matrix weights*, Indiana Univ. Math. J., **50**(1) (2001), 383-410, MR 2002m:47043, Zbl 0992.42006.
- [2] M. Christ, *Weighted norm inequalities and Schur's lemma*, Studia Math., **78** (1984), 309-319, MR 86h:42032, Zbl 0555.42008.
- [3] M. Christ and R. Fefferman, *A note on weighted norm inequalities*, Proc. Amer. Math. Soc., **87** (1983), 447-448, MR 84g:42017, Zbl 0502.42017.
- [4] M. Christ and M. Goldberg, *Vector  $A_2$  weights and a Hardy-Littlewood maximal function*, Trans. Amer. Math. Soc., **353** (2001), 1995-2002, CMP 1 813 604, Zbl 0995.42015.
- [5] R.R. Coifman and C. Fefferman, *Weighted norm inequalities for maximal functions and singular integrals*, Studia Math., **51** (1974), 241-250, MR 50 #10670, Zbl 0291.44007.
- [6] R. Fefferman, C.E. Kenig and J. Pipher, *The theory of weights and the Dirichlet problem for elliptic equations*, Ann. Math., **134** (1991), 65-124, MR 93h:31010, Zbl 0770.35014.
- [7] J. Garnett and P.W. Jones, *BMO from dyadic BMO*, Pacific J. Math., **99** (1982), 351-371, MR 85d:42021, Zbl 0516.46021.
- [8] A. Gillespie, F. Nazarov, S. Pott, S. Treil' and A. Volberg, *Logarithmic growth for matrix martingale transform*. Preprint, Michigan State Univ., 1998.
- [9] ———, *Logarithmic growth for weighted Hilbert transform and vector Hankel operators*. Preprint, Michigan State Univ., 1998.
- [10] R. Hunt, B. Muckenhoupt and R. Wheeden, *Weighted norm inequalities for conjugate function and Hilbert transform*, Trans. Amer. Math. Soc., **176** (1973), 227-251, MR 47 #701, Zbl 0262.44004.
- [11] B. Muckenhoupt, *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc., **165** (1972), 207-226, MR 45 #2461, Zbl 0236.26016.
- [12] F.L. Nazarov and S.R. Treil', *The hunt for a Bellman function: Applications to estimates for singular integral operators and to other classical problems of harmonic analysis*, Algebra i Analiz, **8**(5) (1996), 32-162 (Russian); translation in St. Petersburg Math. J., **8**(5) (1997), 721-824, MR 99d:42026, Zbl 0873.42011.
- [13] ———, *The weighted norm inequalities for Hilbert transform are now trivial*, C.R. Acad. Sci. Paris Sér. I Math., **323**(7) (1996), 717-722, MR 99j:42010, Zbl 0858.44004.
- [14] F. Nazarov, S.R. Treil' and A. Volberg, *Counterexample to the infinite-dimensional Carleson embedding theorem*, C.R. Acad. Sci. Paris Sér. I Math., **325**(4) (1997), 383-388, MR 98d:46039, Zbl 0892.42013.
- [15] ———, *The Bellman functions and two-weight inequalities for Haar multipliers*, J. Amer. Math. Soc., **12**(4) (1999), 909-928, MR 2000k:42009, Zbl 0951.42007.

- [16] E.M. Stein, *Interpolation of linear operators*, Trans. Amer. Math. Soc., **83** (1956), 482-492, MR 18,575d, Zbl 0072.32402.
- [17] ———, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, NJ, 1970, MR 44 #7280, Zbl 0207.13501.
- [18] ———, *Harmonic Analysis*, Princeton University Press, Princeton, NJ, 1993, MR 95c:42002, Zbl 0821.42001.
- [19] S.R. Treil', *The geometric approach to weight estimates of the Hilbert transform*, Funktsional. Anal. i Prilozhen., **17**(4) (1983), 90–91 (Russian), MR 85c:42012.
- [20] ———, *An operator approach to weighted estimates of singular integrals*, Investigations on linear operators and the theory of functions, **XIII**, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov (LOMI) **135** (1984), 150-174 (Russian), MR 86g:42032, Zbl 0577.47009.
- [21] ———, *Geometric methods in spectral theory of vector valued functions: Some recent results*, Operator Theory: Adv. Appl., **42** (1989), 209-280, MR 91j:47036, Zbl 0699.47009.
- [22] S.R. Treil' and A.L. Volberg, *Weighted embeddings and weighted norm inequalities for the Hilbert transform and the maximal operator*, Algebra i Analiz, **7**(6) (1995), 205-226 (Russian); translation in St. Petersburg Math. J., **7**(6) (1996), 1017-1032, MR 97c:42017, Zbl 0852.42006, Zbl 0861.42006.
- [23] ———, *Wavelets and the angle between past and future*, J. Funct. Anal., **143**(2) (1997), 269-308, MR 99k:42073, Zbl 0876.42027.
- [24] ———, *Continuous frame decomposition and a vector Hunt-Muckenhoupt-Wheeden theorem*, Ark. Mat., **35**(2) (1997), 363-386, MR 98m:42026, Zbl 0961.44006.
- [25] A. Volberg, *Matrix  $A_p$  weights via  $S$ -functions*, J. Amer. Math. Soc., **10**(2) (1997), 445-466, MR 98a:42013, Zbl 0877.42003.

Received July 30, 2001 and revised May 28, 2002. The author was supported in part by an NSF graduate research fellowship.

DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CA 94720-3840  
*E-mail address:* mikeg@math.berkeley.edu