

# *Pacific Journal of Mathematics*

REAL PALEY-WIENER THEOREMS FOR THE INVERSE  
FOURIER TRANSFORM ON A RIEMANNIAN  
SYMMETRIC SPACE

NILS BYRIAL ANDERSEN

# REAL PALEY-WIENER THEOREMS FOR THE INVERSE FOURIER TRANSFORM ON A RIEMANNIAN SYMMETRIC SPACE

NILS BYRIAL ANDERSEN

**We prove real Paley-Wiener theorems for the inverse Fourier transform on a semisimple Riemannian symmetric space  $G/K$  of the noncompact type. The functions on  $G/K$  whose Fourier transform has compact support are characterised by a  $L^2$  growth condition. We also obtain real Paley-Wiener theorems for the inverse spherical transform.**

## 1. Introduction.

The classical Fourier transform  $\mathcal{F}_{\text{cl}}$  is an isomorphism of the Schwartz space  $\mathcal{S}(\mathbb{R}^k)$  onto itself. The space  $C_c^\infty(\mathbb{R}^k)$  of smooth functions with compact support is dense in  $\mathcal{S}(\mathbb{R}^k)$ , and the classical Paley-Wiener theorem characterises the image of  $C_c^\infty(\mathbb{R}^k)$  under  $\mathcal{F}_{\text{cl}}$  as rapidly decreasing functions having an holomorphic extension to  $\mathbb{C}^k$  of exponential type. Since  $\mathbb{R}^k$  is self-dual, the same theorem also applies to the inverse Fourier transform.

Let  $G$  be a noncompact semisimple Lie group and  $K$  a maximal compact subgroup of  $G$ . The Fourier transform  $\mathcal{F}$  on the Riemannian symmetric space  $X = G/K$  is an analogue of the classical Fourier transform on  $\mathbb{R}^k$ . A Paley-Wiener theorem for the Fourier transform  $\mathcal{F}$ , which characterises the image of  $C_c^\infty(X)$  under  $\mathcal{F}$  in terms of holomorphic extensions and growth behaviour, as in the classical case, was proved by Helgason, see [7]. Furthermore, the  $L^2$ -Schwartz space  $\mathcal{S}^2(X)$  contains  $C_c^\infty(X)$  as a dense subspace and  $\mathcal{F}$  is an isomorphism of  $\mathcal{S}^2(X)$  onto some generalised Schwartz space, see [4].

Unlike the classical case, however, we can not use a duality argument to deduce a Paley-Wiener theorem for the inverse Fourier transform. So how can we characterise the functions whose Fourier transform  $\mathcal{F}$  has compact support?

The Fourier transform on  $X$  reduces to the spherical transform  $\mathcal{H}$  on  $G$  when restricted to  $K$ -invariant functions. The paper [8] provides an answer to the above question for the spherical transform on Schwartz functions in the rank one and complex cases. The characterisation is in analogy with the classical Paley-Wiener theorem given in terms of meromorphic extensions and growth conditions.

In this paper we prove (real) Paley-Wiener theorems for the inverse Fourier transform for general Riemannian symmetric spaces, i.e., we characterise, as a subset of  $L^2(X)$ , the set of functions  $f$  on  $X$  whose Fourier transform  $\mathcal{F}f$  has compact support. More precisely,  $f \in C^\infty(X)$  has to satisfy

$$\lim_{n \rightarrow \infty} \|\Delta^n f\|_2^{1/2n} < \infty,$$

where  $\Delta$  is the Laplace-Beltrami operator (and  $(1 + |\cdot|)^n f \in L^2(X)$  for all  $n \in \mathbb{N} \cup \{0\}$  if we also want the Fourier image to be smooth). Specialising to bi- $K$ -invariant functions yields (real) Paley-Wiener theorems for the inverse spherical transform for general noncompact semisimple Lie groups.

Our approach is based on real analysis techniques developed by H. H. Bang, see [2] and [3], and V.K. Tuan, see [10]. Also see [11] for a history and overview of (real) Paley-Wiener theorems for certain transforms (Fourier, Mellin, Hankel...) on  $\mathbb{R}$ . In particular we use Parseval's formula, intertwining properties of  $\mathcal{F}$ , and the following characterisation of the radius of the support of a function  $g$  on  $\mathbb{R}^n$ :

$$\sup_{\lambda \in \text{supp } g} \|\lambda\| = \lim_{n \rightarrow \infty} \left\{ \int_{\mathbb{R}^n} \|\lambda\|^{4n} |g(\lambda)|^2 d\lambda \right\}^{1/4n}.$$

For completeness and comparison, we first consider the Fourier transform on  $\mathbb{R}^k$ . The results here are originally due to H.H. Bang, see [2] and [3], and V.K. Tuan, see [10]. Notice the beautiful symmetry between the (statements of the) results for the various transforms.

## 2. The Fourier transform on $\mathbb{R}^k$ .

For background and details, please see [9, Chapter 7]. Let  $\mathcal{F}_{\text{cl}}$  denote the classical Fourier transform on  $\mathbb{R}^k$ :

$$\mathcal{F}_{\text{cl}} f(\lambda) := \int_{\mathbb{R}^k} f(x) e^{-i\lambda \cdot x} dx,$$

defined for nice functions  $f$ , for all  $\lambda \in \mathbb{C}^k$  for which the above integral makes sense. Let  $\Delta = \frac{d^2}{dx_1^2} + \cdots + \frac{d^2}{dx_k^2}$  denote the Laplacian on  $\mathbb{R}^k$  and let  $\mathcal{S}(\mathbb{R}^k)$  denote the Schwartz space of rapidly decreasing differentiable functions. Then  $\mathcal{F}_{\text{cl}}(\Delta f)(\lambda) = -\|\lambda\|^2 \mathcal{F}_{\text{cl}} f(\lambda)$ , ( $\lambda \in \mathbb{R}^k$ ), for all  $f \in \mathcal{S}(\mathbb{R}^k)$ , and the Fourier transform is an isomorphism of  $\mathcal{S}(\mathbb{R}^k)$  onto itself, with inverse given by:

$$\mathcal{F}_{\text{cl}}^{-1} g(x) = (2\pi)^{-k} \int_{\mathbb{R}^k} g(\lambda) e^{i\lambda \cdot x} d\lambda, \quad (x \in \mathbb{R}^k)$$

for  $g \in \mathcal{S}(\mathbb{R}^k)$ . Parseval's formula states that

$$\begin{aligned} \langle f_1, f_2 \rangle &:= \int_{\mathbb{R}^k} f_1(x) \overline{f_2(x)} dx = (2\pi)^{-k} \int_{\mathbb{R}^k} \mathcal{F}_{\text{cl}} f_1(\lambda) \overline{\mathcal{F}_{\text{cl}} f_2(\lambda)} d\lambda \\ &=: \langle \mathcal{F}_{\text{cl}} f_1, \mathcal{F}_{\text{cl}} f_2 \rangle, \end{aligned}$$

for  $f_1, f_2 \in \mathcal{S}(\mathbb{R}^k)$ , which implies that  $\|f\|_2 = \|\mathcal{F}_{\text{cl}} f\|_2$ , for all  $f \in \mathcal{S}(\mathbb{R}^k)$ , and hence that the Fourier transform extends to an isometry from  $L^2(\mathbb{R}^k)$  onto itself.

Let  $f \in C^\infty(\mathbb{R}^k)$  such that  $\Delta^n f \in L^2(\mathbb{R}^k)$  for all  $n \in \mathbb{N} \cup \{0\}$  and let  $f_2 \in C_c^\infty(\mathbb{R}^k)$ . Then:

$$\begin{aligned} \langle \mathcal{F}_{\text{cl}}(\Delta f), \mathcal{F}_{\text{cl}} f_2 \rangle &= \langle \Delta f, f_2 \rangle = \langle f, \Delta f_2 \rangle = \langle \mathcal{F}_{\text{cl}} f, \mathcal{F}_{\text{cl}}(\Delta f_2) \rangle \\ &= \langle \mathcal{F}_{\text{cl}} f, -\|\lambda\|^2 \mathcal{F}_{\text{cl}} f_2 \rangle = \langle -\|\lambda\|^2 \mathcal{F}_{\text{cl}} f, \mathcal{F}_{\text{cl}} f_2 \rangle, \end{aligned}$$

and we conclude that  $\mathcal{F}_{\text{cl}}(\Delta f)(\lambda) = -\|\lambda\|^2 \mathcal{F}_{\text{cl}} f(\lambda)$  a.e., by a density argument, whence  $\mathcal{F}_{\text{cl}}(\Delta^n f)(\lambda) = (-1)^n \|\lambda\|^{2n} \mathcal{F}_{\text{cl}} f(\lambda)$  a.e., and

$$(1) \quad \int_{\mathbb{R}^k} |\Delta^n f(x)|^2 dx = (2\pi)^{-k} \int_{\mathbb{R}^k} \|\lambda\|^{4n} |\mathcal{F}_{\text{cl}} f(\lambda)|^2 d\lambda,$$

for all  $n \in \mathbb{N} \cup \{0\}$ .

We define the support,  $\text{supp } g$ , of  $g \in L^2(\mathbb{R}^k)$  to be the smallest closed set, outside which the function  $g$  vanishes almost everywhere, and  $R_g := \sup_{\lambda \in \text{supp } g} \|\lambda\|$  to be the radius of the support of  $g$ ;  $R_g$  is finite if, and only if,  $g$  has compact support.

**Lemma 2.1.** *Let  $g \in L^2(\mathbb{R}^k)$  such that  $\|\lambda\|^{2n} g(\lambda) \in L^2(\mathbb{R}^k)$  for all  $n \in \mathbb{N} \cup \{0\}$ . Then*

$$R_g = \lim_{n \rightarrow \infty} \left\{ \int_{\mathbb{R}^k} \|\lambda\|^{4n} |g(\lambda)|^2 d\lambda \right\}^{1/4n}.$$

*Proof.* Assume  $g$  has compact support with  $R_g > 0$ . Then:

$$\begin{aligned} &\limsup_{n \rightarrow \infty} \left\{ \int_{\mathbb{R}^k} \|\lambda\|^{4n} |g(\lambda)|^2 d\lambda \right\}^{1/4n} \\ &\leq R_g \limsup_{n \rightarrow \infty} \left\{ \int_{\|\lambda\| \leq R_g} |g(\lambda)|^2 d\lambda \right\}^{1/4n} = R_g. \end{aligned}$$

On the other hand,

$$\int_{R_g - \varepsilon \leq \|\lambda\| \leq R_g} |g(\lambda)|^2 d\lambda > 0,$$

for any  $\varepsilon > 0$ , hence

$$\begin{aligned}
& \liminf_{n \rightarrow \infty} \left\{ \int_{\mathbb{R}^k} \|\lambda\|^{4n} |g(\lambda)|^2 d\lambda \right\}^{1/4n} \\
& \geq \liminf_{n \rightarrow \infty} \left\{ \int_{R_g - \varepsilon \leq \|\lambda\| \leq R_g} \|\lambda\|^{4n} |g(\lambda)|^2 d\lambda \right\}^{1/4n} \\
& \geq (R_g - \varepsilon) \liminf_{n \rightarrow \infty} \left\{ \int_{R_g - \varepsilon \leq \|\lambda\| \leq R_g} |g(\lambda)|^2 d\lambda \right\}^{1/4n} = R_g - \varepsilon,
\end{aligned}$$

and thus

$$\lim_{n \rightarrow \infty} \left\{ \int_{\mathbb{R}^k} \|\lambda\|^{4n} |g(\lambda)|^2 d\lambda \right\}^{1/4n} = R_g.$$

Now assume that  $g$  has unbounded support. Then

$$\int_{\|\lambda\| \geq N} |g(\lambda)|^2 d\lambda > 0,$$

for any  $N > 0$ , so:

$$\begin{aligned}
& \liminf_{n \rightarrow \infty} \left\{ \int_{\mathbb{R}^k} \|\lambda\|^{4n} |g(\lambda)|^2 d\lambda \right\}^{1/4n} \\
& \geq \liminf_{n \rightarrow \infty} \left\{ \int_{\|\lambda\| \geq N} \|\lambda\|^{4n} |g(\lambda)|^2 d\lambda \right\}^{1/4n} \\
& \geq N \liminf_{n \rightarrow \infty} \left\{ \int_{\|\lambda\| \geq N} |g(\lambda)|^2 d\lambda \right\}^{1/4n} = N,
\end{aligned}$$

for arbitrary  $N > 0$ , and we conclude that

$$\liminf_{n \rightarrow \infty} \left\{ \int_{\mathbb{R}^k} \|\lambda\|^{4n} |g(\lambda)|^2 d\lambda \right\}^{1/4n} = \infty.$$

□

Let  $L_c^2(\mathbb{R}^k)$  denote the subspace of  $L^2(\mathbb{R}^k)$  of functions with compact support and let  $L_R^2(\mathbb{R}^k) := \{g \in L_c^2(\mathbb{R}^k) \mid R_g = R\}$ . Let also  $C_R^\infty(\mathbb{R}^k) := \{g \in C_c^\infty(\mathbb{R}^k) \mid R_g = R\}$ .

**Definition 2.2.** We define the  $L^2$ -Paley-Wiener space  $\text{PW}^2(\mathbb{R}^k)$  to be the space of all functions  $f \in C^\infty(\mathbb{R}^k)$  satisfying:

- (a)  $\Delta^n f \in L^2(\mathbb{R}^k)$  for all  $n \in \mathbb{N} \cup \{0\}$ .
- (b)  $R_f^\Delta := \lim_{n \rightarrow \infty} \|\Delta^n f\|_2^{1/2n} < \infty$ .

Let also  $\text{PW}_R^2(\mathbb{R}^k) := \{f \in \text{PW}^2(\mathbb{R}^k) \mid R_f^\Delta = R\}$ , for  $R \geq 0$ .

The proof of Theorem 2.3 below shows that the limit in (b) above is well-defined. The real version of the  $L^2$ -Paley-Wiener theorem for the inverse Fourier transform can now be formulated as follows:

**Theorem 2.3.** *The inverse Fourier transform  $\mathcal{F}_{\text{cl}}^{-1}$  is a bijection of  $L_c^2(\mathbb{R}^k)$  onto  $\text{PW}^2(\mathbb{R}^k)$ , mapping  $L_R^2(\mathbb{R}^k)$  onto  $\text{PW}_R^2(\mathbb{R}^k)$ .*

*Proof.* Let  $g \in L_R^2(\mathbb{R}^k)$ . Then  $\|\lambda\|^n g(\lambda) \in L^1(\mathbb{R}^k)$  for all  $n \in \mathbb{N} \cup \{0\}$ , and  $\mathcal{F}_{\text{cl}}^{-1}g \in C_o^\infty(\mathbb{R}^k)$ . We also have  $\Delta^n(\mathcal{F}_{\text{cl}}^{-1}g) = \mathcal{F}_{\text{cl}}^{-1}((-1)^n \|\lambda\|^{2n} g) \in L^2(\mathbb{R}^k)$  for all  $n \in \mathbb{N} \cup \{0\}$ , by the formula for  $\mathcal{F}_{\text{cl}}^{-1}$ , and (1) thus yields:

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left\{ \int_{\mathbb{R}^k} |\Delta^n(\mathcal{F}_{\text{cl}}^{-1}g)(x)|^2 dx \right\}^{1/4n} \\ &= \lim_{n \rightarrow \infty} \left\{ (2\pi)^{-k} \int_{\mathbb{R}^k} \|\lambda\|^{4n} |g(\lambda)|^2 d\lambda \right\}^{1/4n} = R, \end{aligned}$$

whence  $\mathcal{F}_{\text{cl}}^{-1}g \in \text{PW}_R^2(\mathbb{R}^k)$ .

Let now  $f \in \text{PW}_R^2(\mathbb{R}^k)$ . Then  $\mathcal{F}_{\text{cl}}(\Delta^n f)(\lambda) = (-1)^n \|\lambda\|^{2n} \mathcal{F}_{\text{cl}} f(\lambda) \in L^2(\mathbb{R}^k)$  for all  $n \in \mathbb{N}$ , and another application of (1) shows that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left\{ (2\pi)^{-k} \int_{\mathbb{R}^k} \|\lambda\|^{4n} |\mathcal{F}_{\text{cl}} f(\lambda)|^2 d\lambda \right\}^{1/4n} \\ &= \lim_{n \rightarrow \infty} \left\{ \int_{\mathbb{R}^k} |\Delta^n f(x)|^2 dx \right\}^{1/4n} = R, \end{aligned}$$

and we conclude that  $\mathcal{F}_{\text{cl}} f$  has compact support with  $R_{\mathcal{F}_{\text{cl}} f} = R$ .  $\square$

**Remark 2.4.** The classical (complex)  $L^2$ -Paley-Wiener theorem implies that  $\text{PW}_R^2(\mathbb{R}^k)$  exactly consists of those  $L^2(\mathbb{R}^k)$  functions that can be extended to holomorphic functions of exponential type  $R$  on  $\mathbb{C}^k$ .

**Remark 2.5.** Let  $f \in \text{PW}^2(\mathbb{R})$ . Then  $\frac{d^n}{dx^n} f \in L^p(\mathbb{R})$  for all  $n \in \mathbb{N} \cup \{0\}$ , and:

$$\lim_{n \rightarrow \infty} \left\| \frac{d^n}{dx^n} f \right\|_p^{1/n} = R_{\mathcal{F}_{\text{cl}} f} = R_f^\Delta,$$

for all  $1 \leq p \leq \infty$ . This follows from [2, Theorem 1]. Similar results hold for  $\mathbb{R}^k$ ,  $k > 1$ , see [3, Theorem 3] and [10, Theorem 4].

**Definition 2.6.** We define the Paley-Wiener space  $\text{PW}(\mathbb{R}^k)$  as the space of all functions  $f \in C^\infty(\mathbb{R}^k)$  satisfying:

- (a)  $(1 + |x|)^m \Delta^n f \in L^2(\mathbb{R}^k)$  for all  $m, n \in \mathbb{N} \cup \{0\}$ .
- (b)  $R_f^\Delta := \lim_{n \rightarrow \infty} \|\Delta^n f\|_2^{1/2n} < \infty$ .

Let again  $\text{PW}_R(\mathbb{R}^k) := \{f \in \text{PW}(\mathbb{R}^k) \mid R_f^\Delta = R\}$ , for  $R \geq 0$ .

We notice that the only difference between  $\text{PW}^2(\mathbb{R}^k)$  and  $\text{PW}(\mathbb{R}^k)$  is an extra requirement of polynomial decay, to help ensure that  $\mathcal{F}_{\text{cl}}f \in C^\infty(\mathbb{R}^k)$ .

The real version of the Paley-Wiener theorem for the inverse Fourier transform is the following:

**Theorem 2.7.** *The inverse Fourier transform  $\mathcal{F}_{\text{cl}}^{-1}$  is a bijection of  $C_c^\infty(\mathbb{R}^k)$  onto  $\text{PW}(\mathbb{R}^k)$ , mapping  $C_R^\infty(\mathbb{R}^k)$  onto  $\text{PW}_R(\mathbb{R}^k)$ .*

*Proof.* Let  $g \in C_R^\infty(\mathbb{R}^k)$ , then  $\mathcal{F}_{\text{cl}}^{-1}g \in \mathcal{S}(\mathbb{R}^k)$ , and  $\mathcal{F}_{\text{cl}}^{-1}g \in \text{PW}_R^2(\mathbb{R}^k)$  by Theorem 2.3.

Let  $f \in \text{PW}_R(\mathbb{R}^k) \subset \text{PW}_R^2(\mathbb{R}^k)$ . Then  $\mathcal{F}_{\text{cl}}f \in C^\infty(\mathbb{R}^k)$  since  $f$  has polynomial decay, and  $\mathcal{F}_{\text{cl}}f$  has compact support with  $R_{\mathcal{F}_{\text{cl}}f} = R$  by Theorem 2.3.  $\square$

### 3. Lie group notation.

In this section we introduce the Lie group notation we need in the next sections. We refer to [5], [6] and [7] for further details.

Let  $G$  be a real connected noncompact semisimple Lie group with finite center and let  $\theta$  be a Cartan involution of  $G$ . Then the fixed point group  $K := G^\theta$  is a maximal compact subgroup. Let  $\mathfrak{g}$  and  $\mathfrak{k}$  denote their Lie algebras, and let  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$  be the Cartan decomposition of  $\mathfrak{g}$  into the  $\pm 1$  eigenspaces of  $\theta$ . The Killing form on  $\mathfrak{g}$  induces an  $\text{Ad}K$ -invariant scalar product on  $\mathfrak{p}$  and hence a  $G$ -invariant Riemannian metric on  $X := G/K$ . With this structure,  $X$  becomes a Riemannian globally symmetric space of the noncompact type.

Fix a maximal abelian subspace  $\mathfrak{a}$  of  $\mathfrak{p}$ . Denote its real dual by  $\mathfrak{a}^*$  and its complex dual by  $\mathfrak{a}_{\mathbb{C}}^*$ . The Killing form of  $\mathfrak{g}$  induces a scalar product  $\langle \cdot, \cdot \rangle$  and hence a norm  $\|\cdot\|$  on  $\mathfrak{a}_{\mathbb{C}}$  and  $\mathfrak{a}_{\mathbb{C}}^*$ . Let  $\Sigma \subset \mathfrak{a}^*$  denote the root system of  $(\mathfrak{g}, \mathfrak{a})$  and let  $W$  be the associated Weyl group. Choose a set  $\Sigma_+ \subset \Sigma$  of positive roots, let  $\mathfrak{n} := \bigoplus_{\alpha \in \Sigma_+} \mathfrak{g}_\alpha$  be the corresponding nilpotent subalgebra of  $\mathfrak{g}$  and let  $\mathfrak{a}_+ := \{H \in \mathfrak{a} \mid \alpha(H) > 0 \forall \alpha \in \Sigma_+\}$  be the positive Weyl chamber with  $\overline{\mathfrak{a}_+}$  its closure. Denote by  $\mathfrak{a}_+^*$  and  $\overline{\mathfrak{a}_+^*}$  the similar cones in  $\mathfrak{a}^*$ , and define the element  $\rho \in \mathfrak{a}^*$  by:  $\rho(H) := \frac{1}{2} \sum_{\alpha \in \Sigma_+} m_\alpha \alpha(H)$ ,  $H \in \mathfrak{a}$ , where  $m_\alpha = \dim \mathfrak{g}_\alpha$ .

Let  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{a} \oplus \mathfrak{n}$  be the Iwasawa decomposition of  $\mathfrak{g}$  and  $G = KAN = NAK$  the corresponding Iwasawa decompositions of  $G$ , where  $A$  and  $N$  are the Lie groups generated by  $\mathfrak{a}$  and  $\mathfrak{n}$  respectively. Every  $g \in G$  can be represented as:  $g = K \exp H(g)N = N \exp A(g)K$ , where the projections onto the  $A$ -parts  $A(g) \in \mathfrak{a}$  and  $H(g) \in \mathfrak{a}$  are uniquely determined. We note that  $A(g) = -H(g^{-1})$ . Let  $M := Z_K(\mathfrak{a})$ , then  $B := K/M$  is a compact homogeneous space. We define the vector  $A(x, b) \in \mathfrak{a}$  as  $A(x, b) := A(k^{-1}g)$ , for  $x = gK \in X$  and  $b = kM \in B$ .

Put  $A_+ = \exp(\mathfrak{a}_+)$ , then  $\overline{A_+} = \exp(\overline{\mathfrak{a}_+})$ . The Cartan decomposition implies that the natural mapping from  $K/M \times A_+ \times K$  into  $G = K\overline{A_+}K$  is a

diffeomorphism onto its dense open image. We define the norm of an element  $g \in G$  as:  $|g| = |k_1 \exp(H)k_2| = \|H\|$ , with  $H \in \bar{\mathfrak{a}}_+$ ; this is the  $K$ -invariant geodesic distance to the origin  $eK$ . We denote by  $B_R := \{g \in G \mid |g| \leq R\}$  the  $K$ -invariant ball of radius  $R$  around  $e$ .

We identify functions on  $X$  with right- $K$ -invariant functions on  $G$ . We normalise the invariant measure on  $X$  as:

$$\int_X f(x)dx = \int_K \int_{\mathfrak{a}_+} \int_K f(k_1 \exp(H)k_2) J(H) dk_1 dH dk_2,$$

for  $f \in C_c^\infty(X)$ , where the Jacobian  $J$  is given by:  $J(H) = \prod_{\alpha \in \Sigma_+} (e^{\alpha(H)} - e^{-\alpha(H)})^{m_\alpha}$ ,  $dH$  is the Lebesgue measure on  $\mathfrak{a}$  and  $dk$  is the measure on  $K$  such that  $\int_K dk = 1$ . We notice that  $0 \leq J(H) \leq Ce^{2\rho(H)}$ , for  $H \in \bar{\mathfrak{a}}_+$ , where  $C$  is a positive constant.

The spherical functions  $\varphi_\lambda$ ,  $\lambda \in \mathfrak{a}_\mathbb{C}^*$ , on  $G$  are defined as:

$$\varphi_\lambda(g) := \int_K e^{(i\lambda+\rho)A(k^{-1}g)} dk = \int_K e^{-(i\lambda+\rho)H(g^{-1}k)} dk.$$

We note that  $\varphi_\lambda$  is Weyl group invariant,  $\varphi_{w\lambda} = \varphi_\lambda$ ,  $w \in W$ . Let  $U(\mathfrak{g})$  denote the universal enveloping algebra of  $\mathfrak{g}$ . We write  $Df(g)$  for the action of  $D \in U(\mathfrak{g})$  on  $f \in C^\infty(G)$  from the left at  $g \in G$ . The  $L^p$ -Schwartz space  $\mathcal{S}^p(X)$ ,  $0 < p \leq 2$ , is defined as the space of all functions  $f \in C^\infty(X)$  such that:

$$\sup_{g \in G} (1 + |g|)^m \varphi_o(g)^{-\frac{2}{p}} |Df(g)| < \infty,$$

for all  $D \in U(\mathfrak{g})$  and  $m \in \mathbb{N} \cup \{0\}$ . We can also characterise  $\mathcal{S}^p(X)$  as the space of all functions  $f \in C^\infty(X)$  satisfying:

$$(1 + |g|)^m Df(g) \in L^p(X),$$

for all  $D \in U(\mathfrak{g})$  and  $m \in \mathbb{N} \cup \{0\}$ . We note that  $\mathcal{S}^p(X) \not\subset L^q(X)$  for  $0 < q < p \leq 2$ .

#### 4. The Fourier transform.

In this section, we recall some facts and theorems for the Fourier transform on a noncompact semisimple Riemannian symmetric space, see [7, Chapter 3] for details and references.

The Fourier transform of a function  $f$  on  $X$  is defined as:

$$\mathcal{F}f(\lambda, b) := \int_X f(x) e^{(-i\lambda+\rho)(A(x,b))} dx,$$

for all  $\lambda \in \mathfrak{a}_\mathbb{C}^*$ ,  $b \in B$  for which the integral exists. In particular,  $\mathcal{F}f$  extends to a smooth function on  $\mathfrak{a}_\mathbb{C}^* \times B$ , holomorphic in the first variable, for  $f \in C_c^\infty(X)$ , see also below.



The plane wave eigenfunction

$$(2) \quad e_{\lambda,b}(x) := e^{(i\lambda+\rho)(A(x,b))},$$

is a joint eigenfunction of  $\mathbb{D}(X)$ , the commutative algebra of  $G$ -invariant differential operators on  $X$ , for all  $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$ ,  $b \in B$ , or, more precisely

$$De_{\lambda,b} = \Gamma(D)(i\lambda)e_{\lambda,b}, \quad \forall D \in \mathbb{D}(X), \quad (\lambda \in \mathfrak{a}_{\mathbb{C}}^*, b \in B)$$

where  $\Gamma : \mathbb{D}(X) \rightarrow S(\mathfrak{a}^*)^W$  is the Harish-Chandra isomorphism. In particular,

$$\Delta e_{\lambda,b} = -(\langle \lambda, \lambda \rangle + \|\rho\|^2)e_{\lambda,b}, \quad (\lambda \in \mathfrak{a}_{\mathbb{C}}^*, b \in B)$$

for the Laplace-Beltrami operator  $\Delta$  on  $X$ , and hence

$$\mathcal{F}(\Delta f)(\lambda, b) = -(\langle \lambda, \lambda \rangle + \|\rho\|^2)\mathcal{F}f(\lambda, b), \quad (\lambda \in \mathfrak{a}_{\mathbb{C}}^*, b \in B)$$

for all  $f \in C_c^\infty(X)$ , by self-adjointness of  $\Delta$ , see also (6).

A  $C^\infty$ -function  $\psi(\lambda, b)$  on  $\mathfrak{a}_{\mathbb{C}}^* \times B$ , holomorphic in  $\lambda$ , is called a holomorphic function of uniform exponential type  $R$ , if there exists a constant  $R \geq 0$ , such that, for each  $N \in \mathbb{N}$ , we have:

$$\sup_{\lambda \in \mathfrak{a}_{\mathbb{C}}^*, b \in B} e^{-R|\Im \lambda|}(1 + |\lambda|)^N |\psi(\lambda, b)| < \infty.$$

The space of holomorphic functions of uniform exponential type  $R$  will be denoted  $\mathcal{H}_R(\mathfrak{a}_{\mathbb{C}}^* \times B)$  and we denote by  $\mathcal{H}(\mathfrak{a}_{\mathbb{C}}^* \times B)$  their union over all  $R > 0$ . Let furthermore  $\mathcal{H}(\mathfrak{a}_{\mathbb{C}}^* \times B)^W$  denote the space of all functions  $\psi \in \mathcal{H}(\mathfrak{a}_{\mathbb{C}}^* \times B)$  satisfying the symmetry condition:

$$(3) \quad \int_B e^{(i w \lambda + \rho)(A(x,b))} \psi(w\lambda, b) db = \int_B e^{(i \lambda + \rho)(A(x,b))} \psi(\lambda, b) db,$$

for  $w \in W$  and all  $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$ ,  $x \in X$ .

The Paley-Wiener theorem states that the Fourier transform is a bijection of the space  $C_c^\infty(X)$  onto the space  $\mathcal{H}(\mathfrak{a}_{\mathbb{C}}^* \times B)^W$ , with the following inversion formula:

$$(4) \quad f(x) = \int_{\mathfrak{a}_+^*} \int_B e^{(i \lambda + \rho)(A(x,b))} \mathcal{F}f(\lambda, b) |c(\lambda)|^{-2} d\lambda db, \quad (x \in X)$$

where  $c(\lambda)$  is the Harish-Chandra  $c$ -function, for  $f \in C_c^\infty(X)$ . Moreover,  $\mathcal{F}f \in \mathcal{H}_R(\mathfrak{a}_{\mathbb{C}}^* \times B)^W$  if, and only if,  $\text{supp } f \subset B_R$ . We note that  $|c(\lambda)|^{-2}$  is bounded by some polynomial for  $\lambda \in \mathfrak{a}^*$ .

Let  $f_1, f_2 \in C_c^\infty(X)$ , then Parseval's formula for  $\mathcal{F}$  is given by:

$$(5) \quad \int_X f_1(x) \overline{f_2(x)} dx = \int_{\mathfrak{a}_+^*} \int_B \mathcal{F}f_1(\lambda, b) \overline{\mathcal{F}f_2(\lambda, b)} |c(\lambda)|^{-2} d\lambda db.$$

We conclude that the Fourier transform extends to an isometry of  $L^2(X)$  onto  $L^2(\mathfrak{a}_+^* \times B, |c(\lambda)|^{-2} d\lambda db)$ . In the following we adopt the convention  $L^2(\mathfrak{a}_+^* \times B) := L^2(\mathfrak{a}_+^* \times B, |c(\lambda)|^{-2} d\lambda db)$ .

Let  $f \in C^\infty(X)$  such that  $\Delta^n f \in L^2(X)$  for all  $n \in \mathbb{N} \cup \{0\}$  and let  $f_2 \in C_c^\infty(X)$ . Then self-adjointness of the Laplace-Beltrami operator  $\Delta$ :

$$(6) \quad \int_X \Delta^n f(x) f_2(x) dx = \int_X f(x) \Delta^n f_2(x) dx,$$

Parseval's formula (5) and density of  $C_c^\infty(X)$  imply, as in the classical case, that

$$(7) \quad \mathcal{F}(\Delta^n f)(\lambda, b) = (-1)^n (\|\lambda\|^2 + \|\rho\|^2)^n \mathcal{F}f(\lambda, b),$$

a.e., for all  $n \in \mathbb{N} \cup \{0\}$ .

### 5. The inverse Fourier transform.

We define the inverse Fourier transform  $\mathcal{F}^{-1}g$  of a function  $g$  on  $\mathfrak{a}_+^* \times B$  via (4):

$$\mathcal{F}^{-1}g(x) := \int_{\mathfrak{a}_+^*} \int_B e^{(i\lambda + \rho)(A(x, b))} g(\lambda, b) |c(\lambda)|^{-2} d\lambda db,$$

for all  $x \in X$  for which the integral exists.

We define the support,  $\text{supp } g$ , of  $g \in L^2(\mathfrak{a}_+^* \times B)$  to be the smallest closed set in  $\mathfrak{a}_+^* \times B$ , outside which the function  $g$  vanishes almost everywhere, and  $R_g := \sup_{(\lambda, b) \in \text{supp } g} \|\lambda\|$  to be the ‘radius’ of the support of  $g$ .

**Lemma 5.1.** *Let  $g \in L^2(\mathfrak{a}_+^* \times B)$  such that  $\|\lambda\|^{2n} g(\lambda, b) \in L^2(\mathfrak{a}_+^* \times B)$  for all  $n \in \mathbb{N} \cup \{0\}$ . Then*

$$R_g = \lim_{n \rightarrow \infty} \left\{ \int_{\mathfrak{a}_+^*} \int_B \|\lambda\|^{4n} |g(\lambda, b)|^2 |c(\lambda)|^{-2} d\lambda db \right\}^{1/4n}.$$

*Proof.* As for Lemma 2.1. □

Let  $L_c^2(\mathfrak{a}_+^* \times B)$  denote the subspace of  $L^2(\mathfrak{a}_+^* \times B)$  of functions with bounded support and let  $L_R^2(\mathfrak{a}_+^* \times B) := \{g \in L_c^2(\mathfrak{a}_+^* \times B) \mid R_g = R\}$ .

**Definition 5.2.** We define the  $L^2$ -Paley-Wiener space  $\text{PW}^2(X)$  as the space of all functions  $f \in C^\infty(X)$  satisfying:

- (a)  $\Delta^n f \in L^2(X)$  for all  $n \in \mathbb{N} \cup \{0\}$ .
- (b)  $R_f^\Delta := \lim_{n \rightarrow \infty} \|(\Delta + \|\rho\|^2)^n f\|_2^{1/2n} < \infty$ .

Let also  $\text{PW}_R^2(X) := \{f \in \text{PW}^2(X) \mid R_f^\Delta = R\}$ , for  $R \geq 0$ .

The real  $L^2$ -Paley-Wiener theorem for the inverse Fourier transform can now be formulated as follows:

**Theorem 5.3.** *The inverse Fourier transform  $\mathcal{F}^{-1}$  is a bijection of  $L_c^2(\mathfrak{a}_+^* \times B)$  onto  $\text{PW}^2(X)$ , mapping  $L_R^2(\mathfrak{a}_+^* \times B)$  onto  $\text{PW}_R^2(X)$ .*

*Proof.* Let  $g \in L_R^2(\mathfrak{a}_+^* \times B)$ . Then  $\mathcal{F}^{-1}g \in C^\infty(X)$  by Lebesgue's dominated convergence theorem. Equation (2) gives, for  $D \in \mathbb{D}(X)$ ,

$$D(\mathcal{F}^{-1}g)(x) = \int_{\mathfrak{a}_+^*} \int_B \Gamma(D)(i\lambda) e^{(i\lambda + \rho)(A(x, b))} g(\lambda, b) |c(\lambda)|^{-2} d\lambda db,$$

which in particular shows that  $(\Delta + \|\rho\|)^n \mathcal{F}^{-1}g = \mathcal{F}^{-1}((-1)^n \|\lambda\|^{2n} g) \in L^2(X)$  for all  $n \in \mathbb{N} \cup \{0\}$ . Parseval's formula (5) with

$$f_1 = f_2 = \mathcal{F}^{-1}((-1)^n \|\lambda\|^{2n} g)$$

yields:

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left\{ \int_X |(\Delta + \|\rho\|)^n (\mathcal{F}^{-1}g)(x)|^2 dx \right\}^{1/4n} \\ &= \lim_{n \rightarrow \infty} \left\{ \int_{\mathfrak{a}_+^*} \int_B \|\lambda\|^{4n} |g(\lambda, b)|^2 |c(\lambda)|^{-2} d\lambda db \right\}^{1/4n} = R, \end{aligned}$$

whence  $\mathcal{F}^{-1}g \in \text{PW}_R^2(X)$ .

Let now  $f \in \text{PW}_R^2(X)$ . Then  $\mathcal{F}((\Delta + \|\rho\|)^n f)(\lambda, b) = (-1)^n \|\lambda\|^{2n} \mathcal{F}f(\lambda, b) \in L^2(\mathfrak{a}_+^* \times B)$  for all  $n \in \mathbb{N}$  by (7). Another application of Parseval's formula as above with  $f_1 = f_2 = (\Delta + \|\rho\|)^n f$  shows that  $R_{\mathcal{F}f} = R_f^\Delta = R$ , and we conclude that  $\mathcal{F}f$  has bounded support.  $\square$

**Corollary 5.4.** *Let  $f \in C^\infty(X)$  be such that  $\Delta^n f \in L^2(X)$  for all  $n \in \mathbb{N} \cup \{0\}$ . It then follows that  $\lim_{n \rightarrow \infty} \|\Delta^n f\|_2^{1/2n} < \infty$  if, and only if,  $\lim_{n \rightarrow \infty} \|(\Delta + \|\rho\|^2)^n f\|_2^{1/2n} < \infty$ . Furthermore,  $\lim_{n \rightarrow \infty} \|\Delta^n f\|_2^{1/2n} = (R^2 + \|\rho\|^2)^{1/2}$ , for  $f \in \text{PW}_R^2(X)$  with  $R > 0$ .*

*Proof.* Let  $f \in \text{PW}_R^2(X)$ , with  $R > 0$ , then  $\mathcal{F}f \in L_R^2(\mathfrak{a}_+^* \times B)$ . Parseval's formula and an easy adaption of the proof of Lemma 2.1 shows that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \|\Delta^n f\|_2^{1/2n} \\ &= \lim_{n \rightarrow \infty} \left\{ \int_{\mathfrak{a}_+^*} \int_B (\|\lambda\|^2 + \|\rho\|^2)^{2n} |\mathcal{F}f(\lambda, b)|^2 |c(\lambda)|^{-2} d\lambda db \right\}^{1/4n} \\ &= (R^2 + \|\rho\|^2)^{1/2}. \end{aligned}$$

Assume that  $\lim_{n \rightarrow \infty} \|\Delta^n f\|_2^{1/2n} < \infty$ . Then  $\mathcal{F}(\Delta^n f)(\lambda, b) = (-1)^n (\|\lambda\|^2 + \|\rho\|^2)^n \mathcal{F}f(\lambda, b) \in L^2(\mathfrak{a}_+^* \times B)$ , for all  $n \in \mathbb{N}$ , and

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left\{ \int_{\mathfrak{a}_+^*} \int_B \|\lambda\|^{4n} |\mathcal{F}f(\lambda, b)|^2 |c(\lambda)|^{-2} d\lambda db \right\}^{1/4n} \\ & \leq \lim_{n \rightarrow \infty} \left\{ \int_{\mathfrak{a}_+^*} \int_B (\|\lambda\|^2 + \|\rho\|^2)^{2n} |\mathcal{F}f(\lambda, b)|^2 |c(\lambda)|^{-2} d\lambda db \right\}^{1/4n} \\ & = \lim_{n \rightarrow \infty} \|\Delta^n f\|_2^{1/2n} < \infty, \end{aligned}$$

that is,  $\mathcal{F}f$  has bounded support.  $\square$

**Remark 5.5.** Assume that  $f \in \mathcal{S}^p(X)$ , with  $0 < p < 2$ , then  $\mathcal{F}f$  extends to an analytic function on a small tube domain around  $\mathfrak{a}^* \times B$  in  $\mathfrak{a}_{\mathbb{C}}^* \times B$ . Hence  $\mathcal{F}f$  cannot have compact support on  $\mathfrak{a}^* \times B$  and we conclude that  $\mathcal{S}^p(X) \cap \text{PW}^2(X) = \{0\}$  for any  $0 < p < 2$ .

**Definition 5.6.** We define the Paley-Wiener space  $\text{PW}(X)$  as the space of all functions  $f \in C^\infty(X)$  satisfying:

- (a)  $(1 + |x|)^m \Delta^n f \in L^2(X)$  for all  $m, n \in \mathbb{N} \cup \{0\}$ .
- (b)  $R_f^\Delta = \lim_{n \rightarrow \infty} \|(\Delta + \|\rho\|^2)^n f\|_2^{1/2n} < \infty$ .

Let also  $\text{PW}_R(X) := \{f \in \text{PW}(X) \mid R_f^\Delta = R\}$ , for  $R \geq 0$ .

Here  $|x| := |g|$ , for  $x = gK \in X$ . Again, the only difference between the Paley-Wiener spaces  $\text{PW}(X)$  and  $\text{PW}^2(X)$  is the polynomial decay condition (a), ensuring that  $\mathcal{F}f \in C^\infty(\mathfrak{a}^* \times B)^W$  (see below).

The space  $C_c^\infty(\mathfrak{a}^* \times B)^W$  is defined as the subspace of functions  $\psi \in C_c^\infty(\mathfrak{a}^* \times B)$  satisfying the symmetry condition (3) for all  $w \in W$  and all  $\lambda \in \mathfrak{a}^*$ ,  $x \in X$ . Let finally  $C_R^\infty(\mathfrak{a}^* \times B) := \{F \in C_c^\infty(\mathfrak{a}^* \times B) \mid R_g = R\}$ .

The real Paley-Wiener theorem for the inverse Fourier transform then is:

**Theorem 5.7.** *The inverse Fourier transform  $\mathcal{F}^{-1}$  is a bijection of  $C_c^\infty(\mathfrak{a}^* \times B)^W$  onto  $\text{PW}(X)$ , mapping  $C_R^\infty(\mathfrak{a}^* \times B)^W$  onto  $\text{PW}_R(X)$ .*

*Proof.* Let  $g \in C_R^\infty(\mathfrak{a}^* \times B)^W$ , then  $g \in L_R^2(\mathfrak{a}_+^* \times B)$  and thus  $\mathcal{F}^{-1}g \in \text{PW}_R^2(X)$  by Theorem 5.3. We furthermore see that  $\mathcal{F}^{-1}g \in \mathcal{S}^2(X)$  by [4, Theorem 4.1.1], whence  $\mathcal{F}^{-1}g$  satisfies the polynomial decay condition (a).

Let now  $f \in \text{PW}_R(X)$ . The basic estimate  $\|A(g)\| \leq C|g|$ , for all  $g \in G$ , gives us a polynomial estimate (in  $x$ ) of the derivatives (with respect to  $\lambda$ ) of the plane wave eigenfunctions  $e_{\lambda, b}(x)$ . It is also well-known that  $(1 + |x|)^{-r} \varphi_0 \in L^2(X)$  for some large  $r \in \mathbb{N}$ . All this, the polynomial decay condition (a), the Cauchy-Schwartz theorem and Lebesgue's dominated convergence theorem imply that  $\mathcal{F}f \in C^\infty(\mathfrak{a}^* \times B)^W$ . Furthermore  $\mathcal{F}f$  has the desired compact support by Theorem 5.3.  $\square$

## 6. The inverse spherical transform.

In this section, we specialise our results to bi- $K$ -invariant functions, that is, we consider the (inverse) spherical transform. We refer to [1], [5] and [6] for background concerning Paley-Wiener theorems for the spherical transform. Let  $C^\infty(K \backslash G / K) \subset C^\infty(G)$  denote the subspace of bi- $K$ -invariant differentiable functions on  $G$ . We will use similar notation for the  $L^2$ , Paley-Wiener and Schwartz spaces of  $K$ -invariant differentiable functions.

Let  $f \in C_c^\infty(K \backslash G / K)$ . The spherical transform  $\mathcal{H}f$  of  $f$  is defined as:

$$\mathcal{H}f(\lambda) := \int_G f(x) \varphi_{-\lambda}(x) dx,$$

for  $\lambda \in \mathfrak{a}_\mathbb{C}^*$ . We note that  $\mathcal{F}f(\lambda, b) = \mathcal{H}f(\lambda)$  for all  $\lambda \in \mathfrak{a}_\mathbb{C}^*$  and all  $b \in B$ . This follows from left- $K$ -invariance of  $f$ , the identity  $A(k \cdot x, b) = A(x, k^{-1} \cdot b)$  and integrating over  $K$ .

The spherical transform is an isomorphism of  $\mathcal{S}^2(K \backslash G / K)$  onto  $\mathcal{S}(\mathfrak{a}^*)^W$ , the Weyl group invariant Schwartz functions on  $\mathfrak{a}^*$ . The inversion formula is given by:

$$(8) \quad f(x) = \frac{1}{|W|} \int_{\mathfrak{a}^*} \mathcal{H}f(\lambda) \varphi_\lambda(x) |c(\lambda)|^{-2} d\lambda, \quad (x \in G)$$

for  $f \in \mathcal{S}^2(K \backslash G / K)$ . We use (8) to define the inverse spherical transform  $\mathcal{H}^{-1}g$  for a general function  $g$  on  $\mathfrak{a}^*$ :

$$\mathcal{H}^{-1}g(x) := \frac{1}{|W|} \int_{\mathfrak{a}^*} g(\lambda) \varphi_\lambda(x) |c(\lambda)|^{-2} d\lambda.$$

Let  $f \in C^\infty(K \backslash G / K)$  be such that  $\Delta^n f \in L^2(K \backslash G / K)$  for all  $n \in \mathbb{N} \cup \{0\}$ . Then  $\mathcal{H}((\Delta + \|\rho\|^2)^n f)(\lambda) = (-1)^n \|\lambda\|^{2n} \mathcal{H}f(\lambda)$  a.e., and Parseval's formula for  $\mathcal{H}$  gives:

$$\int_G |(\Delta + \|\rho\|^2)^n f(x)|^2 dx = \frac{1}{|W|} \int_{\mathfrak{a}^*} \|\lambda\|^{4n} |\mathcal{H}f(\lambda)|^2 |c(\lambda)|^{-2} d\lambda,$$

for all  $n \in \mathbb{N} \cup \{0\}$ . It also follows that the spherical transform extends to an isometry from  $L^2(K \backslash G / K)$  onto  $L^2(\mathfrak{a}^*, \frac{1}{|W|} |c(\lambda)|^{-2} d\lambda)^W$ , where superscript  $W$  denotes Weyl group invariance.

Let  $L_c^2(\mathfrak{a}^*)^W$  denote the Weyl group invariant  $L^2$ -functions on  $\mathfrak{a}^*$  with compact support and let subscript  $R$  denote the radius of the support. The real versions of the Paley-Wiener theorems for the inverse spherical transform then becomes:

**Theorem 6.1.** *The inverse spherical transform  $\mathcal{H}^{-1}$  is a bijection of  $L_c^2(\mathfrak{a}^*)^W$  onto  $\text{PW}^2(K \backslash G / K)$ , mapping  $L_R^2(\mathfrak{a}^*)^W$  onto  $\text{PW}_R^2(K \backslash G / K)$ .*

**Theorem 6.2.** *The inverse spherical transform  $\mathcal{H}^{-1}$  is a bijection of  $C_c^\infty(\mathfrak{a}^*)^W$  onto  $\text{PW}(K \backslash G / K)$ , mapping  $C_R^\infty(\mathfrak{a}^*)^W$  onto  $\text{PW}_R(K \backslash G / K)$ .*

*Proof.* The above theorems are special cases of Theorem 5.3 and Theorem 5.7. We note, however, that we can prove them independently using Parseval's formula and intertwining properties of  $\mathcal{H}$ .  $\square$

**Remark 6.3.** Let  $f \in \text{PW}(K \backslash G / K)$  and consider  $f$  as a function on  $\mathfrak{a}$  by the application  $H \mapsto f(\exp(H))$ . Then  $f$  does not extend to an entire function on  $\mathfrak{a}_{\mathbb{C}}$ , due to the poles of the spherical function  $\varphi_{\lambda}(\exp(H))$ . There is, however, a description of the Paley-Wiener space  $\text{PW}(K \backslash G / K)$  as functions having an explicit meromorphic extension and satisfying some exponential growth conditions for the rank 1 and the complex cases, see [8] for details.

## References

- [1] J.-Ph. Anker, *The spherical Fourier transform of rapidly decreasing functions. A simple proof of a characterization due to Harish-Chandra, Helgason, Trombi, and Varadarajan*, J. Funct. Anal., **96** (1991), 331-349, [MR 92d:22008](#), [Zbl 0732.43006](#).
- [2] H.H. Bang, *A property of infinitely differentiable functions*, Proc. Amer. Math. Soc., **108** (1990), 73-76, [Zbl 0707.26015](#).
- [3] ———, *Functions with bounded spectrum*, Trans. Amer. Math. Soc., **347** (1995), 1067-1080, [MR 95e:42008](#), [Zbl 0828.42009](#).
- [4] M. Eguchi, *Asymptotic expansions of Eisenstein integrals and Fourier transform on symmetric spaces*, J. Funct. Anal., **34** (1979), 167-216, [MR 81e:43022](#), [Zbl 0433.43012](#).
- [5] R. Gangolli and V.S. Varadarajan, *Harmonic Analysis of Spherical Functions on Real Reductive Groups*, Springer-Verlag, Berlin, 1988, [MR 89m:22015](#), [Zbl 0675.43004](#).
- [6] S. Helgason, *Groups and Geometric Analysis*, Academic Press, Orlando, 1984, [MR 86c:22017](#), [Zbl 0543.58001](#).
- [7] ———, *Geometric Analysis on Symmetric Spaces*, Mathematical Surveys and Monographs, **39**, American Mathematical Society, Providence, Rhode Island, 1994, [MR 96h:43009](#), [Zbl 0809.53057](#).
- [8] A. Pasquale, *A Paley-Wiener theorem for the inverse spherical transform*, Pacific J. Math., **193** (2000), 143-176, [MR 2001c:22011](#), [Zbl 1014.22010](#).
- [9] W. Rudin, *Functional Analysis*, McGraw-Hill, New York, 1991, [MR 92k:46001](#), [Zbl 0867.46001](#).
- [10] V.K. Tuan, *Paley-Wiener-type theorems*, Fract. Calc. Appl. Anal., **2** (1999), 135-143.
- [11] V.K. Tuan and A.I. Zayed, *Paley-Wiener-type theorems for a class of integral transforms*, J. Math. Anal. Appl., **266** (2002), 200-226, [MR 2002k:44005](#), [Zbl 0998.44001](#).

Received October 14, 2002 and revised February 11, 2003. The author is supported by a research grant from the Australian Research Council.

SCHOOL OF MATHEMATICS  
 UNIVERSITY OF NEW SOUTH WALES  
 SYDNEY NSW 2052  
 AUSTRALIA  
*E-mail address:* [byrial@maths.unsw.edu.au](mailto:byrial@maths.unsw.edu.au)