

# ***Pacific Journal of Mathematics***

**FORWARD GENERATOR FOR PREIMAGE ENTROPY**

WEN-CHIAO CHENG

Volume 223      No. 1

January 2006

# FORWARD GENERATOR FOR PREIMAGE ENTROPY

WEN-CHIAO CHENG

**We investigate preimage entropy and show how to calculate it under the assumption of forward expansiveness. We then define new invariants of non-invertible maps, called the upper preimage entropy and the metric preimage entropy, and obtain variational principles for them. Last, we prove a similar result for the Kolmogorov–Sinai generator.**

## 1. Introduction

The topological entropy  $h(T)$  of a continuous map  $T$  of a compact metric space to itself is a measure of its dynamical complexity. It was first defined by Adler, Konheim and McAndrew, and later was given several equivalent definitions by Bowen and others (see [Bowen 1972] for an exposition). These definitions have led to results connecting topological and measure-theoretic entropies.

More recently, the preimage relation entropy  $h_r(T)$  of a compact metric space was introduced in [Langevin and Walczak 1991] and used as a new tool for studying the topology and dynamics of endomorphisms of compact metric spaces. Hurley [1995] and Nitecki and Przytycki [1999] then introduced several other entropy-like invariants for noninvertible maps. One, the preimage branch entropy  $h_i(T)$ , is closely related to  $h_r(T)$ . Two others are based on how many branches of the inverse of the iterated map  $T^{-n}$  at a point  $x$  can be distinguished by measurements of finite accuracy; they are called pointwise preimage entropies and are denoted by  $h_p(T)$  and  $h_m(T)$ . The following inequalities hold for any continuous map  $T$  on a compact metric space [Hurley 1995; Nitecki and Przytycki 1999]:

$$h_p(T) \leq h_m(T) \leq h(T) \leq h_i(T) + h_m(T) \leq h_r(T) + h_m(T).$$

In this paper we first concentrate on the pointwise preimage entropies  $h_p(T)$  and  $h_m(T)$ , whose definitions are in some sense analogous to (and were motivated by) Bowen’s notion of “local entropy” (see [Bowen 1972]). The definitions and some basic properties of these invariants are reviewed in [Section 2](#).

---

MSC2000: [37B40](#).

Keywords: preimage entropy, forward generator, expansiveness.

In [Section 3](#) we consider forward expansive maps. We show that, as in the case of the standard topological entropy, both  $h_p(T)$  and  $h_m(T)$  can be computed using the forward generator.

In [Section 4](#) we introduce modified preimage entropies in the topological and measure-theoretic contexts. We obtain variational principles and a similar result for generators in the case of a measure-preserving transformation. We also show that the construction of the metric preimage entropy equals the conditional metric preimage entropy with infinite past  $\sigma$ -algebra.

## 2. Pointwise preimage entropy

As in Adler, Konheim and McAndrew's definition of topological entropy, we take an open cover  $U$  of a compact space  $X$  and a continuous map  $T : X \rightarrow X$ , and set

$$\bigvee_{i=0}^n T^{-i}(U) = \{U_{j_0} \cap T^{-1}U_{j_1} \cap \cdots \cap T^{-n}U_{j_n} \mid U_{j_k} \in U, 0 \leq k \leq n\}.$$

This is an open cover and a refinement of  $U$ . If  $x \in U_{j_0} \cap T^{-1}U_{j_1} \cap \cdots \cap T^{-n}U_{j_n}$ , then  $x \in U_{j_0}$ ,  $Tx \in U_{j_1}$ ,  $\dots$ ,  $T^n x \in U_{j_n}$ .

**Definition 2.1.** Let  $T : X \rightarrow X$  be a continuous map of a compact space  $X$  and take  $x \in X$ . For  $N = 1, 2, \dots$ , the  $N$ -th preimage set of  $x$  under  $T$  is the set  $T^{-N}(x) := \{z \in X \mid T^N(z) = x\}$ , and the  $N$ -th branch at  $x$  is the set

$$B_N(x, T) = \{(z_N, z_{N-1}, \dots, z_0) \mid T(z_{i+1}) = z_i, 0 \leq i \leq N-1 \text{ and } z_0 = x\}.$$

Let  $O(X)$  be the collection of all (finite or infinite) open covers of  $X$ . Given  $U \in O(X)$ , let  $U^N$  be the open cover of  $X^N$  by product sets  $U_1 \times U_2 \times \cdots \times U_N$ , where  $U_i \in U$ . For a subset  $S_N \subset X^N$ , define  $\aleph(U, N, S_N)$  to be the least cardinality of a subset of  $U^N$  that covers  $S_N$ .

**Remark 2.2.** The continuity of  $T$  and the compactness of  $X$  insure that  $B_N(x, T)$  is compact, and hence that the numbers  $\aleph(U, N, B_N(x, T))$  are all finite and bounded for fixed  $N$  over  $x \in X$ .

**Definition 2.3** (Pointwise preimage entropies [[Nitecki and Przytycki 1999](#)]). For  $T : X \rightarrow X$  be a continuous mapping from a compact space  $X$  to itself, define

$$h_p(T) = \sup_{x \in X} \sup_{U \in O(X)} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \aleph(U, N, B_N(x, T)),$$

$$h_m(T) = \sup_{U \in O(X)} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \sup_{x \in X} \aleph(U, N, B_N(x, T)).$$

**Remark 2.4.** If  $T$  is a homeomorphism, then  $h_p(T) = h_m(T) = 0$ .

As in the case of the topological entropy, we can give metric definitions for these invariants by reinterpreting the numbers  $\aleph(U, N, S_N)$  in terms of  $\varepsilon$ -spanning and

$\varepsilon$ -separated sets. Given a metric space  $(X, d)$  and some  $\varepsilon > 0$ , we say that a subset  $S \subset X$  is  $\varepsilon$ -separated if any two distinct points of  $S$  are at least  $\varepsilon$  apart; and we say that  $A \subset X$  is  $\varepsilon$ -spanned by  $R \subset X$  (or that  $R$   $\varepsilon$ -spans  $A$ ) if any point of  $A$  is at a distance of at most  $\varepsilon$  from  $R$ . Set

$$r(\varepsilon, d, A) = \min\{\text{card } R \mid R \text{ is } \varepsilon\text{-spans } A\},$$

$$s(\varepsilon, d, A) = \max\{\text{card } S \mid S \subset A \text{ is } \varepsilon\text{-separated}\}.$$

**Theorem 2.5** [Nitecki and Przytycki 1999]. *Let  $(X, d)$  be a compact metric space. For any positive integer  $N$ , let  $d^N$  be the metric on  $X^N$  given by*

$$d^N((x_1, \dots, x_N), (y_1, \dots, y_N)) = \max_{1 \leq i \leq N} d(x_i, y_i).$$

*Then for  $T : X \rightarrow X$  continuous, we have*

$$h_p(T) = \sup_{x \in X} \lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log s(\varepsilon, d^N, B_N(x, T)),$$

$$h_m(T) = \lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \sup_{x \in X} s(\varepsilon, d^N, B_N(x, T)).$$

*In both formulas,  $s(\varepsilon, d^N, B_N(x, T))$  can be replaced by  $r(\varepsilon, d^N, B_N(x, T))$ .*

We define a new metric  $d_n$  on  $X$  by setting

$$d_n(x, y) = \max_{0 \leq i \leq n-1} d(T^i(x), T^i(y)).$$

A subset  $F$  of  $X$  is said to  $(n, \varepsilon)$ -span  $K$  if for all  $x \in K$  there exists  $y \in F$  with  $d_n(x, y) \leq \varepsilon$ . Let  $r_n(\varepsilon, K)$  denote the smallest cardinality of any  $(n, \varepsilon)$ -spanning set for  $K$ . A similar definition holds for  $(n, \varepsilon)$  separated sets and  $s_n(\varepsilon, K)$ .

**Definition 2.6** [Walters 1982]. Let  $U$  be a cover of  $X$  and  $Y$  a subset of  $X$ . We denote by  $\aleph(U)|_Y$  the smallest cardinality of a subcover of  $U$  which can cover  $Y$ .

For  $U \in \mathcal{O}(X)$ , we easily see that  $r(\varepsilon, d^N, B_N(x, T)) = r_N(\varepsilon, T^{-N}(x))$ , that  $s(\varepsilon, d^N, B_N(x, T)) = s_N(\varepsilon, T^{-N}(x))$ , and that

$$\aleph(U, N, B_N(x, T)) = \aleph(\bigvee_{n=0}^N T^{-n}U)|_{T^{-N}(x)}.$$

**Remark 2.7.** The pointwise preimage entropies are also given by

$$h_p(T) = \sup_{x \in X} \lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log s_N(\varepsilon, T^{-N}(x))$$

$$= \sup_{x \in X} \lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log r_N(\varepsilon, T^{-N}(x))$$

$$= \sup_{x \in X} \sup_{U \in \mathcal{O}(X)} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \aleph(\bigvee_{n=0}^N T^{-n}U)|_{T^{-N}(x)}$$

and

$$\begin{aligned}
 h_m(T) &= \lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \sup_{x \in X} s_N(\varepsilon, T^{-N}(x)) \\
 &= \lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \sup_{x \in X} r_N(\varepsilon, T^{-N}(x)) \\
 &= \sup_{U \in O(X)} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \sup_{x \in X} \aleph(\bigvee_{n=0}^N T^{-n}U) |_{T^{-N}(x)}.
 \end{aligned}$$

If  $X$  is the circle or a closed interval,  $h_p(T)$  and  $h_m(T)$  always coincide with the topological entropy of  $T$  [Langevin and Przytycki 1992; Langevin and Walczak 1991]. But there exist  $X$  and  $T$  such that  $h_p(T) = 0$  and  $h_m(T) > 0$ ; in fact,  $X$  can be taken to be a zero-dimensional compact metric space [Nitecki and Przytycki 1999].

**Theorem 2.8** [Nitecki and Przytycki 1999]. *If  $T_1 : X \rightarrow X$  and  $T_2 : Y \rightarrow Y$  are topologically conjugate, then  $h_p(T_1) = h_p(T_2)$  and  $h_m(T_1) = h_m(T_2)$ .*

**Theorem 2.9.** *If  $d, d'$  are metrics on a compact set  $X$  defining the same topology, the pointwise preimage entropies with respect to  $d$  and  $d'$  coincide.*

If  $T_2$  is a factor of  $T_1$  then  $h(T_2) \leq h(T_1)$ , where  $h$  is the topological entropy. This inequality need not hold for pointwise preimage entropies, which can increase when we pass to factors. As an example, consider for a map  $f : X \rightarrow X$  the shift  $\sigma_f$  defined on the sequence space  $\Sigma_f = \{ \{x_i\}_{i=0}^\infty \mid f(x_i) = x_{i-1}, i = 1, 2, \dots \}$  by

$$\sigma_f(x_0, x_1, \dots) = (f(x_0), f(x_1), \dots) = (f(x_0), x_0, x_1, \dots).$$

The product topology on  $\Sigma_f \subset X^\mathbb{N}$  makes  $\Sigma_f$  compact and  $\sigma_f$  a homeomorphism. Furthermore, if  $f$  is surjective,  $f$  is a factor of  $\sigma_f$  via the projection  $\varphi(\{x_i\}_{i=0}^\infty) = x_0$ . By Remark 2.4, we have

$$h_p(\sigma_f) = h_m(\sigma_f) = 0.$$

Taking  $f$  such that  $h_p(f) = h_m(f) > 0$  (for instance, the standard expanding map  $z \mapsto 2z \bmod 1$  of the circle  $S^1 = \mathbb{R}/\mathbb{Z}$ , for which  $h_p(f) = h_m(f) = \log 2$ ) and setting  $g = \sigma_f$ , we see that there exist maps  $f : X \rightarrow X, g : Y \rightarrow Y$  with  $f$  a factor of  $g$  and

$$h_m(f) = h_p(f) > h_m(g) = h_p(g).$$

The pointwise preimage entropies are subadditive under Cartesian products and multiplicative under iteration:

**Lemma 2.10** [Nitecki and Przytycki 1999]. *Let  $X, X'$  be compact metric spaces and  $T : X \rightarrow X, T' : X' \rightarrow X'$  continuous maps. Then*

$$\begin{aligned}
 h_p(T \times T') &\leq h_p(T) + h_p(T'), & h_p(T^k) &= kh_p(T), \\
 h_m(T \times T') &\leq h_m(T) + h_m(T'), & h_m(T^k) &= kh_m(T).
 \end{aligned}$$

### 3. Topological forward generator

Let  $X$  be a compact metric space and  $T : X \rightarrow X$  a continuous map. A finite open cover  $\alpha$  of  $X$  is a *forward generator* for  $T$  if for every sequence  $(A_n)_{n=0}^\infty$  of members of  $\alpha$  the set  $\bigcap_{n=0}^\infty T^{-n} \bar{A}_n$  contains at most one point of  $X$ .

**Lemma 3.1.** *Let  $T : X \rightarrow X$  be a continuous map of a compact metric space  $(X, d)$ . Let  $\alpha$  be a forward generator for  $T$ . For any  $\varepsilon > 0$ , there exists  $N > 0$  such that each set in  $\bigvee_{n=0}^N T^{-n} \alpha$  has diameter less than  $\varepsilon$ .*

*Proof.* Suppose the lemma does not hold. Then we can find a positive  $\varepsilon$  such that, for all  $j > 0$ , there exist  $x_j, y_j$  satisfying the conditions  $d(x_j, y_j) > \varepsilon$  and

$$x_j, y_j \in \bigcap_{i=0}^j T^{-i} A_{j,i},$$

for some sequence  $\{A_{j,i}\}_{i=0}^j$  of sets  $A_{j,i} \in \alpha$ . Using the compactness of  $X$ , we can assume (passing to a subsequence if necessary) that  $x_j \rightarrow x$  and  $y_j \rightarrow y$ . We have  $x \neq y$ . Consider the sets  $A_{j,0}$ . Infinitely many of them coincide, since  $\alpha$  is finite. Thus  $x_j, y_j \in A_0$ , say, for infinitely many  $j$ , and hence  $x, y \in \bar{A}_0$ . Similarly, for each  $n$ , infinitely many  $A_{j,n}$  coincide and we obtain  $A_n \in \alpha$  with  $x, y \in T^{-n} \bar{A}_n$ . Thus

$$x, y \in \bigcap_{n=0}^\infty T^{-n} \bar{A}_n,$$

contradicting the assumption that  $\alpha$  is a forward generator.  $\square$

**Definition 3.2** [Nitecki and Przytycki 1999]. A continuous map  $T$  from a compact metric space  $(X, d)$  to itself is said to be *forward expansive* if there exists  $\delta > 0$  such that, for any distinct  $x \neq y \in X$ , the forward images  $T^n x$  and  $T^n y$  are more than  $\delta$  apart, for some  $n$ .

**Lemma 3.3.** *A continuous map  $T$  from a compact metric space  $(X, d)$  to itself is forward expansive if and only if it has a forward generator.*

*Proof.* Suppose  $T$  is forward expansive. Let  $\delta$  be as in the definition and let  $\alpha$  be a finite cover of  $X$  by open balls of radius  $\delta/2$ . Suppose that  $x, y \in \bigcap_{n=0}^\infty T^{-n} \bar{A}_n$ , where  $A_n \in \alpha$ . Then  $d(T^n(x), T^n(y)) \leq \delta$  for all  $n \in \mathbb{N} \cup \{0\}$  so, by assumption  $x = y$ . Then  $\alpha$  is a forward generator.

Conversely, suppose  $\alpha$  is a forward generator. Let  $\delta$  be a Lebesgue number for  $\alpha$ . If  $d(T^n(x), T^n(y)) \leq \delta$  for all  $n \in \mathbb{N} \cup \{0\}$ , then for all  $n \in \mathbb{N}$  exists  $A_n \in \alpha$  with  $T^n(x), T^n(y) \in A_n$  and so,  $x, y \in \bigcap_{n=0}^\infty T^{-n} A_n$ . Since this intersection contains at most one point we have  $x = y$ . Hence  $T$  is forward expansive.  $\square$

**Example 3.4.** Take  $X = \{1, 2, \dots, m\}^{\mathbb{N}}$  and  $T =$  left shift. Then  $\{[k] : 1 \leq k \leq m\}$  is a forward generator, where  $[k] = \{(kx_1x_2x_3\dots) : x_i \in \{1, 2, \dots, m\}\}$ .

Recall from Definitions 2.1 and 2.6 the notations  $\aleph(U, N, Y)$  and  $\aleph(U|_Y)$ . Also recall that  $O(X)$  is the collection of all open covers of  $X$ . Take  $U \in O(X)$  and  $x \in X$ , and set

$$h_p(T, U) = \sup_{x \in X} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \aleph(\bigvee_{n=0}^N T^{-n} U)|_{T^{-N}(x)},$$

$$h_m(T, U) = \limsup_{N \rightarrow \infty} \frac{1}{N} \log \sup_{x \in X} \aleph(\bigvee_{n=0}^N T^{-n} U)|_{T^{-N}(x)}.$$

**Theorem 3.5.** *Let  $T : X \rightarrow X$  be a forward expansive continuous map of the compact metric space  $(X, d)$ . If  $\alpha$  is a forward generator for  $T$ , then*

$$h_p(T) = h_p(T, \alpha) \quad \text{and} \quad h_m(T) = h_m(T, \alpha).$$

*Proof.* Since  $\alpha$  is a forward generator, for any  $U \in O(X)$ , we can choose  $N$  large enough such that  $\bigvee_{n=0}^N T^{-n} \alpha$  is a refinement of  $U$ . This implies that

$$\log \aleph(\bigvee_{n=0}^k T^{-n} U)|_{T^{-k}(x)} \leq \log \aleph(\bigvee_{n=0}^k T^{-n} \bigvee_{n=0}^N T^{-n} \alpha)|_{T^{-k}(x)} \quad \text{for any } k.$$

Then

$$\begin{aligned} \limsup_{k \rightarrow \infty} \frac{1}{k} \log \aleph(\bigvee_{n=0}^k T^{-n} U)|_{T^{-k}(x)} &\leq \limsup_{k \rightarrow \infty} \frac{1}{k} \log \aleph(\bigvee_{n=0}^k T^{-n} \bigvee_{n=0}^N T^{-n} \alpha)|_{T^{-k}(x)} \\ &= \limsup_{k \rightarrow \infty} \frac{1}{k} \log \aleph(\bigvee_{n=0}^{k+N} T^{-n} \alpha)|_{T^{-k}(x)} \\ &\leq \limsup_{k \rightarrow \infty} \frac{1}{k} \log \aleph(\bigvee_{n=0}^{k+N} T^{-n} \alpha)|_{T^{-(k+N)}(x)} \\ &= \limsup_{k \rightarrow \infty} \frac{k+N}{k} \frac{1}{k+N} \log \aleph(\bigvee_{n=0}^{k+N} T^{-n} \alpha)|_{T^{-(k+N)}(x)} \\ &\leq \limsup_{k \rightarrow \infty} \frac{k+N}{k} \limsup_{k \rightarrow \infty} \frac{1}{k+N} \log \aleph(\bigvee_{n=0}^{k+N} T^{-n} \alpha)|_{T^{-(k+N)}(x)} \\ &= \limsup_{k \rightarrow \infty} \frac{1}{k+N} \log \aleph(\bigvee_{n=0}^{k+N} T^{-n} \alpha)|_{T^{-(k+N)}(x)}. \end{aligned}$$

Let

$$h_p(T, U, x) = \limsup_{k \rightarrow \infty} \frac{1}{k} \log \aleph(\bigvee_{n=0}^k T^{-n} U)|_{T^{-k}(x)}.$$

Then  $h_p(T, U, x) \leq h_p(T, \alpha, x)$  for all open covers  $U$  and any fixed  $x$  in  $X$ , which implies

$$h_p(T) = \sup_{x \in X} \sup_{U \in O(X)} h_p(T, U, x) \leq h_p(T, \alpha) = \sup_{x \in X} h_p(T, \alpha, x).$$

Then  $h_p(T) = h_p(T, \alpha)$ .

Similarly, since  $U$  is refined by  $\bigvee_{n=0}^N T^{-n}\alpha$ , we have

$$\mathfrak{K}(\bigvee_{n=0}^k U)|_{T^{-k}(x)} \leq \mathfrak{K}(\bigvee_{n=0}^k \bigvee_{n=0}^N T^{-n}\alpha)|_{T^{-k}(x)}.$$

Thus

$$\begin{aligned} h_m(T, U) &= \limsup_{k \rightarrow \infty} \frac{1}{k} \log \sup_{x \in X} \mathfrak{K}(\bigvee_{n=0}^k T^{-n}U)|_{T^{-k}(x)} \\ &\leq \limsup_{k \rightarrow \infty} \frac{1}{k} \log \sup_{x \in X} \mathfrak{K}(\bigvee_{n=0}^k T^{-n} \bigvee_{n=0}^N T^{-n}\alpha)|_{T^{-k}(x)}. \end{aligned}$$

A similar calculation yields  $h_m(T, U) \leq h_m(T, \alpha)$  for all open covers  $U$ . Thus  $h_m(T) = h_m(T, \alpha)$ .  $\square$

#### 4. Measure-theoretic forward generator

We continue to consider a continuous self-map  $T$  of a compact metric space  $(X, d)$ . Given a subset  $K \subset X$ , a  $\delta > 0$ , and a positive integer  $n$ , we set

$$r(n, \delta, K) = r(n, \delta, K, T) = \max\{\text{card } E : E \subseteq K \text{ is } (n, \delta)\text{-separated}\}.$$

**Definition 4.1.** The *upper preimage entropy* of a continuous self-map  $T : X \rightarrow X$  of a compact metric space  $X$  is the number

$$\begin{aligned} h_{\text{top}}(T | \xi^-) &= \lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \sup_{k \geq 0, x \in X} r(n, \delta, T^{-k}x) \\ &= \sup_{\alpha \in \mathcal{O}(X)} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \sup_{\substack{k \geq 0 \\ x \in X}} \mathfrak{K}(\bigvee_{i=0}^{n-1} T^{-i}\alpha)|_{T^{-k}(x)}. \end{aligned}$$

One can check that  $h_p(T) \leq h_m(T) \leq h_{\text{top}}(T | \xi^-) \leq h(T)$ .

**Example 4.2.** Consider  $S : \{1, 2\}^{\mathbb{N}} \rightarrow \{1, 2\}^{\mathbb{N}}$  and  $T : \{1, 2\}^{\mathbb{Z}} \rightarrow \{1, 2\}^{\mathbb{Z}}$ , where  $S$  and  $T$  are left shifts. We know that  $h_p(S) = h_m(S) = h_{\text{top}}(S) = \log 2$ , that  $h_p(T) = h_m(T) = h_{\text{top}}(T) = 0$ , and that the product rule holds, i.e.,

$$h_{\text{top}}(S \times T | \xi^-) = h_{\text{top}}(S | \xi^-) + h_{\text{top}}(T | \xi^-)$$

(see [Cheng and Newhouse 2005]) and

$$h(S \times T) = h(S) + h(T).$$

It follows that

$$h_p(S \times T) = h_m(S \times T) = h_{\text{top}}(S \times T | \xi^-) = \log 2 < h(S \times T) = 2 \log 2.$$



Now we introduce conditional entropy. Let  $\zeta = \{A_1, A_2, \dots\}$  be a countable partition of  $X$  into measurable sets. For each  $x \in X$ , denote by  $\zeta(x)$  the element of  $\zeta$  to which  $x$  belongs. The information function associated to  $\zeta$  is defined to be

$$I_\zeta(x) = -\log m(\zeta(x)) = -\sum_{A \in \zeta} \log m(A) \chi_A(x),$$

so that  $I_\zeta(x)$  takes the constant value  $-\log m(A)$  on the cell  $A$  of  $\zeta$ . Clearly

$$H(\zeta) = \int_X I_\zeta(x) dm(x).$$

It is useful to consider conditional information and entropy, which take into account information that may already be in hand. Let  $\mathcal{F}$  be a sub- $\sigma$ -algebra of  $\mathcal{B}$ . Recall that for  $\phi \in L^1(X)$ , the conditional expectation  $E(\phi | \mathcal{F})$  of  $\phi$  given  $\mathcal{F}$  is an  $\mathcal{F}$ -measurable function on  $X$  satisfying

$$\int_F E(\phi | \mathcal{F}) dm = \int_F \phi dm$$

for all  $F \in \mathcal{F}$ ; the name comes from the fact that  $E(\phi | \mathcal{F})(x)$  represents our expected value for  $\phi$  if we are given the foreknowledge  $\mathcal{F}$ . Thus we let  $m(A | \mathcal{F}) = E(\chi_A | \mathcal{F})$  and define the conditional information function of a countable partition  $\zeta$  given a  $\sigma$ -algebra  $\mathcal{F} \subset \mathcal{B}$  to be

$$I_{\zeta|\mathcal{F}}(x) = -\sum_{A \in \zeta} \log m(A | \mathcal{F}) \chi_A(x).$$

The conditional entropy of  $\zeta$  given  $\mathcal{F}$  is defined by

$$H(\zeta | \mathcal{F}) = \int_X I_{\zeta|\mathcal{F}}(x) dm.$$

Next, let  $\xi$  denote the point partition of  $X$ , we also identify with the  $\sigma$ -algebra  $\mathcal{B}$  of Borel measurable sets. For  $n > 0$ , set

$$\xi^{-n} = T^{-n}\xi.$$

Given a finite partition  $\alpha$ , let  $\alpha^n = \bigvee_{i=0}^{n-1} T^{-i}\alpha$ . For a  $T$ -invariant probability  $\mu$ , let

$$H_\mu(\alpha^n | \xi^{-k})$$

denote the conditional entropy of  $\alpha^n$  given the  $\sigma$ -algebra  $T^{-k}\mathcal{B}$ . We call this the *entropy of  $\alpha^n$  given the preimage partition  $\xi^{-k}$* .

Since  $H_\mu(\cdot | \cdot)$  is increasing in the first variable and decreasing in the second, the inequalities  $n \geq m, l \geq k$  imply

$$H_\mu(\alpha^n | \xi^{-l}) \geq H_\mu(\alpha^m | \xi^{-k}).$$

Set

$$H_\mu(\alpha^n | \xi^-) = H_\mu(\alpha^n | \xi^{-\infty}) = \sup_{k \geq 0} H_\mu(\alpha^n | \xi^{-k}) = \lim_{k \rightarrow \infty} H_\mu(\alpha^n | \xi^{-k}).$$

We can also define

$$\xi^- = \bigcap_{k=1}^{\infty} \xi^{-k},$$

and we call this eventual range  $\xi^-$  the infinity past  $\sigma$ -algebra.

**Lemma 4.3** [Bowen 1972]. *The quantity  $a_n = H_\mu(\alpha^n | \xi^-)$  is subadditive.*

**Definition 4.4.** For any finite partition  $\alpha$ , the *entropy of  $\alpha$  given  $\xi^-$*  is the number

$$h_\mu(T | \xi^-, \alpha) = h_\mu(\alpha | \xi^-) = \lim_{n \rightarrow \infty} \frac{1}{n} H_\mu(\alpha^n | \xi^-) = \inf_{n \rightarrow \infty} \frac{1}{n} H_\mu(\alpha^n | \xi^-),$$

and we define the *metric preimage entropy of  $T$  given  $\xi^-$  with respect to  $\mu$*  to be

$$h_\mu(T | \xi^-) = \sup_{\alpha} h_\mu(\alpha | \xi^-) = \sup_{\alpha} h_\mu(T | \xi^-, \alpha).$$

**Lemma 4.5.** *The metric preimage entropy  $h_\mu(T | \xi^-)$  is a measure-theoretic conjugacy invariant. The upper preimage entropy  $h_{\text{top}}(T | \xi^-)$  is a topological conjugacy invariant.*

**Theorem 4.6** (Variational principle [Cheng and Newhouse 2005]). *Let  $T : X \rightarrow X$  be a continuous map of a compact metric space  $X$ . Then*

$$h_{\text{top}}(T | \xi^-) = \sup_{\mu} h_\mu(T | \xi^-),$$

where  $\mu$  runs over all  $T$ -invariant Borel probability measures on  $X$ .

**Theorem 4.7.** *Let  $T : X \rightarrow X$  be a forward expansive continuous function of a compact metric space  $(X, d)$ . If  $\alpha$  is a forward generator for  $T$ , then*

$$h_{\text{top}}(T | \xi^-) = h_{\text{top}}(T | \xi^-, \alpha).$$

The proof is similar to that of [Theorem 3.5](#).

**Lemma 4.8.** *Let  $\zeta$  and  $\eta$  be two finite partitions of  $X$ . Then*

$$h_\mu(\zeta | \xi^-) \leq h_\mu(\eta | \xi^-) + H_\mu(\zeta | \eta).$$

*Proof.* We have

$$\begin{aligned} H_\mu(\bigvee_{i=0}^{n-1} T^{-i} \zeta | \xi^{-k}) &\leq H_\mu((\bigvee_{i=0}^{n-1} T^{-i} \zeta \vee \bigvee_{i=0}^{n-1} T^{-i} \eta) | \xi^{-k}) \\ &= H_\mu(\bigvee_{i=0}^{n-1} T^{-i} \eta | \xi^{-k}) + H_\mu(\bigvee_{i=0}^{n-1} T^{-i} \zeta | \bigvee_{i=0}^{n-1} T^{-i} \eta \vee \xi^{-k}) \\ &\leq H_\mu(\bigvee_{i=0}^{n-1} T^{-i} \eta | \xi^{-k}) + H_\mu(\bigvee_{i=0}^{n-1} T^{-i} \zeta | \bigvee_{i=0}^{n-1} T^{-i} \eta). \end{aligned}$$

Let  $k \rightarrow \infty$ ; since  $H_\mu(\bigvee_{i=0}^{n-1} T^{-i} \zeta \mid \bigvee_{i=0}^{n-1} T^{-i} \eta) \leq n \cdot H_\mu(\zeta \mid \eta)$ , this implies that  $H_\mu(\zeta_0^n \mid \xi^-) \leq H_\mu(\eta_0^n \mid \xi^-) + n \cdot H_\mu(\zeta \mid \eta)$ . Divide by  $n$  and let  $n$  go to infinity; then  $h_\mu(\zeta \mid \xi^-) \leq h_\mu(\eta \mid \xi^-) + H_\mu(\zeta \mid \eta)$ .  $\square$

**Lemma 4.9.** *For any fixed  $k$ ,*

$$h_\mu(T \mid \xi^-, \alpha) = h_\mu(T \mid \xi^-, \bigvee_{i=0}^k T^{-i} \alpha).$$

$$\begin{aligned} \text{Proof. } h_\mu(T \mid \xi^-, \bigvee_{i=0}^k T^{-i} \alpha) &= \lim_{n \rightarrow \infty} \frac{1}{n} H_\mu(\bigvee_{i=0}^{n-1} T^{-i} (\bigvee_{i=0}^k T^{-i} \alpha) \mid \xi^-) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} H_\mu(\bigvee_{i=0}^{k+n-1} T^{-i} \alpha \mid \xi^-) \\ &= \lim_{n \rightarrow \infty} \frac{k+n-1}{n} \frac{1}{k+n-1} H_\mu(\bigvee_{i=0}^{k+n-1} T^{-i} \alpha \mid \xi^-) \\ &= h_\mu(T \mid \xi^-, \alpha). \end{aligned} \quad \square$$

**Lemma 4.10.** *If  $\{A_n\}$  is an increasing sequence of finite partitions of  $X$  and  $C$  is a partition with  $C \leq \bigvee_{i=0}^\infty A_i$ , then  $H_\mu(C \mid A_n) \rightarrow 0$  as  $n \rightarrow \infty$ .*

Let  $C = \{C_i : i = 1, 2, \dots, n\}$  be a finite sub- $\sigma$ -algebra of  $\mathcal{B}$ . The nonempty sets of the form  $B_1 \cap B_2 \cdots \cap B_n$ , where  $B_i = C_i$  or  $X \setminus C_i$ , form a finite partition of  $X$ . We denote it by  $\alpha(C)$  and we define  $h_\mu(T \mid \xi^-, C) = h_\mu(T \mid \xi^-, \alpha(C))$ .

As in the case of measure-theoretic entropy, the main method for calculating  $h_\mu(T \mid \xi^-)$  is supplied by the next theorem.

**Theorem 4.11** (Kolmogorov–Sinai forward generator). *Let  $T$  be a measure-preserving transformation of  $(X, \mathcal{B}, \mu)$  and  $\mathcal{R}$  a finite sub- $\sigma$ -algebra such that*

$$\bigvee_{n=0}^\infty T^{-n}(\alpha(\mathcal{R})) = \mathcal{B}.$$

*Then*

$$h_\mu(T \mid \xi^-) = h_\mu(T \mid \xi^-, \mathcal{R}).$$

*Proof.* Let  $C$  be any partition. We show that  $h_\mu(T \mid \xi^-, C) \leq h_\mu(T \mid \xi^-, \alpha(\mathcal{R}))$ . For  $n \geq 1$ , by Lemmas 4.8 and 4.9,

$$\begin{aligned} h_\mu(T \mid \xi^-, C) &\leq h_\mu(T \mid \xi^-, \bigvee_{i=0}^n T^{-i} \alpha(\mathcal{R})) + H_\mu(C \mid \bigvee_{i=0}^n T^{-i} \alpha(\mathcal{R})) \\ &= h_\mu(T \mid \xi^-, \alpha(\mathcal{R})) + H_\mu(C \mid \bigvee_{i=0}^n T^{-i} \alpha(\mathcal{R})). \end{aligned}$$

Let  $A_n = \bigvee_{i=0}^n T^{-i} \alpha(\mathcal{R})$  be as in Lemma 4.10. Then  $H_\mu(C \mid A_n) \rightarrow 0$  as  $n \rightarrow \infty$ . This implies  $h_\mu(T \mid \xi^-, C) \leq h_\mu(T \mid \xi^-, \alpha(\mathcal{R}))$ . Therefore

$$h_\mu(T \mid \xi^-, C) \leq h_\mu(T \mid \xi^-, \mathcal{R}). \quad \square$$

We end this section with some propositions about  $h_\mu(T \mid \xi^-)$  and from those results we conclude that  $h_\mu(T \mid \xi^-) \leq h_\mu(T)$ , where  $h_\mu(T)$  is the measure-theoretic entropy.

**Lemma 4.12.** *We let  $\mathcal{B}_\infty = \bigvee_{n=1}^\infty \mathcal{B}_n$  if  $\{\mathcal{B}_n\}$  is an increasing sequence of sub- $\sigma$ -algebras of  $X$  and let  $\mathcal{B}_\infty = \bigcap \mathcal{B}_n$  if  $\{\mathcal{B}_n\}$  is a decreasing sequence. If  $\alpha$  is a finite partition, then*

$$\lim_{n \rightarrow \infty} H_\mu(\alpha \mid \mathcal{B}_n) = H_\mu(\alpha \mid \mathcal{B}_\infty).$$

*Proof.* We show the decreasing case; the discussion for the increasing sequence is similar — see [Petersen 1983, Proposition 5.2.11].

Take  $A \in \alpha$ . Because  $E(E(\chi_A \mid \mathcal{B}_{n-1}) \mid \mathcal{B}_n) = E(\chi_A \mid \mathcal{B}_n)$ , by the reverse martingale theorem and [Billingsley 1995, Theorem 35.9], we have

$$\lim_{n \rightarrow \infty} E(\chi_A \mid \mathcal{B}_n) = E(\chi_A \mid \mathcal{B}_\infty).$$

Also  $I_{\alpha \mid \mathcal{B}_n} = -\sum_{A \in \alpha} \log E(\chi_A \mid \mathcal{B}_n) \cdot E(\chi_A \mid \mathcal{B}_n)$  is a bounded continuous function; thus, by the bounded convergence theorem, we get

$$\lim_{n \rightarrow \infty} H(\alpha \mid \mathcal{B}_n) = \lim_{n \rightarrow \infty} \int I_{\alpha \mid \mathcal{B}_n} d\mu = \int \lim_{n \rightarrow \infty} I_{\alpha \mid \mathcal{B}_n} d\mu = H(\alpha \mid \mathcal{B}_\infty). \quad \square$$

**Lemma 4.13.** *Let  $\alpha$  be a finite partition. Then*

$$\begin{aligned} h_\mu(\alpha \mid \xi^-) &= h_\mu(T \mid \xi^-, \alpha) = \lim_{n \rightarrow \infty} \lim_{k \rightarrow \infty} H_\mu(\alpha \mid \bigvee_{l=1}^{n-1} T^{-l} \alpha \vee T^{-k}(\xi)) \\ &= \lim_{n \rightarrow \infty} H_\mu(\alpha \mid \lim_{k \rightarrow \infty} \bigvee_{l=1}^{n-1} T^{-l} \alpha \vee T^{-k}(\xi)) \\ &= \lim_{n \rightarrow \infty} H_\mu(\alpha \mid \bigvee_{l=1}^{n-1} T^{-l} \alpha \vee \xi^-). \end{aligned}$$

*Proof.* We have

$$\begin{aligned} &\lim_{k \rightarrow \infty} H_\mu(\alpha \vee T^{-1} \alpha \vee \dots \vee T^{-(j-1)} \alpha \mid T^{-k}(\xi)) \\ &= \lim_{k \rightarrow \infty} H_\mu(\alpha \mid \bigvee_{l=1}^{j-1} T^{-l} \alpha \vee T^{-k}(\xi)) + \lim_{k \rightarrow \infty} H_\mu(\bigvee_{l=1}^{j-1} T^{-l} \alpha \mid T^{-k}(\xi)), \end{aligned}$$

which implies

$$\begin{aligned} &\lim_{k \rightarrow \infty} H_\mu(\alpha \mid \bigvee_{l=1}^{j-1} T^{-l} \alpha \vee T^{-k}(\xi)) \\ &= \lim_{k \rightarrow \infty} H_\mu(\alpha \vee T^{-1} \alpha \vee \dots \vee T^{-(j-1)} \alpha \mid T^{-k}(\xi)) - \lim_{k \rightarrow \infty} H_\mu(\bigvee_{l=1}^{j-1} T^{-l} \alpha \mid T^{-k}(\xi)) \\ &= \lim_{k \rightarrow \infty} H_\mu(\bigvee_{l=0}^{j-1} T^{-l} \alpha \mid T^{-k}(\xi)) - \lim_{k \rightarrow \infty} H_\mu(\bigvee_{l=0}^{j-2} T^{-l} \alpha \mid T^{-(k-1)}(\xi)). \end{aligned}$$

We thus get

$$\begin{aligned} &\sum_{j=2}^n \lim_{k \rightarrow \infty} H_\mu(\alpha \mid \bigvee_{l=1}^{j-1} T^{-l} \alpha \vee T^{-k}(\xi)) \\ &= \lim_{k \rightarrow \infty} H_\mu(\bigvee_{l=0}^{n-1} T^{-l} \alpha \mid T^{-k}(\xi)) - \lim_{k \rightarrow \infty} H_\mu(\alpha \mid T^{-k}(\xi)) \end{aligned}$$

By Cesàro's Theorem and [Lemma 4.12](#),

$$\begin{aligned}
 h_\mu(\alpha \mid \xi^-) &= \lim_{n \rightarrow \infty} \lim_{k \rightarrow \infty} H_\mu(\alpha \mid \bigvee_{l=1}^{n-1} T^{-l} \alpha \vee T^{-k}(\xi)) \\
 &= \lim_{n \rightarrow \infty} H_\mu(\alpha \mid \lim_{k \rightarrow \infty} \bigvee_{l=1}^{n-1} T^{-l} \alpha \vee T^{-k}(\xi)) \\
 &= \lim_{n \rightarrow \infty} H_\mu(\alpha \mid \bigvee_{l=1}^{n-1} T^{-l} \alpha \vee \xi^-). \quad \square
 \end{aligned}$$

**Lemma 4.14.**  $h_\mu(T \mid \xi^-) \leq h_\mu(T)$ .

*Proof.* For any finite partition  $\alpha$ ,  $H_\mu(\alpha \mid \bigvee_{i=1}^{n-1} T^{-i} \alpha \vee \xi^-) \leq H_\mu(\alpha \mid \bigvee_{i=1}^{n-1} T^{-i} \alpha)$  and  $h_\mu(T, \alpha) = \lim_{n \rightarrow \infty} H_\mu(\alpha \mid \bigvee_{i=1}^{n-1} T^{-i} \alpha)$ . By [Lemma 4.13](#),  $h_\mu(T \mid \xi^-) \leq h_\mu(T)$ .  $\square$

## References

- [Billingsley 1995] P. Billingsley, *Probability and measure*, 3rd ed., John Wiley & Sons, New York, 1995. [MR 95k:60001](#) [Zbl 0822.60002](#)
- [Bowen 1972] R. Bowen, “Entropy-expansive maps”, *Trans. Amer. Math. Soc.* **164** (1972), 323–331. [MR 44 #2907](#) [Zbl 0229.28011](#)
- [Cheng and Newhouse 2005] W.-C. Cheng and S. E. Newhouse, “Pre-image entropy”, *Ergodic Theory Dynam. Systems* **25**:4 (2005), 1091–1113. [MR 2158398](#)
- [Hurley 1995] M. Hurley, “On topological entropy of maps”, *Ergodic Theory Dynam. Systems* **15**:3 (1995), 557–568. [MR 96h:58097](#) [Zbl 0833.54021](#)
- [Langevin and Przytycki 1992] R. Langevin and F. Przytycki, “Entropie de l’image inverse d’une application”, *Bull. Soc. Math. France* **120**:2 (1992), 237–250. [MR 94b:58085](#) [Zbl 0757.58023](#)
- [Langevin and Walczak 1991] R. Langevin and P. Walczak, “Entropie d’une dynamique”, *C. R. Acad. Sci. Paris Sér. I Math.* **312**:1 (1991), 141–144. [MR 92e:58159](#) [Zbl 0723.54020](#)
- [Nitecki and Przytycki 1999] Z. Nitecki and F. Przytycki, “Preimage entropy for mappings”, *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* **9**:9 (1999), 1815–1843. [MR 2000j:37025](#)
- [Petersen 1983] K. Petersen, *Ergodic theory*, Cambridge Studies in Advanced Mathematics **2**, Cambridge University Press, Cambridge, 1983. [MR 87i:28002](#) [Zbl 0507.28010](#)
- [Walters 1982] P. Walters, *An introduction to ergodic theory*, Graduate Texts in Mathematics **79**, Springer, New York, 1982. [MR 84e:28017](#) [Zbl 0475.28009](#)

Received April 25, 2004. Revised February 2, 2005.

WEN-CHIAO CHENG  
 DEPARTMENT OF MATHEMATICS  
 NATIONAL CHUNG CHENG UNIVERSITY  
 MIN-HSIUNG, CHIA-YI  
 TAIWAN, 621  
[wccheng@math.ccu.edu.tw](mailto:wccheng@math.ccu.edu.tw)