

*Pacific
Journal of
Mathematics*

**SCHWARZIAN DERIVATIVES
AND A LINEARLY INVARIANT FAMILY IN \mathbb{C}^n**

RODRIGO HERNÁNDEZ R.

SCHWARZIAN DERIVATIVES AND A LINEARLY INVARIANT FAMILY IN \mathbb{C}^n

RODRIGO HERNÁNDEZ R.

We use Oda's definition of the Schwarzian derivative for locally univalent holomorphic maps F in several complex variables to define a Schwarzian derivative operator $\mathcal{S}F$. We use the Bergman metric to define a norm $\|\mathcal{S}F\|$ for this operator, which in the ball is invariant under composition with automorphisms. We study the linearly invariant family

$$\mathcal{F}_\alpha = \{F : \mathbb{B}^n \rightarrow \mathbb{C}^n \mid F(\mathbf{0}) = \mathbf{0}, DF(\mathbf{0}) = \text{Id}, \|\mathcal{S}F\| \leq \alpha\},$$

estimating its order and norm order.

1. Introduction

The link between the Schwarzian derivative of a locally univalent holomorphic map in one complex variable, given by

$$Sf = \left(\frac{f''}{f'}\right)' - \frac{1}{2} \left(\frac{f''}{f'}\right)^2,$$

with the univalence of f and distortion problems has been studied extensively; see [Chuaqui and Osgood 1993; Epstein 1986; Kraus 1932; Nehari 1949], for example. Sf vanishes identically if and only if f is a Möbius mapping, and we have $S(f \circ g) = (Sf \circ g)(g')^2 + Sg$. An analytic function f with Schwarzian derivative $Sf = 2p$ has the form $f = u/v$, where u and v are any linearly independent solutions of the equation $u'' + pu = 0$. If f is defined in the unit disk \mathbb{D} , the norm

$$\|Sf\| = \sup_{|z|=1} (1 - |z|^2)^2 |Sf(z)|$$

is invariant under precomposition with automorphisms of the disk.

Some analogues of the Schwarzian derivative in several complex variables are available, but results relating it to the aforementioned problems of univalence and

MSC2000: primary 32A17, 32W50; secondary 32H02, 30C35.

Keywords: Several complex variables, Schwarzian derivative, Linearly invariant families, Sturm comparison.

distortion are less satisfactory than in one variable [Molzon and Pinney Mortensen 1997]. Consider the overdetermined system of partial differential equations

$$(1-1) \quad \frac{\partial^2 u}{\partial z_i \partial z_j} = \sum_{k=1}^n P_{ij}^k(z) \frac{\partial u}{\partial z_k} + P_{ij}^0(z)u, \quad i, j = 1, 2, \dots, n,$$

where $z = (z_1, z_2, \dots, z_n) \in \mathbb{C}^n$. The system is called *completely integrable* if (1-1) has $n + 1$ linearly independent solutions. The system (1-1) is said to be in *canonical form* (see [Yoshida 1976]) if the coefficients satisfy

$$\sum_{j=1}^n P_{ij}^j(z) = 0, \quad i = 1, 2, \dots, n.$$

T. Oda [1974] defined the Schwarzian derivative S_{ij}^k of a locally injective holomorphic mapping $F(z_1, z_2, \dots, z_n) = (w_1, w_2, \dots, w_n)$ as

$$S_{ij}^k F = \sum_{l=1}^n \frac{\partial^2 w_l}{\partial z_i \partial z_j} \frac{\partial z_k}{\partial w_l} - \frac{1}{n+1} \left(\delta_i^k \frac{\partial}{\partial z_j} + \delta_j^k \frac{\partial}{\partial z_i} \right) \log \Delta,$$

where $i, j, k = 1, 2, \dots, n$, $\Delta = \det(\partial F / \partial z)$, and δ_i^k is the Kronecker symbol. For $n > 1$ these Schwarzian derivatives satisfy

$$S_{ij}^k F = 0 \quad \text{for all } i, j, k = 1, 2, \dots, n$$

if and only if $F(z)$ is a Möbius transformation, that is, if it has the form

$$F(z) = \left(\frac{l_1(z)}{l_0(z)}, \dots, \frac{l_n(z)}{l_0(z)} \right),$$

where $l_i(z) = a_{i0} + a_{i1}z_1 + \dots + a_{in}z_n$ with $\det(a_{ij}) \neq 0$. For a composition we have

$$(1-2) \quad S_{ij}^k(G \circ F)(z) = S_{ij}^k F(z) + \sum_{l,m,r=1}^n S_{lm}^r G(w) \frac{\partial w_l}{\partial z_i} \frac{\partial w_m}{\partial z_j} \frac{\partial z_k}{\partial w_r}, \quad w = F(z).$$

Thus, precomposition with a Möbius transformation G leads to $S_{ij}^k(G \circ F) = S_{ij}^k F$. The coefficients $S_{ij}^0 F$ are given by

$$S_{ij}^0 F(z) = \Delta^{1/(n+1)} \left(\frac{\partial^2}{\partial z_i \partial z_j} \Delta^{-1/(n+1)} - \sum_{k=1}^n \frac{\partial}{\partial z_k} \Delta^{-1/(n+1)} S_{ij}^k F(z) \right).$$

The function $u = \Delta^{-1/n+1}$ is always a solution of (1-1) with $S_{ij}^k F = P_{ij}^k$.

Remark 1.1. For $n = 1$, $S_{11}^1 f = 0$ for all locally injective f , but $S_{11}^0 f = -\frac{1}{2} S f$.

Proposition 1.2 [Yoshida 1976]. *Let (1-1) be a completely integrable system in canonical form and consider a set $u_0(z), u_1(z), \dots, u_n(z)$ of linearly independent solutions. Then*

$$P_{ij}^k(z) = S_{ij}^k F(z), \quad i, j, k = 1, 2, \dots, n,$$

where $F(z) = (w_1(z), \dots, w_n(z))$ and $w_i(z) = u_i(z)/u_0(z)$.

Remark 1.3. In contrast to the one-dimensional case, when $n > 1$ the Schwarzian derivatives $S_{ij}^k F$ are differential operators of order 2. One way to understand this phenomenon is through a dimensional argument: For $n = 1$ the Möbius group has dimension 3, which allows one to choose $f(z_0), f'(z_0)$ and $f''(z_0)$ for a holomorphic mapping f at a given point z_0 arbitrarily. It would therefore be pointless to seek a Möbius-invariant differential operator of order 2. But for $n > 1$ the number of parameters involved in the value and all derivatives of order 1 and 2 of a locally biholomorphic mapping is $n^2(n + 1)/2 + n^2 + n$, which exceeds the dimension $n^2 + 2n$ of the corresponding Möbius group in \mathbb{C}^n . Moreover, since $S_{ij}^k F = S_{ji}^k F$ for all k and

$$\sum_{j=1}^n S_{ij}^j F = 0,$$

there are exactly $n(n - 1)(n + 2)/2$ independent terms $S_{ij}^k F$, which is equal to the excess mentioned above.

In this paper we employ the Oda Schwarzian derivatives S_{ij}^k to propose a Schwarzian derivative operator $\mathcal{S}F$. Using the Bergman metric, we will define a norm for $\mathcal{S}F$, which for mappings defined in the ball \mathbb{B} turns out to be invariant under the group of automorphisms. We then focus on the study of geometric properties of the linearly invariant family given by bounded Schwarzian norm. We will appeal to the relationship with the completely integrable system (1-1) and Sturm comparison techniques adapted to this special situation.

2. The Schwarzian derivative operator

For $\Omega \subset \mathbb{C}^n$ open, let $F : \Omega \rightarrow \mathbb{C}^n, F(z_1, \dots, z_n) = (w_1, \dots, w_n)$, be a locally univalent holomorphic mapping, and set $\Delta = \det(\partial F / \partial z)$. For $k = 1, \dots, n$, define an $n \times n$ matrix

$$\mathbb{S}^k F = (S_{ij}^k F), \quad i, j = 1, \dots, n.$$

Proposition 2.1. *Let F be a locally injective holomorphic mapping and let $w = G(z)$ be a Möbius transformation. Then*

$$\mathbb{S}^k (F \circ G) = \sum_{r=1}^n \frac{\partial z_k}{\partial w_r} DG^t ((S^r F) \circ G) DG \quad \text{for } k = 1, \dots, n.$$

Proof. From (1-2) and the Möbius property of G we have

$$\begin{aligned} S_{ij}^k(F \circ G)(z) &= \sum_{l,m,r=1}^n S_{lm}^r F(w) \frac{\partial w_l}{\partial z_i} \frac{\partial w_m}{\partial z_j} \frac{\partial z_k}{\partial w_r} + S_{ij}^k G(z) \\ &= \sum_{r=1}^n \frac{\partial z_k}{\partial w_r} \sum_{m,l=1}^n \frac{\partial w_l}{\partial z_i} S_{lm}^r F(w) \frac{\partial w_m}{\partial z_j} + S_{ij}^k G(z) \\ &= \sum_{r=1}^n \frac{\partial z_k}{\partial w_r} \sum_{m,l=1}^n \frac{\partial w_l}{\partial z_i} S_{lm}^r F(w) \frac{\partial w_m}{\partial z_j}. \end{aligned}$$

The proposition follows after rewriting this in terms of matrices. □

Definition 2.2. The Schwarzian derivative operator is the operator $\mathcal{S}F(z) : T_z\Omega \rightarrow T_{F(z)}\Omega$ given by

$$\mathcal{S}F(z)(\vec{v}) = (\vec{v}^t \mathbb{S}^1 F(z) \vec{v}, \vec{v}^t \mathbb{S}^2 F(z) \vec{v}, \dots, \vec{v}^t \mathbb{S}^n F(z) \vec{v}),$$

where $\vec{v} \in T_z\Omega$.

Recall that the Bergman metric on \mathbb{B}^n is the hermitian product defined by

$$(2-1) \quad g_{ij}(z) = \frac{n+1}{(1-|z|^2)^2} ((1-|z|^2)\delta_{ij} + \bar{z}_i z_j).$$

Any automorphism of the ball is an isometry of the Bergman metric.

We define the norm of the Schwarzian derivative operator by

$$\|\mathcal{S}F(z)\| = \sup_{\|\vec{v}\|=1} \|\mathcal{S}F(z)(\vec{v})\|,$$

where $\|\vec{v}\| = (\sum g_{ij} v_i \bar{v}_j)^{1/2}$ is the Bergman norm of $\vec{v} \in T_z\mathbb{B}^n$.

A routine calculation using the fact that $u_0 = \Delta^{-1/n+1}$ is a solution of (1-1) with $P_{ij}^k = S_{ij}^k F$ allows one to rewrite the Schwarzian derivative operator as

$$\mathcal{S}F(z)(\vec{v}, \vec{v}) = (DF(z))^{-1} D^2 F(z)(\vec{v}, \vec{v}) - \frac{2}{n+1} \left(\frac{1}{\Delta} \sum_{j=1}^n \Delta_j(z) v_j \right) \vec{v},$$

or yet

$$(2-2) \quad \mathcal{S}F(z)(\vec{v}, \vec{v}) = (DF(z))^{-1} D^2 F(z)(\vec{v}, \vec{v}) + 2\Delta^{1/n+1} (\nabla u_0 \cdot \vec{v}) \vec{v},$$

where $\Delta_j = \sum_{k=1}^n (-1)^{j-1} \delta_{jk}$ and δ_{jk} is the determinant of $DF(z)$ with the k -th column replaced by the column

$$\left(\frac{\partial^2 f_1}{\partial z_j \partial z_k}, \dots, \frac{\partial^2 f_n}{\partial z_j \partial z_k} \right)^{\text{tr}}(z).$$

The operator $(DF(z))^{-1}D^2F(z)(z, \cdot)$ was considered by Pfaltzgraff [1974] in his generalization of the Becker criterion.

Theorem 2.3. *Let $F : B^n \rightarrow \mathbb{C}^n$ be a locally injective holomorphic mapping and let σ be an automorphism of \mathbb{B}^n . Then*

$$\|\mathcal{S}(F \circ \sigma)(z)\| = \|\mathcal{S}F(\sigma(z))\|.$$

Proof. We know that

$$\begin{aligned} \mathbb{S}^k(F \circ \sigma) &= \sum_{l=1}^n \frac{\partial z_k}{\partial w_l} (D\sigma)^t \mathbb{S}^l F \circ \sigma (D\sigma) \\ &= \left(\frac{\partial z_k}{\partial w_1}, \dots, \frac{\partial z_k}{\partial w_n} \right) \begin{pmatrix} (D\sigma)^t \mathbb{S}^1 F \circ \sigma (D\sigma) \\ \vdots \\ (D\sigma)^t \mathbb{S}^n F \circ \sigma (D\sigma) \end{pmatrix}. \end{aligned}$$

Hence

$$(\mathcal{S}F \circ \sigma)(z)(\vec{v}) = D\sigma^{-1} \begin{pmatrix} \vec{v}^t (D\sigma)^t \mathbb{S}^1 F(\sigma(z))(D\sigma) \vec{v} \\ \vdots \\ \vec{v}^t (D\sigma)^t \mathbb{S}^n F(\sigma(z))(D\sigma) \vec{v} \end{pmatrix} = D\sigma^{-1} \begin{pmatrix} \vec{u}^t \mathbb{S}^1 F(\sigma(z)) \vec{u} \\ \vdots \\ \vec{u}^t \mathbb{S}^n F(\sigma(z)) \vec{u} \end{pmatrix},$$

where $\vec{u} = D\sigma(z)(\vec{v})$. Then

$$\|\mathcal{S}(F \circ \sigma)(z)(\vec{v})\| = \|DG\sigma^{-1}\mathcal{S}F(\sigma(z))(\vec{u})\| = \|\mathcal{S}F(\sigma(z))(\vec{u})\|,$$

and since σ is a isometry in the Bergman metric, the theorem follows after taking supremum over all unit vectors \vec{v} . □

The definition of norm for the Schwarzian operator can be given using any hermitian metric or even a Finsler metric. Since in ball the Bergman metric coincides up to constant multiples with the Kobayashi or the Carathéodory metric, the resulting norm for $\mathcal{S}F$ is the same. This will certainly not be the case on arbitrary domains. [Theorem 2.3](#) will also fail on arbitrary domains because it requires the automorphisms to be Möbius.

3. The family \mathcal{F}_α

Definition 3.1. Consider the family

$$LS = \{F : B^n \rightarrow \mathbb{C}^n \mid F(0) = 0, DF(0) = \text{Id}\}$$

of normalized locally biholomorphic mappings on the ball \mathbb{B}^n , and the Koebe transformations $\Lambda_\sigma(F)$ of the ball, given by

$$\Lambda_\sigma(F)(z) = (D\sigma(0))^{-1} (DF(\sigma(0)))^{-1} (F(\sigma(z)) - F(\sigma(0)))$$

for $F \in LS$ and $\sigma \in \text{Aut } B^n$. A family $\mathcal{F} \subseteq LS$ is called *linearly invariant* (LIF) if $\Lambda_\sigma(F) \in \mathcal{F}$ for all $F \in \mathcal{F}$ and $\sigma \in \text{Aut } B^n$.

This extends the notion of a linearly invariant family in one dimension, that is, a family \mathcal{F} of analytic functions $f(z) = z + a_2z^2 + \dots$ defined on \mathbb{D} that is closed under Koebe transformations

$$g(z) = \frac{f\left(\frac{z_0+z}{1+\bar{z}_0z}\right) - f(z_0)}{(1 - |z_0|^2)f'(z_0)}, \quad z_0 \in \mathbb{D}.$$

In one dimension, several properties such as growth, covering, distortion and compactness are determined by the *order* $\sup_{f \in \mathcal{F}} a_2(f)$ of the family \mathcal{F} . Pommerenke [1964] showed that the linearly invariant family defined by a Schwarzian derivative bound, $\mathcal{F}_\alpha = \{f : \mathbb{D} \rightarrow \mathbb{C} : f(0) = 0, f'(0) = 1, \|Sf\| \leq \alpha\}$, has norm $\sqrt{1+\alpha/2}$.

Definition 3.2. The *order* of a linearly invariant family \mathcal{F} in arbitrary dimension is defined as

$$\text{ord } \mathcal{F} = \sup_{F \in \mathcal{F}} \sup_{|\vec{v}|=1} \left| \text{tr} \left\{ \frac{1}{2} D^2 F(0)(\vec{v}, \cdot) \right\} \right|,$$

where $|\vec{v}|$ is the Euclidean norm of \vec{v} .

The order of an LIF \mathcal{F} can be written equivalently as

$$\text{ord } \mathcal{F} = \sup_{F \in \mathcal{F}} \left| \frac{1}{2} \sum_{j=1}^n \frac{\partial^2 f_j}{\partial z_j \partial z_k}(0) \right|$$

(see [Pfaltzgraff 1997]). For example, for $n = 2$ a straightforward computation shows that the order is

$$\sup_{F \in \mathcal{F}} \left| \frac{1}{2} \frac{\partial^2 f_1}{\partial z_1^2}(0, 0) + \frac{\partial^2 f_2}{\partial z_1 \partial z_2}(0, 0) \right|.$$

Pfaltzgraff and Suffridge [2000] have introduced the notion of *norm order*, which has much broader applicability to the study of geometric properties of locally biholomorphic mappings than does the order. Consider the Taylor expansion

$$F(z) = z + \frac{1}{2} D^2 F(0)(z, z) + \dots = z + A_2(z, z) + A_3(z, z, z) + \dots,$$

where $A_m(\cdot, \dots, \cdot) = (1/m!) D^m F(0)$, for $m = 1, 2, \dots$, is an m -linear symmetric mapping. Then

$$\|A_m\| = \sup_{|\lambda| \leq 1} |A_m(\lambda, \dots, \lambda)|.$$

Definition 3.3. The *norm order* of a linearly invariant family \mathcal{F} is defined as

$$\|\text{Ord}\| \mathcal{F} = \sup_{F \in \mathcal{F}} \|A_2(F)\|.$$

We define

$$\mathcal{F}_\alpha = \{F : \mathbb{B}^n \rightarrow \mathbb{C}^n \mid F(0) = 0, \quad DF(0) = \text{Id}, \quad \|\mathcal{S}F(z)\| \leq \alpha\}.$$

By [Theorem 2.3](#), this is an LIF.

Remark 3.4. The task of calculating the exact value of the norm of $\mathcal{S}F$ is, in general, not easy, especially because the Bergman and the Euclidean metrics are not conformal. For example, define a locally univalent holomorphic mapping in the ball \mathbb{B}^n by $F_\delta = (f(z_1), \hat{z}g(z_1))$, where $\hat{z} = (0, z_2, \dots, z_n)$,

$$g(z_1) = \frac{1}{1-z_1} \quad \text{and} \quad f(z_1) = \frac{1}{2\delta} \left(\left(\frac{1+z_1}{1-z_1} \right)^\delta - 1 \right).$$

For $n = 2$ a direct calculation shows that

$$\mathbb{S}^1 F_\delta(z) = \begin{pmatrix} \frac{2(\delta-1)}{3(1-z_1^2)} & 0 \\ 0 & 0 \end{pmatrix}, \quad \mathbb{S}^2 F_\delta(z) = \begin{pmatrix} \frac{2z_2(1-\delta)}{(1-z_1)^2(1+z_1)} & -\frac{2(\delta-1)}{3(1-z_1^2)} \\ -\frac{2(\delta-1)}{3(1-z_1^2)} & 0 \end{pmatrix}.$$

Then

$$\mathcal{S}F_\delta(z)(\vec{v}) = \frac{2(\delta-1)}{3(1-z_1^2)} \left(v_1^2, \frac{-3z_2}{1-z_1} v_1^2 - 2v_1 v_2 \right).$$

Is easy to see that for $z_2 = 0$ the norm of the Schwarzian operator is

$$\|\mathcal{S}F_\delta(z)\| = \frac{4}{9}(\delta-1), \quad \delta > 1,$$

while for $z_1 = 0$ with a little bit more effort one can show that

$$\|\mathcal{S}F_\delta(z)\| \frac{2}{\sqrt{3}}(\delta-1), \quad \delta > 1.$$

For arbitrary $z \in \mathbb{B}^2$ we had to resort to a numerical calculation in AMPL [[Fourer et al. 2003](#)]. The numerical results show that

$$\|\mathcal{S}F_\delta(z)\| \leq \frac{2}{\sqrt{3}}(\delta-1), \quad \delta > 1.$$

On the other hand, Pfaltzgraff and Suffridge [[2000](#)] have shown that the norm order of the linear family generated for F_δ is equal to δ ; then for $\delta = \frac{\sqrt{3}}{2}\alpha + 1$ the norm of Schwarzian operator of F_δ is α , so that $F_\delta \in \mathcal{F}_\alpha$ and

$$\|\text{Ord}\| \mathcal{F}_\alpha \geq \frac{\sqrt{3}}{2}\alpha + 1.$$

Pfaltzgraff and Suffridge [[2000](#)] show that an LIF is normal if and only if the norm order is bounded. Our aim is to study the family \mathcal{F}_α , and we shall prove that it is normal. We begin with some lemmas.

Lemma 3.5. *Let F be a holomorphic mapping in \mathcal{F}_α . For $z_1 = (z_1, 0, \dots, 0) \in \mathbb{B}^n$,*

- (i) $|S_{11}^1 F(z_1)| \leq \frac{\sqrt{n+1} \alpha}{1 - |z_1|^2},$
- (ii) $|S_{ii}^1 F(z_1)| \leq \sqrt{n+1} \alpha \quad \text{for } i = 2, 3, \dots, n,$
- (iii) $|S_{11}^k F(z_1)| \leq \frac{\sqrt{n+1} \alpha}{(1 - |z_1|^2)^{3/2}} \quad \text{for } k = 2, 3, \dots, n,$
- (iv) $|S_{1j}^k F(z_1)| \leq \frac{2\sqrt{n+1} \alpha}{1 - |z_1|^2} \quad \text{for } k, j = 2, 3, \dots, n,$
- (v) $|S_{1j}^1 F(z_1)| \leq \frac{2\sqrt{n+1} \alpha}{(1 - |z_1|^2)^{1/2}} \quad \text{for } j = 2, 3, \dots, n,$
- (vi) $|S_{ij}^1 F(z_1)| \leq 2\sqrt{n+1} \alpha \quad \text{for } i \neq j \neq 1,$
- (vii) $|S_{ii}^k F(z_1)| \leq \frac{\sqrt{n+1} \alpha}{(1 - |z_1|^2)^{1/2}} \quad \text{for } k, i = 2, 3, \dots, n,$
- (viii) $|S_{ij}^k F(z_1)| \leq \frac{2\sqrt{n+1} \alpha}{(1 - |z_1|^2)^{1/2}} \quad \text{for } k \neq 1, i \neq j \neq 1.$

Proof. From (2-1) we have

$$g_{11}(z_1, 0, 0, \dots, 0) = \frac{n+1}{(1 - |z_1|^2)^2} \quad \text{and} \quad g_{ij}(z_1, 0, 0, \dots, 0) = \frac{n+1}{(1 - |z_1|^2)},$$

for all $i, j \neq 1$. Let \vec{v} be a unit vector in the Bergman metric. Since $\|\mathcal{S}F(z_1)(\vec{v})\| \leq \alpha$, by setting $\vec{v} = (\lambda, 0, \dots, 0)$ with $\lambda = (1 - |z_1|^2)/\sqrt{n+1}$ we obtain

$$\|\mathcal{S}F(z_1, 0, \dots, 0)(\vec{v})\|^2 = (n+1) \left(\frac{|S_{11}^1 \lambda^2|^2}{(1 - |z_1|^2)^2} + \frac{|S_{11}^2 \lambda^2|^2}{1 - |z_1|^2} + \dots + \frac{|S_{11}^n \lambda^2|^2}{1 - |z_1|^2} \right) \leq \alpha^2,$$

whence (i) and (iii) follow. Now consider $\vec{v} = (0, 0, \dots, \lambda_k, 0, \dots, 0)$ with $\lambda_k^2 = (1 - |z_1|^2)/(n+1)$. As above we have that \vec{v} is a unit vector in the Bergman metric. Since $\|\mathcal{S}F(z_1, 0, \dots, 0)(\vec{v})\| \leq \alpha$ then (ii) and (vii) follow. We obtain (vi) and (vii) analogously, by setting $\vec{v} = (0, \dots, \lambda_i, 0, \dots, \lambda_j, 0, \dots, 0)$, where

$$\lambda_i = \lambda_j = \frac{1}{\sqrt{2}} \frac{(1 - |z_1|^2)^{1/2}}{\sqrt{n+1}}.$$

Finally, (iv) and (v) are established by letting $\vec{v} = (\lambda_1, \dots, \lambda_j, 0, \dots, 0)$, with

$$\lambda_1 = \frac{1}{\sqrt{2}} \frac{(1 - |z_1|^2)}{\sqrt{n+1}} \quad \text{and} \quad \lambda_2 = \frac{1}{\sqrt{2}} \frac{(1 - |z_1|^2)^{1/2}}{\sqrt{n+1}}. \quad \square$$

Lemma 3.6. *If $F \in \mathcal{F}_\alpha$ we have*

$$(3-1) \quad |S_{11}^0 F(z_1, 0, \dots, 0)| \leq \frac{C(n, \alpha)}{(1 - |z_1|^2)^2}$$

with

$$C(n, \alpha) = \left(4n^2 + 2n - 2 + \frac{n+1}{n-1}\right)\alpha^2 + \left(4\sqrt{n+1} + 8\frac{\sqrt{n+1}}{n-1}\right)\alpha,$$

and

$$(3-2) \quad |S_{1j}^0 F(z_1, 0, \dots, 0)| \leq \frac{K(n, \alpha)}{(1 - |z_1|^2)^{3/2}},$$

with

$$K(n, \alpha) = (16 + 3\sqrt{2})\sqrt{n+1}\alpha + 6(n^2 - 1)\alpha.$$

Proof. Differentiating (1-1) and using Proposition 1.2 we get

$$S_{ii}^0 F(z) = -\frac{1}{n-1} \sum_{k=1}^n \frac{\partial}{\partial z_k} S_{ii}^k F(z) + \frac{1}{n-1} \sum_{k=1}^n \sum_{j=1}^n S_{ij}^k F(z) S_{ki}^j F(z),$$

$$S_{ij}^0 F(z) = \frac{\partial}{\partial z_j} S_{ii}^i F(z) - \frac{\partial}{\partial z_i} S_{ij}^i F(z) + \sum_{k=1}^n S_{ii}^k F(z) S_{kj}^i F(z) - \sum_{k=1}^n S_{ij}^k F(z) S_{ki}^i F(z)$$

for $i \neq j$. Thus, the coefficients S_{ij}^0 depend on the S_{ij}^k . Let $F(z_1) = F(z_1, 0, \dots, 0)$, so that for all mappings in \mathcal{F}_α we have

$$\begin{aligned} |S_{11}^0 F(z_1)| &\leq \frac{1}{n-1} \sum_{k=1}^n \left| \frac{\partial}{\partial z_k} S_{11}^k F(z_1) \right| + \frac{1}{n-1} \sum_{k=1}^n \sum_{j=1}^n |S_{1j}^k F(z_1)| |S_{k1}^j F(z_1)|, \\ &= \frac{1}{n-1} \sum_{k=1}^n \left| \frac{\partial}{\partial z_k} S_{11}^k F(z_1) \right| + \frac{1}{n-1} \sum_{k=2}^n \sum_{j=2}^n |S_{1j}^k F(z_1)| |S_{k1}^j F(z_1)| \\ &\quad + \frac{1}{n-1} \sum_{k=2}^n |S_{11}^k F(z_1)| |S_{k1}^1 F(z_1)| + \frac{1}{n-1} |S_{11}^1 F(z_1)|^2. \end{aligned}$$

Therefore Lemma 3.5 implies

$$\begin{aligned} &|S_{11}^0 F(z_1, 0, \dots, 0)| \\ &\leq \frac{4(n+1)(n-1)\alpha^2}{(1 - |z_1|^2)^2} + \frac{2(n+1)\alpha^2}{(1 - |z_1|^2)^2} + \frac{\frac{n+1}{n-1}\alpha^2}{(1 - |z_1|^2)^2} + \frac{1}{n-1} \sum_{k=1}^n \left| \frac{\partial}{\partial z_k} S_{11}^k F(z_1) \right|. \end{aligned}$$

Since $F \in \mathcal{F}_\alpha$, by taking the unit vector $\vec{v} = (\lambda, 0, \dots, 0)$ where

$$|\lambda|^2 = \frac{(1 - |z_1|^2 - |z_k|^2)^2}{(n+1)(1 - |z_k|^2)}$$

in the Bergman metric, a straightforward calculation shows that

$$|S_{11}^k F(z_1, 0, \dots, 0, z_k, 0, \dots, 0)| \leq \frac{\sqrt{n+1} \alpha (1 - |z_k|^2)}{(1 - |z_1|^2 - |z_k|^2)^{3/2}} \quad \text{for } k \neq 1.$$

By considering $S_{11}^k F(z_1, 0, \dots, 0, z_k, 0, \dots, 0)$ as a holomorphic function of z_k we deduce from Cauchy's integral formula that

$$\left| \frac{\partial}{\partial z_k} S_{11}^k F(z_1, 0, \dots, 0) \right| \leq \frac{4\sqrt{n+1} \alpha}{(1 - |z_1|^2)^2} \quad \text{for } k \neq 1.$$

Similarly,

$$\left| \frac{\partial}{\partial z_1} S_{11}^1 F(z_1, 0, \dots, 0) \right| \leq \frac{8\sqrt{n+1} \alpha}{(1 - |z_1|^2)^2}.$$

Using these two inequalities we conclude that

$$\begin{aligned} & |S_{11}^0 F(z_1, 0, \dots, 0)| \\ & \leq \frac{(4n^2 + 2n - 2)\alpha^2}{(1 - |z_1|^2)^2} + \frac{\frac{n+1}{n-1} \alpha^2}{(1 - |z_1|^2)^2} + \frac{4\sqrt{n+1} \alpha}{(1 - |z_1|^2)^2} + \frac{1}{n-1} \frac{8\sqrt{n+1} \alpha}{(1 - |z_1|^2)^2}. \end{aligned}$$

For $j \neq 1$ we have

$$\begin{aligned} |S_{1j}^0 F(z_1)| & \leq \left| \frac{\partial}{\partial z_j} S_{11}^1 F(z_1) \right| + \left| \frac{\partial}{\partial z_1} S_{1j}^1 F(z_1) \right| \\ & \quad \sum_{k=1}^n |S_{11}^k F(z_1)| |S_{kj}^1 F(z_1)| + |S_{1j}^k F(z_1)| |S_{k1}^1 F(z_1)|, \end{aligned}$$

The contribution of the last two summands is at most

$$\frac{2\alpha(n+1)(n-1)}{(1 - |z_1|^2)^{3/2}} + \frac{4\alpha(n+1)(n-1)}{(1 - |z_1|^2)^{3/2}},$$

while the first two can be estimated using Cauchy's integral formula:

$$\left| \frac{\partial}{\partial z_1} S_{1j}^1 F(z_1) \right| \leq \frac{16\sqrt{n+1} \alpha}{(1 - |z_1|^2)^{3/2}}, \quad \left| \frac{\partial}{\partial z_j} S_{11}^1 F(z_1) \right| \leq \frac{3\sqrt{2}\sqrt{n+1} \alpha}{(1 - |z_1|^2)^{3/2}}.$$

Putting it all together,

$$|S_{1j}^0 F(z_1)| \leq \frac{6\alpha(n^2 - 1)}{(1 - |z_1|^2)^{3/2}} + \frac{16\sqrt{n+1} \alpha}{(1 - |z_1|^2)^{3/2}} + \frac{3\sqrt{2}\sqrt{n+1} \alpha}{(1 - |z_1|^2)^{3/2}},$$

proving the theorem. \square

It is clear that if $u(z_1, \dots, z_n)$ is a solution of the system (1-1) then $u(z_1) = u(z_1, 0, \dots, 0)$ satisfies

$$u'' = S_{11}^1 u' + \sum_{j=2}^n S_{11}^j \phi_j + S_{11}^0 u \quad \text{and} \quad \phi'_k = S_{1k}^1 u' + \sum_{j=2}^n S_{1k}^j \phi_j + S_{1k}^0 u$$

for $k = 2, 3, \dots, n$, where $\phi_k(z) = \partial u / \partial z_k$.

Lemma 3.7. *Let $P = P(x)$, $Q = Q(x)$ be continuous functions defined on $[0, 1)$, with $Q(x) \geq 0$. Let $u = u(x)$, $v = v(x)$ satisfy*

$$\begin{aligned} u'' + Pu + Q &\geq 0, \quad u(0) = 1, \quad u'(0) = 0, \\ v'' + Pv + Q &= 0, \quad v(0) = 1, \quad v'(0) = 0. \end{aligned}$$

Then $u \geq v$ on $[0, x_0)$, where x_0 is the first zero of v .

Proof. For $\varepsilon > 0$, let $u_\varepsilon = u + \varepsilon y$, where y is solution of $y'' + Py = 0$, $y(0) = 0$, $y'(0) = 1$. Then $w = u'_\varepsilon v - v' u_\varepsilon$ satisfies $w(0) = \varepsilon > 0$ and $w' \geq Q(u_\varepsilon - v)$. Because of the initial conditions of u_ε and v , the function w has $w' > 0$ on an interval $(0, r)$. But then $w > 0$ (in fact, $\geq \varepsilon$) on that interval, which implies that $u'_\varepsilon / u_\varepsilon > v' / v$ if $v > 0$, thus $u_\varepsilon > v$. It follows from this argument that the first zero of u_ε cannot occur before the first zero of v , and the lemma obtains after letting $\varepsilon \rightarrow 0$. \square

Lemma 3.8. *Let u be a solution of the system (1-1) satisfying $u(0, \dots, 0) = 1$, $\nabla u(0, \dots, 0) = 0$ and $P_{ij}^k = S_{ij}^k F$ with $F \in \mathcal{F}_\alpha$. Then there exists $r > 0$ and $\delta > 0$ such that $|u| > \delta > 0$ for $|z| < r$.*

Proof. Let $z_0 \in \mathbb{B}^n$ be a zero of u of smallest euclidean norm, that is, $u(z_0) = 0$ and $u(z) \neq 0$ for $|z| < |z_0| = r_0$. Since \mathcal{F}_α is a linearly invariant family we can assume that $z_0 = (x_0, 0, \dots, 0)$. We shall study the zeros of the function $u(x) = u(x, 0, \dots, 0)$ in $0 < x < 1$. If $F(x) = F(x, 0, \dots, 0)$, then $u(x)$ and $\varphi_k(x) = (\partial u / \partial z_k)(x, 0, \dots, 0)$ satisfies the system

$$\begin{aligned} (3-3) \quad u''(x) &= \sum_{k=1}^n S_{11}^k F(x) \varphi_k(x) + S_{11}^0 F(x) u(x), \\ \varphi'_j(x) &= \sum_{k=1}^n S_{1j}^k F(x) \varphi_k(x) + S_{1j}^0 F(x) u(x), \quad j = 2, \dots, n, \end{aligned}$$

with initial conditions $u(0) = 1$ and $\varphi_k(0) = 0$. With $\theta = (\varphi_1, \dots, \varphi_n, u)$, we can rewrite the system (3-3) as

$$(3-4) \quad \theta'(x) = A(x) \cdot \theta(x), \quad \theta(0) = (0, 0, \dots, 1),$$

where $A(x)$ is the $(n+1) \times (n+1)$ matrix of coefficients of the system. Let $f^2(x) = \|\theta(x)\|^2$ be the square of the Euclidean norm of $\theta(x)$. Using \cdot to represent the

Euclidean inner product of vectors in $\mathbb{C}^{n+1} = \mathbb{R}^{2n+2}$, we have

$$f'(x)f(x) = \theta'(x) \cdot \theta(x) = A(x)\theta(x) \cdot \theta(x);$$

therefore $f'(x)f(x) \leq \|A(x)\|\|\theta(x)\|^2 = \|A(x)\|f^2(x)$, so

$$\frac{f'(x)}{f(x)} \leq \|A(x)\|.$$

Since $f(0) = 1$ we conclude that $f(x) \leq e^{\int_0^x p(s) ds}$, where $p(s)$ stands for the bounds obtained for $\|A(s)\|$ from Lemmas 3.5 and 3.6. In particular, we have

$$|u'(x)| \leq e^{\int_0^x p(s) ds}, \quad |\varphi_k(x)| \leq e^{\int_0^x p(s) ds} \quad \text{for } k = 2, \dots, n.$$

Setting $U^2(x) = |u(x)|^2$, we obtain $2UU' = 2\operatorname{Re}(u'\bar{u})$, hence $(U')^2 + UU'' = \operatorname{Re}(u''\bar{u}) + |u'|^2$, $U(0) = 1$, $U'(0) = 0$. Since $|U'| \leq |u'|$, we have

$$UU'' \geq \operatorname{Re}(u''\bar{u}).$$

Using this in (3-3) we get

$$UU'' \geq \operatorname{Re}\{S_{11}^0 F(x)\}U^2 + \operatorname{Re}(q(x)\bar{u}),$$

where $q(x) = S_{11}^1 F(x)u'(x) + \sum_{k=2}^n S_{11}^k F(x)\varphi_k(x)$; hence

$$U'' \geq -|S_{11}^0 F(x)|U - |q(x)|,$$

or $U'' + P(x)U + Q(x) \geq 0$, where P and Q are the bounds obtained from Lemmas 3.5 and 3.6 for $|S_{11}^0 F(x, 0, \dots, 0)|$ and $|q(x)|$, respectively. It follows now from Lemma 3.7 that $U \geq v$ on $[0, x_0)$, where x_0 is the first zero of v , which is solution of $v'' + Pv + Q = 0$, $v(0) = 1$, $v'(0) = 0$. The lemma follows taking $r < x_0$. \square

Remark 3.9. It is clear that we need to estimate the first zero of the function v . In fact, we proved that $|S_{11}^0 F(x, 0, \dots, 0)| \leq c(n, \alpha)(1 - x^2)^{-2} = P$, where $c = c(n, \alpha)$ is a constant. Also one can obtain from Lemmas 3.5 and 3.6 a bound of $|q(x)|$ of the form

$$|q(x)| \leq \frac{M}{(1 - x^2)^{\delta+1}} = Q,$$

where $M = \sqrt{n(n+1)}\alpha$ and δ also depends on n and α . Then v is a solution of

$$v'' + \frac{c}{(1 - x^2)^2} v + \frac{M}{(1 - x^2)^{\delta+1}} = 0, \quad v(0) = 1, \quad v'(0) = 0.$$

In general, for given constants c, M, δ , one will be able to estimate the first zero of v only numerically. However, if $\delta < 1$ then by comparison, it follows that the first zero of v does not occur before the first zero of the solution w of

$$w'' + \frac{c}{(1 - x^2)^2} w + \frac{M}{(1 - x^2)^2} = 0, \quad w(0) = 1, \quad w'(0) = 0,$$

and this can be determined analytically. Indeed we have $w = (M + 1)y_c - M$, where y_c is the solution of

$$y'' + \frac{c}{(1-x^2)^2} y = 0, \quad y(0) = 1, \quad y'(0) = 0,$$

which can be found, for example, in [Kamke 1930]. Thus the first zero of w is the solution of the (transcendental) equation $y_c(x) = M/(1+M)$.

Theorem 3.10. Fix $\alpha < \infty$. The family

$$\mathcal{F}_\alpha = \left\{ F : \mathbb{B}^n \rightarrow \mathbb{C}^n \mid F(0) = 0, \quad DF(0) = \text{Id}, \quad \|\mathcal{S}F(z)\| \leq \alpha \right\}$$

is a normal family.

Proof. Let $F \in \mathcal{F}_\alpha$. From Proposition 1.2 we have

$$(3-5) \quad F = \left(\frac{u_1}{u_0}, \dots, \frac{u_n}{u_0} \right) = (f_1, \dots, f_n),$$

where u_i and $u_0 = \Delta^{-1/n+1}$ are linearly independent solutions of (1-1) such that $(\partial u_i / \partial z_k)(0) = 0$ for all $k \neq i$ and $(\partial u_i / \partial z_i)(0) = 1$ for $i = 1, \dots, n$; see [Yoshida 1984]. From equation (2-2) we deduce that

$$D^2F(0)(\vec{v}, \vec{v}) = \mathcal{S}F(0)(\vec{v}, \vec{v}) + 2(\nabla u_0(0) \cdot \vec{v}) \vec{v}.$$

Hence $|A_2(z)|$ will be uniformly bounded for F in the family \mathcal{F}_α provided that the same holds for the derivatives $|(\partial u_0 / \partial z_j)(0)|$ for $j = 1, \dots, n$. To show the latter, consider the composition $G = T \circ F$ with the Möbius transformation given by

$$T(z) = \frac{z}{1+z \cdot \bar{a}},$$

where we have introduced the inner product $\langle z, w \rangle = z_1 \bar{w}_1 + \dots + z_n \bar{w}_n$. Using (3-5), we get

$$\begin{aligned} G(z) &= \frac{F(z)}{1 + \langle F(z), \bar{a} \rangle} = \left(\frac{u_1}{u_0 + a_1 u_1 + \dots + a_n u_n}, \dots, \frac{u_n}{u_0 + a_1 u_1 + \dots + a_n u_n} \right) \\ &= \left(\frac{\tilde{u}_1}{\tilde{u}_0}, \dots, \frac{\tilde{u}_n}{\tilde{u}_0} \right), \end{aligned}$$

where $\tilde{u}_0 = u_0 + a_1 u_1 + \dots + a_n u_n$ and $\tilde{u}_i = u_i$ for $i = 1, \dots, n$. Differentiating and setting $a_k = (\partial u_0 / \partial z_k)(0)$ for $k = 1, \dots, n$, we obtain $\nabla(\tilde{u}_0)(0) = 0$. This may introduce a pole of G but away from the origin. The function \tilde{u}_0 satisfies the system

$$\frac{\partial^2 \tilde{u}_0}{\partial z_i \partial z_j}(z) = \sum_{k=1}^n S_{ij}^k F(z) \frac{\partial \tilde{u}_0}{\partial z_k} + S_{ij}^0 F(z) \tilde{u}_0(z), \quad \tilde{u}_0(0) = 1, \quad \nabla \tilde{u}_0(0) = 0,$$

and in view of [Lemma 3.8](#), \tilde{u}_0 does not vanish on B_r for some $r > 0$. At the same time, since satisfies $\tilde{u}_i(0) = 0$ and $|\nabla\tilde{u}_i(0)| = 1$ for each $i = 1, \dots, n$, it is easy to see from (1-1) and the bounds in [Lemmas 3.5](#) and [3.6](#) that the functions \tilde{u}_i will be uniformly bounded on compact subsets. Therefore, the class of mappings G obtained with this normalization is normal on $|z| < r_0$ with $r_0 < r$; then there exists $s_0 > 0$ such that $G(\mathbb{B}_{r_0}^n) \supset \mathbb{B}_{s_0}^n$. Since the image of $G := (\tilde{f}_1, \dots, \tilde{f}_n)$ covers a ball of radius s_0 and

$$F = \frac{G}{1 - \langle a, \tilde{f} \rangle}$$

is holomorphic, we conclude that $|a_1|^2 + \dots + |a_n|^2 \leq 1/s_0^2$. This shows that $|\nabla u_0(0)| = \sqrt{|a_1|^2 + \dots + |a_n|^2}$ is uniformly bounded and the theorem follows. \square

In analogy to the result of Pommerenke cited on page [206](#), we have:

Theorem 3.11. $\|\text{Ord}\| \mathcal{F}_\alpha \leq \frac{\sqrt{n+1}}{2} \alpha + \lambda_\alpha$, where $\lambda_\alpha = \frac{2\sqrt{n}}{n+1} \text{ord } \mathcal{F}_\alpha$.

Proof. Equation (2-2) yields $D^2F(0)(\vec{v}, \vec{v}) = \mathcal{S}F(0)(\vec{v}, \vec{v}) + 2(\nabla u_0(0) \cdot \vec{v}) \vec{v}$. Is not difficult to see that

$$\frac{\partial u_0}{\partial z_k}(0) = -\frac{1}{n+1} \sum_{j=1}^n \frac{\partial^2 f_j}{\partial z_j \partial z_k}(0);$$

hence, taking the Euclidean norm and the supremum over all unit vectors \vec{v} , we obtain

$$|A_2(F)| \leq \frac{\sqrt{n+1}}{2} \|\mathcal{S}F(0)\| + |\nabla u_0(0)|,$$

where $\|\cdot\|$ is the Bergman metric. Therefore

$$\|\text{Ord}\| F_\alpha \leq \frac{\sqrt{n+1}}{2} \alpha + \lambda_\alpha. \quad \square$$

Nehari [[1949](#)] proved that if f belongs to the univalent class in the unit disk, the Schwarzian derivative of f has norm at most 6; but this has no counterpart in higher dimensions, since the norm order of univalent mappings is infinite.

Corollary 3.12. *Let F be a convex holomorphic mapping in \mathbb{B}^2 , then*

$$\|\mathcal{S}F(z)\| \leq \alpha_K, \quad \text{where } \alpha_K = \frac{2}{\sqrt{3}} + \frac{4\sqrt{2}}{3\sqrt{3}} \cdot 1.761.$$

Proof. Barnard, FitzGerald and Gong [[Barnard et al. 1994](#)] established that $\frac{3}{2} \leq \text{ord } K(\mathbb{B}^2) \leq 1.761$ for the family of convex mappings $K(\mathbb{B}^2)$. Using (2-2) and setting the Bergman norm in the origin, we deduce that

$$\|\mathcal{S}F(0)(\vec{v})\| \leq \sqrt{3} |D^2F(0)(\vec{v})| + 2|\nabla u_0(0) \cdot \vec{v}|,$$

where $|\cdot|$ is the Euclidean norm. Thus, taking the supremum over all vectors with $\|\vec{v}\| = 1$, we obtain

$$\|\mathcal{S}F(0)\| \leq \frac{2}{\sqrt{3}} \|\text{Ord}\| K(\mathbb{B}^2) + \frac{4\sqrt{2}}{3\sqrt{3}} \text{ord } K(\mathbb{B}^2) \leq \frac{2}{\sqrt{3}} + \frac{4\sqrt{2}}{3\sqrt{3}} \cdot 1.761.$$

To establish the estimate at an arbitrary point in the ball, apply the appropriate Koebe transform and [Theorem 2.3](#). □

The order of $K(\mathbb{B}^n)$ for $n \geq 2$ is unknown, but Liu [1989] has established an upper bound in any dimension. The conjecture in [Barnard et al. 1994] that $\text{ord } K(\mathbb{B}^n) = \frac{1}{2}(n + 1)$ for $n \geq 2$ was shown to be false by Pfaltzgraft and Suffridge [2000].

Definition 3.13. A holomorphic mapping $F \in \mathcal{F}_\alpha$ is an *extremal order function* for \mathcal{F}_α if its order is equal to the order of family \mathcal{F}_α .

Theorem 3.14. *Let F be a extremal order function for the family \mathcal{F}_α . There exists $\{z_k\} \in \mathbb{B}^n$ with $|z_k| \rightarrow 1$ when $k \rightarrow \infty$, such that*

$$\lim_{k \rightarrow \infty} |F(z_k)| = \infty.$$

Proof. Let $F = (f_1, \dots, f_n) = (u_1/u_0, \dots, u_n/u_0)$ be an extremal order mapping and consider the Möbius transformation

$$G = \left(\frac{f_1}{1 + \varepsilon f_1}, \dots, \frac{f_n}{1 + \varepsilon f_1} \right),$$

for $\varepsilon > 0$. We have $\mathcal{S}F(z) = \mathcal{S}G(z)$, $G(0) = 0$, $DG(0) = \text{Id}$ and we can write $G = (u_1/\tilde{u}_0, \dots, u_n/\tilde{u}_0)$, where $\tilde{u}_0 = u_0 + \varepsilon u_1$. Differentiating with respect to z_1 and evaluating in the origin, we obtain

$$\frac{\partial \tilde{u}_0}{\partial z_1}(0) = \frac{\partial u_0}{\partial z_1}(0) + \varepsilon.$$

But is easy to see that

$$\frac{\partial u_0}{\partial z_1}(0) = \frac{1}{n + 1} \sum_{j=1}^n \frac{\partial^2 f_j}{\partial z_1 \partial z_j}(0) = \frac{2}{n + 1} \text{ord } \mathcal{F}_\alpha.$$

If G were holomorphic in the ball, it would lie in \mathcal{F}_α , contradicting the fact that F is an extremal order function. Hence there must exist a point z_ε such that $1 + \varepsilon f_1(z_\varepsilon) = 0$, that is, $f_1(z_\varepsilon) = -1/\varepsilon$. It is also clear that $|z_\varepsilon| \rightarrow 1$ when $\varepsilon \rightarrow 0$, which finishes the proof. □

4. An estimate for λ_α

To find explicit bounds for λ_α in terms of α we have to estimate the radius s_0 of a ball covered by the function $G = (u_1/u_0, \dots, u_n/u_0)$ considered in the proof of [Theorem 3.10](#). Recall that the u_i formed a set of linearly independent solutions of (1-1) with initial conditions $u_0(0) = 1$, $\nabla u_0(0) = 0$, $u_i(0) = 0$ and $|\nabla u_i(0)| = 1$ for $i = 1, \dots, n$. Set $u(x) = u_k(x, 0, \dots, 0)$ and

$$\theta(x) = \left(\frac{\partial u}{\partial z_1}(x), \frac{\partial u}{\partial z_2}(x)(1-x^2)^{-\frac{1}{2}}, \dots, \frac{\partial u}{\partial z_n}(x)(1-x^2)^{-\frac{1}{2}}, u(x)(1-x^2)^{-1} \right).$$

It follows from [Lemmas 3.5](#) and [3.6](#) that $\theta' = B\theta$ for some modification B of the matrix A of (3-4), such that

$$\|B(x)\| \leq \frac{k}{1-x^2} \quad \text{with } k = \delta(n, \alpha) + 2,$$

where $\delta(n, \alpha) \rightarrow 0$ when $\alpha \rightarrow 0$. As in the proof of [Lemma 3.8](#) we obtain

$$(4-1) \quad \|\theta(x)\| \leq \left(\frac{1+x}{1-x} \right)^{k/2}.$$

In particular, taking $u = u_0$ we get $|u_0(x)| \leq (1-x^2) \left(\frac{1+x}{1-x} \right)^{k/2}$. Now we need to find a lower bound for $|u_i|$, $i = 1, \dots, n$. Consider the real function $U(x) = |u(x)|$, for which

$$U'' \geq - \left| S_{11}^1 G(x) \frac{\partial u}{\partial z_1}(x) + \dots + S_{11}^n G(x) \frac{\partial u}{\partial z_n}(x) + S_{11}^0 G(x) u(x) \right|.$$

Using (4-1) and [Lemmas 3.5](#) and [3.6](#), we obtain

$$U'' + \frac{C}{(1-x^2)^2} U \geq - \frac{\sqrt{n(n+1)}\alpha}{1-x^2} \left(\frac{1+x}{1-x} \right)^{k/2}, \quad U(0) = 0, \quad U'(0) = 1.$$

Then $U \geq y$ until the first zero $x = x_\alpha$ of the solution y of

$$y'' + \frac{C}{(1-x^2)^2} y = - \frac{\sqrt{n(n+1)}\alpha}{1-x^2} \left(\frac{1+x}{1-x} \right)^{k/2}, \quad y(0) = 0, \quad y'(0) = 1.$$

Hence

$$|G(x)| \geq \frac{\sqrt{n} y(x)}{(1-x^2) \left(\frac{1+x}{1-x} \right)^{k/2}} = \phi(x).$$

It follows that $G(\mathbb{B}_{x_\alpha})$ covers a ball of radius $M_\alpha = \max\{\phi(x) : 0 < x \leq x_\alpha\}$. From the proof of [Theorem 3.10](#) we finally see that

$$\lambda_\alpha \leq \frac{1}{M_\alpha}.$$

Acknowledgment

We thank the referee for useful suggestions and an interesting discussion.

References

- [Barnard et al. 1994] R. W. Barnard, C. H. FitzGerald, and S. Gong, “A distortion theorem for biholomorphic mappings in \mathbb{C}^2 ”, *Trans. Amer. Math. Soc.* **344**:2 (1994), 907–924. [MR 94k:32034](#) [Zbl 0814.32004](#)
- [Chuaqui and Osgood 1993] M. Chuaqui and B. Osgood, “Sharp distortion theorems associated with the Schwarzian derivative”, *J. London Math. Soc.* (2) **48**:2 (1993), 289–298. [MR 94g:30005](#) [Zbl 0792.30013](#)
- [Epstein 1986] C. L. Epstein, “The hyperbolic Gauss map and quasiconformal reflections”, *J. Reine Angew. Math.* **372** (1986), 96–135. [MR 88b:30029](#) [Zbl 0591.30018](#)
- [Fourer et al. 2003] R. Fourer, D. M. Gay, and B. W. Kernighan, *AMPL: a modeling language for mathematical programming*, 2nd ed., Duxbury Press and Brooks/Cole, 2003.
- [Kamke 1930] E. Kamke, *Differentialgleichungen reeller Funktionen*, Akademische Verlagsgesellschaft, Leipzig, 1930. Reprinted Chelsea, New York, 1948. [JFM 56.0375.03](#)
- [Kraus 1932] W. Kraus, “Ueber den Zusammenhang einiger Charakteristiken eines einfach zusammenhängenden bereiches mit der Kreisabbildung”, *Mitt. Math. Sem. Giessen* **21** (1932), 1–28. [Zbl 0005.30104](#)
- [Liu 1989] T. Liu, “The distortion theorem for biholomorphic mappings in \mathbb{C}^n ”, Preprint, 1989.
- [Molzon and Pinney Mortensen 1997] R. Molzon and K. Pinney Mortensen, “Univalence of holomorphic mappings”, *Pacific J. Math.* **180**:1 (1997), 125–133. [MR 98k:32034](#) [Zbl 0898.32015](#)
- [Nehari 1949] Z. Nehari, “The Schwarzian derivative and schlicht functions”, *Bull. Amer. Math. Soc.* **55** (1949), 545–551. [MR 10,696e](#) [Zbl 0035.05104](#)
- [Oda 1974] T. Oda, “On Schwarzian derivatives in several variables”, *Sūrikaisekikenkyūsho Kōkyūroku (RIMS, Kyoto)* **226** (1974). In Japanese.
- [Pfaltzgraff 1974] J. A. Pfaltzgraff, “Subordination chains and univalence of holomorphic mappings in \mathbb{C}^n ”, *Math. Ann.* **210** (1974), 55–68. [MR 50 #4997](#) [Zbl 0275.32012](#)
- [Pfaltzgraff 1997] J. A. Pfaltzgraff, “Distortion of locally biholomorphic maps of the n -ball”, *Complex Variables Theory Appl.* **33**:1-4 (1997), 239–253. [MR 99a:32030](#) [Zbl 0912.32017](#)
- [Pfaltzgraff and Suffridge 2000] J. A. Pfaltzgraff and T. J. Suffridge, “Norm order and geometric properties of holomorphic mappings in \mathbb{C}^n ”, *J. Anal. Math.* **82** (2000), 285–313. [MR 2001k:32028](#) [Zbl 0978.32017](#)
- [Pommerenke 1964] C. Pommerenke, “Linear-invariante Familien analytischer Funktionen, I, II”, *Math. Ann.* **155** (1964), 108–154 and **156** (1964), 226–262. [MR 1513275](#) [Zbl 0128.30105](#)
- [Yoshida 1976] M. Yoshida, “Canonical forms of some systems of linear partial differential equations”, *Proc. Japan Acad.* **52**:9 (1976), 473–476. [MR 54 #13962](#) [Zbl 0378.35013](#)
- [Yoshida 1984] M. Yoshida, “Orbifold-uniformizing differential equations”, *Math. Ann.* **267** (1984), 125–142. [MR 85j:14013](#) [Zbl 0521.58052](#)

RODRIGO HERNÁNDEZ R.
UNIVERSIDAD ADOLFO IBÁÑEZ
FACULTAD DE CIENCIA Y TECNOLOGÍA
AVENIDA LAS TORRES 2640
PEÑALOLÉN
CHILE

rodrigo.hernandez@uai.cl