

Pacific Journal of Mathematics

TILINGS DEFINED BY AFFINE WEYL GROUPS

ECKHARD MEINRENKEN

TILINGS DEFINED BY AFFINE WEYL GROUPS

ECKHARD MEINRENKEN

Let W be a Weyl group, presented as a reflection group on a Euclidean vector space V , and $C \subset V$ an open Weyl chamber. In a recent paper, Waldspurger proved that the images $(\text{id} - w)(C)$ for $w \in W$ are all disjoint, with union the closed cone spanned by the positive roots. We prove that similarly, the images $(\text{id} - w)(A)$ of the open Weyl alcove A , for $w \in W^{\text{aff}}$ in the affine Weyl group, are disjoint and their union is V .

1. Introduction

Let W be the Weyl group of a simple Lie algebra, presented as a crystallographic reflection group in a finite-dimensional Euclidean vector space $(V, \langle \cdot, \cdot \rangle)$. Choose a fundamental Weyl chamber $C \subset V$, and let D be its dual cone, that is, the open cone spanned by the corresponding positive roots. Waldspurger [2007] proved the following remarkable result. Consider the linear transformations $(\text{id} - w): V \rightarrow V$ defined by elements $w \in W$.

Theorem 1.1 (Waldspurger). *The images $D_w := (\text{id} - w)(C)$ for $w \in W$ are all disjoint, and their union is the closed cone spanned by the positive roots:*

$$\bar{D} = \bigcup_{w \in W} D_w.$$

For instance, the identity transformation $w = \text{id}$ corresponds to $D_{\text{id}} = \{0\}$ in this decomposition, while the reflection s_α defined by a positive root α corresponds to the open half-line $D_{s_\alpha} = \mathbb{R}_{>0} \cdot \alpha$.

The aim of this note is to prove a similar result for the *affine* Weyl group W^{aff} . Recall that $W^{\text{aff}} = \Lambda \rtimes W$, where the coroot lattice $\Lambda \subset V$ acts by translations. Let $A \subset C$ be the Weyl alcove, with $0 \in \bar{A}$.


Theorem 1.2. *The images $V_w = (\text{id} - w)(A)$ for $w \in W^{\text{aff}}$ are all disjoint, and their union is V :*

$$V = \bigcup_{w \in W^{\text{aff}}} V_w.$$

MSC2000: 20F55, 22E46.

Keywords: affine Weyl group, reflection groups, alcoves.

The author is partially supported by an NSERC Discovery Grant and a Steacie Fellowship.



The figure above is a picture of the resulting tiling of V for the root system \mathbf{G}_2 . Up to translation by elements of the lattice Λ , there are five 2-dimensional tiles, corresponding to the five Weyl group elements with trivial fixed point set. With s_1 and s_2 denoting the simple reflections, the lightly shaded polytopes are labeled by the Coxeter elements s_1s_2 and s_2s_1 , the medium shaded polytopes by $(s_1s_2)^2$ and $(s_2s_1)^2$, and the darkly shaded polytope by the longest Weyl group element $w_0 = (s_1s_2)^3$.

One also has the following related statement.

Theorem 1.3. *Suppose $S \in \text{End}(V)$ with $\|S\| < 1$. Then the sets $V_w^{(S)} = (S - w)(A)$ for $w \in W^{\text{aff}}$ are all disjoint, and their closures cover V :*

$$V = \bigcup_{w \in W^{\text{aff}}} \overline{V}_w^{(S)}.$$

Note that for $S = 0$ the resulting decomposition of V is just the Stiefel diagram, while for $S = \tau \text{ id}$ with $\tau \rightarrow 1$ one recovers the decomposition from [Theorem 1.2](#).

The proof of [Theorem 1.2](#) is in large parts parallel to Waldspurger’s [\[2007\]](#) proof of [Theorem 1.1](#). We will nevertheless give full details so the paper is self-contained.

2. Notation

With no loss of generality we will take W to be irreducible. Let $\mathfrak{R} \subset V$ be the set of roots, $\{\alpha_1, \dots, \alpha_l\} \subset \mathfrak{R}$ a set of simple roots, and

$$C = \{x \mid \langle \alpha_i, x \rangle > 0, \ i = 1, \dots, l\}$$

the corresponding Weyl chamber. We denote by $\alpha_{\max} \in \mathfrak{R}$ the highest root, and $\alpha_0 = -\alpha_{\max}$ the lowest root. The open Weyl alcove is the l -dimensional simplex

defined as

$$A = \{x \mid \langle \alpha_i, x \rangle + \delta_{i,0} > 0, \ i = 0, \dots, l\}.$$

Its faces are indexed by the proper subsets $I \subset \{0, \dots, l\}$, where A_I is given by inequalities $\langle \alpha_i, x \rangle + \delta_{i,0} > 0$ for $i \notin I$ and equalities $\langle \alpha_i, x \rangle + \delta_{i,0} = 0$ for $i \in I$. Each A_I has codimension $|I|$. In particular, $A_i = A_{\{i\}}$ are the codimension 1 faces, with α_i as inward-pointing normal vectors. Let s_i be the affine reflections across the affine hyperplanes supporting A_i , that is,

$$s_i : x \mapsto x - (\langle \alpha_i, x \rangle + \delta_{i,0}) \alpha_i^\vee \quad \text{for } i = 0, \dots, l,$$

where $\alpha_i^\vee = 2\alpha_i / \langle \alpha_i, \alpha_i \rangle$ is the simple coroot corresponding to α_i . The Weyl group W is generated by the reflections s_1, \dots, s_l , while the affine Weyl group W^{aff} is generated by the affine reflections s_0, \dots, s_l . The affine Weyl group is a semidirect product

$$W^{\text{aff}} = \Lambda \rtimes W,$$

where the coroot lattice $\Lambda = \mathbb{Z}[\alpha_1^\vee, \dots, \alpha_l^\vee] \subset V$ acts on V by translations. For any $w \in W^{\text{aff}}$, we will denote by $\tilde{w} \in W$ its image under the quotient map $W^{\text{aff}} \rightarrow W$, that is, $\tilde{w}(x) = w(x) - w(0)$, and by $\lambda_w = w(0) \in \Lambda$ the corresponding lattice vector.

The stabilizer of any element of A_I is the subgroup W_I^{aff} generated by s_i for $i \in I$. It is a finite subgroup of W^{aff} , and the map $w \mapsto \tilde{w}$ induces an isomorphism onto the subgroup W_I generated by \tilde{s}_i for $i \in I$. Recall that W_I is itself a Weyl group (not necessarily irreducible): Its Dynkin diagram is obtained from the extended Dynkin diagram of the root system \mathfrak{A} by removing all vertices that are in I .

3. The top-dimensional polytopes

For any $w \in W^{\text{aff}}$, the subset $V_w = (\text{id} - w)(A)$ is the relative interior of a convex polytope in the affine subspace $\text{ran}(\text{id} - w)$. Let

$$W_{\text{reg}}^{\text{aff}} = \{w \in W^{\text{aff}} \mid (\text{id} - w) \text{ is invertible}\}$$

and $W_{\text{reg}} = W \cap W_{\text{reg}}^{\text{aff}}$, so that $w \in W_{\text{reg}}^{\text{aff}}$ if and only if $\tilde{w} \in W_{\text{reg}}$. The top dimensional polytopes V_w are those indexed by $w \in W_{\text{reg}}^{\text{aff}}$, and the faces of these polytopes are $V_{w,I} := (\text{id} - w)(A_I)$. For $w \in W_{\text{reg}}$ and $i = 0, \dots, l$, let

$$n_{w,i} := (\text{id} - \tilde{w}^{-1})^{-1}(\alpha_i).$$

Lemma 3.1. *For all $w \in W_{\text{reg}}^{\text{aff}}$, the open polytope V_w is given by the inequalities*

$$\langle n_{w,i}, \xi + \lambda_w \rangle + \delta_{i,0} > 0 \quad \text{for } i = 0, \dots, l.$$

The face $V_{w,I} = (\text{id} - w)(A_I)$ is obtained by replacing the inequalities for $i \in I$ by equalities.

Proof. For any $\zeta = (\text{id} - w)x \in V$, we have

$$\langle \alpha_i, x \rangle = \langle (\text{id} - \tilde{w}^{-1})^{-1} \alpha_i, (\text{id} - \tilde{w})x \rangle = \langle n_{w,i}, (\text{id} - \tilde{w})x \rangle = \langle n_{w,i}, \zeta + \lambda_w \rangle,$$

since \tilde{w}^{-1} is the transpose of \tilde{w} under the inner product $\langle \cdot, \cdot \rangle$. This gives the description of V_w and of its faces $V_{w,I}$. \square

Lemma 3.2. *Suppose $w \in W_{\text{reg}}^{\text{aff}}$ for $i \in \{0, \dots, l\}$. Then $V_{w,i} = V_{\sigma,i} \subset \text{ran}(\text{id} - \sigma)$ with $\sigma = ws_i$. In particular, σ is an affine reflection, and $n_{w,i}$ is a vector normal to the affine hyperplane $\text{ran}(\text{id} - \sigma)$. One has $\langle n_{w,i}, \alpha_i^\vee \rangle = 1$.*

Proof. For any orthogonal transformation $g \in O(V)$ and any reflection $s \in O(V)$, the dimension of the fixed point set of the orthogonal transformations g and gs differ by ± 1 . Since \tilde{w} fixes only the origin, it follows that $\tilde{\sigma}$ has a 1-dimensional fixed point set. Hence $\text{ran}(\text{id} - \sigma)$ is an affine hyperplane, and σ is the affine reflection across that hyperplane. Since s_i fixes A_i , we have

$$V_{w,i} = (\text{id} - w)(A_i) = (\text{id} - ws_i)(A_i) = V_{\sigma,i} \subset \text{ran}(\text{id} - \sigma).$$

By definition $n_{w,i} - \tilde{w}^{-1}n_{w,i} = \alpha_i$. Hence

$$-2\langle n_{w,i}, \alpha_i \rangle + \langle \alpha_i, \alpha_i \rangle = \|n_{w,i} - \alpha_i\|^2 - \|n_{w,i}\|^2 = \|\tilde{w}^{-1}n_{w,i}\|^2 - \|n_{w,i}\|^2 = 0. \quad \square$$

The following proposition indicates how the top-dimensional polytopes $V_{w,i}$ are glued along the polytopes of codimension 1.

Proposition 3.3. *Let $\sigma \in W^{\text{aff}}$ be an affine reflection, that is, $\text{ran}(\text{id} - \sigma)$ is an affine hyperplane. Consider*

$$(1) \quad \zeta \in V_\sigma \setminus \bigcup_{|I| \geq 2} V_{\sigma,I}.$$

Then there are two distinct indices $i, i' \in \{0, \dots, l\}$ such that $\zeta \in V_{\sigma,i} \cap V_{\sigma,i'}$. Furthermore, $w = \sigma s_i$ and $w' = \sigma s_{i'}$ are both in $W_{\text{reg}}^{\text{aff}}$, so that $V_{w,i} = V_{\sigma,i}$ and $V_{w',i'} = V_{\sigma,i'}$, and the polytopes $V_w, V_{w'}$ are on opposite sides of the affine hyperplane $\text{ran}(\text{id} - \sigma)$.

Proof. Let n be a generator of the 1-dimensional subspace $\ker(\text{id} - \tilde{\sigma})$. Then n is a vector normal to $\text{ran}(\text{id} - \sigma)$. The preimage $(\text{id} - \sigma)^{-1}(\zeta) \subset V$ is an affine line in the direction of n . Since $\zeta \in V_\sigma$, this line intersects A ; hence it intersects the boundary $\partial \bar{A}$ in exactly two points x and x' . By (1), x and x' are contained in two distinct codimension 1 boundary faces A_i and $A_{i'}$. Since n is inward-pointing at one of the boundary faces, and outward-pointing at the other, the inner products $\langle n, \alpha_i \rangle$ and $\langle n, \alpha_{i'} \rangle$ are both nonzero, with opposite signs. Let $w = \sigma s_i$ and $w' = \sigma s_{i'}$. We will show that $w \in W_{\text{reg}}^{\text{aff}}$, that is, $\tilde{w} \in W_{\text{reg}}$ (the proof for w' is




Figure 1. The set X for the root system \mathbf{B}_2 .

similar). Let $z \in V$ with $\tilde{w}z = z$. Then $\tilde{\sigma}^{-1}z = \tilde{s}_i z$, so

$$(\text{id} - \tilde{\sigma}^{-1})(z) = (\text{id} - \tilde{s}_i)(z) = \langle \alpha_i, z \rangle \alpha_i^\vee.$$

The left side lies in $\text{ran}(\text{id} - \tilde{\sigma})$, which is orthogonal to n , while the right side is proportional to α_i . Since $\langle n, \alpha_i \rangle \neq 0$, this is only possible if both sides are 0. Thus z is fixed under $\tilde{\sigma}$, and hence a multiple of n . On the other hand we have $\langle \alpha_i, z \rangle = 0$; hence using again that $\langle n, \alpha_i \rangle \neq 0$ we obtain $z = 0$. This shows $\ker(\text{id} - \tilde{w}) = 0$.

As we had seen above, $n_{w,i}$ is a vector normal to $\text{ran}(\text{id} - \sigma)$ and hence is a multiple of n . By [Lemma 3.2](#), it is a positive multiple if and only if $\langle n, \alpha_i \rangle > 0$. But then $\langle n, \alpha_{i'} \rangle < 0$, and so $n_{w',i'}$ is a negative multiple of n . This shows that V_w and $V_{w'}$ are on opposite sides of the hyperplane $\text{ran}(\text{id} - \sigma)$. \square

Consider the union

$$(2) \quad X := \bigcup_{w \in W} V_w.$$

over $W \subset W^{\text{aff}}$. Thus $\bigcup_{w \in W^{\text{aff}}} V_w = \bigcup_{\lambda \in \Lambda} (\lambda + X)$. The statement of [Theorem 1.2](#) means in particular that X is a fundamental domain for the action of Λ . [Figure 1](#) and [Figure 2](#) give pictures of X for the root systems \mathbf{B}_2 and \mathbf{G}_2 . The shaded regions are the top-dimensional polytopes (that is, the sets V_w for $\text{id} - w$ invertible), the dark lines are the 1-dimensional polytopes (corresponding to reflections), and the origin corresponds to $w = \text{id}$.

Proposition 3.4. (a) *The sets $\lambda + \text{int}(\bar{X})$ for $\lambda \in \Lambda$ are disjoint, and*

$$\bigcup_{\lambda \in \Lambda} \lambda + \bar{X} = V.$$




Figure 2. The set X for the root system G_2 .

(b) *The open polytopes V_w for $w \in W_{\text{reg}}^{\text{aff}}$ are disjoint, and*

$$\bigcup_{w \in W_{\text{reg}}^{\text{aff}}} \overline{V}_w = V.$$

Proof. Since the collection of closed polytopes \overline{V}_w for $w \in W_{\text{reg}}$ is locally finite, the union $V' := \bigcup_{w \in W_{\text{reg}}^{\text{aff}}} \overline{V}_w$ is a closed polyhedral subset of V . Proposition 3.3 shows that a point $\zeta \in V_{w,i}$ cannot contribute to the boundary of this subset unless it lies in $\bigcup_{\sigma \in W^{\text{aff}}} \bigcup_{|I| \geq 2} V_{\sigma,I}$. We therefore see that the boundary has codimension at least 2, and hence is empty since V' is a closed polyhedron. This proves $V' = V$, and also $\bigcup_{\lambda \in \Lambda} (\lambda + \overline{X}) = V$ with X as defined in (2). Hence the volume $\text{vol}(X)$ (for the Riemannian measure on V defined by the inner product) must be at least the volume of a fundamental domain for the action of Λ :

$$(3) \qquad \text{vol}(X) \geq |W| \text{vol}(A).$$

On the other hand, $\text{vol}(V_w) = \text{vol}((\text{id} - w)(A)) = \det(\text{id} - w) \text{vol}(A)$, so

$$(4) \qquad \text{vol}(X) \leq \sum_{w \in W} \text{vol}(V_w) = \text{vol}(A) \sum_{w \in W} \det(\text{id} - w) = |W| \text{vol}(A),$$

where we used that $\sum_{w \in W} \det(\text{id} - w) = |W|$ from [Bourbaki 1975, page 134]. This confirms $\text{vol}(X) = |W| \text{vol}(A)$. It follows that the sets $\lambda + \text{int}(\overline{X})$ are pairwise disjoint, or else the inequality (3) would be strict. Similarly that the sets V_w for $w \in W_{\text{reg}}$ are disjoint, or else the inequality (4) would be strict. (Of course, this also follows from Waldspurger’s Theorem 1.1 since $C_w \subset D_w$.) Hence all V_w for $w \in W_{\text{reg}}^{\text{aff}}$ are disjoint. □

To proceed, we quote the following result from Waldspurger’s paper, where it is stated in greater generality .

Proposition 3.5 [Waldspurger 2007, Lemme]. *Given $w \in W$ and a proper subset $I \subset \{0, \dots, l\}$, there exists a unique $q \in W_I$ such that*

$$\ker(\text{id} - wq) \cap \{x \in V \mid \langle \alpha_i, x \rangle > 0 \text{ for all } i \in I\} \neq \emptyset.$$

Following [Waldspurger 2007] we use this to prove,

Proposition 3.6. *Every element of V is contained in some V_w for $w \in W^{\text{aff}}$:*

$$(5) \quad \bigcup_{w \in W^{\text{aff}}} V_w = V.$$

Proof. Let $\zeta \in V$ be given. Pick $w \in W_{\text{reg}}^{\text{aff}}$ with $\zeta \in \bar{V}_w$, and let $I \subset \{0, \dots, l\}$ with $\zeta \in V_{w,I}$. Then $x := (\text{id} - w)^{-1}(\zeta) \in A_I$ is fixed under W_I^{aff} . By Proposition 3.5 we may choose $\tilde{q} \in W_I$ and $n \in V$ such that

- (a) $\tilde{w}\tilde{q}(n) = n$,
- (b) $\langle \alpha_i, n \rangle > 0$ for all $i \in I$.

Taking $\|n\|$ sufficiently small, we have $x + n \in A$, and

$$(\text{id} - wq)(x + n) = (\text{id} - wq)(x) + (\text{id} - \tilde{w}\tilde{q})n = (\text{id} - w)(x) = \zeta.$$

This shows $\zeta \in V_{wq}$. □

4. Disjointness of the sets $\lambda + X$

To finish the proof of Theorem 1.2, we have to show that the union (5) is disjoint. Waldspurger's Theorem 1.1 shows that all $D_w = (\text{id} - w)(C)$ for $w \in W$ are disjoint. (We refer to his paper for a very simple proof of this fact.) Hence the same is true of $V_w \subset D_w$ for $w \in W$. It remains to show that the sets $\lambda + X$ for $\lambda \in \Lambda$, with X given by (2), are disjoint.

The closure $\bar{X} = \bigcup_{w \in W} \bar{V}_w$ only involves the top-dimensional polytopes:

Lemma 4.1. *The closure of the set X is the union $\bar{X} = \bigcup_{w \in W_{\text{reg}}} \bar{V}_w$. Furthermore, $\text{int}(\bar{X}) = \text{int}(X)$.*

Proof. We must show that for any $\zeta \in \bar{V}_\sigma$ with $\sigma \in W \setminus W_{\text{reg}}$, there exists a $w \in W_{\text{reg}}$ such that $\zeta \in \bar{V}_w$. Using induction, it suffices to find $\sigma' \in W$ such that $\zeta \in \bar{V}_{\sigma'}$ and $\dim(\ker(\text{id} - \sigma')) = \dim(\ker(\text{id} - \sigma)) - 1$. Let $\pi : V \rightarrow \ker(\text{id} - \sigma)^\perp = \text{ran}(\text{id} - \sigma)$ denote the orthogonal projection. Then $\text{id} - \sigma$ restricts to an invertible transformation of $\pi(V)$, and \bar{V}_σ is the image of $\pi(\bar{A})$ under this transformation. We have

$$\pi(\bar{A}) = \pi(\partial \bar{A}) = \bigcup_{i=0}^l \pi(\bar{A}_i),$$

and this continues to hold if we remove the index $i = 0$ from the right side, as well as all indices i for which $\dim \pi(A_i) < \dim \pi(V)$. That is, for each point $x \in \pi(\bar{A})$ there exists an index $i \neq 0$ such that $x \in \pi(\bar{A}_i)$, with $\dim \pi(A_i) = \dim \pi(V)$. Taking x to be the preimage of ζ under $(\text{id} - \sigma)|_{\pi(V)}$, we have $\zeta \in \bar{V}_{\sigma,i}$ with $i \neq 0$ and $\dim V_{\sigma,i} = \dim \text{ran}(\text{id} - \sigma)$. Let $\sigma' = \sigma s_i \in W$. Then $V_{\sigma,i} = V_{\sigma',i}$; hence $\dim(\text{ran}(\text{id} - \sigma')) \geq \dim V_{\sigma,i} = \dim(\text{ran}(\text{id} - \sigma))$, which shows $\dim \ker(\text{id} - \sigma') \leq \dim \ker(\text{id} - \sigma)$. By elementary properties of reflection groups, the dimensions of the fixed point sets of σ and σ' differ by either $+1$ or -1 . Hence $\dim(\ker(\text{id} - \sigma')) = \dim(\ker(\text{id} - \sigma)) - 1$, proving the first assertion of the lemma.

It follows in particular that the closure of $\text{int}(\bar{X})$ equals that of X . Suppose $\zeta \in \text{int}(\bar{X})$. By [Proposition 3.6](#) there exists $\lambda \in \Lambda$ with $\zeta \in \lambda + X$. It follows that $\text{int}(\bar{X})$ meets $\lambda + X$, and hence also meets $\lambda + \text{int}(\bar{X})$. Since the Λ -translates of $\text{int}(\bar{X})$ are pairwise disjoint (see [Proposition 3.4](#)), it follows that $\lambda = 0$, that is, $\zeta \in X$. This shows $\zeta \in X \cap \text{int}(\bar{X}) = \text{int}(X)$; hence $\text{int}(\bar{X}) \subset \text{int}(X)$. The opposite inclusion is obvious. \square

Since we already know that the sets $\lambda + \text{int}(X)$ are disjoint, we are interested in $X \setminus \text{int}(X) \subset \partial X = \bar{X} \setminus \text{int}(X)$. Let us call a closed codimension 1 boundary face of the polyhedron \bar{X} *horizontal* if its supporting hyperplane contains $V_{w,0}$ for some $w \in W_{\text{reg}}$, and *vertical* if its supporting hyperplane contains $V_{w,i}$ for some $w \in W_{\text{reg}}$ and $i \neq 0$. These two cases are exclusive:

Lemma 4.2. *Let n be the inward-pointing normal vector to a codimension 1 face of \bar{X} . Then $\langle n, \alpha_{\max} \rangle \neq 0$. In fact, $\langle n, \alpha_{\max} \rangle < 0$ for the horizontal faces and $\langle n, \alpha_{\max} \rangle > 0$ for the vertical faces.*

Proof. Given a codimension 1 boundary face of \bar{X} , pick any point ζ in that boundary face not lying in $\bigcup_{w \in W^{\text{aff}}} \bigcup_{|I| \geq 2} V_{w,I}$. Let $w \in W_{\text{reg}}$ and $i \in \{0, \dots, l\}$ such that $\zeta \in V_{w,i}$ and $n_{w,i}$ is an inward-pointing normal vector. By [Proposition 3.3](#), there is a unique $i' \neq i$ such that $\zeta \in V_{w',i'}$, where $w' = w s_i s_{i'}$. Since V_w and $V_{w'}$ lie on opposite sides of the affine hyperplane spanned by $V_{w,i}$, and ζ is a boundary point of \bar{X} , we have $w' \notin W$. Thus one of i and i' must be zero. If $i = 0$ (so that the given boundary face is horizontal) we obtain $\langle n_{w,0}, \alpha_{\max} \rangle = -\langle n_{w,0}, \alpha_0 \rangle < 0$. If $i' = 0$ we similarly obtain $\langle n_{w',0}, \alpha_{\max} \rangle < 0$; hence $\langle n_{w,i}, \alpha_{\max} \rangle > 0$. \square

Lemma 4.3. *Let $\zeta \in X \setminus \text{int}(X)$. Then there exists a vertical boundary face of \bar{X} containing ζ . Equivalently, the complement $\partial \bar{X} \setminus (X \setminus \text{int}(X))$ is contained in the union of horizontal boundary faces.*

Proof. The alcove A is invariant under multiplication by any scalar in $(0, 1)$. Hence, the same is true for the sets V_w for $w \in W$, as well as for X and $\text{int}(X)$. Hence, if $\zeta \in X \setminus \text{int}(X)$ there exists $t_0 > 1$ such that $t\zeta \in X \setminus \text{int}(X)$ for $1 \leq t < t_0$. The closed codimension 1 boundary face containing this line segment is necessarily vertical,

since a line through the origin intersects the affine hyperplane $\{x \mid \langle n_{w,0}, x - \zeta \rangle = 0\}$ in at most one point. \square

Proposition 4.4. *For any $\zeta \in X$, there exists $\epsilon > 0$ such that $\zeta + s\alpha_{\max} \in \text{int}(X)$ for $0 < s < \epsilon$.*

Proof. If $\zeta \in \text{int}(X)$ there is nothing to show; hence suppose $\zeta \in X \setminus \text{int}(X)$. Suppose first that ζ is not in the union of horizontal boundary faces of \bar{X} . Then there exists an open neighborhood U of ζ such that $U \cap X = U \cap \bar{X}$. All boundary faces of \bar{X} meeting ζ are vertical, and their inward-pointing normal vectors n all satisfy $\langle n, \alpha_{\max} \rangle > 0$. Hence, $\zeta + s\alpha_{\max} \in \text{int}(U \cap \bar{X}) = \text{int}(U \cap X) \subset X$ for $s > 0$ sufficiently small.

For the general case, suppose by way of contradiction that for all $\epsilon > 0$, there is $s \in (0, \epsilon)$ with $\zeta + s\alpha_{\max} \notin \text{int}(X)$. Since ζ is contained in some vertical boundary face, one can choose $t > 1$ so that $\zeta' := t\zeta \in X \setminus \text{int}(X)$, but ζ' is not in the closure of the union of horizontal boundary faces. Given $\epsilon > 0$, pick $s \in (0, \epsilon)$ such that $\zeta + (s/t)\alpha_{\max} \notin \text{int}(X)$. Since $\text{int}(X)$ is invariant under multiplication by scalars in $(0, 1)$, the complement $V \setminus \text{int}(X)$ is invariant under multiplication by scalars in $(1, \infty)$; hence we obtain $\zeta' + s\alpha_{\max} \notin \text{int}(X)$. This contradicts what we have shown above, and completes the proof. \square

Proposition 4.5. *The sets $\lambda + X$ for $\lambda \in \Lambda$ are disjoint.*

Proof. Suppose $\zeta \in (\lambda + X) \cap (\lambda' + X)$. By [Proposition 4.4](#), we can choose $s > 0$ so that $\zeta + s\alpha_{\max} \in (\lambda + \text{int}(X)) \cap (\lambda' + \text{int}(X))$. Since the Λ -translates of $\text{int}(X)$ are disjoint, it follows that $\lambda = \lambda'$. \square

This completes the proof of [Theorem 1.2](#). We conclude with some remarks on the properties of the decomposition $V = \bigcup_{w \in W^{\text{aff}}} V_w$.

Remarks 4.6. (a) The group of symmetries τ of the extended Dynkin diagram (that is, the outer automorphisms of the corresponding affine Lie algebra) acts by symmetries of the decomposition $V = \bigcup_{w \in W^{\text{aff}}} V_w$, as follows. Identify the nodes of the extended Dynkin diagram with the simple affine reflections s_0, \dots, s_l . Then τ extends to a group automorphism of W^{aff} , taking s_i to $\tau(s_i)$. This automorphism is implemented by a unique Euclidean transformation $g: V \rightarrow V$, that is,

$$gwg^{-1} = \tau(w) \quad \text{for all } w \in W^{\text{aff}}.$$

Then g preserves A , and consequently

$$g V_w = g(\text{id} - w)(A) = (\text{id} - \tau(w))(A) = V_{\tau(w)} \quad \text{for } w \in W^{\text{aff}}.$$

(b) It is immediate from the definition that $-w: V \rightarrow V$, $x \mapsto -wx$ takes $V_{w^{-1}}$ into V_w :

$$-w(V_{w^{-1}}) = V_w.$$

(c) For any positive root α , let s_α be the corresponding reflection. Then

$$(\text{id} - s_\alpha)(\xi) = \langle \alpha, \xi \rangle \alpha^\vee,$$

where α^\vee is the coroot corresponding to α . Hence D_{s_α} is the relative interior of the line segment from 0 to $\lambda \alpha^\vee$, where λ is the maximum value of the linear functional $\xi \mapsto \langle \alpha, \xi \rangle$ on the closed alcove \bar{A} . This maximum is achieved at one of the vertices. Let $\varpi_1^\vee, \dots, \varpi_l^\vee$ be the fundamental coweights, defined by $\langle \alpha_i, \varpi_j^\vee \rangle = \delta_{ij}$ for $i, j = 1, \dots, l$. Let $c_i \in \mathbb{N}$ be the coefficients of α_{\max} relative to the simple roots: $\alpha_{\max} = \sum_{i=1}^l c_i \alpha_i$. Then the nonzero vertices of A are ϖ_i^\vee / c_i . Similarly let $a_i \in \mathbb{Z}_{\geq 0}$ be the coefficients of α , so that $\alpha = \sum_{i=1}^l a_i \alpha_i$. Then the value of α at the i -th vertex of \bar{A} is a_i / c_i , and λ is the maximum of those values. There are two interesting cases: First, if $\alpha = \alpha_{\max}$, then all $a_i / c_i = 1$, and $\alpha^\vee = \alpha$. That is, the open line segment from the origin to the highest root always appears in the decomposition. Second, if $\alpha = \alpha_i$, then $a_i = 1$ while all other a_j vanish. In this case, one obtains the open line segment from the origin to $(1/c_i) \alpha_i^\vee$.

(d) Every V_w contains a distinguished ‘base point’. Indeed, let $\rho \in V$ be the half-sum of positive roots, and $h^\vee = 1 + \langle \alpha_{\max}, \rho \rangle$ the dual Coxeter number. Then $\rho / h^\vee \in A$, and consequently $\rho / h^\vee - w(\rho / h^\vee) \in V_w$.

5. Proof of Theorem 1.3

The proof is very similar to the proof of Proposition 3.4; hence we will be brief. Each $V_w^{(S)} = (S - w)(A)$ is the interior of a simplex in V , with codimension 1 faces $V_{w,i}^{(S)} = (S - w)(A_i)$. As in the proof of Lemma 3.1, we see that

$$n_{w,i}^{(S)} = (S - \tilde{w}^{-1})^{-1} \alpha_i$$

is an inward-pointing normal vector to the face $V_{w,i}^{(S)}$. For $S = 0$ this simplifies to $n_{w,i}^{(0)} = -w \alpha_i$. If $w' = w s_i$, then $V_{w,i}^{(S)} = V_{w',i}^{(S)}$, so that $n_{w,i}^{(S)}$ and $n_{w',i}^{(S)}$ are proportional. Since $n_{w,i}^{(0)} = -n_{w',i}^{(0)}$, continuity implies that $n_{w,i}^{(S)}$ is a negative multiple of $n_{w',i}^{(S)}$. As a consequence, we see that $V_w^{(S)}$ and $V_{w'}^{(S)}$ are on opposite sides of affine hyperplane supporting $V_{w,i}^{(S)} = V_{w',i}^{(S)}$. Arguing as in the proof of Proposition 3.4, this shows that

$$\bigcup_{w \in W^{\text{aff}}} \bar{V}_w^{(S)} = V.$$

Letting $X^{(S)} = \bigcup_{w \in W} V_w^{(S)}$, it follows that $V = \bigcup_{\lambda \in \Lambda} (\lambda + \bar{X}^{(S)})$. Therefore we have $\text{vol}(X^{(S)}) \geq |W| \text{vol}(A)$. But

$$\begin{aligned} \text{vol}(X^{(S)}) &\leq \sum_{w \in W} \text{vol}((S - w)(A)) \\ &= \text{vol}(A) \sum_{w \in W} |\det(S - w)| \\ &= \text{vol}(A) \sum_{w \in W} \det(\text{id} - Sw^{-1}) = |W| \text{vol}(A), \end{aligned}$$

using [Bourbaki 1975, page 134]. It follows that $\text{vol}(X^{(S)}) = |W| \text{vol}(A)$, which implies (as in the proof of Proposition 3.4) that all $\text{int}(\bar{V}_w^{(S)}) = V_w^{(S)}$ are disjoint. This completes the proof. \square

Remark 5.1. Theorem 1.3 and its proof go through for any S in the component of 0 in the set $\{S \in \text{End}(V) \mid \det(S - w) \neq 0 \text{ for all } w \in W\}$. For instance, the fact that $\det(\text{id} - Sw^{-1}) > 0$ follows by continuity from $S = 0$. On the other hand, the result becomes false if, for example, S is a positive matrix with $S > 2 \text{id}$, since then $\sum_{w \in W} |\det(S - w)| = \sum_{w \in W} \det(S - w) = \det(S)|W|$; see [Bourbaki 1975, page 134].

Acknowledgments

I would like to thank Bert Kostant for telling me about Waldspurger's result, and the referee for helpful comments.

References

- [Bourbaki 1975] N. Bourbaki, *Éléments de mathématique, Fasc. XXXVIII: Groupes et algèbres de Lie. Chapitre VII: Sous-algèbres de Cartan, éléments réguliers. Chapitre VIII: Algèbres de Lie semi-simples déployées*, Actualités Scientifiques et Industrielles **1364**, Hermann, Paris, 1975. [MR 56 #12077](#) [Zbl 0329.17002](#)
- [Waldspurger 2007] J.-L. Waldspurger, “Une remarque sur les systèmes de racines”, *J. Lie Theory* **17**:3 (2007), 597–603. [MR 2008i:20051](#) [Zbl 1133.20029](#)

Received January 12, 2009. Revised February 4, 2009.

ECKHARD MEINRENKEN
UNIVERSITY OF TORONTO
DEPARTMENT OF MATHEMATICS
40 ST. GEORGE STREET
TORONTO, ONTARIO M4S 2E4
CANADA

mein@math.toronto.edu