

*Pacific
Journal of
Mathematics*

**STRONG KÄHLER WITH TORSION STRUCTURES
FROM ALMOST CONTACT MANIFOLDS**

MARISA FERNÁNDEZ, ANNA FINO,
LUIS UGARTE AND RAQUEL VILLACAMPA

Volume 249 No. 1

January 2011

STRONG KÄHLER WITH TORSION STRUCTURES FROM ALMOST CONTACT MANIFOLDS

MARISA FERNÁNDEZ, ANNA FINO,
LUIS UGARTE AND RAQUEL VILLACAMPA

For an almost contact metric manifold N , we find conditions under which either the total space of an S^1 -bundle over N or the Riemannian cone over N admit a strong Kähler with torsion (SKT) structure. In so doing, we construct new 6-dimensional SKT manifolds. Moreover, we study the geometric structure induced on a hypersurface of an SKT manifold and use it to construct new SKT manifolds via appropriate evolution equations. We also study hyper-Kähler with torsion (HKT) structures on the total space of an S^1 -bundle over manifolds with three almost contact structures.

1. Introduction

On any Hermitian manifold (M^{2n}, J, h) there exists a unique Hermitian connection ∇^B with totally skew-symmetric torsion, which is called the Bismut connection after [Bismut 1989]. The torsion 3-form $h(X, T^B(Y, Z))$ of ∇^B can be identified with the 3-form

$$-JdF(\cdot, \cdot, \cdot) = -dF(J\cdot, J\cdot, J\cdot),$$

where $F(\cdot, \cdot) = h(\cdot, J\cdot)$ is the fundamental 2-form associated to the Hermitian structure (J, h) .

Hermitian structures with closed JdF are called *strong Kähler with torsion* (in short, SKT) or *pluriclosed* [Egidi 2001]. Since $\partial\bar{\partial}$ acts as $\frac{1}{2}dJd$ on forms of bidegree $(1, 1)$, the latter condition is equivalent to $\partial\bar{\partial}F = 0$. SKT structures have been recently studied by many authors, and they also have applications in type II string theory and in 2-dimensional supersymmetric σ -models [Gates et al. 1984; Strominger 1986; Ivanov and Papadopoulos 2001].

The class of SKT metrics includes of course the Kähler metrics, but as in [Fino et al. 2004], we are interested on non-Kähler geometry, so by an *SKT metric* we

MSC2000: primary 53C55; secondary 53C26, 22E25, 53C15.

Keywords: Hermitian metric, hypercomplex structure, torsion, almost contact structure, Riemannian cone.

Partially supported through Project MICINN (Spain) MTM2008-06540-C02-01/02, Project MIUR “Riemannian Metrics and Differentiable Manifolds” and by GNSAGA of INdAM.

will mean a Hermitian metric h whose fundamental 2-form F is $\partial\bar{\partial}$ -closed but not d -closed.

Gauduchon [1984] showed that on a compact complex surface, an SKT metric can be found in the conformal class of any given Hermitian metric, but in higher dimensions the situation is more complicated.

SKT structures on 6-dimensional nilmanifolds, that is, on compact quotients of nilpotent Lie groups by discrete subgroups, were classified in [Fino et al. 2004; Ugarte 2007]. Simply connected examples of 6-dimensional SKT manifolds have been found in [Grantcharov et al. 2008] by using torus bundles, and recently Swann [2010] has reproduced them via the twist construction, by extending them to higher dimensions and finding new compact simply connected SKT manifolds. Moreover, Fino and Tomassini [2009] showed that the SKT condition is preserved by the blow-up construction.

The odd-dimensional analogue of a Hermitian structure is given by a normal almost contact metric structure. Indeed, on the product $N^{2n+1} \times \mathbb{R}$ of a $(2n+1)$ -dimensional almost contact metric manifold N^{2n+1} by the real line \mathbb{R} , it is possible to define a natural almost complex structure, which is integrable if and only if the almost contact metric structure on N^{2n+1} is normal [Sasaki and Hatakeyama 1961]. More generally, it is possible to construct Hermitian manifolds starting from an almost contact metric manifold N^{2n+1} by considering a principal fiber bundle P with base space N^{2n+1} and structural group S^1 , that is, an S^1 -bundle over N^{2n+1} ; see [Ogawa 1963]. Indeed, by using the almost contact metric structure on N^{2n+1} and the connection 1-form θ , Ogawa constructed an almost Hermitian structure (J, h) on P and found conditions under which J is integrable and (J, h) is Kähler.

In Section 2, we determine in Theorem 2.3 general conditions under which an S^1 -bundle over an almost contact metric $(2n+1)$ -dimensional manifold N^{2n+1} is SKT. We study the particular case when N^{2n+1} is quasi-Sasakian, that is, when it has an almost contact metric structure for which the fundamental form is closed (Corollary 2.4). In this way, we are able to construct some new 6-dimensional SKT examples, starting from 5-dimensional quasi-Sasakian Lie algebras, and also from Sasakian ones.

A Sasakian structure can be also seen as the analogue, in odd dimensions, of a Kähler structure. Indeed, by [Boyer and Galicki 1999], a Riemannian manifold (N^{2n+1}, g) of odd dimension $2n+1$ admits a compatible Sasakian structure if and only if the Riemannian cone $N^{2n+1} \times \mathbb{R}^+$ is Kähler. In Section 3, Theorem 3.1 gives the conditions that must be satisfied by the compatible almost contact metric structure on a Riemannian manifold (N^{2n+1}, g) in order that the Riemannian cone $N^{2n+1} \times \mathbb{R}^+$ be SKT. We provide an example of an SKT manifold constructed as a Riemannian cone, and in Section 4 we consider the case when the Riemannian cone is 6-dimensional. This case is interesting since one can impose that the SKT

structure is in addition an SKT $SU(3)$ -structure, and one can find relations with the $SU(2)$ -structures studied by Conti and Salamon [2007].

In Section 5, we study the geometric structure induced naturally on any oriented hypersurface N^{2n+1} of a $(2n+2)$ -dimensional manifold M^{2n+2} carrying an SKT structure, and in Section 6, we use such structures in Theorem 6.4 to construct new SKT manifolds via appropriate evolution equations [Hitchin 2001; Conti and Salamon 2007] starting from a 5-dimensional manifold endowed with an $SU(2)$ -structure.

A good quaternionic analogue of Kähler geometry is given by *hyper-Kähler with torsion* (in short, HKT) geometry. An HKT manifold is a hyper-Hermitian manifold $(M^{4n}, J_1, J_2, J_3, h)$ admitting a hyper-Hermitian connection with totally skew-symmetric torsion, that is, one in which the three Bismut connections associated with the three Hermitian structures (J_r, h) coincide for $r = 1, 2, 3$. This geometry was introduced by Howe and Papadopoulos [1996] and later studied in [Grantcharov and Poon 2000; Fino and Grantcharov 2004; Barberis et al. 2009; Barberis and Fino 2008; Swann 2010].

In the interesting special case in which the torsion 3-form of such a hyper-Hermitian connection is closed, the HKT manifold is called *strong*.

In Section 7, Theorem 7.1 gives conditions under which an S^1 -bundle over a $(4n+3)$ -dimensional manifold endowed with three almost contact metric structures is HKT and in particular when it is strong HKT.

2. SKT structures arising from S^1 -bundles

Consider a $(2n+1)$ -manifold N^{2n+1} endowed with an almost contact metric structure (I, ξ, η, g) ; that is, I is a tensor field of type $(1, 1)$, ξ is a vector field, η is a 1-form, and g is a Riemannian metric on N^{2n+1} , satisfying together the conditions

$$I^2 = -\text{Id} + \eta \otimes \xi, \quad \eta(\xi) = 1, \quad g(IU, IV) = g(U, V) - \eta(U)\eta(V)$$

for any vector fields U and V on N^{2n+1} . Denote by ω the fundamental 2-form of (I, ξ, η, g) ; that is, ω is the 2-form on N^{2n+1} given by

$$\omega(\cdot, \cdot) = g(\cdot, I\cdot).$$

Given the tensor field I , consider its Nijenhuis torsion $[I, I]$, defined by

$$(1) \quad [I, I](X, Y) = I^2[X, Y] + [IX, IY] - I[IX, Y] - I[X, IY].$$

On the product $N^{2n+1} \times \mathbb{R}$, one can define a natural almost complex structure

$$J\left(X, f \frac{d}{dt}\right) = \left(IX + f\xi, -\eta(X) \frac{d}{dt}\right),$$

where f is a \mathcal{C}^∞ -function on $N^{2n+1} \times \mathbb{R}$ and t is the coordinate on \mathbb{R} .

Definition 2.1 [Sasaki and Hatakeyama 1961]. *We call an almost contact metric structure (I, ξ, η, g) on N^{2n+1} normal if the almost complex structure J on $N^{2n+1} \times \mathbb{R}$ is integrable, or equivalently if*

$$[I, I](X, Y) + 2d\eta(X, Y)\xi = 0$$

for any vector fields X, Y on N^{2n+1} .

By [Blair 1967, Lemma 2.1], one has $i_\xi d\eta = 0$ for a normal almost contact metric structure (I, ξ, η, g) .

Remark 2.2. The normality of the almost contact structure implies as well that $I d\eta = d\eta$. Indeed, $d(\eta - i dt) = d\eta$ has no $(0, 2)$ -part and therefore has no $(2, 0)$ -part since $d\eta$ is real. Thus, $J d\eta = d\eta$, but $J d\eta = I d\eta$ as well since $i_\xi d\eta = 0$.

We recall that a Hermitian manifold (M, J, h) is SKT if and only if the 3-form $J dF$ is closed, where F is the fundamental 2-form of (J, h) . We will use the convention that J acts on r -forms β by

$$(J\beta)(X_1, \dots, X_r) = \beta(JX_1, \dots, JX_r) \quad \text{for any vector fields } X_1, \dots, X_r.$$

We now show general conditions under which an S^1 -bundle over an almost contact metric $(2n+1)$ -dimensional manifold is SKT.

Let (N^{2n+1}, I, ξ, η) be a $(2n+1)$ -dimensional almost contact manifold, and let Ω be a closed 2-form on N^{2n+1} that represents an integral cohomology class on N^{2n+1} . From the well-known result of Kobayashi [1956], we can consider the circle bundle $S^1 \hookrightarrow P \rightarrow N^{2n+1}$ and the connection 1-form θ on P whose curvature form is $d\theta = \pi^*(\Omega)$, where $\pi : P \rightarrow N^{2n+1}$ is the projection.

By using the almost contact structure (I, ξ, η) and the connection 1-form θ , one can define an almost complex structure J on P as follows [Ogawa 1963]. For any right-invariant vector field X on P , the vector field JX is given by

$$(2) \quad \theta(JX) = -\pi^*(\eta(\pi_*X)) \quad \text{and} \quad \pi_*(JX) = I(\pi_*X) + \tilde{\theta}(X)\xi,$$

where $\tilde{\theta}(X)$ is the unique function on N^{2n+1} such that

$$(3) \quad \pi^*\tilde{\theta}(X) = \theta(X).$$

This definition can be extended to an arbitrary vector field X on P since X can be written in the form $X = \sum_j f_j X_j$, with f_j smooth functions on P , and X_j right-invariant vector fields. Then $JX = \sum_j f_j JX_j$.

Ogawa [1963] showed that when (N^{2n+1}, I, ξ, η) is normal, the almost complex structure J on P defined by (2) is integrable if and only if $d\theta$ is J -invariant, that is, if $J(d\theta) = d\theta$ or equivalently $d\theta(JX, Y) + d\theta(X, JY) = 0$ for any vector fields X and Y on P . That is, $d\theta$ is a complex 2-form on P having bidegree $(1, 1)$ with respect to J .

In terms of the 2-form Ω , whose lift to P is the curvature of the circle bundle $S^1 \hookrightarrow P \rightarrow N^{2n+1}$, the previous condition means that Ω is I -invariant, that is, $I(\Omega) = \Omega$. Therefore $i_\xi \Omega = 0$.

If $\{e^1, \dots, e^{2n}, \eta\}$ is an adapted coframe on a neighborhood U on N^{2n+1} , that is,

$$Ie^{2j-1} = -e^{2j} \quad \text{and} \quad Ie^{2j} = e^{2j-1} \quad \text{for } 1 \leq j \leq n,$$

then we can take $\{\pi^*e^1, \dots, \pi^*e^{2n}, \pi^*\eta, \theta\}$ as a coframe in $\pi^{-1}(U)$. By using the coframe $\{\pi^*e^1, \dots, \pi^*e^{2n}\}$, we may write

$$d\theta = \pi^*\alpha + \pi^*\beta \wedge \pi^*\eta,$$

where α is a 2-form in $\wedge^2\langle e^1, \dots, e^{2n} \rangle$, and $\beta \in \wedge^1\langle e^1, \dots, e^{2n} \rangle$.

Suppose that N^{2n+1} has a normal almost contact metric structure (I, ξ, η, g) . We consider a principal S^1 -bundle P with base space N^{2n+1} and connection 1-form θ , and endow P with the almost complex structure J (associated to θ) defined by (2). Since N^{2n+1} has a Riemannian metric g , a Riemannian metric h on P compatible with J (see [Ogawa 1963]) is given by

$$(4) \quad h(X, Y) = \pi^*g(\pi_*X, \pi_*Y) + \theta(X)\theta(Y)$$

for any right-invariant vector fields X and Y . This definition can be extended to any vector field on P .

Theorem 2.3. *Consider a $(2n+1)$ -dimensional almost contact metric manifold $(N^{2n+1}, I, \xi, \eta, g)$, and let Ω be a closed 2-form on N^{2n+1} that represents an integral cohomology class. Consider the circle bundle $S^1 \hookrightarrow P \rightarrow N^{2n+1}$ with connection 1-form θ , whose curvature form is $d\theta = \pi^*(\Omega)$ for the projection $\pi : P \rightarrow N^{2n+1}$.*

The almost Hermitian structure (J, h) on P defined by (2) and (4) is SKT if and only if (I, ξ, η, g) is normal, $d\theta$ is J -invariant, and

$$(5) \quad \begin{aligned} d(\pi^*(I(i_\xi d\omega))) &= 0, \\ d(\pi^*(I(d\omega) - d\eta \wedge \eta)) &= (-\pi^*(I(i_\xi d\omega)) + \pi^*\Omega) \wedge \pi^*\Omega, \end{aligned}$$

where ω is the fundamental form of the almost contact metric structure (I, ξ, η, g) .

Proof. As we mentioned, a result of Ogawa [1963] asserts that the almost complex structure J is integrable if and only if (I, ξ, η, g) is normal and $J(d\theta) = d\theta$. Thus, (J, h) is SKT if and only if the 3-form JdF is closed. Using the first equality in (2), we find that the fundamental 2-form F on P is

$$\begin{aligned} F(X, Y) &= h(X, JY) \\ &= \pi^*g(\pi_*X, \pi_*JY) + \theta(X)\theta(JY) \\ &= \pi^*g(\pi_*X, \pi_*JY) - \theta(X)\pi^*\eta(\pi_*Y). \end{aligned}$$

Therefore, taking into account that we are working with a circle bundle (whose fiber is thus 1-dimensional), we have

$$\begin{aligned} F &= \pi^* \omega + \pi^* \eta \wedge \theta, \\ dF &= \pi^*(d\omega) + \pi^*(d\eta) \wedge \theta - \pi^* \eta \wedge d\theta, \\ (6) \quad JdF &= J(\pi^*(d\omega)) - J(\pi^*(d\eta)) \wedge \pi^* \eta - \theta \wedge d\theta \end{aligned}$$

since $J(\pi^* \eta) = \theta$ and J is integrable, and so $J(d\theta) = d\theta$. Moreover,

$$(7) \quad J(\pi^*(d\omega)) = \pi^*(I(d\omega)) + \pi^*(I(i_\xi d\omega)) \wedge \theta.$$

Indeed, locally and in terms of the adapted basis $\{e^1, \dots, e^{2n+1}\}$ with

$$Ie^{2j-1} = -e^{2j} \quad \text{for } 1 \leq j \leq n, \quad Ie^{2n+1} = 0, \quad \text{and} \quad \eta = e^{2n+1},$$

we can write $d\omega = \alpha + \beta \wedge \eta$, where the local forms $\alpha \in \bigwedge^3 \langle e^1, \dots, e^{2n} \rangle$ and $\beta \in \bigwedge^2 \langle e^1, \dots, e^{2n} \rangle$ are generated only by e^1, \dots, e^{2n} . Furthermore, we have $I\alpha = I(d\omega)$ and $\beta = i_\xi d\omega$. Thus

$$J(\pi^*(d\omega)) = J(\pi^*(\alpha)) + J(\pi^*(i_\xi d\omega)) \wedge \theta.$$

Now, by using (2) and (3), we see that $J(\pi^*(\alpha)) = \pi^*(I\alpha)$ and $J(\pi^*(i_\xi d\omega)) = \pi^*(I(i_\xi d\omega))$, which proves (7). As a consequence of Remark 2.2,

$$(8) \quad J(\pi^*(d\eta)) = \pi^*(I(d\eta)) - \pi^*(I(i_\xi d\eta)) \wedge \theta = \pi^*(d\eta)$$

since $i_\xi d\eta = 0$ and $I d\eta = d\eta$.

Using (7) and (8), we get

$$(9) \quad JdF = \pi^*(I(d\omega)) + \pi^*(I(i_\xi d\omega)) \wedge \theta - \pi^*(d\eta) \wedge \pi^* \eta - \theta \wedge d\theta.$$

Therefore,

$$\begin{aligned} d(JdF) &= d(\pi^*(I(d\omega))) + d(\pi^*(I(i_\xi d\omega))) \wedge \theta + \pi^*(I(i_\xi d\omega)) \wedge d\theta \\ &\quad - d(\pi^*(d\eta)) \wedge \pi^* \eta - \pi^*(d\eta) \wedge d\pi^* \eta - d\theta \wedge d\theta. \end{aligned}$$

Consequently, $d(JdF) = 0$ if and only if

$$\begin{aligned} d(\pi^*(I(i_\xi d\omega))) &= 0, \\ d(\pi^*(I(d\omega)) - d\eta \wedge \eta) &= (\pi^*(-I(i_\xi d\omega)) + d\theta) \wedge d\theta. \quad \square \end{aligned}$$

An almost contact metric manifold $(N^{2n+1}, I, \xi, \eta, g)$ is *quasi-Sasakian* if it is normal and its fundamental form ω is closed. In particular, if $d\eta = \alpha \omega$, then the almost contact metric structure is called α -*Sasakian*. When $\alpha = -2$, the structure is said to be *Sasakian*.

By [Friedrich and Ivanov 2002, Theorem 8.2], an almost contact metric manifold $(N^{2n+1}, I, \xi, \eta, g)$ admits a connection ∇^c that preserves the almost contact metric

structure and has totally skew-symmetric torsion tensor if and only if the Nijenhuis tensor of I , given by (1), is skew-symmetric and ξ is a Killing vector field. This connection is unique.

In particular, on any quasi-Sasakian manifold $(N^{2n+1}, I, \xi, \eta, g)$ there exists a unique connection ∇^c with totally skew-symmetric torsion, such that

$$\nabla^c I = 0, \quad \nabla^c g = 0, \quad \nabla^c \eta = 0.$$

Such a connection ∇^c is uniquely determined by

$$(10) \quad g(\nabla_X^c Y, Z) = g(\nabla_X^g Y, Z) + \frac{1}{2}(d\eta \wedge \eta)(X, Y, Z),$$

where ∇^g is the Levi-Civita connection and $\frac{1}{2}(d\eta \wedge \eta)$ is the torsion 3-form of ∇^c .

Corollary 2.4. *Let $(N^{2n+1}, I, \xi, \eta, g)$ be a quasi-Sasakian $(2n+1)$ -manifold, and let Ω be a closed 2-form on N^{2n+1} that represents an integral cohomology class. Consider the circle bundle $S^1 \hookrightarrow P \rightarrow N^{2n+1}$ with connection 1-form θ whose curvature form is $d\theta = \pi^*(\Omega)$ for the projection $\pi : P \rightarrow N^{2n+1}$. The almost Hermitian structure (J, h) on P defined by (2) and (4) is SKT if and only if Ω is I -invariant, $i_\xi \Omega = 0$, and*

$$(11) \quad d\eta \wedge d\eta = -\Omega \wedge \Omega.$$

The Bismut connection ∇^B of (J, h) on P and the connection ∇^c on N given by (10) are related by

$$(12) \quad h(\nabla_X^B Y, Z) = \pi^* g(\nabla_{\pi_* X}^c \pi_* Y, \pi_* Z)$$

for any vector fields $X, Y, Z \in \text{Ker } \theta$.

Proof. Since $d\omega = 0$, if we impose the SKT condition, then we get by using the previous theorem the equation (11).

The Bismut connection ∇^B associated to the Hermitian structure (J, h) on P is

$$(13) \quad h(\nabla_X^B Y, Z) = h(\nabla_X^h Y, Z) - \frac{1}{2}dF(JX, JY, JZ)$$

for any vector fields X, Y, Z on P , where ∇^h is the Levi-Civita connection associated to h . Then, for any X, Y, Z in the kernel of θ , we have

$$h(\nabla_X^B Y, Z) = \pi^* g(\nabla_X^h Y, Z) + \frac{1}{2}(\pi^*(d\eta) \wedge \pi^*\eta)(X, Y, Z).$$

By [Ogawa 1963, Lemma 3] and the definition of ∇^c , we get

$$\begin{aligned} h(\nabla_X^B Y, Z) &= \pi^* g(\nabla_{\pi_* X}^g \pi_* Y, \pi_* Z) + \frac{1}{2}(\pi^*(d\eta) \wedge \pi^*\eta)(X, Y, Z) \\ &= \pi^* g(\nabla_{\pi_* X}^c \pi_* Y, \pi_* Z) \end{aligned}$$

for any X, Y, Z in the kernel of θ . □

Remark 2.5. If the structure (I, ξ, η, g) is α -Sasakian, equation (11) reads

$$\Omega \wedge \Omega = -\alpha^2 \omega \wedge \omega.$$

In the case of a trivial S^1 -bundle, that is, if we consider the natural almost Hermitian structure on the product $N^{2n+1} \times \mathbb{R}$, we get this:

Corollary 2.6. *Let $(N^{2n+1}, I, \xi, \eta, g)$ be a $(2n+1)$ -dimensional almost contact metric manifold. Impose on the product $N^{2n+1} \times \mathbb{R}$ the almost complex structure J given by*

$$JX = IX \quad \text{for } X \in \text{Ker } \eta \quad \text{and} \quad J\xi = -\frac{d}{dt},$$

and the metric h given by $h = g + (dt)^2$. The Hermitian structure (J, h) is SKT if and only if (I, ξ, η, g) is normal, $d(I(d\omega)) = d(d\eta \wedge \eta)$ and $d(I(i_\xi d\omega)) = 0$, where ω denotes the fundamental 2-form of the almost contact metric structure (g, I, ξ, η) .

Corollary 2.7. *Let $(N^{2n+1}, I, \xi, \eta, g)$ be a $(2n+1)$ -dimensional quasi-Sasakian manifold with $d\eta \wedge d\eta = 0$. The Hermitian structure (J, h) on $N^{2n+1} \times \mathbb{R}$ is SKT. Moreover, its Bismut connection ∇^B coincides with the unique connection ∇^c on N^{2n+1} given by (10).*

Proof. In this case, since $d\omega = 0$ we get $d(JdF) = -d(d\eta \wedge \eta)$. By using (12), we get $h(\nabla_X^B Y, Z) = g(\nabla_X^c Y, Z)$ for any vector fields X, Y, Z on N^{2n+1} . \square

2.1. Examples. We will present three examples of quasi-Sasakian Lie algebras satisfying the condition $d\eta \wedge d\eta = 0$. By applying Corollary 2.7, one gets an SKT structure on the product of the corresponding simply connected Lie group by \mathbb{R} .

Example 2.8. Let \mathfrak{s} be the 5-dimensional Lie algebra with structure equations

$$\begin{aligned} de^1 &= e^{13} + e^{23} + e^{25} - e^{34} + e^{35}, \\ de^2 &= 2e^{12} - 2e^{13} + e^{14} - e^{15} - e^{24} + e^{34} + e^{45}, \\ de^3 &= -e^{12} + e^{13} + e^{14} - e^{15} + 2e^{24} - 2e^{34} + e^{45}, \\ de^4 &= -e^{12} - e^{23} + e^{24} - e^{25} - e^{35}, \\ de^5 &= e^{12} - e^{13} - e^{24} + e^{34}, \end{aligned}$$

where $e^{ij} = e^i \wedge e^j$. On \mathfrak{s} , take the quasi-Sasakian structure (I, ξ, η, g) given by

$$(14) \quad \eta = e^5, \quad Ie^1 = -e^2, \quad Ie^3 = -e^4, \quad \omega = -e^{12} - e^{34}, \quad g = \sum_{j=1}^5 (e^j)^2.$$

This quasi-Sasakian structure satisfies the condition $d(d\eta \wedge \eta) = 0$. The Lie algebra \mathfrak{s} is 2-step solvable since the commutator

$$\mathfrak{s}^1 = [\mathfrak{s}, \mathfrak{s}] = \mathbb{R}(e_1 - e_4, e_2 + e_3, e_1 - e_2 + 2e_3 - e_5)$$

is abelian, where $\{e_1, \dots, e_5\}$ denotes the dual basis of $\{e^1, \dots, e^5\}$. Moreover, \mathfrak{s} has trivial center, is irreducible and nonunimodular, since the trace of ad_{e_1} is -3 .

Example 2.9. Consider the family of 2-step solvable Lie algebras \mathfrak{s}_a for $a \in \mathbb{R} - \{0\}$, given by

$$\begin{aligned} de^1 &= ae^{23} + 3e^{25}, & de^3 &= ae^{34}, \\ de^2 &= -ae^{13} - 3e^{15}, & de^4 &= 0, \\ & & de^5 &= -\frac{1}{3}a^2e^{34}. \end{aligned}$$

The almost contact metric structure (I, ξ, η, g) defined in (14) is quasi-Sasakian and satisfies the condition $d\eta \wedge d\eta = 0$. The second cohomology group of \mathfrak{s}_a is generated by e^{12} and e^{45} .

Example 2.10. Another family of quasi-Sasakian Lie algebras that satisfies the condition $d\eta \wedge d\eta = 0$ is \mathfrak{g}_b for $b \in \mathbb{R} - \{0\}$, with structure equations

$$\begin{aligned} de^1 &= b(e^{13} + e^{14} - e^{23} + e^{24}) + e^{25}, & de^3 &= 2e^{45}, \\ de^2 &= b(-e^{13} + e^{14} - e^{23} - e^{24}) - e^{15}, & de^4 &= -2e^{35}, \\ & & de^5 &= -4b^2e^{34}, \end{aligned}$$

and endowed with the quasi-Sasakian structure given by (14). The second cohomology group of \mathfrak{g}_b is generated by e^{12} . The Lie algebras \mathfrak{g}_b are not solvable since the commutators are $[\mathfrak{g}_b, \mathfrak{g}_b] = \mathfrak{g}_b$.

The Lie groups underlying Examples 2.9 and 2.10 also satisfy the conditions of Corollary 2.4 with $\Omega \wedge \Omega = 0$, by just taking as connection 1-form the 1-form e^6 such that $de^6 = \lambda e^{12}$. Then, $\Omega = \lambda e^{12}$. With this expression of de^6 , we have

$$d^2e^6 = 0, \quad J(de^6) = de^6, \quad \text{and} \quad de^6 \wedge de^6 = 0.$$

Therefore, equation (11) is satisfied. Observe that $\lambda = 0$ provides examples of trivial S^1 -bundles.

The next example allows us to recover one of the 6-dimensional nilmanifolds found in [Fino et al. 2004]:

Example 2.11. Consider the 5-dimensional nilpotent Lie algebra with structure equations

$$\begin{aligned} de^j &= 0 \quad \text{for } j = 1, \dots, 4, \\ de^5 &= e^{12} + e^{34}, \end{aligned}$$

and endowed with the quasi-Sasakian structure given by (14). If we consider the closed 2-form $\Omega = e^{13} + e^{24}$ and apply Corollary 2.4, we see that there exists a nontrivial S^1 -bundle over the corresponding 5-dimensional nilmanifold. Since $de^5 \wedge de^5 = -\Omega \wedge \Omega \neq 0$, the total space of this S^1 -bundle is an SKT nilmanifold. More precisely, according to the classification given in [Fino et al. 2004] (see also

[Ugarte 2007]), the nilmanifold is the one with underlying Lie algebra isomorphic to $\mathfrak{h}_3 \oplus \mathfrak{h}_3$, where by \mathfrak{h}_3 we denote the real 3-dimensional Heisenberg Lie algebra.

Since the starting Lie algebra from Example 2.11 is Sasakian, it is natural to start with other 5-dimensional Sasakian Lie algebras to construct new SKT structures in dimension 6. A classification of 5-dimensional Sasakian Lie algebras was obtained in [Andrada et al. 2009].

Example 2.12. Consider the 5-dimensional Lie algebra \mathfrak{k}_3 with structure equations

$$\begin{aligned} de^1 &= 0, & de^4 &= 0, \\ de^2 &= -e^{13}, & de^5 &= \lambda e^{14} + \mu e^{23}, \\ de^3 &= e^{12}, \end{aligned}$$

where $\lambda, \mu < 0$. By [Andrada et al. 2009], this algebra admits the Sasakian structure given by

$$\begin{aligned} Ie^1 &= e^4, & Ie^2 &= e^3, & \eta &= e^5, \\ g &= -\frac{1}{2}\lambda(e_1)^2 - \frac{1}{2}\lambda(e_2)^2 - \frac{1}{2}\mu(e_3)^2 - \frac{1}{2}\mu(e_4)^2 + (e_5)^2, \end{aligned}$$

and is isomorphic to $\mathbb{R} \ltimes (\mathfrak{h}_3 \times \mathbb{R})$; moreover, the corresponding solvable simply connected Lie group admits a compact quotient by a discrete subgroup.

Consider on \mathfrak{k}_3 the closed 2-form $\Omega = \lambda e^{14} - \mu e^{23}$. The form Ω is I -invariant and satisfies $\Omega \wedge \Omega = -2\lambda\mu e^{1234}$. Since e^5 is the contact form and $de^5 \wedge de^5 = 2\lambda\mu e^{1234}$, we get again by Corollary 2.4 an SKT structure on a nontrivial S^1 -bundle over the 5-dimensional solvmanifold. We denote by e^6 the connection 1-form.

The orthonormal basis $\{\alpha^1 = e^1, \alpha^2 = e^4, \alpha^3 = e^2, \alpha^4 = e^3, \alpha^5 = e^5, \alpha^6 = \theta\}$ for the SKT metric satisfies the equations

$$\begin{aligned} d\alpha^1 &= d\alpha^2 = 0, & d\alpha^3 &= -\alpha^{14}, & d\alpha^4 &= \alpha^{13}, \\ d\alpha^5 &= \lambda\alpha^{12} + \mu\alpha^{34}, & d\alpha^6 &= \lambda\alpha^{12} - \mu\alpha^{34}, \end{aligned}$$

and the complex structure is given by $J(X_1) = X_2$, $J(X_3) = X_4$ and $J(X_5) = X_6$, where $\{X_i\}_{i=1}^6$ denotes the basis dual to $\{\alpha^i\}_{i=1}^6$. Since the fundamental 2-form is $F = \alpha^{12} + \alpha^{34} + \alpha^{56}$, the 3-form torsion T of the SKT structure is

$$T = \lambda\alpha^{12}(\alpha^5 + \alpha^6) + \mu\alpha^{34}(\alpha^5 - \alpha^6).$$

Moreover, $*T = \lambda\alpha^{12}(\alpha^5 + \alpha^6) - \mu\alpha^{34}(\alpha^5 - \alpha^6)$, where $*$ denotes the metric's Hodge operator; this implies that the torsion form is also coclosed.

The only nonzero curvature forms $(\Omega^B)_j^i$ of the Bismut connection ∇^B are

$$(\Omega^B)_2^1 = -2\lambda^2\alpha^{12} \quad \text{and} \quad (\Omega^B)_4^3 = -2\mu^2\alpha^{34}.$$

A direct calculation shows that the 1-forms α^5 and α^6 and the 2-forms α^{12} and α^{34} are parallel with respect to the Bismut connection, which implies that $\nabla^B T = 0$.

Finally, $\text{Hol}(\nabla^B) = U(1) \times U(1) \subset U(3)$ since $\nabla^B \alpha^i \neq 0$ for $i = 1, 2, 3, 4$.

3. SKT structures arising from Riemannian cones

Let N^{2n+1} be a $(2n+1)$ -dimensional manifold endowed with an almost contact metric structure (I, ξ, η, g) , and denote by ω its fundamental 2-form.

The Riemannian cone of N^{2n+1} is defined as the manifold $N^{2n+1} \times \mathbb{R}^+$ equipped with the cone metric

$$(15) \quad h = t^2 g + (dt)^2.$$

The cone $N^{2n+1} \times \mathbb{R}^+$ has a natural almost Hermitian structure defined by

$$(16) \quad F = t^2 \omega + t \eta \wedge dt.$$

The almost complex structure J on $N^{2n+1} \times \mathbb{R}^+$ defined by (F, h) is given by

$$JX = IX \quad \text{for } X \in \text{Ker } \eta \quad \text{and} \quad J\xi = -t \frac{d}{dt}.$$

In terms of a local orthonormal adapted coframe $\{e^1, \dots, e^{2n}\}$ for g with

$$(17) \quad \omega = - \sum_{j=1}^n e^{2j-1} \wedge e^{2j},$$

we have

$$(18) \quad \begin{aligned} J e^{2j-1} &= -e^{2j}, & J e^{2j} &= e^{2j-1} \quad \text{for } j = 1, \dots, n, \\ J(t e^{2n+1}) &= dt, & J(dt) &= -t e^{2n+1}. \end{aligned}$$

The almost Hermitian structure (J, h) on $N^{2n+1} \times \mathbb{R}^+$ is Kähler if and only if the almost contact metric structure (I, ξ, η, g) on N^{2n+1} is Sasakian, that is, a normal contact metric structure.

If we impose that the almost Hermitian structure (J, h) on $N^{2n+1} \times \mathbb{R}^+$ is SKT, we can prove the following:

Theorem 3.1. *Consider a $(2n+1)$ -dimensional almost contact metric manifold $(N^{2n+1}, I, \xi, \eta, g)$. The almost Hermitian structure (J, h) given by (15) and (16) on the Riemannian cone $(N^{2n+1} \times \mathbb{R}^+, h)$ is SKT if and only if (I, ξ, η, g) is normal and*

$$(19) \quad -4\eta \wedge \omega + 2I(d\omega) - 2d\eta \wedge \eta = d(I(i_\xi d\omega)),$$

where ω denotes the fundamental 2-form of the almost contact metric structure (I, ξ, η, g) .

Proof. J is integrable if and only if the almost contact metric structure is normal. We compute JdF . We have

$$\begin{aligned} dF &= 2tdt \wedge \omega + t^2d\omega + t d\eta \wedge dt, \quad \text{and so} \\ JdF &= -2t^2\eta \wedge \omega + t^2J(d\omega) - t^2d\eta \wedge \eta \end{aligned}$$

since $J\omega = \omega$, $J(dt) = -t\eta$ and $Jd\eta = d\eta$. Moreover, with respect to an adapted basis $\{e^1, \dots, e^{2n+1}\}$ we can get, in a way similar to the proof of Theorem 2.3, that

$$(20) \quad Jd\omega = I(d\omega) + I(i_\xi d\omega) \wedge J\eta.$$

As a consequence, we get $JdF = -2t^2\eta \wedge \omega + t^2I(d\omega) + t dt \wedge I(i_\xi d\omega) - t^2d\eta \wedge \eta$. Therefore, by imposing that $d(JdF) = 0$, we obtain

$$\begin{aligned} -4\eta \wedge \omega + 2I(d\omega) - 2d\eta \wedge \eta - d(I(i_\xi d\omega)) &= 0, \\ -2d(\eta \wedge \omega) + d(I(d\omega)) - d(d\eta \wedge \eta) &= 0. \end{aligned}$$

Since the second equation is a consequence of the first, the Hermitian structure (F, h) on the Riemannian cone $N^{2n+1} \times \mathbb{R}^+$ is SKT if and only if the almost contact metric structure $(I, \eta, \xi, g, \omega)$ on N^{2n+1} satisfies equation (19). \square

Remark 3.2. As a consequence of previous theorem, when $n = 1$, equation (19) is satisfied if and only if the 3-dimensional manifold N is Sasakian. On the other hand, if $n > 1$ and the almost contact metric structure on N^{2n+1} is quasi-Sasakian (that is, $d\omega = 0$), then the structure has to be Sasakian, that is, $d\eta = -2\omega$.

Example 3.3. Consider the 5-dimensional Lie algebras $\mathfrak{g}_{a,b,c}$ with structure equations

$$\begin{aligned} de^1 &= ae^{23} + 2e^{25} + \left(-\frac{1}{2}ab + \frac{b^3}{2a} + 2\frac{b}{a}\right)e^{34} + be^{45}, \\ de^2 &= -ae^{13} - 2e^{15} - \frac{1}{2}bce^{34} - be^{35}, \\ de^3 &= \left(-\frac{4}{a} - \frac{b^2}{a}\right)e^{34}, \\ de^4 &= ce^{34}, \\ de^5 &= 2e^{12} + be^{14} - be^{23} + (2 + b^2)e^{34}, \end{aligned}$$

where $a, b, c \in \mathbb{R}$ and $a \neq 0$. They are endowed with the normal almost contact metric structure $(I, \xi, \eta, g, \omega)$ with

$$Ie^1 = -e^2, \quad Ie^3 = -e^4, \quad \eta = e^5, \quad \omega = -e^{12} - e^{34}.$$

This structure satisfies (19), and therefore the Riemannian cones over the corresponding simply connected Lie groups are SKT.

4. SKT SU(3)-structures

Let (M^6, J, h) be a 6-dimensional almost Hermitian manifold. An SU(3)-structure on M^6 is determined by the choice of a $(3, 0)$ -form $\Psi = \Psi_+ + i\Psi_-$ of unit norm. If Ψ is closed, then the underlying almost complex structure J is integrable and the manifold is Hermitian. We will denote the SU(3)-structure (J, h, Ψ) simply by (F, Ψ) , where F is the fundamental 2-form, since from F and Ψ we can reconstruct the almost Hermitian structure.

Definition 4.1. *We say that an SU(3)-structure (F, Ψ) on M^6 is SKT if*

$$(21) \quad d\Psi = 0 \quad \text{and} \quad d(JdF) = 0,$$

where J is the associated complex structure.

We will see the relation between SKT SU(3)-structures in dimension 6 and SU(2)-structures in dimension 5.

First, we recall some facts about SU(2)-structures on a 5-dimensional manifold. An SU(2)-structure on a 5-dimensional manifold N^5 is an SU(2)-reduction of the principal bundle of linear frames on N^5 . By [Conti and Salamon 2007, Proposition 1], these structures are in one-to-one correspondence with quadruplets $(\eta, \omega_1, \omega_2, \omega_3)$, where η is a 1-form and ω_i are 2-forms on N^5 satisfying $\omega_i \wedge \omega_j = \delta_{ij}v$ and $v \wedge \eta \neq 0$ for some 4-form v , and $\omega_2(X, Y) \geq 0$ if $i_X \omega_3 = i_Y \omega_1$, where i_X denotes the contraction by X . Equivalently, an SU(2)-structure on N^5 can be viewed as the datum of (η, ω_1, Φ) , where η is a 1-form, ω_1 is a 2-form, and $\Phi = \omega_2 + i\omega_3$ is a complex 2-form such that

$$\eta \wedge \omega_1 \wedge \omega_1 \neq 0, \quad \Phi \wedge \Phi = 0, \quad \omega_1 \wedge \Phi = 0, \quad \Phi \wedge \bar{\Phi} = 2\omega_1 \wedge \omega_1,$$

and Φ is of type $(2, 0)$ with respect to ω_1 .

Conti and Salamon [2007] locally characterize an SU(2)-structure as follows. If $(\eta, \omega_1, \omega_2, \omega_3)$ is an SU(2)-structure on a 5-manifold N^5 , then locally there exists an orthonormal basis of 1-forms $\{e^1, \dots, e^5\}$ such that

$$\omega_1 = e^{12} + e^{34}, \quad \omega_2 = e^{13} - e^{24}, \quad \omega_3 = e^{14} + e^{23}, \quad \eta = e^5.$$

We can also consider the local tensor field I given by

$$Ie^1 = -e^2, \quad Ie^2 = e^1, \quad Ie^3 = -e^4, \quad Ie^4 = e^3, \quad Ie^5 = 0.$$

This tensor gives rise to a global tensor field of type $(1, 1)$ on the manifold N^5 , defined by $\omega_1(X, Y) = g(X, IY)$ for any vector fields X and Y on N^5 , where g is the Riemannian metric on N^5 underlying the SU(2)-structure. The tensor field I satisfies $I^2 = -\text{Id} + \eta \otimes \xi$, where ξ is the vector field on N^5 dual to the 1-form η .

Therefore, given an SU(2)-structure $(\eta, \omega_1, \omega_2, \omega_3)$ we also have an almost contact metric structure (I, ξ, η, g) on the manifold, where ω_1 is its fundamental form.

Remark 4.2. Notice that we have two more almost contact metric structures when we consider ω_2 and ω_3 as fundamental forms.

If N^5 has an $SU(2)$ -structure $(\eta, \omega_1, \omega_2, \omega_3)$, the product $N^5 \times \mathbb{R}$ has a natural $SU(3)$ -structure given by

$$(22) \quad F = \omega_1 + \eta \wedge dt \quad \text{and} \quad \Psi = (\omega_2 + i\omega_3) \wedge (\eta - i dt).$$

By Corollary 2.6, the previous $SU(3)$ -structure is SKT if and only if

$$(23) \quad \begin{aligned} d(I(d\omega_1)) &= d(d\eta \wedge \eta), & d\omega_2 &= -3\omega_3 \wedge \eta, \\ d(I(i_\xi d\omega_1)) &= 0, & d\omega_3 &= 3\omega_2 \wedge \eta, \end{aligned}$$

proving this:

Theorem 4.3. *Suppose N^5 is a 5-dimensional manifold endowed with an $SU(2)$ -structure $(\eta, \omega_1, \omega_2, \omega_3)$. The $SU(3)$ -structure (F, Ψ) given by (22) on the product $N^5 \times \mathbb{R}$ is SKT if and only if the equations (23) are satisfied.*

Example 4.4. On the 5-dimensional Lie algebras introduced in Examples 2.8, 2.9 and 2.10, consider the $SU(2)$ -structure given by

$$\omega = \omega_1 = e^{12} + e^{34}, \quad \omega_2 = e^{13} - e^{24}, \quad \omega_3 = e^{14} + e^{23}.$$

For Example 2.8, we have

$$d\omega_2 = -2\omega_3 \wedge \eta - 4(e^{124} - e^{134}) \quad \text{and} \quad d\omega_3 = 2\omega_2 \wedge \eta + 4(e^{123} + e^{234}).$$

For Examples 2.9 and 2.10, we get $d\omega_2 = -3\omega_3 \wedge \eta$ and $d\omega_3 = 3\omega_2 \wedge \eta$. Therefore one gets an SKT $SU(3)$ -structure on the product of the corresponding simply connected Lie groups by \mathbb{R} .

We will study the existence of SKT $SU(3)$ -structures on a Riemannian cone over a 5-dimensional manifold N^5 endowed with an $SU(2)$ -structure $(\eta, \omega_1, \omega_2, \omega_3)$. Then N^5 has an induced almost contact metric structure (I, ξ, η, g) , and ω_1 is its fundamental form.

The Riemannian cone $(N^5 \times \mathbb{R}^+, h)$ of (N^5, g) has a natural $SU(3)$ -structure defined by

$$F = t^2 \omega_1 + t\eta \wedge dt \quad \text{and} \quad \Psi = t^2(\omega_2 + i\omega_3) \wedge (t\eta - i dt).$$

In terms of a local orthonormal coframe $\{e^1, \dots, e^5\}$ for g such that

$$\omega_1 = -e^{12} - e^{34}, \quad \omega_2 = -e^{13} + e^{24}, \quad \omega_3 = -e^{14} - e^{23}, \quad \eta = e^5,$$

we have

$$\begin{aligned} J e^1 &= -e^2, & J e^2 &= e^1, & J e^3 &= -e^4, \\ J e^4 &= e^3, & J(t e^5) &= dt, & J(dt) &= -t e^5. \end{aligned}$$

We recall that the $SU(3)$ -structure (F, Ψ) on $N^5 \times \mathbb{R}^+$ is integrable if and only if the $SU(2)$ -structure $(\eta, \omega_1, \omega_2, \omega_3)$ on N^5 is Sasaki–Einstein, or equivalently if and only if

$$d\eta = -2\omega_1, \quad d\omega_2 = -3\omega_3 \wedge \eta, \quad d\omega_3 = 3\omega_2 \wedge \eta.$$

For the Riemannian cones, we can prove the following

Corollary 4.5. *Let N^5 be a 5-dimensional manifold endowed with an $SU(2)$ -structure $(\eta, \omega_1, \omega_2, \omega_3)$. The $SU(3)$ -structure (F, Ψ) on the Riemannian cone $(N^5 \times \mathbb{R}^+, h)$ is SKT if and only if*

$$(24) \quad \begin{aligned} -4\eta \wedge \omega_1 + 2I(d\omega_1) - 2d\eta \wedge \eta &= d(I(i_\xi d\omega_1)), \\ d\omega_2 &= 3\omega_3 \wedge \eta, \\ d\omega_3 &= -3\omega_2 \wedge \eta. \end{aligned}$$

Proof. By imposing that $d\Psi = 0$, we get the conditions $d\omega_2 = -3\omega_3 \wedge \eta$ and $d\omega_3 = 3\omega_2 \wedge \eta$. By imposing $d(JdF) = 0$, we get, as in the proof of Theorem 3.1, equation (19) for $\omega = \omega_1$. \square

5. Almost contact metric structure induced on a hypersurface

We study the almost contact metric structure induced naturally on any oriented hypersurface N^{2n+1} of a $(2n+2)$ -manifold M^{2n+2} equipped with an SKT structure.

Let $f: N^{2n+1} \rightarrow M^{2n+2}$ be an oriented hypersurface of a $(2n+2)$ -dimensional manifold M^{2n+2} endowed with an SKT structure (J, h, F) , and denote by \mathbb{U} the unitary normal vector field. It is well known that N^{2n+1} inherits an almost contact metric structure (I, ξ, η, g) such that η and the fundamental 2-form ω are given by

$$(25) \quad \eta = -f^*(i_{\mathbb{U}}F) \quad \text{and} \quad \omega = f^*F,$$

where F is the fundamental 2-form of the almost Hermitian structure; see, for instance, [Blair 2002].

Proposition 5.1. *Suppose $f: N^{2n+1} \rightarrow M^{2n+2}$ is an immersion of an oriented $(2n+1)$ -dimensional manifold into a $(2n+2)$ -dimensional Hermitian manifold. If the Hermitian structure (J, h) on M^{2n+2} is SKT, then the induced almost contact metric structure (I, ξ, η, g) on N^{2n+1} , with η and ω given by (25), satisfies*

$$(26) \quad d(Id\omega - I(f^*(i_{\mathbb{U}}dF)) \wedge \eta) = 0.$$

Proof. Locally we can choose an adapted coframe $\{e^1, \dots, e^{2n+2}\}$ for the Hermitian structure so that the unitary normal vector field \mathbb{U} is dual to e^{2n+2} . Since the

almost complex structure J is given in this adapted basis by

$$\begin{aligned} J e^{2j-1} &= -e^{2j}, & J e^{2j} &= e^{2j-1} \quad \text{for } j = 1, \dots, n, \\ J e^{2n+1} &= e^{2n+2}, & J e^{2n+2} &= -e^{2n+1}, \end{aligned}$$

it follows that the tensor field I on N^{2n+1} has $I f^* e^i = f^* J e^i$ for $i = 1, \dots, 2n+1$. That is,

$$I f^* e^{2j-1} = -f^* e^{2j}, \quad I f^* e^{2j} = f^* e^{2j-1} \quad \text{for } j = 1, \dots, n, \quad I f^* e^{2n+1} = 0.$$

However, $I f^* e^{2n+2} = 0 \neq f^* e^{2n+1} = -f^* J e^{2n+2}$.

We compute $f^* J dF$. First, we decompose (locally and in terms of the adapted basis) the differential of F as

$$dF = \alpha + \beta \wedge e^{2n+1} + \gamma \wedge e^{2n+2} + \mu \wedge e^{2n+1} \wedge e^{2n+2},$$

where the local forms

$$\alpha \in \bigwedge^3 \langle e^1, \dots, e^{2n} \rangle, \quad \beta, \gamma \in \bigwedge^2 \langle e^1, \dots, e^{2n} \rangle, \quad \mu \in \bigwedge^1 \langle e^1, \dots, e^{2n} \rangle$$

are generated only by e^1, \dots, e^{2n} . Then,

$$J dF = J \alpha + J \beta \wedge e^{2n+2} - J \gamma \wedge e^{2n+1} + J \mu \wedge e^{2n+1} \wedge e^{2n+2}.$$

Since $f^* e^{2n+2} = 0$ and $f^* e^{2n+1} = \eta$, we have $f^* J dF = f^* J \alpha - (f^* J \gamma) \wedge \eta$. However, $f^*(i_{\cup} dF) = f^* \gamma + f^* \mu \wedge \eta$, which implies that

$$I(f^*(i_{\cup} dF)) = I f^* \gamma = f^* J \gamma.$$

On the other hand, $I d\omega = I d f^* F = I f^* dF = I f^* \alpha = f^* J \alpha$. We conclude that

$$f^* J dF = f^* J \alpha - (f^* J \gamma) \wedge \eta = I d\omega - I(f^*(i_{\cup} dF)) \wedge \eta.$$

Now, if the Hermitian structure is SKT, then $J dF$ is closed and the induced structure satisfies (26). \square

Remark 5.2. Notice that, using $i_{\cup} dF = \mathcal{L}_{\cup} F - d i_{\cup} F$, we can write (26) as

$$d(I d\omega - I(f^*(\mathcal{L}_{\cup} F) + d\eta) \wedge \eta) = 0.$$

Therefore, if $f^*(\mathcal{L}_{\cup} F) = 0$, then the induced almost contact metric structure has to satisfy the equation $d(I d\omega - I(d\eta) \wedge \eta) = 0$. In the case of the product $N^{2n+1} \times \mathbb{R}$, the condition $f^*(\mathcal{L}_{\cup} F) = 0$ is satisfied. In the case of the Riemannian cone, we have $\mathcal{L}_{d/dt} F = 2t\omega + dt \wedge \eta$ and therefore $f^*(\mathcal{L}_{d/dt} F) = 2\omega$. In this way, we recover some of the equations obtained in Corollary 2.6 and Theorem 3.1.

Now we study the structure that is induced naturally on any oriented hypersurface N^5 of a 6-manifold M^6 equipped with an SKT $SU(3)$ -structure.

Let $f: N^5 \rightarrow M^6$ be an oriented hypersurface of a 6-manifold M^6 endowed with an $SU(3)$ -structure $(F, \Psi = \Psi_+ + i\Psi_-)$, and denote by \mathbb{U} the unitary normal vector field. Then N^5 inherits an $SU(2)$ -structure $(\eta, \omega_1, \omega_2, \omega_3)$ given by

$$(27) \quad \eta = -f^*(i_{\mathbb{U}}F), \quad \omega_1 = f^*F, \quad \omega_2 = -f^*(i_{\mathbb{U}}\Psi_-), \quad \omega_3 = f^*(i_{\mathbb{U}}\Psi_+).$$

Corollary 5.3. *Let $f: N^5 \rightarrow M^6$ be an immersion of an oriented 5-dimensional manifold into a 6-dimensional manifold with an $SU(3)$ -structure. If the $SU(3)$ -structure is SKT, then the induced $SU(2)$ -structure on N^5 given by (27) satisfies*

$$(28) \quad d(Id\omega_1 - If^*(i_{\mathbb{U}}dF) \wedge \eta) = 0,$$

$$(29) \quad d(\omega_2 \wedge \eta) = 0 \quad \text{and} \quad d(\omega_3 \wedge \eta) = 0.$$

Proof. Equation (28) follows from Proposition 5.1 by taking $\omega = \omega_1$. Locally, we can choose an adapted coframe $\{e^1, \dots, e^5, e^6\}$ for the $SU(3)$ -structure such that the unitary normal vector field \mathbb{U} is dual to e^6 . From (27), it follows that $\omega_2 \wedge \eta = f^*\Psi_+$ and $\omega_3 \wedge \eta = f^*\Psi_-$. If $\Psi = \Psi_+ + i\Psi_-$ is closed, then the induced structure satisfies (29). \square

5.1. A simple example. Consider the 6-dimensional nilmanifold M^6 whose underlying nilpotent Lie algebra has structure equations

$$de^j = 0 \quad \text{for } j = 1, 2, 3, 6, \quad de^4 = e^{12}, \quad de^5 = e^{14},$$

and is endowed with the $SU(3)$ -structure given by

$$F = -e^{14} - e^{26} - e^{53} \quad \text{and} \quad \Psi = (e^1 - ie^4) \wedge (e^2 - ie^6) \wedge (e^5 - ie^3).$$

The oriented hypersurface with normal vector field dual to e^2 is a 5-dimensional nilmanifold N^5 that by [Conti and Salamon 2007] has no invariant hypo structures. However, the $SU(2)$ -structure on N^5 , namely,

$$(30) \quad \eta = e^2, \quad \omega_1 = -e^{14} - e^{53}, \quad \omega_2 = -e^{15} - e^{34}, \quad \omega_3 = -e^{13} - e^{45},$$

satisfies (28) and (29). In Section 6, we will show that by using this $SU(2)$ -structure and appropriate evolution equations, we can construct an SKT $SU(3)$ -structure on the product of N^5 with an open interval.

6. SKT evolution equations

The goal here is to construct SKT $SU(3)$ -structures by using appropriate evolution equations, starting from a suitable $SU(2)$ -structure on a 5-dimensional manifold. We follow ideas of [Hitchin 2001] and [Conti and Salamon 2007].

Lemma 6.1. *Let $(\eta(t), \omega_1(t), \omega_2(t), \omega_3(t))$ be a family of $SU(2)$ -structures on a 5-dimensional manifold N^5 for $t \in (a, b)$. The $SU(3)$ -structure on $M^6 = N^5 \times (a, b)$*

given by $F = \omega_1(t) + \eta(t) \wedge dt$ and $\Psi = (\omega_2(t) + i\omega_3(t)) \wedge (\eta(t) - i dt)$ satisfies the condition $d\Psi = 0$ if and only if $(\eta(t), \omega_1(t), \omega_2(t), \omega_3(t))$ is an $SU(2)$ -structure such that, for any t in the open interval (a, b) ,

$$(31) \quad \begin{aligned} \hat{d}(\omega_2(t) \wedge \eta(t)) &= 0, & \partial_t(\omega_2(t) \wedge \eta(t)) &= -\hat{d}\omega_3(t), \\ \hat{d}(\omega_3(t) \wedge \eta(t)) &= 0, & \partial_t(\omega_3(t) \wedge \eta(t)) &= \hat{d}\omega_2(t). \end{aligned}$$

Here, \hat{d} denotes the exterior differential on N^5 and d is the exterior differential on M^6 . We now present the additional evolution equations to be added to the last two of (31) in order to ensure that $dJdF = 0$.

Proposition 6.2. *Let $(\eta(t), \omega_1(t), \omega_2(t), \omega_3(t))$ be a family of $SU(2)$ -structures on N^5 for $t \in (a, b)$. The $SU(3)$ -structure on $M^6 = N^5 \times (a, b)$ given by*

$$(32) \quad F = \omega_1(t) + \eta(t) \wedge dt \quad \text{and} \quad \Psi = (\omega_2(t) + i\omega_3(t)) \wedge (\eta(t) - i dt),$$

has JdF closed if and only if $(\eta(t), \omega_1(t), \omega_2(t), \omega_3(t))$ satisfies the evolution equations

$$(33) \quad \begin{aligned} \hat{d}(I_t \hat{d}\omega_1(t) - I_t(\partial_t \omega_1(t) + \hat{d}\eta(t)) \wedge \eta(t)) &= 0, \\ \partial_t(I_t \hat{d}\omega_1(t) - I_t(\partial_t \omega_1(t) + \hat{d}\eta(t)) \wedge \eta(t)) \\ &= -\hat{d}(I_t(i_\xi \hat{d}\omega_1(t)) - I_t(i_\xi(\partial_t \omega_1(t) + \hat{d}\eta(t)))) \wedge \eta(t), \end{aligned}$$

where $\xi(t)$ denotes the vector field on N^5 dual to $\eta(t)$ for each $t \in (a, b)$.

Proof. Since $F = \omega_1(t) + \eta(t) \wedge dt$, we have $dF = \hat{d}\omega_1 + (\partial_t \omega_1 + \hat{d}\eta) \wedge dt$. Define $\{e^1(t), \dots, e^4(t), \eta(t)\}$ to be a local adapted basis for the $SU(2)$ -structure $(\eta(t), \omega_1(t), \omega_2(t), \omega_3(t))$. Then, $\{e^1(t), \dots, e^4(t), \eta(t), dt\}$ is an adapted basis for the $SU(3)$ -structure (32), and J is given by

$$\begin{aligned} J e^1(t) &= -e^2(t), & J e^2(t) &= e^1(t), & J \eta(t) &= dt, \\ J e^3(t) &= -e^4(t), & J e^4(t) &= e^3(t), & J dt &= -\eta(t). \end{aligned}$$

For each t , the structures I_t induced on N^5 are given by

$$\begin{aligned} I_t e^1(t) &= -e^2(t), & I_t e^2(t) &= e^1(t), \\ I_t e^3(t) &= -e^4(t), & I_t e^4(t) &= e^3(t), & I_t \eta(t) &= 0. \end{aligned}$$

Now, we can locally decompose a given $\tau(t) \in \Omega^k(N^5)$ for $t \in (a, b)$ as

$$\tau(t) = \alpha(t) + \beta(t) \wedge \eta(t),$$

where $\alpha(t) \in \bigwedge^k \langle e^1(t), \dots, e^4(t) \rangle$ and $\beta(t) \in \bigwedge^{k-1} \langle e^1(t), \dots, e^4(t) \rangle$. Therefore,

$$\begin{aligned} J\tau(t) &= J\alpha(t) + J\beta(t) \wedge J\eta(t) = I_t \alpha(t) + I_t \beta(t) \wedge dt \\ &= I_t \tau(t) - (-1)^k I_t(i_{\xi(t)} \tau(t)) \wedge dt. \end{aligned}$$

Applying this to JdF , we get

$$\begin{aligned} JdF &= J\hat{d}\omega_1 - J(\partial_t\omega_1 + \hat{d}\eta) \wedge \eta(t) \\ &= I_t\hat{d}\omega_1 - I_t(\partial_t\omega_1 + \hat{d}\eta) \wedge \eta(t) + I_t(i_{\xi(t)}\hat{d}\omega_1) \wedge dt \\ &\quad - I_t(i_{\xi}(\partial_t\omega_1 + \hat{d}\eta)) \wedge \eta(t) \wedge dt. \end{aligned}$$

Finally, taking the differential of JdF , we get

$$\begin{aligned} dJdF &= \hat{d}(I_t\hat{d}\omega_1 - I_t(\partial_t\omega_1 + \hat{d}\eta) \wedge \eta(t)) \\ &\quad + \partial_t(I_t\hat{d}\omega_1 - I_t(\partial_t\omega_1 + \hat{d}\eta) \wedge \eta(t)) \wedge dt \\ &\quad + \hat{d}(I_t(i_{\xi(t)}\hat{d}\omega_1) - I_t(i_{\xi}(\partial_t\omega_1 + \hat{d}\eta))) \wedge \eta(t) \wedge dt. \quad \square \end{aligned}$$

Remark 6.3. Observe that the first equation in (33) is exactly condition (28) for $F = \omega_1(t) + \eta(t) \wedge dt$. See Remark 5.2.

As a consequence of Lemma 6.1 and Proposition 6.2, we get the following:

Theorem 6.4. For $t \in (a, b)$, let $(\eta(t), \omega_1(t), \omega_2(t), \omega_3(t))$ be a family of $SU(2)$ -structures on a 5-dimensional manifold N^5 such that

$$(34) \quad \hat{d}(\omega_2(t) \wedge \eta(t)) = 0 \quad \text{and} \quad \hat{d}(\omega_3(t) \wedge \eta(t)) = 0$$

for any t . If the evolution equations

$$\begin{aligned} &\hat{d}(I_t\hat{d}\omega_1(t) - I_t(\partial_t\omega_1(t) + \hat{d}\eta(t)) \wedge \eta(t)) = 0, \\ &\partial_t(I_t\hat{d}\omega_1(t) - I_t(\partial_t\omega_1(t) + \hat{d}\eta(t)) \wedge \eta(t)) \\ (35) \quad &= -\hat{d}(I_t(i_{\xi}\hat{d}\omega_1(t)) - I_t(i_{\xi}(\partial_t\omega_1(t) + \hat{d}\eta(t)))) \wedge \eta(t), \\ &\partial_t(\omega_2(t) \wedge \eta(t)) = -\hat{d}\omega_3(t), \\ &\partial_t(\omega_3(t) \wedge \eta(t)) = \hat{d}\omega_2(t), \end{aligned}$$

are satisfied, then the $SU(3)$ -structure on $M = N \times (a, b)$ given by

$$(36) \quad F = \omega_1(t) + \eta(t) \wedge dt \quad \text{and} \quad \Psi = (\omega_2(t) + i\omega_3(t)) \wedge (\eta(t) - idt)$$

is SKT.

Example 6.5. Consider the Lie algebra with structure equations

$$de^j = 0 \quad \text{for } j = 1, 2, 3, \quad de^4 = e^{12}, \quad de^5 = e^{14},$$

which underlies the 5-dimensional nilmanifold N^5 considered in Section 5.1. We endow it with the $SU(2)$ -structure given by (30). It is easy to verify that

$$d(\omega_2 \wedge \eta) = d(\omega_3 \wedge \eta) = d(\omega_1 \wedge \omega_1) = 0.$$

We evolve this $SU(2)$ -structure by

$$\begin{aligned}\omega_1(t) &= -e^{14} - e^{53}, & \omega_2(t) &= -(1 + \frac{3}{2}t)^{1/3}e^{15} - (1 + \frac{3}{2}t)^{-1/3}e^{34}, \\ \eta(t) &= (1 + \frac{3}{2}t)^{1/3}e^2, & \omega_3(t) &= -(1 + \frac{3}{2}t)^{1/3}e^{13} - (1 + \frac{3}{2}t)^{-1/3}e^{45},\end{aligned}$$

where $t \in (-2/3, \infty)$.

For any $t \in (-2/3, \infty)$, the family $(\omega_1(t), \omega_2(t), \omega_3(t), \eta(t))$ satisfies equations (34) and the last two equations of (35). Moreover, it satisfies the conditions

$$\partial_t \omega_1(t) = 0, \quad \hat{d}(\eta(t)) = 0, \quad i_{\xi}(\hat{d}(\omega_1(t))) = 0, \quad \partial_t(I_t(\hat{d}\omega_1(t))) = 0,$$

which implies that the evolution equations (33) are also satisfied.

On the product $N^5 \times \mathbb{R}$, we consider the local basis of 1-forms

$$\begin{aligned}\beta^1 &= (1 + \frac{3}{2}t)^{1/3}e^1, & \beta^2 &= (1 + \frac{3}{2}t)^{-1/3}e^4, & \beta^3 &= e^5, \\ \beta^4 &= e^3, & \beta^5 &= (1 + \frac{3}{2}t)^{1/3}e^2, & \beta^6 &= dt.\end{aligned}$$

The structure equations are

$$\begin{aligned}d\beta^1 &= -\frac{1}{2}(1 + \frac{3}{2}t)^{-1}\beta^{16}, & d\beta^4 &= 0, \\ d\beta^2 &= (1 + \frac{3}{2}t)^{-1}(\beta^{15} + \frac{1}{2}\beta^{26}), & d\beta^5 &= -\frac{1}{2}(1 + \frac{3}{2}t)^{-1}\beta^{56}, \\ d\beta^3 &= \beta^{12}, & d\beta^6 &= 0.\end{aligned}$$

Locally, J is given by $J\beta^1 = -\beta^2$, $J\beta^3 = -\beta^4$ and $J\beta^5 = \beta^6$. The fundamental form $F = -\beta^{12} - \beta^{34} + \beta^{56}$ satisfies $d(JdF) = 0$, and the $(3, 0)$ -form $\Psi = (\beta^1 + i\beta^2) \wedge (\beta^3 + i\beta^4) \wedge (\beta^5 - i\beta^6)$ is closed. Therefore, (F, Ψ) is a local SKT $SU(3)$ -structure on $N^5 \times \mathbb{R}$.

Remark 6.6. A Hermitian structure (J, h) on a 6-dimensional manifold M^6 is called *balanced* if $F \wedge F$ is closed, F being the associated fundamental 2-form. The paper [Fernández et al. 2009] introduced the notion of balanced $SU(2)$ -structures on 5-dimensional manifolds, together with appropriate evolution equations whose solution gives rise to a balanced $SU(3)$ -structure in six dimensions.

If M^6 is compact, then a balanced structure cannot be SKT; see, for instance, [Fino et al. 2004].

The $SU(2)$ -structure (30) from the previous example is also balanced, and it gives rise to a balanced metric on the product of N^5 with a open interval; see [Fernández et al. 2009, (11)]. However, one can check directly that this solution is not SKT.

If G is the nilpotent Lie group underlying N^5 , the product $G \times \mathbb{R}$ has no left-invariant SKT structures and does not admit any left-invariant complex structures; however, we can find a local SKT $SU(3)$ -structure on it.

7. HKT structures

We will now find conditions under which an S^1 -bundle over a $(4n+3)$ -dimensional manifold endowed with three almost contact metric structures is hyper-Kähler with torsion (HKT, for short). Recall that a $4n$ -dimensional hyper-Hermitian manifold $(M^{4n}, J_1, J_2, J_3, h)$ is a hypercomplex manifold (M^{4n}, J_1, J_2, J_3) endowed with a Riemannian metric h compatible with the complex structures J_r for $r = 1, 2, 3$; that is, h satisfies

$$h(J_r X, J_r Y) = h(X, Y)$$

for any $r = 1, 2, 3$ and any vector fields X and Y on M^{4n} .

A hyper-Hermitian manifold $(M^{4n}, J_1, J_2, J_3, h)$ is called HKT if and only if

$$(37) \quad J_1 dF_1 = J_2 dF_2 = J_3 dF_3,$$

where F_r denotes the fundamental 2-form associated to the Hermitian structure (J_r, h) ; see [Grantcharov and Poon 2000].

We consider a $(4n+3)$ -dimensional manifold N^{4n+3} endowed with three almost contact metric structures (I_r, ξ_r, η_r, g) for $r = 1, 2, 3$, and satisfying

$$(38) \quad \begin{aligned} I_k &= I_i I_j - \eta_j \otimes \xi_i = -I_j I_i + \eta_i \otimes \xi_j, \\ \xi_k &= I_i \xi_j = -I_j \xi_i, \quad \eta_k = \eta_i I_j = -\eta_j I_i. \end{aligned}$$

By applying Theorem 2.3, we can construct hyper-Hermitian structures on S^1 -bundles over N^{4n+3} and study when they are strong HKT.

Theorem 7.1. *Let N^{4n+3} be a $(4n+3)$ -dimensional manifold with three normal almost contact metric structures (I_r, ξ_r, η_r, g) for $r = 1, 2, 3$, and satisfying (38). Let Ω be a closed 2-form on N^{4n+3} that represents an integral cohomology class, and that is I_r -invariant for every $r = 1, 2, 3$. Consider the circle bundle $S^1 \hookrightarrow P \rightarrow N^{4n+3}$ with a connection 1-form θ whose curvature form is $d\theta = \pi^*(\Omega)$, where $\pi : P \rightarrow N$ is the projection. The hyper-Hermitian structure (J_1, J_2, J_3, h) on P defined by (2) and (4) is HKT if and only if*

$$(39) \quad \begin{aligned} \pi^*(I_1(d\omega_1)) - \pi^*(d\eta_1) \wedge \pi^*\eta_1 &= \pi^*(I_2(d\omega_2)) - \pi^*(d\eta_2) \wedge \pi^*\eta_2 \\ &= \pi^*(I_3(d\omega_3)) - \pi^*(d\eta_3) \wedge \pi^*\eta_3, \end{aligned}$$

$$\pi^*(I_1(i_{\xi_1} d\omega_1)) = \pi^*(I_2(i_{\xi_2} d\omega_2)) = \pi^*(I_3(i_{\xi_3} d\omega_3)),$$

where ω_r is the fundamental form of the almost contact structure (I_r, ξ_r, η_r, g) . Moreover, the HKT structure is strong if and only if

$$(40) \quad \begin{aligned} d(\pi^*(I_r(i_{\xi_r} d\omega_r))) &= 0, \\ d(\pi^*(I_r(d\omega_r) - d\eta_r \wedge \eta_r)) &= (\pi^*(-I_r(i_{\xi_r} d\omega_r)) + \pi^*\Omega) \wedge \pi^*\Omega \end{aligned}$$

for every $r = 1, 2, 3$.

Proof. The almost hyper-Hermitian structure (J_1, J_2, J_3, h) on P defined by (2) and (4) is hyper-Hermitian if and only if (I_r, ξ_r, η_r, g) is normal and $d\theta$ is J_r -invariant for every $r = 1, 2, 3$. The HKT condition is equivalent to (37). By (9), we have

$$J_r dF_r = \pi^*(I_r(d\omega_r)) + \pi^*(I_r(i_{\xi_r}d\omega_r)) \wedge \theta - \pi^*(d\eta_r) \wedge \pi^*\eta_r - \theta \wedge d\theta,$$

where F_r is the fundamental 2-form of (J_r, h) . Therefore, condition (37) is satisfied if and only if (39) holds. Finally, the $J_r dF_r$ are closed if and only if (40) holds. \square

On $N^{4n+3} \times \mathbb{R}$, consider the almost Hermitian structures (J_r, F_r, h) defined by

$$(41) \quad \begin{aligned} h &= g + (dt)^2, & F_r &= \omega_r + \eta_r \wedge dt, \\ J_r(\eta_r) &= dt, & J_r(X) &= I_r(X) \quad \text{for } X \in \text{Ker } \eta_r. \end{aligned}$$

By (38), we have

$$\begin{aligned} J_1 J_2 &= J_3 = -J_2 J_1, \\ J_1 \eta_2 &= I_1 \eta_2 = -\eta_3, & J_2 \eta_3 &= I_2 \eta_3 = -\eta_1, & J_3 \eta_1 &= I_3 \eta_1 = -\eta_2. \end{aligned}$$

Therefore, (J_r, F_r, h) for $r = 1, 2, 3$ is a hyper-Hermitian structure on $N^{4n+3} \times \mathbb{R}$ if and only if the structures (I_r, ξ_r, η_r, g) are normal.

Corollary 7.2. *Let N^{4n+3} be a $(4n+3)$ -dimensional manifold endowed with three normal almost contact metric structures (I_r, ξ_r, η_r, g) for $r = 1, 2, 3$. On the product $N^{4n+3} \times \mathbb{R}$, consider the hyper-Hermitian structure (J_1, J_2, J_3, h) defined by (41). Then, (J_1, J_2, J_3, h) is HKT if and only if*

$$\begin{aligned} I_1(d\omega_1) - d\eta_1 \wedge \eta_1 &= I_2(d\omega_2) - d\eta_2 \wedge \eta_2 = I_3(d\omega_3) - d\eta_3 \wedge \eta_3, \\ I_1(i_{\xi_1}d\omega_1) &= I_2(i_{\xi_2}d\omega_2) = I_3(i_{\xi_3}d\omega_3). \end{aligned}$$

The HKT structure is strong if and only if

$$d(I_r(i_{\xi_r}d\omega_r)) = 0 \quad \text{and} \quad d(I_r(d\omega_r) - d\eta_r \wedge \eta_r) = 0 \quad \text{for every } r = 1, 2, 3.$$

Moreover, if (J_1, J_2, J_3, h) is such that $d\eta_1 \wedge \eta_1 = d\eta_2 \wedge \eta_2 = d\eta_3 \wedge \eta_3$ and one of the conditions

- (a) $d\omega_r = 0$ for any $r = 1, 2, 3$, that is, (I_r, ξ_r, η_r, g) is quasi-Sasakian for any $r = 1, 2, 3$; or
- (b) $d\omega_i \wedge \eta_j \wedge \eta_k \neq 0$, where (i, j, k) is a permutation of $(1, 2, 3)$, as well as

$$I_1(d\omega_1) = I_2(d\omega_2) = I_3(d\omega_3) \quad \text{and} \quad I_1(i_{\xi_1}d\omega_1) = I_2(i_{\xi_2}d\omega_2) = I_3(i_{\xi_3}d\omega_3),$$

is satisfied, then (J_1, J_2, J_3, h) is HKT. In case (a), the HKT structure is strong. In case (b), the HKT structure is strong if and only if $d(I_1(d\omega_1)) = d(I_1(i_{\xi_1}d\omega_1)) = 0$.

Proof. By Theorem 7.1, the hyper-Hermitian structure (J_r, F_r, h) for $r = 1, 2, 3$ is HKT if and only if

$$(42) \quad \begin{aligned} I_1(d\omega_1) - d\eta_1 \wedge \eta_1 &= I_2(d\omega_2) - d\eta_2 \wedge \eta_2 = I_3(d\omega_3) - d\eta_3 \wedge \eta_3, \\ I_1(i_{\xi_1}d\omega_1) &= I_2(i_{\xi_2}d\omega_2) = I_3(i_{\xi_3}d\omega_3). \end{aligned}$$

Locally, we write

$$(43) \quad d\omega_r = \alpha_r + \sum_{i=1}^3 \beta_i^r \wedge \eta_i + \sum_{i<j=1}^3 \gamma_{ij}^r \wedge \eta_i \wedge \eta_j + \rho_r \eta_1 \wedge \eta_2 \wedge \eta_3,$$

where ρ_r are smooth functions, while α_r , β_i^r , and γ_{ij}^r are respectively 3-forms, 2-forms, and 1-forms in $\bigcap_{i=1}^3 \text{Ker } \eta_i$.

By first using the normality of the three almost contact metric structures, and then that $i_{\xi_r}d\eta_r = 0$ and $I_r(d\eta_r) = d\eta_r$, locally we can write

$$(44) \quad \begin{aligned} d\eta_1 &= A_1 + B_1 \wedge \eta_2 - I_1 B_1 \wedge \eta_3 + C_1 \eta_2 \wedge \eta_3, \\ d\eta_2 &= A_2 + B_2 \wedge \eta_1 + I_2 B_2 \wedge \eta_3 + C_2 \eta_1 \wedge \eta_3, \\ d\eta_3 &= A_3 + B_3 \wedge \eta_1 - I_3 B_3 \wedge \eta_2 + C_3 \eta_1 \wedge \eta_2, \end{aligned}$$

where $I_r A_r = A_r$. Here, the A_r and B_r are respectively 2-forms and 1-forms in $\bigcap_{i=1}^3 \text{Ker } \eta_i$, while the C_r are smooth functions. We have

$$J_r(dF_r) = J_r(d\omega_r) + J_r(d\eta_r \wedge dt) = J_r(d\omega_r) - d\eta_r \wedge \eta_r.$$

Therefore, by using (43) and (44), we obtain

$$\begin{aligned} J_1(dF_1) &= I_1\alpha_1 + I_1\beta_1^1 \wedge dt - A_1 \wedge \eta_1 - I_1\beta_3^1 \wedge \eta_2 - I_1\beta_2^1 \wedge \eta_3 \\ &\quad - I_1\gamma_{13}^1 \wedge \eta_2 \wedge dt + I_1\gamma_{12}^1 \wedge \eta_3 \wedge dt + B_1 \wedge \eta_1 \wedge \eta_2 - I_1 B_1 \wedge \eta_1 \wedge \eta_3 \\ &\quad + I_1\gamma_{23}^1 \wedge \eta_2 \wedge \eta_3 + \rho_1 \eta_2 \wedge \eta_3 \wedge dt - C_1 \eta_1 \wedge \eta_2 \wedge \eta_3, \\ J_2(dF_2) &= I_2\alpha_2 + I_2\beta_2^2 \wedge dt - I_2\beta_3^2 \wedge \eta_1 - A_2 \wedge \eta_2 + I_2\beta_1^2 \wedge \eta_3 \\ &\quad + I_2\gamma_{23}^2 \wedge \eta_1 \wedge dt + I_2\gamma_{12}^2 \wedge \eta_3 \wedge dt - B_2 \wedge \eta_1 \wedge \eta_2 + I_2\gamma_{13}^2 \wedge \eta_1 \wedge \eta_3 \\ &\quad + I_2 B_2 \wedge \eta_2 \wedge \eta_3 - \rho_2 \eta_1 \wedge \eta_3 \wedge dt + C_2 \eta_1 \wedge \eta_2 \wedge \eta_3, \\ J_3(dF_3) &= I_3\alpha_3 + I_3\beta_3^3 \wedge dt + I_3\beta_2^3 \wedge \eta_1 - I_3\beta_1^3 \wedge \eta_2 - A_3 \wedge \eta_3 \\ &\quad + I_3\gamma_{23}^3 \wedge \eta_1 \wedge dt - I_3\gamma_{13}^3 \wedge \eta_2 \wedge dt + I_3\gamma_{12}^3 \wedge \eta_1 \wedge \eta_2 - B_3 \wedge \eta_1 \wedge \eta_3 \\ &\quad + I_3 B_3 \wedge \eta_2 \wedge \eta_3 + \rho_3 \eta_1 \wedge \eta_2 \wedge dt - C_3 \eta_1 \wedge \eta_2 \wedge \eta_3. \end{aligned}$$

The conditions (42) are satisfied if and only if

$$\begin{aligned}
 & \gamma_{12}^1 = \gamma_{13}^1 = \gamma_{12}^2 = \gamma_{23}^2 = \gamma_{13}^3 = \gamma_{23}^3 = 0, \\
 & \rho_r = 0, \quad C_1 = -C_2 = C_3, \\
 (45) \quad & I_1\alpha_1 = I_2\alpha_2 = I_3\alpha_3, \quad I_1\beta_1^1 = I_2\beta_2^2 = I_3\beta_3^3, \\
 & A_1 = I_2\beta_3^2 = -I_3\beta_2^3, \quad A_2 = -I_1\beta_3^1 = I_3\beta_1^3, \quad A_3 = I_1\beta_2^1 = -I_2\beta_1^2, \\
 & B_1 = -B_2 = I_3\gamma_{12}^3, \quad -I_1B_1 = -B_3 = I_2\gamma_{13}^2, \quad I_2B_2 = I_3B_3 = I_1\gamma_{23}^1.
 \end{aligned}$$

Since $I_r A_r = A_r$ the coefficients β_i^r for $r \neq i = 1, 2, 3$ must satisfy the conditions

$$I_i(\beta_j^i - I_k\beta_j^i) = 0 \quad \text{for all } i, j, k = 1, 2, 3 \text{ with } i \neq j, j \neq k \text{ and } k \neq i.$$

The last three equations in (45) are satisfied if and only if $\gamma_{23}^1 = \gamma_{13}^2 = \gamma_{12}^3 = 0$.

Thus, finally, we obtain

$$\begin{aligned}
 & d\omega_r = \alpha_r + \sum_{i=1}^3 \beta_i^r \wedge \eta_i, \quad d\eta_i = A_i + \lambda \eta_j \wedge \eta_k, \\
 (46) \quad & 0 = I_i(\beta_j^i - I_k\beta_j^i) \quad \text{for all } i, j, k = 1, 2, 3 \text{ with } i \neq j, j \neq k \text{ and } k \neq i, \\
 & I_1\alpha_1 = I_2\alpha_2 = I_3\alpha_3, \\
 & A_1 = I_2\beta_3^2 = -I_3\beta_2^3, \quad A_2 = -I_1\beta_3^1 = I_3\beta_1^3, \quad A_3 = I_1\beta_2^1 = -I_2\beta_1^2
 \end{aligned}$$

for any even permutation of $(1, 2, 3)$.

The expression for $d(J_1 dF_1)$ is

$$\begin{aligned}
 d(J_1 dF_1) &= d(I_1(d\omega_1) + I_1(i_{\xi_1} d\omega_1) \wedge dt) - d((d\eta_1) \wedge \eta_1) \\
 &= d(I_1(d\omega_1)) + d(I_1(i_{\xi_1} d\omega_1)) \wedge dt - d\eta_1 \wedge d\eta_1 \\
 &= d(I_1(d\omega_1) - d\eta_1 \wedge \eta_1) + d(I_1(i_{\xi_1} d\omega_1)) \wedge dt,
 \end{aligned}$$

and thus the HKT structure is strong if and only if

$$d(I_1(d\omega_1) - d\eta_1 \wedge \eta_1) = 0 \quad \text{and} \quad d(I_1(i_{\xi_1} d\omega_1)) = 0.$$

To prove the last part of the corollary it suffices to consider coefficients $\beta_r^i = 0$ if $r \neq i$ in expression (43). \square

Example 7.3. Consider the 7-dimensional Lie group $G = \text{SU}(2) \times \mathbb{R}^4$, with structure equations

$$\begin{aligned}
 de^1 &= -\frac{1}{2}e^{25} - \frac{1}{2}e^{36} - \frac{1}{2}e^{47}, & de^5 &= e^{67}, \\
 de^2 &= \frac{1}{2}e^{15} + \frac{1}{2}e^{37} - \frac{1}{2}e^{46}, & de^6 &= -e^{57}, \\
 de^3 &= \frac{1}{2}e^{16} - \frac{1}{2}e^{27} + \frac{1}{2}e^{45}, & de^7 &= e^{56}, \\
 de^4 &= \frac{1}{2}e^{17} + \frac{1}{2}e^{26} - \frac{1}{2}e^{35},
 \end{aligned}$$

By [Fino and Tomassini 2008], G admits a compact quotient $M^7 = \Gamma \backslash G$ by a uniform discrete subgroup Γ , and is endowed with a weakly generalized G_2 -structure. By [Barberis and Fino 2008], $M^7 \times S^1$ admits a strong HKT structure. We can show that M^7 has three normal almost contact metric structures (I_r, ξ_r, η_r, g) for $r = 1, 2, 3$ that are given by

$$\begin{aligned} I_1 e^1 &= e^2, & I_1 e^3 &= e^4, & I_1 e^5 &= e^6, & \eta_1 &= e^7, \\ I_2 e^1 &= e^3, & I_2 e^2 &= -e^4, & I_2 e^5 &= -e^7, & \eta_2 &= e^6, \\ I_3 e^1 &= e^4, & I_3 e^2 &= e^3, & I_3 e^6 &= e^7, & \eta_3 &= e^5 \end{aligned}$$

and that satisfy the conditions of Corollary 7.2(a).

References

- [Andrada et al. 2009] A. Andrada, A. Fino, and L. Vezzoni, “A class of Sasakian 5-manifolds”, *Transformation Groups* **14**:3 (2009), 493–512. MR 2534796 Zbl 1185.53046 arXiv 0807.1800
- [Barberis and Fino 2008] M. L. Barberis and A. Fino, “New HKT manifolds arising from quaternionic representations”, preprint, 2008. To appear in *Math. Z.* arXiv 0805.2335
- [Barberis et al. 2009] M. L. Barberis, I. G. Dotti, and M. Verbitsky, “Canonical bundles of complex nilmanifolds, with applications to hypercomplex geometry”, *Math. Res. Lett.* **16**:2 (2009), 331–347. MR 2010g:53083 Zbl 1178.32014
- [Bismut 1989] J.-M. Bismut, “A local index theorem for non-Kähler manifolds”, *Math. Ann.* **284**:4 (1989), 681–699. MR 91i:58140
- [Blair 1967] D. E. Blair, “The theory of quasi-Sasakian structures”, *J. Differential Geometry* **1** (1967), 331–345. MR 37 #2127 Zbl 0163.43903
- [Blair 2002] D. E. Blair, *Riemannian geometry of contact and symplectic manifolds*, Progress in Mathematics **203**, Birkhäuser, Boston, MA, 2002. MR 2002m:53120 Zbl 1011.53001
- [Boyer and Galicki 1999] C. Boyer and K. Galicki, “3-Sasakian manifolds”, pp. 123–184 in *Surveys in differential geometry: Essays on Einstein manifolds*, edited by C. LeBrun and M. Wang, Surv. Differ. Geom. **6**, Int. Press, Boston, 1999. MR 2001m:53076 Zbl 1008.53047
- [Conti and Salamon 2007] D. Conti and S. Salamon, “Generalized Killing spinors in dimension 5”, *Trans. Amer. Math. Soc.* **359**:11 (2007), 5319–5343. MR 2008h:53077 Zbl 1130.53033
- [Egidi 2001] N. Egidi, “Special metrics on compact complex manifolds”, *Differential Geom. Appl.* **14**:3 (2001), 217–234. MR 2002b:32038 Zbl 0981.32014
- [Fernández et al. 2009] M. Fernández, A. Tomassini, L. Ugarte, and R. Villacampa, “Balanced Hermitian metrics from SU(2)-structures”, *J. Math. Phys.* **50**:3 (2009), 033507, 15. MR 2010f:53124 Zbl 05646541
- [Fino and Grantcharov 2004] A. Fino and G. Grantcharov, “Properties of manifolds with skew-symmetric torsion and special holonomy”, *Adv. Math.* **189**:2 (2004), 439–450. MR 2005k:53062 Zbl 1114.53043
- [Fino and Tomassini 2008] A. Fino and A. Tomassini, “Generalized G_2 -manifolds and SU(3)-structures”, *Internat. J. Math.* **19**:10 (2008), 1147–1165. MR 2009m:53054 Zbl 1168.53012
- [Fino and Tomassini 2009] A. Fino and A. Tomassini, “Blow-ups and resolutions of strong Kähler with torsion metrics”, *Adv. Math.* **221**:3 (2009), 914–935. MR 2010d:32012

- [Fino et al. 2004] A. Fino, M. Parton, and S. Salamon, “Families of strong KT structures in six dimensions”, *Comment. Math. Helv.* **79**:2 (2004), 317–340. MR 2005f:53061 Zbl 1062.53062
- [Friedrich and Ivanov 2002] T. Friedrich and S. Ivanov, “Parallel spinors and connections with skew-symmetric torsion in string theory”, *Asian J. Math.* **6**:2 (2002), 303–335. MR 2003m:53070 Zbl 1127.53304
- [Gates et al. 1984] S. J. Gates, Jr., C. M. Hull, and M. Roček, “Twisted multiplets and new supersymmetric nonlinear σ -models”, *Nuclear Phys. B* **248**:1 (1984), 157–186. MR 87b:81108
- [Gauduchon 1984] P. Gauduchon, “La 1-forme de torsion d’une variété hermitienne compacte”, *Math. Ann.* **267**:4 (1984), 495–518. MR 87a:53101 Zbl 0523.53059
- [Grantcharov and Poon 2000] G. Grantcharov and Y. S. Poon, “Geometry of hyper-Kähler connections with torsion”, *Comm. Math. Phys.* **213**:1 (2000), 19–37. MR 2002a:53059
- [Grantcharov et al. 2008] D. Grantcharov, G. Grantcharov, and Y. S. Poon, “Calabi–Yau connections with torsion on toric bundles”, *J. Differential Geom.* **78**:1 (2008), 13–32. MR 2009g:32052 Zbl 1171.53044
- [Hitchin 2001] N. Hitchin, “Stable forms and special metrics”, pp. 70–89 in *Global differential geometry: The mathematical legacy of Alfred Gray* (Bilbao, 2000), edited by M. Fernández and J. A. Wolf, Contemp. Math. **288**, Amer. Math. Soc., Providence, RI, 2001. MR 2003f:53065 Zbl 1004.53034
- [Howe and Papadopoulos 1996] P. S. Howe and G. Papadopoulos, “Twistor spaces for hyper-Kähler manifolds with torsion”, *Phys. Lett. B* **379**:1-4 (1996), 80–86. MR 97h:53073
- [Ivanov and Papadopoulos 2001] S. Ivanov and G. Papadopoulos, “Vanishing theorems and string backgrounds”, *Classical Quantum Gravity* **18**:6 (2001), 1089–1110. MR 2002h:53076 Zbl 0990.53078
- [Kobayashi 1956] S. Kobayashi, “Principal fibre bundles with the 1-dimensional toroidal group”, *Tôhoku Math. J. (2)* **8** (1956), 29–45. MR 18,328a Zbl 0075.32103
- [Ogawa 1963] Y. Ogawa, “Some properties on manifolds with almost contact structures”, *Tôhoku Math. J. (2)* **15** (1963), 148–161. MR 27 #704 Zbl 0147.41002
- [Sasaki and Hatakeyama 1961] S. Sasaki and Y. Hatakeyama, “On differentiable manifolds with certain structures which are closely related to almost contact structure, II”, *Tôhoku Math. J. (2)* **13** (1961), 281–294. MR 25 #1513 Zbl 0112.14002
- [Strominger 1986] A. Strominger, “Superstrings with torsion”, *Nuclear Phys. B* **274**:2 (1986), 253–284. MR 87m:81177
- [Swann 2010] A. Swann, “Twisting Hermitian and hypercomplex geometries”, *Duke Math. J.* **155** (2010), 403–431.
- [Ugarte 2007] L. Ugarte, “Hermitian structures on six-dimensional nilmanifolds”, *Transformation Groups* **12**:1 (2007), 175–202. MR 2008e:53139 Zbl 1129.53052

Received September 23, 2009.

MARISA FERNÁNDEZ
 UNIVERSIDAD DEL PAÍS VASCO
 DEPARTAMENTO DE MATEMÁTICAS
 FACULTAD DE CIENCIA Y TECNOLOGÍA
 APARTADO 644
 48080 BILBAO
 SPAIN
 marisa.fernandez@ehu.es

ANNA FINO
UNIVERSITÀ DI TORINO
DIPARTIMENTO DI MATEMATICA
VIA CARLO ALBERTO 10
I-10123 TORINO
ITALY
annamaria.fino@unito.it

LUIS UGARTE
UNIVERSIDAD DE ZARAGOZA
DEPARTAMENTO DE MATEMÁTICAS-IUMA
CAMPUS PLAZA SAN FRANCISCO
50009 ZARAGOZA
SPAIN
ugarte@unizar.es

RAQUEL VILLACAMPA
CENTRO UNIVERSITARIO DE LA DEFENSA
ACADEMIA GENERAL MILITAR
CRTA. DE HUESCA S/N
50090 ZARAGOZA
SPAIN
raquelvg@unizar.es

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 249 No. 1 January 2011

Metabelian $SL(n, \mathbb{C})$ representations of knot groups, II: Fixed points	1
HANS U. BODEN and STEFAN FRIEDL	
Lewis–Zagier correspondence for higher-order forms	11
ANTON DEITMAR	
Topology of positively curved 8-dimensional manifolds with symmetry	23
ANAND DESSAI	
Strong Kähler with torsion structures from almost contact manifolds	49
MARISA FERNÁNDEZ, ANNA FINO, LUIS UGARTE and RAQUEL VILLACAMPA	
Connections between Floer-type invariants and Morse-type invariants of Legendrian knots	77
MICHAEL B. HENRY	
A functional calculus for unbounded generalized scalar operators on Banach spaces	135
DRAGOLJUB KEČKIĆ and ĐORĐE KRTINIĆ	
Geometric formality of homogeneous spaces and of biquotients	157
D. KOTSCHICK and S. TERZIĆ	
Positive solutions for a nonlinear third order multipoint boundary value problem	177
YANG LIU, ZHANG WEIGUO, LIU XIPING, SHEN CHUNFANG and CHEN HUA	
The braid group surjects onto G_2 tensor space	189
SCOTT MORRISON	
Analogues of the Wiener Tauberian and Schwartz theorems for radial functions on symmetric spaces	199
E. K. NARAYANAN and ALLADI SITARAM	
Semidirect products of representations up to homotopy	211
YUNHE SHENG and CHENCHANG ZHU	
Homology sequence and excision theorem for Euler class group	237
YONG YANG	
Acknowledgement	255



0030-8730(201101)249:1;1-D