

Pacific Journal of Mathematics

**ON THE NUMBER OF PAIRS OF POSITIVE INTEGERS
 $x_1, x_2 \leq H$ SUCH THAT x_1x_2 IS A k -TH POWER**

DOYCHIN I. TOLEV

ON THE NUMBER OF PAIRS OF POSITIVE INTEGERS $x_1, x_2 \leq H$ SUCH THAT x_1x_2 IS A k -TH POWER

DOYCHIN I. TOLEV

We find an asymptotic formula for the number of pairs of positive integers $x_1, x_2 \leq H$ such that the product x_1x_2 is a k -th power.

1. Notation

Let H be a sufficiently large positive number and $k \geq 2$ be a fixed integer. By the letters $j, l, m, n, u, v, x, y, z$ we denote positive integers. The letter p is reserved for primes, and \prod_p denotes a product over all primes. By the letters s and w , we denote complex numbers, and $i = \sqrt{-1}$. By ε we denote an arbitrary small positive number. The constants in the Vinogradov and Landau symbols are absolute or depend on ε and k . As usual, $\zeta(s)$ is the Riemann zeta function. By V_k we denote the set of k -free numbers (that is, positive integers not divided by a k -th power of a prime), and N_k is the set of k -th powers of natural numbers. We denote by $\mu(n)$ the Möbius function and by $\tau(n)$ the number of positive divisors of n . Further, we define $\eta(n) = \prod_{p|n} p$. We write (u, v) for the greatest common divisor of u and v . We assume that $\min(1, 0^{-1}) = 1$.

2. Introduction and statement of the result

Let $S_k(H)$ be the number of pairs of positive integers $x_1, x_2 \leq H$ whose product x_1x_2 is in N_k . We will establish an asymptotic formula for $S_k(H)$. This problem is related to a result of Heath-Brown and Moroz [1999]. They considered the diophantine equation $x_1x_2x_3 = x_0^3$ and found an asymptotic formula for the number of primitive solutions such that $1 \leq x_1, x_2, x_3 \leq H$.

It is easy to find an asymptotic formula for the quantity

$$S_k^*(H) = \#\{x_1, x_2 \mid x_1, x_2 \leq H, (x_1, x_2) = 1, x_1x_2 \in N_k\}.$$

Supported by Sofia University grant 028/2009.

MSC2000: 11D45.

Keywords: counting solutions of Diophantine equations.

Indeed, if $(x_1, x_2) = 1$, then $x_1 x_2 \in N_k$ exactly when $x_1 \in N_k$ and $x_2 \in N_k$. Hence

$$S_k^*(H) = \#\{x_1, x_2 \mid x_1, x_2 \leq H, (x_1, x_2) = 1, x_1 \in N_k, x_2 \in N_k\}$$

$$= \sum_{\substack{z_1, z_2 \leq H^{1/k}, \\ (z_1, z_2) = 1}} 1,$$

and using the well-known property of the Möbius function we get

$$S_k^*(H) = \sum_{z_1, z_2 \leq H^{1/k}} \sum_{d \mid (z_1, z_2)} \mu(d) = \sum_{d \leq H^{1/k}} \mu(d) \left(\frac{H^{1/k}}{d} + O(1) \right)^2.$$

Therefore

$$(1) \quad \begin{aligned} S_k^*(H) &= H^{2/k} \sum_{d \leq H^{1/k}} \frac{\mu(d)}{d^2} + O(H^{1/k} \log H) \\ &= \zeta(2)^{-1} H^{2/k} + O(H^{1/k} \log H). \end{aligned}$$

It is also easy to evaluate $S_2(H)$. Indeed, we have

$$S_2(H) = \sum_{d \leq H} \sum_{\substack{x_1, x_2 \leq H, \\ (x_1, x_2) = d, \\ x_1 x_2 \in N_2}} 1 = \sum_{d \leq H} \sum_{\substack{y_1, y_2 \leq H/d, \\ (y_1, y_2) = 1, \\ y_1 y_2 d^2 \in N_2}} 1 = \sum_{d \leq H} S_2^*(H/d).$$

Now we apply (1) and after calculations that we leave to the reader, we find

$$S_2(H) = \zeta(2)^{-1} H \log H + O(H).$$

However it is not clear how to apply (1) in order to evaluate $S_k(H)$ for $k \geq 3$.

Another quantity related to $S_k(H)$ is

$$T_k(H) = \#\{x_1, x_2 \mid x_1 x_2 \leq H^2, x_1 x_2 \in N_k\} = \sum_{n \leq H^{2/k}} \tau(n^k).$$

Using well-known analytic methods, based on Perron’s formula and the simplest properties of $\zeta(s)$, we are able to prove the asymptotic formula

$$T_k(H) \sim \gamma_k H^{2/k} (\log H)^k,$$

where $\gamma_k > 0$ depends only on k . In this paper we show that using the same analytic tools, as well as an idea of Heath-Brown and Moroz [1999], we may find an asymptotic formula for $S_k(H)$ for any $k \geq 2$:

Theorem. *For any integer $k \geq 2$, we have*

$$(2) \quad S_k(H) = c_k H^{2/k} (\log H)^{k-1} + O(H^{2/k} (\log H)^{k-2}),$$

where

$$(3) \quad c_k = \frac{\mathcal{P}_k}{((k-1)!)^2} \left(1 + \frac{1}{k^{k-2}} \sum_{k/2 < m \leq k-1} \frac{(-1)^{k-m} (2m-k)^{k-1} \binom{k-1}{m}}{k-m} \right),$$

$$(4) \quad \mathcal{P}_k = \prod_p \left(1 - \frac{1}{p} \right)^{k-1} \left(1 + \frac{k-1}{p} \right).$$

3. Some lemmas

Lemma 1. (i) Every positive integer x can be represented uniquely in the form $x = yz$, where $y \in V_k$ and $z \in N_k$.

(ii) Every integer $y \in V_k$ can be written uniquely in the form $y = u_1 u_2^2 u_3^3 \cdots u_{k-1}^{k-1}$, where $u_j \in V_2$ for $1 \leq j \leq k-1$ and $(u_i, u_j) = 1$ for $1 \leq i, j \leq k-1, i \neq j$.

(iii) If $y_1, y_2 \in V_k$ and $y_1 y_2 \in N_k$, then $\eta(y_1) = \eta(y_2) = (y_1 y_2)^{1/k}$.

Proof. The proofs of (i) and (ii) can be obtained easily from the fundamental theorem of arithmetic and we leave this to the reader. Let us prove (iii). By our assumption, any prime in the factorization of $y_1 y_2$ occurs with exponent at most $2k-2$, and hence with exponent exactly k . Since the exponent of each prime in y_1 and y_2 is $\leq k-1$, the integers y_1 and y_2 have the same prime factors. \square

The next lemma is a version of the Perron formula. Denote

$$(5) \quad E(\gamma) = \begin{cases} 1 & \text{if } \gamma \geq 1, \\ 0 & \text{if } 0 < \gamma < 1. \end{cases}$$

Lemma 2. If $\gamma > 0, 0 < c < c_0$ and $T > 1$, then

$$E(\gamma) = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \frac{\gamma^s}{s} ds + O(\gamma^c \min(1, T^{-1} |\log \gamma|^{-1})).$$

The constant in the Landau symbol depends only on c_0 .

Proof. This is a slightly simplified version of a lemma from [Davenport 2000, Section 17]. \square

Some of the basic properties of Riemann’s zeta function are presented in the next lemma.

Lemma 3. (i) $\zeta(s)$ is meromorphic in the complex plane and has a pole only at $s = 1$. It is simple and with a residue equal to 1.

(ii) If $\text{Re}(s) > 1$, then $\zeta(s) = \prod_p (1 - p^{-s})^{-1}$.

(iii) If $\text{Re}(s) \geq \sigma > 1$, then $\zeta(s) \ll (\sigma - 1)^{-1} + 1$.

(iv) If $1/2 \leq \sigma_0 \leq 1, \sigma \geq \sigma_0$ and $|t| \geq 2$, then $\zeta(\sigma + it) \ll |t|^{(1-\sigma_0)/2+\varepsilon}$.

(v) *There exist $\lambda_0 > 0$ such that if $X \geq 2$, $|t| \leq X$ and $\sigma \geq 1 - \lambda_0/\log X$, then $\zeta(\sigma + it) \neq 0$.*

Proof. See [Titchmarsh 1986, Chapters 1–3 and 5]. □

4. Proof of the theorem

4.1. We already considered the case $k = 2$, so we may assume that $k \geq 3$.

Working as in [Heath-Brown and Moroz 1999] we apply Lemma 1(i) and find that $S_k(H)$ is equal to the number of quadruples y_1, y_2, z_1, z_2 such that

$$y_1, y_2 \in V_k, \quad z_1, z_2 \in N_k, \quad y_1 z_1 \leq H, \quad y_2 z_2 \leq H, \quad y_1 z_1 y_2 z_2 \in N_k.$$

Obviously the last of the above conditions is equivalent to $y_1 y_2 \in N_k$ because z_1 and z_2 are k -th powers. Hence

$$S_k(H) = \sum_{\substack{y_1, y_2 \leq H, \\ y_1, y_2 \in V_k, \\ y_1 y_2 \in N_k}} \sum_{\substack{m_j \leq (H/y_j)^{1/k}, \\ j=1,2}} 1 = \sum_{\substack{y_1, y_2 \leq H, \\ y_1, y_2 \in V_k, \\ y_1 y_2 \in N_k}} ((H/y_1)^{1/k} + O(1))((H/y_2)^{1/k} + O(1)).$$

Expanding brackets, we get

$$(6) \quad S_k(H) = H^{2/k} U_k(H) + O(H^{1/k} W_k(H)),$$

where

$$U_k(H) = \sum_{\substack{y_1, y_2 \leq H, \\ y_1, y_2 \in V_k, \\ y_1 y_2 \in N_k}} (y_1 y_2)^{-1/k} \quad \text{and} \quad W_k(H) = \sum_{\substack{y_1, y_2 \leq H, \\ y_1, y_2 \in V_k, \\ y_1 y_2 \in N_k}} y_1^{-1/k}.$$

Using Lemma 1(iii), we see that for a given y_1 the integer y_2 is determined uniquely. Therefore we have

$$(7) \quad U_k(H) = \sum_{\substack{y \leq H, \\ y \in V_k, \\ \eta(y)^k \leq Hy}} \eta(y)^{-1} \quad \text{and} \quad W_k(H) = \sum_{\substack{y \leq H, \\ y \in V_k, \\ \eta(y)^k \leq Hy}} y^{1/k} \eta(y)^{-1}.$$

To prove the theorem we have to find an asymptotic formula for $U_k(H)$ and to estimate $W_k(H)$.

4.2. Consider first $W_k(H)$. Applying Lemma 1(ii), we get

$$\begin{aligned} W_k(H) &\leq \sum_{u_1 u_2^2 \cdots u_{k-1}^{k-1} \leq H} \frac{(u_1 u_2^2 \cdots u_{k-1}^{k-1})^{1/k}}{u_1 u_2 \cdots u_{k-1}} \\ &= \sum_{u_1 u_2^2 \cdots u_{k-2}^{k-2} \leq H} u_1^{-1+1/k} u_2^{-1+2/k} \cdots u_{k-2}^{-1+(k-2)/k} \sum_{u_{k-1} \leq \left(\frac{H}{u_1 u_2^2 \cdots u_{k-2}^{k-2}}\right)^{1/(k-1)}} u_{k-1}^{-1/k}. \end{aligned}$$

The inner sum is $\ll H^{1/k} (u_1 u_2^2 \dots u_{k-2}^{k-2})^{-1/k}$; hence

$$(8) \quad W_k(H) \ll H^{1/k} \sum_{u_1 u_2^2 \dots u_{k-2}^{k-2} \leq H} (u_1 u_2 \dots u_{k-2})^{-1} \ll H^{1/k} (\log H)^{k-2}.$$

It remains to show that

$$(9) \quad U_k(H) = c_k (\log H)^{k-1} + O((\log H)^{k-2}).$$

Formula (2) is a consequence of (6), (8) and (9).

4.3. Using (5) and (7), we write $U_k(H)$ in the form

$$U_k(H) = \sum_{\substack{y \leq H, \\ y \in V_k}} \eta(y)^{-1} E(Hy\eta(y)^{-k}).$$

We put

$$(10) \quad c = (\log H)^{-1} \quad \text{and} \quad T = (\log H)^{100k^3}$$

and applying Lemma 2 we find that

$$(11) \quad U_k(H) = U^{(1)} + O(\Delta),$$

where

$$(12) \quad U^{(1)} = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \frac{H^s}{s} \Phi(s) ds, \quad \text{and} \quad \Phi(s) = \sum_{\substack{y \leq H, \\ y \in V_k}} y^s \eta(y)^{-ks-1}$$

$$\text{and } \Delta = \sum_{\substack{y \leq H, \\ y \in V_k}} \eta(y)^{-1} \min(1, T^{-1} |\log(Hy\eta(y)^{-k})|^{-1}).$$

4.4. Consider first the sum Δ . We put

$$(13) \quad \varkappa = T^{-1/2}$$

and write

$$(14) \quad \Delta = \Delta_1 + \Delta_2,$$

where in Δ_1 the summation is taken over y satisfying $|\log(Hy\eta(y)^{-k})| \geq \varkappa$ and in Δ_2 over the other y . To estimate Δ_1 we apply Lemma 1(iii), (10) and (13) to find

$$(15) \quad \begin{aligned} \Delta_1 &\ll T^{-1/2} \sum_{\substack{y \leq H, \\ y \in V_k}} \eta(y)^{-1} \ll T^{-1/2} \sum_{u_1 u_2^2 \dots u_{k-1}^{k-1} \leq H} (u_1 u_2 \dots u_{k-1})^{-1} \\ &\ll \frac{(\log H)^{k-1}}{T^{1/2}} \ll 1. \end{aligned}$$

Consider Δ_2 . Using its definition and [Lemma 1\(iii\)](#), we find

$$\begin{aligned} \Delta_2 &\ll \sum_{\substack{u_1, u_2, \dots, u_{k-1}: \\ |\log(H/(u_1^{k-1}u_2^{k-2}\dots u_{k-2}^2u_{k-1}))| < \varkappa}} (u_1u_2 \dots u_{k-1})^{-1} \\ &\ll \sum_{He^{-\varkappa} < u_1^{k-1}u_2^{k-2}\dots u_{k-2}^2u_{k-1} < He^\varkappa} (u_1u_2 \dots u_{k-1})^{-1} \\ &\ll \sum_{u_1^{k-1}u_2^{k-2}\dots u_{k-2}^2 < 2H} (u_1u_2 \dots u_{k-2})^{-1} \sum_{\frac{He^{-\varkappa}}{u_1^{k-1}u_2^{k-2}\dots u_{k-2}^2} < u_{k-1} < \frac{He^\varkappa}{u_1^{k-1}u_2^{k-2}\dots u_{k-2}^2}} u_{k-1}^{-1}. \end{aligned}$$

To estimate the inner sum we apply the obvious inequality

$$(16) \quad \sum_{a < n \leq b} n^{-1} \leq a^{-1} + \log(b/a) \quad \text{for } 0 < a < b$$

and find that

$$(17) \quad \Delta_2 \ll \sum_{u_1^{k-1}u_2^{k-2}\dots u_{k-2}^2 < 2H} \frac{H^{-1}u_1^{k-1}u_2^{k-2}\dots u_{k-2}^2 + \varkappa}{u_1u_2 \dots u_{k-2}} \ll H^{-1}\Delta_3 + \varkappa(\log H)^{k-2},$$

where

$$(18) \quad \Delta_3 = \sum_{u_1^{k-1}u_2^{k-2}\dots u_{k-2}^2 < 2H} u_1^{k-2}u_2^{k-3}\dots u_{k-2}.$$

If $k > 3$, then

$$\begin{aligned} \Delta_3 &\ll \sum_{u_1^{k-1}u_2^{k-2}\dots u_{k-3}^3 < 2H} u_1^{k-2}u_2^{k-3}\dots u_{k-3}^2 \sum_{u_{k-2} < (2H/(u_1^{k-1}u_2^{k-2}\dots u_{k-3}^3))^{1/2}} u_{k-2} \\ (19) \quad &\ll H \sum_{u_1^{k-1}u_2^{k-2}\dots u_{k-3}^3 < 2H} (u_1u_2 \dots u_{k-3})^{-1} \ll H(\log H)^{k-3}. \end{aligned}$$

The last estimate for Δ_3 is obviously true also for $k = 3$. From [\(10\)](#), [\(13\)–\(15\)](#), [\(17\)](#) and [\(19\)](#), we get

$$(20) \quad \Delta \ll (\log H)^{k-3}.$$

4.5. Consider the expression $\Phi(s)$ defined by [\(12\)](#). Let c and T be specified by [\(10\)](#) and

$$(21) \quad T_1 = 2kT.$$

We apply [Lemma 2](#) again and show that if $\text{Re}(s) = c$, then

$$(22) \quad \Phi(s) = \frac{1}{2\pi i} \int_{c-iT_1}^{c+iT_1} \frac{H^w}{w} \mathcal{M}(s, w) dw + O(\Delta^*),$$

where

$$(23) \quad \mathcal{M}(s, w) = \sum_{y=1, y \in V_k}^{\infty} y^{s-w} \eta(y)^{-ks-1},$$

$$(24) \quad \Delta^* = \sum_{y=1, y \in V_k}^{\infty} \eta(y)^{-kc-1} \min(1, T_1^{-1} |\log(H/y)|^{-1}).$$

To justify (22) we note from Euler’s identity, (10) and parts (ii) and (iii) of [Lemma 3](#) it follows that

$$(25) \quad \sum_{\substack{y=1, \\ y \in V_k}}^{\infty} \eta(y)^{-kc-1} = \prod_p \left(1 + \frac{k-1}{p^{kc+1}}\right) \ll \zeta^{k-1}(kc+1) \ll c^{-k+1} \ll (\log H)^{k-1}.$$

Hence $\mathcal{M}(s, w)$ is absolutely and uniformly convergent in $\text{Re}(s) = \text{Re}(w) = c$ because under this assumption we have $\mathcal{M}(s, w) \ll \sum_{y=1, y \in V_k}^{\infty} \eta(y)^{-kc-1}$. This completes the verification of (22).

4.6. Consider the expression Δ^* defined by (24). We write it in the form

$$(26) \quad \Delta^* = \Delta_1^* + \Delta_2^*,$$

where the summation in Δ_1^* is taken over y such that $|\log(H/y)| \geq \varkappa$ and in Δ_2^* over the other y . Using (10), (13), (21) and (25), we find

$$(27) \quad \Delta_1^* \ll T^{-1/2} \sum_{y=1, y \in V_k}^{\infty} \eta(y)^{-kc-1} \ll (\log H)^{k-1-50k^3} \ll 1.$$

To estimate Δ_2^* we apply [Lemma 1\(iii\)](#) and (10), (13), (16) to get

$$\begin{aligned} \Delta_2^* &\ll \sum_{\substack{He^{-\varkappa} < y < He^{\varkappa}, \\ y \in V_k}} \eta(y)^{-1} \ll \sum_{He^{-\varkappa} < u_1 u_2^3 \cdots u_{k-1}^{k-1} < He^{\varkappa}} (u_1 u_2 \cdots u_{k-1})^{-1} \\ &\ll \sum_{u_2^2 u_3^3 \cdots u_{k-1}^{k-1} < 2H} (u_2 u_3 \cdots u_{k-1})^{-1} \sum_{\substack{\frac{He^{-\varkappa}}{u_2^2 u_3^3 \cdots u_{k-1}^{k-1}} < u_1 < \frac{He^{\varkappa}}{u_2^2 u_3^3 \cdots u_{k-1}^{k-1}}}} u_1^{-1} \\ &\ll \sum_{u_2^2 u_3^3 \cdots u_{k-1}^{k-1} < 2H} \frac{H^{-1} u_2^2 u_3^3 \cdots u_{k-1}^{k-1} + \varkappa}{u_2 u_3 \cdots u_{k-1}} \\ (28) \quad &\ll H^{-1} \Delta_3 + 1, \end{aligned}$$

where Δ_3 is given by (18). Applying (19), (26)–(28) we find

$$(29) \quad \Delta^* \ll (\log H)^{k-3}.$$

We substitute in formula (12) the expression for $\Phi(s)$ given by (22) and find a new form of $U^{(1)}$. Using (10) and (29) we see that the contribution to $U^{(1)}$ coming from Δ^* is

$$\ll (\log H)^{k-3} \int_{-T}^T \frac{dt}{\sqrt{c^2+t^2}} \ll (\log H)^{k-2}.$$

Therefore, taking also into account (11) and (20), we find

$$(30) \quad U_k(H) = \frac{1}{(2\pi i)^2} \int_{c-iT}^{c+iT} \frac{H^s}{s} \int_{c-iT_1}^{c+iT_1} \frac{H^w}{w} \mathcal{M}(s, w) dw ds + O((\log H)^{k-2}).$$

4.7. For a fixed s satisfying $\text{Re}(s) = c$ the infinite series $\mathcal{M}(s, w)$ defined by (23) is absolutely and uniformly convergent for $\text{Re}(w) \geq c$ and represents a holomorphic function in $\text{Re}(w) > c$. Applying Euler’s identity we find

$$\begin{aligned} \mathcal{M}(s, w) &= \prod_p (1 + p^{-ks-1}(p^{s-w} + p^{2(s-w)} + \dots + p^{(k-1)(s-w)})) \\ &= \prod_p \left(1 + \sum_{j=1}^{k-1} p^{-(k-j)s-jw-1}\right). \end{aligned}$$

Using Lemma 3(ii), we conclude that for $\text{Re}(s) = c$ and $\text{Re}(w) \geq c$, we have

$$(31) \quad \mathcal{M}(s, w) = \mathcal{H}(s, w) \prod_{j=1}^{k-1} \zeta((k-j)s + jw + 1),$$

where

$$\mathcal{H}(s, w) = \prod_p \left(\left(1 + \sum_{j=1}^{k-1} p^{-(k-j)s-jw-1}\right) \prod_{j=1}^{k-1} (1 - p^{-(k-j)s-jw-1}) \right).$$

It is clear that there exists $\delta = \delta(k) \in (0, 1/100)$ such that in the region

$$(32) \quad \text{Re}(s) > -\delta \quad \text{and} \quad \text{Re}(w) > -\delta$$

the function $\mathcal{H}(s, w)$ is holomorphic with respect to s as well as to w and satisfies

$$(33) \quad 0 < |\mathcal{H}(s, w)| \ll 1.$$

We have also

$$(34) \quad \mathcal{H}(0, 0) = \mathcal{P}_k,$$

where \mathcal{P}_k is given by (4).

Suppose that we have a fixed $s = c + it$ with $-T \leq t \leq T$. From (31), (33) and Lemma 3(i), we conclude that the function $H^w w^{-1} \mathcal{M}(s, w)$ has a meromorphic continuation to $\operatorname{Re}(w) > -\delta$ and that poles may occur only at the points

$$(35) \quad w = 0 \quad \text{and} \quad w = (1 - k/m)s \quad \text{for } 1 \leq m \leq k - 1.$$

All these points are actually simple poles. Indeed, for $w = 0$ this follows immediately from (33) and parts (i) and (v) of Lemma 3. In the case $1 \leq m \leq k - 1$, the point $w = (1 - k/m)s$ is a simple pole of $\zeta((k - m)s + mw + 1)$ and, due to Lemma 3(v) and (10), it cannot be a pole or zero of $\zeta((k - j)s + jw + 1)$ for $1 \leq j \leq k - 1$ with $j \neq m$.

For $1 \leq m \leq k - 1$, we denote by $\mathcal{R}_m(s)$ the residue of $H^w w^{-1} \mathcal{M}(s, w)$ at $w = (1 - k/m)s$ and let $\mathcal{R}_0(s)$ be the residue at $w = 0$. A straightforward calculation, based on the arguments above, (33) and Lemma 3(i), leads to

$$(36) \quad \begin{aligned} \mathcal{R}_0(s) &= \mathcal{H}(s, 0) \prod_{j=1}^{k-1} \zeta(js + 1), \\ \mathcal{R}_m(s) &= \frac{H^{(1-k/m)s}}{(m-k)s} \mathcal{H}\left(s, \left(1 - \frac{k}{m}\right)s\right) \prod_{\substack{j=1, \\ j \neq m}}^{k-1} \zeta\left(k\left(1 - \frac{j}{m}\right)s + 1\right) \end{aligned} \quad \text{for } 1 \leq m \leq k - 1.$$

4.8. Let us define

$$(37) \quad \theta = \frac{\delta}{2k^3}.$$

By (10) and (21) and since $s = c + it$, where $-T \leq t \leq T$, we see that all points (35) are inside the rectangle with vertices $c - iT_1$, $-\theta - iT_1$, $-\theta + iT_1$, $c + iT_1$. Applying the residue theorem we find that

$$\int_{c-iT_1}^{c+iT_1} \frac{H^w}{w} \mathcal{M}(s, w) dw = 2\pi i \sum_{m=0}^{k-1} \mathcal{R}_m(s) + I_1 + I_2 + I_3,$$

where

$$\begin{aligned} I_1 &= \int_{c-iT_1}^{-\theta-iT_1} \frac{H^w}{w} \mathcal{M}(s, w) dw, & I_2 &= \int_{-\theta-iT_1}^{-\theta+iT_1} \frac{H^w}{w} \mathcal{M}(s, w) dw, \\ I_3 &= \int_{-\theta+iT_1}^{c+iT_1} \frac{H^w}{w} \mathcal{M}(s, w) dw. \end{aligned}$$

From the formula above and (30) we get

$$(38) \quad U_k(H) = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \frac{H^s}{s} \sum_{m=0}^{k-1} \mathcal{R}_m(s) ds + J_1 + J_2 + J_3 + O((\log H)^{k-2}).$$

Here J_μ are the contributions coming from I_μ for $\mu = 1, 2, 3$ and we will see that we may neglect them.

To estimate J_μ we will first show that if $s = c + it$, where $|t| \leq T$, and if w belongs to some of the sets of integration of I_1, I_2 or I_3 , then

$$(39) \quad \mathcal{M}(s, w) \ll T^{k^2\theta}.$$

Having in mind (31) and (33), we see that in order to verify this it is enough to establish that for s and w satisfying the conditions above, we have

$$(40) \quad \zeta(\lambda) \ll T^{k\theta}, \quad \text{where } \lambda = (k-j)s + jw + 1 \quad \text{for } 1 \leq j \leq k-1.$$

If $w = \beta + iT_1$ (or $w = \beta - iT_1$), where $-\theta \leq \beta \leq c$, then from (10), (21), (37) it follows that for the number λ given by (40), we have $\text{Re}(\lambda) \geq 1 - k\theta$ and $T \ll |\text{Im}(\lambda)| \ll T$. Hence the estimate (40) is a consequence of Lemma 3(iv). Suppose now that $w = -\theta + it_1$, where $|t_1| \leq T_1$. From (10), (21) and (37), we get $\text{Re}(\lambda) \geq 1 - k\theta$ and $|\text{Im}(\lambda)| \ll T$. If $|\text{Im}(\lambda)| \geq 2$, then the estimate (40) follows again from Lemma 3(iv). In the case $|\text{Im}(\lambda)| < 2$ we use also the inequality $\text{Re}(\lambda) \leq 1 - \theta/2$ to conclude that $\zeta(\lambda) \ll 1$, so the estimate (40) is true again.

From the definitions of J_μ and (10), (21), (37) and (39), we find

$$J_1, J_3 \ll \int_{-T}^T \frac{1}{\sqrt{c^2+t^2}} \int_{-\theta}^c \frac{T^{k^2\theta}}{\sqrt{\beta^2+T_1^2}} d\beta dt \ll c^{-1} + \log T \ll \log H,$$

$$J_2 \ll \int_{-T}^T \frac{1}{\sqrt{c^2+t^2}} \int_{-T_1}^{T_1} \frac{H^{-\theta} T^{k^2\theta}}{\sqrt{\theta^2+t_1^2}} dt_1 dt \ll H^{-\theta} (c^{-1} + \log T) T^{k^2\theta} \log T \ll 1.$$

This means that the terms J_μ in formula (38) can be omitted. Then using (36), we get

$$(41) \quad U_k(H) = \frac{1}{2\pi i} \left(\mathfrak{N}_0 + \sum_{m=1}^{k-1} \frac{1}{m-k} \mathfrak{N}_m \right) + O((\log H)^{k-2}),$$

where

$$(42) \quad \mathfrak{N}_m = \int_{c-iT}^{c+iT} \Xi_m(s) ds$$

and

$$(43) \quad \Xi_0(s) = s^{-1} H^s \mathfrak{H}(s, 0) \prod_{j=1}^{k-1} \zeta(js + 1),$$

$$(44) \quad \Xi_m(s) = s^{-2} H^{(2-k/m)s} \mathfrak{H}\left(s, \left(1 - \frac{k}{m}\right)s\right) \prod_{\substack{j=1, \\ j \neq m}}^{k-1} \zeta\left(k\left(1 - \frac{j}{m}\right)s + 1\right) \quad \text{for } 1 \leq m \leq k-1.$$

4.9. Consider first \mathfrak{N}_m for $1 \leq m \leq k/2$. Since $\Xi_m(s)$ is a holomorphic function in the rectangle with vertices $c - iT$, $\theta - iT$, $\theta + iT$ and $c + iT$, we have

$$(45) \quad \begin{aligned} \mathfrak{N}_m &= \int_{c-iT}^{\theta-iT} \Xi_m(s) ds + \int_{\theta-iT}^{\theta+iT} \Xi_m(s) ds + \int_{\theta+iT}^{c+iT} \Xi_m(s) ds \\ &= \mathfrak{N}_m^{(1)} + \mathfrak{N}_m^{(2)} + \mathfrak{N}_m^{(3)}, \end{aligned}$$

say. If s belongs to the sets of integration of $\mathfrak{N}_m^{(1)}$ or $\mathfrak{N}_m^{(3)}$ and if $1 \leq j \leq k-1$, $j \neq m$, then from [Lemma 3\(iv\)](#), it follows that

$$\zeta(k(1-j/m)s+1) \ll T^{k^2\theta}.$$

Hence, using [\(33\)](#), [\(37\)](#) and our assumption $1 \leq m \leq k/2$, we find

$$(46) \quad \mathfrak{N}_m^{(1)}, \mathfrak{N}_m^{(3)} \ll \int_c^\theta \frac{H^{(2-k/m)\beta}}{\beta^2 + T^2} T^{k^3\theta} d\beta \ll T^{k^3\theta-2} \ll 1.$$

Suppose now that s belongs to the set of integration of $\mathfrak{N}_m^{(2)}$ (that is, $s = \theta + it$ for $|t| \leq T$) and consider the number $\tilde{\lambda} = k(1-j/m)s + 1$. It is easy to see that for each j that occurs in [\(44\)](#), we have

$$\operatorname{Re}(\tilde{\lambda}) \geq 1 - k^2\theta, \quad |\operatorname{Re}(\tilde{\lambda}) - 1| \geq \theta, \quad |\operatorname{Im}(\tilde{\lambda})| \leq k^2|t|.$$

Hence an application of [Lemma 3\(iv\)](#) gives

$$\zeta(\tilde{\lambda}) \ll (1 + |t|)^{k^2\theta}.$$

Therefore

$$(47) \quad \mathfrak{N}_m^{(2)} \ll \int_{-T}^T \frac{H^{(2-k/m)\theta}}{\theta^2 + t^2} (1 + |t|)^{k^3\theta} dt \ll 1.$$

From [\(45\)](#)–[\(47\)](#), we get $\mathfrak{N}_m \ll 1$ for $1 \leq m \leq k/2$ and using [\(41\)](#) we find

$$(48) \quad U_k(H) = \frac{1}{2\pi i} \left(\mathfrak{N}_0 + \sum_{k/2 < m \leq k-1} \frac{1}{m-k} \mathfrak{N}_m \right) + O((\log H)^{k-2}).$$

4.10. Consider now \mathfrak{N}_m for $k/2 < m \leq k-1$. The function $\Xi_m(s)$ has a pole only at $s = 0$ and it is not difficult to compute that the corresponding residue is equal to

$$\mathcal{L}_m(\log H)^{k-1} + O((\log H)^{k-2}),$$

where

$$(49) \quad \mathcal{L}_m = \frac{(2m-k)^{k-1} (-1)^{k-m-1} \binom{k-1}{m} \mathcal{P}_k}{((k-1)!)^2 k^{k-2}}.$$

We leave the standard verification to the reader. From (42) and the residue theorem we get

$$(50) \quad \mathfrak{N}_m = 2\pi i \mathcal{L}_m(\log H)^{k-1} + \mathfrak{N}'_m + \mathfrak{N}''_m + \mathfrak{N}'''_m + O((\log H)^{k-2}),$$

where

$$\mathfrak{N}'_m = \int_{c-iT}^{-\theta-iT} \Xi_m(s) ds, \quad \mathfrak{N}''_m = \int_{-\theta-iT}^{-\theta+iT} \Xi_m(s) ds, \quad \mathfrak{N}'''_m = \int_{-\theta+iT}^{c+iT} \Xi_m(s) ds.$$

Using Lemma 3(iv), we find that if s belongs to the set of integration of some of the integrals above, then the product of the values of the zeta-function in the definition (44) is $\ll T^{k^3\theta}$. Hence from (10), (33), (37) and our assumption $k/2 < m \leq k-1$, it follows that

$$(51) \quad \begin{aligned} \mathfrak{N}'_m, \mathfrak{N}'''_m &\ll \int_{-\theta}^c \frac{T^{k^3\theta}}{\beta^2 + T^2} d\beta \ll 1 \\ \mathfrak{N}''_m &\ll \int_{-T}^T \frac{H^{-(2-k/m)\theta}}{\theta^2 + t^2} T^{k^3\theta} dt \ll H^{-(2-k/m)\theta} T^{k^3\theta} \ll 1. \end{aligned}$$

From (50) and (51), we find

$$(52) \quad \mathfrak{N}_m = 2\pi i \mathcal{L}_m(\log H)^{k-1} + O((\log H)^{k-2}) \quad \text{for } k/2 < m \leq k-1.$$

4.11. It remains to consider \mathfrak{N}_0 . It is not difficult to see that the function $\Xi_0(s)$ specified by (43) has a pole only at $s = 0$, with residue equal to

$$\mathcal{L}_0(\log H)^{k-1} + O((\log H)^{k-2}),$$

where

$$(53) \quad \mathcal{L}_0 = \frac{\mathcal{P}_k}{((k-1)!)^2}.$$

From (42) and the residue theorem we find

$$\mathfrak{N}_0 = 2\pi i \mathcal{L}_0(\log H)^{k-1} + \mathfrak{N}'_0 + \mathfrak{N}''_0 + \mathfrak{N}'''_0 + O((\log H)^{k-2}),$$

where

$$\mathfrak{N}'_0 = \int_{c-iT}^{-\theta-iT} \Xi_0(s) ds, \quad \mathfrak{N}''_0 = \int_{-\theta-iT}^{-\theta+iT} \Xi_0(s) ds, \quad \mathfrak{N}'''_0 = \int_{-\theta+iT}^{c+iT} \Xi_0(s) ds.$$

Arguing as above, we conclude that $\mathfrak{N}'_0, \mathfrak{N}''_0, \mathfrak{N}'''_0 \ll 1$ (we leave the verification to the reader). Hence

$$(54) \quad \mathfrak{N}_0 = 2\pi i \mathcal{L}_0(\log H)^{k-1} + O((\log H)^{k-2}).$$

From (3), (34), (48), (49), and (52)–(54), we obtain (9), and the proof of the theorem is complete. \square

References

- [Davenport 2000] H. Davenport, *Multiplicative number theory*, 3rd ed., Graduate Texts in Mathematics **74**, Springer, New York, 2000. [MR 2001f:11001](#) [Zbl 1002.11001](#)
- [Heath-Brown and Moroz 1999] D. R. Heath-Brown and B. Z. Moroz, “The density of rational points on the cubic surface $X_0^3 = X_1 X_2 X_3$ ”, *Math. Proc. Cambridge Philos. Soc.* **125**:3 (1999), 385–395. [MR 2000f:11080](#) [Zbl 0938.11016](#)
- [Titchmarsh 1986] E. C. Titchmarsh, *The theory of the Riemann zeta-function*, 2nd ed., Oxford University Press, 1986. [MR 88c:11049](#) [Zbl 0601.10026](#)

Received September 17, 2009.

DOYCHIN I. TOLEV
FACULTY OF MATHEMATICS AND INFORMATICS
SOFIA UNIVERSITY “ST. KL. OHRIDSKY”
5 J.BOURCHIER
1164 SOFIA
BULGARIA
dtolev@fmi.uni-sofia.bg
<http://www.tolev.org/>

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Mathew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 249 No. 2 February 2011

A gluing construction for prescribed mean curvature	257
ADRIAN BUTSCHER	
Large eigenvalues and concentration	271
BRUNO COLBOIS and ALESSANDRO SAVO	
Sur les conditions d'existence des faisceaux semi-stables sur les courbes multiples primitives	291
JEAN-MARC DRÉZET	
A quantitative estimate for quasiintegral points in orbits	321
LIANG-CHUNG HSIA and JOSEPH H. SILVERMAN	
Möbius isoparametric hypersurfaces with three distinct principal curvatures, II	343
ZEJUN HU and SHUIJIE ZHAI	
Discrete Morse theory and Hopf bundles	371
DMITRY N. KOZLOV	
Regularity of canonical and deficiency modules for monomial ideals	377
MANOJ KUMMINI and SATOSHI MURAI	
$SL_2(\mathbb{C})$ -character variety of a hyperbolic link and regulator	385
WEIPING LI and QINGXUE WANG	
Hypergeometric evaluation identities and supercongruences	405
LING LONG	
Necessary and sufficient conditions for unit graphs to be Hamiltonian	419
H. R. MAIMANI, M. R. POURNAKI and S. YASSEMI	
Instability of the geodesic flow for the energy functional	431
DOMENICO PERRONE	
String structures and canonical 3-forms	447
CORBETT REDDEN	
Dual pairs and contragredients of irreducible representations	485
BINYONG SUN	
On the number of pairs of positive integers $x_1, x_2 \leq H$ such that $x_1 x_2$ is a k -th power	495
DOYCHIN I. TOLEV	
Correction to the article A Floer homology for exact contact embeddings	509
KAI CIELIEBAK and URS ADRIAN FRAUENFELDER	