

Pacific Journal of Mathematics

**THE SECOND VARIATION
OF THE RICCI EXPANDER ENTROPY**

MENG ZHU

THE SECOND VARIATION OF THE RICCI EXPANDER ENTROPY

MENG ZHU

The critical points of the \mathcal{W}_+ functional introduced by M. Feldman, T. Ilmanen and L. Ni are the expanding Ricci solitons, which are special solutions of the Ricci flow. On compact manifolds, expanding solitons coincide with Einstein metrics. In this paper, we compute the first and second variations of the entropy functional of the \mathcal{W}_+ functional, and briefly discuss the linear stability of compact hyperbolic space forms.

1. Introduction

Perelman [2002] introduced two important functionals, denoted by \mathcal{F} and \mathcal{W} . The corresponding entropy functionals λ and ν are monotone along the Ricci flow $\partial g_{ij}/\partial t = -2R_{ij}$ and are constant precisely on steady and shrinking solitons. H.-D. Cao, R. Hamilton and T. Ilmanen [Cao et al. 2004] presented the second variations of both entropy functionals and studied the linear stabilities of certain closed Einstein manifolds of nonnegative scalar curvature.

To find the corresponding variational structure for the expanding case, M. Feldman, T. Ilmanen and L. Ni [Feldman et al. 2005] introduced the functional \mathcal{W}_+ . Let (M^n, g) be a compact Riemannian manifold, f a smooth function on M , and $\sigma > 0$. Define

$$\mathcal{W}_+(g, f, \sigma) = (4\pi\sigma)^{-n/2} \int_M e^{-f} (\sigma(|\nabla f|^2 + R) - f + n) dV,$$

$$\mu_+(g, \sigma) = \inf \left\{ \mathcal{W}_+(g, f, \sigma) \mid f \in C^\infty(M) \text{ with } (4\pi\sigma)^{-n/2} \int_M e^{-f} dV = 1 \right\},$$

$$\nu_+(g) = \sup_{\sigma > 0} \mu_+(g, \sigma).$$

Then ν_+ is nondecreasing along the Ricci flow and constant precisely on expanding solitons.

Research is partially supported by NSF grant DMS-0354621.

MSC2000: 53C21, 53C25, 53C44, 58J60.

Keywords: entropy functional, ν_+ functional, \mathcal{W}_+ functional, linear stability, linear variation, negative Einstein manifold, second variation.

In this note, analogous to [Cao et al. 2004], we present the first and second variations of the entropy ν_+ . By computing the first variation of ν_+ , one can see that the critical points are expanding solitons, which are actually negative Einstein manifolds (see [Cao and Zhu 2006], for example). Our main result is this:

Theorem 1.1. *Let (M^n, g) be a compact negative Einstein manifold. Let h be a symmetric 2-tensor. Consider the variation of metric $g(s) = g + sh$. Then the second variation of ν_+ is*

$$\frac{d^2\nu_+(g(s))}{ds^2}\Big|_{s=0} = \frac{\sigma}{\text{Vol } g} \int_M \langle N_+ h, h \rangle,$$

where

$$N_+ h := \frac{1}{2} \Delta h + \text{div}^* \text{div } h + \frac{1}{2} \nabla^2 v_h + \text{Rm}(h, \cdot) + \frac{g}{2n\sigma \text{Vol } g} \int_M \text{tr } h;$$

here tr is the trace with respect to g and v_h is the unique solution of

$$\Delta v_h - \frac{v_h}{2\sigma} = \text{div}(\text{div } h), \quad \int_M v_h = 0.$$

In this case, we may still define the concept of linear stability. We say that an expanding soliton is *linearly stable* if $N_+ \leq 0$; otherwise it is *linearly unstable*. Similar to [Cao et al. 2004], the N_+ operator is nonpositive definite if and only if the maximal eigenvalue of the Lichnerowicz Laplacian acting on the space of transverse traceless 2-tensors has a certain upper bound. Using the results in [Delay 2002; 2008] or [Lee 2006], one can then see that compact hyperbolic spaces are linearly stable. But unlike the positive Einstein case, it seems hard to find other examples of negative Einstein manifolds which are either linear stable or linear unstable.

2. The first variation of the expander entropy

Recall that in [Perelman 2002], the \mathcal{F} functional is defined by

$$\mathcal{F}(f, g) = \int_M (|\nabla f|^2 + R) e^{-f} dV,$$

and its entropy $\lambda(g)$ is

$$\lambda(g) = \inf \left\{ \mathcal{F}(f, g) \mid f \in C^\infty(M) \text{ with } \int_M e^{-f} = 1 \right\},$$

where R is the scalar curvature. By [Feldman et al. 2005, Theorem 1.7], we know that $\mu_+(g, \sigma)$ is attained by some function f . Moreover, if $\lambda(g) < 0$, then $\nu_+(g)$ can be attained by some positive number σ .

Lemma 2.1. *If $v_+(g)$ is realized by some f and σ , it is necessary that the pair (f, σ) solves the equations*

$$(1) \quad \sigma(-2\Delta f + |\nabla f|^2 - R) + f - n + v_+ = 0$$

and

$$(2) \quad (4\pi\sigma)^{-n/2} \int_M f e^{-f} dV = \frac{n}{2} - v_+.$$

Proof. For fixed $\sigma > 0$, suppose that $\mu_+(g, \sigma)$ is attained by some function f . Using the Lagrange multiplier method, consider the following functional

$$\begin{aligned} L(g, f, \sigma, \lambda) \\ = (4\pi\sigma)^{-n/2} \int_M e^{-f} (\sigma(|\nabla f|^2 + R) - f + n) dV + \lambda \left((4\pi\sigma)^{-n/2} \int_M e^{-f} dV - 1 \right). \end{aligned}$$

Denote by δf the variation of f . Then the variation of L is

$$\begin{aligned} 0 = \delta L \\ = (4\pi\sigma)^{-n/2} \int_M e^{-f} (-\delta f) (\sigma(|\nabla f|^2 + R) - f + n) dV \\ + (4\pi\sigma)^{-n/2} \int_M e^{-f} (2\sigma \nabla f \nabla(\delta f) - \delta f) dV - (4\pi\sigma)^{-n/2} \int_M \lambda(\delta f) e^{-f} dV \\ = (4\pi\sigma)^{-n/2} \int_M e^{-f} (\delta f) (\sigma(-2\Delta f + |\nabla f|^2 - R)) dV \\ + (4\pi\sigma)^{-n/2} \int_M e^{-f} (\delta f) (f - n - 1 - \lambda) dV \end{aligned}$$

Therefore,

$$\sigma(-2\Delta f + |\nabla f|^2 - R) + f - n - 1 - \lambda = 0.$$

Integrating both sides with respect to the measure $(4\pi\sigma)^{-n/2} e^{-f} dV$, we get

$$-\lambda - 1 = (4\pi\sigma)^{-n/2} \int_M e^{-f} (\sigma(|\nabla f|^2 + R) - f + n) dV = \mu_+(g, \sigma).$$

When σ and f realize $v_+(g)$, this is just [Equation \(1\)](#).

Now we consider the variations $\delta\sigma$ and δf of both σ and f . We have

$$\begin{aligned} (3) \quad 0 = (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(-\frac{n}{2\sigma} \delta\sigma - \delta f \right) (\sigma(|\nabla f|^2 + R) - f + n) dV \\ + (4\pi\sigma)^{-n/2} \int_M e^{-f} (\delta\sigma (|\nabla f|^2 + R) + 2\sigma \nabla f \nabla(\delta f) - \delta f) dV \end{aligned}$$

and

$$(4) \quad (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(-\frac{n}{2\sigma} \delta\sigma - \delta f \right) dV = 0.$$

Using (1) and (4), we can write (3) as

$$\begin{aligned} 0 &= (4\pi\sigma)^{-n/2} \int_M e^{-f} (\delta\sigma(|\nabla f|^2 + R) - \delta f) dV \\ &= (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(\frac{1}{\sigma} \delta\sigma(v_+ + f - n) + \frac{n}{2\sigma} \delta\sigma \right) dV \\ &= (\delta\sigma) \frac{1}{\sigma} (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(v_+ + f - \frac{n}{2} \right) dV, \end{aligned}$$

which gives (2). \square

Before computing the variations of the v_+ functional, let's recall some variation formulas for curvatures. By direct computation, we have:

Lemma 2.2. *Suppose that h is a symmetric 2-tensor and $g(s) = g + sh$ is a variation of g . Then*

$$(5) \quad \frac{\partial R}{\partial s} \Big|_{s=0} = -h_{kl} R_{kl} + \nabla_p \nabla_k h_{pk} - \Delta \operatorname{tr} h$$

and

$$\begin{aligned} (6) \quad \frac{\partial^2 R}{\partial s^2} \Big|_{s=0} &= 2h_{kp} h_{pl} R_{kl} - 2h_{kl} \frac{\partial R_{kl}}{\partial s} \Big|_{s=0} + g^{kl} \frac{\partial^2 R_{kl}}{\partial s^2} \Big|_{s=0} \\ &= 2h_{kp} h_{pl} R_{kl} - h_{kl} (2\nabla_p \nabla_k h_{pl} - \Delta h_{kl} - \nabla_k \nabla_l \operatorname{tr} h) \\ &\quad - \nabla_p (h_{pq} (2\nabla_k h_{kq} - \nabla_q \operatorname{tr} h)) + \nabla_k (h_{pq} \nabla_k h_{pq}) \\ &\quad + \frac{1}{2} \nabla_p \operatorname{tr} h (2\nabla_k h_{kp} - \nabla_p \operatorname{tr} h) + \frac{1}{2} (\nabla_k h_{pq} \nabla_k h_{pq} - 2\nabla_p h_{kq} \nabla_q h_{kp}), \end{aligned}$$

where ∇ is the Levi-Civita connection of g and $\operatorname{tr} h$ is the trace of h taken with respect to g .

Now we are ready to compute the first variation of $v_+(g)$.

Proposition 2.3. *Let (M^n, g) be a compact Riemannian manifold with $\lambda(g) < 0$. Let h be any symmetric covariant 2-tensor on M , and consider the variation*

$$g(s) = g + sh.$$

Then the first variation of $v_+(g(s))$ is

$$\frac{d v_+(g(s))}{ds} \Big|_{s=0} = (4\pi\sigma)^{-n/2} \int_M \sigma e^{-f} \left(-R_{ij} - \nabla_i \nabla_j f - \frac{1}{2\sigma} g_{ij} \right) h_{ij} dV,$$

where the smooth function f and $\sigma > 0$ realize $v_+(g)$.

Proof. By taking derivatives directly, we have

$$(7) \quad \begin{aligned} \frac{\partial \nu_+}{\partial s} &= (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(-\frac{n}{2\sigma} \frac{\partial \sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} g^{ij} h_{ij} \right) (\sigma(|\nabla f|^2 + R)) dV \\ &\quad + (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(-\frac{n}{2\sigma} \frac{\partial \sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} g^{ij} h_{ij} \right) (-f + n) dV \\ &\quad + (4\pi\sigma)^{-n/2} \int_M e^{-f} \frac{\partial \sigma}{\partial s} (|\nabla f|^2 + R) dV \\ &\quad - (4\pi\sigma)^{-n/2} \int_M e^{-f} (\sigma g^{ip} g^{jq} h_{pq} \nabla_i f \nabla_j f) dV \\ &\quad + (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(\sigma \left(2g^{ij} \nabla_i f \nabla_j \frac{\partial f}{\partial s} + \frac{\partial R}{\partial s} \right) - \frac{\partial f}{\partial s} \right) dV. \end{aligned}$$

Since $(4\pi\sigma)^{-n/2} \int_M e^{-f} dV = 1$, we have

$$(8) \quad (4\pi\sigma)^{-n/2} \int_M \left(-\frac{n}{2\sigma} \frac{\partial \sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} g^{ij} h_{ij} \right) e^{-f} dV = 0.$$

Substituting (1), (2) and (8) in (7), we obtain

$$\begin{aligned} \frac{\partial \nu_+(s)}{\partial s}|_{s=0} &= (4\pi\sigma)^{-n/2} \int_M (2\sigma(|\nabla f|^2 - \Delta f) + \nu_+(0)) \left(-\frac{n}{2\sigma} \frac{\partial \sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} g^{ij} h_{ij} \right) e^{-f} dV \\ &\quad + (4\pi\sigma)^{-n/2} \int_M \left(\frac{\partial \sigma}{\partial s} (|\nabla f|^2 + R) - \frac{\partial f}{\partial s} - \sigma h_{ij} \nabla_i f \nabla_j f \right) e^{-f} dV \\ &\quad + (4\pi\sigma)^{-n/2} \int_M \sigma \left(2 \frac{\partial f}{\partial s} (|\nabla f|^2 - \Delta f) + \nabla_i \nabla_j h_{ij} - \Delta \operatorname{tr} h - h_{ij} R_{ij} \right) e^{-f} dV \\ &= (4\pi\sigma)^{-n/2} \int_M \left(\frac{\partial \sigma}{\partial s} (|\nabla f|^2 + R) - \frac{\partial f}{\partial s} - \sigma (h_{ij} \nabla_i \nabla_j f + h_{ij} R_{ij}) \right) e^{-f} dV \\ &= (4\pi\sigma)^{-n/2} \int_M \left(\frac{\partial \sigma}{\partial s} (|\nabla f|^2 + R) + \frac{n}{2\sigma} \frac{\partial \sigma}{\partial s} \right) e^{-f} dV \\ &\quad - (4\pi\sigma)^{-n/2} \int_M \sigma h_{ij} \left(R_{ij} + \nabla_i \nabla_j f + \frac{1}{2\sigma} g_{ij} \right) e^{-f} dV \\ &= (4\pi\sigma)^{-n/2} \int_M \frac{1}{\sigma} \frac{\partial \sigma}{\partial s} \left(f(0) - \frac{n}{2} + \nu_+(0) - 2\sigma(|\nabla f|^2 - \Delta f) \right) e^{-f} dV \\ &\quad - (4\pi\sigma)^{-n/2} \int_M \sigma h_{ij} \left(R_{ij} + \nabla_i \nabla_j f + \frac{1}{2\sigma} g_{ij} \right) e^{-f} dV \\ &= -(4\pi\sigma)^{-n/2} \int_M \sigma h_{ij} \left(R_{ij} + \nabla_i \nabla_j f + \frac{1}{2\sigma} g_{ij} \right) e^{-f} dV. \end{aligned}$$

Hence, the first variation of ν_+ is

$$\frac{d\nu_+(g(s))}{ds}\Big|_{s=0} = (4\pi\sigma)^{-n/2} \int_M \sigma e^{-f} \left(-R_{ij} - \nabla_i \nabla_j f - \frac{1}{2\sigma} g_{ij} \right) h_{ij} dV. \quad \square$$

From the proposition, we can see that a critical point of $\nu_+(g)$ satisfies

$$\text{Rc} + \nabla^2 f + \frac{1}{2\sigma} g = 0,$$

which means that (M, g) is a gradient expanding soliton.

3. The second variation

Now we compute the second variation of ν_+ . Since any compact expanding soliton is Einstein (see [Cao and Zhu 2006], for example), f is a constant. After adding a constant to f we may assume that $f = n/2$.

In the following, as in [Cao et al. 2004], we set $\text{Rm}(h, h) = R_{ijkl}h_{ik}h_{jl}$, $\text{div } \omega = \nabla_i \omega_i$, $(\text{div } h)_i = \nabla_j h_{ji}$, and $(\text{div}^* \omega)_{ij} = -(\nabla_i \omega_j + \nabla_j \omega_i) = -\frac{1}{2}L_{\omega^\#}g_{ij}$, where h is a symmetric 2-tensor, ω is a 1-tensor, $\omega^\#$ is the dual vector field of ω , and $L_{\omega^\#}$ is the Lie derivative.

Proof of Theorem 1.1. Let (M, g) be a compact negative Einstein manifold with $f = n/2$ and $R_{ij} = -1/(2\sigma)g_{ij}$. For any symmetric 2-tensor h , consider the variation $g(s) = g + sh$. By Proposition 2.3, we know that $(d\nu_+/ds)|_{s=0} = 0$.

From (1) and (2), we get

$$(9) \quad \frac{n}{2\sigma} \frac{\partial \sigma}{\partial s}(0) - 2\sigma \Delta \frac{\partial f}{\partial s}(0) - \sigma \frac{\partial R}{\partial s}(0) + \frac{\partial f}{\partial s}(0) = 0,$$

and

$$(4\pi\sigma)^{-n/2} \int_M e^{-n/2} \left(\frac{n}{2} \left(-\frac{n}{2\sigma} \frac{\partial \sigma}{\partial s}(0) - \frac{\partial f}{\partial s}(0) + \frac{1}{2} \text{tr } h \right) + \frac{\partial f}{\partial s}(0) \right) dV = 0.$$

It follows by (8) that

$$(10) \quad (4\pi\sigma)^{-n/2} \int_M \frac{\partial f}{\partial s}(0) e^{-n/2} dV = 0$$

and

$$(11) \quad \frac{n}{2\sigma} \frac{\partial \sigma}{\partial s}(0) = \frac{1}{\text{Vol } g} \int_M \frac{1}{2} \text{tr } h dV,$$

where $(4\pi\sigma)^{-n/2}e^{-n/2} = \frac{1}{\text{Vol } g}$. Thus

$$\begin{aligned}
\frac{d\nu_+}{ds} &= (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(-\frac{n}{2\sigma} \frac{\partial\sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} g^{ij} h_{ij} \right) (\sigma(|\nabla f|^2 + R) - f + n) dV \\
&\quad + (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(\frac{\partial\sigma}{\partial s} (|\nabla f|^2 + R) - \frac{\partial f}{\partial s} \right) dV \\
&\quad + (4\pi\sigma)^{-n/2} \int_M \sigma e^{-f} \left(-g^{ip} g^{jq} h_{pq} \nabla_i f \nabla_j f + 2g^{ij} \nabla_i f \nabla_j \frac{\partial f}{\partial s} + \frac{\partial R}{\partial s} \right) dV \\
&= (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(-\frac{n}{2\sigma} \frac{\partial\sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} g^{ij} h_{ij} \right) (2\sigma(|\nabla f|^2 - \Delta f) + \nu_+) dV \\
&\quad + (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(\frac{\partial\sigma}{\partial s} (|\nabla f|^2 + R) - \frac{\partial f}{\partial s} \right) dV \\
&\quad + (4\pi\sigma)^{-n/2} \int_M \sigma e^{-f} \left(-g^{ip} g^{jq} h_{pq} \nabla_i f \nabla_j f + 2g^{ij} \nabla_i f \nabla_j \frac{\partial f}{\partial s} + \frac{\partial R}{\partial s} \right) dV \\
&= (4\pi\sigma)^{-n/2} \int_M \sigma e^{-f} g^{ij} h_{ij} (|\nabla f|^2 - \Delta f) dV \\
&\quad + (4\pi\sigma)^{-n/2} \int_M e^{-f} \left(\sigma \left(-g^{ip} g^{jq} h_{pq} \nabla_i f \nabla_j f + \frac{\partial R}{\partial s} \right) - \frac{1}{2} g^{ij} h_{ij} \right) dV,
\end{aligned}$$

where we note that

$$\int_M 2\sigma e^{-f} g^{ij} \nabla_i f \nabla_j \frac{\partial f}{\partial s} dV = \int_M 2\sigma e^{-f} \frac{\partial f}{\partial s} (|\nabla f|^2 - \Delta f) dV$$

and

$$\begin{aligned}
&\int_M e^{-f} \left(\frac{\partial\sigma}{\partial s} (|\nabla f|^2 + R) - \frac{\partial f}{\partial s} \right) dV \\
&= \int_M e^{-f} \left(\frac{\partial\sigma}{\partial s} (|\nabla f|^2 + R) + \frac{n}{2\sigma} \frac{\partial\sigma}{\partial s} - \frac{1}{2} g^{ij} h_{ij} \right) dV \\
&= \int_M e^{-f} \left(\frac{1}{\sigma} \frac{\partial\sigma}{\partial s} \left(\sigma (|\nabla f|^2 + R) + \frac{n}{2} \right) - \frac{1}{2} g^{ij} h_{ij} \right) dV \\
&= \int_M e^{-f} \frac{1}{\sigma} \frac{\partial\sigma}{\partial s} \left(\sigma (2|\nabla f|^2 - 2\Delta f) + f - \frac{n}{2} + \nu_+ \right) dV - \int_M e^{-f} \cdot \frac{1}{2} g^{ij} h_{ij} dV \\
&= - \int_M e^{-f} \cdot \frac{1}{2} g^{ij} h_{ij} dV.
\end{aligned}$$

Since $f(0) = \frac{n}{2}$, we have

$$(12) \quad \begin{aligned} \frac{d^2 v_+}{ds^2} \Big|_{s=0} &= -\frac{1}{\text{Vol } g} \int_M \sigma \operatorname{tr} h \Delta \frac{\partial f}{\partial s} dV \\ &\quad + \frac{1}{\text{Vol } g} \int_M \left(-\frac{n}{2\sigma} \frac{\partial \sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} \operatorname{tr} h \right) \left(\sigma \frac{\partial R}{\partial s} - \frac{1}{2} \operatorname{tr} h \right) dV \\ &\quad + \frac{1}{\text{Vol } g} \int_M \left(\frac{\partial \sigma}{\partial s} \frac{\partial R}{\partial s} + \sigma \frac{\partial^2 R}{\partial s^2} + \frac{1}{2} |h_{ij}|^2 \right) dV. \end{aligned}$$

In the following, all quantities are evaluated at $s = 0$. First, we have

$$(13) \quad \begin{aligned} &\frac{1}{\text{Vol } g} \int_M \sigma \frac{\partial^2 R}{\partial s^2} dV \\ &= \frac{\sigma}{\text{Vol } g} \int_M \left(-\frac{1}{\sigma} |h_{ij}|^2 - h_{kl} (2\nabla_p \nabla_k h_{pl} - \Delta h_{kl} - \nabla_k \nabla_l \operatorname{tr} h) \right. \\ &\quad \left. - \nabla_p (h_{pq} (2\nabla_k h_{kq} - \nabla_q \operatorname{tr} h)) + \nabla_k (h_{pq} \nabla_k h_{pq}) \right. \\ &\quad \left. + \frac{1}{2} \nabla_p \operatorname{tr} h (2\nabla_k h_{kp} - \nabla_p \operatorname{tr} h) + \frac{1}{2} (\nabla_k h_{pq} \nabla_k h_{pq} - 2\nabla_p h_{kq} \nabla_q h_{kp}) \right) dV \\ &= \frac{\sigma}{\text{Vol } g} \int_M \left(-\frac{1}{\sigma} |h_{ij}|^2 - h_{kl} \nabla_p \nabla_k h_{pl} - \frac{1}{2} |\nabla h|^2 - \frac{1}{2} |\nabla \operatorname{tr} h|^2 \right) dV \\ &= \frac{\sigma}{\text{Vol } g} \int_M \left(-\frac{1}{\sigma} |h_{ij}|^2 - \frac{1}{2} |\nabla h|^2 - \frac{1}{2} |\nabla \operatorname{tr} h|^2 \right) dV \\ &\quad - \frac{\sigma}{\text{Vol } g} \int_M h_{kl} (\nabla_k \nabla_p h_{pl} + R_{kq} h_{ql} + R_{pkql} h_{pq}) dV \\ &= -\frac{1}{\text{Vol } g} \int_M \frac{1}{2} |h_{ij}|^2 dV \\ &\quad + \frac{\sigma}{\text{Vol } g} \int_M (|\operatorname{div} h|^2 + \operatorname{Rm}(h, h) - \frac{1}{2} |\nabla h|^2 - \frac{1}{2} |\nabla \operatorname{tr} h|^2) dV. \end{aligned}$$

Moreover,

$$(14) \quad \begin{aligned} \frac{1}{\text{Vol } g} \int_M \frac{\partial \sigma}{\partial s} \frac{\partial R}{\partial s} dV &= \frac{\sigma}{n} \frac{1}{\text{Vol } g} \int_M \operatorname{tr} h dV \frac{1}{\text{Vol } g} \int_M \frac{\partial R}{\partial s} dV \\ &= \frac{1}{2n} \left(\frac{1}{\text{Vol } g} \int_M \operatorname{tr} h dV \right)^2. \end{aligned}$$

Let v_h be the solution to the equation

$$\Delta v_h - \frac{v_h}{2\sigma} = \operatorname{div} \operatorname{div} h = \nabla_p \nabla_q h_{pq}, \quad \int_M v_h = 0.$$

Then

$$\begin{aligned}
& \frac{1}{\text{Vol } g} \int_M \left(-\frac{n}{2\sigma} \frac{\partial \sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} \text{tr } h \right) \sigma \frac{\partial R}{\partial s} dV \\
&= \frac{\sigma}{\text{Vol } g} \int_M \left(-\frac{n}{2\sigma} \frac{\partial \sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} \text{tr } h \right) \left(\Delta v_h - \frac{v_h}{2\sigma} + \frac{1}{2\sigma} \text{tr } h - \Delta \text{tr } h \right) dV \\
&= - \left(\frac{1}{\text{Vol } g} \int_M \frac{1}{2} \text{tr } h dV \right)^2 + \frac{\sigma}{\text{Vol } g} \int_M v_h \left(-\Delta \frac{\partial f}{\partial s} + \frac{1}{2\sigma} \frac{\partial f}{\partial s} \right) dV \\
&\quad + \frac{\sigma}{\text{Vol } g} \int_M \text{tr } h \left(\Delta \frac{\partial f}{\partial s} - \frac{1}{2\sigma} \frac{\partial f}{\partial s} \right) dV \\
&\quad + \frac{\sigma}{\text{Vol } g} \int_M \frac{1}{2} \text{tr } h \left(\Delta v_h - \frac{v_h}{2\sigma} + \frac{1}{2\sigma} \text{tr } h - \Delta \text{tr } h \right) dV,
\end{aligned}$$

where we have used (11) to derive the first term in the last equality. Meanwhile,

$$\begin{aligned}
& -\frac{1}{\text{Vol } g} \int_M \frac{1}{2} \text{tr } h \left(-\frac{n}{2\sigma} \frac{\partial \sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} \text{tr } h \right) \\
&= -\frac{1}{\text{Vol } g} \int_M \frac{1}{2} \text{tr } h \left(-2\sigma \Delta \frac{\partial f}{\partial s} - \sigma \frac{\partial R}{\partial s} + \frac{1}{2} \text{tr } h \right).
\end{aligned}$$

It follows that

$$\begin{aligned}
& \frac{1}{\text{Vol } g} \int_M \left(-\frac{n}{2\sigma} \frac{\partial \sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} \text{tr } h \right) \left(\sigma \frac{\partial R}{\partial s} - \frac{1}{2} \text{tr } h \right) dV \\
&= \frac{1}{\text{Vol } g} \int_M \sigma \text{tr } h \Delta \frac{\partial f}{\partial s} dV - \frac{1}{\text{Vol } g} \int_M \frac{1}{4} (\text{tr } h)^2 dV \\
&\quad - \left(\frac{1}{\text{Vol } g} \int_M \frac{1}{2} \text{tr } h dV \right)^2 + \frac{\sigma}{\text{Vol } g} \int_M v_h \left(-\Delta \frac{\partial f}{\partial s} + \frac{1}{2\sigma} \frac{\partial f}{\partial s} \right) dV \\
&\quad + \frac{\sigma}{\text{Vol } g} \int_M \text{tr } h \left(\Delta \frac{\partial f}{\partial s} - \frac{1}{2\sigma} \frac{\partial f}{\partial s} \right) dV \\
&\quad + \frac{\sigma}{\text{Vol } g} \int_M \text{tr } h \left(\Delta v_h - \frac{v_h}{2\sigma} + \frac{1}{2\sigma} \text{tr } h - \Delta \text{tr } h \right) dV.
\end{aligned}$$

Now since

$$\begin{aligned}
& \frac{\sigma}{\text{Vol } g} \int_M v_h \left(-\Delta \frac{\partial f}{\partial s} + \frac{1}{2\sigma} \frac{\partial f}{\partial s} \right) dV = \frac{\sigma}{\text{Vol } g} \int_M v_h \left(-\frac{n}{4\sigma^2} \frac{\partial \sigma}{\partial s} + \frac{1}{2} \frac{\partial R}{\partial s} \right) dV \\
&= \frac{\sigma}{\text{Vol } g} \int_M \frac{1}{2} v_h \left(\Delta v_h - \frac{v_h}{2\sigma} + \frac{1}{2\sigma} \text{tr } h - \Delta \text{tr } h \right) dV \\
&= \frac{\sigma}{\text{Vol } g} \int_M -\frac{1}{2} |\nabla v_h|^2 - \frac{v_h^2}{4\sigma} + \frac{v_h}{4\sigma} \text{tr } h - \frac{1}{2} v_h \Delta \text{tr } h dV
\end{aligned}$$

and

$$\begin{aligned} & \frac{\sigma}{\text{Vol } g} \int_M \text{tr } h \left(\Delta \frac{\partial f}{\partial s} - \frac{1}{2\sigma} \frac{\partial f}{\partial s} \right) dV \\ &= \frac{\sigma}{\text{Vol } g} \int_M \text{tr } h \left(\frac{n}{4\sigma^2} \frac{\partial \sigma}{\partial s} - \frac{1}{2} \frac{\partial R}{\partial s} \right) dV \\ &= \left(\frac{1}{\text{Vol } g} \int_M \frac{1}{2} \text{tr } h dV \right)^2 - \frac{\sigma}{\text{Vol } g} \int_M \frac{1}{2} \text{tr } h \left(\Delta v_h - \frac{v_h}{2\sigma} + \frac{1}{2\sigma} \text{tr } h - \Delta \text{tr } h \right) dV, \end{aligned}$$

we have

$$\begin{aligned} (15) \quad & \frac{1}{\text{Vol } g} \int_M \left(-\frac{n}{2\sigma} \frac{\partial \sigma}{\partial s} - \frac{\partial f}{\partial s} + \frac{1}{2} \text{tr } h \right) \left(\sigma \frac{\partial R}{\partial s} - \frac{1}{2} \text{tr } h \right) dV \\ &= \frac{1}{\text{Vol } g} \int_M \sigma \text{tr } h \Delta \frac{\partial f}{\partial s} dV + \frac{\sigma}{\text{Vol } g} \int_M \left(-\frac{1}{2} |\nabla v_h|^2 - \frac{v_h^2}{4\sigma} + \frac{1}{2} |\nabla \text{tr } h|^2 \right) dV. \end{aligned}$$

Substituting (13), (14) and (15) in (12), we get

$$\begin{aligned} \frac{d^2 \nu_+}{ds^2} \Big|_{s=0} &= \frac{\sigma}{\text{Vol } g} \left(\int_M \left(|\text{div } h|^2 + \text{Rm}(h, h) - \frac{1}{2} |\nabla h|^2 - \frac{1}{2} |\nabla v_h|^2 - \frac{v_h^2}{4\sigma} \right) dV \right) \\ &\quad + \frac{1}{2n} \left(\frac{1}{\text{Vol } g} \int_M \text{tr } h dV \right)^2 \\ &= \frac{\sigma}{\text{Vol } g} \int_M \langle N_+ h, h \rangle. \end{aligned} \quad \square$$

As a simple application, we discuss briefly the linear stability of negative Einstein manifolds. In analogy with [Cao et al. 2004], we say that a negative Einstein manifold is linearly stable if $N_+ \leq 0$, otherwise it is linearly unstable. As in that paper, decompose the space of symmetric 2-tensors as

$$\ker \text{div} \oplus \text{im } \text{div}^*,$$

and further decompose $\ker \text{div}$ as

$$(\ker \text{div})_0 \oplus \mathbb{R}g,$$

where $(\ker \text{div})_0$ is the space of divergence free 2-tensors h with $\int_M \text{tr } h = 0$. It is easy to see that N_+ vanishes on $\text{im } \text{div}^*$, and on $(\ker \text{div})_0$

$$N_+ = \frac{1}{2} \left(\Delta_L - \frac{1}{\sigma} \right),$$

where $\Delta_L = \Delta + 2 \text{Rm}(\cdot, \cdot) - 2 \text{Rc}$ is the Lichnerowicz Laplacian on symmetric 2-tensors.

Moreover, we may write $(\ker \operatorname{div})_0$ as

$$(\ker \operatorname{div})_0 = S_0 \oplus S_1,$$

where S_0 is the subspace of trace free 2-tensors and

$$S_1 = \left\{ h \in (\ker \operatorname{div})_0 \mid h_{ij} = \left(-\frac{1}{2\sigma} u + \Delta u \right) g_{ij} - \nabla_i \nabla_j u, u \in C^\infty(M) \text{ and } \int_M u = 0 \right\};$$

see [Buzzanca 1984], for example.

Define

$$Tu := \left(-\frac{1}{2\sigma} u + \Delta u \right) g_{ij} - \nabla_i \nabla_j u.$$

Since $\Delta_L(Tu) = T(\Delta u)$ for all smooth functions u and $\ker T = \{0\}$, we can see that the Lichnerowicz Laplacian and the Laplacian on function space have the same eigenvalues. Thus N_+ is always negative on S_1 . Therefore, to study the linear stability of negative Einstein manifolds, it remains to look at the behavior of Δ_L acting on S_0 which is the space of transverse traceless 2-tensors.

Example. Suppose that M is an n dimensional compact real hyperbolic space with $n \geq 3$. By [Delay 2002] or [Lee 2006], the biggest eigenvalue of Δ_L on trace free symmetric 2-tensors on real hyperbolic space is $-\frac{1}{4}(n-1)(n-9)$. Since on M we have $\operatorname{Rc} = -(n-1)g$, we obtain

$$\frac{1}{\sigma} = 2(n-1).$$

Thus the biggest eigenvalue of N_+ on S_0 is not greater than $-\frac{1}{8}(n-1)^2$. This implies that M is linearly stable for $n \geq 3$.

Remarks. (1) When $n = 3$, D. Knopf and A. Young [2009] proved that closed 3-folds with constant negative curvature are geometrically stable under certain normalized Ricci flow. R. Ye [1993] had obtained a more powerful stability result earlier.

- (2) For $n = 2$, R. Hamilton [1988] proved that when the average scalar curvature is negative, the solution of the normalized Ricci flow with any initial metric converges to a metric with constant negative curvature. In particular, they are linearly stable. On the other hand, in [Delay 2008] we see that the biggest eigenvalue of the Lichnerowicz Laplacian on trace free symmetric 2-tensors is 2. Thus N_+ is nonpositive definite on $(\ker \operatorname{div})_0$, which also implies the linear stability.
- (3) For the noncompact case, V. Suneeta [2009] proved certain geometric stability of \mathbb{H}^n using different methods.

Acknowledgement

The author thanks his advisor, Professor Huai-Dong Cao, for encouragement and suggestions.

References

- [Buzzanca 1984] C. Buzzanca, “The Lichnerowicz Laplacian on tensors”, *Boll. Un. Mat. Ital. B* (6) **3**:2 (1984), 531–541. [MR 86c:58132](#)
- [Cao and Zhu 2006] H.-D. Cao and X.-P. Zhu, “A complete proof of the Poincaré and geometrization conjectures: application of the Hamilton–Perelman theory of the Ricci flow”, *Asian J. Math.* **10**:2 (2006), 165–492. [MR 2008d:53090](#) [Zbl 1200.53057](#)
- [Cao et al. 2004] H.-D. Cao, R. Hamilton, and T. Ilmanen, “Gaussian densities and stability for some Ricci solitons”, preprint, 2004. [arXiv math/0404165](#)
- [Delay 2002] E. Delay, “Essential spectrum of the Lichnerowicz Laplacian on two tensors on asymptotically hyperbolic manifolds”, *J. Geom. Phys.* **43**:1 (2002), 33–44. [MR 2003i:58055](#) [Zbl 1146.58301](#)
- [Delay 2008] E. Delay, “Spectrum of the Lichnerowicz Laplacian on asymptotically hyperbolic surfaces”, *Math. Phys. Anal. Geom.* **11**:3-4 (2008), 365–379. [MR 2010e:58031](#) [Zbl 1178.35276](#)
- [Feldman et al. 2005] M. Feldman, T. Ilmanen, and L. Ni, “Entropy and reduced distance for Ricci expanders”, *J. Geom. Anal.* **15**:1 (2005), 49–62. [MR 2006b:53091](#) [Zbl 1071.53040](#)
- [Hamilton 1988] R. S. Hamilton, “The Ricci flow on surfaces”, pp. 237–262 in *Mathematics and general relativity* (Santa Cruz, CA, 1986), edited by J. A. Isenberg, Contemp. Math. **71**, Amer. Math. Soc., Providence, RI, 1988. [MR 89i:53029](#) [Zbl 0663.53031](#)
- [Knopf and Young 2009] D. Knopf and A. Young, “Asymptotic stability of the cross curvature flow at a hyperbolic metric”, *Proc. Amer. Math. Soc.* **137**:2 (2009), 699–709. [MR 2009j:53085](#) [Zbl 1168.53034](#)
- [Lee 2006] J. M. Lee, *Fredholm operators and Einstein metrics on conformally compact manifolds*, Mem. Amer. Math. Soc. **183**:864, Amer. Math. Soc., Providence, RI, 2006. [MR 2007m:53047](#) [Zbl 1112.53002](#)
- [Perelman 2002] G. Perelman, “The entropy formula for the Ricci flow and its geometric applications”, preprint, 2002. [Zbl 1130.53001](#) [arXiv math.DG/0211159](#)
- [Suneeta 2009] V. Suneeta, “Investigating the off-shell stability of anti-de Sitter space in string theory”, *Classical Quantum Gravity* **26**:3 (2009), 035023. [MR 2010c:83119](#) [Zbl 1159.83023](#)
- [Ye 1993] R. Ye, “Ricci flow, Einstein metrics and space forms”, *Trans. Amer. Math. Soc.* **338**:2 (1993), 871–896. [MR 93j:58029](#) [Zbl 0804.53054](#)

Received January 30, 2010.

MENG ZHU
DEPARTMENT OF MATHEMATICS
LEHIGH UNIVERSITY
BETHLEHEM, PA 18015
UNITED STATES
mez206@lehigh.edu

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)

Department of Mathematics

University of California

Los Angeles, CA 90095-1555

pacific@math.ucla.edu

Vyjayanthi Chari

Department of Mathematics

University of California

Riverside, CA 92521-0135

chari@math.ucr.edu

Darren Long

Department of Mathematics

University of California

Santa Barbara, CA 93106-3080

long@math.ucsb.edu

Sorin Popa

Department of Mathematics

University of California

Los Angeles, CA 90095-1555

popa@math.ucla.edu

Robert Finn

Department of Mathematics

Stanford University

Stanford, CA 94305-2125

finn@math.stanford.edu

Jiang-Hua Lu

Department of Mathematics

The University of Hong Kong

Pokfulam Rd., Hong Kong

jhlu@maths.hku.hk

Jie Qing

Department of Mathematics

University of California

Santa Cruz, CA 95064

qing@cats.ucsc.edu

Kefeng Liu

Department of Mathematics

University of California

Los Angeles, CA 90095-1555

liu@math.ucla.edu

Alexander Merkurjev

Department of Mathematics

University of California

Los Angeles, CA 90095-1555

merkurev@math.ucla.edu

Jonathan Rogawski

Department of Mathematics

University of California

Los Angeles, CA 90095-1555

jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI

CALIFORNIA INST. OF TECHNOLOGY

INST. DE MATEMÁTICA PURA E APLICADA

KEIO UNIVERSITY

MATH. SCIENCES RESEARCH INSTITUTE

NEW MEXICO STATE UNIV.

OREGON STATE UNIV.

STANFORD UNIVERSITY

UNIV. OF BRITISH COLUMBIA

UNIV. OF CALIFORNIA, BERKELEY

UNIV. OF CALIFORNIA, DAVIS

UNIV. OF CALIFORNIA, LOS ANGELES

UNIV. OF CALIFORNIA, RIVERSIDE

UNIV. OF CALIFORNIA, SAN DIEGO

UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ

UNIV. OF MONTANA

UNIV. OF OREGON

UNIV. OF SOUTHERN CALIFORNIA

UNIV. OF UTAH

UNIV. OF WASHINGTON

WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic.

Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from [Periodicals Service Company](#), 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in LATEX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 251 No. 2 June 2011

Two Kazdan–Warner-type identities for the renormalized volume coefficients and the Gauss–Bonnet curvatures of a Riemannian metric BIN GUO, ZHENG-CHAO HAN and HAIZHONG LI	257
Gonality of a general ACM curve in \mathbb{P}^3 ROBIN HARTSHORNE and ENRICO SCHLESINGER	269
Universal inequalities for the eigenvalues of the biharmonic operator on submanifolds SAÏD ILIAS and OLA MAKHOUL	315
Multigraded Fujita approximation SHIN-YAO JOW	331
Some Dirichlet problems arising from conformal geometry QI-RUI LI and WEIMIN SHENG	337
Polycyclic quasiconformal mapping class subgroups KATSUHIKO MATSUZAKI	361
On zero-divisor graphs of Boolean rings ALI MOHAMMADIAN	375
Rational certificates of positivity on compact semialgebraic sets VICTORIA POWERS	385
Quiver grassmannians, quiver varieties and the preprojective algebra ALISTAIR SAVAGE and PETER TINGLEY	393
Nonautonomous second order Hamiltonian systems MARTIN SCHECHTER	431
Generic fundamental polygons for Fuchsian groups AKIRA USHIJIMA	453
Stability of the Kähler–Ricci flow in the space of Kähler metrics KAI ZHENG	469
The second variation of the Ricci expander entropy MENG ZHU	499