

*Pacific
Journal of
Mathematics*

**AVERAGE MAHLER'S MEASURE AND L_p NORMS
OF UNIMODULAR POLYNOMIALS**

KWOK-KWONG STEPHEN CHOI AND MICHAEL J. MOSSINGHOFF

AVERAGE MAHLER'S MEASURE AND L_p NORMS OF UNIMODULAR POLYNOMIALS

KWOK-KWONG STEPHEN CHOI AND MICHAEL J. MOSSINGHOFF

A polynomial $f \in \mathbb{C}[z]$ is unimodular if all its coefficients have unit modulus. Let \mathfrak{U}_n denote the set of unimodular polynomials of degree $n - 1$, and let \mathfrak{U}_n^* denote the subset of reciprocal unimodular polynomials, which have the property that $f(z) = \omega z^{n-1} \overline{f(1/\bar{z})}$ for some complex number ω with $|\omega| = 1$. We study the geometric and arithmetic mean values of both the normalized Mahler's measure $M(f)/\sqrt{n}$ and L_p norm $\|f\|_p/\sqrt{n}$ over the sets \mathfrak{U}_n and \mathfrak{U}_n^* , and compute asymptotic values in each case. We show for example that both the geometric and arithmetic mean of the normalized Mahler's measure approach $e^{-\gamma/2} = 0.749306 \dots$ as $n \rightarrow \infty$ for unimodular polynomials, and $e^{-\gamma/2}/\sqrt{2} = 0.529839 \dots$ for reciprocal unimodular polynomials. We also show that for large n , almost all polynomials in these sets have normalized Mahler's measure or L_p norm very close to the respective limiting mean value.

1. Introduction

Mahler's measure of a polynomial $f(z) \in \mathbb{C}[z]$, denoted $M(f)$, is defined by

$$\log M(f) = \int_0^1 \log |f(e(t))| dt,$$

where $e(t)$ denotes the function $e^{2\pi i t}$. If $f(z) = a_{n-1} \prod_{k=1}^{n-1} (z - \alpha_k)$, then it follows from Jensen's formula that

$$M(f) = |a_{n-1}| \prod_{k=1}^{n-1} \max\{1, |\alpha_k|\}.$$

The first author's research was supported in part by NSERC of Canada. The second author's research was supported in part by NSA grant number H98230-08-1-0052.

MSC2010: primary 30C10, 11R06; secondary 11C08, 60G99.

Keywords: mean Mahler's measure, mean L_p norm, unimodular polynomial, Littlewood polynomial.

Mahler's measure is also equal to the limit as p approaches zero of the integral defining the L_p norm of f over the unit circle,

$$M(f) = \lim_{p \rightarrow 0^+} \|f\|_p,$$

where $\|f\|_p$ is defined for $p > 0$ by

$$\|f\|_p = \left(\int_0^1 |f(e(t))|^p dt \right)^{1/p}.$$

Of course, this is only a norm for $p \geq 1$, but the restriction $p > 0$ is more natural in this article.

We say a polynomial

$$f(z) = \sum_{k=0}^{n-1} a_k z^k \in \mathbb{C}[z]$$

is *unimodular* if $|a_k| = 1$ for $0 \leq k < n$, and we call f a *Littlewood polynomial* if in fact $a_k = \pm 1$ for each k . Let \mathfrak{U}_n denote the set of unimodular polynomials of degree $n - 1$, and let \mathfrak{L}_n denote the set of Littlewood polynomials of degree $n - 1$. Note that by Parseval's formula, if $f \in \mathfrak{U}_n$ then $\|f\|_2 = \sqrt{n}$.

Littlewood posed a number of problems regarding the behavior of unimodular and Littlewood polynomials on the unit disk; see, for instance, [Littlewood 1968, problem 19]. He asked, for example, if there exist positive constants c_1 and c_2 and a sequence of polynomials $\{f_n\}$ with $f_n \in \mathfrak{U}_n$ (or, more strictly, $f_n \in \mathfrak{L}_n$) such that $c_1 \sqrt{n} \leq |f_n(z)| \leq c_2 \sqrt{n}$ for every z with modulus 1. Kahane [1980] proved this for unimodular polynomials, showing in fact that one may choose c_1 and c_2 arbitrarily close to 1 for sufficiently large n . Further, Beck [1991] showed that constants c_1 and c_2 exist for the set of polynomials whose coefficients are 1200th roots of unity. The problem for Littlewood polynomials however remains open — the Rudin–Shapiro polynomials satisfy the upper bound, and no sequence is known that satisfies the lower bound. Erdős [1962] conjectured that the constant c_2 must remain bounded away from 1 for large n for the case of the Littlewood polynomials.

Littlewood's problem has been investigated by using L_p norms and Mahler's measure. Since $\|f\|_2 < \|f\|_p < \|f\|_\infty$ when $2 < p < \infty$ and f is not a monomial, Erdős' conjecture regarding c_2 would follow if one could establish for some such p that $\|f\|_p / \|f\|_2$ is bounded away from 1, for every Littlewood polynomial f of positive degree. The special case $p = 4$ is equivalent to the well-known *merit factor problem* of Golay [Borwein et al. 2008b]. In the same way, one may ask for unimodular or Littlewood polynomials f whose normalized Mahler's measure $M(f) / \|f\|_2$ or L_p norm $\|f\|_p / \|f\|_2$ with $p < 2$ is especially large. For the case of the measure, Mahler [1963] proved that the maximum value of the measure of a polynomial with fixed degree having all its coefficients in the closed unit disk is

attained by a unimodular polynomial. Fielding [1970] reported that

$$(1-1) \quad \lim_{n \rightarrow \infty} \sup_{f \in \mathfrak{L}_n} \frac{M(f)}{\sqrt{n}} = 1,$$

attributing the proof to an anonymous referee. Beller and Newman [1973] also obtained a stronger version of this result.

For Littlewood polynomials, the problem of determining whether there exists a positive number ϵ such that $M(f)/\|f\|_2 < 1 - \epsilon$ for every $f \in \mathfrak{L}$ with $\deg(f) > 0$ is commonly known as *Mahler's problem* [Borwein 2002], and it remains open. The largest known value of $M(f)/\|f\|_2$ for a Littlewood polynomial of positive degree is $0.98636\dots$, achieved by $f(x) = x^{12} + x^{11} + x^{10} + x^9 + x^8 - x^7 - x^6 + x^5 + x^4 - x^3 + x^2 - x + 1$. The best known asymptotic result in Mahler's problem is due to Erdélyi and Lubinsky [2007], who proved that the Fekete polynomials, defined by $f_q(x) = \sum_{k=1}^{q-1} \left(\frac{k}{q}\right) x^{k-1}$, where q is a prime number and $\left(\frac{\cdot}{q}\right)$ denotes the Legendre symbol, satisfy

$$(1-2) \quad \frac{M(f_q)}{\sqrt{q-1}} > \frac{1}{2} - \epsilon,$$

for arbitrarily small $\epsilon > 0$ when q is sufficiently large.

Statistical properties of L_p norms and Mahler's measure are also of interest for the unimodular and Littlewood polynomials. Newman and Byrnes [1990] determined the mean value of the L_4 norm of a Littlewood polynomial, proving that

$$\text{mean}_{f \in \mathfrak{L}_n} \|f\|_4^4 = 2n^2 - n.$$

Borwein and Choi [2007] found similar formulas for the average value of the L_6 and L_8 norms, showing that

$$\text{mean}_{f \in \mathfrak{L}_n} \|f\|_6^6 = 6n^3 - 9n^2 + 4n$$

and

$$\text{mean}_{f \in \mathfrak{L}_n} \|f\|_8^8 = 24n^4 - 66n^3 + 58n^2 - 9n + 3((-1)^n - 1).$$

Borwein and Lockhart [2001] investigated the asymptotic behavior of the mean value of normalized L_p norms of Littlewood polynomials for arbitrary $p > 0$. Using the Lindeberg Central Limit Theorem and dominated convergence, they proved that

$$\lim_{n \rightarrow \infty} \text{mean}_{f \in \mathfrak{L}_n} \frac{\|f\|_p^p}{n^{p/2}} = \Gamma\left(1 + \frac{p}{2}\right).$$

Their result in fact holds for any class of polynomials whose coefficients are independent, identically distributed random variables with mean 0, variance 1, and

finite m -th moment for each $m > 2$, so this holds for the set of unimodular polynomials as well. By identifying a polynomial $f \in \mathfrak{U}_n$ by its vector of coefficients, we may consider \mathfrak{U}_n as an n -dimensional torus, and we let μ_n denote Haar measure on this space, normalized so that $\int_{\mathfrak{U}_n} d\mu_n = 1$. Using this notation, Borwein and Lockhart thus established that

$$(1-3) \quad \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \frac{\|f\|_p^p}{n^{p/2}} d\mu_n = \Gamma\left(1 + \frac{p}{2}\right),$$

for $p > 0$.

Fielding [1970] established a lower bound on the expected value of the normalized Mahler's measure for unimodular polynomials, proving that

$$(1-4) \quad \int_{\mathfrak{U}_n} \frac{M(f)}{\sqrt{n}} d\mu_n \geq e^{-\gamma/2} (1 + O(n^{-1/4+\epsilon}))$$

for arbitrary $\epsilon > 0$ and sufficiently large n , where γ denotes Euler's constant. Fielding's proof in fact established the precise limiting value for the geometric mean of the normalized Mahler's measure:

$$(1-5) \quad \int_{\mathfrak{U}_n} \log\left(\frac{M(f)}{\sqrt{n}}\right) d\mu_n = -\frac{\gamma}{2} + O(n^{-1/4+\epsilon}).$$

In this paper, we strengthen (1-4) by proving that the limiting value of the arithmetic mean of the normalized measure of unimodular polynomials of fixed degree is precisely $e^{-\gamma/2} = 0.749306\dots$. We also recover (1-3) and (1-5) by using a different method, and we establish the asymptotic value of the geometric mean of certain L_p norms of the unimodular polynomials as well. We prove the following theorem in Section 2.

Theorem 1. *Let \mathfrak{U}_n denote the set of unimodular polynomials of degree $n - 1$. Then*

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \log\left(\frac{M(f)}{\sqrt{n}}\right) d\mu_n = -\frac{\gamma}{2},$$

and for any even positive integer p ,

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \log\left(\frac{\|f\|_p}{\sqrt{n}}\right) d\mu_n = \frac{1}{p} \log \Gamma\left(1 + \frac{p}{2}\right).$$

Also,

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \frac{M(f)}{\sqrt{n}} d\mu_n = e^{-\gamma/2},$$

and for any positive real number p ,

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \frac{\|f\|_p^p}{n^{p/2}} d\mu_n = \Gamma\left(1 + \frac{p}{2}\right).$$

A polynomial $f(z) \in \mathbb{C}[z]$ of degree d is said to be *reciprocal* if $f(z) = \omega z^d \overline{f(1/\bar{z})}$, for some complex number ω of modulus 1. Reciprocal polynomials often arise in a natural way in problems involving Mahler’s measure. For example, Smyth [1971] proved that if $f(z) \in \mathbb{Z}[z]$ is nonreciprocal and $f(0) \neq 0$, then

$$M(f) \geq M(z^3 - z - 1) = 1.32471 \dots$$

Let \mathfrak{U}_n^* denote the set of reciprocal unimodular polynomials of degree $n - 1$, and let \mathfrak{L}_n^* denote the set of reciprocal Littlewood polynomials of degree $n - 1$. Borwein and Choi [2007] determined the expected value of the L_4 norm of a polynomial $f \in \mathfrak{L}_n^*$, showing that

$$\text{mean}_{f \in \mathfrak{L}_n^*} \|f\|_4^4 = 3n^2 - 3n + \frac{1 - (-1)^n}{2}.$$

In this article, we prove that both the geometric mean and the arithmetic mean of the normalized Mahler’s measure of the reciprocal unimodular polynomials approach

$$\frac{e^{-\gamma/2}}{\sqrt{2}} = 0.529839 \dots,$$

and we determine limiting values for the mean normalized L_p norms of these polynomials. Since \mathfrak{U}_n^* can be identified with a $(1 + \lfloor n/2 \rfloor)$ -dimensional real torus, we let μ_n^* denote Haar measure on the set \mathfrak{U}_n^* , normalized so that $\int_{\mathfrak{U}_n^*} d\mu_n^* = 1$. In Section 3 we prove the following theorem.

Theorem 2. *Let \mathfrak{U}_n^* denote the set of reciprocal unimodular polynomials of degree $n - 1$. Then*

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n^*} \log\left(\frac{M(f)}{\sqrt{n}}\right) d\mu_n^* = -\frac{\gamma + \log 2}{2},$$

and for any even positive integer p ,

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n^*} \log\left(\frac{\|f\|_p}{\sqrt{n}}\right) d\mu_n^* = \frac{1}{2} \log(2\pi^{1/p}) + \frac{1}{p} \log \Gamma\left(\frac{p+1}{2}\right).$$

Also,

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n^*} \frac{M(f)}{\sqrt{n}} d\mu_n^* = \frac{1}{\sqrt{2e^\gamma}},$$

and for any positive number p ,

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n^*} \frac{\|f\|_p^p}{n^{p/2}} d\mu_n^* = \frac{2^{p/2}}{\sqrt{\pi}} \Gamma\left(\frac{p+1}{2}\right).$$

We may obtain more information on the distribution of values of L_p norms and the Mahler measure by studying their moments, and to this end we establish the following result in Section 4.

Theorem 3. *Let $p \geq 2$ be an even integer and m a positive integer. Then*

$$(1-6) \quad \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \left(\frac{\|f\|_p^p}{n^{p/2}} \right)^m d\mu_n = \left(\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \frac{\|f\|_p^p}{n^{p/2}} d\mu_n \right)^m$$

and

$$(1-7) \quad \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \left(\frac{M(f)}{\sqrt{n}} \right)^m d\mu_n = \left(\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \frac{M(f)}{\sqrt{n}} d\mu_n \right)^m.$$

Also, (1-6) holds for all positive real numbers p when $m = 2$. The same statements hold for the set \mathfrak{U}_n^* .

Theorem 3 establishes that the limiting value of the variance of each distribution approaches 0 as the degree grows large. It follows that the normalized Mahler's measure or L_p norm (with $p \geq 2$ an even integer) of a polynomial in one of these sets lies arbitrarily close to the corresponding mean value, with probability approaching 1 as the degree grows large. We thus obtain the following corollary regarding convergence in probability of the measure and L_p norms. (The result on L_p norms for \mathfrak{U}_n was also obtained by Borwein and Lockhart [2001] for all $p > 0$.)

Corollary 4. *For any $\epsilon > 0$ and $\delta > 0$, there exists a positive integer $N = N(\epsilon, \delta)$ such that if $n \geq N$, then*

$$\Pr\left(f \in \mathfrak{U}_n : \left| \frac{M(f)}{\sqrt{n}} - e^{-\gamma/2} \right| > \delta\right) < \epsilon,$$

and, if $p \geq 2$ is an even integer, then

$$\Pr\left(f \in \mathfrak{U}_n : \left| \frac{\|f\|_p^p}{n^{p/2}} - \Gamma\left(1 + \frac{p}{2}\right) \right| > \delta\right) < \epsilon.$$

Analogous statements hold for the set \mathfrak{U}_n^* .

Section 5 adds some remarks on related problems.

2. Unimodular polynomials

We require the following notation. Let $Q_{n,k}$ denote the set of all sequences of length k selected without repetition from the set $\{0, 1, \dots, n-1\}$, so $|Q_{n,k}| = n!/(n-k)!$ for $0 \leq k \leq n$. Let $R_{n,k}$ denote the set of such sequences if repetition is allowed, so $|R_{n,k}| = n^k$. For two sequences I and J in $R_{n,k}$, we write $I \sim J$ if I is a permutation of J . Let P_k denote the set of permutations of $\{1, 2, \dots, k\}$, so $|P_k| = k!$. Finally, for a vector $\mathbf{t} \in \mathbb{R}^k$, let $N_{\mathbf{t}}$ denote the number of permutations of P_k that fix \mathbf{t} , so

$$N_{\mathbf{t}} = |\{\pi \in P_k : t_\ell = t_{\pi(\ell)} \text{ for } 1 \leq \ell \leq k\}|.$$

Lemma 1. *Let $\mathbf{t} = (t_1, \dots, t_k)$ denote a point in $[0, 1)^k$. If $Q_{n,k}$ and $N_{\mathbf{t}}$ are defined as above, then*

$$(2-1) \quad \sum_{\substack{I, J \in Q_{n,k} \\ I \sim J}} \prod_{\ell=1}^k e(t_{\ell}(i_{\ell} - j_{\ell})) = n^k N_{\mathbf{t}} + O(n^{k-1}),$$

where the implicit constant may depend on k and \mathbf{t} but is independent of n .

Proof. We compute

$$\begin{aligned} \sum_{\substack{I, J \in Q_{n,k} \\ I \sim J}} \prod_{\ell=1}^k e(t_{\ell}(i_{\ell} - j_{\ell})) &= \sum_{\pi \in P_k} \sum_{I \in Q_{n,k}} \prod_{\ell=1}^k e(t_{\ell}(i_{\ell} - i_{\pi(\ell)})) \\ &= \sum_{\pi \in P_k} \sum_{I \in R_{n,k}} \prod_{\ell=1}^k e(t_{\ell}(i_{\ell} - i_{\pi(\ell)})) + O(n^{k-1}) \\ &= \sum_{\pi \in P_k} \sum_{I \in R_{n,k}} \prod_{\ell=1}^k e((t_{\ell} - t_{\pi^{-1}(\ell)})i_{\ell}) + O(n^{k-1}) \\ &= \sum_{\pi \in P_k} \prod_{\ell=1}^k \sum_{0 \leq i_{\ell} < n} e((t_{\ell} - t_{\pi^{-1}(\ell)})i_{\ell}) + O(n^{k-1}). \end{aligned}$$

If α is an integer, then certainly

$$\sum_{0 \leq j < n} e(\alpha j) = n,$$

and if α is not an integer, then

$$\sum_{0 \leq j < n} e(\alpha j) \ll \frac{1}{\|\alpha\|},$$

where $\|\alpha\|$ denotes the distance from α to the nearest integer, $\|\alpha\| = \min_{n \in \mathbb{Z}} |\alpha - n|$.

Thus

$$\sum_{\substack{I, J \in Q_{n,k} \\ I \sim J}} \prod_{\ell=1}^k e(t_{\ell}(i_{\ell} - j_{\ell})) = n^k N_{\mathbf{t}} + O(n^{k-1}). \quad \square$$

In particular, suppose m_1, \dots, m_r are positive integers, and let $M_j = \sum_{i=1}^j m_i$ for each j with $0 \leq j \leq r$. Suppose also that t_{M_1}, \dots, t_{M_r} are distinct real numbers in $[0, 1)$, and that for each j with $1 \leq j \leq r$, we have

$$t_{M_{j-1}+i} = t_{M_j},$$

for $1 \leq i < m_j$. Let $M = M_r$. Then by [Lemma 1](#), we have

$$(2-2) \quad \sum_{\substack{I, J \in \mathcal{Q}_{n, M} \\ I \sim J}} \prod_{\ell=1}^M e(t_\ell(i_\ell - j_\ell)) = n^M \prod_{\ell=1}^r m_\ell! + O(n^{M-1}).$$

We use this to establish the following result regarding the mean value of a particular expression over the set of unimodular polynomials.

Lemma 2. *Suppose t_1, \dots, t_r are distinct real numbers in $[0, 1)$, and m_1, \dots, m_r are positive integers. Then*

$$\int_{\mathfrak{A}_n} \prod_{\ell=1}^r \frac{|f(e(t_\ell))|^{2m_\ell}}{n^{m_\ell}} d\mu_n = \prod_{\ell=1}^r m_\ell! + O\left(\frac{1}{n}\right),$$

where the implicit constant may depend on the m_ℓ and the t_ℓ but is independent of n . Thus,

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \prod_{\ell=1}^r \frac{|f(e(t_\ell))|^{2m_\ell}}{n^{m_\ell}} d\mu_n = \prod_{\ell=1}^r m_\ell!.$$

Proof. For clarity, first consider the case $r = 1$. Write m in place of m_1 , let $f(z) = \sum_{k=0}^n a_k z^k$, and let $I = (i_1, \dots, i_m)$ and $J = (j_1, \dots, j_m)$. Then

$$\begin{aligned} \int_{\mathfrak{A}_n} \frac{|f(e(t))|^{2m}}{n^m} d\mu_n &= \frac{1}{n^m} \int_{\mathfrak{A}_n} \left| \sum_{k=1}^n a_k e(kt) \right|^{2m} d\mu_n \\ &= \frac{1}{n^m} \sum_{\substack{I, J \in R_{n, m} \\ I \sim J}} e\left(t \sum_{k=1}^m (i_k - j_k)\right) \int_{\mathfrak{A}_n} \prod_{k=1}^m a_{i_k} \overline{a_{j_k}} d\mu_n \\ &= \frac{1}{n^m} \sum_{\substack{I, J \in R_{n, m} \\ I \sim J}} 1 \\ &= \frac{1}{n^m} \sum_{\substack{I, J \in \mathcal{Q}_{n, m} \\ I \sim J}} 1 + O\left(\frac{1}{n}\right) \\ &= m! + O\left(\frac{1}{n}\right). \end{aligned}$$

The general case is similar. Let $M = \sum_{\ell=1}^r m_\ell$. We find

$$\int_{\mathfrak{A}_n} \prod_{\ell=1}^r \frac{|f(e(t_\ell))|^{2m_\ell}}{n^{m_\ell}} d\mu_n = \frac{1}{n^M} \sum_{\substack{I_\ell, J_\ell \in R_{n, m_\ell} \\ (I_1, \dots, I_r) \sim (J_1, \dots, J_r)}} \prod_{\ell=1}^r e\left(t_\ell \sum_{k=1}^{m_\ell} (i_{\ell, k} - j_{\ell, k})\right),$$

where $I_\ell = (i_{\ell, 1}, \dots, i_{\ell, m_\ell})$ and $J_\ell = (j_{\ell, 1}, \dots, j_{\ell, m_\ell})$. Next, restricting from R_{n, m_ℓ} to \mathcal{Q}_{n, m_ℓ} changes the value here by $O(1/n)$, since the terms in the sum over I_ℓ and

J_ℓ having repeated elements possess fewer degrees of freedom and contribute only $O(n^{m_\ell-1})$. Thus

$$\int_{\mathfrak{A}_n} \prod_{\ell=1}^r \frac{|f(e(t_\ell))|^{2m_\ell}}{n^{m_\ell}} d\mu_n = \frac{1}{n^M} \sum_{\substack{I_\ell, J_\ell \in \mathcal{Q}_{n, m_\ell} \\ (I_1, \dots, I_r) \sim (J_1, \dots, J_r)}} \prod_{\ell=1}^r e\left(t_\ell \sum_{k=1}^{m_\ell} (i_{\ell, k} - j_{\ell, k})\right) + O\left(\frac{1}{n}\right),$$

where the implicit constant depends on the t_ℓ and the m_ℓ . Similarly, we may restrict further to the case where the sequences I_1, \dots, I_r are disjoint, so by (2-2),

$$\begin{aligned} \int_{\mathfrak{A}_n} \prod_{\ell=1}^r \frac{|f(e(t_\ell))|^{2m_\ell}}{n^{m_\ell}} d\mu_n &= \frac{1}{n^M} \sum_{\substack{I, J \in \mathcal{Q}_{n, M} \\ I \sim J}} \prod_{\ell=1}^r \prod_{k=1}^{m_\ell} e(t_\ell (i_k - j_k)) + O\left(\frac{1}{n}\right) \\ &= \prod_{\ell=1}^r m_\ell! + O\left(\frac{1}{n}\right). \end{aligned} \quad \square$$

The next lemma establishes another mean value over the unimodular polynomials.

Lemma 3. *Let t_1, \dots, t_r be distinct real numbers in $[0, 1)$, and let s_1, \dots, s_r be complex numbers satisfying $\Re(s_\ell) > 0$ for $1 \leq \ell \leq r$. Then*

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \exp\left(-\sum_{\ell=1}^r s_\ell \frac{|f(e(t_\ell))|^2}{n}\right) d\mu_n = \prod_{\ell=1}^r \frac{1}{1+s_\ell}.$$

Proof. Let $F(s_1, \dots, s_r)$ denote the function

$$F(s_1, \dots, s_r) = \lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \exp\left(-\sum_{\ell=1}^r s_\ell \frac{|f(e(t_\ell))|^2}{n}\right) d\mu_n,$$

so this defines an analytic function on the region

$$\{(s_1, \dots, s_r) \in \mathbb{C}^r : \Re(s_\ell) > 0 \text{ for } 1 \leq \ell \leq r\}.$$

For real numbers x_1, \dots, x_r in $[0, 1)$, use Lemma 2 to compute

$$\begin{aligned} F(x_1, \dots, x_r) &= \lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \prod_{\ell=1}^r \sum_{m_\ell \geq 0} \frac{(-x_\ell)^{m_\ell} |f(e(t_\ell))|^{2m_\ell}}{n^{m_\ell} m_\ell!} d\mu_n \\ &= \sum_{\substack{m_\ell \geq 0 \\ 1 \leq \ell \leq r}} \left(\prod_{\ell=1}^r \frac{(-x_\ell)^{m_\ell}}{m_\ell!} \right) \lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \prod_{\ell=1}^r \frac{|f(e(t_\ell))|^{2m_\ell}}{n^{m_\ell}} d\mu_n \\ &= \prod_{\ell=1}^r \sum_{m_\ell \geq 0} (-x_\ell)^{m_\ell} = \prod_{\ell=1}^r \frac{1}{1+x_\ell}. \end{aligned}$$

The statement then follows by analytic continuation. □

Next, recall that a sequence of random variables $\{X_n\}$ is *uniformly integrable* if

$$(2-3) \quad \lim_{c \rightarrow \infty} \sup_n \int_{|X_n| > c} |X_n| d\mu_n = 0.$$

For a fixed real number $t \in [0, 1)$, let $U_{t,n}$ denote the random variable whose value is $|f(e(t))|^2/n$, where each coefficient of f is selected uniformly over the unit circle. For convenience, let U_n denote the random variable $U_{0,n}$. We show that the sequence of random variables $X_n = \log U_{t,n}$ has the property (2-3).

Lemma 4. *Let t be a fixed real number with $0 \leq t < 1$. The sequence of random variables $\{\log U_{t,n}\}$, with $n \geq 1$, is uniformly integrable.*

Proof. By symmetry, we may assume that $t = 0$. Let c be a positive real number, let $F_n(x)$ denote the distribution function of U_n , and let $G_n(x) = 1 - F_n(x)$. Certainly U_n is supported on $[0, n]$, so $F_n(x) = 1$ for $x \geq n$. Thus

$$\begin{aligned} \int_{U_n > e^c} \log U_n d\mu_n &= \int_{e^c}^n \log x dF_n(x) \\ &= \log n - cF_n(c) - \int_{e^c}^n \frac{F_n(x)}{x} dx \\ &= cG_n(e^c) + \int_{e^c}^n \frac{G_n(x)}{x} dx. \end{aligned}$$

By Markov's inequality, we have $G_n(x) \leq E(U_n)/x = 1/x$, so

$$\int_{U_n > e^c} \log U_n d\mu_n \leq (c+1)e^{-c}.$$

For $U_n < e^{-c}$, we require some additional information. Using a result of Kluyver [1906] on the distribution of the modulus of sums of complex numbers of unit modulus, Fielding [1970] determined an asymptotic formula for the density function of the random variable Y_n that measures the value of $|f(1)|$, for a randomly selected polynomial $f \in \mathfrak{U}_n$. This random variable is well-approximated by a Weibull distribution with shape parameter 2 and scale parameter \sqrt{n} ; its density function is

$$h_n(y) = \begin{cases} \frac{2y}{n} e^{-y^2/n} + O\left(\frac{\sqrt{y}}{n^{7/4-\epsilon}}\right), & \text{if } 0 \leq y \leq n, \\ 0, & \text{if } y > n. \end{cases}$$

Since $U_n = Y_n^2/n$, it follows that U_n has approximately an exponential distribution, with density function

$$(2-4) \quad u_n(x) = \begin{cases} e^{-x} + O(x^{-1/4}n^{-1+\epsilon}), & \text{if } 0 \leq x \leq n, \\ 0, & \text{if } x > n. \end{cases}$$

Thus

$$\begin{aligned} \int_{U_n < e^{-c}} |\log U_n| d\mu_n &= \int_0^{e^{-c}} (e^{-x} + O(x^{-1/4}n^{-1+\epsilon})) |\log x| dx \\ &\leq \int_0^{e^{-c}} |\log x| dx + O\left(\frac{1}{n^{1-\epsilon}} \int_0^{e^{-c}} \frac{|\log x|}{x^{1/4}} dx\right) \\ &= (c+1)e^{-c} + O\left(\frac{ce^{-3c/4}}{n^{1-\epsilon}}\right), \end{aligned}$$

and so

$$\sup_n \int_{U_n < e^{-c}} |\log U_n| d\mu_n \ll ce^{-3c/4}. \quad \square$$

The final three lemmata record results about certain integrals.

Lemma 5. *If t_1, \dots, t_r are real numbers in $[0, 1)$, then*

$$\int_{\mathfrak{A}_n} \prod_{\ell=1}^r \left| \log \left(\frac{|f(e(t_\ell))|^2}{n} \right) \right| d\mu_n < \infty.$$

Proof. Let $\|\cdot\|_r$ denote the L_r norm with respect to the measure μ_n , so

$$\|f\|_r = \left(\int_{\mathfrak{A}_n} |f|^r d\mu_n \right)^{1/r}.$$

Using the arithmetic-geometric mean inequality, followed by the triangle inequality for $\|\cdot\|_r$, we obtain

$$\begin{aligned} \int_{\mathfrak{A}_n} \prod_{\ell=1}^r \left| \log \left(\frac{|f(e(t_\ell))|^2}{n} \right) \right| d\mu_n &\leq \int_{\mathfrak{A}_n} \left(\frac{1}{r} \sum_{\ell=1}^r \left| \log \left(\frac{|f(e(t_\ell))|^2}{n} \right) \right| \right)^r d\mu_n \\ &= \frac{1}{r^r} \left\| \sum_{\ell=1}^r \left| \log \left(\frac{|f(e(t_\ell))|^2}{n} \right) \right| \right\|_r^r \\ &\leq \frac{1}{r^r} \left(\sum_{\ell=1}^r \left\| \left| \log \left(\frac{|f(e(t_\ell))|^2}{n} \right) \right| \right\|_r \right)^r \\ &= E(|\log U_n|^r). \end{aligned}$$

From (2-4), we have

$$E(|\log U_n|^r) = \int_0^n |\log x|^r (e^{-x} + O(x^{-1/4}n^{-1+\epsilon})) dx.$$

We compute

$$\begin{aligned} \int_0^1 |\log x|^r e^{-x} dx &\leq \int_0^1 |\log x|^r dx = r!, \\ \int_1^\infty |\log x|^r e^{-x} dx &\leq \int_1^\infty (x-1)^r e^{-x} dx = \frac{r!}{e}, \end{aligned}$$

and

$$\frac{1}{n^{1-\epsilon}} \int_0^n \frac{|\log x|^r}{x^{1/4}} dx \ll_r n^{-1/4+2\epsilon},$$

where the symbol \ll_r indicates that the implicit constant depends on r . \square

Lemma 6. *If $y > 0$, then $\log y = \int_0^\infty \frac{1}{x}(e^{-x} - e^{-xy}) dx$.*

Proof. The left side is $\int_1^y \int_0^\infty e^{-xt} dx dt$; the right side is $\int_0^\infty \int_1^y e^{-xt} dt dx$. \square

Lemma 7. *If t_1, \dots, t_r are real numbers in $[0, 1]$, then*

$$\int_{\mathfrak{U}_n} \prod_{\ell=1}^r \int_0^\infty \frac{1}{x_\ell} |e^{-x_\ell} - e^{-x_\ell |f(e(t_\ell))|^2/n}| dx_\ell d\mu_n < \infty.$$

Proof. Using a strategy similar to that of the proof of [Lemma 5](#), and adopting the notation used there, we find

$$\begin{aligned} & \int_{\mathfrak{U}_n} \prod_{\ell=1}^r \int_0^\infty \frac{1}{x_\ell} |e^{-x_\ell} - e^{-x_\ell |f(e(t_\ell))|^2/n}| dx_\ell d\mu_n \\ & \leq \left\| \frac{1}{r} \sum_{\ell=1}^r \int_0^\infty \frac{1}{x_\ell} |e^{-x_\ell} - e^{-x_\ell |f(e(t_\ell))|^2/n}| dx_\ell \right\|_r^r \\ & \leq \left(\frac{1}{r} \sum_{\ell=1}^r \left\| \int_0^\infty \frac{1}{x_\ell} |e^{-x_\ell} - e^{-x_\ell |f(e(t_\ell))|^2/n}| dx_\ell \right\|_r \right)^r \\ & = \left\| \int_0^\infty \frac{1}{x} |e^{-x} - e^{-x |f(e(1))|^2/n}| dx \right\|_r^r = E(|\log U_n|^r) \end{aligned}$$

by [Lemma 6](#), and $E(|\log U_n|^r) < \infty$ from the proof of [Lemma 5](#). \square

We now prove our first main result on unimodular polynomials.

Proof of [Theorem 1](#). For distinct real numbers t_1, \dots, t_r in $[0, 1]$, let $W_{\mathbf{t},n}$ denote the random variable whose value is $(1/n^r) \prod_{\ell=1}^r |f(e(t_\ell))|^2$, where $f \in \mathfrak{U}_n$ is selected at random. Selecting $m_1 = \dots = m_r = m$ in [Lemma 2](#), we see that the limiting value of the m -th moment of $W_{\mathbf{t},n}$ as n grows large is

$$(2-5) \quad \lim_{n \rightarrow \infty} E(W_{\mathbf{t},n}^m) = (m!)^r.$$

Since $\sum_{m \geq 1} m!^{-r/2m} = \infty$ for $r = 1$ or 2 , by Carleman's test [[Shiryayev 1996](#), p. 296] there is a unique distribution function F_r having these moments, and it also follows from [[Chung 1974](#), Theorem 4.5.5] that $W_{\mathbf{t},n}$ converges in distribution to a

random variable V_r with distribution function F_r as $n \rightarrow \infty$. It is straightforward to verify that the density function of V_r is

$$(2-6) \quad v_r(x) = \begin{cases} e^{-x} & \text{if } r = 1, \\ \int_0^\infty \frac{1}{t} e^{-t-x/t} dt & \text{if } r = 2, \end{cases}$$

up to a set of measure zero. Note that $v_2(x) = 2K_0(2\sqrt{x})$, where $K_\nu(x)$ is the modified Bessel function of the second kind.

In particular, for $r = 1$ (and writing t for t_1) it follows from [Chung 1974, Theorem 4.5.2] that $E(U_{t,n}^p) \rightarrow E(V_1^p)$ for all $p > 0$, so

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \left(\frac{|f(e(t))|}{\sqrt{n}} \right)^p d\mu_n = \int_0^\infty x^{p/2} e^{-x} dx = \Gamma\left(1 + \frac{p}{2}\right)$$

for each t . The density function for $U_{t,n}$ is independent of t , so this convergence is uniform in t . Therefore

$$(2-7) \quad \lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \frac{\|f\|_p^p}{n^{p/2}} d\mu_n = \Gamma\left(1 + \frac{p}{2}\right).$$

This establishes the asymptotic value for the arithmetic mean of the normalized L_p norm of the unimodular polynomials.

For the geometric mean of the normalized measure, the sequence of random variables $\{\log U_{t,n}\}_{n \geq 1}$ is uniformly integrable by Lemma 4, so $E(\log U_{t,n}) \rightarrow E(\log V_1)$ as $n \rightarrow \infty$ for each t . Therefore

$$\lim_{n \rightarrow \infty} E(\log \sqrt{U_{t,n}}) = \int_0^\infty e^{-x} \log \sqrt{x} dx = -\frac{\gamma}{2},$$

and so

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \log\left(\frac{M(f)}{\sqrt{n}}\right) d\mu_n = -\frac{\gamma}{2}.$$

To calculate the arithmetic mean for the measure, we first observe that

$$(2-8) \quad \begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \frac{M(f)}{\sqrt{n}} d\mu_n &= \lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \exp\left(\int_0^1 \log\left(\frac{|f(e(t))|}{\sqrt{n}}\right) dt\right) d\mu_n \\ &= \lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \sum_{r \geq 0} \frac{1}{r!} \left(\int_0^1 \log\left(\frac{|f(e(t))|}{\sqrt{n}}\right) dt\right)^r d\mu_n \\ &= \lim_{n \rightarrow \infty} \sum_{r \geq 0} \frac{1}{r!} \left(\int_0^1 \dots \int_0^1 \int_{\mathfrak{A}_n} \prod_{\ell=1}^r \log\left(\frac{|f(e(t_\ell))|}{\sqrt{n}}\right) d\mu_n dt_1 \dots dt_r\right) \\ &= \sum_{r \geq 0} \frac{1}{r!} \left(\int_0^1 \dots \int_0^1 \lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n} \prod_{\ell=1}^r \log\left(\frac{|f(e(t_\ell))|}{\sqrt{n}}\right) d\mu_n dt_1 \dots dt_r\right). \end{aligned}$$

The validity of switching the order of integration follows from [Lemma 5](#) and Fubini's theorem. Next, using [Lemmata 3, 6, and 7](#), and uniformity in t , and noting that the subset of $[0, 1)^n$ where the t_1, \dots, t_r are not distinct forms a set of measure 0 in \mathfrak{U}_n and so does not affect the integral, we find that

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \prod_{\ell=1}^r \log\left(\frac{|f(e(t_\ell))|}{\sqrt{n}}\right) d\mu_n \\
&= \frac{1}{2^r} \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \prod_{\ell=1}^r \left(\int_0^\infty \frac{1}{x_\ell} \left(e^{-x_\ell} - e^{-x_\ell |f(e(t_\ell))|^2/n} \right) dx_\ell \right) d\mu_n \\
&= \frac{1}{2^r} \int_0^\infty \dots \int_0^\infty \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \prod_{\ell=1}^r \frac{1}{x_\ell} \left(e^{-x_\ell} - e^{-x_\ell |f(e(t_\ell))|^2/n} \right) d\mu_n dx_1 \dots dx_r \\
&= \frac{1}{2^r} \int_0^\infty \dots \int_0^\infty \prod_{\ell=1}^r \frac{1}{x_\ell} \left(e^{-x_\ell} - \frac{1}{1+x_\ell} \right) dx_1 \dots dx_r \\
&= \frac{1}{2^r} \left(\int_0^\infty \frac{1}{x} \left(e^{-x} - \frac{1}{1+x} \right) dx \right)^r = \left(-\frac{\gamma}{2} \right)^r.
\end{aligned}$$

Therefore, from [\(2-8\)](#) we conclude that

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \frac{M(f)}{\sqrt{n}} d\mu_n = \sum_{r=0}^{\infty} \frac{1}{r!} \int_0^1 \dots \int_0^1 \left(-\frac{\gamma}{2} \right)^r dt_1 \dots dt_r = e^{-\gamma/2}.$$

For the geometric mean of the L_p norm when p is an even integer, we use a similar strategy and [\(2-5\)](#) to obtain

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \log\left(\int_0^1 \frac{|f(e(t))|^p}{n^{p/2}} dt \right) d\mu_n \\
&= \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \int_0^\infty \frac{1}{x} \left(\exp(-x) - \exp\left(-x \int_0^1 \frac{|f(e(t))|^p}{n^{p/2}} dt\right) \right) dx d\mu_n \\
&= \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \int_0^\infty \frac{1}{x} \left(\exp(-x) - \sum_{r \geq 0} \frac{(-x)^r}{r!} \prod_{\ell=1}^r \left(\int_0^1 \frac{|f(e(t_\ell))|^p}{n^{p/2}} dt_\ell \right) \right) dx d\mu_n \\
&= \int_0^\infty \frac{1}{x} \left(\exp(-x) - \sum_{r \geq 0} \frac{(-x)^r}{r!} \int_0^1 \dots \int_0^1 \lim_{n \rightarrow \infty} E(W_{\mathbf{t},n}^{p/2}) dt_1 \dots dt_r \right) dx \\
&= \int_0^\infty \frac{1}{x} \left(\exp(-x) - \sum_{r \geq 0} \frac{(-x)^r}{r!} \Gamma\left(1 + \frac{p}{2}\right)^r \right) dx \\
&= \int_0^\infty \frac{1}{x} \left(e^{-x} - e^{-x \Gamma(1+p/2)} \right) dx = \log \Gamma\left(1 + \frac{p}{2}\right). \quad \square
\end{aligned}$$

The limiting distribution of $W_{t,n}$ for $r \geq 3$ does not follow from the computation of the moments in (2-5), since in this case the sum $\sum_{m \geq 1} m!^{-r/2m}$ converges, and so Carleman's criterion is not satisfied. In fact, Berg [2005] has shown that for any fixed $r \geq 3$, the sequence of the moments $\{(m!)^r\}_{m \geq 1}$ does not determine a unique distribution function. We suspect that the density function $v_r(x)$ in question is given by

$$(2-9) \quad v_r(x) = \int_0^\infty \dots \int_0^\infty \exp\left(-x \prod_{\ell=1}^{r-1} x_\ell^{-1}\right) \prod_{\ell=1}^{r-1} \frac{e^{-x_\ell}}{x_\ell} dx_1 \dots dx_{r-1},$$

although we are unable to prove this.

3. Reciprocal polynomials

We use a similar strategy to prove Theorem 2. For a fixed positive integer n , for convenience we say that two nonnegative integers i and j are *complementary* if $i + j = n - 1$. Thus, if $f(x) = \sum_k a_k x^k \in \mathfrak{A}_n^*$ and i and j are complementary, then the coefficients a_i and a_j of f satisfy $a_i = \omega \overline{a_j}$, where ω is a fixed complex number of modulus 1. Also, for an odd integer $2k - 1$, let $(2k - 1)!!$ denote the product $1 \cdot 3 \cdot 5 \dots (2k - 1) = (2k)!/2^k k!$.

Lemma 8. *Suppose t_1, \dots, t_r are distinct real numbers in $(0, 1)$, and m_1, \dots, m_r are positive integers. Let $M = \sum_{\ell=1}^r m_\ell$. Then*

$$\int_{\mathfrak{A}_n^*} \prod_{\ell=1}^r \frac{|f(e(t_\ell))|^{2m_\ell}}{n^{m_\ell}} d\mu_n^* = \prod_{\ell=1}^r (2m_\ell - 1)!! + O\left(\frac{1}{n}\right);$$

the implicit constant may depend on the m_ℓ and the t_ℓ but is independent of n . Thus,

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{A}_n^*} \prod_{\ell=1}^r \frac{|f(e(t_\ell))|^{2m_\ell}}{n^{m_\ell}} d\mu_n^* = \prod_{\ell=1}^r (2m_\ell - 1)!!.$$

Proof. We compute

$$\begin{aligned} \int_{\mathfrak{A}_n^*} \prod_{\ell=1}^r \frac{|f(e(t_\ell))|^{2m_\ell}}{n^{m_\ell}} d\mu_n^* &= \frac{1}{n^M} \int_{\mathfrak{A}_n^*} \prod_{\ell=1}^r \left| \sum_{k_\ell=0}^{n-1} a_{k_\ell} e(k_\ell t_\ell) \right|^{2m_\ell} d\mu_n^* \\ &= \frac{1}{n^M} \sum_{\substack{I_\ell, J_\ell \in R_{n, m_\ell} \\ 1 \leq \ell \leq r}} \prod_{\ell=1}^r e\left(t_\ell \sum_{k=1}^{m_\ell} (i_{\ell, k} - j_{\ell, k})\right) \int_{\mathfrak{A}_n^*} \prod_{\ell=1}^r \prod_{k=1}^{m_\ell} a_{i_{\ell, k}} \overline{a_{j_{\ell, k}}} d\mu_n^* \\ &= \frac{1}{n^M} \sum_{\substack{I_\ell, J_\ell \in R_{n, m_\ell}^* \\ 1 \leq \ell \leq r}} \prod_{\ell=1}^r e\left(t_\ell \sum_{k=1}^{m_\ell} (i_{\ell, k} - j_{\ell, k})\right). \end{aligned}$$

Here \sum^* denotes the sum over only those $2M$ -tuples $(I_1, \dots, I_r, J_1, \dots, J_r)$ where the value of each index $i_{\ell,k}$ or $j_{\ell,k}$ either matches that of some other index $i_{\ell,k'}$ or $j_{\ell,k'}$, or complements the value of some other such index. We again restrict our consideration to those terms where $(I_1, \dots, I_r, J_1, \dots, J_r)$ can be grouped into M distinct pairs such that each grouping consists of some number $s \equiv M \pmod{2}$ of pairs $(i_{\ell,k}, j_{\ell,k'})$ of matching integers, together with $(M-s)/2$ pairs $(i_{\ell,k}, i_{\ell,k'})$ and $(M-s)/2$ pairs $(j_{\ell,k}, j_{\ell,k'})$ of complementary integers. This way, we have

$$\prod_{k=1}^{m_\ell} a_{i_{\ell,k}} \overline{a_{j_{\ell,k}}} = 1$$

for each ℓ . The number of possible pairings is $\prod_{\ell=1}^r (2m_\ell - 1)!!$, and the statement follows in the same way as [Lemma 2](#). \square

We now prove our main result on reciprocal unimodular polynomials.

Proof of Theorem 2. It is well known that the $2m$ -th moment of the standard normal distribution is $(2m - 1)!!$, so

$$\sqrt{\frac{2}{\pi}} \int_0^\infty x^{2m} e^{-x^2/2} dx = (2m - 1)!!.$$

Therefore,

$$\frac{1}{\sqrt{2\pi}} \int_0^\infty y^{m-1/2} e^{-y/2} dy = (2m - 1)!!,$$

so by Carleman's test the limiting density function as $n \rightarrow \infty$ for the random variables $|f(e(t))|^2/n$ on \mathfrak{U}_n^* for $t \neq 0$ is

$$\rho_t(x) = \frac{e^{-x/2}}{\sqrt{2\pi x}}$$

for almost every x in $[0, \infty)$. Thus, for $t \neq 0$, we have

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n^*} \log\left(\frac{|f(e(t))|}{\sqrt{n}}\right) d\mu_n^* = \int_0^\infty \frac{e^{-x/2}}{\sqrt{2\pi x}} \log \sqrt{x} dx = -\frac{\gamma + \log 2}{2}$$

and

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n^*} \left(\frac{|f(e(t))|}{\sqrt{n}}\right)^p d\mu_n^* = \int_0^\infty \frac{x^{p/2} e^{-x/2}}{\sqrt{2\pi x}} dx = \frac{2^{p/2}}{\sqrt{\pi}} \Gamma\left(\frac{p+1}{2}\right),$$

and the results on the geometric mean of the normalized measure and the arithmetic mean of the normalized L_p norm follow as in the proof of [Theorem 1](#). The remaining statements may be proved in the same way as [Theorem 1](#). \square

4. Moments of the distributions

We now establish the result on the moments of the distributions of the L_p norms and Mahler's measure of unimodular polynomials.

Proof of Theorem 3. For any even integer $p \geq 2$, we compute

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \left(\frac{\|f\|_p^p}{n^{p/2}} \right)^m d\mu_n &= \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \left(\prod_{\ell=1}^m \int_0^1 \frac{|f(e(t))|^p}{n^{p/2}} dt \right) d\mu_n \\ &= \int_0^1 \dots \int_0^1 \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \prod_{\ell=1}^m \frac{|f(e(t_\ell))|^p}{n^{p/2}} d\mu_n dt_1 \dots dt_m \\ &= \int_0^1 \dots \int_0^1 \Gamma\left(1 + \frac{p}{2}\right)^m dt_1 \dots dt_m \\ &= \left(\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \frac{\|f\|_p^p}{n^{p/2}} d\mu_n \right)^m, \end{aligned}$$

by Lemma 2. The computation for Mahler's measure is similar. When $m = 2$, we may use the density function (2-6) to determine that

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \left(\frac{\|f\|_p^p}{n^{p/2}} \right)^2 d\mu_n &= \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \int_0^1 \int_0^1 \frac{|f(e(t_1))|^p}{n^{p/2}} \frac{|f(e(t_2))|^p}{n^{p/2}} dt_1 dt_2 d\mu_n \\ &= \int_0^1 \int_0^1 \lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \frac{|f(e(t_1))|^p}{n^{p/2}} \frac{|f(e(t_2))|^p}{n^{p/2}} d\mu_n dt_1 dt_2 \\ &= \int_0^1 \int_0^1 \int_0^\infty x^{p/2} \int_0^\infty \frac{1}{t} e^{-t-x/t} dt dx dt_1 dt_2 \\ &= \Gamma\left(1 + \frac{p}{2}\right)^2 \\ &= \left(\lim_{n \rightarrow \infty} \int_{\mathfrak{U}_n} \frac{\|f\|_p^p}{n^{p/2}} d\mu_n \right)^2 \end{aligned}$$

for $p > 0$. The computations for \mathfrak{U}_n^* are similar. (The density function for the case $m = 2$ in this case is $K_0(\sqrt{x})/\pi\sqrt{x}$.) □

5. Concluding remarks

It would be of interest to apply this method to determine the expected value in the limit for the normalized Mahler's measure of the Littlewood polynomials, and to study the reciprocal Littlewood polynomials in the same way. We conjecture that the limiting mean values for the Littlewood polynomials match the ones stated in Theorems 1 and 2 for the unimodular polynomials. (This was established already in [Borwein and Lockhart 2001] for the arithmetic mean of the L_p norms with

$p > 0$ in \mathcal{L}_n .) Verifying this for Mahler’s measure in \mathcal{L}_n would thus improve the best known result (1-2) in Mahler’s problem. Indeed, by calculating sequences of moments, one may establish analogues of Lemmata 2 and 8 for the Littlewood polynomials. However, the moment sequences are no longer independent of t in this case, as $t = 0$ and $t = \frac{1}{2}$ behave differently from other $t \in (0, 1)$.

Lower bounds on Mahler’s measure of noncyclotomic Littlewood polynomials are found in [Borwein et al. 2007; Dubickas and Mossinghoff 2005; Borwein et al. 2004; Garza et al. 2010], and a list of the maximum values of Mahler’s measure for polynomials in \mathcal{L}_n for $n \leq 25$ can be found in [Borwein and Mossinghoff 2008]. Also, the distribution of the values of Mahler’s measure for polynomials with arbitrary real or complex coefficients is studied in [Chern and Vaaler 2001; Sinclair 2004], and average L_p norms of other families of polynomials are considered in [Borwein and Choi 2007; Borwein et al. 2008a; Mansour 2004].

Acknowledgment

We thank Andrew Granville for spurring this research by using some calculations of moments to conjecture that the mean value of the normalized Mahler’s measure of Littlewood polynomials approaches $e^{-\gamma/2}$ as the degree grows large.

References

- [Beck 1991] J. Beck, “Flat polynomials on the unit circle — note on a problem of Littlewood”, *Bull. London Math. Soc.* **23**:3 (1991), 269–277. [MR 93b:42002](#) [Zbl 0748.30006](#)
- [Beller and Newman 1973] E. Beller and D. J. Newman, “An extremal problem for the geometric mean of polynomials”, *Proc. Amer. Math. Soc.* **39** (1973), 313–317. [MR 47 #5233](#) [Zbl 0267.30001](#)
- [Berg 2005] C. Berg, “On powers of Stieltjes moment sequences, I”, *J. Theoret. Probab.* **18**:4 (2005), 871–889. [MR 2008i:44008](#) [Zbl 1086.44003](#)
- [Borwein 2002] P. Borwein, *Computational excursions in analysis and number theory*, CMS Books in Mathematics **10**, Springer, New York, 2002. [MR 2003m:11045](#) [Zbl 1020.12001](#)
- [Borwein and Choi 2007] P. Borwein and K.-K. S. Choi, “The average norm of polynomials of fixed height”, *Trans. Amer. Math. Soc.* **359**:2 (2007), 923–936. [MR 2007h:11036](#) [Zbl 1198.11030](#)
- [Borwein and Lockhart 2001] P. Borwein and R. Lockhart, “The expected L_p norm of random polynomials”, *Proc. Amer. Math. Soc.* **129**:5 (2001), 1463–1472. [MR 2001m:60124](#) [Zbl 0999.30004](#)
- [Borwein and Mossinghoff 2008] P. Borwein and M. J. Mossinghoff, “Barker sequences and flat polynomials”, pp. 71–88 in *Number theory and polynomials*, London Math. Soc. Lecture Note Ser. **352**, Cambridge Univ. Press, 2008. [MR 2009k:11040](#)
- [Borwein et al. 2004] P. Borwein, K. G. Hare, and M. J. Mossinghoff, “The Mahler measure of polynomials with odd coefficients”, *Bull. London Math. Soc.* **36**:3 (2004), 332–338. [MR 2004m:11177](#) [Zbl 1143.11349](#)
- [Borwein et al. 2007] P. Borwein, E. Dobrowolski, and M. J. Mossinghoff, “Lehmer’s problem for polynomials with odd coefficients”, *Ann. of Math. (2)* **166**:2 (2007), 347–366. [MR 2008j:11153](#) [Zbl 1172.11034](#)

- [Borwein et al. 2008a] P. Borwein, K.-K. S. Choi, and I. Mercer, “Expected norms of zero-one polynomials”, *Canad. Math. Bull.* **51**:4 (2008), 497–507. [MR 2009m:11036](#) [Zbl 1173.11015](#)
- [Borwein et al. 2008b] P. Borwein, R. Ferguson, and J. Knauer, “The merit factor problem”, pp. 52–70 in *Number theory and polynomials*, London Math. Soc. Lecture Note Ser. **352**, Cambridge Univ. Press, 2008. [MR 2010b:11030](#)
- [Chern and Vaaler 2001] S.-J. Chern and J. D. Vaaler, “The distribution of values of Mahler’s measure”, *J. Reine Angew. Math.* **540** (2001), 1–47. [MR 2003a:11137](#) [Zbl 0986.11017](#)
- [Chung 1974] K. L. Chung, *A course in probability theory*, 2nd ed., Probability and Mathematical Statistics **21**, Academic Press, New York, 1974. [MR 49 #11579](#) [Zbl 0345.60003](#)
- [Dubickas and Mossinghoff 2005] A. Dubickas and M. J. Mossinghoff, “Auxiliary polynomials for some problems regarding Mahler’s measure”, *Acta Arith.* **119**:1 (2005), 65–79. [MR 2006e:11162](#) [Zbl 1074.11018](#)
- [Erdélyi and Lubinsky 2007] T. Erdélyi and D. S. Lubinsky, “Large sieve inequalities via subharmonic methods and the Mahler measure of the Fekete polynomials”, *Canad. J. Math.* **59**:4 (2007), 730–741. [MR 2009e:41030](#) [Zbl 1166.41004](#)
- [Erdős 1962] P. Erdős, “An inequality for the maximum of trigonometric polynomials”, *Ann. Polon. Math.* **12** (1962), 151–154. [MR 25 #5330](#) [Zbl 0106.27702](#)
- [Fielding 1970] G. T. Fielding, “The expected value of the integral around the unit circle of a certain class of polynomials”, *Bull. London Math. Soc.* **2** (1970), 301–306. [MR 43 #6408](#) [Zbl 0208.09402](#)
- [Garza et al. 2010] J. Garza, M. I. M. Ishak, M. J. Mossinghoff, C. G. Pinner, and B. Wiles, “Heights of roots of polynomials with odd coefficients”, *J. Théor. Nombres Bordeaux* **22**:2 (2010), 369–381. [Zbl 05862105](#)
- [Kahane 1980] J.-P. Kahane, “Sur les polynômes à coefficients unimodulaires”, *Bull. London Math. Soc.* **12**:5 (1980), 321–342. [MR 82a:30003](#) [Zbl 0443.30005](#)
- [Kluyver 1906] J. C. Kluyver, “A local probability problem”, *Verh. Natuurkd. K. Nederl. Akad. Wetensch.* **8** (1906), 341–350.
- [Littlewood 1968] J. E. Littlewood, *Some problems in real and complex analysis*, D. C. Heath, Lexington, MA, 1968. [MR 39 #5777](#) [Zbl 0185.11502](#)
- [Mahler 1963] K. Mahler, “On two extremum properties of polynomials”, *Illinois J. Math.* **7** (1963), 681–701. [MR 28 #192](#) [Zbl 0117.04003](#)
- [Mansour 2004] T. Mansour, “Average norms of polynomials”, *Adv. in Appl. Math.* **32**:4 (2004), 698–708. [MR 2004m:30008](#) [Zbl 1066.30005](#)
- [Newman and Byrnes 1990] D. J. Newman and J. S. Byrnes, “The L^4 norm of a polynomial with coefficients ± 1 ”, *Amer. Math. Monthly* **97**:1 (1990), 42–45. [MR 91d:30006](#) [Zbl 0722.11018](#)
- [Shiryayev 1996] A. N. Shiryayev, *Probability*, 2nd ed., Graduate Texts in Mathematics **95**, Springer, New York, 1996. [MR 97c:60003](#) [Zbl 0835.60002](#)
- [Sinclair 2004] C. D. Sinclair, “The distribution of Mahler’s measures of reciprocal polynomials”, *Int. J. Math. Math. Sci.* 49-52 (2004), 2773–2786. [MR 2006i:33034](#) [Zbl 1097.33018](#)
- [Smyth 1971] C. J. Smyth, “On the product of the conjugates outside the unit circle of an algebraic integer”, *Bull. London Math. Soc.* **3** (1971), 169–175. [MR 44 #6641](#) [Zbl 0235.12003](#)

KWOK-KWONG STEPHEN CHOI
DEPARTMENT OF MATHEMATICS
SIMON FRASER UNIVERSITY
BURNABY, BRITISH COLUMBIA V5A 1S6
CANADA

kkchoi@math.sfu.ca

<http://www.math.sfu.ca/~kkchoi>

MICHAEL J. MOSSINGHOFF
DEPARTMENT OF MATHEMATICS
DAVIDSON COLLEGE
DAVIDSON, NORTH CAROLINA 28035-6996
UNITED STATES

mimossinghoff@davidsen.edu

<http://www.davidson.edu/math/mossinghoff>

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Mathew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 252 No. 1 July 2011

Some dynamic Wirtinger-type inequalities and their applications	1
RAVI P. AGARWAL, MARTIN BOHNER, DONAL O'REGAN and SAMIR H. SAKER	
Splitting criteria for vector bundles on higher-dimensional varieties	19
PARSA BAKHTARY	
Average Mahler's measure and L_p norms of unimodular polynomials	31
KWOK-KWONG STEPHEN CHOI and MICHAEL J. MOSSINGHOFF	
Tate resolutions and Weyman complexes	51
DAVID A. COX and EVGENY MATEROV	
On pointed Hopf algebras over dihedral groups	69
FERNANDO FANTINO and GASTON ANDRÉS GARCIA	
Integral topological quantum field theory for a one-holed torus	93
PATRICK M. GILMER and GREGOR MASBAUM	
Knot 4-genus and the rank of classes in $W(\mathbb{Q}(t))$	113
CHARLES LIVINGSTON	
Roots of Toeplitz operators on the Bergman space	127
ISSAM LOUHICHI and NAGISETTY V. RAO	
Uniqueness of the foliation of constant mean curvature spheres in asymptotically flat 3-manifolds	145
SHIGUANG MA	
On the multiplicity of non-iterated periodic billiard trajectories	181
MARCO MAZZUCHELLI	
A remark on Einstein warped products	207
MICHELE RIMOLDI	
Exceptional Dehn surgery on large arborescent knots	219
YING-QING WU	
Harnack estimates for the linear heat equation under the Ricci flow	245
XIAORUI ZHU	