

# *Pacific Journal of Mathematics*

**HARNACK ESTIMATES FOR THE LINEAR HEAT EQUATION  
UNDER THE RICCI FLOW**

XIAORUI ZHU

# HARNACK ESTIMATES FOR THE LINEAR HEAT EQUATION UNDER THE RICCI FLOW

XIAORUI ZHU

**We consider the linear heat equation on a manifold that evolves under the Ricci flow. The gradient estimates for positive solutions as well as Li–Yau type inequalities are given in this paper. Both the case where  $M$  is a complete manifold without boundary and the case where  $M$  is compact are considered. We have also obtained the Harnack inequalities for the heat equation on  $M$  by previous results.**

## 1. Introduction

The heat equation is a classical subject that has been extensively studied and has lead to many important results, especially in studies of differential geometry. One of the important techniques used in studying the heat equation is the differential Harnack inequality developed by Li and Yau [1986]. This is also applied to Ricci flow by Hamilton [1993], and plays an important role in solving the Poincaré conjecture.

We consider the positive solutions of the linear heat equation on a manifold  $M$  that evolves under the Ricci flow. A series of gradient estimates are obtained for such solutions, including several Li–Yau-type inequalities. The manifold  $M$  considered here is a complete manifold without boundary.

Let  $M$  be a manifold without boundary, and  $(M, g(x, t))_{t \in [0, T]}$  be a complete solution to the Ricci flow

$$(1) \quad \frac{\partial}{\partial t} g(x, t) = -2 \operatorname{Ric}(x, t), \quad x \in M, t \in [0, T].$$

We assume that its curvature remains uniformly bounded for all  $t \in [0, T]$ . Consider a positive function  $u(x, t)$  defined on  $M \times [0, T]$  solving the equation

$$(2) \quad \frac{\partial u}{\partial t} = \Delta u - q(x, t)u, \quad x \in M, \quad t \in [0, T],$$

where  $\Delta$  stands for the Laplacian given by  $g(x, t)$  and  $q(x, t)$  is a  $C^2$  function defined on  $M \times [0, T]$ . Noticing that  $\Delta$  depends on the parameter  $t$ , we study the

---

MSC2000: 53C44.

Keywords: linear heat equation, Harnack estimate.

linear heat equation (2) along with the Ricci flow (1). Equation (1) provides us with additional information about the coefficients of the operator  $\Delta$  appearing in (2), but is itself fully independent of (2).

## 2. Gradient estimates

Firstly, we introduce a cutoff function on  $B_{\rho,T}$ . the notation  $B_{\rho,T}$  stands for the set  $\{(\chi, t) \in M \times [0, T] \mid \text{dist}(\chi, x_0, t) < \rho\}$ , which satisfies the basic analytical results stated in the following lemma.

**Lemma 2.1.** *Given  $\tau \in (0, T]$ , there exists a smooth function*

$$\bar{\Psi} : [0, \infty) \times [0, T] \rightarrow \mathbb{R}$$

*satisfying the following requirements:*

1. *The support of  $\bar{\Psi}(r, t)$  is a subset of  $[0, \rho] \times [0, T]$ , and  $0 \leq \bar{\Psi}(r, t) \leq 1$  in  $[0, \rho] \times [0, T]$ .*

2. *The equalities*

$$\bar{\Psi}(r, t) = 1 \quad \text{and} \quad \frac{\partial \bar{\Psi}}{\partial r}(r, t) = 0$$

*hold on  $[0, \rho/2] \times [\tau, T]$  and  $[0, \rho/2] \times [0, T]$ , respectively.*

3. *The estimate*

$$\left| \frac{\partial \bar{\Psi}}{\partial t} \right| \leq \frac{\bar{C} \bar{\Psi}^{1/2}}{\tau}$$

*is satisfied on  $[0, \infty) \times [0, T]$  for some  $\bar{C} > 0$ , and  $\bar{\Psi}(r, 0) = 0$  for all  $r \in [0, \infty)$ .*

4. *The inequalities*

$$-\frac{C_\alpha \bar{\Psi}^\alpha}{\rho} \leq \frac{\partial \bar{\Psi}}{\partial r} \leq 0 \quad \text{and} \quad \left| \frac{\partial^2 \bar{\Psi}}{\partial r^2} \right| \leq \frac{C_\alpha \bar{\Psi}^\alpha}{\rho^2}$$

*hold on  $[0, \infty) \times [0, T]$  for every  $a \in (0, 1)$ , with a constant  $C_\alpha$  dependent on  $a$ .*

This lemma was first introduced in [Bailesteanu et al. 2010]. In the following part of this section, we establish Li–Yau-type inequalities for system (1)–(2) and obtain a local and a global estimate. To this end, we must introduce an auxiliary function to apply the maximum principle on it. The following lemma deals with the evolution equation of the auxiliary function.

**Lemma 2.2.** *Suppose  $(M, g(x, t))_{t \in [0, T]}$  is a complete solution to the Ricci flow (1). Assume that  $-k_1 g(x, t) \leq \text{Ric}(x, t) \leq k_2 g(x, t)$  for some  $k_1, k_2 > 0$  and all  $(x, t) \in B_{\rho,T}$ . Suppose  $u : M \times [0, T] \rightarrow \mathbb{R}$  is a smooth positive function satisfying the heat equation (2), and  $q(x, t)$  is a  $C^2$  function defined on  $M \times [0, T]$ . Given*

$\alpha \geq 1$ , define  $f = \log u$  and  $F = t(|\nabla f|^2 - \alpha f_t - \alpha q)$ . The estimate

$$(3) \quad \left( \Delta - \frac{\partial}{\partial t} \right) F \geq -2 \langle \nabla f, \nabla F \rangle - F t^{-1} - 2k_1 \alpha t |\nabla f|^2 \\ + \frac{2a\alpha t}{n} (|\nabla f|^2 - f_t - q)^2 - \frac{\alpha t n}{2b} \max\{k_1^2, k_2^2\} - t\alpha \Delta q - 2t(\alpha - 1) \nabla f \nabla q$$

holds for any  $a, b > 0$ , such that  $a + b = 1/\alpha$ .

*Proof.* We begin with finding a convenient bound on  $\Delta F$ . Notice that

$$\Delta F = t(2f_{ji}^2 + 2f_j f_{jii} - \alpha \Delta(f_t) - a \Delta q), \quad x \in M, \quad t \in [0, T].$$

By the assumption on the Ricci curvature of  $M$ , it follows that

$$f_j f_{jii} = f_j f_{iij} + R_{ij} f_i f_j \geq \langle \nabla f, \nabla \Delta f \rangle - k_1 |\nabla f|^2$$

at an arbitrary point  $(x, t) \in B_{\rho, T}$ . Using (1), we get that

$$\Delta(f_t) = (\Delta f)_t - 2 \sum_{i,j=1}^n R_{ij} f_{ij}.$$

Thus, the estimate

$$\Delta F \geq t[2f_{ji}^2 + 2 \langle \nabla f, \nabla \Delta f \rangle - 2k_1 |\nabla f|^2 - \alpha (\Delta f)_t + 2\alpha R_{ij} f_{ij} - \alpha \Delta q]$$

holds at  $(x, t) \in B_{\rho, T}$ . The next step is to find a suitable bound on those terms in the right side involving  $f_{ij}$  by completing the square. Specifically, we find that

$$\sum_{i,j=1}^n (f_{ij}^2 + \alpha R_{ij} f_{ij}) = \sum_{i,j=1}^n [(a\alpha + b\alpha) f_{ij}^2 + \alpha R_{ij} f_{ij}] \\ = \sum_{i,j=1}^n \left( a\alpha f_{ij}^2 + \alpha \left( \sqrt{b} f_{ij} + \frac{R_{ij}}{2\sqrt{b}} \right)^2 - \frac{\alpha}{4b} R_{ij}^2 \right) \\ \geq \sum_{i,j=1}^n \left( a\alpha f_{ij}^2 - \frac{\alpha}{4b} R_{ij}^2 \right)$$

at  $(x, t) \in B_{\rho, T}$  for any  $a, b > 0$ , such that  $a + b = 1/\alpha$ . Using the assumptions in the lemma and the standard inequality

$$\sum_{i,j=1}^n f_{ij}^2 \geq \frac{(\Delta f)^2}{n},$$

we obtain the estimate

$$\sum_{i,j=1}^n (f_{ij}^2 + \alpha R_{ij} f_{ij}) \geq \frac{a\alpha}{n} (\Delta f)^2 - \frac{\alpha}{4b} \max\{k_1^2, k_2^2\}, \quad (x, t) \in B_{\rho, T}.$$

Obviously, we conclude that

$$\begin{aligned}
 (4) \quad \Delta F &\geq t \left[ \frac{2a\alpha}{n} (\Delta f)^2 + 2 \langle \nabla f, \nabla \Delta f \rangle - 2k_1 |\nabla f|^2 - \alpha (\Delta f)_t \right. \\
 &\quad \left. - \alpha \Delta q - \frac{\alpha n}{2b} \max\{k_1^2, k_2^2\} \right] \\
 &= \frac{2a\alpha t}{n} (|\nabla f|^2 - f_t - q)^2 - 2t \nabla f \nabla (|\nabla f|^2 - f_t - q) \\
 &\quad - 2k_1 t |\nabla f|^2 - t\alpha \Delta q + \alpha t (|\nabla f|^2 - f_t - q)_t - \frac{\alpha n t}{2b} \max\{k_1^2, k_2^2\}
 \end{aligned}$$

in the set  $B_{\rho, T}$ .

This gives a convenient bound for  $\Delta F$ . Now we consider the derivative of  $F$  in  $t \in [0, T]$ . It is not hard to compute that

$$\frac{\partial F}{\partial t} = \frac{F}{t} + t (|\nabla f|^2 - \alpha f_t - \alpha q)_t.$$

Subtracting this from (4), we see that the inequality

$$\begin{aligned}
 \left( \Delta - \frac{\partial}{\partial t} \right) F &\geq \frac{2a\alpha t}{n} (|\nabla f|^2 - f_t - q)^2 - 2t \nabla f \nabla (|\nabla f|^2 - f_t - q) - 2k_1 t |\nabla f|^2 \\
 &\quad - t\alpha \Delta q - \frac{\alpha n t}{2b} \max\{k_1^2, k_2^2\} - \frac{F}{t} + (\alpha - 1)t (|\nabla f|^2)_t
 \end{aligned}$$

holds in the set  $B_{\rho, T}$ , with  $t > 0$ . We need the estimate on  $|\nabla f|_t^2$  in order to obtain (3) from this inequality. The Ricci flow equation (1) and the assumptions of the lemma imply

$$|\nabla f|_t^2 = 2\nabla f \nabla (f_t) + 2 \operatorname{Ric}(\nabla f, \nabla f) \geq 2\nabla f \nabla (f_t) - 2k_1 |\nabla f|^2$$

at  $(x, t) \in B_{\rho, T}$ . As a consequence,

$$\begin{aligned}
 \left( \Delta - \frac{\partial}{\partial t} \right) F &\geq \frac{2a\alpha t}{n} (|\nabla f|^2 - f_t - q)^2 - 2k_1 \alpha |\nabla f|^2 - t\alpha \Delta q \\
 &\quad - \frac{\alpha n t}{2b} \max\{k_1^2, k_2^2\} - \frac{F}{t} - 2\nabla f \nabla F - 2(\alpha - 1)t \nabla f \nabla q
 \end{aligned}$$

in  $B_{\rho, T}$ . The desired assertion follows.  $\square$

Now we can consider the local space-time gradient estimate with Lemma 2.2. In the following part,  $n$  is the dimension of  $M$ .

**Theorem 2.3.** *Let  $(M, g(x, t))_{t \in [0, T]}$  be a complete solution to the Ricci flow (1). Suppose  $-k_1 g(x, t) \leq \operatorname{Ric}(x, t) \leq k_2 g(x, t)$  for some  $k_1, k_2 > 0$  and all  $(x, t) \in B_{\rho, T}$ . Suppose  $u : M \times [0, T] \rightarrow \mathbb{R}$  is a smooth positive function solving the heat equation (2), and  $q(x, t)$  is a  $C^2$  function defined on  $M \times [0, T]$ ,  $|\nabla q| \leq \gamma$ ,  $|\Delta q| \leq \theta$ . There exists a constant  $C'$  that depends only on the dimension of  $M$  and*

satisfies the estimate

$$(5) \quad \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} - \alpha q \leq C' \alpha^2 \left( \frac{\alpha^2}{\rho^2(\alpha-1)} + \frac{1}{t} + \max\{k_1, k_2\} \right) + \frac{nk_1\alpha^3}{\alpha-1} + \frac{\gamma(\alpha-1)}{\alpha} \sqrt{\frac{2\alpha k_1}{n}} + 2\alpha\sqrt{\alpha n\theta}$$

for all  $\alpha > 1$  and all  $(x, t) \in B_{\rho/2, T}$  with  $t \neq 0$ .

*Proof.* We will use the same notation,  $f = \log u$  and  $F = t(|\nabla f|^2 - \alpha f_t - \alpha q)$ , as in [Lemma 2.2](#) and denote  $\max\{k_1, k_2\}$  as  $\bar{k}$ . For a fixed  $\tau \in (0, T]$  and fixed  $\bar{\Psi}(r, t)$  satisfying the conditions in [Lemma 2.2](#), define  $\Psi : M \times [0, T] \rightarrow \mathbb{R}$  by setting

$$\Psi(x, t) = \bar{\Psi}(\text{dist}(x, x_0, t), t).$$

We will establish the inequality in [Theorem 2.3](#) at  $(x, \tau)$  for  $x \in M$ , such that  $\text{dist}(x, x_0, \tau) < \rho/2$ . This will complete the proof.

From [Lemma 2.2](#), some straightforward computations lead to

$$(6) \quad \left( \Delta - \frac{\partial}{\partial t} \right) (\Psi F) \geq -2\nabla f \nabla (\Psi F) + 2F \nabla f \nabla \Psi + \left( \frac{2\alpha\alpha t}{n} (|\nabla f|^2 - f_t - q)^2 - 2k_1\alpha t |\nabla f|^2 - \frac{\alpha n t}{2b} \bar{k}^2 - t\alpha \Delta q - \frac{F}{t} - 2(\alpha-1)t \nabla f \nabla q \right) \Psi + (\Delta \Psi) F + 2 \frac{\nabla \Psi}{\Psi} \nabla (\Psi F) - 2 \frac{|\nabla \Psi|^2}{\Psi} F - \frac{\partial \Psi}{\partial t} F.$$

This inequality holds in the part of  $B_{\rho, T}$  where  $\Psi(x, t)$  is smooth and strictly positive. Let  $(x_1, t_1)$  be a maximum point for the function  $\Psi F$  in the set

$$\{(x, t) \mid 0 \leq t \leq \tau, d(x, x_0, t) \leq \rho\}.$$

Then we have

$$(7) \quad 0 \geq 2F \nabla f \nabla \Psi + \left( \frac{2\alpha\alpha t_1}{n} (|\nabla f|^2 - f_t - q)^2 - 2k_1\alpha t_1 |\nabla f|^2 - \frac{\alpha n t_1}{2b} \bar{k}^2 - t_1\alpha \Delta q - \frac{F}{t_1} - 2(\alpha-1)t_1 \nabla f \nabla q \right) \Psi + (\Delta \Psi) F - 2 \frac{|\nabla \Psi|^2}{\Psi} F - \frac{\partial \Psi}{\partial t} F$$

at  $(x_1, t_1)$ . We will now use (7) to show that a certain quadratic expression in  $\Psi F$  is nonpositive. The desired result will then follow.

We recall [Lemma 2.1](#) and apply the Laplacian comparison theorem to obtain

$$\frac{|\nabla \Psi|^2}{\Psi} \leq \frac{C_{1/2}^2}{\rho^2},$$

$$\Delta \Psi \geq -\frac{C_{1/2}\Psi^{1/2}}{\rho^2} - \frac{C_{1/2}\Psi^{1/2}}{\rho}(n-1)\sqrt{k_1} \coth(\sqrt{k_1}\rho) \geq -\frac{d_1}{\rho^2} - \frac{d_1\Psi^{1/2}}{\rho}\sqrt{k_1}$$

at the point  $(x_1, t_1)$ , where  $d_1$  is a positive constant depending on  $n$ . There exists  $C > 0$  such that

$$-\frac{\partial \Psi}{\partial t} \geq -\frac{\bar{C}\Psi^{1/2}}{\tau} - C_{1/2}\bar{k}\Psi^{1/2}.$$

Using these observations along with (6), we get the estimate

$$\begin{aligned} 0 \geq & -2F |\nabla f| |\nabla \Psi| + \left( \frac{2a\alpha t_1}{n} (|\nabla f|^2 - f_t - q)^2 - 2k_1\alpha t_1 |\nabla f|^2 \right. \\ & \left. - t_1\alpha \Delta q - \frac{\alpha n t_1}{2b} \bar{k}^2 - \frac{F}{t_1} - 2(\alpha - 1)t_1 \nabla f \nabla q \right) \Psi \\ & + d_2 \left( -\frac{1}{\rho^2} - \frac{\Psi^{1/2}}{\rho} \sqrt{k_1} - \frac{\Psi^{1/2}}{\tau} - \bar{k} \Psi^{1/2} \right) F \end{aligned}$$

at  $(x_1, t_1)$ , where

$$d_2 = \{3d_1, C_{1/2}, 3C_{1/2}^2, \bar{C}\}.$$

Multiplying by  $t\Psi$  and making a few elementary manipulations, we obtain

$$\begin{aligned} (8) \quad 0 \geq & -2t_1 F \frac{C_{1/2}\Psi^{3/2}}{\rho} |\nabla f| \\ & + \frac{2t_1^2}{n} \left( a\alpha \Psi^2 (|\nabla f|^2 - f_t - q)^2 - nk_1\alpha \Psi^2 |\nabla f|^2 \right. \\ & \left. - \frac{n^2\alpha}{4b} \bar{k}^2 \Psi^2 - \frac{n}{2} \alpha \Psi^2 \Delta q - n(\alpha - 1) \nabla f \nabla q \Psi^2 \right) \\ & + d_2 \left( -\frac{1}{\rho^2} - \frac{1}{\rho} \sqrt{k_1} - \frac{1}{\tau} - \bar{k} \right) t_1 (\Psi F) - \Psi F \end{aligned}$$

at  $(x_1, t_1)$ . Our next step is to estimate the first two terms on the right side. In order to finish, we introduce the following notations.

Define

$$y = \Psi |\nabla f|^2 \quad \text{and} \quad z = \Psi(f_t + q).$$

It is clear that

$$y^{1/2}(y - \alpha z) = \frac{\Psi^{3/2} F |\nabla f|}{t}$$

when  $t \neq 0$ , which yields

$$\begin{aligned}
 (9) \quad & -2t_1 F \frac{C_{1/2} \Psi^{3/2}}{\rho} |\nabla f| + \frac{2t_1^2}{n} \left( a\alpha \Psi^2 (|\nabla f|^2 - f_t - q)^2 \right. \\
 & \quad \left. - nk_1 \alpha \Psi^2 |\nabla f|^2 - \frac{n^2 \alpha}{4b} \bar{k}^2 \Psi^2 - \frac{n}{2} \alpha \Psi^2 \Delta q - n(\alpha - 1) \nabla f \nabla q \Psi^2 \right) \\
 & \geq \frac{2t_1^2}{n} \left( a\alpha (y - z)^2 - nk_1 \alpha y - \frac{n^2 \alpha}{4b} \bar{k}^2 \Psi^2 \right. \\
 & \quad \left. - \frac{n C_{1/2}}{\rho} y^{1/2} (y - \alpha z) - \frac{n}{2} \alpha \Psi^2 \Delta q - n(\alpha - 1) \nabla f \nabla q \Psi^2 \right).
 \end{aligned}$$

Let us observe that

$$(y - z)^2 = \frac{1}{\alpha^2} (y - \alpha z)^2 + \frac{(\alpha - 1)^2}{\alpha^2} y^2 + \frac{2(\alpha - 1)}{\alpha^2} y(y - \alpha z)$$

and substitute this into the previous estimate. Regrouping the terms and applying the inequality  $mv^2 - nv \geq -n^2/4m$ , which is valid for  $m, n > 0$  and  $v \in \mathbb{R}$ , we get

$$\begin{aligned}
 & -2t_1 F \frac{C_{1/2} \Psi^{3/2}}{\rho} |\nabla f| + \frac{2t_1^2}{n} \left( a\alpha \Psi^2 (|\nabla f|^2 - f_t - q)^2 - nk_1 \alpha \Psi^2 |\nabla f|^2 \right. \\
 & \quad \left. - \frac{n^2 \alpha}{4b} \bar{k}^2 \Psi^2 - \frac{n}{2} \alpha \Psi^2 \Delta q - n(\alpha - 1) \nabla f \nabla q \Psi^2 \right) \\
 & \geq \frac{2t_1^2}{n} \left( \frac{a}{\alpha} (y - \alpha z)^2 - 2nk_1 \alpha y + nk_1 \alpha y - \frac{n^2 \alpha}{4b} \bar{k}^2 \Psi^2 - \frac{n C_{1/2}}{\rho} y^{1/2} (y - \alpha z) \right. \\
 & \quad \left. - \frac{n}{2} \alpha \Psi^2 \theta - n(\alpha - 1) \gamma \Psi^{1/2} y^{1/2} + \frac{a}{\alpha} (\alpha - 1)^2 y^2 + \frac{2a(\alpha - 1)}{\alpha} y(y - \alpha z) \right) \\
 & \geq \frac{2t_1^2}{n} \left( \frac{a}{\alpha} (y - \alpha z)^2 - \frac{n^2 \alpha}{4b} \bar{k}^2 \Psi^2 \right. \\
 & \quad \left. - \frac{n}{2} \alpha \Psi^2 \theta - \frac{n^2 d_2 \alpha}{8a\rho^2(\alpha - 1)} (y - \alpha z) - \frac{n^2 k_1^2 \alpha^3}{a(\alpha - 1)^2} - \frac{\gamma^2 n(\alpha - 1)^2 \Psi}{4k_1 \alpha} \right).
 \end{aligned}$$

Because  $t(y - \alpha z) = \Psi F$  by definition, (8) now implies that

$$\begin{aligned}
 0 & \geq \frac{2a}{n\alpha} (\Psi F)^2 + \left( -\frac{nd_2 t_1}{\rho^2} \left( \frac{\alpha}{a(\alpha - 1)} + 1 + \rho \sqrt{k_1} + \frac{\rho^2}{\tau} + \rho^2 \bar{k} \right) (\Psi F) \right. \\
 & \quad \left. - \Psi F - \frac{2nk_1^2 \alpha^3}{a(\alpha - 1)^2} t_1^2 - \frac{\alpha n}{2b} t_1^2 \bar{k}^2 \Psi^2 - \frac{\gamma^2 (\alpha - 1)^2 t_1^2 \Psi}{2k_1 \alpha} - \alpha t_1^2 \Psi^2 \theta \right) \\
 & \geq \frac{2a}{n\alpha} (\Psi F)^2 + \left( -\frac{d_3 t_1}{\rho^2} \left( \frac{\alpha}{a(\alpha - 1)} + \frac{\rho^2}{\tau} + \rho^2 \bar{k} \right) - 1 \right) (\Psi F) - \frac{2nk_1^2 \alpha^3}{a(\alpha - 1)^2} t_1^2 \\
 & \quad - \frac{\alpha n}{2b} t_1^2 \bar{k}^2 \Psi^2 - \frac{\gamma^2 (\alpha - 1)^2 t_1^2 \Psi}{2k_1 \alpha} - \alpha t_1^2 \Psi^2 \theta
 \end{aligned}$$



at  $(x_1, t_1)$  with  $d_3 = 4nd_2$ . The expression in the last two lines is a polynomial in  $\Psi F$  of degree 2. Consequently, in accordance with the quadratic formula,

$$\begin{aligned} \Psi F \leq \frac{n\alpha}{2a} \left[ \frac{d_3 t_1}{\rho^2} \left( \frac{\alpha}{a(\alpha-1)} + \frac{\rho^2}{\tau} + \rho^2 \bar{k} \right) + 1 + \frac{2k_1 \alpha}{\alpha-1} t_1 \right. \\ \left. + \sqrt{\frac{a}{b}} t_1 \bar{k} \Psi + \frac{\gamma(\alpha-1)t_1}{\alpha} \sqrt{\frac{a}{k_1 n}} + t_1 \sqrt{\frac{2a\theta}{n}} \Psi \right] \end{aligned}$$

at  $(x_1, t_1)$ . We will now use this conclusion to obtain a bound on  $F(x, \tau)$  for an appropriate range of  $x \in M$ .

Recall that  $\Psi(x, \tau) = 1$  whenever  $\text{dist}(x, x_0, t) < \rho/2$ . Also,  $(x_1, t_1)$  is a maximum point for  $\Psi F$  in the set  $\{(x, t) \in M \times [0, \tau] \mid \text{dist}(x, x_0, \tau) < \rho\}$ . Hence,

$$\begin{aligned} F(x, \tau) = (\Psi F)(x, \tau) &\leq (\Psi F)(x_1, t_1) \\ &\leq \frac{n\alpha d_3 \tau}{2a\rho^2} \left( \frac{\alpha}{a(\alpha-1)} + \frac{\rho^2}{\tau} + \rho^2 \bar{k} \right) + \frac{n\alpha}{2a} + \frac{nk_1 \alpha^2}{a(\alpha-1)} \tau \\ &\quad + \frac{\alpha \tau n \bar{k}}{2} \sqrt{\frac{1}{ab}} + \frac{\gamma(\alpha-1)\tau}{\alpha} \sqrt{\frac{k_1}{an}} + \alpha \tau \sqrt{\frac{2n\theta}{a}} \end{aligned}$$

for all  $x \in M$ , such that  $\text{dist}(x, x_0, \tau) < \rho/2$ . Since  $\tau \in (0, T]$  is chosen arbitrarily, this formula implies that

$$\begin{aligned} (|\nabla f|^2 - \alpha f_t - \alpha q)(x, t) \\ \leq \frac{\alpha d_4}{a\rho^2} \left( \frac{\alpha}{a(\alpha-1)} + \frac{\rho^2}{t} + \rho^2 \bar{k} \right) + \frac{nk_1 \alpha^2}{a(\alpha-1)} + \frac{\alpha n \bar{k}}{2} \sqrt{\frac{1}{ab}} + \frac{\gamma(\alpha-1)}{\alpha} \sqrt{\frac{k_1}{an}} + \alpha \sqrt{\frac{2n\theta}{a}}, \end{aligned}$$

$(x, t) \in B_{\rho/2, T}$ , with  $d_4 = \max\{nd_3, n\}$ , as long as  $t \neq 0$ . If we set  $a = 1/(2\alpha)$ , note that  $b = 1/\alpha - a$ , and define the constant  $C'$  appropriately, estimate (5) will follow by a straightforward computation.  $\square$

Now we consider the case where the manifold  $M$  is compact. Assume  $M$  has nonnegative Ricci curvature, we will deduce a global estimate on  $u(x, t)$ . A related inequality for (1)–(2) can be found in [Hamilton 1993].

**Theorem 2.4.** *Suppose the manifold  $M$  is a solution to the Ricci flow (1). Assume that  $0 \leq \text{Ric}(x, t) \leq kg(x, t)$  for some  $k > 0$  and all  $(x, t) \in M \times [0, T]$ . Suppose  $u : M \times [0, T] \rightarrow \mathbb{R}$  is a smooth positive function satisfying the heat equation (2), and  $q(x, t)$  is a  $C^2$  function defined on  $M \times (0, T)$  and  $|\Delta q| \leq \theta$ . The estimate*

$$(10) \quad \frac{|\nabla u|^2}{u^2} - \frac{u_t}{u} - q \leq kn + \frac{n}{2t} + 2\sqrt{n\theta}$$

holds for all  $(x, t) \in M \times [0, T]$ .

*Proof.* As before, we write  $f$  instead of  $\log u$ . It will be convenient for us to denote  $F_1 = t(|\nabla f|^2 - f_t - q)$ . Fix  $\tau \in (0, T]$ , and choose a point  $(x_0, t_0) \in M \times [0, \tau]$  where  $F_1$  attains its maximum on  $M \times [0, \tau]$ . Our first step is to show that

$$F_1(x_0, t_0) \leq t_0 kn + \frac{n}{2} + 2\sqrt{n\theta}t_0.$$

Then the theorem will follow.

If  $t_0 = 0$ , then  $F_1(x, t_0)$  is equal to 0 for every  $x \in M$ , and estimate (10) becomes evident. Therefore, we can assume  $t_0 > 0$  without loss of generality. By Lemma 2.2 and assumptions on the Ricci curvature of  $M$ , we can deduce that

$$\left(\Delta - \frac{\partial}{\partial t}\right)F_1 \geq \frac{2a}{n} \frac{F^2}{t_0} - \frac{F}{t_0} - 2\nabla f \nabla F - \frac{nt_0 k^2}{2(1-a)} - t_0 \Delta q$$

for all  $a \in (0, 1)$  at the point  $(x_0, t_0)$ . Recall that  $F_1$  attains its maximum at  $(x_0, t_0)$ . This tells us that

$$\Delta F_1(x_0, t_0) \leq 0, \quad \frac{\partial}{\partial t} F_1(x_0, t_0) \geq 0, \quad \text{and } \nabla F_1(x_0, t_0) = 0.$$

In consequence, the estimate

$$0 \geq \frac{2a}{n} \frac{F^2}{t_0} - \frac{F}{t_0} - \frac{nt_0 k^2}{2(1-a)} - t_0 \theta$$

holds at  $(x_0, t_0)$ , and the quadratic formula yields

$$F_1 \leq \frac{n}{4a} \left( 1 + \sqrt{1 + \frac{4at_0^2}{1-a} k^2 + \frac{8a\theta}{n} t_0^2} \right).$$

Letting  $a = (1 + kt_0)/(1 + 2kt_0)$ , and substituting this into the above inequality, we arrive at (10).

We now need only a simple argument to complete the proof. Since  $(x_0, t_0)$  is the maximum point for  $F_1$  on  $M \times [0, \tau]$ , we are then able to conclude that

$$F_1(x, \tau) \leq F_1(x_0, t_0) \leq t_0 kn + \frac{n}{2} + 2\sqrt{n\theta}t_0 \leq \tau kn + \frac{n}{2} + 2\sqrt{n\theta}\tau$$

for all  $x \in M$ . Therefore, the estimate

$$\frac{|\nabla u|^2}{u^2} - \frac{u_t}{u} - q \leq kn + \frac{n}{2\tau} + 2\sqrt{n\theta}$$

holds at  $(x, \tau)$ . Because  $\tau \in (0, T]$  can be chosen arbitrarily, the assertion of the theorem follows.  $\square$

### 3. Harnack Inequalities

The last goal is to get two Harnack inequalities. These may be viewed as applications of Theorems 2.3 and 2.4.

**Lemma 3.1.** *Let  $(M, g(x, t))_{t \in [0, T]}$  be a complete solution to the Ricci flow (1). Suppose  $u : M \times [0, T] \rightarrow \mathbb{R}$  is a smooth positive function satisfying the heat equation (2), and  $q(x, t)$  is a  $C^2$  function defined on  $M \times [0, T]$ . Define  $f = \log u$ , and assume that*

$$\frac{\partial f}{\partial t} \geq \frac{1}{A_1} \left( |\nabla f|^2 - A_2 - \frac{A_3}{t} \right) - q, \quad x \in M, t \in (0, T]$$

for some  $A_1, A_2, A_3 > 0$ . Then the inequality

$$u(x_2, t_2) \geq u(x_1, t_1) \left( \frac{t_2}{t_1} \right)^{-A_3/A_1} \exp \left( -\frac{A_1}{4} \Gamma(x_1, t_1, x_2, t_2) - \frac{A_2}{A_1} (t_2 - t_1) \right)$$

holds for all  $(x_1, t_1) \in M \times (0, T)$ , and  $(x_2, t_2) \in M \times (0, T)$  such that  $t_1 < t_2$ , where

$$\Gamma(x_1, t_1, x_2, t_2) = \inf \int_{t_1}^{t_2} \left[ \left| \frac{d}{dt} \gamma(t) \right|^2 + \frac{4}{A_1} q \right] dt,$$

and the infimum is taken over the set  $\Theta(x_1, t_1, x_2, t_2)$  of all the smooth paths  $\gamma : [t_1, t_2] \rightarrow M$  that connect  $x_1$  to  $x_2$ . We remind the reader that the norm  $|\cdot|$  depends on  $t$ .

*Proof.* Consider a path  $\gamma \in \Theta(x_1, t_1, x_2, t_2)$ . We begin by computing

$$\begin{aligned} \frac{d}{dt} f(\gamma(t), t) &= \nabla f(\gamma(t), t) \frac{d}{dt} \gamma(t) + \frac{\partial}{\partial s} f((\gamma(t), s)) \Big|_{s=t} \\ &\geq -|\nabla f(\gamma(t), t)| \left| \frac{d}{dt} \gamma(t) \right| + \frac{1}{A_1} \left( |\nabla f|^2 - A_2 - \frac{A_3}{t} \right) - q \\ &\geq -\frac{A_1}{4} \left| \frac{d}{dt} \gamma(t) \right|^2 + \frac{1}{A_1} \left( -A_2 - \frac{A_3}{t} \right) - q, \end{aligned}$$

$t \in [t_1, t_2]$ .

By the inequality  $mv^2 - nv \geq -n^2/(4m)$ , valid for  $m, n > 0$  and  $v \in \mathbb{R}$ , we get the last step. It then follows that

$$\begin{aligned} f(x_2, t_2) - f(x_1, t_1) &= \int_{t_1}^{t_2} \frac{d}{dt} f(\gamma(t), t) dt \\ &\geq -\frac{A_1}{4} \int_{t_1}^{t_2} \left| \frac{d}{dt} \gamma(t) \right|^2 - \frac{A_2}{A_1} (t_2 - t_1) - \frac{A_3}{A_1} \ln \frac{t_2}{t_1} - \int_{t_1}^{t_2} q dt. \end{aligned}$$

The assertion of the lemma follows by exponentiating.  $\square$

We are ready to formulate our Harnack inequalities for (1)–(2). The first one applies on noncompact manifolds. The second one does not, but provides a more explicit estimate.

**Theorem 3.2.** *Let  $(M, g(x, t))_{t \in [0, T]}$  be a complete solution to the Ricci flow (1). Assume that  $-k_1 g(x, t) \leq \text{Ric}(x, t) \leq k_2 g(x, t)$  for some  $k_1, k_2 > 0$  and all  $(x, t) \in M \times [0, T]$ . Suppose  $u : M \times [0, T] \rightarrow \mathbb{R}$  is a smooth positive function solving the heat equation (2), and  $q(x, t)$  is a  $C^2$  function defined on  $M \times [0, T]$ ,  $|\nabla q| \leq \gamma$ , and  $|\Delta q| \leq \theta$ . Given  $\alpha > 1$ , the estimate*

$$(11) \quad u(x_2, t_2) \geq u(x_1, t_1) \left( \frac{t_2}{t_1} \right)^{-C'\alpha} \exp \left( -\frac{\alpha}{4} \Gamma(x_1, t_1, x_2, t_2) \right. \\ \left. - \frac{1}{\alpha} \left( C' \alpha^2 \bar{k} + \frac{nk_1 \alpha^3}{\alpha - 1} + \frac{\gamma(\alpha - 1)}{\alpha} \sqrt{\frac{2\alpha k_1}{n}} + 2\alpha \sqrt{\alpha n \theta} \right) (t_2 - t_1) \right)$$

holds for all  $(x_1, t_1) \in M \times [0, T]$ , and  $(x_2, t_2) \in M \times [0, T]$ , such that  $t_1 < t_2$ . The constant  $C'$  comes from Theorem 2.3, where

$$\Gamma(x_1, t_1, x_2, t_2) = \inf \int_{t_1}^{t_2} \left[ \left| \frac{d}{dt} \gamma(t) \right|^2 + \frac{4}{A_1} q \right] dt,$$

and the infimum is taken over the set  $\Theta(x_1, t_1, x_2, t_2)$  of all the smooth paths  $\gamma : [t_1, t_2] \rightarrow M$  that connect  $x_1$  to  $x_2$ .

*Proof.* Letting  $\rho$  go to infinity in (5), we conclude that

$$\frac{u_t}{u} \geq \frac{1}{\alpha} \left( \frac{|\nabla u|^2}{u^2} - C' \alpha^2 \left( \frac{1}{t} + \bar{k} \right) - \frac{nk_1 \alpha^3}{\alpha - 1} - \frac{\gamma(\alpha - 1)}{\alpha} \sqrt{\frac{2\alpha k_1}{n}} - 2\alpha \sqrt{\alpha n \theta} \right) - q$$

on  $M \times (0, T]$ . The desired assertion is now a consequence of Lemma 3.1.  $\square$

**Theorem 3.3.** *Suppose the manifold  $M$  is a solution to the Ricci flow (1). Assume that  $0 \leq \text{Ric}(x, t) \leq k g(x, t)$  for some  $k > 0$  and all  $(x, t) \in M \times [0, T]$ . Suppose  $u : M \times [0, T] \rightarrow \mathbb{R}$  is a smooth positive function satisfying the heat equation (2), and  $q(x, t)$  is a  $C^2$  function defined on  $M \times [0, T]$ ,  $|\Delta q| \leq \theta$ . The estimate*

$$u(x_2, t_2) \geq u(x_1, t_1) \left( \frac{t_2}{t_1} \right)^{-n/2} \exp \left( -\frac{\alpha}{4} \Gamma(x_1, t_1, x_2, t_2) - (kn + 2\sqrt{n\theta})(t_2 - t_1) \right)$$

holds for all  $(x_1, t_1) \in M \times [0, T]$  and  $(x_2, t_2) \in M \times (0, T)$  as long as  $t_1 < t_2$ .

*Proof.* Theorem 2.4 implies that

$$\frac{|\nabla u|^2}{u^2} - q - \left( kn + \frac{n}{2t} + 2\sqrt{n\theta} \right) \leq \frac{u_t}{u}, \quad x \in M, t \in (0, T].$$

We now use Lemma 3.1 to complete the proof.  $\square$

## References

- [Bailesteanu et al. 2010] M. Bailesteanu, X. Cao, and A. Pulemotov, “Gradient estimates for the heat equation under the Ricci flow”, *J. Funct. Anal.* **258**:10 (2010), 3517–3542. [MR 2011b:53153](#) [Zbl 1193.53139](#)
- [Hamilton 1993] R. S. Hamilton, “The Harnack estimate for the Ricci flow”, *J. Differential Geom.* **37**:1 (1993), 225–243. [MR 93k:58052](#) [Zbl 0804.53023](#)
- [Li and Yau 1986] P. Li and S.-T. Yau, “On the parabolic kernel of the Schrödinger operator”, *Acta Math.* **156**:3-4 (1986), 153–201. [MR 87f:58156](#) [Zbl 0611.58045](#)

Received August 16, 2010. Revised November 30, 2010.

XIAORUI ZHU  
PUBLIC SECURITY MARINE POLICE ACADEMY  
NINGBO 315801  
CHINA

and

CENTER OF MATHEMATICAL SCIENCES  
ZHEJIANG UNIVERSITY  
HANGZHOU 310027  
CHINA

[zjuzxr@gmail.com](mailto:zjuzxr@gmail.com)

# PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)

Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pacific@math.ucla.edu](mailto:pacific@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Darren Long  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[long@math.ucsb.edu](mailto:long@math.ucsb.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Alexander Merkurjev  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[merkurev@math.ucla.edu](mailto:merkurev@math.ucla.edu)

Jonathan Rogawski  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[jonr@math.ucla.edu](mailto:jonr@math.ucla.edu)

## PRODUCTION

[pacific@math.berkeley.edu](mailto:pacific@math.berkeley.edu)

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [www.pjmath.org](http://www.pjmath.org) for submission instructions.

---

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from [Periodicals Service Company](#), 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFlow™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2011 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 252      No. 1      July 2011

---

Some dynamic Wirtinger-type inequalities and their applications	1
RAVI P. AGARWAL, MARTIN BOHNER, DONAL O'REGAN and SAMIR H. SAKER	
Splitting criteria for vector bundles on higher-dimensional varieties	19
PARSA BAKHTARY	
Average Mahler's measure and $L_p$ norms of unimodular polynomials	31
KWOK-KWONG STEPHEN CHOI and MICHAEL J. MOSSINGHOFF	
Tate resolutions and Weyman complexes	51
DAVID A. COX and EVGENY MATEROV	
On pointed Hopf algebras over dihedral groups	69
FERNANDO FANTINO and GASTON ANDRÉS GARCIA	
Integral topological quantum field theory for a one-holed torus	93
PATRICK M. GILMER and GREGOR MASBAUM	
Knot 4-genus and the rank of classes in $W(\mathbb{Q}(t))$	113
CHARLES LIVINGSTON	
Roots of Toeplitz operators on the Bergman space	127
ISSAM LOUHICHI and NAGISETTY V. RAO	
Uniqueness of the foliation of constant mean curvature spheres in asymptotically flat 3-manifolds	145
SHIGUANG MA	
On the multiplicity of non-iterated periodic billiard trajectories	181
MARCO MAZZUCHELLI	
A remark on Einstein warped products	207
MICHELE RIMOLDI	
Exceptional Dehn surgery on large arborescent knots	219
YING-QING WU	
Harnack estimates for the linear heat equation under the Ricci flow	245
XIAORUI ZHU	