

*Pacific
Journal of
Mathematics*

WHITNEY UMBRELLAS AND SWALLOWTAILS

TAKASHI NISHIMURA

Volume 252 No. 2

August 2011

WHITNEY UMBRELLAS AND SWALLOWTAILS

TAKASHI NISHIMURA

Dedicated to Professor Shyuichi Izumiya on the occasion of his sixtieth birthday

We introduce map germs of pedal unfolding type and the notion of normalized Legendrian map germs. We show that the fundamental theorem of calculus provides a natural one-to-one correspondence between Whitney umbrellas of pedal unfolding type and normalized swallowtails.

1. Introduction

The map germ

$$(1) \quad f(x, y) = (xy, x^2, y)$$

is known as the normal form of Whitney umbrella, after Whitney's pioneering works [1943; 1944]. Compose the germ (1) with the coordinate transformations

$$h_s(x, y) = (x, x^2 + y) \quad \text{and} \quad h_t(X, Y, Z) = (X, -Z, -Y + Z),$$

where (X, Y, Z) are the standard coordinates of the target space \mathbb{R}^3 . This leads to the map germ

$$(2) \quad g(x, y) = h_t \circ f \circ h_s(x, y) = (x^3 + xy, -x^2 - y, y).$$

Set

$$(3) \quad G(x, y) = \left(\int_0^x (x^3 + xy) dx, \int_0^x (-x^2 - y) dx, y \right) \\ = \left(\frac{1}{4}x^4 + \frac{1}{2}x^2y, -\frac{1}{3}x^3 - xy, y \right).$$

Compose the map germ (3) with the scaling transformations

$$H_s(x, y) = \left(x, \frac{1}{6}y\right) \quad \text{and} \quad H_t(X, Y, Z) = (12X, 12Y, 6Z)$$

to obtain the map germ

$$(4) \quad H_t \circ G \circ H_s(x, y) = (3x^4 + x^2y, -4x^3 - 2xy, y),$$

MSC2010: 53A05, 57R45, 58K25.

Keywords: Whitney umbrella, pedal unfolding type, normalized swallowtail, swallowtail.

known as the normal form of the swallowtail [Bruce and Giblin 1992, page 129].

Two C^∞ map germs $\varphi, \psi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ are said to be \mathcal{A} -equivalent if there exist germs of C^∞ diffeomorphisms

$$h_s : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0) \quad \text{and} \quad h_t : (\mathbb{R}^3, 0) \rightarrow (\mathbb{R}^3, 0),$$

such that $\psi = h_t \circ \varphi \circ h_s$. A C^∞ map germ $\varphi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ is called a *Whitney umbrella* if it is \mathcal{A} -equivalent to (1); it is called a *swallowtail* if it is \mathcal{A} -equivalent to (4). As seen above, the Whitney umbrella (1) produces the swallowtail (4) via (2) and (3). By the converse procedure, the swallowtail (4) produces the Whitney umbrella (1).

It is impossible to produce a swallowtail by integrating (1) directly. This is because the discriminant set of (4) is not diffeomorphic to the discriminant set of

$$(5) \quad (x, y) \mapsto \left(\int_0^x xy dx, \int_0^x x^2 dx, y \right).$$

Note that the form (2) may be written as follows:

$$g(x, y) = (x(x^2 + y), -(x^2 + y), y) = (b(-x, -(x^2 + y)), y),$$

where $b(X, Y) = (XY, Y)$ (b stands for “blowdown”).

Definition 1.1. (i) A C^∞ map germ $\varphi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ having the following form is said to be of *pedal unfolding type*.

$$(6) \quad \varphi(x, y) = (n(x, y)p(x, y), p(x, y), y) = (b(n(x, y), p(x, y)), y),$$

where $n : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0)$ is a C^∞ function germ, such that

$$\frac{\partial n}{\partial x}(0, 0) \neq 0 \quad \text{and} \quad p : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0) \text{ is a } C^\infty \text{ function germ.}$$

(ii) For a C^∞ map germ of pedal unfolding type

$$\varphi(x, y) = (n(x, y)p(x, y), p(x, y), y),$$

set

$$\mathcal{I}(\varphi)(x, y) = \left(\int_0^x n(x, y)p(x, y) dx, \int_0^x p(x, y) dx, y \right).$$

The map germ $\mathcal{I}(\varphi) : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ is called the *integration of φ* .

(iii) A C^∞ map germ $\Phi : (\mathbb{R}^m, 0) \rightarrow (\mathbb{R}^{m+1}, 0)$ is called a *Legendrian map germ* if there exists a germ of C^∞ vector field $\nu_\Phi : (\mathbb{R}^m, 0) \rightarrow T_1\mathbb{R}^{m+1}$ along Φ such that

$$\frac{\partial \Phi}{\partial x_1}(x_1, \dots, x_m) \cdot \nu_\Phi(x_1, \dots, x_m) = \dots = \frac{\partial \Phi}{\partial x_m}(x_1, \dots, x_m) \cdot \nu_\Phi(x_1, \dots, x_m) = 0$$

and the map germ $L_\Phi : (\mathbb{R}^m, 0) \rightarrow T_1\mathbb{R}^{m+1}$ defined by

$$L_\Phi(x_1, \dots, x_m) = (\Phi(x_1, \dots, x_m), \nu_\Phi(x_1, \dots, x_m))$$

is nonsingular. L_Φ is called a *Legendrian lift of Φ* . (Here the dot stands for the scalar product of two vectors of $T_{\Phi(x,y)}\mathbb{R}^{m+1}$, and $T_1\mathbb{R}^{m+1}$ is the unit tangent bundle of \mathbb{R}^{m+1} .) The C^∞ vector field ν_Φ is called a *unit normal vector field* of Φ .

(iv) A Legendrian map germ $\Phi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ is said to be *normalized* if it has the form

$$(7) \quad \Phi(x, y) = (\Phi_1(x, y), \Phi_2(x, y), y)$$

with

$$(8) \quad \frac{\partial \Phi_2}{\partial x}(0, 0) = 0$$

and if, furthermore,

$$(9) \quad \nu_\Phi(0, 0) = \frac{\partial}{\partial X} \quad \text{or} \quad \nu_\Phi(0, 0) = -\frac{\partial}{\partial X}.$$

(v) For a normalized Legendrian map germ $\Phi(x, y) = (\Phi_1(x, y), \Phi_2(x, y), y)$, set

$$\mathcal{D}(\Phi)(x, y) = \left(\frac{\partial \Phi_1}{\partial x}(x, y), \frac{\partial \Phi_2}{\partial x}(x, y), y \right).$$

The map germ $\mathcal{D}(\Phi) : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ is called the *differential* of Φ .

We showed in [Nishimura 2010] that any germ $\varphi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ of a one-parameter pedal unfolding of a spherical pedal curve has the form (6). Hence, a map germ φ having the form (6) is said to be of *pedal unfolding type*. As shown in [Nishimura 2010], not only nonsingular map germs, but also Whitney umbrellas may be realized as germs of one-parameter pedal unfoldings of spherical pedal curves. For more information on Legendrian map germs, see [Arnold et al. 1985; Izumiya 1987; Zakalyukin 1976; 1983]. Note that both (3) and (10) are normalized Legendrian map germs.

Proposition 1.2. (i) *If $\varphi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ is a C^∞ map germ of pedal unfolding type, $\mathcal{I}(\varphi)$ is a normalized Legendrian map germ.*

(ii) *If $\Phi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ is a normalized Legendrian map germ, $\mathcal{D}(\Phi)$ is a map germ of pedal unfolding type.*

Set

$$\mathcal{W} = \{\varphi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0) \text{ Whitney umbrella of pedal unfolding type}\},$$

$$\mathcal{S} = \{\Phi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0) \text{ normalized swallowtail}\}.$$

The main purpose of this paper is to show the following:

- Theorem 1.3.** (i) *The map $\mathcal{I} : \mathcal{W} \rightarrow \mathcal{S}$ defined by $\mathcal{W} \ni \varphi \mapsto \mathcal{I}(\varphi) \in \mathcal{S}$ is well-defined and bijective.*
 (ii) *The map $\mathcal{D} : \mathcal{S} \rightarrow \mathcal{W}$ defined by $\mathcal{S} \ni \Phi \mapsto \mathcal{D}(\Phi) \in \mathcal{W}$ is well-defined and bijective.*

Incidentally, we show Theorem 1.4. A C^∞ map germ $\Phi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ is called a *cuspidal edge* if Φ is \mathcal{A} -equivalent to the following:

$$(10) \quad (x, y) \mapsto \left(\frac{1}{3}x^3, \frac{1}{2}x^2, y\right).$$

Set

- $\mathcal{N} = \{\varphi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0) \text{ nonsingular map germ of pedal unfolding type}\},$
 $\mathcal{C} = \{\Phi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0) \text{ normalized cuspidal edge}\}.$

- Theorem 1.4.** (i) *The map $\mathcal{I} : \mathcal{N} \rightarrow \mathcal{C}$ defined by $\mathcal{N} \ni \varphi \mapsto \mathcal{I}(\varphi) \in \mathcal{C}$ is well-defined and bijective.*
 (ii) *The map $\mathcal{D} : \mathcal{C} \rightarrow \mathcal{N}$ defined by $\mathcal{C} \ni \Phi \mapsto \mathcal{D}(\Phi) \in \mathcal{N}$ is well-defined and bijective.*

Any stable map germ $(\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ is either a Whitney umbrella or nonsingular, and any Legendrian stable singularity $(\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ is either a cuspidal edge or a swallowtail (see [Arnold et al. 1985], for example). Theorems 1.3 and 1.4 can thus be regarded as a “fundamental theorem of calculus” for stable map germs $(\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ and Legendrian stable singularities $(\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$.

Based on Theorems 1.3 and 1.4, it is natural to ask:

- Question 1.5.** (i) Let $\varphi_1, \varphi_2 : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ be two C^∞ map germs of pedal unfolding type. Suppose that φ_1 is \mathcal{A} -equivalent to φ_2 . Is $\mathcal{I}(\varphi_1)$ necessarily \mathcal{A} -equivalent to $\mathcal{I}(\varphi_2)$?
 (ii) Let $\Phi_1, \Phi_2 : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ be two normalized Legendrian map germs. Suppose that Φ_1 is \mathcal{A} -equivalent to Φ_2 . Is $\mathcal{D}(\Phi_1)$ necessarily \mathcal{A} -equivalent to $\mathcal{D}(\Phi_2)$?

In Section 2, several preparations for the proofs of Theorems 1.3 and 1.4 and the proof of Proposition 1.2 are given. Theorems 1.3 and 1.4 are proved in Section 3 and Section 4 respectively.

2. Preliminaries

Function germs with two variables and map germs with two variables. Let \mathcal{E}_2 be the set of C^∞ function germs $(\mathbb{R}^2, 0) \rightarrow \mathbb{R}$, and let m_2 be the subset of \mathcal{E}_2

consisting of C^∞ function germs $(\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0)$. The set \mathcal{E}_2 has a natural \mathbb{R} -algebra structure. For a C^∞ map germ $\varphi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$, let $\varphi^* : \mathcal{E}_2 \rightarrow \mathcal{E}_2$ be the \mathbb{R} -algebra homomorphism defined by $\varphi^*(u) = u \circ \varphi$. Set $Q(\varphi) = \mathcal{E}_2 / \varphi^* m_2 \mathcal{E}_2$. Then, $Q(\varphi)$ is an \mathbb{R} -algebra. A special case of [Mather 1969, Theorem 2.1] follows.

Proposition 2.1. *Let $p : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0)$ be a C^∞ function germ.*

(i) *The \mathbb{R} -algebra $Q(p(x, y), y)$ is isomorphic to $Q(x^2, y)$ if and only if*

$$\frac{\partial p}{\partial x}(0, 0) = 0 \quad \text{and} \quad \frac{\partial^2 p}{\partial x^2}(0, 0) \neq 0.$$

(ii) *The \mathbb{R} -algebra $Q(p(x, y), y)$ is isomorphic to $Q(x, y)$ if and only if*

$$(x, y) \mapsto (p(x, y), y)$$

is a germ of C^∞ diffeomorphism.

Definition 2.2 [Mond 1985]. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation of the form $T(s, \lambda) = (-s, \lambda)$. Two C^∞ function germs $p_1, p_2 : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0)$ are said to be \mathcal{H}^T -equivalent if there exists a germ of C^∞ diffeomorphism

$$h : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$$

of the form $h \circ T = T \circ h$, and a C^∞ function germ $M : (\mathbb{R}^2, (0, 0)) \rightarrow \mathbb{R} - \{0\}$ of the form $M \circ T = M$, such that $p_1 \circ h(x, y) = M(x, y)p_2(x, y)$.

Theorem 2.3 [Mond 1985]. *Two C^∞ map germs $\varphi_i : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ ($i = 1, 2$) of the form*

$$\varphi_i(x, y) = (xp_i(x^2, y), x^2, y), \quad \text{where } p_i(x^2, y) \notin m_2^\infty \quad (i = 1, 2)$$

are \mathcal{A} -equivalent if and only if the function germs $p_i(x^2, y)$ are \mathcal{H}^T -equivalent. Here,

$$m_2^\infty = \left\{ q : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0) \mid \frac{\partial^{i+j} q}{\partial x^i \partial y^j}(0, 0) = 0 \text{ for all } i, j \in \{0\} \cup \mathbb{N} \right\}.$$

From this and the Malgrange preparation theorem [Arnold et al. 1985], we have:

Corollary 2.4. *Two C^∞ map germs $\varphi_i : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ ($i = 1, 2$) of the form*

$$\varphi_i(x, y) = (n_i(x, y)p_i(x^2, y), x^2, y),$$

where $p_i(x^2, y) \notin m_2^\infty$ and $(\partial n_i / \partial x)(0, 0) \neq 0$ for $i = 1, 2$, are \mathcal{A} -equivalent if and only if the function germs $p_i(x^2, y)$ are \mathcal{H}^T -equivalent.

Map germs of pedal unfolding type. Let $\varphi : I \times J \rightarrow \mathbb{R}^3$ be a representative of a given C^∞ map germ of pedal unfolding type, where I, J are sufficiently small intervals containing the origin of \mathbb{R} . We may put $\varphi(x, y) = (n(x, y)p(x, y), p(x, y))$. Set

$$\Phi(x, y) = (\Phi_1(x, y), \Phi_2(x, y), y) = \left(\int_0^x n(x, y)p(x, y)dx, \int_0^x p(x, y)dx, y \right)$$

and

$$\tilde{\mu}_\Phi(x, y) = \frac{\partial}{\partial X} - n(x, y) \frac{\partial}{\partial Y}.$$

Since $\tilde{\mu}_\Phi(x, y) \neq 0$ for any $x \in I$ and $y \in J$, for any fixed $y \in J$ we may define the map germ $L_{\Phi, y} : (\mathbb{R}, 0) \rightarrow T_1\mathbb{R}^2$ as

$$L_{\Phi, y}(x) = \left((\Phi_1(x, y), \Phi_2(x, y)), \frac{\tilde{\mu}_\Phi(x, y)}{\|\tilde{\mu}_\Phi(x, y)\|} \right),$$

where $T_1\mathbb{R}^2$ is the unit tangent bundle of \mathbb{R}^2 . Then, since φ is a representative of a map germ of pedal unfolding type, we have:

Lemma 2.5. *For any $y \in J$, $L_{\Phi, y} : (\mathbb{R}, 0) \rightarrow T_1\mathbb{R}^2$ is a Legendrian lift of the map germ $x \mapsto (\Phi_1(x, y), \Phi_2(x, y))$.*

This implies:

Lemma 2.6. *For any $y \in J$, the map germ $\tilde{\Phi}_y : (\mathbb{R}, 0) \rightarrow (\mathbb{R}^2, 0)$ defined by $\tilde{\Phi}_y(x) = (\Phi_1(x, y), \Phi_2(x, y))$ is a Legendrian map germ.*

Next, set

$$\tilde{v}_\Phi(x, y) = \tilde{\mu}_\Phi(x, y) - \left(\frac{\partial \Phi_1}{\partial y}(x, y) - n(x, y) \frac{\partial \Phi_2}{\partial y}(x, y) \right) \frac{\partial}{\partial Z}.$$

Lemma 2.7. *For any $x \in I$ and $y \in J$,*

$$\tilde{v}_\Phi(x, y) \cdot \frac{\partial \Phi}{\partial x}(x, y) = 0, \quad \tilde{v}_\Phi(x, y) \cdot \frac{\partial \Phi}{\partial y}(x, y) = 0.$$

Since $\tilde{v}_\Phi(x, y) \neq 0$ for any $x \in I$ and $y \in J$, we may define the map germ

$$L_\Phi : (\mathbb{R}^2, 0) \rightarrow T_1\mathbb{R}^3$$

as

$$L_\Phi(x, y) = \left(\Phi(x, y), \frac{\tilde{v}_\Phi(x, y)}{\|\tilde{v}_\Phi(x, y)\|} \right).$$

Then Lemma 2.7 implies successively:

Lemma 2.8. $L_\Phi : (\mathbb{R}^2, 0) \rightarrow T_1\mathbb{R}^3$ is a Legendrian lift of Φ .

Lemma 2.9. $\Phi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ is a Legendrian map germ.

Normalized Legendrian map germs. Let $\Phi : U \rightarrow \mathbb{R}^3$ be a representative of a given normalized Legendrian map germ $(\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$, where U is a sufficiently small neighborhood of the origin of \mathbb{R}^2 . We assume that the origin of \mathbb{R}^2 is a singular point of Φ . By condition (7) of the definition of normalized Legendrian map germs, we may assume that Φ has the form

$$\Phi(x, y) = (\Phi_1(x, y), \Phi_2(x, y), y).$$

Since Φ is a representative of a Legendrian map germ, we have the following:

Lemma 2.10. *There exists a C^∞ vector field v_Φ along Φ ,*

$$v_\Phi(x, y) = n_1(x, y) \frac{\partial}{\partial X} + n_2(x, y) \frac{\partial}{\partial Y} + n_3(x, y) \frac{\partial}{\partial Z},$$

such that

- (i) $n_1(x, y) \frac{\partial \Phi_1}{\partial x}(x, y) + n_2(x, y) \frac{\partial \Phi_2}{\partial x}(x, y) = 0;$
- (ii) $n_1(x, y) \frac{\partial \Phi_1}{\partial y}(x, y) + n_2(x, y) \frac{\partial \Phi_2}{\partial y}(x, y) + n_3(x, y) = 0;$
- (iii) *the map $L_\Phi : U \rightarrow T_1\mathbb{R}^3$ defined by $L_\Phi(x, y) = (\Phi(x, y), v_\Phi(x, y))$ is an immersion.*

Condition (9) in the definition of normalized Legendrian map germs gives:

Lemma 2.11. *For the vector field v_Φ , $n_1(0, 0) \neq 0$ and $n_2(0, 0) = n_3(0, 0) = 0$.*

By Lemma 2.10(i) and Lemma 2.11, we have the following equality of function germs:

$$(11) \quad \frac{\partial \Phi_1}{\partial x}(x, y) = -\frac{n_2(x, y)}{n_1(x, y)} \frac{\partial \Phi_2}{\partial x}(x, y).$$

This, together with condition (8) in the definition of normalized Legendrian maps, implies that

$$(12) \quad \mathfrak{D}(\Phi)(0, 0) = (0, 0, 0).$$

The next lemma is clear:

Lemma 2.12. *The function germs n and p given by*

$$n(x, y) = -\frac{n_2(x, y)}{n_1(x, y)} \quad \text{and} \quad p(x, y) = \frac{\partial \Phi_2}{\partial x}(x, y)$$

are of class C^∞ , and satisfy $\mathfrak{D}(\Phi)(x, y) = (n(x, y)p(x, y), p(x, y), y)$.

Lemma 2.13. *The function germ n satisfies $n(0, 0) = 0$ and $\frac{\partial n}{\partial x}(0, 0) \neq 0$.*

Proof. By Lemma 2.11, we have $n(0, 0) = 0$. Assume, for a contradiction, that $\partial n/\partial x$ vanishes at the origin; then so does $\partial n_2/\partial x$. At the same time, by differentiating both sides of the equality in Lemma 2.10(ii) with respect to x , we have

$$(13) \quad n_1(0, 0) \frac{\partial^2 \Phi_1}{\partial x \partial y}(0, 0) + \frac{\partial n_3}{\partial x}(0, 0) = 0.$$

Because Φ is a normalized Legendrian map germ such that the origin of \mathbb{R}^2 is a singular point of Φ , we obtain $(\partial n_3/\partial x)(0, 0) \neq 0$, which together with (13) gives

$$\frac{\partial^2 \Phi_1}{\partial x \partial y}(0, 0) \neq 0.$$

From (8), (11), and Lemma 2.11 we have a contradiction. □

Definition 2.14. Let $\Phi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ be a Legendrian map germ, and let ν_Φ be a unit normal vector field of Φ given in the definition of Legendrian map germs. The C^∞ function germ $LJ_\Phi : (\mathbb{R}^2, 0) \rightarrow \mathbb{R}$ defined by

$$LJ_\Phi(x, y) = \det\left(\frac{\partial \Phi}{\partial x}(x, y), \frac{\partial \Phi}{\partial y}(x, y), \nu_\Phi(x, y)\right)$$

is called the *Legendrian Jacobian* of Φ .

Note that if ν_Φ satisfies the conditions of unit normal vector field of Φ , then $-\nu_\Phi$ also satisfies them. Thus, the sign of $LJ_\Phi(x, y)$ depends on the particular choice of unit normal vector field ν_Φ . The Legendrian Jacobian of Φ is also called the *signed area density function* [Saji et al. 2009b]. Although it is reasonable to call LJ_Φ the area density function from the viewpoint of investigating the singular surface $\Phi(U)$ (U is a sufficiently small neighborhood of the origin of \mathbb{R}^2), it is also reasonable to call it the Legendrian Jacobian from the viewpoint of investigating the singular map germ Φ .

Let $\Phi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0)$ be a normalized Legendrian map germ and ν_Φ a unit normal vector field of Φ . Write

$$\begin{aligned} \Phi(x, y) &= (\Phi_1(x, y), \Phi_2(x, y), y), \\ \nu_\Phi(x, y) &= n_1(x, y) \frac{\partial}{\partial X} + n_2(x, y) \frac{\partial}{\partial Y} + n_3(x, y) \frac{\partial}{\partial Z}. \end{aligned}$$

By Lemma 2.11, we may set

$$\tilde{\nu}_\Phi(x, y) = \frac{\partial}{\partial X} + \frac{n_2(x, y)}{n_1(x, y)} \frac{\partial}{\partial Y} + \frac{n_3(x, y)}{n_1(x, y)} \frac{\partial}{\partial Z}.$$

We now give a formula for the Legendrian Jacobian. We start with the cross product (vector product)

$$\frac{\partial \Phi}{\partial x}(x, y) \times \frac{\partial \Phi}{\partial y}(x, y) = \frac{\partial \Phi_2}{\partial x}(x, y) \tilde{\nu}_\Phi(x, y).$$

This gives

$$(14) \quad LJ_{\Phi}(x, y) = \frac{(\partial\Phi_2/\partial x)(x, y)}{n_1(x, y)}.$$

Proof of Proposition 1.2. (i) Set

$$\mathcal{F}(\varphi) = \Phi(x, y) = (\Phi_1(x, y), \Phi_2(x, y), y).$$

By Lemma 2.9, Φ is a Legendrian map germ. Thus, it is sufficient to show that (8) and (9) in the definition of normalized Legendrian map germs are satisfied.

Set $\varphi(x, y) = (n(x, y)p(x, y), p(x, y), y)$. By the definition of map germs of pedal unfolding type, we have $n(0, 0) = 0$ and $p(0, 0) = 0$. It follows that

$$\frac{\partial\Phi_2}{\partial x}(0, 0) = p(0, 0) = 0.$$

Thus, condition (8) is satisfied. By Lemma 2.8, the germ L_{Φ} given by

$$L_{\Phi}(x, y) = \left(\Phi(x, y), \frac{\tilde{v}_{\Phi}(x, y)}{\|\tilde{v}_{\Phi}(x, y)\|} \right)$$

is a germ of Legendrian lift of Φ , where

$$\tilde{v}_{\Phi}(x, y) = \frac{\partial}{\partial X} - n(x, y) \frac{\partial}{\partial Y} - \left(\frac{\partial\Phi_1}{\partial y}(x, y) - n(x, y) \frac{\partial\Phi_2}{\partial y}(x, y) \right) \frac{\partial}{\partial Z}.$$

Since $n(0, 0) = 0$ and

$$\frac{\partial\Phi_1}{\partial y}(0, 0) = \int_0^0 \frac{\partial np}{\partial y}(x, 0) dx = 0,$$

we have

$$\frac{\tilde{v}_{\Phi}(0, 0)}{\|\tilde{v}_{\Phi}(0, 0)\|} = \frac{\partial}{\partial X}.$$

Thus, condition (9) is satisfied, proving part (i) of the proposition.

Proposition 1.2(ii) follows from (12), Lemma 2.12, and Lemma 2.13. □

3. Proof of Theorem 1.3

Suppose that both $\mathcal{F} : \mathcal{W} \rightarrow \mathcal{S}$ and $\mathcal{D} : \mathcal{S} \rightarrow \mathcal{W}$ are well-defined. By the fundamental theorem of calculus, we have $\mathcal{D} \circ \mathcal{F}(\varphi) = \varphi$ for all $\varphi \in \mathcal{W}$, and $\mathcal{F} \circ \mathcal{D}(\Phi) = \Phi$ for all $\Phi \in \mathcal{S}$. That is, both \mathcal{F} and \mathcal{D} are bijective. Therefore, in order to complete the proof, it is sufficient to show that both \mathcal{F} and \mathcal{D} are well-defined.

Proof that $\mathcal{F} : \mathcal{W} \rightarrow \mathcal{S}$ is well-defined. Let $\varphi(x, y) = (n(x, y)p_{\varphi}(x, y), p_{\varphi}(x, y), y)$ be an element of \mathcal{W} . Set $\Phi = \mathcal{F}(\varphi)$. Then Φ is a normalized Legendrian map germ

by Proposition 1.2. Let g be the Whitney umbrella of pedal unfolding type — see (2) in Section 1:

$$g(x, y) = (xp_g(x, y), p_g(x, y), y) = (x(x^2 + y), -x^2 - y, y).$$

Lemma 3.1. *There exists a germ of C^∞ diffeomorphism $h : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$ such that h has the form $h(x, y) = (h_1(x, y), h_2(y))$ and $p_\varphi \circ h(x, y)$ is $x^2 + y$ or $-(x^2 + y)$.*

Proof. Since φ is a Whitney umbrella of pedal unfolding type, we have

$$Q(p_\varphi(x, y), y) \cong Q(\varphi) \cong Q(g) \cong Q(x^2, y).$$

Thus, we may set $p_\varphi(x, 0) = a_2x^2 + o(x^2)$ ($a_2 \neq 0$) by Proposition 2.1. By the Morse lemma with parameters [Bruce and Giblin 1992], there exists a germ of C^∞ diffeomorphism $h : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$ such that h has the form

$$h(x, y) = (h_1(x, y), h_2(y)) \quad \text{and} \quad p_\varphi \circ h(x, y) = \pm(x^2 + q(y))$$

by a certain C^∞ function germ $q : (\mathbb{R}, 0) \rightarrow (\mathbb{R}, 0)$. Since φ is \mathcal{A} -equivalent to g , by Corollary 2.4, $\pm(x^2 + q(y))$ is \mathcal{H}^T -equivalent to p_g and thus $q : (\mathbb{R}, 0) \rightarrow (\mathbb{R}, 0)$ is a germ of C^∞ diffeomorphism. Lemma 3.1 follows. \square

Set $G = \mathcal{F}(g)$. Then, G has the form of (3) from Section 1, which is a normalized swallowtail. Since G is normalized, $\partial/\partial x$ is the null vector field for G defined in [Kokubu et al. 2005; Saji et al. 2009a], that is,

$$\frac{\partial G}{\partial x}(x, y) = 0$$

holds for any (x, y) which is a singular point of G . Since G is a swallowtail, we have, by [Saji et al. 2009a, Corollary 2.5],

$$LJ_G(0, 0) = \frac{\partial LJ_G}{\partial x}(0, 0) = 0, \quad \frac{\partial^2 LJ_G}{\partial x^2}(0, 0) \neq 0, \quad Q\left(LJ_G, \frac{\partial LJ_G}{\partial x}\right) \cong Q(x, y).$$

On the other hand, by (14) and Lemma 3.1, there exists a germ of C^∞ diffeomorphism $h : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$ and a C^∞ function germ $\xi : (\mathbb{R}^2, 0) \rightarrow \mathbb{R}$, such that h has the form $h(x, y) = (h_1(x, y), h_2(y))$, $\xi(0, 0) \neq 0$ and we have

$$LJ_\Phi \circ h(x, y) = \xi(x, y)LJ_G(x, y).$$

Because $\partial/\partial x$ is the null vector field for Φ (this is because Φ is normalized), LJ_Φ satisfies

$$LJ_\Phi(0, 0) = \frac{\partial LJ_\Phi}{\partial x}(0, 0) = 0, \quad \frac{\partial^2 LJ_\Phi}{\partial x^2}(0, 0) \neq 0, \quad Q\left(LJ_\Phi, \frac{\partial LJ_\Phi}{\partial x}\right) \cong Q(x, y).$$

Hence, Φ is a swallowtail by [Saji et al. 2009a, Corollary 2.5], and we have proved that $\mathcal{F} : \mathcal{W} \rightarrow \mathcal{S}$ is well-defined. \square

Proof that $\mathfrak{D} : \mathcal{S} \rightarrow \mathfrak{W}$ is well-defined.. Let Φ be an element of \mathcal{S} . Then, by Proposition 2.1, $\mathfrak{D}(\Phi)$ is of pedal unfolding type; we must show that it is a Whitney umbrella.

Lemma 3.2. *For the Legendrian Jacobian LJ_Φ , we have*

$$LJ_\Phi(0, 0) = \frac{\partial LJ_\Phi}{\partial x}(0, 0) = 0, \quad \frac{\partial^2 LJ_\Phi}{\partial x^2}(0, 0) \neq 0, \quad Q\left(LJ_\Phi, \frac{\partial LJ_\Phi}{\partial x}\right) \cong Q(x, y).$$

Proof. Since Φ is normalized, $\partial/\partial x$ is the null vector field. Since Φ is a swallowtail, Lemma 3.2 follows from [Saji et al. 2009a, Corollary 2.5]. \square

Since $\mathfrak{D}(\Phi)$ is of pedal unfolding type, there exists a C^∞ function germ $n : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0)$ such that

$$\frac{\partial n}{\partial x}(0, 0) \neq 0 \quad \text{and} \quad \frac{\partial \Phi_1}{\partial x}(x, y) = n(x, y) \frac{\partial \Phi_2}{\partial x}(x, y),$$

where $\Phi(x, y) = (\Phi_1(x, y), \Phi_2(x, y), y)$. Set $p_\varphi = \partial \Phi_2 / \partial x$. By (14), Lemma 3.2, and the Morse lemma with parameters, there is a germ of C^∞ diffeomorphism $h : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$ such that h has the form $h(x, y) = (h_1(x, y), h_2(y))$ and $p_\varphi \circ h(x, y) = \pm(x^2 + y)$. Then, by Corollary 2.4, $\mathfrak{D}(\Phi)$ is \mathcal{A} -equivalent to g . \square

4. Proof of Theorem 1.4

As with Theorem 1.3, it is sufficient to show that both $\mathcal{F} : \mathcal{N} \rightarrow \mathcal{C}$ and $\mathfrak{D} : \mathcal{C} \rightarrow \mathcal{N}$ are well-defined.

Proof that $\mathcal{F} : \mathcal{N} \rightarrow \mathcal{C}$ is well-defined. Let $\varphi(x, y) = (n(x, y)p_\varphi(x, y), p_\varphi(x, y), y)$ be an element of \mathcal{N} . Set $\Phi = \mathcal{F}(\varphi)$. Then, since φ is of pedal unfolding type, Φ is a normalized Legendrian map germ by Proposition 1.2. Let g be the nonsingular map germ of pedal unfolding type defined by $g(x, y) = (x^2, x, y)$.

Lemma 4.1. *There exists a germ of C^∞ diffeomorphism $h : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$ having the form $h(x, y) = (h_1(x, y), h_2(y))$ and such that $p_\varphi \circ h(x, y) = x$.*

Proof. Since φ is nonsingular and of pedal unfolding type, we have

$$Q(p_\varphi(x, y), y) \cong Q(\varphi) \cong Q(g) \cong Q(x, y).$$

Thus, $(p_\varphi(x, y), y)$ is a germ of C^∞ diffeomorphism by Proposition 2.1. From the form of $(p_\varphi(x, y), y)$, its inverse map germ $h : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$ has the form $h(x, y) = (h_1(x, y), h_2(y))$. Since h is the inverse map germ of $(p_\varphi(x, y), y)$, it follows that $p_\varphi \circ h(x, y) = x$. \square

Since Φ is normalized, $\partial/\partial x$ is the null vector field for Φ . By Lemma 4.1 and (14), we have

$$\frac{\partial LJ_\Phi}{\partial x}(0, 0) \neq 0.$$

Thus, the null vector field $\partial/\partial x$ is transverse to $\{(x, y) \mid LJ_\Phi(x, y) = 0\}$ at $(0, 0) \in \mathbb{R}^2$. Hence, Φ is a cuspidal edge by [Kokubu et al. 2005, Proposition 1.3], showing that $\mathcal{D} : \mathcal{N} \rightarrow \mathcal{C}$ is well-defined. \square

Proof that $\mathcal{D} : \mathcal{C} \rightarrow \mathcal{N}$ is well-defined. Let Φ be an element of \mathcal{C} . By Proposition 1.2, $\mathcal{D}(\Phi)$ is of pedal unfolding type.

Lemma 4.2. *The Legendrian Jacobian LJ_Φ satisfies*

$$LJ_\Phi(0, 0) = 0 \quad \text{and} \quad \frac{\partial LJ_\Phi}{\partial x}(0, 0) \neq 0.$$

Proof. Since $\partial/\partial x$ is the null vector field for Φ and Φ is a cuspidal edge, Lemma 4.2 follows from [Saji et al. 2009a, Corollary 2.5]. \square

Since $\mathcal{D}(\Phi)$ is of pedal unfolding type, there exists a C^∞ function germ $n : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0)$ such that

$$\frac{\partial n}{\partial x}(0, 0) \neq 0 \quad \text{and} \quad \frac{\partial \Phi_1}{\partial x}(x, y) = n(x, y) \frac{\partial \Phi_2}{\partial x}(x, y),$$

where $\Phi(x, y) = (\Phi_1(x, y), \Phi_2(x, y), y)$. Set $p_\varphi = \partial \Phi_2 / \partial x$. By Lemma 4.2 and (14), the map germ $(x, y) \mapsto (p_\varphi(x, y), y)$ is a germ of a C^∞ diffeomorphism. Thus, $\mathcal{D}(\Phi)$ is nonsingular. Since $\mathcal{D}(\Phi)(0, 0) = (0, 0, 0)$, we have proved that $\mathcal{D} : \mathcal{C} \rightarrow \mathcal{N}$ is well-defined. \square

Acknowledgement

The author thanks Kentaro Saji for valuable comments.

References

- [Arnold et al. 1985] V. I. Arnold, S. M. Gusein-Zade, and A. N. Varchenko, *Singularities of differentiable maps*, vol. 1, Monographs in Mathematics **82**, Birkhäuser, Boston, 1985. MR 86f:58018 Zbl 0554.58001
- [Bruce and Giblin 1992] J. W. Bruce and P. J. Giblin, *Curves and singularities*, 2nd ed., Cambridge University Press, 1992. MR 93k:58020 Zbl 0770.53002
- [Izumiya 1987] S. Izumiya, “On Legendrian singularities”, *Proc. Amer. Math. Soc.* **101**:4 (1987), 748–752. MR 89c:58016 Zbl 0639.58006
- [Kokubu et al. 2005] M. Kokubu, W. Rossman, K. Saji, M. Umehara, and K. Yamada, “Singularities of flat fronts in hyperbolic space”, *Pacific J. Math.* **221**:2 (2005), 303–351. MR 2006k:53102 Zbl 1110.53044
- [Mather 1969] J. N. Mather, “Stability of C^∞ mappings, IV: Classification of stable germs by R -algebras”, *Inst. Hautes Études Sci. Publ. Math.* **37** (1969), 223–248. MR 43 #1215b Zbl 0202.55102
- [Mond 1985] D. Mond, “On the classification of germs of maps from \mathbf{R}^2 to \mathbf{R}^3 ”, *Proc. London Math. Soc.* (3) **50**:2 (1985), 333–369. MR 86g:58021 Zbl 0557.58006
- [Nishimura 2010] T. Nishimura, “Singularities of one-parameter pedal unfoldings of spherical pedal curves”, *J. Singul.* **2** (2010), 160–169. MR 2763024

- [Saji et al. 2009a] K. Saji, M. Umehara, and K. Yamada, “ A_k singularities of wave fronts”, *Math. Proc. Cambridge Philos. Soc.* **146**:3 (2009), 731–746. MR 2010h:58060
- [Saji et al. 2009b] K. Saji, M. Umehara, and K. Yamada, “The geometry of fronts”, *Ann. of Math.* (2) **169**:2 (2009), 491–529. MR 2010e:58042 Zbl 1177.53014
- [Whitney 1943] H. Whitney, “The general type of singularity of a set of $2n - 1$ smooth functions of n variables”, *Duke Math. J.* **10** (1943), 161–172. MR 4,193b Zbl 0061.37207
- [Whitney 1944] H. Whitney, “The singularities of a smooth n -manifold in $(2n - 1)$ -space”, *Ann. of Math.* (2) **45** (1944), 247–293. MR 5,274a Zbl 0063.08238
- [Zakalyukin 1976] V. Zakalyukin, “Lagrangian and Legendrian singularities”, *Funktsional. Anal. i Prilozhen.* **10**:1 (1976), 26–36. In Russian; translated in *Funct. Anal. Appl.* **10**:1 (1976), 23–31. Zbl 0331.58007
- [Zakalyukin 1983] V. M. Zakalyukin, “Reconstructions of fronts and caustics depending on a parameter, and versality of mappings”, pp. 56–93 in *Современные проблемы математики*, vol. 22, edited by R. V. Gamkrelidze, Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1983. In Russian; translated in *J. Sov. Math.* **27** (1984), 2713–2735. MR 85h:58029 Zbl 0534.58014

Received July 30, 2011.

TAKASHI NISHIMURA
GROUP OF MATHEMATICAL SCIENCES
RESEARCH INSTITUTE OF ENVIRONMENT AND INFORMATION SCIENCES
YOKOHAMA NATIONAL UNIVERSITY
YOKOHAMA 240-8501
JAPAN
nishimura-takashi-yx@ynu.jp

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)

Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 252 No. 2 August 2011

Remarks on a Künneth formula for foliated de Rham cohomology	257
MÉLANIE BERTELSON	
K -groups of the quantum homogeneous space $SU_q(n)/SU_q(n-2)$	275
PARTHA SARATHI CHAKRABORTY and S. SUNDAR	
A class of irreducible integrable modules for the extended baby TKK algebra	293
XUEWU CHANG and SHAOBIN TAN	
Duality properties for quantum groups	313
SOPHIE CHEMLA	
Representations of the category of modules over pointed Hopf algebras over S_3 and S_4	343
AGUSTÍN GARCÍA IGLESIAS and MARTÍN MOMBELLI	
(p, p) -Galois representations attached to automorphic forms on GL_n	379
EKNATH GHATE and NARASIMHA KUMAR	
On intrinsically knotted or completely 3-linked graphs	407
RYO HANAKI, RYO NIKKUNI, KOUKI TANIYAMA and AKIKO YAMAZAKI	
Connection relations and expansions	427
MOURAD E. H. ISMAIL and MIZAN RAHMAN	
Characterizing almost Prüfer v -multiplication domains in pullbacks	447
QING LI	
Whitney umbrellas and swallowtails	459
TAKASHI NISHIMURA	
The Koszul property as a topological invariant and measure of singularities	473
HAL SADOFSKY and BRAD SHELTON	
A completely positive map associated with a positive map	487
ERLING STØRMER	
Classification of embedded projective manifolds swept out by rational homogeneous varieties of codimension one	493
KIWAMU WATANABE	
Note on the relations in the tautological ring of \mathcal{M}_g	499
SHENGMAO ZHU	



0030-8730(201108)252:2;1-9