

Pacific Journal of Mathematics

**EIGENVALUE ESTIMATES FOR HYPERSURFACES IN $\mathbb{H}^m \times \mathbb{R}$
AND APPLICATIONS**

PIERRE BÉRARD, PHILIPPE CASTILLON AND MARCOS CAVALCANTE

EIGENVALUE ESTIMATES FOR HYPERSURFACES IN $\mathbb{H}^m \times \mathbb{R}$ AND APPLICATIONS

PIERRE BÉRARD, PHILIPPE CASTILLON AND MARCOS CAVALCANTE

In the first part of this paper, we give a lower bound for the spectrum of the Laplacian on minimal hypersurfaces immersed into $\mathbb{H}^m \times \mathbb{R}$. As an application, in dimension 2, we prove that a complete minimal surface with finite total extrinsic curvature has finite index. In the second part, we consider the operator $L = \Delta_g + a + bK_g$ on a complete noncompact surface (M^2, g) . Assuming that L is nonnegative for some constants $a > 0$ and $b > 1/4$, we show that the infimum of the spectrum of M^2 is bounded from above by $a/(4b - 1)$. We apply this result to stable minimal surfaces immersed into homogeneous 3-manifolds.

1. Introduction

In this paper, we mainly consider orientable, minimal hypersurfaces immersed into $\mathbb{H}^m \times \mathbb{R}$. Let ν denote a unit normal field along M and let $v = \hat{g}(\nu, \partial_t)$ be the component of ν with respect to the unit vector field ∂_t tangent to the \mathbb{R} -direction in the ambient space.

In [Section 3](#), we give a lower bound of the spectrum of Δ_g which relies on the inequality $-\Delta_g b \geq (m - 2) + v^2$ satisfied by a “horizontal” Busemann function b (see [Theorem 3.1](#)). As an application, in [Section 4](#), we get a finiteness result for the index of a minimal surface in $\mathbb{H}^2 \times \mathbb{R}$ with finite total *extrinsic* curvature (answering a question raised in [[Bérard and Sá Earp 2008](#)]):

Theorem 1.1. *Let $(M^2, g) \looparrowright (\mathbb{H}^2 \times \mathbb{R}, \hat{g})$ be a complete, orientable, minimal surface with second fundamental form A . If $\int_M |A|^2 dv_g$ is finite, then the immersion has finite index.*

We also obtain a lower bound for the spectrum of the Laplacian on a complete minimal surface contained in a slab ([Proposition 4.3](#)).

Bérard was partially supported by the French cooperation programs ARCUS Rhône-Alpes–Brésil and Math-AmSud. Cavalcante gratefully acknowledges financial support from CAPES and FA-PEAL, and thanks the Institut Fourier for its hospitality during the preparation of this paper.

MSC2000: 53C42, 58C40.

Keywords: minimal hypersurface, eigenvalue estimate, stability, index.

The stability operator of a minimal surface in $\mathbb{H}^2 \times \mathbb{R}$ has the form $\Delta_g + a + bK_g$ with $a \geq 0$ and $b > \frac{1}{4}$. In [Section 5A](#), we consider such operators on an arbitrary complete Riemannian surface, and we show that their positivity implies an upper bound on the infimum of the spectrum of Δ_g .

Theorem 1.2. *Let (M, g) be a complete noncompact Riemannian surface. Let $a \geq 0$ and $b > \frac{1}{4}$. Denote by Δ_g the (nonnegative) Laplacian and by K_g the Gaussian curvature of (M, g) . Denote by $\lambda_\sigma(\Delta_g)$ the infimum of the spectrum of Δ_g and by $\lambda_e(\Delta_g)$ the infimum of the essential spectrum of Δ_g .*

(i) *If the operator $\Delta_g + a + bK_g$ is nonnegative on $C_0^\infty(M)$, then*

$$\lambda_\sigma(\Delta_g) \leq \frac{a}{4b-1}.$$

(ii) *If the operator $\Delta_g + a + bK_g$ has finite index on $C_0^\infty(M)$ and if M has infinite volume, then*

$$\lambda_e(\Delta_g) \leq \frac{a}{4b-1}.$$

We also prove that the positivity of such operators implies an upper bound on the volume growth of the surface ([Proposition 5.3](#)). In [Section 5B](#), applying this result to stable minimal surfaces in \mathbb{H}^3 or in $\mathbb{H}^2 \times \mathbb{R}$, we generalize and extend results of [[Candel 2007](#)]. While Candel used Pogorelov's method [[1981](#)], we use the method of Colding and Minicozzi [[2002](#)] (see also [[Castillon 2006](#)]).

In [Section 6](#), we give some applications of our general lower bounds on the spectrum to higher dimensional hypersurfaces. In [Section 2](#), we provide some preliminary technical lemmas.

2. Preliminary computations

In this section, we make some preliminary computations for later reference. For the sake of simplicity, we work in the following model for the hyperbolic space \mathbb{H}^{m+1} :

$$(1) \quad \begin{cases} \mathbb{H}^{m+1} = \mathbb{R}^m \times \mathbb{R}, \\ h = e^{2s}(dx_1^2 + \cdots + dx_m^2) + ds^2 \text{ at the point } (x, s) \in \mathbb{H}^{m+1}. \end{cases}$$

These coordinates are called *horocyclic* because the slices $\mathbb{R}^m \times \{s\}$ are horospheres and the coordinate function s is a Busemann function. They are quite natural when some Busemann function plays a special role, as will be the case in the sequel. Let

$$(2) \quad \gamma_0 : [0, \infty) \rightarrow \mathbb{H}^{m+1}, \quad u \mapsto \gamma_0(u) = (0, \dots, 0, u).$$

be a geodesic ray. The Busemann function (see [[Ballmann et al. 1985](#), page 23]) associated with γ_0 is the function

$$(3) \quad B : \mathbb{H}^{m+1} \rightarrow \mathbb{R}, \quad (x, s) \mapsto B(x, s) = s.$$

In the sequel, we denote by

$$(4) \quad \begin{cases} D^h & \text{the Levi-Civita connection,} \\ \Delta_h & \text{the geometric (that is, nonnegative) Laplacian,} \end{cases}$$

for the hyperbolic metric h on \mathbb{H}^{m+1} .

Lemma 2.1. *With the notation above, we have the formulas*

$$(5) \quad \Delta_h B = -m,$$

$$(6) \quad \text{Hess}_h B = e^{2s}(dx_1^2 + \cdots + dx_m^2)$$

at the point $(x, s) \in \mathbb{H}^{m+1}$. In particular, if we decompose the vector $u \in T_{(x,s)}\mathbb{H}^{m+1}$ h -orthogonally as $u = (u_x, u_s)$, we have

$$(7) \quad \text{Hess}_h B(u, u) = h(u_x, u_x).$$

The proof is straightforward.

Recall the following general lemmas.

Lemma 2.2. *Let $(M^m, g) \looparrowright (\widehat{M}^{m+1}, \hat{g})$ be an orientable, isometric immersion with unit normal field ν and corresponding normalized mean curvature H . Let $\hat{F} : \widehat{M} \rightarrow \mathbb{R}$ be a smooth function and let $F := \hat{F}|_M$ be its restriction to M . Then, on M ,*

$$\Delta_g F = \Delta_{\hat{g}} \hat{F}|_M + \text{Hess}_{\hat{g}} \hat{F}(\nu, \nu) - mHd\hat{F}(\nu).$$

Proof. See for example [Choe and Gulliver 1992, Lemma 2]. □

Lemma 2.3. *Assume that the manifold (M, g) carries a function f which satisfies*

$$|df|_g \leq 1 \quad \text{and} \quad -\Delta_g f \geq c \text{ for some constant } c > 0.$$

Then, any smooth, relatively compact domain $\Omega \subset M$ satisfies the isoperimetric inequalities

$$\text{Vol}_{m-1}(\partial\Omega) \geq c \text{Vol}_m(\Omega) \quad \text{and} \quad \lambda_1(\Omega) \geq \frac{c^2}{4},$$

where $\lambda_1(\Omega)$ is the least eigenvalue of Δ_g in Ω , with Dirichlet boundary condition.

Proof. Use integration by parts and Cauchy–Schwarz. □

3. Hypersurfaces in $\mathbb{H}^m \times \mathbb{R}$

We consider orientable, isometric immersions $(M^m, g) \looparrowright (\widehat{M}^{m+1}, \hat{g})$ with unit normal ν , where $\widehat{M} = \mathbb{H}^m \times \mathbb{R}$ with the product metric $\hat{g} = h + dt^2$. We take the model (1) for the hyperbolic space (here with dimension m), so that \widehat{M} is the product $\mathbb{R}^{m-1} \times \mathbb{R} \times \mathbb{R}$ with the Riemannian metric \hat{g} given by

$$\hat{g} = e^{2s}(dx_1^2 + \cdots + dx_{m-1}^2) + ds^2 + dt^2.$$

We define the function \hat{b} on \widehat{M} by

$$(8) \quad \hat{b}(x_1, \dots, x_{m-1}, s, t) = s.$$

This function is in fact a Busemann function of \widehat{M} (seen as a Cartan–Hadamard manifold) associated with a “horizontal” geodesic (justifying the name “horizontal” Busemann function used in the introduction).

Denote by $b := \hat{b}|_M$ the restriction of \hat{b} to M . Decompose the unit vector v according to the product structure $\mathbb{R}^{m-1} \times \mathbb{R} \times \mathbb{R}$, orthogonally with respect to \hat{g} , as

$$(9) \quad v = v_x + w \partial_s + v \partial_t.$$

Applying [Lemma 2.2](#), we obtain the equation

$$(10) \quad \Delta_g b = \Delta_{\hat{g}} \hat{b}|_M + \text{Hess}_{\hat{g}} \hat{b}(v, v) - m H \hat{g}(v, \partial_s).$$

Using [\(7\)](#) and [\(9\)](#), this inequality can be rewritten as

$$(11) \quad -\Delta_g b = (m-1) - |v_x|^2 + m H w.$$

From $|v_x|^2 + v^2 + w^2 = 1$, it follows that

$$(12) \quad -\Delta_g b \geq (m-2) + v^2 + w^2 - m H |w|.$$

Theorem 3.1. *Let $(M^m, g) \looparrowright (\mathbb{H}^m \times \mathbb{R}, \hat{g})$ be a complete, orientable, minimal hypersurface with normal vector v . Let v be the vertical component of the normal vector, that is, $v = \hat{g}(v, \partial_t)$.*

(i) *The function b satisfies the inequality*

$$(13) \quad -\Delta_g b \geq (m-2) + v^2,$$

and the infimum $\lambda_\sigma(\Delta_g)$ of the spectrum of the Laplacian Δ_g of M satisfies

$$(14) \quad \lambda_\sigma(\Delta_g) \geq \left(\frac{m-2 + \inf_M v^2}{2} \right)^2 \geq \left(\frac{m-2}{2} \right)^2.$$

In particular, if $m \geq 3$, (M, g) is nonparabolic.

(ii) *The spectrum of the operator $\Delta_g + (m-1)(1-v^2)$ is bounded from below by $((m-1)/2)^2$.*

Proof. Assertion (i) follows immediately from [\(12\)](#) with $H = 0$. For the last statement, apply [\(14\)](#) and [[Grigoryan 1999](#), Proposition 10.1]. To prove (ii), we start from the inequality [\(13\)](#), $-\Delta_g b \geq (m-2) + v^2$. Multiplying this inequality by f^2 , where $f \in C_0^\infty(M)$, and integrating by parts using the fact that $|db|_g \leq 1$ gives (all integrals are taken with respect to the Riemannian measure dv_g)

$$(m-2) \int_M f^2 + \int_M v^2 f^2 \leq \int_M |df|^2 \leq 2 \int_M |f| |df|.$$

Rewrite this inequality as

$$(m-1) \int_M f^2 \leq 2 \int_M |f| |df| + \int_M (1-v^2) f^2.$$

Using the Cauchy–Schwarz inequality $2|f| \cdot |df| \leq \frac{1}{a} |df|^2 + a f^2$ for $a > 0$, we obtain

$$a(m-1-a) \int_M f^2 \leq \int_M (|df|^2 + a(1-v^2) f^2) \leq \int_M (|df|^2 + (m-1)(1-v^2) f^2),$$

provided that $0 \leq a \leq m-1$. We can now maximize the constant on the left-hand side by choosing $a = (m-1)/2$. \square

When the mean curvature H is nonzero, we also obtain the following result from inequality (12):

Proposition 3.2. *Let $(M^m, g) \looparrowright (\mathbb{H}^m \times \mathbb{R}, \hat{g})$ be a complete, orientable hypersurface with normal vector v and constant mean curvature H for $0 \leq H \leq (m-1)/m$. Recall that $v = \hat{g}(v, \partial_t)$. Then,*

$$(15) \quad -\Delta_g b \geq (m-2)(1 - \sqrt{1-v^2}) + (m-2) \left(1 - \frac{mH}{m-2}\right) \sqrt{1-v^2}.$$

Remarks. (i) Inequalities (13), (14) and Theorem 3.1(ii) are sharp. Indeed, taking the horizontal slice $M = \mathbb{H}^m \times \{0\}$ gives the case $v = 1$, and taking $M = \mathbb{P} \times \mathbb{R}$, where \mathbb{P} is some totally geodesic $(m-1)$ -space in \mathbb{H}^m , gives the case $v = 0$.

- (ii) If we assume that $v^2 \leq \alpha^2 < 1$, the spectrum of $\Delta_g + (m-1)(1-v^2)$ is bounded from below by $(m-1)(1-\alpha^2)$.
- (iii) Inequality (15) generalizes an earlier result of [Castillon 1997] for submanifolds immersed into Hadamard manifolds. For other estimates, see [Bessa and Costa 2009]. We point out that it is more convenient in our context to use the “horizontal” Busemann function rather than the hyperbolic distance function as in [Castillon 1997].
- (iv) Inequality (15) still holds if M^m is only assumed to have mean curvature bounded from above by H .

4. Applications to minimal hypersurfaces in $\mathbb{H}^m \times \mathbb{R}$

4A. Index of minimal surfaces immersed in $\mathbb{H}^m \times \mathbb{R}$. The Jacobi (or stability) operator of a minimal hypersurface $M^m \looparrowright \mathbb{H}^m \times \mathbb{R}$ is given by

$$(16) \quad J_M = \Delta + (m-1)(1-v^2) - |A|^2,$$

where v is the vertical component of the unit normal ν and A the second fundamental form of the immersion (see [Bérard and Sá Earp 2008]). As a corollary of Theorem 3.1, we obtain:

Theorem 4.1. *Let $(M^2, g) \looparrowright (\mathbb{H}^2 \times \mathbb{R}, \hat{g})$ be a complete, orientable, minimal surface with second fundamental form A . If $\int_M |A|^2 dv_g$ is finite, then the immersion has finite index.*

Proof. When $\int_M |A|^2$ is finite, the second fundamental form tends to zero uniformly at infinity (see [Bérard and Sá Earp 2008, Theorem 4.1]). Theorem 3.1(ii) with $m = 2$ implies that the essential spectrum of the Jacobi operator J_M is bounded from below by $\frac{1}{4}$. Since the operator J_M is also bounded from below, it has only finitely many negative eigenvalues (see [Bérard et al. 1997, Proposition 1]). \square

Remark. This theorem answers a question raised in [Bérard and Sá Earp 2008], where the finiteness of the index of J_M is proved in dimension $m \geq 3$ under the assumption that $\int_M |A|^m$ is finite, and in dimension 2 under the assumption that both $\int_M v^2$ and $\int_M |A|^2$ are finite. In dimension $m \geq 3$, the index of J_M is bounded from above by a constant times $\int_M |A|^m$ (see [Bérard and Sá Earp 2008]). In the next section, we investigate bounds on the index in dimension 2.

4B. Bounds on the index of minimal surfaces immersed in $\mathbb{H}^m \times \mathbb{R}$.

Proposition 4.2. *Let $(M^2, g) \looparrowright (\mathbb{H}^2 \times \mathbb{R}, \hat{g})$ be a complete, orientable, minimal surface with second fundamental form A . If $\int_M |A|^2 dv_g$ is finite, then for any $r > 1$, there exists a constant C_r such that the index of the immersion is bounded from above by $C_r \int_M |A|^{2r} dv_g$.*

Remarks. (i) The assumption that $\int_M |A|^2 dv_g$ is finite implies that A tends to zero uniformly at infinity. Thus the integrals $\int_M |A|^{2r} dv_g$ are all finite.

(ii) Our proof provides a constant C_r that tends to infinity when r tends to 1. We do not know whether there is a bound for the index in terms of $\int_M |A|^2 dv_g$, as is the case for minimal surfaces in \mathbb{R}^3 [Tysk 1987].

Proof. As in Section 4A, we write the Jacobi operator as $J = \Delta_g + 1 - v^2 - |A|^2$. The closure \tilde{Q} of the quadratic form

$$Q[f] = \int_M (|df|^2 + (1 - v^2)f^2) dv_g$$

with domain $C_0^1(M)$ satisfies the Beurling–Deny condition (if f is in the domain of \tilde{Q} , then so is $|f|$ and $\tilde{Q}[|f|] = \tilde{Q}[f]$; see [Davies 1989, Theorem 1.3.2]) and, by Theorem 3.1, the Cheeger inequality

$$(17) \quad \int_M f^2 dv_g \leq 4Q[f] \quad \text{for all } f \in C_0^1(M).$$

On the other hand, the surface M satisfies the Sobolev inequality

$$(18) \quad \int_M f^2 dv_g \leq S \left(\int_M |df|_g^2 dv_g \right)^2 \quad \text{for all } f \in C_0^1(M),$$

for some constant $S > 0$. Indeed, this follows from the Sobolev inequality for minimal surfaces in $\mathbb{H}^2 \times \mathbb{R}$, using the fact that the ambient space has nonpositive curvature and infinite injectivity radius (see [Hoffman and Spruck 1974]).

From the above Cheeger and Sobolev inequalities, we can establish that for any $q \geq 1$, there exists a constant D_q such that for any $f \in C_0^1(M)$,

$$(19) \quad \left(\int_M |f|^{2q} dv_g \right)^{1/q} \leq D_q Q[f].$$

When q is an integer, the inequality follows from an induction argument and we can conclude by interpolation.

We can then apply [Levin and Solomyak 1997, Theorem 1.2] to conclude that the index is less than $e^p D_q^p \int_M |A|^{2p} dv_g$, where $p = q/(q-1)$. \square

4C. Hypersurfaces in a slab. In this section, we use the computations of Section 3 to give a lower bound on the spectrum of the Laplacian on a complete minimal surface immersed in a slab $\mathbb{H}^2 \times [-a, a]$, where $a > 0$.

First, consider functions on $\mathbb{H}^m \times \mathbb{R}$ depending only on the height t , namely $\hat{\beta}(x, s, t) = f(t)$. In this case, $d\hat{\beta} = f'(t) dt$, and

$$\text{Hess}_{\hat{\beta}} \hat{\beta}(X, Y) = f''(t) \hat{g}(X, \partial_t) \hat{g}(Y, \partial_t).$$

In particular,

$$\Delta_{\hat{\beta}} \hat{\beta} = -f''(t) \quad \text{and} \quad \text{Hess}_{\hat{\beta}} \hat{\beta}(v, v) = v^2 f''(t).$$

Define $\beta = \hat{\beta}|_M$. Using Lemma 2.2, we have

$$(20) \quad -\Delta_g \beta = (1 - v^2) f''(t) + m H v f'(t).$$

To estimate the first eigenvalue of a *minimal* hypersurface $M^m \looparrowright \mathbb{H}^m \times \mathbb{R}$, use the identity (20) with some particular choice of f . For instance, let

$$\hat{\beta}(x, s, t) = \tfrac{1}{2} t^2.$$

In this case,

$$-\Delta_g \beta = (1 - v^2).$$

Assume now that $M^m \looparrowright \mathbb{H}^m \times [-a, a]$, for some $a > 0$. Then,

$$-\Delta_g \beta = (1 - v^2) \quad \text{and} \quad |d\beta| \leq a.$$

Defining $Z = b + \beta$, where b is the restriction of the Busemann function \hat{b} to M^m , we can use the last inequality in (12) to obtain

$$(21) \quad -\Delta Z \geq m - 1 \quad \text{and} \quad |dZ| \leq \sqrt{1 + a^2}.$$

Using the above notation and Lemma 2.3, we have the following estimate:

Proposition 4.3. *Given $a > 0$, let $(M^m, g) \looparrowright (\mathbb{H}^m \times [-a, a], \hat{g})$ be a complete, immersed, orientable, minimal hypersurface. Then, the infimum of the spectrum of Δ_g on M is positive. More precisely,*

$$(22) \quad \lambda_\sigma(\Delta_g) \geq \frac{(m-1)^2}{4(1+a^2)}.$$

5. Bounds derived from a stability assumption

Let (M, g) be a complete Riemannian surface with (nonnegative) Laplace operator Δ_g and Gaussian curvature K_g . Let a, b be real numbers with $a \geq 0$ and $b > 1/4$. Let L be the operator $L = \Delta_g + a + bK_g$.

Let $\text{Ind}(L, \Omega)$ denote the number of negative eigenvalues of the operator L in Ω with Dirichlet boundary conditions on $\partial\Omega$. The index $\text{Ind}(L)$ of the operator L is defined to be the supremum

$$\text{Ind}(L) = \sup\{\text{Ind}(L, \Omega) \mid \Omega \Subset M\}$$

taken over the relatively compact subdomains Ω in M .

In Section 5A, we state two intrinsic consequences of the assumption that the operator L has finite index. In Sections 5B and 5C, we consider applications to minimal and CMC surfaces.

5A. Intrinsic results. We prove the next theorem using the method of [Colding and Minicozzi 2002], and more precisely [Castillon 2006, Lemma 1.8].

Theorem 5.1. *Let (M, g) be a complete noncompact Riemannian surface. Let $a \geq 0$ and $b > \frac{1}{4}$. Denote by Δ_g the (nonnegative) Laplacian and by K_g the Gaussian curvature of (M, g) . Denote by $\lambda_\sigma(\Delta_g)$ the infimum of the spectrum of Δ_g and by $\lambda_e(\Delta_g)$ the infimum of the essential spectrum of Δ_g .*

(i) *If the operator $\Delta_g + a + bK_g$ is nonnegative on $C_0^\infty(M)$, then*

$$\lambda_\sigma(\Delta_g) \leq \frac{a}{4b-1}.$$

(ii) *If the operator $\Delta_g + a + bK_g$ has finite index on $C_0^\infty(M)$ and if M has infinite volume, then*

$$\lambda_e(\Delta_g) \leq \frac{a}{4b-1}.$$

Proof. (i) We can assume the surface M has infinite volume (otherwise $\lambda_\sigma(\Delta_g) = 0$ because the function 1 is in $L^2(M, v_g)$, and the estimate is trivial). Fix a point x_0 in M and let $r(x)$ denote the Riemannian distance to the point x_0 . Given $S > R > 0$, let $B(R)$ denote the open geodesic ball in M with center x_0 and radius R . Let $C(R, S)$ denote the open annulus $B(S) \setminus \bar{B}(R)$. Let $V(R)$ denote the volume of $B(R)$ and $L(R)$ the length of its boundary $\partial B(R)$. Let $G(R)$ denote the integral curvature of $B(R)$, that is, $G(R) = \int_{B(R)} K_g(x) dv_g(x)$, where dv_g is the Riemannian measure. The main idea in [Castillon 2006] is to use the work of [Shiohama and Tanaka 1989; 1993] on the length of geodesic circles, which shows that the function $L(r)$ is differentiable almost everywhere and is related to the Euler characteristic and to the integral curvature of geodesic balls by the formula

$$L'(r) \leq 2\pi \chi(B(r)) - G(r) \leq 2\pi - G(r)$$

(see [Castillon 2006, Theorem 1.7]; the second inequality comes from the fact that the Euler characteristic of balls is less than or equal to 1).

Now choose ξ as in the following lemma:

Lemma 5.2 [Castillon 2006, Lemma 1.8]. *For $0 < R < S$, let $\xi : [R, S] \rightarrow \mathbb{R}$ be such that $\xi \geq 0$, $\xi' \leq 0$, $\xi'' \geq 0$ and $\xi(S) = 0$. Then*

$$\begin{aligned} & \int_{C(R,S)} K_g \xi^2(r) dv_g \\ & \leq -\xi^2(R)G(R) + 2\pi \xi^2(R) - 2\xi(R)\xi'(R)L(R) - \int_{C(R,S)} (\xi^2)''(r) dv_g. \quad \square \end{aligned}$$

Choose a function $f : B(S) \rightarrow \mathbb{R}$ such that $f(r) \equiv \xi(R)$ on $B(R)$, $f(r) = \xi(r)$ on $C(R, S)$, and write the positivity assumption

$$0 \leq \int_M |df|_g^2 dv_g + a \int_M f^2 dv_g + b \int_M K_g f^2 dv_g.$$

On the ball $B(R)$, we have $\int_{B(R)} K_g f^2 dv_g = \xi^2(R)G(R)$ and $\int_{B(R)} |df|^2 dv_g = 0$. Using Lemma 5.2, we obtain

$$\begin{aligned} 0 & \leq \int_{C(R,S)} (\xi')^2(r) dv_g + a \int_M f^2 dv_g + b \xi^2(R)G(R) - b \xi^2(R)G(R) \\ & \quad + 2\pi b \xi^2(R) - 2b \xi(R)\xi'(R)L(R) - b \int_{C(R,S)} (\xi^2)''(r) dv_g, \end{aligned}$$

and hence,

$$\begin{aligned} (23) \quad 0 & \leq (1 - 2b) \int_{C(R,S)} (\xi')^2(r) dv_g + a \int_M f^2 dv_g + 2\pi b \xi^2(R) \\ & \quad - 2b \xi(R)\xi'(R)L(R) - 2b \int_{C(R,S)} \xi(r)\xi''(r) dv_g. \end{aligned}$$

Choose $\xi(r) = (S - r)^k$ in $[R, S]$ for $k \geq 1$ big enough (we will eventually let k tend to infinity). Then $\xi(r)\xi''(r) = (1 - \frac{1}{k})(\xi'(r))^2$. It follows that

$$0 \leq \left(1 - 4b + \frac{2b}{k}\right) \int_M |df|^2 dv_g + a \int_M f^2 dv_g + 2b(\pi(S - R)^{2k} + kL(R)(S - R)^{2k-1}).$$

Using the fact that $\int_M f^2 dv_g \geq (S - R)^{2k} V(R)$, we obtain

$$(24) \quad \lambda_\sigma(\Delta_g) \leq \frac{\int_M |df|^2 dv_g}{\int_M f^2 dv_g} \leq \frac{a}{4b - 1 - 2b/k} + \frac{2b}{(4b - 1 - 2b/k)V(R)} \left(\pi + \frac{kL(R)}{S - R} \right).$$

First letting S tend to infinity, then letting R tend to infinity, using the fact that M has infinite volume, and finally letting k tend to infinity gives

$$\lambda_\sigma(\Delta_g) \leq \frac{a}{4b - 1}.$$

(ii) The finiteness of the index of the operator $\Delta_g + a + bK_g$ implies that it is nonnegative outside a compact set (see [Fischer-Colbrie 1985, Proposition 1], for instance). Choose R_0 big enough for $\Delta_g + a + bK_g$ to be nonnegative in $M \setminus B(R_0)$. Next, for $S > R > R_1 + 1 > R_0 + 1$, choose ξ as in Lemma 5.2 and a test function

$$(25) \quad f(r) = \begin{cases} 0 & \text{in } B(R_1), \\ \xi(R)(r - R_1) & \text{in } C(R_1, R_1 + 1), \\ \xi(R) & \text{in } C(R_1 + 1, R), \\ \xi(r) & \text{in } C(R, S). \end{cases}$$

Following the same scheme as for (i), and under the assumption that the volume of M is infinite, we can prove that the bottom of the spectrum of Δ_g in $M \setminus B(R_1)$ with Dirichlet boundary conditions on $\partial B(R_1)$ satisfies the inequality

$$\lambda_\sigma(\Delta_g, M \setminus B(R_1)) \leq \frac{a}{4b - 1}.$$

To conclude the proof of Theorem 5.1, we use the fact that

$$\lambda_e(\Delta_g) = \lim_{R \rightarrow \infty} \lambda_\sigma(\Delta_g, M \setminus B(R)). \quad \square$$

Proposition 5.3. *Let (M, g) be a complete Riemannian surface with (nonnegative) Laplace operator Δ_g and Gaussian curvature K_g . Let $V(r)$ denote the volume of the geodesic ball of radius r in M centered at some point. Let a, b be positive real numbers with $b > \frac{1}{4}$. Let $\alpha_0 = \sqrt{a/(4b - 1)}$. If the operator $L := \Delta_g + a + bK_g$ has finite index, then*

$$\int_0^\infty e^{-2\alpha r} V(r) dr < \infty \quad \text{for all } \alpha > \alpha_0,$$

and hence, the lower volume growth of M satisfies

$$\liminf_{r \rightarrow \infty} \frac{\ln V(r)}{r} \leq 2\alpha_0.$$

Proof. It follows from our assumptions that the operator L is positive outside some compact set (see [Fischer-Colbrie 1985, Proposition 1]). In particular, it is positive on $M \setminus B(R_0)$ for some radius R_0 . Choose $R > R_0 + 1$ and define the function

$$(26) \quad \xi(r) = \begin{cases} 0 & \text{for } r \leq R_0, \\ (1 - (R_0 + 1)/R)^{\alpha R} (r - R_0) & \text{for } R_0 \leq r \leq R_0 + 1, \\ (1 - r/R)^{\alpha R} & \text{for } R_0 + 1 \leq r \leq R, \end{cases}$$

where the parameter α will be chosen later on. The positivity of the operator L on $M \setminus B(R_0)$ implies that

$$0 \leq \int_M ((\xi'(r))^2 + a\xi^2(r) + bK_g \xi^2(r)) dv_g.$$

Write this integral as the sum of $\int_{C(R_0, R_0+1)}$ and $\int_{C(R_0+1, R)}$. The first integral can be written as

$$\int_{C(R_0, R_0+1)} = \left(1 - \frac{R_0+1}{R}\right)^{\alpha R} C(B(R_0)),$$

where $C(B(R_0))$ is a constant which only depends on the geometry of M on the ball $B(R_0)$. Using Lemma 5.2 and the fact that $\chi(B(r)) \leq 1$ for all r , the second integral can be estimated as follows:

$$\begin{aligned} \int_{C(R_0+1, R)} &\leq \int_{C(R_0+1, R)} ((\xi')^2 + a\xi^2 - b(\xi^2)'') dv_g \\ &\quad + 2\pi b - \xi^2(R_0 + 1)G(R_0 + 1) + 2\alpha L(R_0 + 1). \end{aligned}$$

Using (26), the definition for the function ξ , the integral in the first line of the above inequality can be written as

$$-\left((4b - 1)\alpha^2 - \frac{2b\alpha}{R} - a\right) \int_{R_0+1}^R \left(1 - \frac{r}{R}\right)^{2\alpha R - 2} L(r) dr.$$

Taking α big enough so that the constant is positive, and using $L(r) = V'(r)$, we obtain the inequality

$$\frac{2\alpha R - 2}{R} \left((4b - 1)\alpha^2 - \frac{2b\alpha}{R} - a\right) \int_{R_0+1}^R \left(1 - \frac{r}{R}\right)^{2\alpha R - 3} V(r) dr \leq D(B(R_0), \alpha),$$

where $D(B(R_0), \alpha)$ is a constant which only depends on the geometry of M in the ball $B(R_0)$ and α . Letting R tend to infinity finally shows

$$2\alpha \left((4b - 1)\alpha^2 - a\right) \int_{R_0+1}^{\infty} e^{-2\alpha r} V(r) dr < \infty,$$

provided that $\alpha > \alpha_0$, which proves the first assertion in the proposition. The second assertion follows easily. \square

Remark. In [Proposition 5.3](#), we assumed that $a > 0$. In the case $a = 0$, the volume growth is at most quadratic (see [\[Castillon 2006, Proposition 2.2\]](#)).

5B. Applications to stable minimal surfaces in \mathbb{H}^3 or $\mathbb{H}^2 \times \mathbb{R}$. Let M be a complete, orientable, minimal immersion into either 3-dimensional hyperbolic space \mathbb{H}^3 or into $\mathbb{H}^2 \times \mathbb{R}$. Let J_M denote the Jacobi operator of the immersion.

In the case of a minimal immersion $M \looparrowright \mathbb{H}^3(-1)$, the operator J_M takes the form $J_M = \Delta_M + 2 - |A|^2$, where A is the second fundamental form. Using the Gauss equation, we have that $K_M = -1 - \frac{1}{2}|A|^2$, so that we can rewrite the Jacobi operator of $M \looparrowright \mathbb{H}^3(-1)$ as

$$(27) \quad J_M = \Delta_M + 4 + 2K_M.$$

In the case of a minimal immersion $M \looparrowright \mathbb{H}^2(-1) \times \mathbb{R}$, the Jacobi operator is given by $J_M = \Delta_M + 1 - v^2 - |A|^2$, where v is the vertical component of the unit normal vector to the surface. Using the Gauss equation, we have that $K_M = -v^2 - \frac{1}{2}|A|^2$, so that we can rewrite the Jacobi operator of $M \looparrowright \mathbb{H}^2(-1) \times \mathbb{R}$ as

$$(28) \quad J_M = \Delta_M + 2 + 2K_M - (1 - v^2) \leq \tilde{J}_M := \Delta_M + 2 + 2K_M.$$

In this case, the positivity of the operator J_M implies the positivity of the operator \tilde{J}_M .

Applying [Theorem 5.1](#) to the operator J_M in the form (27) when M is a minimal surface in \mathbb{H}^3 , respectively, to the operator \tilde{J}_M in the form (28) when M is a minimal surface in $\mathbb{H}^2 \times \mathbb{R}$, we obtain the following corollary.

Corollary 5.4. *Let $(M, g) \looparrowright (\hat{M}, \hat{g})$ be a complete, orientable, minimal immersion. Assume that the immersion is stable.*

- (i) *If $\hat{M} = \mathbb{H}^3$, then $\lambda_\sigma(\Delta_g) \leq \frac{4}{7}$.*
- (ii) *If $\hat{M} = \mathbb{H}^2 \times \mathbb{R}$, then $\lambda_\sigma(\Delta_g) \leq \frac{2}{7}$.*

If the immersion is only assumed to have finite index, then the same inequalities hold with $\lambda_\sigma(\Delta_g)$ replaced by $\lambda_e(\Delta_g)$, the infimum of the essential spectrum.

Remarks. (i) The first assertion improves an earlier result from [\[Candel 2007\]](#), to the effect that $\lambda_\sigma(M) \leq \frac{4}{3}$, provided that M is a complete, simply connected, stable minimal surface in \mathbb{H}^3 .

- (ii) In both cases, the bottom of the spectrum of a totally geodesic \mathbb{H}^2 is $\frac{1}{4}$.

Applying [Proposition 5.3](#), we have the following corollary.

Corollary 5.5. *Let $(M, g) \looparrowright (\widehat{M}, \widehat{g})$ be a complete, orientable, minimal immersion. Let μ denote the lower volume growth rate of M , given by*

$$\mu = \liminf_{r \rightarrow \infty} \frac{\ln V(r)}{r},$$

where $V(r)$ is the volume of the geodesic ball $B(x_0, r)$ about some point. Assume that the immersion has finite index.

(i) *If $\widehat{M} = \mathbb{H}^3$, then $\mu \leq 2\sqrt{\frac{4}{7}}$.*

(ii) *If $\widehat{M} = \mathbb{H}^2 \times \mathbb{R}$, then $\mu \leq 2\sqrt{\frac{2}{7}}$.*

Remarks. (i) Assertion (i) in [Corollary 5.5](#) improves an earlier upper bound on μ given in [\[Candel 2007\]](#) under the assumption that M is simply connected.

(ii) Recall from [\[Brooks 1981; Kumura 2007\]](#) that the volume growth is related to the infimum of the essential spectrum by the formula

$$\lambda_e(\Delta_g) \leq \left(\frac{1}{2} \liminf_{r \rightarrow \infty} \frac{\ln V(r)}{r} \right)^2.$$

5C. Further applications. The argument above also works for surfaces with constant mean curvature $|H| \leq 1$ in hyperbolic space. In that case,

$$K_M = -(1 - H^2) - \frac{1}{2}|A|^2 \quad \text{and} \quad J_M = \Delta_M + 4(1 - H^2) + 2K_M.$$

Proposition 5.6. *Let $(M, g) \looparrowright \mathbb{H}^3$ be a complete, orientable, stable CMC immersion with $|H| \leq 1$. Then*

$$\lambda_\sigma(\Delta_g) \leq \frac{4(1 - H^2)}{7}.$$

The space $\mathbb{H}^2 \times \mathbb{R}$ is a simply connected 3-dimensional homogeneous manifold, whose isometry group has dimension 4. Such manifolds have been well studied (see for instance [\[Daniel 2007\]](#) and references therein) and can be parametrized by two real parameters, say κ and τ with $\kappa \neq 4\tau^2$. Denote these manifolds by $\mathbb{E}^3(\kappa, \tau)$. When $\tau = 0$, the manifold $\mathbb{E}^3(\kappa, 0)$ is the product space $\mathbb{E}^2(\kappa) \times \mathbb{R}$, where $\mathbb{E}^2(\kappa)$ is the space form of constant curvature κ . In particular, $\mathbb{H}^2 \times \mathbb{R} = \mathbb{E}^3(-1, 0)$.

If $(M, g) \looparrowright \mathbb{E}^3(\kappa, \tau)$ is an immersed CMC H surface, then its Jacobi operator is given by (see [\[Daniel 2007, Proposition 5.11\]](#))

$$J_M := \Delta_g + 2K - 4H^2 - \kappa - (\kappa - 4\tau^2)v^2.$$

In the next proposition we give an upper bound for the bottom of the spectrum in this general framework.

Proposition 5.7. *Let $(M, g) \looparrowright \mathbb{E}^3(\kappa, \tau)$ be a complete, orientable, stable CMC H immersion such that $\kappa < 4\tau^2$. Assume furthermore that $2H^2 \leq (2\tau^2 - \kappa)$. Then*

$$\lambda_\sigma(\Delta_g) \leq \frac{4\tau^2 - 2\kappa - 4H^2}{7}.$$

Proof. Under the hypotheses we have the inequalities

$$0 \leq \Delta_g + 2K - 4H^2 - \kappa - (\kappa - 4\tau^2)v^2 \leq \Delta_g + 2K - 4H^2 - 2(\kappa - 2\tau^2),$$

and we may apply [Theorem 5.1](#) again. \square

6. Applications in higher dimensions

In this section, we give some further applications of the inequalities proved in [Section 3](#). In the following proposition, we give a structure theorem for minimal hypersurfaces in $\mathbb{H}^m \times \mathbb{R}$.

Let $M^m \looparrowright \mathbb{H}^m \times \mathbb{R}$ be a complete, orientable minimal hypersurface with second fundamental form A , where $\|A\|_m < \infty$ and $m \geq 3$. By [[Bérard and Sá Earp 2008](#)], the Ricci curvature of M^m satisfies the inequality (30) below and furthermore $|A|$ tends to zero uniformly at infinity. It follows that M satisfies the assumption of [[Li and Wang 2001](#), Theorem 3.1] provided that $m \geq 7$ and hence that M has only finitely many infinite volume ends. On the other-hand, all ends of M must have infinite volume [[Cheng et al. 2008](#), Proposition 2.1]. Thus M has only finitely many ends provided that $m \geq 7$. The next result gives a sufficient condition for M to have only one end.

Proposition 6.1. *Let $M^m \looparrowright \mathbb{H}^m \times \mathbb{R}$, $m \geq 3$, be a complete, orientable minimal hypersurface with unit normal field ν and second fundamental form A . Let v denote the component of ν along ∂_t . Assume that*

- (i) $m \geq 7$ and $0 \leq \alpha \leq 1$, or
- (ii) $m = 6$ and $0.083 \leq \alpha \leq 1$, or
- (iii) $m = 5$ and $0.578 \leq \alpha \leq 1$.

There exists a constant $c(m, \alpha) > 0$ such that, if M satisfies $\|A\|_m \leq c(m, \alpha)$ and $v^2 \geq \alpha^2$, then M carries no nontrivial L^2 -harmonic 1-form and hence has at most one end.

Sketch of proof. Step 1: According to [[Hoffman and Spruck 1974](#)], the manifold M^m satisfies the Sobolev inequality

$$(29) \quad \|\varphi\|_{2m/(m-2)}^2 \leq S(2, m) \|d\varphi\|_2^2 \quad \text{for all } \varphi \in C_0^1(M).$$

Step 2: Let $u \in T_1 M$ be a unit tangent vector to M . By the Gauss equation, we have

$$\text{Ric}(u, u) = \widehat{\text{Ric}}(u, u) - \hat{R}(u, \nu, u, \nu) - |A(u)|^2,$$

where Ric denotes the Ricci curvature of M , $\widehat{\text{Ric}}$ the Ricci curvature of $\widehat{M} = \mathbb{H}^m \times \mathbb{R}$ and \widehat{R} the curvature tensor of \widehat{M} , and where A denotes the Weingarten operator of the immersion. Using the curvature computations in [Bérard and Sá Earp 2008] and the fact that A has trace zero, we obtain the inequality

$$(30) \quad \text{Ric}(u, u) \geq -(m-1) - \frac{m-1}{m}|A|^2.$$

Step 3: Let ω be an L^2 harmonic 1-form on M . Using the Weitzenböck formula for 1-forms, the improved Kato inequality

$$(31) \quad \frac{1}{m-1}|d|\omega||^2 \leq |D\omega|^2 - |d|\omega||^2,$$

and inequality (30), we find that ω satisfies the inequality

$$(32) \quad \frac{1}{m-1}|d|\omega||^2 + |\omega|\Delta|\omega| \leq (m-1)|\omega|^2 + \frac{m-1}{m}|A|^2|\omega|^2$$

in the weak sense. The following formal calculation can easily be made rigorous by using cut-off functions. Integrate (32) over M using integration by parts and using the notation $f := |\omega|$, we obtain

$$\frac{m}{m-1} \int_M |df|^2 \leq (m-1) \int_M f^2 + \frac{m-1}{m} \int_M |A|^2 f^2.$$

Plug the assumption $|v| \geq \alpha$ and the inequality (14) into the preceding inequality. Use Hölder's inequality to estimate the integral $\int_M |A|^2 f^2$ and the Sobolev inequality (29). It follows that

$$\left(\frac{m}{m-1} - \frac{4(m-1)}{(m-2+\alpha)^2} \right) \|f\|_{2m/(m-2)}^2 \leq S(2, m) \frac{m-1}{m} \|A\|_m^2 \|f\|_{2m/(m-2)}^2$$

and we can conclude the proof with the constant

$$c(m, \alpha) = \frac{m}{(m-1)S(2, m)} \frac{m(m-2+\alpha)^2 - 4(m-1)^2}{(m-1)(m-2+\alpha)^2}. \quad \square$$

Proposition 6.2. *Let $M^m \looparrowright \mathbb{H}^m \times \mathbb{R}$, $m \geq 3$, be a complete, orientable minimal hypersurface with unit normal field v and second fundamental form A . Let v denote the component of v along ∂_t . The immersion M is stable if*

- (i) $\|A\|_\infty \leq \frac{1}{4}(m-1)^2$, or
- (ii) $\|A\|_\infty \leq \frac{1}{4}(m-2+\alpha)^2$ and $v^2 \geq \alpha$, or
- (iii) $|A|^2 + (m-1)v^2 \leq \frac{1}{4}m^2$ on M .

Proof. Recall that the Jacobi operator J_M of the immersion M is given by $J_M = \Delta_g + (m-1)(1-v^2) - |A|^2$. The proposition follows from Theorem 3.1. \square

Remark 1. Condition (ii) is not so interesting. Indeed, if M is connected, we may assume that $v > 0$ and it follows that M is stable because v is a Jacobi field.

Remark 2. We can write the operator J_M as

$$J_M = \Delta_g - \left(\frac{m-2}{2}\right)^2 + \left(\left(\frac{m}{2}\right)^2 - |A|^2\right).$$

In view of the results à la Lieb or Li and Yau, one can show that if the integral

$$\int_M \left(\left(\frac{m}{2}\right)^2 - |A|^2\right)_-^{m/2}$$

is small enough, then M is stable.

Acknowledgements

The authors thank the referee for useful comments.

References

- [Ballmann et al. 1985] W. Ballmann, M. Gromov, and V. Schroeder, *Manifolds of nonpositive curvature*, Progress in Math. **61**, Birkhäuser, Boston, 1985. [MR 87h:53050](#) [Zbl 0591.53001](#)
- [Bérard and Sá Earp 2008] P. Bérard and R. Sá Earp, “Minimal hypersurfaces in $\mathbb{H}^n \times \mathbb{R}$, total curvature and index”, preprint, 2008. [arXiv 0808.3838v3](#)
- [Bérard et al. 1997] P. Bérard, M. do Carmo, and W. Santos, “The index of constant mean curvature surfaces in hyperbolic 3-space”, *Math. Z.* **224**:2 (1997), 313–326. [MR 98a:53008](#) [Zbl 0868.53010](#)
- [Bessa and Costa 2009] G. P. Bessa and M. S. Costa, “Eigenvalue estimates for submanifolds with locally bounded mean curvature in $N \times \mathbb{R}$ ”, *Proc. Amer. Math. Soc.* **137**:3 (2009), 1093–1102. [MR 2010b:53099](#) [Zbl 1169.53045](#)
- [Brooks 1981] R. Brooks, “A relation between growth and the spectrum of the Laplacian”, *Math. Z.* **178**:4 (1981), 501–508. [MR 83a:58089](#) [Zbl 0458.58024](#)
- [Candel 2007] A. Candel, “Eigenvalue estimates for minimal surfaces in hyperbolic space”, *Trans. Amer. Math. Soc.* **359**:8 (2007), 3567–3575. [MR 2007m:53076](#) [Zbl 1115.53005](#)
- [Castillon 1997] P. Castillon, “Sur l’opérateur de stabilité des sous-variétés à courbure moyenne constante dans l’espace hyperbolique”, *Manuscripta Math.* **94**:3 (1997), 385–400. [MR 99c:53062](#) [Zbl 0902.53044](#)
- [Castillon 2006] P. Castillon, “An inverse spectral problem on surfaces”, *Comment. Math. Helv.* **81**:2 (2006), 271–286. [MR 2007b:58042](#) [Zbl 1114.58025](#)
- [Cheng et al. 2008] X. Cheng, L.-f. Cheung, and D. Zhou, “The structure of weakly stable constant mean curvature hypersurfaces”, *Tohoku Math. J. (2)* **60**:1 (2008), 101–121. [MR 2009i:53053](#) [Zbl 1154.53036](#)
- [Choe and Gulliver 1992] J. Choe and R. Gulliver, “Isoperimetric inequalities on minimal submanifolds of space forms”, *Manuscripta Math.* **77**:2-3 (1992), 169–189. [MR 93k:53059](#) [Zbl 0777.53063](#)
- [Colding and Minicozzi 2002] T. H. Colding and W. P. Minicozzi, II, “Estimates for parametric elliptic integrands”, *Int. Math. Res. Not.* **2002**:6 (2002), 291–297. [MR 2002k:53060](#) [Zbl 1002.53035](#)
- [Daniel 2007] B. Daniel, “Isometric immersions into 3-dimensional homogeneous manifolds”, *Comment. Math. Helv.* **82**:1 (2007), 87–131. [MR 2008a:53058](#) [Zbl 1123.53029](#)
- [Davies 1989] E. B. Davies, *Heat kernels and spectral theory*, Cambridge Tracts in Math. **92**, Cambridge University Press, Cambridge, 1989. [MR 90e:35123](#) [Zbl 0699.35006](#)

- [Fischer-Colbrie 1985] D. Fischer-Colbrie, “On complete minimal surfaces with finite Morse index in three-manifolds”, *Invent. Math.* **82**:1 (1985), 121–132. [MR 87b:53090](#) [Zbl 0573.53038](#)
- [Grigoryan 1999] A. Grigoryan, “Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds”, *Bull. Amer. Math. Soc. (N.S.)* **36**:2 (1999), 135–249. [MR 99k:58195](#) [Zbl 0927.58019](#)
- [Hoffman and Spruck 1974] D. Hoffman and J. Spruck, “Sobolev and isoperimetric inequalities for Riemannian submanifolds”, *Comm. Pure Appl. Math.* **27**:6 (1974), 715–727. [MR 51 #1676](#) [Zbl 0295.53025](#)
- [Kumura 2007] H. Kumura, “Infimum of the exponential volume growth and the bottom of the essential spectrum of the Laplacian”, preprint, 2007. [arXiv 0707.0185v1](#)
- [Levin and Solomyak 1997] D. Levin and M. Solomyak, “The Rozenblum–Lieb–Cwikel inequality for Markov generators”, *J. Anal. Math.* **71** (1997), 173–193. [MR 98j:47090](#) [Zbl 0910.47017](#)
- [Li and Wang 2001] P. Li and J. Wang, “Complete manifolds with positive spectrum”, *J. Differential Geom.* **58**:3 (2001), 501–534. [MR 2003e:58046](#) [Zbl 1032.58016](#)
- [Pogorelov 1981] A. V. Pogorelov, “On the stability of minimal surfaces”, *Dokl. Akad. Nauk SSSR* **260**:2 (1981), 293–295. In Russian. [MR 83b:49043](#) [Zbl 0495.53005](#)
- [Shiohama and Tanaka 1989] K. Shiohama and M. Tanaka, “An isoperimetric problem for infinitely connected complete open surfaces”, pp. 317–343 in *Geometry of manifolds* (Matsumoto, 1988), edited by K. Shiohama, Academic Press, Boston, 1989. [MR 91b:53049](#) [Zbl 0697.53040](#)
- [Shiohama and Tanaka 1993] K. Shiohama and M. Tanaka, “The length function of geodesic parallel circles”, pp. 299–308 in *Progress in differential geometry*, edited by K. Shiohama, Adv. Stud. Pure Math. **22**, Math. Soc. Japan, Tokyo, 1993. [MR 95b:53054](#) [Zbl 0799.53052](#)
- [Tysk 1987] J. Tysk, “Eigenvalue estimates with applications to minimal surfaces”, *Pacific J. Math.* **128**:2 (1987), 361–366. [MR 88i:53102](#) [Zbl 0594.58018](#)

Received July 23, 2010. Revised March 9, 2011.

PIERRE BÉRARD
UNIVERSITÉ GRENOBLE 1
INSTITUT FOURIER (UJF-CNRS), BP 74
38402 SAINT MARTIN D’HÈRES CEDEX
FRANCE
pierre.berard@ujf-grenoble.fr

PHILIPPE CASTILLON
DÉPARTEMENT DES SCIENCES MATHÉMATIQUES CC 51
UNIVERSITÉ MONTPELLIER II, I3M (UMR 5149)
34095 MONTPELLIER CEDEX 5
FRANCE
cast@math.univ-montp2.fr

MARCOS CAVALCANTE
INSTITUTO DE MATEMÁTICA
UNIVERSIDADE FEDERAL DE ALAGOAS
57072-900 MACEIÓ AL
BRAZIL
marcos.petrucio@pq.cnpq.br

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from [Periodicals Service Company](#), 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 253 No. 1 September 2011

Singularities of the projective dual variety	1
ROLAND ABUAF	
Eigenvalue estimates for hypersurfaces in $\mathbb{H}^m \times \mathbb{R}$ and applications	19
PIERRE BÉRARD, PHILIPPE CASTILLON and MARCOS CAVALCANTE	
Conformal Invariants associated to a measure: Conformally covariant operators	37
SUN-YUNG A. CHANG, MATTHEW J. GURSKY and PAUL YANG	
Compact symmetric spaces, triangular factorization, and Cayley coordinates	57
DEREK HABERMAS	
Automorphisms of the three-torus preserving a genus-three Heegaard splitting	75
JESSE JOHNSON	
The rationality problem for purely monomial group actions	95
HIDETAKA KITAYAMA	
On a Neumann problem with p -Laplacian and noncoercive resonant nonlinearity	103
SALVATORE A. MARANO and NIKOLAOS S. PAPAGEORGIOU	
Minimal ramification in nilpotent extensions	125
NADYA MARKIN and STEPHEN V. ULLOM	
Regularity of weakly harmonic maps from a Finsler surface into an n -sphere	145
XIAOHUAN MO and LIANG ZHAO	
On the sum of powered distances to certain sets of points on the circle	157
NIKOLAI NIKOLOV and RAFAEL RAFAILOV	
Formal geometric quantization II	169
PAUL-ÉMILE PARADAN	
Embedded constant-curvature curves on convex surfaces	213
HAROLD ROSENBERG and MATTHIAS SCHNEIDER	
A topological construction for all two-row Springer varieties	221
HEATHER M. RUSSELL	