

*Pacific
Journal of
Mathematics*

ISOPERIMETRIC SURFACES WITH BOUNDARY, II

ABRAHAM FRANDSEN, DONALD SAMPSON AND NEIL STEINBURG

ISOPERIMETRIC SURFACES WITH BOUNDARY, II

ABRAHAM FRANDBSEN, DONALD SAMPSON AND NEIL STEINBURG

Following our previous work with Dorff and Lawlor, we extend results for the so-called equitent problem of fixed boundary and fixed volume. We define sufficient conditions, which in \mathbb{R}^2 and \mathbb{R}^3 are also necessary, for local minima to be piecewise spherical, and we show that these are area-minimizing in their homotopy class. We also give new examples of these surfaces in \mathbb{R}^2 and \mathbb{R}^3 .

1. Introduction

Equitent problems, first introduced in the paper “Isoperimetric surfaces with boundary” [Dorff et al. 2011], ask what is the area-minimizing surface enclosing a given volume and spanning a given boundary. In this way, equitent problems represent a combination of isoperimetry and boundary conditions, such as in Steiner problems and minimal surfaces. Our previous approach, which we extend here, uses the technique of metacalibration. *Metacalibration* is a version of the calibration methods popularized by Harvey and Lawson [1982], adapted to use on isoperimetric problems. In particular, we use a combination of the mapping of [Gromov 1986], after [Knothe 1957], and the paired calibrations of [Lawlor and Morgan 1994].

In our original results, we construct various classes of surfaces bounded by the dual figures of uniform polytopes and enclosing a prescribed volume and prove that these surfaces are minimizing in their homotopy class. The results, however, turn out to be limited in scope, as shown in [Ross et al. 2011]. Consequently, there remains much ground to be covered.

In this paper, we will extend previous results by considering equitent systems generated by polytopes whose edges are all of a given length. This results in a much wider range of equitent surfaces than those bounded by uniform polytopes. We construct the conjectured minimizing surface using a refinement of previous methods and prove that this surface is indeed area-minimizing in its homotopy class.

Further, we show that any homotopically area-minimizing equitent surface with piecewise spherical faces and simplex vertex figures is equivalent to one generated

MSC2010: primary 53C38; secondary 49Q10.

Keywords: metacalibration, calibration, isoperimetric, equitent.

by our construction. We conclude with a discussion of new equitent surfaces and a survey of open problems.

2. The surfaces


Let Γ be a convex polytope of dimension $m \leq n$ with equal edge lengths, r , embedded in \mathbb{R}^n . Let p_1, \dots, p_k be the vertices of Γ . For each p_i let R_i be the region farthest from p_i :

$$R_i = \{x \in \mathbb{R}^n : r < \|x - p_i\| \text{ and } \|x - p_j\| < \|x - p_i\| \text{ for all } j \neq i\}.$$

Note that if p_i and p_j share an edge in Γ , $\partial R_i \cap \partial R_j$ is a subset of the perpendicular bisecting hyperplane of that edge. Now, define

$$R_0 = \{x \in \mathbb{R}^n : \|x - p_i\| < r \text{ for } 1 \leq i \leq k\}.$$

See figure.



This region represents the enclosed volume. We suppose that $R_0 \neq \emptyset$ and $\mathcal{H}^{n-1}(\partial R_0 \cap \partial R_i) \neq \emptyset$ for all $i > 0$. Let $V_0 = \mathcal{H}^n(R_0)$. Then let

$$M = \bigcup_{i=0}^k \partial R_i.$$

Notice that ∂R_0 is the portion of the surface that encloses the volume R_0 . In order to have a nontrivial result, we require $R_0 \neq \emptyset$. The condition

$$\mathcal{H}^{n-1}(\partial R_0 \cap \partial R_i) \neq \emptyset$$

for all $i > 0$ ensures that all smooth subsurfaces of M meet at 120 degree angles. For $m = 2$, the only viable generating figures, Γ , are equilateral triangles, rhombi with interior angles strictly greater than 60° , and small perturbations of regular pentagons. For $m = 3$, the valid generating figures include all but two of the eight convex deltahedra (polyhedra where all faces are equilateral triangles), as well as

other polytopes with faces of higher degree. It is worth noting, however, that those generating figures, Γ , whose faces are not equilateral triangles produce surfaces which are locally minimal within their homotopy class, but not globally minimal. As will be seen in the proof, this construction gives sufficient conditions for minimizing surfaces to be piecewise spherical. Furthermore, due to the regularity properties of soap films proved by Taylor and Almgren [Taylor 1976], these are also necessary conditions in \mathbb{R}^2 and \mathbb{R}^3 . In higher dimensions nonsimplicial vertex figures may be minimizing, but are not considered in this paper. See for example [Brakke 1991].

3. The minimization theorem

In this section we prove that the surfaces constructed are homotopically minimizing in the following sense: Let U be a bounded open set that contains $\overline{R_0}$, and let $M_0 = M \cap U$.

Theorem 1. *The surface M_0 is area-minimizing among all compact surfaces (rectifiable sets) in U with boundary $\partial U \cap M$ that enclose the fixed volume $\mathcal{H}^n(R_0)$ and are homotopically equivalent to M_0 . This also holds with the weaker assumption that competitor surfaces are not necessarily homotopic to M_0 but separate space into the same regions as M_0 and these regions share boundary nontrivially (on a set of positive \mathcal{H}^{n-1} measure) only if the corresponding regions do in M_0 .*

Our proof uses a metacalibration argument that compares figures according to their flux on specially crafted vector fields. In particular, we use a paired-calibration approach with one vector field defined for each separated region.

Let N be any competitor surface and let S_i be the separated regions that correspond to each R_i respectively. (Then $\mathcal{H}^n(S_0) = \mathcal{H}^n(R_0)$ by the volume condition.) Define $v_i : S_i \rightarrow \mathbb{R}^n$ for $1 \leq i \leq m$ to be the constant vector field $-p_i/r$. Let $\phi : S_0 \rightarrow R_0$ be the Knothe–Rosenblatt rearrangement and let $v_0 : S_0 \rightarrow \mathbb{R}^n$ be given by $v_0 = \phi/r$.

At this point a few simple results would be useful:

Proposition 2. *If S_i and S_j share boundary nontrivially, then $v_i - v_j$ is a unit vector. If $N = M_0$ then $v_i - v_j$ is the unit normal to $\partial R_i \cap \partial R_j$.*

Proof. Note that S_i and S_j share boundary nontrivially if and only if p_i and p_j are adjacent in Γ . Thus $\|v_i - v_j\| = (1/r)\|p_i - p_j\| = 1$. Also if $N = M_0$, $v_i - v_j$ is the unit normal to $\partial R_i \cap \partial R_j$ since $\partial R_i \cap \partial R_j$ lies on the hyperplane equidistant to p_i and p_j . \square

Proposition 3. *The matrix Dv_0 is triangular. If $N = M$ then v_0 is the identity scaled by $1/r$.*

Proof. Follows from the definition of v_0 . See [Dorff et al. 2011] for details. \square

Proposition 4. For $i \neq 0$, $\int_{N \cap \partial S_i} v_i \cdot n \, d\mathcal{H}^{n-1} = \int_{M \cap \partial R_i} v_i \cdot n \, d\mathcal{H}^{n-1}$, where n is the unit normal to the surface of integration, outward pointing with respect to S_i or R_i .

Proof. Follows from the divergence theorem since v_i is divergence free and $\partial(M \cap \partial R_i) = \partial(N \cap \partial S_i)$. \square

Proof of Theorem 1. For any competitor surface N , let $G(N) = \sum_i \int_{N \cap \partial S_i} v_i \cdot n \, d\mathcal{H}^{n-1}$. Letting $P(N) = \sum_i \int_{N \cap \partial S_i} d\mathcal{H}^{n-1}$ be our objective function, we find that

$$\begin{aligned} G(N) &= \sum_i \int_{N \cap \partial S_i} v_i \cdot n \, d\mathcal{H}^{n-1} \\ &= \sum_{i \neq j} \int_{N \cap (\partial S_i \cap \partial S_j)} (v_i - v_j) \cdot n \, d\mathcal{H}^{n-1} \\ &\leq \sum_{i \neq j} \int_{N \cap (\partial S_i \cap \partial S_j)} \|v_i - v_j\| \|n\| \, d\mathcal{H}^{n-1} \\ &\leq \sum_{i \neq j} \int_{N \cap (\partial S_i \cap \partial S_j)} d\mathcal{H}^{n-1} \\ &= \sum_i \int_{N \cap \partial S_i} d\mathcal{H}^{n-1} = P(N), \end{aligned}$$

with equality if $N = M_0$.

Now also note that

$$\begin{aligned} \int_{N \cap \partial S_0} v_0 \cdot n \, d\mathcal{H}^{n-1} &= \int_{S_0} \operatorname{div} v_0 \, d\mathcal{H}^{n-1} \\ &= \int_{S_0} \frac{1}{r} \left(\frac{\partial \phi_1}{\partial x_1} + \frac{\partial \phi_2}{\partial x_2} + \cdots + \frac{\partial \phi_n}{\partial x_n} \right) d\mathcal{H}^{n-1} \\ &\geq \int_{S_0} \frac{n}{r} \sqrt{\frac{\partial \phi_1}{\partial x_1} \frac{\partial \phi_2}{\partial x_2} \cdots \frac{\partial \phi_n}{\partial x_n}} \, d\mathcal{H}^{n-1} \\ &= \int_{S_0} \frac{n}{r} \sqrt[1]{1} \, d\mathcal{H}^{n-1} \\ &= \frac{n}{r} \mathcal{H}^{n-1}(S_0) = \frac{n}{r} \mathcal{H}^{n-1}(R_0), \end{aligned}$$

with equality if $N = M_0$. This follows from the AM-GM inequality and the equality

$$\frac{\partial \phi_1}{\partial x_1} \frac{\partial \phi_2}{\partial x_2} \cdots \frac{\partial \phi_n}{\partial x_n} = \det(D\phi) = 1,$$


which is valid since ϕ is volume-preserving.

Combining these results we find

$$\begin{aligned}
 P(M_0) = G(M_0) &= \sum_{i \neq 0} \int_{M_0 \cap \partial R_i} v_i \cdot n \, d\mathcal{H}^{n-1} + \int_{M_0 \cap \partial R_0} v_0 \cdot n \, d\mathcal{H}^{n-1} \\
 &= \sum_{i \neq 0} \int_{N \cap \partial S_i} v_i \cdot n \, d\mathcal{H}^{n-1} + \frac{n}{r} \mathcal{H}^{n-1}(R_0) \\
 &\leq \sum_{i \neq 0} \int_{N \cap \partial S_i} v_i \cdot n \, d\mathcal{H}^{n-1} + \int_{N \cap \partial S_0} v_0 \cdot n \, d\mathcal{H}^{n-1} \\
 &= G(N) \leq P(N). \quad \square
 \end{aligned}$$

4. Soap films in \mathbb{R}^3

In [Dorff et al. 2011] and [Ross et al. 2011] we identified the regular tetrahedron, the regular octahedron, and the regular icosahedron as polytopes that generate realizable soap films. The only other three-dimensional polytopes, Γ , that generate surfaces realizable as soap films are the triangular dipyramid, the pentagonal dipyramid, and the snub disphenoid. The generated soap films are shown below.



This is due to the conditions proven by Jean Taylor [1976], namely that each face not intersecting with the boundary must meet with exactly two other faces in 120-degree angles. Thus, any generating figure, Γ , with nontriangular faces will not yield a surface realizable as a soap film. The remaining two deltahedra fail to meet the conditions of our construction because of their large circumradius.

Every surface generated by our construction will have piecewise spherical faces and simplicial vertex figures. The converse is also true. Given any area-minimizing equitent surface with piecewise spherical faces and simplicial vertex figures, we can recover the generating polytope, Γ , as the set of centers of each spherical face. In higher dimensions there may exist area-minimizing equitent surfaces with nonsimplicial vertex figures.

5. Conclusion

We have characterized all piecewise spherical equitent surfaces in two and three dimensions, and proven them to be area minimizing. Several interesting and open problems arise. An especially intriguing question deals with equitent surfaces that have negatively curved bubbles, meaning each face of the bubble region bends inward. Such a surface can be created using soap films, but our methods are not yet able to address this case. Similarly, equitent surfaces with nonspherical faces fall outside the scope of our approach. Finally, our construction generates surfaces whose vertices are cones over simplices. In spaces of dimension greater than three, however, minimal surfaces need not have simplicial vertex figures, and we may yet find interesting new equitent surfaces. Extensions of the metacalibration methods outlined in this paper show great promise in solving these open problems.

References

- [Brakke 1991] K. A. Brakke, “Minimal cones on hypercubes”, *J. Geom. Anal.* **1**:4 (1991), 329–338. MR 92k:49082 Zbl 0738.49022
- [Dorff et al. 2011] R. Dorff, D. Johnson, G. R. Lawlor, and D. Sampson, “Isoperimetric surfaces with boundary”, *Proc. Amer. Math. Soc.* **139**:12 (2011), 4467–4473. MR 2012f:53101 Zbl 05988131
- [Gromov 1986] M. Gromov, “Appendix I: Isoperimetric inequalities in Riemannian manifolds”, pp. 114–129 in *Asymptotic theory of finite-dimensional normed spaces*, by V. D. Milman and G. Schechtman, Lecture Notes in Mathematics **1200**, Springer, Berlin, 1986.
- [Harvey and Lawson 1982] R. Harvey and H. B. Lawson, Jr., “Calibrated geometries”, *Acta Math.* **148** (1982), 47–157. MR 666108 (85i:53058) Zbl 0584.53021
- [Knothe 1957] H. Knothe, “Contributions to the theory of convex bodies”, *Michigan Math. J.* **4** (1957), 39–52. MR 18,757b Zbl 0077.35803
- [Lawlor and Morgan 1994] G. Lawlor and F. Morgan, “Paired calibrations applied to soap films, immiscible fluids, and surfaces or networks minimizing other norms”, *Pacific J. Math.* **166**:1 (1994), 55–83. MR 95i:58051 Zbl 0830.49028

[Ross et al. 2011] J. Ross, D. Sampson, and N. Steinburg, “Soap film realization of isoperimetric surfaces with boundary”, *Involve* **4**:3 (2011), 271–276.

[Taylor 1976] J. E. Taylor, “The structure of singularities in soap-bubble-like and soap-film-like minimal surfaces”, *Ann. of Math. (2)* **103**:3 (1976), 489–539. MR 55 #1208a Zbl 0335.49032

Received January 26, 2012.

ABRAHAM FRANDSEN
BRIGHAM YOUNG UNIVERSITY
PROVO, UT 84604
UNITED STATES
abefrandsen@gmail.com

DONALD SAMPSON
BRIGHAM YOUNG UNIVERSITY
PROVO, UT 84604
UNITED STATES
dsampson@byu.net

NEIL STEINBURG
BRIGHAM YOUNG UNIVERSITY
PROVO, UT 84604
UNITED STATES
neil.steinburg@gmail.com

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 259 No. 2 October 2012

Flag subdivisions and γ -vectors	257
CHRISTOS A. ATHANASIADIS	
Rays and souls in von Mangoldt planes	279
IGOR BELEGRADEK, ERIC CHOI and NOBUHIRO INNAMI	
Isoperimetric surfaces with boundary, II	307
ABRAHAM FRANDBSEN, DONALD SAMPSON and NEIL STEINBURG	
Cyclic branched coverings of knots and quandle homology	315
YUICHI KABAYA	
On a class of semihereditary crossed-product orders	349
JOHN S. KAUTA	
An explicit formula for spherical curves with constant torsion	361
DEMETRE KAZARAS and IVAN STERLING	
Comparing seminorms on homology	373
JEAN-FRANÇOIS LAFONT and CHRISTOPHE PITTET	
Relatively maximum volume rigidity in Alexandrov geometry	387
NAN LI and XIAOCHUN RONG	
Properness, Cauchy indivisibility and the Weil completion of a group of isometries	421
ANTONIOS MANOUSSOS and POLYCHRONIS STRANTZALOS	
Theta lifts of strongly positive discrete series: the case of $(\tilde{Sp}(n), O(V))$	445
IVAN MATIĆ	
Tunnel one, fibered links	473
MATT RATHBUN	
Fusion symmetric spaces and subfactors	483
HANS WENZL	



0030-8730(201210)259:2;1-6