

*Pacific  
Journal of  
Mathematics*

**AN EXPLICIT FORMULA FOR  
SPHERICAL CURVES WITH CONSTANT TORSION**

DEMETRE KAZARAS AND IVAN STERLING

Volume 259    No. 2

October 2012


## AN EXPLICIT FORMULA FOR SPHERICAL CURVES WITH CONSTANT TORSION

DEMETRE KAZARAS AND IVAN STERLING

**We give an explicit formula for all curves of constant torsion in the unit two-sphere. Our approach uses hypergeometric functions to solve relevant ordinary differential equations.**

### 1. Introduction

The purpose of this article is to give an explicit formula for all curves of constant torsion  $\tau$  in the unit two-sphere  $S^2(1)$ . These curves and their basic properties have been known since the 1890s, and some of these properties are discussed in the Appendix. Some example curves, computed with a standard ODE package, with  $\tau = 0.1, 0.5, 1, 2$  are shown in Figure 1. Though their existence and some



**Figure 1.** The curves of torsion  $\tau = 0.1, 0.5, 1, 2$  on the unit sphere.

*MSC2010:* 53A04.

*Keywords:* constant torsion curves, spherical curves.

of their general properties were known, our explicit formulas for them, in terms of hypergeometric functions, are new.

Curves of constant torsion are also of interest because all asymptotic curves on pseudospherical surfaces (that is, surfaces in  $\mathbb{R}^3$  with constant negative Gauss curvature) are of constant torsion. Furthermore, any pair of curves with constant torsion  $\pm\tau$ , intersecting at one point, define an essentially unique pseudospherical surface. A complete classification of curves of constant torsion in  $\mathbb{R}^3$ , in the context of integrable geometry, is a work in progress and is related to the corresponding unfinished classification of pseudospherical surfaces.

The authors would like to thank the referees and the editor for their suggestions to improve the original version of this paper.

## 2. General setting

**2.1. General curves in  $\mathbb{R}^3$ .** Let

$$\gamma : (a, b) \longrightarrow \mathbb{R}^3$$

be a regular (that is, nonzero speed)  $C^\infty$  curve in  $\mathbb{R}^3$  with nonzero curvature. The speed  $v$ , curvature  $\kappa$  and torsion  $\tau$  of  $\gamma$  are given by

$$v = \|\gamma'\|, \quad \kappa = \frac{\|\gamma' \times \gamma''\|}{\|\gamma'\|^3}, \quad \tau = \frac{[\gamma' \gamma'' \gamma''']}{\|\gamma' \times \gamma''\|^2}.$$

The unit tangent  $T$  is given by

$$(1) \quad T = \frac{\gamma'}{v}.$$

The unit normal and unit binormal are given by

$$N = \frac{T'}{v\kappa} \quad \text{and} \quad B = T \times N.$$

These are related by the Frenet formulas

$$(2) \quad \begin{aligned} T' &= v\kappa N \\ N' &= -v\kappa T + v\tau B \\ B' &= -v\tau N. \end{aligned}$$

Curves  $\gamma$  with prescribed differentiable curvature  $\kappa > 0$  and torsion  $\tau$  can be found by integrating (2) and (1). Up to reparametrization (see below) a curve in  $\mathbb{R}^3$  is determined, up to a rigid motion of  $\mathbb{R}^3$ , by its curvature  $\kappa$  and torsion  $\tau$ .

**2.2. Changing parametrizations.** Given  $\gamma(t)$ , the arc length function  $s(t)$  of  $\gamma(t)$  is given by

$$s(t) = \int_a^t \|\gamma'(u)\| du.$$

Note that since  $s(t)$  is increasing, it has an inverse  $t(s)$ . To obtain a unit-speed reparametrization  $\gamma^{\text{unit}}$  of  $\gamma$  we let

$$\gamma^{\text{unit}}(s) := \gamma(t(s)).$$

We denote the parameter of a unit-speed curve by the letter  $s$ .

On the other hand, if we are given a unit-speed curve  $\gamma^{\text{unit}}(s)$ , we may wish to find a reparametrization  $\gamma$  of  $\gamma^{\text{unit}}$  by letting  $t(s)$  be some special monotone function. In that case we have

$$\gamma(t) := \gamma^{\text{unit}}(s(t)),$$

where  $s(t)$  is the inverse of  $t(s)$ .

**2.3. Spherical curves.** We call  $\gamma$  a *spherical curve* if  $\gamma(t) \in S^2(r)$  for all  $t$  (for fixed  $r > 0$ ). One can show (see [Gray et al. 2006], for example) that the speed, curvature and torsion of a spherical curve satisfy

$$(3) \quad \kappa^2 \tau^2 (\kappa^2 r^2 - 1) = \kappa'^2 v^2.$$

**2.4. Effect of homothety on curvature and torsion.** If two curves  $\gamma, \tilde{\gamma}$  are related by  $\tilde{\gamma}(t) := \lambda \gamma(t)$ , then  $\tilde{\kappa}(t) = \kappa(t)/\lambda$  and  $\tilde{\tau}(t) = \tau(t)/\lambda$ . Thus, a curve of constant torsion  $\tau_1$  on a sphere of radius  $r_1$  corresponds by homothety to a curve of constant torsion  $\tau_2 = (r_1/r_2) \tau_1$  on a sphere of radius  $r_2$ .

In other words, any spherical curve of constant positive torsion corresponds to precisely one spherical curve with  $\tau = 1$  as well as to precisely one curve of constant positive torsion on the unit sphere. Without loss of generality, we consider only spherical curves of constant positive torsion on the unit sphere.

**2.5. Constant torsion unit-speed curves on the unit sphere.** Let  $r = 1$ . If  $\tau$  is a positive constant and  $\gamma : (a, b) \rightarrow S^2(1)$  is of unit speed, then (3) is an ordinary differential equation in  $\kappa$ :

$$(4) \quad \kappa'^2 = \kappa^2 \tau^2 (\kappa^2 - 1).$$

(Notice that  $\kappa \geq 1$  holds for any curve on the unit sphere.) The general solution to (4) is given by

$$(5) \quad \kappa = \csc(\tau s + C), \quad \frac{-C}{\tau} < s < \frac{-C + \pi}{\tau}.$$

Notice we use the parameter  $s$  instead of  $t$  since  $\gamma$  is a unit-speed curve. Furthermore,  $\kappa(s)$  is decreasing on

$$\left( \frac{-C}{\tau}, \frac{-C + \pi/2}{\tau} \right).$$

**2.6. Our goal.** As mentioned, a unit-speed curve  $\gamma$  is determined up to rigid motion by its curvature and torsion. However, in general it is not possible to explicitly solve for  $\gamma$  given  $\kappa > 0$  and  $\tau > 0$ . Spherical curves of constant torsion provide an interesting and natural example to study. They were considered by classical geometers and the formula (5) was known. Even though the formula for  $\kappa$  is so simple, no explicit solutions for  $\gamma$  were found. This is most likely because the integration methods that we found necessary were not developed until decades later. By choosing a special reparametrization and using functions defined in the 1940s, we were successful in obtaining an explicit formula for  $\gamma$  involving hypergeometric functions.

### 3. Explicit formulas

**3.1. The radius of curvature parametrization.** In curve theory parametrization by the curvature is called the “natural parametrization”. In our case, when the natural parametrization is used, the domain of definition lies outside the radius of convergence of the resulting hypergeometric solutions. To avoid having to deal with the problem of analytically continuing hypergeometric functions beyond their radii of convergence we instead parametrize by the reciprocal of the curvature, which is called the radius of curvature.

We seek unit-speed curves  $\gamma^{\text{unit}} : (-C/\tau, (-C + \frac{\pi}{2})/\tau) \rightarrow S^2(1)$  of constant torsion  $\tau > 0$  on the unit sphere. In order to simplify the Frenet equations, we reparametrize  $\gamma^{\text{unit}}$  by  $t(s) = 1/\kappa(s) = \sin(\tau s + C)$ . Since  $1/\kappa(s)$  is increasing on its domain, the inverse  $s(t), s : (0, 1) \rightarrow (-C/\tau, (-C + \frac{\pi}{2})/\tau)$ , exists and we have

$$\gamma(t) = \gamma^{\text{unit}}(s(t)) = \gamma^{\text{unit}}\left(\frac{\sin^{-1}(t) - C}{\tau}\right), \quad 0 < t < 1.$$

One can recover  $\gamma^{\text{unit}}$  from  $\gamma$  by reversing the process. Note that

$$v = \|\gamma'\| = \|\gamma^{\text{unit}'}\| |s'(t)| = |s'(t)| = \frac{1}{\tau\sqrt{1-t^2}}.$$

With  $\kappa = 1/t$  the Frenet equations (2) become

$$\begin{aligned} (6a) \quad T' &= vN/t \\ (6b) \quad N' &= -vT/t + v\tau B \\ (6c) \quad B' &= -v\tau N. \end{aligned}$$

Recall  $\gamma' = vT$ . Thus as a preliminary step we will compute  $T$ . Namely, we want to solve (6) for  $T$ .

From (6a) and (6b) we have

$$(7) \quad N = t\sqrt{1-t^2}\tau T',$$

$$(8) \quad B = \sqrt{1-t^2}N' + \frac{1}{t\tau}T.$$

Equations (7) and (6c) yield

$$(9) \quad B' = -\tau t T'.$$

On the other hand differentiating (7) yields

$$(10) \quad N' = \frac{\tau}{\sqrt{1-t^2}}(t(1-t^2)T'' + (1-2t^2)T').$$

Plugging this into (8) yields

$$(11) \quad B = \frac{-1}{t\tau}(t^3(t^2-1)\tau^2T'' + t(2t^2-1)\tau^2T' - T).$$

Hence

$$(12) \quad B' = \frac{-1}{t^2\tau}(t^3(t^2-1)\tau^2T''' + t^2(5t^2-2)\tau^2T'' + t(4t^2\tau^2-1)T' + T).$$

Equating (9) and (12) and simplifying we arrive at

$$(13) \quad t^3(t^2-1)\tau^2T''' + t^2(5t^2-2)\tau^2T'' + t(3t^2\tau^2-1)T' + T = 0.$$

This is a third-order linear homogeneous differential equation with nonconstant coefficients. In general it is not possible to find a closed form solution for such an equation. However, this is one of the special cases where one can find hypergeometric type solutions. These methods were developed in the 1940s, and hence were not available to the classical (1890s) geometers.

**3.2. Initial conditions.** To arrive at initial conditions for our ODE, we find initial conditions for  $T$ ,  $N$ , and  $B$  and use the Frenet equations (6) to arrive at initial conditions for  $T$ ,  $T'$ , and  $T''$ . We let  $T = (T_1, T_2, T_3)$ ,  $N = (N_1, N_2, N_3)$ , and  $B = (B_1, B_2, B_3)$ . For  $V = T, N, B, T'$ , or  $T''$  we use the notation  $V_{i_0} := V_i(t_0)$ .

$T$  and  $N$  are unit vectors ( $\|T\| = 1$  and  $\|N\| = 1$ ) so we have

$$\begin{aligned} |T_{1_0}| \leq 1, \quad |T_{2_0}| \leq \sqrt{1-T_{1_0}}, \quad T_{3_0} &= \sqrt{1-T_{1_0}^2-T_{2_0}^2}, \\ |N_{1_0}| \leq 1, \quad |N_{2_0}| \leq \sqrt{1-N_{1_0}}, \quad N_{3_0} &= \sqrt{1-N_{1_0}^2-N_{2_0}^2}. \end{aligned}$$

Also  $T$  is orthogonal to  $N$  ( $T \cdot N = 0$ ),

$$N_{1_0}T_{1_0} + N_{2_0}T_{2_0} + N_{3_0}T_{3_0} = 0.$$

Since  $B = T \times N$ , we have

$$B_{10} = T_{20}N_{30} - T_{30}N_{20}, \quad B_{20} = T_{30}N_{10} - T_{10}N_{30}, \quad B_{30} = T_{10}N_{20} - T_{20}N_{10}.$$

We will, without loss of generality and up to rigid motion, choose  $N_0 = (0, 1, 0)$ ,  $T_0 = (T_{10}, T_{20}, T_{30}) = (1, 0, 0)$ , and  $t_0 = \frac{1}{2}$ . Now that we have initial conditions for  $T$ ,  $N$ , and  $B$ , we will use the Frenet equations to express  $T'_0$  and  $T''_0$  in terms of  $T_0$ ,  $N_0$ , and  $B_0$ ,

$$T'_0 = \frac{v_0 N_0}{t_0}, \quad T''_0 = (v'_0 t_0 + v_0)N_0 - v_0^2 t_0^2 T_0 + v_0^2 \tau t_0 B_0.$$

The set of initial conditions  $t_0 = \frac{1}{2}$ ,  $T_0 = (T_{10}, T_{20}, T_{30}) = (1, 0, 0)$ , and  $N_0 = (0, 1, 0)$  yields

$$T'_0 = \left(0, \frac{4}{\sqrt{3}\tau}, 0\right) \quad \text{and} \quad T''_0 = \left(\frac{-16}{3\tau^2}, \frac{-16}{3\sqrt{3}\tau}, \frac{8}{3\tau}\right).$$

**3.3. Solving for  $T$  via hypergeometric functions.** The Barnes generalized hypergeometric function [Itō 1987] is defined by

$${}_pF_q(a_1, a_2, \dots, a_p; b_1, b_2, \dots, b_q; t^a) := \sum_{n=1}^{\infty} \frac{(a_1)_n \dots (a_p)_n t^{an}}{(b_1)_n \dots (b_q)_n n!}.$$

Note the use of the Pochhammer symbols  $(x)_n := \Gamma(x + n)/\Gamma(x)$ . We will also use

$${}_2F_1^{\text{reg}}(a, b, c, t^a) := \frac{{}_2F_1(a, b; c; t^a)}{\Gamma[c]}.$$

By direct substitution (see for example Section 46 of [Rainville 1971]) it is straightforward to check that the following is a solution to (13):

$$T = (T_1, T_2, T_3), \quad T_j = \sum_{l=1}^3 c_{jl} S_l,$$

where

$$\begin{aligned} S_1 &= it \, {}_3F_2\left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}; \frac{3}{2} - \frac{i}{2\tau}, \frac{3}{2} + \frac{i}{2\tau}; t^2\right), \\ S_2 &= (-1)^{-i/(2\tau)} t^{-i/(2\tau)} \, {}_3F_2\left(1 - \frac{i}{2\tau}, -\frac{i}{2\tau}, -\frac{i}{2\tau}; \frac{1}{2} - \frac{i}{2\tau}, 1 - \frac{i}{\tau}; t^2\right), \\ S_3 &= (-1)^{i/(2\tau)} t^{i/(2\tau)} \, {}_3F_2\left(1 + \frac{i}{2\tau}, \frac{i}{2\tau}, \frac{i}{2\tau}; \frac{1}{2} + \frac{i}{2\tau}, 1 + \frac{i}{\tau}; t^2\right), \end{aligned}$$

the  $c_{jl}$  are constants, and  $i = \sqrt{-1}$ . Note that  $S_1$  is pure imaginary and that  $S_3$  is the complex conjugate of  $S_2$ . For proper complex constants  $c_{jl}$ ,  $T$  is a real valued vector function. By plugging in the initial conditions of the last section we can solve for the  $c_{jl}$ .

**3.4. Solving for  $\gamma$ .** Recall that  $\gamma(t) = \int vT dt$ . Since we have found  $T$  in terms of hypergeometric functions, we must compute the following type of integrals:

$$\int \frac{h(t) {}_pF_q(a_1, a_2, \dots, a_p; b_1, b_2, \dots, b_q; , t^2)}{\tau \sqrt{1-t^2}} dt$$

$$:= \int \frac{h(t) \sum_{n=1}^{\infty} \frac{(a_1)_n \dots (a_p)_n t^{2n}}{(b_1)_n \dots (b_q)_n n!}}{\tau \sqrt{1-t^2}} dt = \frac{\alpha}{\tau} \sum_{n=1}^{\infty} \frac{(a_1)_n \dots (a_p)_n}{(b_1)_n \dots (b_q)_n n!} \int \frac{t^{\beta n}}{\sqrt{1-t^2}} dt.$$

Where  $\alpha, \beta$  are constants. We repeat this process for each  $S_l$ , using the notation  $\gamma = (U_1, U_2, U_3)$ . For  $S_1$ ,

$$U_1 := \int \frac{S_1}{\tau \sqrt{1-t^2}} dt = \int \frac{it {}_3F_2(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}; \frac{3}{2} - \frac{i}{2\tau}, \frac{3}{2} + \frac{i}{2\tau}; t^2)}{\tau \sqrt{1-t^2}} dt$$

$$= \sum_{n=0}^{\infty} d_{1n} \int \frac{t^{2n+1}}{\sqrt{1-t^2}} dt,$$

where

$$d_{1n} = \frac{i(1+\tau^2)\Gamma(\frac{1}{2}+n)^2\Gamma(\frac{3}{2}+n)\operatorname{sech}(\frac{\pi}{2\tau})}{2\sqrt{\pi}\tau^3 n! \Gamma(\frac{3}{2}+n-\frac{i}{2\tau})\Gamma(\frac{3}{2}+n+\frac{i}{2\tau})}.$$

For  $S_2$ ,

$$U_2 := \int \frac{S_2}{\tau \sqrt{1-t^2}} dt$$

$$= \int \frac{(-1)^{-i/(2\tau)} t^{-i/(2\tau)} {}_3F_2(1-\frac{i}{2\tau}, -\frac{i}{2\tau}, -\frac{i}{2\tau}; \frac{1}{2}-\frac{i}{2\tau}, 1-\frac{i}{\tau}; t^2)}{\tau \sqrt{1-t^2}} dt$$

$$= \sum_{n=0}^{\infty} d_{2n} \int \frac{t^{2n-i/\tau}}{\sqrt{1-t^2}} dt,$$

where

$$d_{2n} = \frac{e^{\frac{\pi}{2\tau}} 2^{-\frac{i}{\tau}} \Gamma(n-\frac{i}{2\tau})^2 \Gamma(1+n-\frac{i}{2\tau}) \Gamma(\frac{-i+\tau}{2\tau})^2}{\sqrt{\pi}\tau \Gamma(1+n) \Gamma(1+n-\frac{i}{\tau}) \Gamma(-\frac{i}{2\tau})^2 \Gamma(n+\frac{-i+\tau}{2\tau})}.$$

For  $S_3$ ,

$$U_3 := \int \frac{S_3}{\tau \sqrt{1-t^2}} dt$$

$$= \int \frac{(-1)^{i/(2\tau)} t^{i/(2\tau)} {}_3F_2(1+\frac{i}{2\tau}, \frac{i}{2\tau}, \frac{i}{2\tau}; \frac{1}{2}+\frac{i}{2\tau}, 1+\frac{i}{\tau}; t^2)}{\tau \sqrt{1-t^2}} dt$$

$$= \sum_{n=0}^{\infty} d_{3n} \int \frac{t^{2n+\frac{i}{\tau}}}{\sqrt{1-t^2}} dt,$$



where

$$d_{3n} = \frac{e^{-\frac{\pi}{2\tau}} 2^{\frac{i}{\tau}} \Gamma\left(n + \frac{i}{2\tau}\right)^2 \Gamma\left(1 + n + \frac{i}{2\tau}\right) \Gamma\left(\frac{i+\tau}{2\tau}\right)^2}{\sqrt{\pi} \tau \Gamma(1+n) \Gamma\left(1 + n + \frac{i}{\tau}\right) \Gamma\left(\frac{i}{2\tau}\right)^2 \Gamma\left(n + \frac{i+\tau}{2\tau}\right)}.$$

Once again we are lucky and for each  $U_l$  we can evaluate the integrals. In each case they are hypergeometric,

$$\begin{aligned} U_1 &= \sum_{n=0}^{\infty} \frac{1}{2} n! d_{1n} {}_2F_1^{\text{reg}}\left(\frac{1}{2}, 1+n, 2+n, t^2\right), \\ U_2 &= \sum_{n=0}^{\infty} \frac{1}{2} \Gamma\left(n + \frac{-i+\tau}{2\tau}\right) d_{2n} {}_2F_1^{\text{reg}}\left(\frac{1}{2}, n + \frac{-i+\tau}{2\tau}, \frac{3}{2} + n - \frac{i}{2\tau}, t^2\right), \\ U_3 &= \sum_{n=0}^{\infty} \frac{1}{2} \Gamma\left(n + \frac{i+\tau}{2\tau}\right) d_{3n} {}_2F_1^{\text{reg}}\left(\frac{1}{2}, n + \frac{i+\tau}{2\tau}, \frac{3}{2} + n + \frac{i}{2\tau}, t^2\right). \end{aligned}$$

Each  ${}_2F_1^{\text{reg}}$  also has a power series,


$$\begin{aligned} {}_2F_1^{\text{reg}}\left(\frac{1}{2}, 1+n, 2+n, t^2\right) &= \sum_{m=0}^{\infty} e_{1m} t^{2m}, \\ {}_2F_1^{\text{reg}}\left(\frac{1}{2}, n + \frac{-i+\tau}{2\tau}, \frac{3}{2} + n - \frac{i}{2\tau}, t^2\right) &= \sum_{m=0}^{\infty} e_{2m} t^{2m}, \\ {}_2F_1^{\text{reg}}\left(\frac{1}{2}, n + \frac{i+\tau}{2\tau}, \frac{3}{2} + n + \frac{i}{2\tau}, t^2\right) &= \sum_{m=0}^{\infty} e_{3m} t^{2m}, \end{aligned}$$

where

$$\begin{aligned} e_{1m} &= \frac{\Gamma\left(\frac{1}{2} + m\right)}{(n+m+1)\sqrt{\pi}\Gamma(1+n)\Gamma(1+m)}, \\ e_{2m} &= \frac{2\tau\Gamma\left(\frac{1}{2} + m\right)}{\sqrt{\pi}\Gamma(2n\tau + 2m\tau + \tau - i)\Gamma(1+m)\Gamma\left(n + \frac{-i+\tau}{2\tau}\right)}, \\ e_{3m} &= \frac{2\tau\Gamma\left(\frac{1}{2} + m\right)}{\sqrt{\pi}\Gamma(2n\tau + 2m\tau + \tau + i)\Gamma(1+m)\Gamma\left(n + \frac{i+\tau}{2\tau}\right)}. \end{aligned}$$

Thus

$$\begin{aligned} U_1 &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{1}{2} n! d_{1n} e_{1m} t^{2m+2n+2}, \\ U_2 &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{1}{2} \Gamma\left(n + \frac{-i+\tau}{2\tau}\right) d_{2n} e_{2m} t^{2m+2n+2}, \\ U_3 &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{1}{2} \Gamma\left(n + \frac{i+\tau}{2\tau}\right) d_{3n} e_{3m} t^{2m+2n+2}. \end{aligned}$$



**Figure 2.** The curve of torsion  $\tau = 1$  on the unit sphere.

These complicated double sums combine nicely and simplify as

$$\begin{aligned}
 U_1 &= \frac{i}{2\sqrt{\pi}\tau} \sum_{k=0}^{\infty} \frac{\Gamma(\frac{1}{2} + k)}{\Gamma(2 + k)} \\
 &\quad \times {}_4F_3\left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, -k; \frac{1}{2} - k, \frac{3}{2} - \frac{i}{2\tau}, \frac{3}{2} + \frac{i}{2\tau}; 1\right) t^{2+2k}, \\
 U_2 &= \frac{e^{\pi/(2\tau)}}{\sqrt{\pi}} \sum_{k=0}^{\infty} \frac{\Gamma(\frac{1}{2} + k)}{(-i + (1 + 2k)\tau)\Gamma(1 + k)} \\
 &\quad \times {}_4F_3\left(-k, 1 - \frac{i}{2\tau}, -\frac{i}{2\tau}, -\frac{i}{2\tau}; \frac{1}{2} - k, \frac{1}{2} - \frac{i}{2\tau}, 1 - \frac{i}{\tau}; 1\right) t^{1-i/\tau+2k}, \\
 U_3 &= \frac{e^{-\pi/(2\tau)}}{\sqrt{\pi}} \sum_{k=0}^{\infty} \frac{\Gamma(\frac{1}{2} + k)}{(i + (1 + 2k)\tau)\Gamma(1 + k)} \\
 &\quad \times {}_4F_3\left(-k, 1 + \frac{i}{2\tau}, \frac{i}{2\tau}, \frac{i}{2\tau}; \frac{1}{2} - k, \frac{1}{2} + \frac{i}{2\tau}, 1 + \frac{i}{\tau}; 1\right) t^{1+i/\tau+2k}.
 \end{aligned}$$

Thus we can write  $\gamma = (\gamma_1, \gamma_2, \gamma_3)$  as a power series in  $t$ , where

$$\gamma_j = \sum_{l=1}^3 c_{jl} U_l.$$

The curve with  $\tau = 1$  is given in Figure 2, this time using the explicit formula.

### Appendix

The purpose of this appendix is to address questions and issues about the curves raised by the referees and the editor.

For  $\tau = 0$ , the curves are also planar and are precisely the set of circles lying on the sphere. If we consider curves corresponding to solutions (5) with  $C = 0$ , and  $k(\frac{\pi}{2}/\tau) = \csc((\frac{\pi}{2}/\tau)\tau) = 1$ , then as  $\tau$  varies from 0 to  $\infty$  the curves numerically appear to vary (in a nonuniform way) from an infinitely covered great circle, through a family of spiral “clothoid” like curves.



**Figure 3.** Curves of constant torsion approaching a circle.

The editor pointed out that if we consider those solutions to (5) with  $C = 0$ , and  $k(s_0) = \csc(s_0\tau) > 1$ , then the corresponding curves approach a “small circle” on  $S^2(1)$  of constant curvature  $k(s_0)$ . This is an interesting example of nonuniform convergence. The curves as a whole converge pointwise to an infinitely covered great circle, while it is still possible to find sequences of “tails” that converge to infinitely covered small circles. This phenomenon is indicated in Figure 3, where one sees a sequence of curves converging to a small circle. More details of this simple yet interesting behavior will be written up elsewhere.

We will mention a few of the qualitative properties of these curves. Let us consider the case of curves in  $S^2(1)$  with a fixed initial point and varying  $\tau$ . All curves of constant torsion differ from one of these by a rigid motion. In [Cesàro 1926, page 185], it is shown that the curves are embedded, spiral infinitely often about a limiting endpoint, and are reflectionally symmetric through the initial point. (In Figure 1 we show only the upper half of the curves.) Cesàro [1926, page 185] also shows that as  $\tau$  varies from 0 to  $\infty$ , the length varies from  $\infty$  to 0; see also Equation (5).

One referee asked if it would be possible to foliate  $S^2(1)$  with curves of constant torsion (other than by the just using circles). It may be possible to foliate  $S^2(1)$ , in some convoluted way, by packing  $S^2(1)$  with pieces of curves of constant torsion; however our conjecture would be that it is not possible to foliate it in any “reasonable” way.

The reasoning is as follows. It seems to be a difficult problem to find an explicit formula for the upper endpoint in terms of the  $\tau$  and the initial point. Nevertheless, numerically as  $\tau$  varies from 0 to  $\infty$  the upper endpoint steadily moves downward from the north pole to the initial point. In particular this would imply that the curves corresponding to an infinitesimal change in  $\tau$  would (repeatedly) intersect. It would follow that any foliation of the  $S^2(1)$  by curves of constant torsion would have to include curves with common endpoints that differ by a rigid motion; a

rotation about the upper endpoint. This type of foliation could only work in some radius about the upper endpoint, because the effect of a rotation on the opposite lower endpoint would result in (repeated) intersections. In summary, the numerics strongly indicate that there is no foliation (singular or not) of  $S^2(1)$  by curves of constant torsion.

Weiner [1977] proved that there exist arbitrarily short closed constant torsion curves in  $\mathbb{R}^3$ . More recently, Musso [2001] studied those curves of constant torsion in  $\mathbb{R}^3$  whose normal vectors sweep out elastic curves in  $S^2(1)$ . Ivey [2000] generalizes Musso's results and gives examples of closed constant torsion curves of various knot types. The examples in the current paper complement these known examples.

### References

- [Cesàro 1926] E. Cesàro, *Vorlesungen über natürlicher Geometrie*, 2nd ed., Teubner, Leipzig, 1926. JFM 52.0686.03
- [Gray et al. 2006] A. Gray, E. Abbena, and S. Salamon, *Modern differential geometry of curves and surfaces with Mathematica*, 3rd ed., Chapman & Hall/CRC, Boca Raton, 2006. MR 2007d:53001 Zbl 0212.31802
- [Itō 1987] K. Itō (editor), *Encyclopedic dictionary of mathematics*, 2nd ed., MIT Press, Cambridge, MA, 1987. MR 89b:00033 Zbl 0674.00025
- [Ivey 2000] T. A. Ivey, "Minimal curves of constant torsion", *Proc. Amer. Math. Soc.* **128**:7 (2000), 2095–2103. MR 2001j:58021 Zbl 0954.53004
- [Musso 2001] E. Musso, "Elastic curves and the Delaunay problem for curves with constant torsion", *Rend. Circ. Mat. Palermo (2)* **50**:2 (2001), 285–298. MR 2002k:58028 Zbl 1072.37043
- [Rainville 1971] E. D. Rainville, *Special functions*, Chelsea Publishing, New York, 1971. MR 52 #14399 Zbl 0231.33001
- [Weiner 1977] J. L. Weiner, "Closed curves of constant torsion, II", *Proc. Amer. Math. Soc.* **67**:2 (1977), 306–308. MR 57 #1370 Zbl 0382.53001

### Editor's note

I was pleased to receive this interesting and unusual new work submitted to the *Pacific Journal of Mathematics*. It has also found referee approval for the main substance of the new material it offers. I am however uneasy with some assertions in the Appendix; to the best I can interpret them, I find myself in some disagreement. The authors have indicated that they prefer not to alter the statements, and have correctly pointed out that the impact of the appendix material on the main thrust of the paper is peripheral. In the view that differences over ancillary matters should not impede publication of new discoveries, I have recommended publication coupled with this subordinate comment.

Robert Finn

Received October 22, 2011. Revised July 3, 2012.

DEMETRE KAZARAS  
DEPARTMENT OF MATHEMATICAL SCIENCES  
UNIVERSITY OF OREGON  
FENTON HALL  
EUGENE, OR 97403-1222  
UNITED STATES  
dkazaras@gmail.com

IVAN STERLING  
ST. MARY'S COLLEGE OF MARYLAND  
18952 E. FISHER ROAD  
ST. MARY'S CITY, MD 20686-3001  
UNITED STATES  
isterling@smcm.edu

# PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by  
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pacific@math.ucla.edu](mailto:pacific@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Darren Long  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[long@math.ucsb.edu](mailto:long@math.ucsb.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Alexander Merkurjev  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[merkurev@math.ucla.edu](mailto:merkurev@math.ucla.edu)

Jonathan Rogawski  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[jonr@math.ucla.edu](mailto:jonr@math.ucla.edu)

## PRODUCTION

[pacific@math.berkeley.edu](mailto:pacific@math.berkeley.edu)

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [pacificmath.org](http://pacificmath.org) for submission instructions.

---

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2012 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 259    No. 2    October 2012

---

Flag subdivisions and $\gamma$ -vectors	257
CHRISTOS A. ATHANASIADIS	
Rays and souls in von Mangoldt planes	279
IGOR BELEGRADEK, ERIC CHOI and NOBUHIRO INNAMI	
Isoperimetric surfaces with boundary, II	307
ABRAHAM FRANDBSEN, DONALD SAMPSON and NEIL STEINBURG	
Cyclic branched coverings of knots and quandle homology	315
YUICHI KABAYA	
On a class of semihereditary crossed-product orders	349
JOHN S. KAUTA	
An explicit formula for spherical curves with constant torsion	361
DEMETRE KAZARAS and IVAN STERLING	
Comparing seminorms on homology	373
JEAN-FRANÇOIS LAFONT and CHRISTOPHE PITTET	
Relatively maximum volume rigidity in Alexandrov geometry	387
NAN LI and XIAOCHUN RONG	
Properness, Cauchy indivisibility and the Weil completion of a group of isometries	421
ANTONIOS MANOUSSOS and POLYCHRONIS STRANTZALOS	
Theta lifts of strongly positive discrete series: the case of $(\tilde{Sp}(n), O(V))$	445
IVAN MATIĆ	
Tunnel one, fibered links	473
MATT RATHBUN	
Fusion symmetric spaces and subfactors	483
HANS WENZL	



0030-8730(201210)259:2;1-6