

*Pacific
Journal of
Mathematics*

SINGULARITIES OF FREE GROUP CHARACTER VARIETIES

CARLOS FLORENTINO AND SEAN LAWTON

Volume 260 No. 1

November 2012

SINGULARITIES OF FREE GROUP CHARACTER VARIETIES

CARLOS FLORENTINO AND SEAN LAWTON

Let \mathfrak{X}_r be the moduli space of SL_n , SU_n , GL_n , or U_n -valued representations of a rank r free group. We classify the algebraic singular stratification of \mathfrak{X}_r . This comes down to showing that the singular locus corresponds exactly to reducible representations if there exist singularities at all. Then by relating algebraic singularities to topological singularities, we show the moduli spaces \mathfrak{X}_r generally are not topological manifolds, except for a few examples we explicitly describe.

1. Introduction

During the last few decades, character varieties have played important roles in knot theory, hyperbolic geometry, Higgs and vector bundle theory, and quantum field theory. However, many of their fundamental properties and structure are not completely understood.

In this article, we first classify the (algebraic) singular locus of SL_n and GL_n -character varieties of free groups by relating the existence of a singularity with the reducibility of the corresponding representation. We then classify all such character varieties that arise as manifolds by explicitly describing the topological neighborhoods of generic singularities. The results we obtain do not necessarily extend to general G -character varieties of finitely generated groups Γ , if G is not one of SL_n , SU_n , GL_n , or U_n and Γ is not free; explicit counterexamples can be obtained via methods different from those considered in this paper (see Section 3I). Our first main theorem generalizes results in [Heusener and Porti 2004], and our second main theorem generalizes results in [Bratholdt and Cooper 2001]. They may be described more precisely as follows.

Let F_r be a rank r free group and let G be a reductive complex algebraic group with K a maximal compact subgroup (see Section 2). Let $\mathfrak{R}_r(G) = \mathrm{Hom}(F_r, G)$ and $\mathfrak{R}_r(K) = \mathrm{Hom}(F_r, K)$ be varieties of representations, and let G , respectively

This work was partially supported by the Center for Mathematical Analysis, Geometry and Dynamical Systems at I.S.T., and by the Fundação para a Ciência e a Tecnologia through the programs Praxis XXI, POCI/MAT/58549/2004 and FEDER..

MSC2010: primary 14B05, 32S05; secondary 14L30, 20CXX, 14P10.

Keywords: character variety, singularities, local topology.

K , act by conjugation on these representation spaces.

Consider the space $\mathfrak{X}_r(K) := \mathfrak{R}_r(K)/K$ which is the conjugation orbit space of $\mathfrak{R}_r(K)$ where $\rho \sim \psi$ if and only if there exists $k \in K$ so $\rho = k\psi k^{-1}$. Let $\mathbb{C}[\mathfrak{R}_r(G)]$ be the affine coordinate ring of $\mathfrak{R}_r(G)$ and let $\mathbb{C}[\mathfrak{R}_r(G)]^G$ be the subring of G -conjugation invariants. Then define

$$\mathfrak{X}_r(G) := \text{Spec}_{\max}(\mathbb{C}[\mathfrak{R}_r(G)]^G),$$

which parametrizes unions of conjugation orbits where two orbits are in the same union if and only if their closures have a nonempty intersection.

The space $\mathfrak{X}_r(G)$, called the G -character variety of F_r , is a complex affine variety and so has a well-defined (algebraic) singular locus (a proper subvariety) which we denote by $\mathfrak{X}_r(G)^{\text{sing}}$. Similarly, $\mathfrak{X}_r(K)$ is a semialgebraic set and so has a real algebraic coordinate ring which likewise determines an algebraic singular locus $\mathfrak{X}_r(K)^{\text{sing}}$. For simplicity, despite the fact it is generally not an algebraic set, we will also refer to $\mathfrak{X}_r(K)$ as a character variety.

We will be mainly concerned with the cases when G is the general linear group GL_n or the special linear group SL_n (over \mathbb{C}), for which K is the unitary group U_n or the special unitary group SU_n , respectively. In these cases a representation ρ is called *irreducible* if with respect to the standard action of G , respectively K , on \mathbb{C}^n the induced action of $\rho(F_r)$ does not have any nontrivial proper invariant subspaces. Otherwise ρ is called *reducible*. This allows one to define the sets $\mathfrak{X}_r(G)^{\text{red}}$ and $\mathfrak{X}_r(K)^{\text{red}}$ which correspond to the spaces of equivalence classes in $\mathfrak{X}_r(G)$, respectively $\mathfrak{X}_r(K)$, that have a representative which is reducible.

In Section 2, we show that the (algebraic) singular locus of $\mathfrak{X}_r(\text{SL}_n)$ and $\mathfrak{X}_r(\text{GL}_n)$ respectively determines the (algebraic) singular locus of $\mathfrak{X}_r(\text{SU}_n)$ and $\mathfrak{X}_r(\text{U}_n)$. We then show $\mathfrak{X}_r(\text{SL}_n) \subset \mathfrak{X}_r(\text{GL}_n)$ has its singular locus determined by the singular locus of $\mathfrak{X}_r(\text{GL}_n)$. This reduces the classification of the singular loci of these four families of moduli spaces to $\mathfrak{X}_r(\text{GL}_n)$ alone. We end Section 2 with examples of $\mathfrak{X}_r(G)$ that are homeomorphic to manifolds with boundary; we conjectured in [Florentino and Lawton 2009] that these were the *only* examples.

It is straightforward to establish that $\mathfrak{X}_1(\text{SL}_n) \cong \mathbb{C}^{n-1}$ and $\mathfrak{X}_2(\text{SL}_2) \cong \mathbb{C}^3$ are affine spaces and so smooth, and $\mathfrak{X}_1(\text{SL}_n)^{\text{red}} = \mathfrak{X}_1(\text{SL}_n)$. In [Heusener and Porti 2004], it is shown that $\mathfrak{X}_r(\text{SL}_2)^{\text{sing}} = \mathfrak{X}_r(\text{SL}_2)^{\text{red}}$ for $r \geq 3$. More generally, one can establish that all irreducible representations in SL_n -character varieties of free groups are in fact smooth; that is $\mathfrak{X}_r(\text{SL}_n)^{\text{sing}} \subset \mathfrak{X}_r(\text{SL}_n)^{\text{red}}$. In [Lawton 2007] it is shown that the singular locus of $\mathfrak{X}_2(\text{SL}_3)$ corresponds exactly to the set of equivalence classes of reducible representations; that is, $\mathfrak{X}_2(\text{SL}_3)^{\text{red}} = \mathfrak{X}_2(\text{SL}_3)^{\text{sing}}$. These examples generalize to our first main result:

Theorem 1.1. *Let $r, n \geq 2$. Let G be SL_n or GL_n and K be SU_n or U_n . Then $\mathfrak{X}_r(G)^{\text{red}} = \mathfrak{X}_r(G)^{\text{sing}}$ and $\mathfrak{X}_r(K)^{\text{red}} = \mathfrak{X}_r(K)^{\text{sing}}$ if and only if $(r, n) \neq (2, 2)$.*

In fact we are able to use an induction argument to completely classify the singular stratification of these semialgebraic spaces. The proof and development of this result constitutes Section 3, including a brief review of a weak version of the celebrated Luna slice theorem.

Theorem 1.1 is sharper than it might appear at first. Replacing F_r by a general finitely presented group Γ one can find examples where irreducibles are singular and examples where reducibles are smooth. On the other hand, changing G to a general reductive complex algebraic group, we find there are examples where irreducibles are singular. In Section 3I, we discuss this in further detail.

A locally Euclidean Hausdorff space M with a countable basis is called a *topological manifold*. More generally, if the neighborhoods are permitted to be Euclidean half-spaces then M is said to be a *topological manifold with boundary*. In [Florentino and Lawton 2009] we determined the homeomorphism type of $\mathfrak{X}_r(\mathrm{SU}_n)$ in the cases $(r, n) = (r, 1), (1, n), (2, 2), (2, 3)$, and $(3, 2)$ where we showed all were topological manifolds with boundary; this is reviewed in Section 2B. In [Bratholdt and Cooper 2001] it is established that the $\mathfrak{X}_r(\mathrm{SU}_2)$ are not topological manifolds when $r \geq 4$.

Motivated by this we conjectured in [Florentino and Lawton 2009] that the examples computed there are the only cases where a topological manifold with boundary arise. Our second main theorem in this paper establishes that conjecture:

Theorem 1.2. *Let $r, n \geq 2$. Let G be SL_n or GL_n and K be SU_n or U_n . $\mathfrak{X}_r(G)$ is a topological manifold with boundary if and only if $(r, n) = (2, 2)$. $\mathfrak{X}_r(K)$ is a topological manifold with boundary if and only if $(r, n) = (2, 2), (2, 3)$, or $(3, 2)$.*

Theorem 1.1 and the observation that the reducible locus is nonempty for $n \geq 2$, does not immediately imply Theorem 1.2 since algebraic singularities may or may not be an obstruction to the existence of a Euclidean neighborhood (topological singularities). For example, both the varieties given by $xy = 0$ and $y^2 = x^3$ in \mathbb{C}^2 (or \mathbb{R}^2) are (algebraically) singular at the point $(0, 0)$ but only the latter has a Euclidean neighborhood at the origin. So, only the former is topologically singular. The variety $xy = 0$ is reducible; an example of an irreducible variety that has an algebraic singularity that is also a topological singularity is the affine cone over $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ discussed in Section 3E.

The proof of Theorem 1.2 constitutes Section 4. To prove our main theorems we use slice theorems and explicitly describe the homeomorphism type of neighborhoods (showing them to be non-Euclidean) for a family of examples. It is interesting to note that since $\mathfrak{X}_r(\mathrm{SL}_n)$ deformation retracts to $\mathfrak{X}_r(\mathrm{SU}_n)$, by [Florentino and Lawton 2009], it must be the case that for $(r, n) = (2, 3)$ and $(3, 2)$ the non-Euclidean neighborhoods deformation retract to Euclidean neighborhoods. Curiously, these are the only cases ($n \geq 2$) where $\mathfrak{X}_r(\mathrm{SU}_n)$ is a topological manifold, and both are homeomorphic to spheres (see [Florentino and Lawton 2009] or Section 2B).

2. Character varieties

Let G be a complex affine reductive algebraic group and let K be a maximal compact subgroup. Then, $G = K_{\mathbb{C}}$ is the complexification of K (the set complex zeros of K as a real algebraic set). For instance, $K_{\mathbb{C}} = \mathrm{SL}_n$ is the complexification of $K = \mathrm{SU}_n$, and $K_{\mathbb{C}} = \mathrm{GL}_n$ is the complexification of $K = \mathrm{U}_n$.

Let Γ be a finitely generated group and let $\mathfrak{R}_{\Gamma}(G) = \mathrm{Hom}(\Gamma, G)$ be the G -valued representations of Γ . We call $\mathfrak{R}_{\Gamma}(G)$ the G -representation variety of Γ , although it is generally only an affine algebraic set.

In the category of affine varieties, $\mathfrak{R}_{\Gamma}(G)$ has a quotient by the conjugation action of G , a regular action, given by $\rho \mapsto g\rho g^{-1}$. This quotient is realized as $\mathfrak{X}_{\Gamma}(G) = \mathrm{Spec}_{\max}(\mathbb{C}[\mathfrak{R}_{\Gamma}(G)]^G)$, where $\mathbb{C}[\mathfrak{R}_{\Gamma}(G)]^G$ is the subring of invariant polynomials in the affine coordinate ring $\mathbb{C}[\mathfrak{R}_{\Gamma}(G)]$. We call $\mathfrak{X}_{\Gamma}(G)$ the G -character variety of Γ . Concretely, it parametrizes unions of conjugation orbits where two orbits are in the same union if and only if their closures intersect nontrivially. Within each union of orbits, denoted $[\rho]$ and called an extended orbit equivalence class, there is a unique closed orbit (having minimal dimension). Any representative from this closed orbit is called a *polystable point*. For SL_n and GL_n the polystable points will have the property that with respect to the action of $\rho(\Gamma)$ on \mathbb{C}^n , they are completely reducible; that is, each decomposes into a finite direct sum of irreducible subactions (on nonzero subspaces).

Let $F_r = \langle x_1, \dots, x_r \rangle$ be a rank r free group. The G -representation variety of F_r , and the G -character variety of F_r will simply be denoted by $\mathfrak{R}_r(G)$ and $\mathfrak{X}_r(G)$, respectively. The evaluation mapping $\mathfrak{R}_r(G) \rightarrow G^r$ defined by sending $\rho \mapsto (\rho(x_1), \dots, \rho(x_r))$ is a bijection and since G is a smooth affine variety, $\mathfrak{R}_r(G)$ naturally inherits the structure of a smooth affine variety as well. Note that we are not assuming that an algebraic variety is irreducible. Whenever G is an irreducible algebraic set however, $\mathfrak{R}_r(G)$ is irreducible, and consequently $\mathfrak{X}_r(G)$ is irreducible as well.

Since an algebraic reductive group over \mathbb{C} is always linear, we can assume that G is a subgroup of GL_N , for some N , and hence $\mathfrak{R}_r(G) \subset \mathbb{C}^{rN^2}$. So, $\mathfrak{R}_r(G)$ inherits the *ball topology*. Given a set of generators f_1, \dots, f_k of the ring of invariants $\mathbb{C}[\mathfrak{R}_r(G)]^G$, $\mathfrak{X}_r(G)$ also inherits the ball topology from the embedding of $\mathfrak{X}_r(G)$ into \mathbb{C}^k given by $[\rho] \mapsto (f_1(\rho), \dots, f_k(\rho))$. In this topology $\mathfrak{X}_r(G)$ is Hausdorff and has a countable basis. Although the ball topology is dependent on an embedding a priori, an affine embedding corresponds exactly to a set of generators for the associated ring, but all choices result in the same homeomorphism type, so the ball topology is intrinsic. Also, in the ball topology, at each point in $\mathfrak{X}_r(G)$ there is a neighborhood homeomorphic to a real cone over a space with Euler characteristic 0 [Sullivan 1971].

Given a compact Lie group K , for brevity we also call the orbit space $\mathfrak{X}_r(K) = \mathfrak{R}_r(K)/K$ a K -character variety of F_r . Note however that $\mathfrak{X}_r(K)$ is generally only a semialgebraic set, and does not equal, in general, the set of real points of a complex variety. In this case, the topology, also Hausdorff with a countable basis, is the quotient topology. $\mathfrak{X}_r(K)$ is compact since K is compact. Likewise, it is path-connected whenever K is path-connected.

Definition 2.1. Let $\rho : \Gamma \rightarrow G$ be a representation into a reductive complex algebraic group. If the image of ρ does not lie in a parabolic subgroup of G , then ρ is called *irreducible*. If, for every parabolic P containing $\rho(\Gamma)$ there is a Levi factor $L \subset P$ such that $\rho(\Gamma) \subset L$, then ρ is called *completely reducible*.

For SL_n and GL_n the irreducible representations are exactly those that, with respect to their actions on \mathbb{C}^n , do not admit any proper (nontrivial) invariant subspaces. Any representation that is not irreducible is called *reducible*. Denote the set of reducible representations by $\mathfrak{R}_\Gamma(G)^{\text{red}}$. A point is called *stable* if the stabilizer is finite and if the orbit is closed.

The following theorem can be found in [Sikora 2012], building on earlier work in [Johnson and Millson 1987, pp. 54–57]. Let $PG = G/Z(G)$ where $Z(G)$ is the center. Note that the action of PG and G define the same GIT quotients and the same orbit spaces and thus, since the PG action is effective, we will sometimes consider this action.

Theorem 2.2 [Johnson and Millson 1987; Sikora 2012]. *Let G be reductive. The irreducibles are exactly the stable points under the action of PG on $\mathfrak{R}_\Gamma(G)$. The completely reducibles are the polystable points.*

Definition 2.3. The reducibles $\mathfrak{X}_\Gamma(G)^{\text{red}}$ are the image of the projection

$$\mathfrak{R}_\Gamma(G)^{\text{red}} \subset \mathfrak{R}_\Gamma(G) \longrightarrow \mathfrak{X}_\Gamma(G).$$

Since $\mathfrak{R}_r(G) \cong G^r$ all points are smooth, and since $\mathfrak{X}_r(G)$ is an affine quotient of a reductive group, there exists $\rho^{\text{ss}} \in [\rho]$ which has a closed orbit and corresponds to a completely reducible representation. Thus, for G either SL_n or GL_n we can assume it is in block diagonal form. In other words, $\rho^{\text{ss}} \leftrightarrow (X_1, \dots, X_r)$ where X_i all have the same block diagonal form (if they are irreducible then there would be only one block). These representations induce a semisimple module structure on \mathbb{C}^n . We denote the set of semisimple representations by $\mathfrak{R}_r(G)^{\text{ss}}$. We note that $\mathfrak{R}_r(G)^{\text{ss}}/G \cong \mathfrak{X}_r(G)$ since all extended orbits have a semisimple representative, and that the semisimple representations are also the completely reducible representations which are also the polystable representations. Likewise, we denote the irreducible representations (those giving simple actions on \mathbb{C}^n) by $\mathfrak{R}_r(G)^s$ and their quotient by $\mathfrak{X}_r(G)^s$.

2A. The determinant fibration. In order to compare SL_n -character varieties to GL_n -character varieties, the following setup will be useful. The usual exact sequence of groups given by the determinant of an invertible matrix

$$(1) \quad SL_n \rightarrow GL_n \xrightarrow{\det} \mathbb{C}^*$$

induces (by fixing generators of F_r , as before) what we will call the *determinant map*:

$$\det : \mathfrak{X}_r(GL_n) \rightarrow \text{Hom}(F_r, \mathbb{C}^*) \cong (\mathbb{C}^*)^r, \quad [\rho] \mapsto \det(\rho),$$

where $\det(\rho) = (\det(X_1), \dots, \det(X_r))$, for $\rho = (X_1, \dots, X_r) \in \mathfrak{A}_r(GL_n)$. Note that the map is clearly well-defined on conjugation classes. Considering the algebraic torus $(\mathbb{C}^*)^r = \text{Hom}(F_r, \mathbb{C}^*) = \mathfrak{X}_r(\mathbb{C}^*)$ as an algebraic group (with identity $\mathbf{1} = (1, \dots, 1)$ and componentwise multiplication) it is immediate that the SL_n -character variety is the “kernel” of the determinant map, $\mathfrak{X}_r(SL_n) = \det^{-1}(\mathbf{1})$. Therefore, the sequence (1) induces another exact sequence

$$(2) \quad \mathfrak{X}_r(SL_n) \rightarrow \mathfrak{X}_r(GL_n) \xrightarrow{\det} (\mathbb{C}^*)^r.$$

In this way, SL_n -character varieties appear naturally as subvarieties of GL_n -character varieties.

Note also that $\mathfrak{X}_r(GL_n)$ can be viewed as a $\mathfrak{X}_r(\mathbb{C}^*)$ -space, as it admits a well-defined action of this torus. That is, we can naturally define $\rho \cdot \lambda \in \mathfrak{X}_r(GL_n)$, given $\rho \in \mathfrak{X}_r(GL_n)$ and $\lambda \in \mathfrak{X}_r(\mathbb{C}^*)$. Given that $\text{PSL}_n = GL_n // \mathbb{C}^*$, it is easy to see that the corresponding quotient is the PSL_n -character variety:

$$\mathfrak{X}_r(\text{PSL}_n) = \mathfrak{X}_r(GL_n) // \mathfrak{X}_r(\mathbb{C}^*).$$

Also, GL_n^r is a quasiaffine subvariety of $\mathfrak{gl}(n, \mathbb{C})^r$. In fact, it is the principal open set defined by the product of the determinants of generic matrices. Since the determinant is an invariant function and taking invariants commutes with localizing at those invariants, we have

$$\mathbb{C}[GL_n^r]^{GL_n} \approx \mathbb{C}[\mathfrak{gl}(n, \mathbb{C})^r // GL_n] \left[\frac{1}{\det(X_1) \cdots \det(X_r)} \right],$$

where the expression on the right is the localization at the product of determinants.

We now prove how the fixed determinant character varieties, complex and compact, relate to the nonfixed determinant character varieties. Identify the cyclic group of order n , $\mathbb{Z}_n := \mathbb{Z}/n\mathbb{Z}$, with $Z(SL_n) \cong Z(SU_n)$.

Theorem 2.4. *The following are isomorphisms of semialgebraic sets:*

- (i) $\mathfrak{X}_r(GL_n) \cong \mathfrak{X}_r(SL_n) \times_{\mathfrak{X}_r(\mathbb{Z}_n)} \mathfrak{X}_r(GL_1)$.
- (ii) $\mathfrak{X}_r(U_n) \cong \mathfrak{X}_r(SU_n) \times_{\mathfrak{X}_r(\mathbb{Z}_n)} \mathfrak{X}_r(U_1)$.

Proof. We first note that $\mathfrak{X}_r(U_1) \cong (S^1)^r$ and $\mathfrak{X}_r(\mathrm{GL}_1) \cong (\mathbb{C}^*)^r$, and thus $\mathfrak{X}_r(\mathbb{Z}_n) \cong \mathbb{Z}_n^r$, as the groups involved are abelian.

The determinant map (1) defines a principal SL_n -bundle $\mathrm{SL}_n \hookrightarrow \mathrm{GL}_n \rightarrow \mathbb{C}^*$, which also expresses $\mathrm{GL}_n \cong \mathrm{SL}_n \rtimes \mathbb{C}^*$ as a semidirect product since there exists a homomorphic section (SL_n is a normal subgroup).

Let \mathbb{Z}_n correspond to n -th roots of unity $\omega_k = e^{\frac{2\pi i k}{n}}$. As algebraic sets one can show directly, by the mapping $(A, \lambda) \mapsto \lambda A$, that $\mathrm{GL}_n \cong (\mathrm{SL}_n \times \mathbb{C}^*) // \mathbb{Z}_n := \mathrm{SL}_n \times_{\mathbb{Z}_n} \mathbb{C}^*$ where \mathbb{Z}_n acts by $\omega_k \cdot (g, \lambda) = (g\omega_k, \omega_k^{-1}\lambda)$ and \mathbb{C}^* is the center of GL_n . This implies that, as algebraic sets,

$$\begin{aligned} (3) \quad \mathfrak{X}_r(\mathrm{GL}_n) &\cong ((\mathrm{SL}_n \times \mathbb{C}^*) // \mathbb{Z}_n)^r // \mathrm{SL}_n \\ &\cong ((\mathrm{SL}_n^r \times (\mathbb{C}^*)^r) // \mathbb{Z}_n^r) // \mathrm{SL}_n \\ &\cong ((\mathrm{SL}_n^r \times (\mathbb{C}^*)^r) // \mathrm{SL}_n) // \mathbb{Z}_n^r \cong \mathfrak{X}_r(\mathrm{SL}_n) \times_{\mathbb{Z}_n^r} (\mathbb{C}^*)^r, \end{aligned}$$

since the action of \mathbb{Z}_n^r commutes with the action of SL_n which is trivial on $(\mathbb{C}^*)^r$.

In the same way we obtain the other “twisted product” isomorphism $\mathfrak{X}_r(U_n) \cong \mathfrak{X}_r(\mathrm{SU}_n) \times_{\mathbb{Z}_n^r} (S^1)^r$. \square

This result provides an explicit way to write $\mathfrak{X}_r(\mathrm{GL}_n)$ as a $\mathfrak{X}_r(\mathrm{SL}_n)$ -bundle over the algebraic r -torus $(\mathbb{C}^*)^r$ and $\mathfrak{X}_r(U_n)$ as a $\mathfrak{X}_r(\mathrm{SU}_n)$ -bundle over the geometric r -torus $(S^1)^r$.

There are a number of consequences to Theorem 2.4.

Corollary 2.5. $\mathfrak{X}_r(U_n)$, respectively $\mathfrak{X}_r(\mathrm{GL}_n)$, is a manifold whenever $\mathfrak{X}_r(\mathrm{SU}_n)$, respectively $\mathfrak{X}_r(\mathrm{SL}_n)$, is a manifold.

Proof. The action of \mathbb{Z}_n^r is free and proper. \square

Corollary 2.6. $\mathfrak{X}_r(\mathrm{GL}_n)$ and $\mathfrak{X}_r(\mathrm{SL}_n) \times (\mathbb{C}^*)^r$ are étale equivalent.

Proof. First note that $\mathrm{SL}_n^r \times (\mathbb{C}^*)^r$ is smooth and hence a normal variety. This implies (see [Drézet 2004]) that $(\mathrm{SL}_n^r \times (\mathbb{C}^*)^r) // \mathrm{SL}_n = \mathfrak{X}_r(\mathrm{SL}_n) \times (\mathbb{C}^*)^r$ is also normal. However, the GIT projection

$$\mathfrak{X}_r(\mathrm{SL}_n) \times (\mathbb{C}^*)^r \rightarrow \mathfrak{X}_r(\mathrm{SL}_n) \times_{\mathbb{Z}_n^r} (\mathbb{C}^*)^r$$

is then étale because \mathbb{Z}_n^r is finite and acts freely [ibid.]. Then by Theorem 2.4 $\mathfrak{X}_r(\mathrm{GL}_n) \cong \mathfrak{X}_r(\mathrm{SL}_n) \times_{\mathbb{Z}_n^r} (\mathbb{C}^*)^r$ which establishes the result. \square

Corollary 2.7. Let $[\rho] \in \mathfrak{X}_r(\mathrm{SL}_n)$ and let $[\psi] \in \mathfrak{X}_r(\mathrm{SU}_n)$. Then:

- (i) $[\rho] \in \mathfrak{X}_r(\mathrm{SL}_n)^{\mathrm{sing}}$ if and only if $[\rho] \in \mathfrak{X}_r(\mathrm{GL}_n)^{\mathrm{sing}}$.
- (ii) $[\rho] \in \mathfrak{X}_r(\mathrm{SL}_n)^{\mathrm{sm}}$ if and only if $[\rho] \in \mathfrak{X}_r(\mathrm{GL}_n)^{\mathrm{sm}}$.
- (iii) $[\psi] \in \mathfrak{X}_r(\mathrm{SU}_n)^{\mathrm{sing}}$ if and only if $[\psi] \in \mathfrak{X}_r(U_n)^{\mathrm{sing}}$.
- (iv) $[\psi] \in \mathfrak{X}_r(\mathrm{SU}_n)^{\mathrm{sm}}$ if and only if $[\psi] \in \mathfrak{X}_r(U_n)^{\mathrm{sm}}$.

Proof. First let $[\rho] \in \mathfrak{X}_r(\mathrm{SL}_n)$. Corollary 2.6 tells that $\mathfrak{X}_r(\mathrm{SL}_n) \times (\mathbb{C}^*)^r \rightarrow \mathfrak{X}_r(\mathrm{GL}_n)$ is an étale equivalence and such mappings preserve tangent spaces, we conclude

$$T_{[\rho]}(\mathfrak{X}_r(\mathrm{GL}_n)) \cong T_{[\rho]}(\mathfrak{X}_r(\mathrm{SL}_n) \times (\mathbb{C}^*)^r) \cong T_{[\rho]}(\mathfrak{X}_r(\mathrm{SL}_n)) \oplus \mathbb{C}^r.$$

By counting dimensions and noticing that

$$\dim_{\mathbb{C}}(\mathfrak{X}_r(\mathrm{GL}_n)) = \dim_{\mathbb{C}}(\mathfrak{X}_r(\mathrm{SL}_n)) + r,$$

results (i) and (ii) follow.

Results (iii) and (iv) follow from (i) and (ii) and the additional observations that $\dim_{\mathbb{C}}(\mathfrak{X}_r(K_{\mathbb{C}})) = \dim_{\mathbb{R}}(\mathfrak{X}_r(K))$ and $\dim_{\mathbb{C}}(T_{[\psi]}(\mathfrak{X}_r(K_{\mathbb{C}}))) = \dim_{\mathbb{R}}(T_{[\psi]}(\mathfrak{X}_r(K)))$. \square

Corollary 2.8. *We have the following isomorphisms of character varieties:*

- (i) $\mathfrak{X}_r(\mathrm{PSL}_n) \cong \mathfrak{X}_r(\mathrm{SL}_n) // \mathbb{Z}_n^r$ in the category of algebraic varieties.
- (ii) $\mathfrak{X}_r(\mathrm{PU}_n) \cong \mathfrak{X}_r(\mathrm{SU}_n) / \mathbb{Z}_n^r$ in the category of semialgebraic sets.

Proof. From the previous theorem we have

$$\mathfrak{X}_r(\mathrm{GL}_n) \cong \mathfrak{X}_r(\mathrm{SL}_n) \times_{\mathbb{Z}_n^r} (\mathbb{C}^*)^r.$$

Taking the quotient of both sides by $(\mathbb{C}^*)^r$ we can conclude $\mathfrak{X}_r(\mathrm{PSL}_n) \cong \mathfrak{X}_r(\mathrm{SL}_n) // \mathbb{Z}_n^r$. More precisely letting $\mu = (\mu_1, \dots, \mu_r) \in (\mathbb{C}^*)^r$ act only on the second factor of $\mathfrak{X}_r(\mathrm{SL}_n) \times (\mathbb{C}^*)^r$,

$$\mu \cdot ([A_1, \dots, A_r], (\lambda_1, \dots, \lambda_r)) = ([A_1, \dots, A_r], (\mu_1 \lambda_1, \dots, \mu_r \lambda_r)),$$

and going through the isomorphisms in (3), one gets that the action on $\mathfrak{X}_r(\mathrm{GL}_n)$ corresponds to scalar multiplication of each entry, so we obtain

$$\begin{aligned} \mathfrak{X}_r(\mathrm{PSL}_n) &\cong \mathfrak{X}_r(\mathrm{GL}_n) // (\mathbb{C}^*)^r \cong (\mathfrak{X}_r(\mathrm{SL}_n) \times_{\mathbb{Z}_n^r} (\mathbb{C}^*)^r) // (\mathbb{C}^*)^r \\ &\cong ((\mathfrak{X}_r(\mathrm{SL}_n) \times (\mathbb{C}^*)^r) // \mathbb{Z}_n^r) // (\mathbb{C}^*)^r \\ &\cong ((\mathfrak{X}_r(\mathrm{SL}_n) \times (\mathbb{C}^*)^r) // (\mathbb{C}^*)^r) // \mathbb{Z}_n^r \cong \mathfrak{X}_r(\mathrm{SL}_n) // \mathbb{Z}_n^r, \end{aligned}$$

as wanted. The other statement is analogous. \square

2B. Examples. We use the results in Section 2A and the theorems from [Florentino and Lawton 2009] to describe the homeomorphism types of the examples of G -character varieties of F_r known to be manifolds with boundary. Let \bar{B}_n denote a closed real ball of indicated dimension, and let $\{*\}$ denote the space consisting of one point.

One can show that whenever $\phi : \mathfrak{X}_r(\mathrm{SL}_n) \rightarrow M$ is an isomorphism (as affine varieties), then $\mathfrak{X}_r(\mathrm{SU}_n) \cong \phi(\mathfrak{X}_r(\mathrm{SU}_n)) \subset M$ (as semialgebraic sets) by restricting ϕ to $\mathfrak{X}_r(\mathrm{SU}_n) \subset \mathfrak{X}_r(\mathrm{SL}_n)$ [Procesi and Schwarz 1985].

We first consider the trivial case $(r, n) = (r, 1)$. In this case the conjugation action

is trivial, and thus we deduce the following table of moduli of $(r, 1)$ -representations:

| | fixed determinant | nonfixed determinant |
|---------|---|--|
| complex | $\mathfrak{X}_r(\mathrm{SL}_1) \cong \{*\}$ | $\mathfrak{X}_r(\mathrm{GL}_1) \cong (\mathbb{C}^*)^r$ |
| compact | $\mathfrak{X}_r(\mathrm{SU}_1) \cong \{*\}$ | $\mathfrak{X}_r(\mathrm{U}_1) \cong (S^1)^r$ |

We next consider the case $r = 1$. The coefficients of the characteristic polynomial of a matrix X , $\{c_1(X), \dots, c_{n-1}(X), \det(X)\}$, define conjugate invariant regular mappings $\mathfrak{X}_1(\mathrm{SL}_n) \rightarrow \mathbb{C}^{n-1}$ and $\mathfrak{X}_1(\mathrm{GL}_n) \rightarrow \mathbb{C}^{n-1} \times \mathbb{C}^*$ which are isomorphisms. Thus we get this table of moduli of $(1, n)$ -representations:

| | fixed determinant | nonfixed determinant |
|---------|--|--|
| complex | $\mathfrak{X}_1(\mathrm{SL}_n) \cong \mathbb{C}^{n-1}$ | $\mathfrak{X}_1(\mathrm{GL}_n) \cong \mathbb{C}^{n-1} \times \mathbb{C}^*$ |
| compact | $\mathfrak{X}_1(\mathrm{SU}_n) \cong \bar{B}_{n-1}$ | $\mathfrak{X}_1(\mathrm{U}_n) \cong \bar{B}_{n-1} \times S^1$ |

Remark 2.9. With respect to the second table, each of the four families of moduli spaces contains no irreducible representations, yet each space is smooth. For this reason these moduli spaces should perhaps be regarded as everywhere singular, since we will see that irreducibles will generally be smooth points for $r \geq 2$.

In the $r = 2$ case we have a well-known isomorphism $\mathfrak{X}_2(\mathrm{SL}_2) \rightarrow \mathbb{C}^3$ given by $[(A, B)] \mapsto (\mathrm{tr}(A), \mathrm{tr}(B), \mathrm{tr}(AB))$; see [Goldman 2009; Vogt 1889; Fricke and Klein 1912]. More generally there is an isomorphism $\mathfrak{gl}(2, \mathbb{C})^2 // \mathrm{PGL}_2 \rightarrow \mathbb{C}^5$ given by

$$[(A, B)] \mapsto (\mathrm{tr}(A), \mathrm{tr}(B), \mathrm{tr}(AB), \det(A), \det(B)).$$

Therefore we get the following moduli of $(2, 2)$ -representations:

| | fixed determinant | nonfixed determinant |
|---------|--|--|
| complex | $\mathfrak{X}_2(\mathrm{SL}_2) \cong \mathbb{C}^3$ | $\mathfrak{X}_2(\mathrm{GL}_2) \cong \mathbb{C}^3 \times (\mathbb{C}^* \times \mathbb{C}^*)$ |
| compact | $\mathfrak{X}_2(\mathrm{SU}_2) \cong \bar{B}_3$ | $\mathfrak{X}_2(\mathrm{U}_2) \cong \bar{B}_3 \times (S^1 \times S^1)$ |

In [Florentino and Lawton 2009] the following fixed determinant cases are established:

| | fixed determinant | nonfixed determinant |
|------------------|---|--|
| compact $(3, 2)$ | $\mathfrak{X}_3(\mathrm{SU}_2) \cong S^6$ | $\mathfrak{X}_3(\mathrm{U}_2) \cong S^6 \times_{\mathbb{Z}_2^3} (S^1 \times S^1 \times S^1)$ |
| compact $(2, 3)$ | $\mathfrak{X}_2(\mathrm{SU}_3) \cong S^8$ | $\mathfrak{X}_2(\mathrm{U}_3) \cong S^8 \times_{\mathbb{Z}_3^2} (S^1 \times S^1)$ |

Remark 2.10. The complex $(3, 2)$ and $(2, 3)$ cases are left out in this last table since we will show they are not manifolds. In each of these cases, the complex

moduli space of fixed determinant is a branched double cover of complex affine space which deformation retract to a sphere. The explicit scheme structures are known as well. See [Florentino and Lawton 2009; Lawton 2007].

We conjectured in [Florentino and Lawton 2009] that this covers all the cases where a topological manifold with boundary can arise. We will prove this conjecture in Section 4.

3. Singularities

3A. *Algebraic-geometric singularities.* There are a number of equivalent ways to describe smoothness for affine varieties.

Let $X = V(f_1, \dots, f_k) \subset \mathbb{C}^n$ be an affine variety. Then its tangent space at the point $\mathbf{p} = (p_1, \dots, p_n) \in X$ is the vector space

$$T_{\mathbf{p}}(X) = \left\{ (v_1, \dots, v_n) \in \mathbb{C}^n \mid \sum_{j=1}^n \frac{\partial f_i}{\partial x_j} \Big|_{\mathbf{p}} (v_j - p_j) = 0 \text{ for all } i \right\}.$$

This coincides with the more general definition $T_{\mathbf{p}}(X) = (\mathfrak{m}_{\mathbf{p}}/\mathfrak{m}_{\mathbf{p}}^2)^*$, which is the dual to the cotangent space $\mathfrak{m}_{\mathbf{p}}/\mathfrak{m}_{\mathbf{p}}^2$, where $\mathfrak{m}_{\mathbf{p}}$ is a maximal ideal in $\mathbb{C}[X]$ corresponding to \mathbf{p} by Hilbert’s Nullstellensatz.

Definition 3.1. The singular locus of X is defined to be

$$X^{\text{sing}} = \{ \mathbf{p} \in X \mid \dim_{\mathbb{C}} T_{\mathbf{p}}(X) > \dim_{\mathbb{K}} \text{rull } X \}.$$

The complement of this set, $X - X^{\text{sing}}$, is a complex manifold. If X is irreducible, then X is path-connected and furthermore $X - X^{\text{sing}}$ is likewise path-connected. See [Shafarevich 1994].

Let $c = n - \dim_{\mathbb{K}} \text{rull } X$. And let J be the $k \times n$ Jacobian matrix of partial derivatives of the k relations defining $X \subset \mathbb{C}^n$. We can assume n is minimal. Then X^{sing} is concretely realized as the affine variety determined by the determinant of the $c \times c$ minors of J . This ideal is referred to as the Jacobian ideal. In this way, X^{sing} is seen to be a proper subvariety of X .

For example, in [Heusener and Porti 2004] it is shown (for $r \geq 3$) that

$$\mathfrak{X}_r(\text{SL}_2)^{\text{sing}} = \mathfrak{X}_r(\text{SL}_2)^{\text{red}}.$$

In [Lawton 2007], explicitly computing the Jacobian ideal, a similar result is also shown: $\mathfrak{X}_2(\text{SL}_3)^{\text{red}} = \mathfrak{X}_2(\text{SL}_3)^{\text{sing}}$.

3B. *Tangent spaces.* Let \mathfrak{g} be the Lie algebra of G . Having addressed the $r = 1$ and $n = 1$ cases, we now assume that $r, n \geq 2$.

The following two lemmas are classical, and in fact are true for any algebraic Lie group over \mathbb{R} or \mathbb{C} . See [Weil 1964]. For a representation $\rho : F_r \rightarrow G$, let us

denote by $\mathfrak{g}_{\text{Ad}_\rho}$ the F_r -module \mathfrak{g} with the adjoint action via ρ . That is, any word $w \in F_r$ acts as $w \cdot X = \text{Ad}_{\rho(w)} X = \rho(w)X\rho(w)^{-1}$, for $X \in \mathfrak{g}$. Consider the cocycles, coboundaries and cohomology of F_r with coefficients in this module. Explicitly:

$$\begin{aligned} Z^1(F_r; \mathfrak{g}_{\text{Ad}_\rho}) &:= \{u : F_r \rightarrow \mathfrak{g} \mid u(xy) = u(x) + \text{Ad}_{\rho(x)} u(y)\}, \\ B^1(F_r; \mathfrak{g}_{\text{Ad}_\rho}) &:= \{u : F_r \rightarrow \mathfrak{g} \mid u(x) = \text{Ad}_{\rho(x)} X - X \text{ for some } X \in \mathfrak{g}\}, \\ H^1(F_r; \mathfrak{g}_{\text{Ad}_\rho}) &:= Z^1(F_r; \mathfrak{g}_{\text{Ad}_\rho})/B^1(F_r; \mathfrak{g}_{\text{Ad}_\rho}). \end{aligned}$$

Lemma 3.2. *Let G be any algebraic Lie group over \mathbb{R} or \mathbb{C} .*

$$T_\rho(\mathfrak{R}_r(G)) \cong \mathfrak{g}^r \cong Z^1(F_r; \mathfrak{g}_{\text{Ad}_\rho}).$$

Let $\text{Orb}_\rho = \{g\rho g^{-1} \mid g \in G\}$ be the G -orbit of ρ , and let $\text{Stab}_\rho = \{g \in G \mid g\rho g^{-1} = \rho\}$ be the G -stabilizer (or isotropy subgroup).

Lemma 3.3. *Let G be any algebraic Lie group over \mathbb{R} or \mathbb{C} .*

$$T_\rho(\text{Orb}_\rho) \cong \mathfrak{g}/\{X \in \mathfrak{g} \mid \text{Ad}_{\rho(x)} X = X\} \cong B^1(F_r; \mathfrak{g}_{\text{Ad}_\rho}).$$

It is not always the case that the tangent space to the quotient is the quotient of tangent spaces. Consider representations from the free group of rank 1 into SL_3 . The ring of invariants is two dimensional and the ring is generated by $\text{tr}(X)$ and $\text{tr}(X^{-1})$. So the ideal is zero and the ring is free. Consequently it is smooth and the representation sending everything to the identity (having maximal stabilizer) is a nonsingular point. This illustrates that there can be smooth points in the quotient that have positive-dimensional stabilizer. At these points, $T_\rho(\mathfrak{R}_r(G)//G) \not\cong T_\rho(\mathfrak{R}_r(G))/T_\rho(\text{Orb}_\rho)$, seen by simply comparing dimensions.

We also note that if we replace free groups by finitely generated groups Γ then the above isomorphisms require a more careful treatment due to the possible existence of nilpotents in the coordinate ring of the scheme associated to $\mathfrak{R}_\Gamma(G)$ [Sikora 2012].

Recall that $\mathfrak{R}_r(G)^s$ is the set of irreducible representations, and $\mathfrak{X}_r(G)^s = \mathfrak{R}_r(G)^s/G$. An action is called *locally free* if the stabilizer is finite, and is called *proper* if the action $G \times X \rightarrow X \times X$ is a proper mapping. In general, the quotient by a proper locally free action of a reductive group on a smooth manifold is an orbifold (a space locally modeled on finite quotients of \mathbb{R}^n).

The following lemma can be found in [Johnson and Millson 1987, pp. 54–57]. See also [Goldman 1990; 1984].

Lemma 3.4. *Let G be reductive and $r, n \geq 2$. The PG action on $\mathfrak{R}_r(G)^s$ is locally free and proper.*

Therefore, $\mathfrak{R}_r(G)^s/G = \mathfrak{R}_r(G)^s/PG$ are orbifolds.

Lemma 3.5. *For G equal to SL_n , GL_n , SU_n , or U_n and $r, n \geq 2$, the associated PG action on $\mathfrak{X}_r(G)^s$ is free. Therefore, in these cases $\mathfrak{X}_r(G)^s/G$ is a smooth manifold.*

Proof. Let $\rho = (X_1, \dots, X_r) \in \mathfrak{X}_r(G)^s$. Then by Burnside's theorem [Lang 2002] the collection $\{X_1, \dots, X_r\}$ generates all of $n \times n$ matrices $M_{n \times n}$ as an algebra, since $r > 1$ and they form an irreducible set of matrices. Suppose there exists $g \in G$ so that for all $1 \leq k \leq r$ we have $gX_kg^{-1} = X_k$. Then g stabilizes all of $M_{n \times n}$.

Consider $M = \mathbb{C}^n$ as a module over $R = M_{n \times n}$. Clearly, M is a simple module since no nontrivial proper subspaces are left invariant by all matrices. Let f_g be the automorphism of \mathbb{C}^n defined by mapping $v \mapsto gv$. Then f_g defines an R -module automorphism of M since g stabilizes all of R . Thus by Schur's lemma the action of g is equal to the action of a scalar; that is, g is central. \square

Lemma 3.5 and Lemma 3.16 (see Section 3D) together immediately imply the following corollary.

Corollary 3.6. *Let $G = \mathrm{SL}_n$, GL_n , SU_n , or U_n . If $[\rho] \in \mathfrak{X}_r(G)^s$ and $r, n \geq 2$, then*

$$T_{[\rho]}(\mathfrak{X}_r(G)) \cong H^1(\mathbb{F}_r; \mathfrak{g}_{\mathrm{Ad}_\rho}).$$

For $G = \mathrm{SL}_n$ we can calculate that $\dim_{\mathbb{C}} \mathfrak{X}_r(G)^s = (n^2 - 1)(r - 1)$ and for $K = \mathrm{SU}_n$, we have $\dim_{\mathbb{R}} \mathfrak{X}_r(K)^s = (n^2 - 1)(r - 1)$. Likewise, for $G = \mathrm{GL}_n$ we calculate $\dim_{\mathbb{C}} \mathfrak{X}_r(G)^s = n^2(r - 1) + 1$ and for $K = \mathrm{U}_n$, $\dim_{\mathbb{R}} \mathfrak{X}_r(K)^s = n^2(r - 1) + 1$.

Let $\mathfrak{X}_r(G)^{\mathrm{sm}} = \mathfrak{X}_r(G) - \mathfrak{X}_r(G)^{\mathrm{sing}}$ be the smooth stratum, which is a complex manifold, open and dense as a subspace of $\mathfrak{X}_r(G)$. The calculation of dimensions above and Corollary 3.6 imply the following lemma which expresses the fact that the irreducibles not only form a smooth manifold but are naturally contained in the smooth stratum of the variety.

Lemma 3.7. *Let $r, n \geq 2$ and G be one of SL_n or GL_n . Then the following equivalent statements hold:*

- (i) $\mathfrak{X}_r(G)^s \subset \mathfrak{X}_r(G)^{\mathrm{sm}}$.
- (ii) $\mathfrak{X}_r(G)^{\mathrm{sing}} \subset \mathfrak{X}_r(G)^{\mathrm{red}}$.

The next lemmas address important technical points that we will need in our proofs.

Lemma 3.8. *$\mathfrak{X}_r(G)^{\mathrm{red}}$ is an algebraic set; that is, a subvariety of $\mathfrak{X}_r(G)$.*

Proof. The irreducibles are exactly the GIT stable points (zero dimensional stabilizer and closed orbits) and in general these are Zariski open, which implies the complement is an algebraic set [Dolgachev 2003]. \square

Lemma 3.9. *Suppose there exists a set $\mathbb{O} \subset \mathfrak{X}_r(G)^{\mathrm{sing}} \cap \mathfrak{X}_r(G)^{\mathrm{red}}$ that is dense with respect to the ball topology in $\mathfrak{X}_r(G)^{\mathrm{red}}$. Then $\mathfrak{X}_r(G)^{\mathrm{sing}} = \mathfrak{X}_r(G)^{\mathrm{red}}$.*

Proof. Since both $\mathfrak{X}_r(G)^{\text{sing}} \subset \mathfrak{X}_r(G)^{\text{red}}$ are subvarieties (by Lemmas 3.7 and 3.8), \mathbb{O} is dense in both with respect to the ball topology. This follows since \mathbb{O} is dense in $\mathfrak{X}_r(G)^{\text{red}}$ with respect to the ball topology and $\mathbb{O} \subset \mathfrak{X}_r(G)^{\text{sing}} \cap \mathfrak{X}_r(G)^{\text{red}}$. Thus $\mathfrak{X}_r(G)^{\text{sing}} = \overline{\mathbb{O}} = \mathfrak{X}_r(G)^{\text{red}}$, where $\overline{\mathbb{O}}$ is the closure of \mathbb{O} in $\mathfrak{X}_r(G)$ with respect to the (metric) ball topology. \square

A set as in Lemma 3.9 was called an *adherence set* in [Heusener and Porti 2004].

3C. Denseness of reducibles with minimal stabilizer. Now consider the following subvarieties of reducibles. Recall that the 0 vector space is not considered to be an irreducible subrepresentation.

Definition 3.10. Define $U_{r,n} \subset \mathfrak{X}_r(\text{GL}_n)^{\text{red}}$ and $W_{r,n} \subset \mathfrak{X}_r(\text{SL}_n)^{\text{red}}$ by:

$$\begin{aligned} U_{r,n} &= \{[\rho_1 \oplus \rho_2] \in \mathfrak{X}_r(\text{GL}_n) : \rho_i : \mathbb{F}_r \rightarrow \text{GL}_{n_i} \text{ are irreducible}\} \\ W_{r,n} &= \{[\rho_1 \oplus \rho_2] \in \mathfrak{X}_r(\text{SL}_n) : \rho_i : \mathbb{F}_r \rightarrow \text{GL}_{n_i} \text{ are irreducible}\}, \end{aligned}$$

where we consider all possible decompositions $n = n_1 + n_2$, with $n_i > 0$.

Note that a given $\rho \in U_{r,n}$ uniquely determines the integers n_1 and n_2 , up to permutation. We will refer to this situation by saying that ρ is of *reduced type* $[n_1, n_2]$. Similar remarks and terminology apply to $W_{r,n}$.

It is clear that

$$(4) \quad \mathfrak{X}_r(\text{SL}_n)^{\text{red}} = \mathfrak{X}_r(\text{GL}_n)^{\text{red}} \cap \mathfrak{X}_r(\text{SL}_n)$$

and that

$$W_{r,n} = U_{r,n} \cap \mathfrak{X}_r(\text{SL}_n).$$

The following lemma is likewise clear by the proof of Lemma 3.5.

Lemma 3.11. *A representation ρ is in $U_{r,n}$ if and only if $\text{Stab}_\rho \cong (\mathbb{C}^*)^2$. Also, $\rho \in W_{r,n}$ if and only if $\text{Stab}_\rho \cong \mathbb{C}^*$.*

The strategy is now to show that $U_{r,n}$ and $W_{r,n}$ contain only singularities. However, we must first establish the following lemma.

Lemma 3.12. *Let $r, n \geq 2$. $U_{r,n}$ is dense in $\mathfrak{X}_r(\text{GL}_n)^{\text{red}}$ with respect to the ball topology.*

Proof. When $n = 2$, $U_{r,n}$ coincides with $\mathfrak{X}_r(\text{GL}_n)^{\text{red}}$, since any completely reducible representation is of reduced type $[1, 1]$. So we assume $n \geq 3$. Let $\rho \in [\rho] \in \mathfrak{X}_r(\text{GL}_n)^{\text{red}}$ have at least three irreducible blocks; that is, $\rho = \rho_1 \oplus \rho_2 \oplus \rho_3$ where ρ_1 and ρ_2 are irreducible and ρ_3 is semisimple. In other words, $[\rho] \in \mathfrak{X}_r(\text{GL}_n)^{\text{red}} - U_{r,n}$.

Then $\rho_2 \oplus \rho_3$ is a semisimple representation into GL_k for some k . Since the irreducible representations $\mathbb{F}_r \rightarrow \text{GL}_k$ are dense (here we use $r > 1$), there exists an

irreducible sequence $\sigma_j \in \text{Hom}(F_r, \text{GL}_k)$ satisfying

$$\lim_{j \rightarrow \infty} \sigma_j = \rho_2 \oplus \rho_3,$$

which in turn implies

$$\lim_{j \rightarrow \infty} \rho_1 \oplus \sigma_j = \rho_1 \oplus \rho_2 \oplus \rho_3 = \rho,$$

where $\rho_1 \oplus \sigma_j$ is in $U_{r,n}$. Thus we have a sequence $[\rho_1 \oplus \sigma_j] \in U_{r,n} \subset \mathfrak{X}_r(\text{GL}_n)^{\text{red}}$ whose limit is $[\rho_1 \oplus \rho_2 \oplus \rho_3]$. This shows that $U_{r,n}$ is dense in $\mathfrak{X}_r(\text{GL}_n)^{\text{red}}$ and proves the lemma. \square

Corollary 3.13. *Let $r, n \geq 2$. Then $W_{r,n}$ is dense in $\mathfrak{X}_r(\text{SL}_n)^{\text{red}}$ with respect to the ball topology.*

Proof. First we show that $\mathfrak{X}_r(\text{SL}_n)^{\text{red}} \subset \overline{W_{r,n}}$. Using the previous lemma and Equation (4), let

$$\begin{aligned} [\rho] \in \mathfrak{X}_r(\text{SL}_n)^{\text{red}} &= \mathfrak{X}_r(\text{SL}_n) \cap \mathfrak{X}_r(\text{GL}_n)^{\text{red}} \\ &= \mathfrak{X}_r(\text{SL}_n) \cap \overline{U_{r,n}}. \end{aligned}$$

Then, we can write $\rho = \lim \sigma_j$, where $\sigma_j = \rho_1^{(j)} \oplus \rho_2^{(j)} \in U_{r,n}$ is of reduced type $[n_1, n_2]$. Let us write $\lambda_j := \det \rho_1^{(j)} \det \rho_2^{(j)}$. Since the limit is a well-defined point $[\rho] \in \mathfrak{X}_r(\text{SL}_n)^{\text{red}}$, we can arrange for the sequence to be in $W_{r,n}$ as follows. Letting $\alpha_j = (1/\lambda_j)^{1/n_1}$ (for any choice of branch cut), we can also write $\rho = \lim \eta_j$ where $\eta_j = (\rho_1^{(j)} \alpha_j) \oplus \rho_2^{(j)} \in W_{r,n}$, (since now η_j has unit determinant), from which one sees that $\rho \in \overline{W_{r,n}}$, as wanted. Finally, we get:

$$\begin{aligned} \mathfrak{X}_r(\text{SL}_n)^{\text{red}} \subset \overline{W_{r,n}} &= \overline{\mathfrak{X}_r(\text{SL}_n) \cap U_{r,n}} \\ &\subset \overline{\mathfrak{X}_r(\text{SL}_n)} \cap \overline{U_{r,n}} \\ &= \mathfrak{X}_r(\text{SL}_n)^{\text{red}}, \end{aligned}$$

which implies all these sets coincide, finishing the proof. Here, we used the standard fact that the closure of an intersection is contained in the intersection of the closures, and that $\mathfrak{X}_r(\text{SL}_n)$ is closed in $\mathfrak{X}_r(\text{GL}_n)$. \square

3D. The Luna slice theorem and the Zariski tangent space. We now prove a strong lemma, first proved in [Heusener and Porti 2004] and later in more generality in [Sikora 2012], which tells exactly how to understand the Zariski tangent space at a general free group representation. For a similar result see also [Drézé 2004, p. 45]. To that end, we review the Luna slice theorem [1973]. We recommend [Drézé 2004] for a good exposition.

Following [Schwarz 2004], we define an étale map between complex affine varieties as a local analytic isomorphism in the ball topology.

Theorem 3.14 (weak Luna slice theorem at smooth points). *Let G be a reductive algebraic group acting on an affine variety X . Let $x \in X$ be a smooth point with Orb_x closed. Then there exists a subvariety $x \in V \subset X$, and Stab_x -invariant étale morphism $\phi : V \rightarrow T_x V$ satisfying:*

- (i) V is locally closed, affine, smooth, and Stab_x -stable.
- (ii) $V \hookrightarrow X \rightarrow X//G$ induces $T_{[x]}(V//\text{Stab}_x) \cong T_{[x]}(X//G)$.
- (iii) $\phi(x) = 0$ and $d\phi_x = \text{Id}$.
- (iv) $T_x X = T_x(\text{Orb}_x) \oplus T_x V$ with respect to the Stab_x -action.
- (v) ϕ induces $T_{[x]}(V//\text{Stab}_x) \cong T_0(T_x V//\text{Stab}_x)$.

Remark 3.15. The reader familiar with Luna's slice theorem may be wondering how Theorem 3.14 is implied. Firstly, note that ψ is an étale mapping if and only if the completion of the local rings satisfy $\widehat{\mathcal{O}}_x \cong \widehat{\mathcal{O}}_{\psi(x)}$ which implies the subset of derivations are isomorphic, the latter being isomorphic to the Zariski tangent spaces. The usual Luna slice theorem implies $\phi : V//\text{Stab}_x \rightarrow \phi(V)//\text{Stab}_x$ is étale, $(G \times V)//\text{Stab}_x \cong U \subset X$ is saturated and open, and $V//\text{Stab}_x \rightarrow U//G$ is étale. We thus respectively conclude lines (v), (iv), and (ii) in the above theorem.

Lemma 3.16. *Let G be a complex algebraic reductive Lie group. For any $[\rho] \in \mathfrak{X}_r(G)$,*

$$T_{[\rho]}\mathfrak{X}_r(G) \cong T_0(H^1(\mathbb{F}_r; \mathfrak{g}_{\text{Ad}_{\rho^{\text{ss}}}})//\text{Stab}_{\rho^{\text{ss}}}),$$

where ρ^{ss} is a polystable representative from the extended orbit $[\rho]$.

Proof. Any $\rho^{\text{ss}} \in [\rho]$ has a closed orbit and is a smooth point of $\mathfrak{X}_r(G)$, and every point $[\rho] \in \mathfrak{X}_r(G)$ contains such a ρ^{ss} .

By the Luna slice theorem, there exists an algebraic set $\rho^{\text{ss}} \in V_{\rho^{\text{ss}}} \subset \mathfrak{X}_r(G)$ such that:

- (i) $\text{Stab}_{\rho^{\text{ss}}}(V_{\rho^{\text{ss}}}) \subset V_{\rho^{\text{ss}}}$
- (ii) With respect to the reductive action of $\text{Stab}_{\rho^{\text{ss}}}$,

$$\begin{aligned} Z^1(\mathbb{F}_r; \mathfrak{g}_{\text{Ad}_{\rho^{\text{ss}}}}) &\cong T_{\rho^{\text{ss}}}(\mathfrak{X}_r(G)) \cong T_{\rho^{\text{ss}}}(\text{Orb}_{\rho^{\text{ss}}}) \oplus T_{\rho^{\text{ss}}}(V_{\rho^{\text{ss}}}) \\ &\cong B^1(\mathbb{F}_r; \mathfrak{g}_{\text{Ad}_{\rho^{\text{ss}}}}) \oplus T_{\rho^{\text{ss}}}(V_{\rho^{\text{ss}}}), \end{aligned}$$

since ρ^{ss} is smooth.

- (iii) Thus, $H^1(\mathbb{F}_r; \mathfrak{g}_{\text{Ad}_{\rho^{\text{ss}}}}) \cong T_{\rho^{\text{ss}}}(V_{\rho^{\text{ss}}})$, as $\text{Stab}_{\rho^{\text{ss}}}$ -spaces.
- (iv) $V_{\rho^{\text{ss}}} \hookrightarrow \mathfrak{X}_r(G) \rightarrow \mathfrak{X}_r(G)$ induces $T_{[\rho^{\text{ss}}]}(V_{\rho^{\text{ss}}}//\text{Stab}_{\rho^{\text{ss}}}) \cong T_{[\rho]}\mathfrak{X}_r(G)$.
- (v) $T_{[\rho^{\text{ss}}]}(V_{\rho^{\text{ss}}}//\text{Stab}_{\rho^{\text{ss}}}) \cong T_0(T_{\rho^{\text{ss}}}(V_{\rho^{\text{ss}}})//\text{Stab}_{\rho^{\text{ss}}})$, since ρ^{ss} is smooth.

Putting these steps together we conclude

$$T_{[\rho]}\mathfrak{X}_r(G) \cong T_0(T_{\rho^{\text{ss}}}(V_{\rho^{\text{ss}}})//\text{Stab}_{\rho^{\text{ss}}}) \cong T_0(H^1(\mathbb{F}_r; \mathfrak{g}_{\text{Ad}_{\rho^{\text{ss}}}})//\text{Stab}_{\rho^{\text{ss}}}). \quad \square$$

Remark 3.17. Upon closer examination we find $H^1(F_r; \mathfrak{g}_{\text{Ad}_{\rho^{\text{ss}}}} // \text{Stab}_{\rho^{\text{ss}}})$ to be an étale neighborhood; that is, an algebraic set that maps, via an étale mapping, to an open set (in the ball topology) of $\mathfrak{X}_r(G)$ [Schwarz 2004, p. 223].

3E. The \mathbb{C}^* -action on cohomology. As we saw in Corollary 3.13, the generic singularity will occur when Stab_{ρ} is the smallest possible torus group, namely \mathbb{C}^* or $\mathbb{C}^* \times \mathbb{C}^*$, for the cases $G = \text{SL}_n$ or $G = \text{GL}_n$, respectively.

To study the \mathbb{C}^* -action on cohomology, the following setup will be relevant.

Fix two integers $n, k \geq 1$. Consider the vector space $\mathbb{C}^{2n} = \mathbb{C}^n \times \mathbb{C}^n$ with variables $(\mathbf{z}, \mathbf{w}) = (z_1, \dots, z_n, w_1, \dots, w_n)$ and the action of \mathbb{C}^* given by

$$(5) \quad \lambda \cdot (\mathbf{z}, \mathbf{w}) = (\lambda^k \mathbf{z}, \lambda^{-k} \mathbf{w}).$$

Let us denote by $\mathbb{C}^{2n} //_k \mathbb{C}^*$ the corresponding affine GIT quotient. It is the spectrum of the ring $\mathbb{C}[\mathbf{z}, \mathbf{w}]^{\mathbb{C}^*}$ of polynomial invariants under this action. To describe this ring, let

$$p(\mathbf{z}, \mathbf{w}) = z_1^{a_1} \dots z_n^{a_n} w_1^{b_1} \dots w_n^{b_n}$$

be a monomial, with $a_i, b_i \in \mathbb{N}$, and define

$$\partial p := \sum_{j=1}^n a_j - b_j.$$

Any polynomial invariant under the action is a sum of monomials p such that $\partial p = 0$. Considering the monomials with smallest degree, we are led to conclude that

$$\mathbb{C}[\mathbf{z}, \mathbf{w}]^{\mathbb{C}^*} = \mathbb{C}[z_1 w_1, \dots, z_1 w_n, \dots, z_n w_1, \dots, z_n w_n].$$

Note that this shows that the quotient is independent of k . By viewing these n^2 generators as elements of an $n \times n$ matrix $X = (x_{ij})$, $x_{ij} = z_i w_j$, which necessarily has rank at most one, we conclude that this is the ring of polynomial functions on the variety $V \subset M_{n \times n}(\mathbb{C})$ of matrices of rank ≤ 1 :

$$\mathbb{C}[\mathbf{z}, \mathbf{w}]^{\mathbb{C}^*} = \mathbb{C}[V].$$

The variety V is called a determinantal variety [Harris 1992] and one can show that $\mathbb{C}[V] = \mathbb{C}[x_{ij}] / I$ where I is the ideal of 2×2 minors of X . By simple computations, V has a unique singularity, the zero matrix, which corresponds to the orbit of zero in \mathbb{C}^{2n} .

Now, observe that all orbits of the action (5) are closed except those contained in

$$Z := \{0\} \times \mathbb{C}^n \cup \mathbb{C}^n \times \{0\},$$

and moreover there is only one closed orbit in Z , which is easily seen to be the only singular point of $\mathbb{C}^{2n} //_k \mathbb{C}^*$. Therefore, by GIT, the quotient

$$(\mathbb{C}^{2n} \setminus Z) / \mathbb{C}^*$$

is a geometric quotient. We summarize these results as follows.

Lemma 3.18. *Let $n \geq 2$.*

- (i) $\mathbb{C}^{2n} //_k \mathbb{C}^*$ is isomorphic to the determinantal variety of $n \times n$ square matrices of rank ≤ 1 . Its unique singularity is the orbit of the origin.
- (ii) $(\mathbb{C}^{2n} \setminus Z) / \mathbb{C}^*$ is isomorphic to $\mathbb{C}^* \times \mathbb{C}P^{n-1} \times \mathbb{C}P^{n-1}$.

Because of the fact that the GIT quotient is obtained from $(\mathbb{C}^{2n} \setminus Z) / \mathbb{C}^*$ by adding just one point, the singular point, and because of (ii) above, we will refer to $\mathbb{C}^{2n} //_k \mathbb{C}^*$ as an affine cone over $\mathbb{C}P^{n-1} \times \mathbb{C}P^{n-1}$, and denote it by $\mathcal{C}_{\mathbb{C}}(\mathbb{C}P^{n-1} \times \mathbb{C}P^{n-1})$. It is called the affine cone over the Segre variety in [Mukai 2003].

Now consider the following antiholomorphic involution of $\mathbb{C}^{2n} = \mathbb{C}^n \times \mathbb{C}^n$:

$$j : (z, w) \mapsto -(\bar{w}, \bar{z}),$$

and consider the same action as above, but restrict it to $S^1 \subset \mathbb{C}^*$. This will be relevant in the study of the compact quotients. The fixed point set of the involution j is the set

$$F := \{(z, -\bar{z}) : z \in \mathbb{C}\} \subset \mathbb{C}^n \times \mathbb{C}^n,$$

which is canonically identified with the first copy of \mathbb{C}^n (as real vector spaces).

Lemma 3.19. (i) *The S^1 -action on \mathbb{C}^{2n} commutes with j .*

- (ii) *The quotient F/S^1 of its restriction to F is homeomorphic to a real open cone over $\mathbb{C}P^{n-1}$ denoted by $\mathcal{C}(\mathbb{C}P^{n-1})$.*

Proof. Proving (i) is straightforward, and we leave it to the reader.

To prove (ii) first observe that on the fixed point set, the S^1 -action just gives

$$\lambda \cdot (z, -\bar{z}) = (\lambda z, -\bar{\lambda} \bar{z}), \quad \lambda \in S^1$$

so we can describe it as an action of S^1 on the first copy of \mathbb{C}^n . Since the action is free except for the origin, all orbits are circles and the quotient \mathbb{C}^n / S^1 is the union of $\mathbb{C}^n \setminus \{0\} / S^1$ with a single point. Since $\mathbb{C}^n \setminus \{0\} / S^1$ is homeomorphic to $(S^{2n-1} / S^1) \times \mathbb{R}$, we obtain that F/S^1 is the real cone over S^{2n-1} / S^1 , the latter being well known to be $\mathbb{C}P^{n-1}$. \square

These singularity types will be encountered in SL_n and SU_n -character varieties. In fact, the same singularities will also appear in GL_n and U_n -character varieties, because the actions in these cases are very similar.

Indeed one can easily show the following:

Proposition 3.20. *Let $n \geq 2$. Let $T = \mathbb{C}^* \times \mathbb{C}^*$ act on a vector space $V = \mathbb{C}^{2n} = \mathbb{C}^n \times \mathbb{C}^n$ as follows:*

$$(\lambda, \mu) \cdot (\mathbf{z}, \mathbf{w}) = (\lambda\mu^{-1}\mathbf{z}, \mu\lambda^{-1}\mathbf{w}).$$

Then, $\mathbb{C}^{2n} // T$ is isomorphic to $\mathbb{C}^{2n} //_2 \mathbb{C}^$. In particular, as before, this quotient is the determinantal variety of $n \times n$ square matrices of rank ≤ 1 , which has dimension $2n - 1$. Its unique singularity is the orbit of the origin.*

Proof. We just need to argue, as before, that the invariant polynomials are generated by the same monomials, those of the form $z_j w_k$, for any indices $j, k \in \{1, \dots, n\}$, so they form an $n \times n$ matrix with rank one. □

Finally, note that for $n = 1$ we get a smooth variety: $\mathbb{C}^2 //_2 \mathbb{C}^* \cong \mathbb{C}$.

3F. Proof of Theorem 1.1, Case 1: GL_n or SL_n .

Theorem 3.21. *Let $r, n \geq 2$ and $G = GL_n$ or SL_n . Then $\mathfrak{X}_r(G)^{\text{sing}} = \mathfrak{X}_r(G)^{\text{red}}$ if and only if $(r, n) \neq (2, 2)$.*

Remark 3.22. If $n = 1$ the statement is vacuously true since in these cases there are no reducibles, nor are there singularities. We have already seen that there are smooth reducibles in the cases $r = 1, n \geq 2$, and $(r, n) = (2, 2)$ since there always exist reducibles in these cases and the entire moduli spaces are smooth.

Proof. Let $G = GL_n$. By Lemma 3.7 it is enough to show $\mathfrak{X}_r(G)^{\text{red}} \subset \mathfrak{X}_r(G)^{\text{sing}}$.

Let $\rho \in U_{r,n} \subset \mathfrak{X}_r(G)^{\text{red}}$ be of reduced type $[n_1, n_2]$ with $n_1, n_2 > 0$ and $n = n_1 + n_2$ (see Definition 3.10) and write it in the form

$$\rho = \rho_1 \oplus \rho_2 = \begin{pmatrix} \vec{X} & \vec{0}_{n_1 \times n_2} \\ \vec{0}_{n_2 \times n_1} & \vec{Y} \end{pmatrix},$$

where $\vec{X} = (X_1, \dots, X_r) \in M_{n_1 \times n_1}^r$ and $\vec{Y} = (Y_1, \dots, Y_r) \in M_{n_2 \times n_2}^r$ and $\vec{0}_{k \times l} = (0_{k \times l}, \dots, 0_{k \times l})$ where $0_{k \times l}$ is the k by l matrix of zeros and the vector has r entries. Recall that these representations form a dense set in $\mathfrak{X}_r(G)^{\text{red}}$, by Lemma 3.12.

Let $\text{diag}(a_1, \dots, a_n)$ be an $n \times n$ matrix whose (i, j) -entry is 0 if $i \neq j$ and is equal to a_i otherwise. Then $\text{Stab}_\rho = \mathbb{C}^* \times \mathbb{C}^*$ is given by

$$\text{diag}(\underbrace{\lambda, \dots, \lambda}_{n_1}, \underbrace{\mu, \dots, \mu}_{n_2}).$$

We note that the action of the center is trivial so we often consider the stabilizer with respect to the action of G modulo its center.

Then the cocycles satisfy

$$Z^1(\mathbb{F}_r; \text{Ad}_\rho) \cong \mathfrak{g}^r \\ = \left\{ \begin{pmatrix} \vec{A} & \vec{B} \\ \vec{C} & \vec{D} \end{pmatrix} \mid \vec{A} \in M_{n_1 \times n_1}^r, \vec{B} \in M_{n_1 \times n_2}^r, \vec{C} \in M_{n_2 \times n_1}^r, \vec{D} \in M_{n_2 \times n_2}^r \right\},$$

which has dimension $n^2 r$ since it is the tangent space to a representation and the representation variety is smooth. The coboundaries are given by

$$B^1(\mathbb{F}_r; \text{Ad}_\rho) \cong \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} - \begin{pmatrix} \vec{X} & \vec{0}_{n_1 \times n_2} \\ \vec{0}_{n_2 \times n_1} & \vec{Y} \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} \vec{X}^{-1} & \vec{0}_{n_1 \times n_2} \\ \vec{0}_{n_2 \times n_1} & \vec{Y}^{-1} \end{pmatrix} \right\} \\ \cong \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} - \begin{pmatrix} \vec{X}A\vec{X}^{-1} & \vec{X}B\vec{Y}^{-1} \\ \vec{Y}C\vec{X}^{-1} & \vec{Y}D\vec{Y}^{-1} \end{pmatrix} \right\},$$

for a fixed element $\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathfrak{g}$. This has dimension $n^2 - 2$ since it is the tangent space to the G -orbit of ρ which has dimension equal to that of the group minus its stabilizer.

Thus with respect to the torus action,

$$(6) \quad H^1(\mathbb{F}_r; \text{Ad}_\rho) \cong H^1(\mathbb{F}_r; \text{Ad}_{\rho_1}) \oplus H^1(\mathbb{F}_r; \text{Ad}_{\rho_2}) \oplus W,$$

where W exists since the torus action is reductive. By considering the Euler characteristic, one has that

$$\dim_{\mathbb{C}} H^0(\mathbb{F}_r; \text{Ad}_\rho) - \dim_{\mathbb{C}} H^1(\mathbb{F}_r; \text{Ad}_\rho) = (1 - r) \dim_{\mathbb{C}} \mathfrak{gl}(n, \mathbb{C}).$$

Then since $H^0(\mathbb{F}_r; \text{Ad}_\rho) = Z^0(\mathbb{F}_r; \text{Ad}_\rho)$ is the centralizer in \mathfrak{g} of the image of ρ , we calculate

$$\dim_{\mathbb{C}} H^1(\mathbb{F}_r; \text{Ad}_\rho) = n^2(r - 1) + 2, \\ \dim_{\mathbb{C}} H^1(\mathbb{F}_r; \text{Ad}_{\rho_i}) = n_i^2(r - 1) + 1, \quad i = 1, 2.$$

This then implies $\dim_{\mathbb{C}} H^1(\mathbb{F}_r; \text{Ad}_\rho) // (\mathbb{C}^* \times \mathbb{C}^*) = n^2(r - 1) + 1 = \dim_{\mathbb{C}} \mathfrak{X}_r(G)$, since the diagonal of the $\mathbb{C}^* \times \mathbb{C}^*$ -action is the center which acts trivially. We conclude that

$$\dim_{\mathbb{C}} W = (n^2 - n_1^2 - n_2^2)(r - 1) = 2n_1n_2(r - 1).$$

Explicitly, the Stab_ρ action on $H^1(\mathbb{F}_r; \text{Ad}_\rho)$ is given by

$$\text{diag}(\underbrace{\lambda, \dots, \lambda}_{n_1}, \underbrace{\mu, \dots, \mu}_{n_2}) \cdot \left[\begin{pmatrix} \vec{A} & \vec{B} \\ \vec{C} & \vec{D} \end{pmatrix} \right] \mapsto \left[\begin{pmatrix} \vec{A} & \lambda \vec{B} \mu^{-1} \\ \mu \vec{C} \lambda^{-1} & \vec{D} \end{pmatrix} \right]$$

which respects representatives up to coboundary.

So, the action on $H^1(\mathbb{F}_r; \text{Ad}_{\rho_1}) \oplus H^1(\mathbb{F}_r; \text{Ad}_{\rho_2})$ is trivial (but not so on W) and we conclude

$$H^1(\mathbb{F}_r; \text{Ad}_\rho) // (\mathbb{C}^* \times \mathbb{C}^*) \cong H^1(\mathbb{F}_r; \text{Ad}_{\rho_1}) \oplus H^1(\mathbb{F}_r; \text{Ad}_{\rho_2}) \oplus (W // (\mathbb{C}^* \times \mathbb{C}^*)).$$

Therefore, by Proposition 3.20, we have established that 0 is a singularity (solution to the generators of the singular locus) of $W // (\mathbb{C}^* \times \mathbb{C}^*)$ which then implies it is a singularity to $H^1(\mathbb{F}_r; \text{Ad}_\rho) // (\mathbb{C}^* \times \mathbb{C}^*)$ (whenever $\dim_{\mathbb{C}} W > 2$) which then in turn implies any $\rho \in U_{r,n}$ is a singularity in $\mathfrak{X}_r(G)$ by Lemma 3.16 (note $\rho = \rho^{\text{ss}}$ here). $U_{r,n}$ is dense in $\mathfrak{X}_r(\text{GL}_n)^{\text{red}}$ by Lemma 3.12. Then Lemma 3.9 applies to show that $\mathfrak{X}_r(\text{GL}_n)^{\text{sing}} = \mathfrak{X}_r(\text{GL}_n)^{\text{red}}$ whenever $\dim_{\mathbb{C}} W = 2n_1n_2(r-1) > 2$; that is, whenever $(r, n) \neq (2, 2)$.

Now let $[\rho] \in \mathfrak{X}_r(\text{SL}_n)$. Then it is easy to see that $[\rho] \in \mathfrak{X}_r(\text{SL}_n)^{\text{red}}$ if and only if $[\rho] \in \mathfrak{X}_r(\text{GL}_n)^{\text{red}}$. Then Corollary 2.7 and the previously established case together imply $\mathfrak{X}_r(\text{SL}_n)^{\text{red}} = \mathfrak{X}_r(\text{SL}_n)^{\text{sing}}$.

This finishes the proof of Theorem 1.1 for the groups SL_n and GL_n . □

Remark 3.23. We note that the cohomology decomposition used in the proof depends on the decomposition of ρ . For instance, in the 2×2 determinant 1 case, the reducible representation takes values in $\text{SL}_1 \times \text{GL}_1 = \mathbb{C}^0 \times \mathbb{C}^*$, where \mathbb{C}^0 is a point. Then by Lemma 3.18:

$$\begin{aligned} H^1(\mathbb{F}_r, \text{Ad}_\rho) // \mathbb{C}^* &\cong H^1(\mathbb{F}_r, \text{Ad}_{\rho_1}) \oplus H^1(\mathbb{F}_r, \text{Ad}_{\rho_2}) \oplus (W // \mathbb{C}^*) \\ &\cong \mathbb{C}^0 \times \mathbb{C}^r \times ((\mathbb{C}^{2r} / \mathbb{C}^2) // \mathbb{C}^*) \\ &\cong \mathbb{C}^r \times \mathbb{C}^{2r-2} //_2 \mathbb{C}^* \\ &\cong \mathbb{C}^r \times \mathcal{C}_{\mathbb{C}}(\mathbb{C}P^{r-2} \times \mathbb{C}P^{r-2}). \end{aligned}$$

Remark 3.24. The proof above works directly, with suitable modifications for the case $G = \text{SL}_n$. For instance the action of the stabilizer in this case is $\text{Stab}_\rho = \mathbb{C}^*$ given by

$$\text{diag}(\underbrace{\lambda, \dots, \lambda}_{n_1}, \underbrace{\mu, \dots, \mu}_{n_2}),$$

where $\lambda^{n_1} \mu^{n_2} = 1$ which is equivalent to $\mu = \lambda^{-\frac{n_1}{n_2}}$. The cocycles satisfy

$$\begin{aligned} Z^1(\mathbb{F}_r; \text{Ad}_\rho) \\ \cong \mathfrak{g}^r = \left\{ \begin{pmatrix} \vec{A} & \vec{B} \\ \vec{C} & \vec{D} \end{pmatrix} \mid \vec{A} \in M_{n_1 \times n_1}^r, \vec{B} \in M_{n_1 \times n_2}^r, \vec{C} \in M_{n_2 \times n_1}^r, \vec{D} \in M_{n_2 \times n_2}^r, \right. \\ \left. \text{tr}(A_i) = -\text{tr}(D_i), 1 \leq i \leq r \right\}, \end{aligned}$$

which has dimension $(n^2 - 1)r$. The rest carries over without significant change.

Remark 3.25. Similar results for the moduli of tuples of generic matrices have been obtained in [Le Bruyn and Procesi 1987], and with respect to the moduli of vector bundles similar results have been obtained in [Laszlo 1996].

3G. Proof of Theorem 1.1, Case 2: SU_n or U_n . Let $K = SU_n$ or U_n and let \mathfrak{k} be its Lie algebra in either case.

The tangent space at a point $[\rho] \in \mathfrak{X}_r(K)$ is defined from the semialgebraic structure; that is, any real semialgebraic set has a well-defined coordinate ring which allows one to define the Zariski tangent space as we did at the start of this section [Bochnak et al. 1998]. At smooth points this corresponds to the usual tangent space defined by differentials. It is not hard to see that the semialgebraic set $\mathfrak{X}_r(K)$ is a subset of the real points of $\mathfrak{X}_r(K_{\mathbb{C}})$. Then, the Zariski tangent space of $\mathfrak{X}_r(K)$ at $[\rho]$, $T_{[\rho]}(\mathfrak{X}_r(K))$, consists of the real points of the complex Zariski tangent space $T_{[\rho]}(\mathfrak{X}_r(K_{\mathbb{C}}))$.

As is true for $K_{\mathbb{C}}$ -representations, we define a K -representation to be irreducible if it does not admit any proper (nontrivial) invariant subspaces with respect to the standard action on \mathbb{C}^n . As with $K_{\mathbb{C}}$ -valued representations, we call a K -valued representation reducible if it is not irreducible.

Lemma 3.26. $\mathfrak{X}_r(K_{\mathbb{C}})^{\text{red}} \cap \mathfrak{X}_r(K) = \mathfrak{X}_r(K)^{\text{red}}$.

Proof. First note that $\mathfrak{X}_r(K) \subset \mathfrak{X}_r(K_{\mathbb{C}})$ (see [Florentino and Lawton 2009]). So it suffices to prove that every K -valued representation is K -conjugate to a reducible representation if and only if it is $K_{\mathbb{C}}$ -conjugate to a reducible representation.

Let ρ be a K -representation and suppose that it is K -conjugate to a representation that admits a nontrivial proper invariant subspace of \mathbb{C}^n , then since $K \subset K_{\mathbb{C}}$ it is true that ρ is $K_{\mathbb{C}}$ -conjugate to a reducible representation. Conversely, suppose that a K -representation ρ is $K_{\mathbb{C}}$ -conjugate to a reducible representation. However, conjugating by $K_{\mathbb{C}}$ is simply a change-of-basis, and such a change-of-basis is always possible by conjugating by K by using the Gram–Schmidt algorithm. \square

Lemma 3.27. $\mathfrak{X}_r(K_{\mathbb{C}})^{\text{sing}} \cap \mathfrak{X}_r(K) = \mathfrak{X}_r(K)^{\text{sing}}$

Proof. Let

$$[\rho] \in \mathfrak{X}_r(K) \subset \mathfrak{X}_r(K_{\mathbb{C}}).$$

Then $[\rho] \in \mathfrak{X}_r(K)^{\text{sing}}$ if and only if $\dim_{\mathbb{R}} T_{[\rho]}\mathfrak{X}_r(K) = \dim_{\mathbb{C}} T_{[\rho]}\mathfrak{X}_r(K_{\mathbb{C}})$ exceeds $\dim_{\mathbb{R}} \mathfrak{X}_r(K) = \dim_{\mathbb{C}} \mathfrak{X}_r(K_{\mathbb{C}})$, the latter occurring if and only if $[\rho] \in \mathfrak{X}_r(K_{\mathbb{C}})^{\text{sing}}$. \square

The last case to consider to finish the proof of Theorem 1.1 is $\mathfrak{X}_r(K)$ in terms of SU_n and U_n .

Theorem 3.28. *Let K be either U_n or SU_n . Then $\mathfrak{X}_r(K)^{\text{red}} = \mathfrak{X}_r(K)^{\text{sing}}$ if*

$$\mathfrak{X}_r(K_{\mathbb{C}})^{\text{red}} = \mathfrak{X}_r(K_{\mathbb{C}})^{\text{sing}}.$$

Proof. This follows directly by Lemmas 3.26 and 3.27. \square

Since we have already established in Theorem 3.21 that, for $r, n \geq 2$ and $K \in \{U_n, SU_n\}$, we have $\mathfrak{X}_r(K_{\mathbb{C}})^{\text{red}} = \mathfrak{X}_r(K_{\mathbb{C}})^{\text{sing}}$ if and only if $(r, n) \neq (2, 2)$, Theorem 3.28 is enough to finish the proof of Theorem 1.1.

3H. Iterative reducibles and the singular stratification. As above let K be either U_n or SU_n and $G = K_{\mathbb{C}}$, and let the N -th singular stratum be defined by

$$\text{Sing}_N(\mathfrak{X}_r(G)) = (\cdots ((\mathfrak{X}_r(G))^{\text{sing}})^{\text{sing}\cdots})^{\text{sing}},$$

which is well-defined since each singular locus is a variety and as such has a singular locus itself.

The N -th level reducibles

$$\text{Red}_N(\mathfrak{X}_r(G)) = (\cdots ((\mathfrak{X}_r(G))^{\text{red}})^{\text{red}\cdots})^{\text{red}}$$

is defined inductively in the following way.

Let $\text{Red}_1(\mathfrak{X}_r(G)) = \mathfrak{X}_r(G)^{\text{red}}$. For $k \geq 1$ define $\text{Red}_k(\mathfrak{X}_r(G))^{(k+1)}$ to be the set of $\rho \in \text{Red}_k(\mathfrak{X}_r(G))$ which is minimally reducible, that is has a decomposition into irreducible subrepresentations that has minimal summands. We define $\text{Red}_{k+1}(\mathfrak{X}_r(G)) = \text{Red}_k(\mathfrak{X}_r(G)) - \text{Red}_k(\mathfrak{X}_r(G))^{(k+1)}$ to be the complement of $\text{Red}_k(\mathfrak{X}_r(G))^{(k+1)}$ in $\text{Red}_k(\mathfrak{X}_r(G))$. Thus, $\text{Red}_1(\mathfrak{X}_r(G))^{(2)}$ is always the reducibles that have exactly 2 irreducible subrepresentations — these are exactly the ones we considered in the proof of Theorem 3.21. More generally, $\text{Red}_k(\mathfrak{X}_r(G))^{(k+1)}$ are the representations which decompose into exactly $k + 1$ irreducible subrepresentations. For example, $\text{Red}_2(\mathfrak{X}_r(\text{SL}_3))$ are the representations conjugate to a representation that has its semisimplification diagonal, and $\text{Red}_3(\mathfrak{X}_r(\text{SL}_3)) = \emptyset$.

Likewise we have $\text{Red}_N(\mathfrak{X}_r(K))$ and $\text{Sing}_N(\mathfrak{X}_r(K))$.

Theorem 3.29. *Let $r, n \geq 2$ and $(r, n) \neq (2, 2)$. If $N \geq 1$, then*

$$\text{Sing}_N(\mathfrak{X}_r(G)) \cong \text{Red}_N(\mathfrak{X}_r(G)) \quad \text{and} \quad \text{Sing}_N(\mathfrak{X}_r(K)) \cong \text{Red}_N(\mathfrak{X}_r(K)).$$

The result follows by induction on the irreducible block forms and observing that each block form now corresponds to GL_k , or U_k in the compact cases.

3I. Remarks about other groups.

3I.1. General reductive groups. Let G be a reductive complex algebraic group. It can be shown [Sikora 2012] that the definition given before of an irreducible representation $\rho : \Gamma \rightarrow G$ corresponds exactly to the quotient group $\text{Stab}_{\rho}/Z(G)$ being finite.

Proposition 3.30. *If the adjoint action of ρ is irreducible on \mathfrak{g} , then $[\rho]$ is smooth in $\mathfrak{X}_r(G)$.*

Proof. If Ad_ρ is irreducible, then $\text{Stab}_{\rho^{ss}}$ is central and so $\text{Stab}_{\rho^{ss}}$ acts trivially on $H^1(F_r; \mathfrak{g}_{\text{Ad}_{\rho^{ss}}})$. Hence 0 is not in the Jacobian ideal of $H^1(F_r; \mathfrak{g}_{\text{Ad}_{\rho^{ss}}})//\text{Stab}_{\rho^{ss}}$. So, by Lemma 3.16, $[\rho]$ is smooth in $\mathfrak{X}_r(G)$. \square

From the proof of Proposition 3.30, we obtain:

Corollary 3.31. *Let G be a complex reductive algebraic group and $\rho \in \mathfrak{R}_r(G)$ is irreducible with central stabilizer. Then $[\rho]$ is smooth in $\mathfrak{X}_r(G)$.*

A representation satisfying the conditions of this corollary is called *good*. In other words, $\rho \in \mathfrak{R}_r(G)^s$ is good if and only if $\text{Stab}_\rho/Z(G)$ is trivial. Letting $\mathfrak{R}_r(G)^{\text{good}}$ be the open subset of good representations, it easily follows that

$$\mathfrak{X}_r(G)^{\text{good}} := \mathfrak{R}_r(G)^{\text{good}}/G \subset \mathfrak{X}_r(G)^s \subset \mathfrak{X}_r(G)$$

is always a smooth manifold.

[Heusener and Porti 2004] shows that our main theorem, i.e.,

$$\mathfrak{X}_r(G)^{\text{red}} = \mathfrak{X}_r(G)^{\text{sing}},$$

is not true for all reductive Lie groups G and free groups F_r since for PSL_2 there are irreducible representations which are singular. The issue is that the stabilizer of an irreducible representation, modulo the center of G , may not be trivial in general. This is not an issue for GL_n or SL_n since Lemma 3.5 shows the action is free on the set of irreducibles; that is, in these cases a representation is good if and only if it is irreducible.

Let O_n be the group of $n \times n$ complex orthogonal matrices, and let Sp_{2n} be the group of $2n \times 2n$ complex symplectic matrices.

Proposition 3.32. *There exists irreducible representations $\rho : F_r \rightarrow G$ for G any of O_n , PSL_n , and Sp_{2n} such that ρ is not good.*

Proof. It is sufficient in each case to find, for some n , a nonparabolic subgroup of G whose centralizer contains a noncentral element.

First consider a SL_2 -representation ρ contained in the subgroup of diagonal and antidiagonal matrices (containing at least one nondiagonal element and one noncentral element). Then $\text{Stab}_\rho/Z(\text{SL}_2)$ is trivial, and so such a representation is irreducible. However ρ also determines an irreducible PSL_2 -valued representation consisting of diagonal and antidiagonal matrices. However, its stabilizer now contains $\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$, since this element acts as scalar multiplication by -1 on the antidiagonal components and trivially on the diagonal components, so the action is trivial for PSL_2 -representations but nontrivial for SL_2 -representations. This element is not central in SL_2 . Thus ρ defines an irreducible representation into PSL_2 that has finite noncentral stabilizer, and thus is not good.

For O_n representations consider any representation whose image consists of all

matrices of the form

$$\left\{ \begin{pmatrix} \pm 1 & 0 & 0 & 0 \\ 0 & \pm 1 & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & \pm 1 \end{pmatrix} \right\}.$$

One easily computes that the stabilizer is finite and not trivial and thus they are irreducible with $\text{Stab}_\rho/Z(O_n)$ not trivial and thus are not good.

For Sp_{2n} representations we can likewise find examples like the following for $n = 2$. Let the representation have its image generated by

$$\left\{ \pm \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}, \pm \begin{pmatrix} i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \\ 0 & 0 & i & 0 \\ 0 & 0 & 0 & -i \end{pmatrix}, \pm \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \right\}.$$

We get a subgroup of order 16 with finite stabilizer; as such, this group is an irreducible with finite noncentral stabilizer. Again we see that $\text{Stab}_\rho/Z(\text{Sp}_{2n})$ is not trivial and thus this representation is not good. \square

Remark 3.33. In the case of PSL_2 representations (and consequently for SL_2 -valued representations) there are irreducible representations that act reducibly on \mathfrak{g} . However, for PSL_2 these are singular points, but for SL_2 they are smooth. This shows that Ad-reducibility does not imply nonsmoothness in general. In fact, in $\mathfrak{X}_2(\text{PSL}_2)$ there are simultaneously reducibles that are smooth points and irreducibles that are singular. See [Heusener and Porti 2004].

Conjecture 3.34. Let G be a complex reductive algebraic group, and suppose $r \geq 3$. Then $\mathfrak{X}_r(G)^{\text{red}} \subset \mathfrak{X}_r(G)^{\text{sing}}$, and if G is semisimple equality holds if and only if G is a Cartesian product of SL_n 's.

We leave the exploration of this interesting conjecture and the description of singular irreducibles to future work.

3I.2. What if Γ is not free? One may wonder what the relationships exist, if any, between reducible representations and singular points in $\mathfrak{X}_\Gamma(G)$ for a general finitely generated group Γ .

With a given presentation of Γ as $\Gamma = \langle x_1, \dots, x_r \mid r_1, \dots, r_k \rangle$ we can naturally associate the canonical epimorphism $F_r \rightarrow \Gamma = F_r/\langle r_1, \dots, r_k \rangle$ which induces the inclusion $\mathfrak{X}_\Gamma(G) \subset \mathfrak{X}_{F_r}(G)$ providing $\mathfrak{X}_\Gamma(G)$ with the structure of an affine subvariety. As such, ρ is irreducible (resp. completely reducible) in $\mathfrak{X}_\Gamma(G)$ if and only if ρ is irreducible (resp. completely reducible) in $\mathfrak{X}_{F_r}(G)$.

However, the notion of singularity is very far from being well behaved:

- (i) If Γ is free abelian then all representations are reducible and thus the singularities cannot equal the reducibles since the singularities are a proper subset. So reducibles can be smooth; in fact this example shows all smooth points can be reducible.
- (ii) The irreducibles are not generally all smooth in the representation variety let alone in the quotient variety; see [Sikora 2012, Example 38]. Such representations can project to singular points in the quotient (as one might hope is the general situation). Therefore, there can be representations in $\mathfrak{X}_\Gamma(G)^{\text{sing}} \subset \mathfrak{X}_\Gamma(G) \subset \mathfrak{X}_{F_r}(G)$ which are smooth in $\mathfrak{X}_{F_r}(G)$.
- (iii) Singularities in the quotient do not necessarily arise from singularities in the representation space. For example, if Γ is the fundamental group of a genus 2 surface there exist representations in $\mathfrak{R}_\Gamma(\text{SU}_2)$ that are singular but the quotient $\mathfrak{X}_\Gamma(\text{SU}_2) \approx \mathbb{CP}^3$ is smooth. See [Narasimhan and Seshadri 1965; Narasimhan and Ramanan 1969].
- (iv) Lemma 3.16 and its generalizations [Sikora 2012] do not necessarily apply in general.

Therefore, when Γ is not free there is little one can say in general.

4. Local structure and classification of manifold cases

Having completed the proof of Theorem 1.1, we now move on to prove Theorem 1.2. As stated earlier, in [Bratholdt and Cooper 2001] it is established that $\mathfrak{X}_r(\text{SU}_2)$ are not topological manifolds when $r \geq 4$. They compute explicit examples where the representations (abelian, nontrivial) are contained in a neighborhood homeomorphic to $\mathcal{C}(\mathbb{CP}^{r-2}) \times \mathbb{R}^r$, where $\mathcal{C}(X) = (X \times [0, 1]) / (X \times \{0\})$ is the real open cone over a topological space X . From this characterization, simple arguments imply that $\mathfrak{X}_r(\text{SU}_2)$ is not a manifold for $r \geq 4$. It is also a consequence of the following criterion, which will be useful later.

Lemma 4.1. *Let X be a manifold of dimension n and let $d \geq 0$. If $\mathcal{C}(X) \times \mathbb{R}^d$ is Euclidean (i.e, homeomorphic to \mathbb{R}^{d+n+1}) then X is homotopically equivalent to S^n (a sphere of dimension n). Also, if $\mathcal{C}(X) \times \mathbb{R}^d$ is half-Euclidean (i.e, homeomorphic to a closed half-space in \mathbb{R}^{d+n+1}) then X is homotopically equivalent to either a point or S^n .*

Proof. Let p be the cone point of $\mathcal{C}(X)$. Using the natural deformation retraction from $\mathcal{C}(X) - \{p\}$ to X , we see that

$$\mathcal{C}(X) \times \mathbb{R}^d - (\{p\} \times \mathbb{R}^d) = X \times (0, 1) \times \mathbb{R}^d \simeq X,$$

where $Y \simeq X$ symbolizes Y being homotopic to X . On the other hand, if $\mathcal{C}(X) \times \mathbb{R}^d = \mathbb{R}^{n+d+1}$ then $\mathcal{C}(X) \times \mathbb{R}^d - (\{p\} \times \mathbb{R}^d) = \mathbb{R}^{n+d+1} - \mathbb{R}^d \simeq S^n$.

The other statement follows in a similar fashion if the cone point is not on the boundary of the half-space. Otherwise, $\{p\} \times \mathbb{R}^d$ is contained in the boundary so extracting it results in a contractible space. \square

4A. $\mathfrak{X}_r(\mathrm{SU}_n)$ and $\mathfrak{X}_r(\mathrm{U}_n)$. In this subsection we establish the compact cases of Theorem 1.2.

Let $K = \mathrm{SU}_n$ and let \mathfrak{k} be its Lie algebra. Let $d_{r,n} = (n^2 - 1)(r - 1) = \dim_{\mathbb{C}} \mathfrak{X}_r(K) = \dim_{\mathbb{R}} \mathfrak{X}_r(K)$. Whenever $\mathfrak{X}_r(K)$ is not a topological manifold, there exists a point $[\rho] \in \mathfrak{X}_r(K)$ and a neighborhood \mathcal{N} containing $[\rho]$ that is not locally homeomorphic to $\mathbb{R}^{d_{r,n}}$, or $\mathbb{R}_+^{d_{r,n}}$ in the case of a boundary point.

We need a smooth version of Mostow's slice theorem [Mostow 1957; Bredon 1972]. Let \mathcal{N}_x denote a neighborhood at x .

Lemma 4.2. *For any $[\rho] \in \mathfrak{X}_r(K)$, there is a neighborhood $\mathcal{N}_{[\rho]}$ homeomorphic to $H^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho})/\mathrm{Stab}_\rho$. Moreover,*

$$T_{[\rho]}\mathfrak{X}_r(K) \cong T_0(H^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho})/\mathrm{Stab}_\rho).$$

Proof. Let $\mathfrak{R}_r(K) = \mathrm{Hom}(\mathbb{F}_r, K)$. Since $\rho \in \mathfrak{R}_r(K)$ is a smooth point, $T_\rho\mathfrak{R}_r(K) \cong Z^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho})$. Moreover, $T_\rho\mathrm{Orb}_\rho \cong B^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho}) \subset Z^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho})$. Since Stab_ρ is compact and acts on $B^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho})$, there exists a Stab_ρ -invariant complement W . Thus $Z^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho}) \cong T_\rho\mathfrak{R}_r(K) \cong B^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho}) \oplus W$, which respects the action of the stabilizer. Since $\mathfrak{R}_r(K)$ is a smooth compact Riemannian manifold we can invariantly exponentiate W to obtain a slice $\exp(W) = S \subset \mathfrak{R}_r(K)$ such that $T_\rho S = W$. Therefore, $T_\rho S \cong H^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho})$ as Stab_ρ -spaces.

Saturating S by K we obtain an open K -invariant space, which contains the orbit of ρ since $\rho \in S$; namely $U = K(S)$. Since U is open $T_\rho U = T_\rho\mathfrak{R}_r(K)$, and since it is saturated $U/K \cong S/\mathrm{Stab}_\rho$ is an open subset of $\mathfrak{X}_r(K)$.

Putting these observations together we conclude S is locally diffeomorphic to $T_\rho S$ which implies the neighborhood $U/K \cong H^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho})/\mathrm{Stab}_\rho$, which establishes our first claim.

Then S/Stab_ρ is locally homeomorphic to $T_\rho S/\mathrm{Stab}_\rho$, which then implies

$$(7) \quad T_{[\rho]}(S/\mathrm{Stab}_\rho) \cong T_0(T_\rho S/\mathrm{Stab}_\rho).$$

But

$$(8) \quad T_{[\rho]}\mathfrak{X}_r(K) = T_{[\rho]}(U/K) \cong T_{[\rho]}(S/\mathrm{Stab}_\rho)$$

and

$$(9) \quad T_0(T_\rho S/\mathrm{Stab}_\rho) \cong T_0(H^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho})/\mathrm{Stab}_\rho).$$

Equations (7), (8), and (9) together complete the proof. \square

Remark 4.3. The above lemma holds for all compact Lie groups K .

Theorem 4.4. *Let $r, n \geq 2$ and let $\rho \in \mathfrak{R}_r(\mathrm{SU}_n)$ be of reduced type $[n_1, n_2]$. Then, there exists a neighborhood $[\rho] \in \mathcal{N} \subset \mathfrak{X}_r(\mathrm{SU}_n)$ that is homeomorphic to $\mathbb{R}^{d_{\mathrm{SU}_n}} \times \mathcal{C}(\mathbb{CP}^{(r-1)n_1n_2-1})$, where $d_{\mathrm{SU}_n} = (r-1)(n_1^2 + n_2^2 - 1) + 1$. Also, if $\rho \in \mathfrak{R}_r(\mathrm{U}_n)$ is of reduced type $[n_1, n_2]$, there exists a neighborhood $[\rho] \in \mathcal{N} \subset \mathfrak{X}_r(\mathrm{U}_n)$ that is homeomorphic to $\mathbb{R}^{d_{\mathrm{U}_n}} \times \mathcal{C}(\mathbb{CP}^{(r-1)n_1n_2-1})$, where $d_{\mathrm{U}_n} = (r-1)(n_1^2 + n_2^2) + 2$.*

Corollary 4.5. *If $K = \mathrm{U}_n$ or $K = \mathrm{SU}_n$, both $r, n \geq 2$, and $(r, n) \neq (2, 2), (2, 3),$ or $(3, 2)$, then $\mathfrak{X}_r(K)$ is not a manifold with boundary.*

Proof. Theorem 4.4 implies that $\mathfrak{X}_r(\mathrm{U}_n)$ and $\mathfrak{X}_r(\mathrm{SU}_n)$ are manifolds only if $\mathbb{R}^{d_{\mathrm{U}_n}} \times \mathcal{C}(\mathbb{CP}^{(r-1)n_1n_2-1})$ and $\mathbb{R}^{d_{\mathrm{SU}_n}} \times \mathcal{C}(\mathbb{CP}^{(r-1)n_1n_2-1})$, respectively, are locally Euclidean. By Lemma 4.1, this can only be the case if $n_1n_2(r-1) - 1 \in \{0, 1\}$, with $n = n_1 + n_2$ and $n_1, n_2 > 0$. In the first case, $n_1n_2(r-1) = 1$, which implies $n_1 = n_2 = 1$ and $r = 2$, so $(r, n) = (2, 2)$. From Section 2B we know $\mathfrak{X}_2(\mathrm{U}_2)$ and $\mathfrak{X}_2(\mathrm{SU}_2)$ are manifolds with boundary, and we conclude the neighborhood in this case is half-Euclidean since $\mathcal{N} = \mathbb{R}^d \times [0, 1)$, for appropriate d .

The other possibility is $n_1n_2(r-1) = 2$ so that $n_1 = 2$ and $n_2 = 1$, or $n_1 = 1$ and $n_2 = 2$, and $r = 2$. This is the case $(r, n) = (2, 3)$. Otherwise, $r = 3$ and $n_1 = n_2 = 1$, which is the case $(r, n) = (3, 2)$. Moreover, from Section 2B these two are the only cases which are manifolds.

Having exhausted all possibilities, the proof is complete. \square

We now prove Theorem 4.4.

Proof of Theorem 4.4. Similar to Theorem 3.21, there is a direct computational proof of Theorem 4.4. However, using Theorem 3.21, Lemma 3.19 and the relation between K and its complexification, we can provide a shorter argument.

Let τ be the Cartan involution on $\mathfrak{g} = \mathfrak{gl}(n, \mathbb{C})$, the Lie algebra of GL_n , which is just the linear map $A \mapsto -\bar{A}^T$, acting on a matrix $A \in \mathfrak{gl}(n, \mathbb{C})$. By definition, the fixed point subspace of τ is \mathfrak{k} , the Lie algebra \mathfrak{u}_n of U_n . One easily checks that τ induces an involution on $Z^1(\mathbb{F}_r; \mathfrak{g}_{\mathrm{Ad}_\rho}) \cong \mathfrak{g}^r$, whose fixed subspace is $Z^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho}) \cong \mathfrak{k}^r$, and similarly $B^1(\mathbb{F}_r; \mathfrak{g}_{\mathrm{Ad}_\rho})^\tau = B^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho})$. This implies that τ induces an involution, also denoted τ , on the first cohomology, and that $H^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho})$ is naturally isomorphic to $H^1(\mathbb{F}_r; \mathfrak{g}_{\mathrm{Ad}_\rho})^\tau$.

Now, assume that $\rho = \rho_1 \oplus \rho_2 \in U_{r,n} \cap \mathfrak{R}_r(\mathrm{U}_n)$, is of reduced type $[n_1, n_2]$ ($n_1, n_2 > 0, n_1 + n_2 = n$). Note that ρ_1 and ρ_2 are irreducible representations in $\mathfrak{R}_r(\mathrm{U}_{n_1})$ and $\mathfrak{R}_r(\mathrm{U}_{n_2})$, respectively, and with respect to the PU_n conjugation action $\mathrm{Stab}_\rho \cong S^1$. Then a cocycle $\phi \in Z^1(\mathbb{F}_r; \mathfrak{k}_{\mathrm{Ad}_\rho}) \cong \mathfrak{k}^r$ has the form

$$\phi = \begin{pmatrix} \phi_1 & A \\ -\bar{A}^T & \phi_2 \end{pmatrix},$$

where $\phi_i \in Z^1(\mathbb{F}_r; \mathfrak{k}_{\text{Ad}_{\rho_i}})$, and as in Theorem 3.21, A is now an arbitrary r -tuple of $n_1 \times n_2$ matrices. This shows that τ respects the decomposition in Equation (6), so we get

$$\begin{aligned} H^1(\mathbb{F}_r; \mathfrak{k}_{\text{Ad}_\rho}) &= H^1(\mathbb{F}_r; \mathfrak{g}_{\text{Ad}_\rho})^\tau = H^1(\mathbb{F}_r; \mathfrak{g}_{\text{Ad}_{\rho_1}})^\tau \oplus H^1(\mathbb{F}_r; \mathfrak{g}_{\text{Ad}_{\rho_2}})^\tau \oplus W^\tau \\ &= H^1(\mathbb{F}_r; \mathfrak{k}_{\text{Ad}_{\rho_1}}) \oplus H^1(\mathbb{F}_r; \mathfrak{k}_{\text{Ad}_{\rho_2}}) \oplus F \end{aligned}$$

where, by the form of the cocycles above, we can write

$$F := W^\tau = \{(z, -\bar{z}) : z \in \mathbb{C}^{n_1 n_2 (r-1)}\};$$

using also $\dim_{\mathbb{C}} W = 2n_1 n_2 (r - 1)$.

It follows from Lemma 4.2 that a neighborhood of ρ is locally homeomorphic to $H^1(\mathbb{F}_r; \mathfrak{k}_{\text{Ad}_\rho})/\text{Stab}_\rho$. As in the proof of Theorem 3.21, the action of $\text{Stab}_\rho = S^1$ does not affect $H^1(\mathbb{F}_r; \mathfrak{k}_{\text{Ad}_{\rho_i}})$, $i = 1, 2$, and we conclude that

$$\begin{aligned} H^1(\mathbb{F}_r; \mathfrak{k}_{\text{Ad}_\rho})/\text{Stab}_\rho &= H^1(\mathbb{F}_r; \mathfrak{k}_{\text{Ad}_{\rho_1}}) \oplus H^1(\mathbb{F}_r; \mathfrak{k}_{\text{Ad}_{\rho_2}}) \oplus F/S^1 \\ &\cong \mathbb{R}^{d_{U_n}} \oplus \mathcal{C}(\mathbb{C}P^{n_1 n_2 (r-1)-1}), \end{aligned}$$

by using Lemma 3.19. The dimension d_{U_n} is computed by:

$$d_{U_n} = \sum_{i=1}^2 \dim_{\mathbb{R}} H^1(\mathbb{F}_r; \mathfrak{k}_{\text{Ad}_{\rho_i}}) = (n_1^2 + n_2^2)(r - 1) + 2.$$

The case of $K = \text{SU}_n$ is similar. □

Remark 4.6. In [Le Bruyn and Teranishi 1990] it is shown that the cases (2, 2), (2, 3), and (3, 2) are also the only examples which are complete intersections.

Remark 4.7. Note that using the identity representation (maximal stabilizer) results in $H^1(\mathbb{F}_r; \mathfrak{k}_{\text{Ad}_{\text{id}}})/\text{Stab}_{\text{id}} = \mathfrak{k}^r/\text{SU}_n$ since the coboundaries are trivial. Removing a point results in a homological sphere quotient $S^{(n^2-1)(r-1)-1}/\text{SU}_n$. If there was a Euclidean neighborhood about the identity, then this sphere quotient would be a homology sphere $S^{(n^2-1)(r-2)-1}$. We find this quite likely to give a different obstruction. At the other extreme (central stabilizer) the points are smooth and thus admit Euclidean neighborhoods.

Conjecture 4.8. If K is equal to SU_n or U_n , $[\rho] \in \mathfrak{X}_r(K)^{\text{red}}$, $r, n \geq 2$, and $(r, n) \neq (2, 2), (2, 3)$ or $(3, 2)$, then there does not exist a neighborhood of $[\rho]$ that is Euclidean.

We proved this conjecture for representations of reduced type $[n_1, n_2]$ above. In fact, it seems likely that the neighborhoods around most singularities do not even admit an orbifold structure (not homeomorphic to a finite quotient of a Euclidean ball).

4B. $\mathfrak{X}_r(\mathrm{SL}_n)$ and $\mathfrak{X}_r(\mathrm{GL}_n)$. In this last subsection, we complete the proof of Theorem 1.2 by proving the following result.

Theorem 4.9. *Let $r, n \geq 2$ and let G be SL_n or GL_n . $\mathfrak{X}_r(G)$ is a topological manifold with boundary if and only if $(r, n) = (2, 2)$.*

Proof. By Remark 3.17, $H^1(\mathbb{F}_r; \mathfrak{g}_{\mathrm{Ad}_{\rho^{\mathrm{ss}}}}) // \mathrm{Stab}_{\rho^{\mathrm{ss}}}$ is an étale neighborhood; that is, an algebraic set that maps, via an étale mapping, to an open set (in the ball topology) of $\mathfrak{X}_r(G)$. Thus we see that at a reducible representation with minimal stabilizer (\mathbb{C}^* for SL_n and $\mathbb{C}^* \times \mathbb{C}^*$ for GL_n), that this neighborhood is étale equivalent to $\mathbb{C}^{(n_1^2+n_2^2)(r-1)+2} \times \mathcal{C}_{\mathbb{C}}(\mathbb{C}\mathbb{P}^{(r-1)n_1n_2-1} \times \mathbb{C}\mathbb{P}^{(r-1)n_1n_2-1})$ in $\mathfrak{X}_r(\mathrm{GL}_n)$, where the cone here is the affine cone defined over \mathbb{C}^* . In $\mathfrak{X}_r(\mathrm{SL}_n)$ we have a similar neighborhood. Either way, these spaces are not locally Euclidean neighborhoods for $r, n \geq 2$ unless $n = 2 = r$ which implies that $n_1 = 1 = n_2$. This is seen by similar arguments given above in the compact cases. \square

Acknowledgements

The second author thanks W. Goldman for helpful discussions, the Instituto Superior Técnico for its support during the 2007–2009 academic years, and A. Sikora for useful conversations. We also thank M. Reineke and O. Serman for pointing out relevant references.

References

- [Bochnak et al. 1998] J. Bochnak, M. Coste, and M.-F. Roy, *Real algebraic geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **36**, Springer, Berlin, 1998. MR 2000a:14067 Zbl 0912.14023
- [Bratholdt and Cooper 2001] S. Bratholdt and D. Cooper, “On the topology of the character variety of a free group”, *Rend. Istit. Mat. Univ. Trieste* **32**:Suppl. 1 (2001), 45–53. MR 2003d:14072 Zbl 1062.14505
- [Bredon 1972] G. E. Bredon, *Introduction to compact transformation groups*, Pure and Applied Mathematics **46**, Academic Press, New York, 1972. MR 54 #1265 Zbl 0246.57017
- [Dolgachev 2003] I. Dolgachev, *Lectures on invariant theory*, London Mathematical Society Lecture Note Series **296**, Cambridge University Press, Cambridge, 2003. MR 2004g:14051 Zbl 1023.13006
- [Drézet 2004] J.-M. Drézet, “Luna’s slice theorem and applications”, pp. 39–89 in *Algebraic group actions and quotients* (Wykno, 2000), edited by J. Wiśniewski, Hindawi, Cairo, 2004. MR 2006k:14082 Zbl 1109.14307
- [Florentino and Lawton 2009] C. Florentino and S. Lawton, “The topology of moduli spaces of free group representations”, *Math. Ann.* **345**:2 (2009), 453–489. MR 2010h:14075 Zbl 1200.14093
- [Fricke and Klein 1912] R. Fricke and F. Klein, *Vorlesungen über die Theorie der automorphen Funktionen, II: Die funktionentheoretischen Ausführungen und die Anwendungen, 3: Direkte Beweismethoden der Fundamentaltheoreme und Anhang*, Teubner, Leipzig, 1912. Reprinted in *Bibliotheca Mathematica Teubneriana*, vol. 4, by Johnson Reprint, New York, 1965. MR 32 #1348 JFM 43.0529.08

- [Goldman 1984] W. M. Goldman, “The symplectic nature of fundamental groups of surfaces”, *Adv. in Math.* **54**:2 (1984), 200–225. MR 86i:32042 Zbl 0574.32032
- [Goldman 1990] W. M. Goldman, “Convex real projective structures on compact surfaces”, *J. Differential Geom.* **31**:3 (1990), 791–845. MR 91b:57001 Zbl 0711.53033
- [Goldman 2009] W. M. Goldman, “Trace coordinates on Fricke spaces of some simple hyperbolic surfaces”, pp. 611–684 in *Handbook of Teichmüller theory, II*, edited by A. Papadopoulos, IRMA Lect. Math. Theor. Phys. **13**, Eur. Math. Soc., Zürich, 2009. MR 2010j:30093 Zbl 1175.30043
- [Harris 1992] J. Harris, *Algebraic geometry: a first course*, Graduate Texts in Mathematics **133**, Springer, New York, 1992. MR 93j:14001 Zbl 0779.14001
- [Heusener and Porti 2004] M. Heusener and J. Porti, “The variety of characters in $\mathrm{PSL}_2(\mathbb{C})$ ”, *Bol. Soc. Mat. Mexicana* (3) **10**:Special Issue (2004), 221–237. MR 2006m:57020 Zbl 1100.57014
- [Johnson and Millson 1987] D. Johnson and J. J. Millson, “Deformation spaces associated to compact hyperbolic manifolds”, pp. 48–106 in *Discrete groups in geometry and analysis* (New Haven, CT, 1986), edited by R. Howe, Progr. Math. **67**, Birkhäuser, Boston, 1987. MR 88j:22010 Zbl 0664.53023
- [Lang 2002] S. Lang, *Algebra*, 3rd ed., Graduate Texts in Mathematics **211**, Springer, New York, 2002. MR 2003e:00003 Zbl 0984.00001
- [Laszlo 1996] Y. Laszlo, “Local structure of the moduli space of vector bundles over curves”, *Comment. Math. Helv.* **71**:3 (1996), 373–401. MR 97j:14012 Zbl 0949.14015
- [Lawton 2007] S. Lawton, “Generators, relations and symmetries in pairs of 3×3 unimodular matrices”, *J. Algebra* **313**:2 (2007), 782–801. MR 2008k:16039 Zbl 1119.13004
- [Le Bruyn and Procesi 1987] L. Le Bruyn and C. Procesi, “Étale local structure of matrix invariants and concomitants”, pp. 143–175 in *Algebraic groups* (Utrecht, 1986), edited by A. M. Cohen et al., Lecture Notes in Math. **1271**, Springer, Berlin, 1987. MR 89b:16042 Zbl 0634.14034
- [Le Bruyn and Teranishi 1990] L. Le Bruyn and Y. Teranishi, “Matrix invariants and complete intersections”, *Glasgow Math. J.* **32**:2 (1990), 227–229. MR 91d:15059 Zbl 0714.16016
- [Luna 1973] D. Luna, “Slices étales”, pp. 81–105 in *Sur les groupes algébriques*, Bull. Soc. Math. France Mém. **33**, Soc. Math. France, Paris, 1973. MR 49 #7269 Zbl 0286.14014
- [Mostow 1957] G. D. Mostow, “Equivariant embeddings in Euclidean space”, *Ann. of Math. (2)* **65** (1957), 432–446. MR 19,291c Zbl 0080.16701
- [Mukai 2003] S. Mukai, *An introduction to invariants and moduli*, Cambridge Studies in Advanced Mathematics **81**, Cambridge University Press, Cambridge, 2003. MR 2004g:14002 Zbl 1033.14008
- [Narasimhan and Ramanan 1969] M. S. Narasimhan and S. Ramanan, “Moduli of vector bundles on a compact Riemann surface”, *Ann. of Math. (2)* **89** (1969), 14–51. MR 39 #3518 Zbl 0186.54902
- [Narasimhan and Seshadri 1965] M. S. Narasimhan and C. S. Seshadri, “Stable and unitary vector bundles on a compact Riemann surface”, *Ann. of Math. (2)* **82** (1965), 540–567. MR 32 #1725 Zbl 0171.04803
- [Procesi and Schwarz 1985] C. Procesi and G. Schwarz, “Inequalities defining orbit spaces”, *Invent. Math.* **81**:3 (1985), 539–554. MR 87h:20078 Zbl 0578.14010
- [Schwarz 2004] G. W. Schwarz, “Group actions and quotients for compact Lie groups and algebraic groups”, pp. 209–227 in *Invariant theory in all characteristics* (Kingston, ON, 2002), edited by H. E. A. E. Campbell and D. L. Wehlau, CRM Proc. Lecture Notes **35**, Amer. Math. Soc., Providence, RI, 2004. MR 2005d:20083 Zbl 1131.14312
- [Shafarevich 1994] I. R. Shafarevich, *Basic algebraic geometry, 2: Schemes and complex manifolds*, 2nd ed., Springer, Berlin, 1994. MR 95m:14002 Zbl 0797.14002

- [Sikora 2012] A. S. Sikora, “Character varieties”, *Trans. Amer. Math. Soc.* **364**:10 (2012), 5173–5208. MR 2931326 arXiv 0902.2589
- [Sullivan 1971] D. Sullivan, “Combinatorial invariants of analytic spaces”, pp. 165–168 in *Singularities — Symposium I* (Liverpool, 1969–1970), edited by C. T. C. Hall, Lecture Notes in Math. **192**, Springer, Berlin, 1971. MR 43 #4063 Zbl 0227.32005
- [Vogt 1889] H. Vogt, “Sur les invariants fondamentaux des équations différentielles linéaires du second ordre”, *Ann. Sci. École Norm. Sup. (3)* **6** (1889), 3–71. MR 1508833 JFM 21.0314.01
- [Weil 1964] A. Weil, “Remarks on the cohomology of groups”, *Ann. of Math. (2)* **80** (1964), 149–157. MR 30 #199 Zbl 0192.12802

Received May 30, 2012.

CARLOS FLORENTINO
DEPARTAMENTO DE MATEMÁTICA
INSTITUTO SUPERIOR TÉCNICO
1049-001 LISBOA
PORTUGAL
carlos.florentino@math.ist.utl.pt

SEAN LAWTON
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF TEXAS-PAN AMERICAN
1201 WEST UNIVERSITY DRIVE
EDINBURG, TX 78539
UNITED STATES
lawtonsd@utpa.edu

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 260 No. 1 November 2012

| | |
|---|-----|
| The decomposition of global conformal invariants: Some technical proofs II | 1 |
| SPYROS ALEXAKIS | |
| On deformation quantizations of hypertoric varieties | 89 |
| GWYN BELLAMY and TOSHIRO KUWABARA | |
| Almost factoriality of integral domains and Krull-like domains | 129 |
| GYU WHAN CHANG, HWANKOO KIM and JUNG WOOK LIM | |
| Singularities of free group character varieties | 149 |
| CARLOS FLORENTINO and SEAN LAWTON | |
| Energy identity for the maps from a surface with tension field bounded in L^p | 181 |
| LI JIAYU and ZHU XIANGRONG | |
| Remarks on some isoperimetric properties of the $k - 1$ flow | 197 |
| YU-CHU LIN and DONG-HO TSAI | |
| Demystifying a divisibility property of the Kostant partition function | 215 |
| KAROLA MÉSZÁROS | |
| Exceptional Lie algebras, $SU(3)$, and Jordan pairs | 227 |
| PIERO TRUINI | |
| Lower estimate of Milnor number and characterization of isolated homogeneous hypersurface singularities | 245 |
| STEPHEN S.-T. YAU and HUAIQING ZUO | |



0030-8730(201211)260:1;1-C