

Pacific Journal of Mathematics

**DEMYSTIFYING A DIVISIBILITY PROPERTY OF THE
KOSTANT PARTITION FUNCTION**

KAROLA MÉSZÁROS

DEMYSTIFYING A DIVISIBILITY PROPERTY OF THE KOSTANT PARTITION FUNCTION

KAROLA MÉSZÁROS

We prove a family of identities connected to a divisibility property of the Kostant partition function. A special case of these identities first appeared in a paper of Baldoni and Vergne. To prove their identities, Baldoni and Vergne used residue techniques, and called the resulting divisibility property “mysterious.” Our proofs are entirely combinatorial and provide a natural explanation for why divisibility occurs, both in the Baldoni and Vergne identities and in their generalizations.

1. Introduction

The objective of this paper is to provide a natural combinatorial explanation of a divisibility property of the Kostant partition function. The question of evaluating Kostant partition functions has been the subject of much interest, without a satisfactory combinatorial answer. To mention perhaps the most famous such case: it is known that

$$K_{A_n^+}\left(1, 2, \dots, n, -\binom{n+1}{2}\right) = \prod_{k=1}^n C_k, \quad \text{where } C_k = \frac{1}{k+1} \binom{2k}{k}$$

denotes the Catalan numbers, yet there is no combinatorial proof of this identity! While endowed with combinatorial meaning, Kostant partition functions were introduced in and are a vital part of representation theory: weight multiplicities and tensor product multiplicities can be expressed in terms of the Kostant partition function. Kostant partition functions also come up in toric geometry and analytic residue theory.

Given the lack of understanding of the evaluation of the Kostant partition function, it seems a worthy proposition to provide a simple explanation for certain of its divisibility properties. We explore divisibility properties of Kostant partition functions of types A_n and C_{n+1} , noting that such properties in types B_{n+1} and D_{n+1} are easy consequences of the type C_{n+1} case. A significant part of the type A_n

The author is supported by a National Science Foundation Postdoctoral Research Fellowship.
 MSC2010: 05E10.

Keywords: Kostant partition function, divisibility, flow.

family of identities we study first appeared in [Baldoni and Vergne 2008], where the authors prove the identities using residues, and where they call the divisibility property “mysterious.” It is our hope that the combinatorial argument we provide successfully demystifies the divisibility property of the Kostant partition function and provides a natural explanation for why things happen the way they do.

The outline of the paper is as follows. In Section 2 we define Kostant partition functions of type A_n and prove a family of identities, including the Baldoni–Vergne identities, combinatorially. Our proof is bijective, and as such it also yields an affirmative answer to a question of Stanley [2000] regarding a possible bijective proof of a special case of the Baldoni–Vergne identities. In Section 3 we define Kostant partition functions of type C_{n+1} , relate them to flows, and show how to modify our proof of the identities from Section 2 to obtain their analogues for type C_{n+1} .

2. A family of Kostant partition function identities

In this section we prove a family of Kostant partition function identities exhibiting divisibility properties. We start by proving Baldoni–Vergne identities, our proof of which clearly points to several generalizations of these identities. We provide some of these generalizations in this section, and some in the next.

The Baldoni–Vergne identities. Before stating the Baldoni–Vergne identities, we need a few definitions. Throughout this section the graphs G we consider are on the vertex set $[n + 1]$, possibly with multiple edges, but no loops. Denote by m_{ij} the multiplicity of edge (i, j) , $i < j$, in G . To each edge (i, j) , $i < j$, of G , associate the positive type A_n root $e_i - e_j$, where e_i is the i -th standard basis vector. Let $\{\alpha_1, \dots, \alpha_N\}$ be the multiset of vectors corresponding to the multiset of edges of G as described above. Note that $N = \sum_{1 \leq i < j \leq n+1} m_{ij}$.

The *Kostant partition function* K_G evaluated at the vector $\mathbf{a} \in \mathbb{Z}^{n+1}$ is defined as

$$(1) \quad K_G(\mathbf{a}) = \# \left\{ (b_i)_{i \in [N]} \left| \sum_{i \in [N]} b_i \alpha_i = \mathbf{a} \text{ and } b_i \in \mathbb{Z}_{\geq 0} \right. \right\}.$$

That is, $K_G(\mathbf{a})$ is the number of ways to write the vector \mathbf{a} as a nonnegative linear combination of the positive type A_n roots corresponding to the edges of G , without regard to order. Note that in order for $K_G(\mathbf{a})$ to be nonzero, the partial sums of the coordinates of \mathbf{a} have to satisfy $a_1 + \dots + a_i \geq 0$, $i \in [n]$, and $a_1 + \dots + a_{n+1} = 0$.

We now proceed to state and prove Theorem 1, which first appeared in [Baldoni and Vergne 2008]. Baldoni and Vergne gave a proof of it using residues, and called the result “mysterious”. We provide a natural combinatorial explanation of the result. Our explanation also answers a question of Stanley [2000] in the

affirmative, regarding a possible bijective proof of a special case of the Baldoni–Vergne identities.

For brevity, we write $G - e$, or $G - \{e_1, \dots, e_k\}$, to mean a graph obtained from G with the edge e , or the edges e_1, \dots, e_k , deleted.

Theorem 1 [Baldoni and Vergne 2008]. *Given a connected graph G on the vertex set $[n + 1]$ with $m_{n-1,n} = m_{n-1,n+1} = m_{n,n+1} = 1$, and such that*

$$\frac{m_{j,n-1} + m_{j,n} + m_{j,n+1}}{m_{j,n-1}} = c \quad \text{for all } j \in [n - 2],$$

for some constant c independent of j , we have

$$(2) \quad K_G(\mathbf{a}) = \left(\frac{a_1 + \dots + a_{n-2}}{c} + a_{n-1} + 1 \right) K_{G-(n-1,n)}(\mathbf{a}),$$

for any $\mathbf{a} = (a_1, \dots, a_n, -\sum_{i=1}^n a_i) \in \mathbb{Z}^{n+1}$.

Before proceeding to the formal proof of [Theorem 1](#) we outline it, to fully expose the underlying combinatorics. We introduce the notation

$$(3) \quad Q(\mathbf{a}) := \frac{a_1 + \dots + a_{n-2}}{c} + a_{n-1} + 1$$

for the factor in (2). Rephrasing [Equation \(1\)](#), $K_G(\mathbf{a})$ counts the number of *flows* $\mathbf{f}_G = (b_i)_{i \in N}$ on G satisfying

$$\sum_{i \in [N]} b_i \alpha_i = \mathbf{a} \quad \text{and} \quad b_i \in \mathbb{Z}_{\geq 0}.$$

In the proof of [Theorem 1](#), we introduce the concept of *partial flows* \mathbf{f}_H , about which we prove two key statements:

- The elements of the set of partial flows are in bijection with the flows on $G - (n - 1, n)$ that the Kostant partition function $K_{G-(n-1,n)}(\mathbf{a})$ counts. That is,

$$\# \text{ partial flows} = K_{G-(n-1,n)}(\mathbf{a}).$$

- The elements of the multiset of partial flows \mathbf{f}_H — whose cardinality is $Q(\mathbf{a})$ times the cardinality of the set of partial flows — are in bijection with the flows on G that the Kostant partition function $K_G(\mathbf{a})$ counts. That is,

$$Q(\mathbf{a}) (\# \text{ partial flows}) = K_G(\mathbf{a}).$$

From these two statements we see that the two Kostant partition functions $K_G(\mathbf{a})$ and $K_{G-(n-1,n)}(\mathbf{a})$ are connected by

$$K_G(\mathbf{a}) = Q(\mathbf{a}) K_{G-(n-1,n)}(\mathbf{a}),$$

which is [Equation \(2\)](#).

Proof of Theorem 1. Let $\{\alpha_1, \dots, \alpha_N\}$ be the multiset of vectors corresponding to the edges of G . Let $\alpha_N = e_{n-1} - e_n$, $\alpha_{N-1} = e_{n-1} - e_{n+1}$, and $\alpha_{N-2} = e_n - e_{n+1}$. Then (2) can be rewritten as

$$(4) \quad \# \left\{ (b_i)_{i \in [N]} \left| \sum_{i=1}^N b_i \alpha_i = \mathbf{a} \right. \right\} = Q(\mathbf{a}) \# \left\{ (b_i)_{i \in [N-1]} \left| \sum_{i=1}^{N-1} b_i \alpha_i = \mathbf{a} \right. \right\},$$

where $Q(\mathbf{a})$ is defined in (3).

We proceed to prove the equality (4) bijectively. The key concept we use is that of a partial flow, which we now define.

Consider a flow $\mathbf{f}_H = (b_i)_{i \in [N-3]}$ ($b_i \in \mathbb{Z}_{\geq 0}$) on the edges of the graph

$$H := G - \{(n-1, n), (n-1, n+1), (n, n+1)\}.$$

We call \mathbf{f}_H *partial* if

$$(5) \quad \sum_{i=1}^{N-3} b_i \alpha_i = (a_1, \dots, a_{n-2}, x_{n-1}, x_n, x_{n+1}),$$

for some $x_{n-1}, x_n, x_{n+1} \in \mathbb{Z}$ satisfying $x_{n-1} \leq a_{n-1}$ and $x_n \leq a_n$.

Note that a given partial flow $\mathbf{f}_H = (b_i)_{i \in [N-3]}$ can be extended uniquely to a flow $\mathbf{f}_{G-\{(n-1, n)\}} = (b_i)_{i \in [N-1]}$ ($b_i \in \mathbb{Z}_{\geq 0}$) on $G - \{(n-1, n)\}$ such that $\sum_{i=1}^{N-1} b_i \alpha_i = \mathbf{a}$. Furthermore, each such flow arises from a uniquely determined partial flow. Denote by f the map that takes a partial flow \mathbf{f}_H into a flow $\mathbf{f}_{G-\{(n-1, n)\}}$ as just described. See Figure 1 for an example. The observations above imply:

Claim 1. *The map f establishes a bijection between the set of partial flows \mathbf{f}_H on H and the set of flows on $G - \{(n-1, n)\}$ such that $\sum_{i=1}^{N-1} b_i \alpha_i = \mathbf{a}$. In symbols,*

$$(6) \quad \# \left\{ (b_i)_{i \in [N-1]} \left| \sum_{i=1}^{N-1} b_i \alpha_i = \mathbf{a} \right. \right\} = \sum_{\mathbf{f}_H} 1,$$

where the summation runs over all partial flows \mathbf{f}_H .




Figure 1. Here G is the complete graph on 5 vertices and $\mathbf{a} = (4, 3, -1, 1, -7)$. The flows are written immediately below the corresponding edges. On the left is a partial flow \mathbf{f}_H and on the right is its image under f .

Note that the left-hand side of (6) is the same as the cardinality on the right-hand side of (4).

Given a partial flow f_H , denote by $Y_i(f_H)$, for $i \in \{n-1, n, n+1\}$, the total inflow into vertex $i \in \{n-1, n, n+1\}$ in H , that is, the sum of all the flows b_i on edges of H incident to $i \in \{n-1, n, n+1\}$. Note that a partial flow f_H can be extended in $Y_{n-1}(f_H) + a_{n-1} + 1$ ways to a flow $f_G = (b_i)_{i \in [N]}$ ($b_i \in \mathbb{Z}_{\geq 0}$) of G such that $\sum_{i=1}^N b_i \alpha_i = \mathbf{a}$. Furthermore, given a flow $f_G = (b_i)_{i \in [N]}$ ($b_i \in \mathbb{Z}_{\geq 0}$) such that $\sum_{i=1}^N b_i \alpha_i = \mathbf{a}$, there is a unique partial flow $f_H = (b_i)_{i \in [N-3]}$ from which it can be obtained. Therefore, the above establishes a map g which is a bijection between the multiset of partial flows \mathcal{M} such that each partial flow f_H appears exactly $Y_{n-1}(f_H) + a_{n-1} + 1$ times in \mathcal{M} , and the (multi)set of flows $f_G = (b_i)_{i \in [N]}$ ($b_i \in \mathbb{Z}_{\geq 0}$) on G such that $\sum_{i=1}^N b_i \alpha_i = \mathbf{a}$. See Figure 2 for an example. We thus obtain

$$(7) \quad \# \left\{ (b_i)_{i \in [N]} \mid \sum_{i=1}^N b_i \alpha_i = \mathbf{a} \right\} = \sum_{f_H} (Y_{n-1}(f_H) + a_{n-1} + 1) = \#\mathcal{M},$$

where the second summation runs over the set of partial flows f_H .




Figure 2. Again G is the complete graph on 5 vertices and $\mathbf{a} = (4, 3, -1, 1, -7)$. The flows are written immediately below the corresponding edges. On the left is a partial flow f_H and on the right are the $Y_{n-1}(f_H) + a_{n-1} + 1 = 4$ images under g of the 4 copies of f_H appearing in \mathcal{M} .

Claim 2. *The map g just described establishes a bijection between the multiset \mathcal{M} of partial flows $\mathbf{f}_H = (b_i)_{i \in [N-3]}$ and the (multi)set of flows $\mathbf{f}_G = (b_i)_{i \in [N]}$ ($b_i \in \mathbb{Z}_{\geq 0}$) on G such that $\sum_{i=1}^N b_i \alpha_i = \mathbf{a}$. Moreover,*

$$(8) \quad \#\mathcal{M} = Q(\mathbf{a}) \sum_{\mathbf{f}_H} 1,$$

where the summation runs over all partial flows \mathbf{f}_H .

We already showed that g is a bijection; there remains to show (8). Now, by assumption we have

$$(9) \quad \frac{m_{j,n-1} + m_{j,r} + m_{j,n+1}}{m_{j,n-1}} = c,$$

where c is independent of $j \in [n-2]$; hence

$$(10) \quad \begin{aligned} c \sum_{\mathbf{f}_H} Y_{n-1}(\mathbf{f}_H) &= \sum_{\mathbf{f}_H} (Y_{n-1}(\mathbf{f}_H) + Y_n(\mathbf{f}_H) + Y_{n+1}(\mathbf{f}_H)) \\ &= \sum_{\mathbf{f}_H} (a_1 + \cdots + a_{n-2}), \end{aligned}$$

that is,

$$(11) \quad \sum_{\mathbf{f}_H} Y_{n-1}(\mathbf{f}_H) = \sum_{\mathbf{f}_H} \frac{a_1 + \cdots + a_{n-2}}{c}.$$

Adding $a_{n-1} + 1$ to both summands and recalling (3) and the second equality in (7) yields (8), completing the proof of Claim 2.

In light of (4) it is clear that Claims 1 and 2 together provide a bijective proof of Theorem 1. In terms of equations this can also be seen as follows. Rewrite (7) as

$$(12) \quad \begin{aligned} \#\left\{ (b_i)_{i \in [N]} \left| \sum_{i=1}^N b_i \alpha_i = \mathbf{a} \right. \right\} &= Q(\mathbf{a}) \sum_{\mathbf{f}_H} 1 \\ &= Q(\mathbf{a}) \#\left\{ (b_i)_{i \in [N-1]} \left| \sum_{i=1}^{N-1} b_i \alpha_i = \mathbf{a} \right. \right\}, \end{aligned}$$

where the first equality uses (7) and (8), and the second equality uses (6). \square

Further Kostant partition function identities. Several generalizations are suggested by our proof of the Baldoni–Vergne identities. Here we present an immediate one, Theorem 2; to ensure clarity we include its proof, which follows that of Theorem 1. For brevity, we do not explicitly state the analogues of Claims 1 and 2 from the proof of Theorem 1, though it is easy to obtain them from our proof. In the next section we present further generalizations of other types, but we omit the proofs.

Theorem 2. *Let G be a connected graph on the vertex set $[n + 1]$. Given $k \leq n$, suppose the graph $S = (V(S), E(S))$ defined by*

$$V(S) = \{k, k + 1, \dots, n, n + 1\} \subset [n + 1]$$

and

$$E(S) = \{(i, j) \in E(G) \mid i < j, i \in V(S)\}$$

satisfies these conditions: that

- $\text{outdeg}_S(j) = 2$ and $\text{indeg}_S(j) = 0$, for some $k \leq j \leq n$, and $\text{outdeg}_S(i) = 1$ for $k \leq i \leq n, i \neq j$.
- For some constant c independent of l ,

$$\frac{\sum_{i=k}^{n+1} m_{l,i}}{m_{l,j}} = c \quad \text{for all } l \in [k - 1].$$

Let (j, z) be one of the outgoing edges from j in S . Then

$$(13) \quad K_G(\mathbf{a}) = Q'(\mathbf{a}) K_{G-(j,z)}(\mathbf{a}),$$

for any $\mathbf{a} = (a_1, \dots, a_n, -\sum_{i=1}^n a_i) \in \mathbb{Z}^{n+1}$, where $Q'(\mathbf{a}) = \frac{a_1 + \dots + a_{k-1}}{c} + a_j + 1$.

The most important cases of [Theorem 2](#) are for $k \geq n - 1$, since for $k < n - 1$ several of the edges of G can be “contracted” and reduced to the case $k \geq n - 1$. For $k \geq n - 1$ we obtain four interesting cases from [Theorem 2](#) depending on the form of the graph S and the edge (j, z) . If we take

$$V(S) = \{n - 1, n, n + 1\} \quad \text{and} \quad E(S) = \{(n - 1, n), (n - 1, n + 1), (n, n + 1)\}$$

and $(j, z) = (n - 1, n)$, then [Theorem 2](#) specializes to the original Baldoni–Vergne identities, while other choices of S and (j, z) lead to new identities.

Proof of [Theorem 2](#). Let $\{\alpha_1, \dots, \alpha_N\}$ be the multiset of vectors corresponding to the edges of G such that the multiset of vectors corresponding to the edges of S are $\{\alpha_N - (n + 1 - k), \dots, \alpha_N\}$. Also, let α_N correspond to the edge (j, z) .

Consider a flow $f_H = (b_i)_{i \in [N-(n+2-k)]}$ ($b_i \in \mathbb{Z}_{\geq 0}$) of the edges of the graph $H := ([n + 1], E(G) \setminus E(S))$. We call f_H *partial* if

$$\sum_{i=1}^{N-(n+2-k)} b_i \alpha_i = (a_1, \dots, a_{k-1}, x_k, x_{k+1}, \dots, x_{n+1}),$$

for some $x_i \in \mathbb{Z}$, $k \leq i \leq n + 1$ satisfying $x_i \leq a_i$, for $k \leq i \leq n$.

A given partial flow $\mathbf{f}_H = (b_i)_{i \in [N-(n+2-k)]}$, it can be extended uniquely to a flow $\mathbf{f}_{G-\{(j,z)\}} = (b_i)_{i \in [N-1]}$ ($b_i \in \mathbb{Z}_{\geq 0}$) on $G - \{(j, z)\}$ such that $\sum_{i=1}^{N-1} b_i \alpha_i = \mathbf{a}$. Furthermore, this correspondence is a bijection. Therefore,

$$(14) \quad \# \left\{ (b_i)_{i \in [N-1]} \left| \sum_{i=1}^{N-1} b_i \alpha_i = \mathbf{a} \right. \right\} = \sum_{\mathbf{f}_H} 1,$$

where the summation runs over all partial flows \mathbf{f}_H .

Given a partial flow \mathbf{f}_H in H , we denote by $Y_i(\mathbf{f}_H)$, where $i \in \{k, k+1, \dots, n+1\}$, the total *inflow* into vertex i , that is, the sum of all the flows b_i on edges of H incident to i . The partial flow \mathbf{f}_H can be extended in $Y_j(\mathbf{f}_H) + a_j + 1$ ways to a flow $\mathbf{f}_G = (b_i)_{i \in [N]}$ ($b_i \in \mathbb{Z}_{\geq 0}$) of G such that $\sum_{i=1}^N b_i \alpha_i = \mathbf{a}$. Furthermore, given a flow $\mathbf{f}_G = (b_i)_{i \in [N]}$ ($b_i \in \mathbb{Z}_{\geq 0}$) such that $\sum_{i=1}^N b_i \alpha_i = \mathbf{a}$, there is a unique partial flow $\mathbf{f}_H = (b_i)_{i \in [N-(n+2-k)]}$ from which it can be obtained. Therefore,

$$(15) \quad \# \left\{ (b_i)_{i \in [N]} \left| \sum_{i=1}^N b_i \alpha_i = \mathbf{a} \right. \right\} = \sum_{\mathbf{f}_H} (Y_j(\mathbf{f}_H) + a_j + 1),$$

where the summation runs over all partial flows \mathbf{f}_H .

By assumption, we have

$$\frac{\sum_{i=k}^{n+1} m_{l,i}}{m_{l,j}} = c,$$

where c is independent of $l \in [k-1]$; hence

$$c \sum_{\mathbf{f}_H} Y_j(\mathbf{f}_H) = \sum_{\mathbf{f}_H} \sum_{i=k}^{n+1} Y_i(\mathbf{f}_H) = \sum_{\mathbf{f}_H} (a_1 + \dots + a_{k-1}),$$

that is,

$$(16) \quad \sum_{\mathbf{f}_H} Y_j(\mathbf{f}_H) = \sum_{\mathbf{f}_H} \frac{a_1 + \dots + a_{k-1}}{c}.$$

Thus, (15) can be rewritten as

$$\begin{aligned} \# \left\{ (b_i)_{i \in [N]} \left| \sum_{i=1}^N b_i \alpha_i = \mathbf{a} \right. \right\} &= \sum_{\mathbf{f}_H} \left(\frac{a_1 + \dots + a_{k-1}}{c} + a_j + 1 \right) \\ &= Q'(\mathbf{a}) \sum_{\mathbf{f}_H} 1 = Q'(\mathbf{a}) \# \left\{ (b_i)_{i \in [N-1]} \left| \sum_{i=1}^{N-1} b_i \alpha_i = \mathbf{a} \right. \right\}, \end{aligned}$$

where the first equality uses (15) and (16), and the last equality uses (14). \square

3. Type C_{n+1} Kostant partition functions and the Baldoni–Vergne identities

We now show two generalizations of [Theorem 1](#) in the type C_{n+1} case. We first give the necessary definitions and explain the notion of flow in the context of signed graphs. Throughout this section, the graphs G on the vertex set $[n+1]$ we consider are signed, that is there is a sign $\epsilon \in \{+, -\}$ assigned to each of its edges, with possible multiple edges, and all loops labeled positive. Denote by $(i, j, -)$ and $(i, j, +)$, $i \leq j$, a negative and a positive edge, respectively. Denote by m_{ij}^ϵ the multiplicity of edge (i, j, ϵ) in G , $i \leq j$, $\epsilon \in \{+, -\}$. To each edge (i, j, ϵ) , $i \leq j$, of G , associate the positive type C_{n+1} root $v(i, j, \epsilon)$, where $v(i, j, -) = e_i - e_j$ and $v(i, j, +) = e_i + e_j$. Let $\{\alpha_1, \dots, \alpha_N\}$ be the multiset of vectors corresponding to the multiset of edges of G as described above. Note that $N = \sum_{1 \leq i < j \leq n+1} (m_{ij}^- + m_{ij}^+)$. The *Kostant partition function* K_G evaluated at the vector $\mathbf{a} \in \mathbb{Z}^{n+1}$ is defined as

$$K_G(\mathbf{a}) = \# \left\{ (b_i)_{i \in [N]} \mid \sum_{i \in [N]} b_i \alpha_i = \mathbf{a} \text{ and } b_i \in \mathbb{Z}_{\geq 0} \right\}.$$

That is, $K_G(\mathbf{a})$ is the number of ways to write the vector \mathbf{a} as a nonnegative linear combination of the positive type C_{n+1} roots corresponding to the edges of G , without regard to order.

Just like in the type A_n case, we would like to think of the vector $(b_i)_{i \in [N]}$ as a *flow*. For this we here give a precise definition of flows in the type C_{n+1} case, of which type A_n is of course a special case.

Let G be a signed graph on the vertex set $[n+1]$. Let $\{e_1, \dots, e_N\}$ be the multiset of edges of G , and $\{\alpha_1, \dots, \alpha_N\}$ the multiset of vectors corresponding to the multiset of edges of G . Fix an integer vector $\mathbf{a} = (a_1, \dots, a_n, a_{n+1}) \in \mathbb{Z}^{n+1}$. A *nonnegative integer \mathbf{a} -flow* \mathbf{f}_G on G is a vector $\mathbf{f}_G = (b_i)_{i \in [N]}$ ($b_i \in \mathbb{Z}_{\geq 0}$) such that for all $1 \leq i \leq n+1$, we have

$$(17) \quad \sum_{\substack{e \in E \\ \text{inc}(e, v) = -}} b(e) + a_v = \sum_{\substack{e \in E \\ \text{inc}(e, v) = +}} b(e) + \sum_{e=(v, v, +)} b(e),$$

where $b(e_i) = b_i$, $\text{inc}(e, v) = -$ if edge $e = (g, v, -)$, $g < v$, and $\text{inc}(e, v) = +$ if $e = (g, v, +)$, $g < v$, or $e = (v, j, \epsilon)$, $v < j$, and $\epsilon \in \{+, -\}$.

Call $b(e)$ the *flow* assigned to edge e of G . If the edge e is negative, one can think of $b(e)$ units of fluid flowing on e from its smaller to its bigger vertex. If the edge e is positive, then one can think of $b(e)$ units of fluid flowing away both from e 's smaller and bigger vertex to infinity. Edge e is then a “leak” taking away $2b(e)$ units of fluid.

From the above explanation it is clear that if we are given an \mathbf{a} -flow \mathbf{f}_G such that $\sum_{e=(i, j, +), i \leq j} b(e) = y$, then $\mathbf{a} = (a_1, \dots, a_n, 2y - \sum_{i=1}^n a_i)$. It is then a matter of checking the definitions to see that for a signed graph G on the vertex set $[n+1]$ and

vector $\mathbf{a} = (a_1, \dots, a_n, 2y - \sum_{i=1}^n a_i) \in \mathbb{Z}^{n+1}$, the number of nonnegative integer \mathbf{a} -flows on G is equal to $K_G(\mathbf{a})$.

Thinking of $K_G(\mathbf{a})$ as the number of nonnegative integer \mathbf{a} -flows on G , there is a straightforward generalization of [Theorem 1](#) in the type C_{n+1} case:

Theorem 3. *Given a connected signed graph G on the vertex set $[n+1]$ with $m_{n-1,n}^- = m_{n-1,n+1}^- = m_{n,n+1}^- = 1$, $m_{j,n-1}^+ = m_{j,n}^+ = m_{j,n+1}^+ = 0$, for $j \in [n+1]$, and such that*

$$\frac{m_{j,n-1}^- + m_{j,n}^- + m_{j,n+1}^-}{m_{j,n-1}^-} = c \quad \text{for all } j \in [n-2],$$

for some constant c independent of j , we have

$$(18) \quad K_G(\mathbf{a}) = Q''(\mathbf{a}) K_{G-(n-1,n)}(\mathbf{a})$$

for any $\mathbf{a} = (a_1, \dots, a_n, 2y - \sum_{i=1}^n a_i) \in \mathbb{Z}^{n+1}$, where

$$Q''(\mathbf{a}) = \frac{a_1 + \dots + a_{n-2} - 2y}{c} + a_{n-1} + 1.$$

The proof of [Theorem 3](#) proceeds analogously to that of [Theorem 1](#). Namely, define *partial flows* $\mathbf{f}_H = (b_i)_{i \in [N-3]}$ on

$$H := G - \{(n-1, n, -), (n-1, n+1, -), (n, n+1, -)\}$$

such that

$$\sum_{i=1}^{N-3} b_i \alpha_i = (a_1, \dots, a_{n-2}, x_{n-1}, x_n, x_{n+1}),$$

for some $x_{n-1}, x_n, x_{n+1} \in \mathbb{Z}$, such that $x_{n-1} \leq a_{n-1}$, $x_n \leq a_n$ and the sum of flows on positive edges is y .

Then, one can prove:

- The elements of the set partial flows are in bijection with the nonnegative integer \mathbf{a} -flows on $G - (n-1, n)$. That is,

$$\# \text{ partial flows} = K_{G-(n-1,n)}(\mathbf{a}).$$

- The elements of the multiset of partial flows \mathbf{f}_H , whose cardinality is $Q''(\mathbf{a})$ times the cardinality of the set of partial flows, are in bijection with the nonnegative integer \mathbf{a} -flows on G . That is,

$$Q''(\mathbf{a}) (\# \text{ partial flows}) = K_G(\mathbf{a}).$$

Thus,

$$K_G(\mathbf{a}) = Q''(\mathbf{a}) K_{G-(n-1,n)}(\mathbf{a}).$$

We do not have to require that only negative edges are incident to the vertices $n-1, n, n+1$ in G , as the following theorem shows. The proof is analogous to earlier ones.

Theorem 4. *Given a connected signed graph G on the vertex set $[n+1]$ with $m_{n-1,n}^- = m_{n-1,n+1}^- = m_{n,n+1}^- = 1$, $m_{i,j}^+ = 0$, for $i, j \in \{n-1, n, n+1\}$, and such that*

$$\frac{m_{j,n-1}^\epsilon + m_{j,n}^\epsilon + m_{j,n+1}^\epsilon}{m_{j,n-1}^\epsilon} = c \quad \text{for all } j \in [n-2] \text{ and } \epsilon \in \{+, -\},$$

for some constant c independent of j , we have

$$(19) \quad K_G(\mathbf{a}) = \left(\frac{a_1 + \cdots + a_{n-2} - 2y}{c} + a_{n-1} + 1 \right) K_{G-(n-1,n)}(\mathbf{a})$$

for any $\mathbf{a} = (a_1, \dots, a_n, 2y - \sum_{i=1}^n a_i) \in \mathbb{Z}^{n+1}$.

Of course, [Theorem 2](#) also has type C_{n+1} generalizations and variations. We invite the reader to write these out and check each step of the proof of [Theorem 1](#) and see how they can be adapted.

Acknowledgement

I thank Richard Stanley for his intriguing slides and for pointing out the work of Baldoni and Vergne. I also thank Alejandro Morales and Alex Postnikov for numerous conversations about flows and the Kostant partition function.

References

- [Baldoni and Vergne 2008] W. Baldoni and M. Vergne, “Kostant partitions functions and flow polytopes,” *Transform. Groups* **13**:3-4 (2008), 447–469. [MR 2009k:52029](#) [Zbl 1200.52008](#)
- [Stanley 2000] R. P. Stanley, “Acyclic flow polytopes and Kostant’s partition function,” conference transparencies, 2000, Available at <http://www-math.mit.edu/~rstan/transparencies/kostant.ps>.

Received November 27, 2011. Revised April 4, 2012.

KAROLA MÉSZÁROS
DEPARTMENT OF MATHEMATICS
CORNELL UNIVERSITY
ITHACA, NY 14853
UNITED STATES
karola@math.cornell.edu

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)

Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from [Periodicals Service Company](#), 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 260 No. 1 November 2012

The decomposition of global conformal invariants: Some technical proofs II	1
SPYROS ALEXAKIS	
On deformation quantizations of hypertoric varieties	89
GWYN BELLAMY and TOSHIRO KUWABARA	
Almost factoriality of integral domains and Krull-like domains	129
GYU WHAN CHANG, HWANKOO KIM and JUNG WOOK LIM	
Singularities of free group character varieties	149
CARLOS FLORENTINO and SEAN LAWTON	
Energy identity for the maps from a surface with tension field bounded in L^p	181
LI JIAYU and ZHU XIANGRONG	
Remarks on some isoperimetric properties of the $k - 1$ flow	197
YU-CHU LIN and DONG-HO TSAI	
Demystifying a divisibility property of the Kostant partition function	215
KAROLA MÉSZÁROS	
Exceptional Lie algebras, $SU(3)$, and Jordan pairs	227
PIERO TRUINI	
Lower estimate of Milnor number and characterization of isolated homogeneous hypersurface singularities	245
STEPHEN S.-T. YAU and HUAIQING ZUO	