

*Pacific  
Journal of  
Mathematics*

**CALOGERO–MOSER VERSUS  
KAZHDAN–LUSZTIG CELLS**

CÉDRIC BONNAFÉ AND RAPHAËL ROUQUIER

Volume 261 No. 1

January 2013

## CALOGERO–MOSER VERSUS KAZHDAN–LUSZTIG CELLS

CÉDRIC BONNAFÉ AND RAPHAËL ROUQUIER

**In 1979, Kazhdan and Lusztig developed a combinatorial theory associated with Coxeter groups, defining in particular partitions of the group in left and two-sided cells. In 1983, Lusztig generalized this theory to Hecke algebras of Coxeter groups with unequal parameters. We propose a definition of left cells and two-sided cells for complex reflection groups, based on ramification theory for Calogero–Moser spaces. These spaces have been defined via rational Cherednik algebras by Etingof and Ginzburg. We conjecture that these coincide with Kazhdan–Lusztig cells, for real reflection groups. Counterparts of families of irreducible characters have been studied by Gordon and Martino, and we provide here a version of left cell representations. The Calogero–Moser cells will be studied in details in a forthcoming paper, providing thus several results supporting our conjecture.**

### 1. Introduction

Kazhdan and Lusztig [1979] developed a combinatorial theory associated with Coxeter groups. They defined in particular partitions of the group in left and two-sided cells. For Weyl groups, these have a representation-theoretic interpretation in terms of primitive ideals, and they play a key role in Lusztig’s description [1984] of unipotent characters for finite groups of Lie type. Lusztig [1983; 2003] generalized this theory to Hecke algebras of Coxeter groups with unequal parameters.

We propose a definition of left cells and two-sided cells for complex reflection groups, based on ramification theory for Calogero–Moser spaces. These spaces have been defined via rational Cherednik algebras by Etingof and Ginzburg [2002]. We conjecture that these coincide with Kazhdan–Lusztig cells, for real reflection groups. Counterparts of families of irreducible characters have been studied by Gordon and Martino [2009], and we provide here a version of left cell representations. The Calogero–Moser cells are studied in detail in [Bonnafé and Rouquier  $\geq$  2013].

---

*MSC2010:* 20C08.

*Keywords:* Hecke algebra, reflection group, Cherednik algebra, Kazhdan–Lusztig theory.

## 2. Calogero–Moser spaces and cells

**Rational Cherednik algebras at  $t = 0$ .** Let us recall some constructions and results from [Etingof and Ginzburg 2002]. Let  $V$  be a finite-dimensional complex vector space and  $W$  a finite subgroup of  $\mathrm{GL}(V)$ . Let  $\mathcal{S}$  be the set of reflections of  $W$ , that is, elements  $g$  such that  $\ker(g - 1)$  is a hyperplane. We assume that  $W$  is a reflection group, that is, it is generated by  $\mathcal{S}$ .

We denote by  $\mathcal{S}/\sim$  the quotient of  $\mathcal{S}$  by the conjugation action of  $W$  and we let  $\{\underline{c}_s\}_{s \in \mathcal{S}/\sim}$  be a set of indeterminates. We put  $A = \mathbb{C}[\mathbb{C}^{\mathcal{S}/\sim}] = \mathbb{C}[\{\underline{c}_s\}_{s \in \mathcal{S}/\sim}]$ . Given  $s \in \mathcal{S}$ , let  $v_s \in V$  and  $\alpha_s \in V^*$  be eigenvectors for  $s$  associated to the nontrivial eigenvalue.

The 0-rational Cherednik algebra  $\mathbf{H}$  is the quotient of  $A \otimes T(V \oplus V^*) \rtimes W$  by the relations

$$\begin{aligned} [x, x'] &= [\xi, \xi'] = 0, \\ [\xi, x] &= \sum_{s \in \mathcal{S}} \underline{c}_s \frac{\langle v_s, x \rangle \cdot \langle \xi, \alpha_s \rangle}{\langle v_s, \alpha_s \rangle} s \text{ for } x, x' \in V^* \text{ and } \xi, \xi' \in V. \end{aligned}$$

We put  $Q = Z(\mathbf{H})$  and  $P = A \otimes S(V^*)^W \otimes S(V)^W \subset Q$ . The ring  $Q$  is normal. It is a free  $P$ -module of rank  $|W|$ .

**Galois closure.** Let  $K = \mathrm{Frac}(P)$  and  $L = \mathrm{Frac}(Q)$ . Let  $M$  be a Galois closure of the extension  $L/K$  and  $R$  the integral closure of  $Q$  in  $M$ . Let  $G = \mathrm{Gal}(M/K)$  and  $H = \mathrm{Gal}(M/L)$ . Let  $\mathcal{P} = \mathrm{Spec} P = \mathbb{A}_{\mathbb{C}}^{\mathcal{S}/\sim} \times V/W \times V^*/W$ ,  $\mathcal{Q} = \mathrm{Spec} Q$  the Calogero–Moser space, and  $\mathcal{R} = \mathrm{Spec} R$ .

We denote by  $\pi : \mathcal{R} \rightarrow \mathcal{Q}$  the quotient by  $H$ , and by  $\Upsilon : \mathcal{Q} \rightarrow \mathcal{P}$  and  $\phi : \mathcal{P} \rightarrow \mathbb{A}_{\mathbb{C}}^{\mathcal{S}/\sim}$  the canonical maps. We put  $p = \Upsilon\pi : \mathcal{R} \rightarrow \mathcal{P}$  the quotient by  $G$ .

**Ramification.** Let  $\mathfrak{r} \in \mathcal{R}$  be a prime ideal of  $R$ . We denote by  $D(\mathfrak{r}) \subset G$  its decomposition group and by  $I(\mathfrak{r}) \subset D(\mathfrak{r})$  its inertia group.

We have a decomposition into irreducible components

$$\mathcal{R} \times_{\mathcal{P}} \mathcal{Q} = \bigcup_{g \in G/H} \mathbb{O}_g, \text{ where } \mathbb{O}_g = \{(x, \pi(g^{-1}(x))) \mid x \in \mathcal{R}\},$$

inducing a decomposition into irreducible components

$$V(\mathfrak{r}) \times_{\mathcal{P}} \mathcal{Q} = \bigsqcup_{g \in I(\mathfrak{r}) \backslash G/H} \mathbb{O}_g(\mathfrak{r}), \text{ where } \mathbb{O}_g(\mathfrak{r}) = \{(x, \pi(g^{-1}g'(x))) \mid x \in V(\mathfrak{r}), g' \in I(\mathfrak{r})\}.$$

**Undeformed case.** Let  $\mathfrak{p}_0 = \phi^{-1}(0) = \sum_{s \in \mathcal{S}/\sim} P_{\underline{c}_s}$ . We have

$$P/\mathfrak{p}_0 = \mathbb{C}[V \oplus V^*]^{W \times W}, \quad Q/\mathfrak{p}_0 Q = \mathbb{C}[V \oplus V^*]^{\Delta W},$$

where  $\Delta(W) = \{(w, w) \mid w \in W\} \subset W \times W$ . A Galois closure of the extension of  $\mathbb{C}(\mathfrak{p}_0 Q) = \mathbb{C}(V \oplus V^*)^{\Delta W}$  over  $\mathbb{C}(\mathfrak{p}_0) = \mathbb{C}(V \oplus V^*)^{W \times W}$  is  $\mathbb{C}(V \oplus V^*)^{\Delta Z(W)}$ .

Let  $\tau_0 \in \mathcal{R}$  above  $\mathfrak{p}_0$ . Since  $\mathfrak{p}_0 Q$  is prime, we have  $G = D(\tau_0)H = HD(\tau_0)$ ,  $I(\tau_0) = 1$ , and  $\mathbb{C}(r_0)$  is a Galois closure of the extension  $\mathbb{C}(\mathfrak{p}_0 Q)/\mathbb{C}(\mathfrak{p}_0)$ . Fix an isomorphism  $\iota : \mathbb{C}(\tau_0) \xrightarrow{\sim} \mathbb{C}(V \oplus V^*)^{\Delta Z(W)}$  extending the canonical isomorphism of  $\mathbb{C}(\mathfrak{p}_0 Q)$  with  $\mathbb{C}(V \oplus V^*)^{\Delta W}$ .

The application  $\iota$  induces an isomorphism  $D(\tau_0) \xrightarrow{\sim} (W \times W)/\Delta Z(W)$ , that restricts to an isomorphism  $D(\tau_0) \cap H \xrightarrow{\sim} \Delta W/\Delta Z(W)$ . This provides a bijection  $G/H \xrightarrow{\sim} (W \times W)/\Delta W$ . Composing with the inverse of the bijection

$$W \xrightarrow{\sim} (W \times W)/\Delta W, \quad w \mapsto (1, w),$$

we obtain a bijection  $G/H \xrightarrow{\sim} W$ .

From now on, we identify the sets  $G/H$  and  $W$  through this bijection. Note that this bijection depends on the choices of  $\tau_0$  and of  $\iota$ . Since  $M$  is the Galois closure of  $L/K$ , we have  $\bigcap_{g \in G} H^g = 1$ , hence the left action of  $G$  on  $W$  induces an injection  $G \subset \mathfrak{S}(W)$ .

### *Calogero–Moser cells.*

**Definition 2.1.** Let  $\tau \in \mathcal{R}$ . The  $\tau$ -cells of  $W$  are the orbits of  $I(\tau)$  in its action on  $W$ .

Let  $c \in \mathbb{A}_{\mathbb{C}}^{\mathcal{P}/\sim}$ . Choose  $\tau_c \in \mathcal{R}$  with  $\overline{p(\tau_c)} = \bar{c} \times 0 \times 0$ . The  $\tau_c$ -cells are called the *two-sided Calogero–Moser  $c$ -cells* of  $W$ . Choose now  $\tau_c^{\text{left}} \in \mathcal{R}$  contained in  $\tau_c$  with  $\overline{p(\tau_c^{\text{left}})} = \bar{c} \times V/W \times 0 \in \mathcal{P}$ . The  $\tau_c^{\text{left}}$ -cells are called the *left Calogero–Moser  $c$ -cells* of  $W$ . We have  $I(\tau_c^{\text{left}}) \subset I(\tau_c)$ . Consequently, every left cell is contained in a unique two-sided cell.

The map sending  $w \in W$  to  $\pi(w^{-1}(\tau_c))$  induces a bijection from the set of two-sided cells to  $\Upsilon^{-1}(c \times 0 \times 0)$ .

**Families and cell multiplicities.** Let  $E$  be an irreducible representation of  $\mathbb{C}[W]$ . We extend it to a representation of  $S(V) \rtimes W$  by letting  $V$  act by 0. Let

$$\Delta(E) = e \cdot \text{Ind}_{S(V) \rtimes W}^{\mathbf{H}}(A \otimes_{\mathbb{C}} E), \quad \text{where } e = \frac{1}{|W|} \sum_{w \in W} w,$$

be the spherical Verma module associated with  $E$ . It is a  $Q$ -module.

Let  $c \in \mathbb{A}_{\mathbb{C}}^{\mathcal{P}/\sim}$  and let  $\Delta^{\text{left}}(E) = (R/\tau_c^{\text{left}}) \otimes_P \Delta(E)$ .

**Definition 2.2.** Given a left cell  $\Gamma$ , we define the cell multiplicity  $m_{\Gamma}(E)$  of  $E$  as the length of  $\Delta^{\text{left}}(E)$  at the component  $\mathbb{C}_{\Gamma}(\tau_c^{\text{left}})$ .

Note that  $\sum_{\Gamma} m_{\Gamma}(E) \cdot [\mathbb{C}_{\Gamma}(\tau_c^{\text{left}})]$  is the support cycle of  $\Delta^{\text{left}}(E)$ .

There is a unique two-sided cell  $\Lambda$  containing all left cells  $\Gamma$  such that  $m_{\Gamma}(E) \neq 0$ . Its image in  $\mathcal{Q}$  is the unique  $\mathfrak{q} \in \Upsilon^{-1}(c \times 0 \times 0)$  such that  $(Q/\mathfrak{q}) \otimes_Q \Delta(E) \neq 0$ . The corresponding map  $\text{Irr}(W) \rightarrow \Upsilon^{-1}(c \times 0 \times 0)$  is surjective, and its fibers are the *Calogero–Moser families* of  $\text{Irr}(W)$ , as defined by Gordon [2003].

**Dimension 1.** Let  $V$  be a one-dimensional complex vector space, let  $d \geq 2$  and let  $W$  be the group of  $d$ -th roots of unity acting on  $V$ . Let  $\zeta = \exp(2i\pi/d)$ , let  $s = \zeta \in W$  and  $\underline{c}_i = \underline{c}_{s^i}$  for  $1 \leq i \leq d-1$ . We have  $A = \mathbb{C}[\underline{c}_1, \dots, \underline{c}_{d-1}]$  and

$$\mathbf{H} = A \left\langle x, \xi, s \mid sxs^{-1} = \zeta^{-1}x, \quad s\xi s^{-1} = \zeta\xi \text{ and } [\xi, x] = \sum_{i=1}^{d-1} \underline{c}_i s^i \right\rangle.$$

Let  $\text{eu} = \xi x - \sum_{i=1}^{d-1} (1 - \zeta^i)^{-1} \underline{c}_i s^i$ . We have  $P = A[x^d, \xi^d]$  and  $Q = A[x^d, \xi^d, \text{eu}]$ . Define  $\underline{\kappa}_1, \dots, \underline{\kappa}_d = \underline{\kappa}_0$  by  $\underline{\kappa}_1 + \dots + \underline{\kappa}_d = 0$  and  $\sum_{i=1}^{d-1} \underline{c}_i s^i = \sum_{i=0}^{d-1} (\underline{\kappa}_i - \underline{\kappa}_{i+1}) \varepsilon_i$ , where  $\varepsilon_i = \frac{1}{d} \sum_{j=0}^{d-1} \zeta^{ij} s^j$ . We have  $A = \mathbb{C}[\underline{\kappa}_1, \dots, \underline{\kappa}_d] / (\underline{\kappa}_1 + \dots + \underline{\kappa}_d)$ .

The normalization of the Galois closure is described as follows. There is an isomorphism of  $A$ -algebras

$$A[X, Y, Z] / \left( XY - \prod_{i=1}^d (Z - \underline{\kappa}_i) \right) \xrightarrow{\sim} Q, \quad X \mapsto x^d, \quad Y \mapsto \xi^d \quad \text{and} \quad Z \mapsto \text{eu}.$$

We have an isomorphism of  $A$ -algebras

$$A[X, Y, \lambda_1, \dots, \lambda_d] / \left( \begin{array}{l} e_i(\lambda) = e_i(\underline{\kappa}), \quad i = 1, \dots, d-1 \\ e_d(\lambda) = e_d(\underline{\kappa}) + (-1)^{d+1} XY \end{array} \right) \xrightarrow{\sim} R,$$

where  $Z = \lambda_d$  and where  $e_i$  denotes the  $i$ -th elementary symmetric function. We have  $G = \mathfrak{S}_d$ , acting by permuting the  $\lambda_i$ , and  $H = \mathfrak{S}_{d-1}$ .

Let  $\mathfrak{p}_0 = (\underline{\kappa}_1, \dots, \underline{\kappa}_d) \in \text{Spec } P$  and

$$\mathfrak{t}_0 = (\underline{\kappa}_1, \dots, \underline{\kappa}_d, \lambda_1 - \zeta \lambda_d, \dots, \lambda_{d-1} - \zeta^{d-1} \lambda_d) \in \text{Spec } R.$$

We have  $D(\mathfrak{t}_0) = \langle (1, 2, \dots, d) \rangle \subset \mathfrak{S}_d$  and

$$\mathbb{C}(\mathfrak{t}_0) = \mathbb{C}(X, Y, \lambda_d = \sqrt[d]{XY}) = \mathbb{C}(X, Y, Z = \sqrt[d]{XY}).$$

The composite bijection  $D(\mathfrak{t}_0) \xrightarrow{\sim} G/H \xrightarrow{\sim} W$  is an isomorphism of groups given by  $(1, \dots, d) \mapsto s$ .

Fix  $c \in \mathbb{C}^{d-1}$  and let  $\kappa_1, \dots, \kappa_d \in \mathbb{C}$  corresponding to  $c$ . Consider  $\mathfrak{t} = \mathfrak{t}_c$  or  $\mathfrak{t}_c^{\text{left}}$  as in Section 2 (see right after Definition 2.1). Then  $I(\mathfrak{t})$  is the subgroup of  $\mathfrak{S}_d$  stabilizing  $(\kappa_1, \dots, \kappa_d)$ . The left  $c$ -cells coincide with the two-sided  $c$ -cells and two elements  $s^i$  and  $s^j$  are in the same cell if and only if  $\kappa_i = \kappa_j$ . Finally, the multiplicity  $m_\Gamma(\det^j)$  is 1 if  $s^j \in \Gamma$  and 0 otherwise.

### 3. Coxeter groups

**Kazhdan–Lusztig cells.** Following [Kazhdan and Lusztig 1979; Lusztig 1983; 2003], let us recall the construction of cells.

We assume here  $V$  is the complexification of a real vector space  $V_{\mathbb{R}}$  acted on by  $W$ . We choose a connected component  $C$  of  $V_{\mathbb{R}} - \bigcup_{s \in \mathcal{S}} \ker(s - 1)$  and we

denote by  $S$  the set of  $s \in \mathcal{S}$  such that  $\ker(s - 1) \cap \bar{C}$  has codimension 1 in  $\bar{C}$ . This makes  $(W, S)$  into a Coxeter group, and we denote by  $l$  the length function.

Let  $\Gamma$  be a totally ordered free abelian group and let  $L : W \rightarrow \Gamma$  be a weight function, that is, a function such that

$$L(ww') = L(w) + L(w') \quad \text{if } l(ww') = l(w) + l(w').$$

We denote by  $v^\gamma$  the element of the group algebra  $\mathbb{Z}[\Gamma]$  corresponding to  $\gamma \in \Gamma$ .

We denote by  $H$  the Hecke algebra of  $W$ : this is the  $\mathbb{Z}[\Gamma]$ -algebra generated by elements  $T_s$  with  $s \in S$  subject to the relations

$$(T_s - v^{L(s)})(T_s + v^{-L(s)}) = 0 \quad \text{and} \quad \underbrace{T_s T_t T_s \cdots}_{m_{st} \text{ terms}} = \underbrace{T_t T_s T_t \cdots}_{m_{st} \text{ terms}},$$

for  $s, t \in S$  with  $m_{st} \neq \infty$ , where  $m_{st}$  is the order of  $st$ . Given  $w \in W$ , we put  $T_w = T_{s_1} \cdots T_{s_n}$ , where  $w = s_1 \cdots s_n$  is a reduced decomposition.

Let  $i$  be the ring involution of  $H$  given by  $i(v^\gamma) = v^{-\gamma}$  for  $\gamma \in \Gamma$  and  $i(T_s) = T_s^{-1}$ . We denote by  $\{C_w\}_{w \in W}$  the Kazhdan–Lusztig basis of  $H$ . It is uniquely defined by the properties that  $i(C_w) = C_w$  and  $C_w - T_w \in \bigoplus_{w' \in W} \mathbb{Z}[\Gamma_{<0}] T_{w'}$ .

We introduce the partial order  $<_L$  on  $W$ . It is the transitive closure of the relation given by  $w' <_L w$  if there is  $s \in S$  such that the coefficient of  $C_{w'}$  in the decomposition of  $C_s C_w$  in the Kazhdan–Lusztig basis is nonzero. We define  $w \sim_L w'$  to be the corresponding equivalence relation:  $w \sim_L w'$  if and only if  $w <_L w'$  and  $w' <_L w$ . The equivalence classes are the left cells. We define  $<_{LR}$  as the partial order generated by  $w <_{LR} w'$  if  $w <_L w'$  or  $w^{-1} <_L w'^{-1}$ . As above, we define an associated equivalence relation  $\sim_{LR}$ . Its equivalence classes are the two-sided cells.

When  $\Gamma = \mathbb{Z}$ ,  $L = l$ , and  $W$  is a Weyl group, a definition of left cells based on primitive ideals in enveloping algebras was proposed by Joseph [1980]: let  $\mathfrak{g}$  be a complex semisimple Lie algebra with Weyl group  $W$ . Let  $\rho$  be the half-sum of the positive roots. Given  $w \in W$ , let  $I_w$  be the annihilator in  $U(\mathfrak{g})$  of the simple module with highest weight  $-w(\rho) - \rho$ . Then,  $w$  and  $w'$  are in the same left cell if and only if  $I_w = I_{w'}$ .

**Representations and families.** Let  $\Gamma$  be a left cell. Let  $W_{\leq \Gamma}$  and  $W_{< \Gamma}$  be the sets of  $w \in W$  such that there is  $w' \in \Gamma$  with  $w <_L w'$  and, respectively,  $w <_L w'$  and  $w \notin \Gamma$ . The left cell representation of  $W$  over  $\mathbb{C}$  associated with  $\Gamma$  [Kazhdan and Lusztig 1979; Lusztig 2003] is the unique representation, up to isomorphism, that deforms into the left  $H$ -module

$$\left( \bigoplus_{w \in W_{\leq \Gamma}} \mathbb{Z}[\Gamma] C_w \right) / \left( \bigoplus_{w \in W_{< \Gamma}} \mathbb{Z}[\Gamma] C_w \right).$$

Lusztig [1982; 2003] has defined the set of constructible characters of  $W$  inductively as the smallest set of characters with the following properties: it contains the trivial character, it is stable under tensoring by the sign representation and it is stable under  $J$ -induction from a parabolic subgroup. Lusztig's families are the equivalence classes of irreducible characters of  $W$  for the relation generated by  $\chi \sim \chi'$  if  $\chi$  and  $\chi'$  occur in the same constructible character. Lusztig has determined constructible characters and families for all  $W$  and all parameters.

Lusztig has shown for equal parameters, and conjectured in general, that the set of left cell characters coincides with the set of constructible characters.

**A conjecture.** Let  $c \in \mathbb{R}^{\mathcal{G}/\sim}$ . Let  $\Gamma$  be the subgroup of  $\mathbb{R}$  generated by  $\mathbb{Z}$  and  $\{c_s\}_{s \in \mathcal{G}}$ . We endow it with the natural order on  $\mathbb{R}$ . Let  $L : W \rightarrow \Gamma$  be the weight function determined by  $L(s) = c_s$  if  $s \in S$ .

The following conjecture is due to Gordon and Martino [2009]. A similar conjecture has been proposed independently by the second author.<sup>1</sup> It is known to hold for types  $A_n$ ,  $B_n$ ,  $D_n$  and  $I_2(n)$  [Gordon 2008; Gordon and Martino 2009; Bellamy 2011; Martino 2010a; 2010b].

**Conjecture 3.1.** *The Calogero–Moser families of irreducible characters of  $W$  coincide with the Lusztig families.*

We propose now a conjecture involving partitions of elements of  $W$ , via ramification. The part dealing with left cell characters could be stated in a weaker way, using  $Q$  and not  $R$ , and thus not needing the choice of prime ideals, by involving constructible characters.

**Conjecture 3.2.** *There is a choice of  $\tau_c^{\text{left}} \subset \tau_c$  such that*

- *the Calogero–Moser two-sided cells and left cells coincide with the Kazhdan–Lusztig two-sided cells and left cells, respectively, and*
- *the representation  $\sum_{E \in \text{Irr}(W)} m_\Gamma(E)E$ , where  $\Gamma$  is a Calogero–Moser left cell, coincide with the left cell representation of the corresponding Kazhdan–Lusztig cell.*

Various particular cases and general results supporting Conjecture 3.2 are provided in [Bonnafé and Rouquier  $\geq$  2013]. In particular, the conjecture holds for  $W = B_2$ , for all choices of parameters.

## References

- [Bellamy 2011] G. Bellamy, “The Calogero–Moser partition for  $G(m, d, n)$ ”, preprint, 2011. arXiv 0911.0066
- [Bonnafé and Rouquier  $\geq$  2013] C. Bonnafé and R. Rouquier, “Cellules de Calogero–Moser”, in preparation.

<sup>1</sup>Talk at the Enveloping algebra seminar, Paris, December 2004.

- [Etingof and Ginzburg 2002] P. Etingof and V. Ginzburg, “Symplectic reflection algebras, Calogero–Moser space, and deformed Harish-Chandra homomorphism”, *Invent. Math.* **147**:2 (2002), 243–348. MR 2003b:16021 Zbl 1061.16032
- [Gordon 2003] I. Gordon, “Baby Verma modules for rational Cherednik algebras”, *Bull. London Math. Soc.* **35**:3 (2003), 321–336. MR 2004c:16050 Zbl 1042.16017
- [Gordon 2008] I. G. Gordon, “Quiver varieties, category  $\mathcal{O}$  for rational Cherednik algebras, and Hecke algebras”, *Int. Math. Res. Pap.* **2008**:3 (2008), Art. ID rpn006. MR 2010c:16032 Zbl 1168.16015
- [Gordon and Martino 2009] I. G. Gordon and M. Martino, “Calogero–Moser space, restricted rational Cherednik algebras and two-sided cells”, *Math. Res. Lett.* **16**:2 (2009), 255–262. MR 2010g:16052 Zbl 1178.16030
- [Joseph 1980] A. Joseph, “Goldie rank in the enveloping algebra of a semisimple Lie algebra, I, II”, *J. Algebra* **65**:2 (1980), 269–283, 284–306. MR 82f:17009 Zbl 0441.17004
- [Kazhdan and Lusztig 1979] D. Kazhdan and G. Lusztig, “Representations of Coxeter groups and Hecke algebras”, *Invent. Math.* **53**:2 (1979), 165–184. MR 81j:20066 Zbl 0499.20035
- [Lusztig 1982] G. Lusztig, “A class of irreducible representations of a Weyl group, II”, *Nederl. Akad. Wetensch. Indag. Math.* **44**:2 (1982), 219–226. MR 83h:20018 Zbl 0511.20034
- [Lusztig 1983] G. Lusztig, “Left cells in Weyl groups”, pp. 99–111 in *Lie group representations, I* (College Park, Md. 1982/1983), edited by R. Herb et al., Lecture Notes in Math. **1024**, Springer, Berlin, 1983. MR 85f:20035 Zbl 0537.20019
- [Lusztig 1984] G. Lusztig, *Characters of reductive groups over a finite field*, Annals of Mathematics Studies **107**, Princeton University Press, 1984. MR 86j:20038 Zbl 0556.20033
- [Lusztig 2003] G. Lusztig, *Hecke algebras with unequal parameters*, CRM Monograph Series **18**, American Mathematical Society, Providence, RI, 2003. MR 2004k:20011 Zbl 1051.20003
- [Martino 2010a] M. Martino, “Blocks of restricted rational Cherednik algebras for  $G(m, d, n)$ ”, preprint, 2010. arXiv 1009.3200
- [Martino 2010b] M. Martino, “The Calogero–Moser partition and Rouquier families for complex reflection groups”, *J. Algebra* **323**:1 (2010), 193–205. MR 2010j:16071 Zbl 1219.20006

Received March 6, 2012. Revised May 18, 2012.

CÉDRIC BONNAFÉ

INSTITUT DE MATHÉMATIQUES ET DE MODÉLISATION DE MONTPELLIER

UNIVERSITÉ MONTPELLIER 2

CASE COURRIER 051

34095 MONTPELLIER

FRANCE

cedric.bonnafe@math.univ-montp2.fr

<http://www.math.univ-montp2.fr/~bonnafe/>

RAPHAËL ROUQUIER

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF CALIFORNIA

BOX 951555

LOS ANGELES, CA 90095-1555

UNITED STATES

rouquier@math.ucla.edu

<http://www.math.ucla.edu>

MATHEMATICAL INSTITUTE

UNIVERSITY OF OXFORD

24-29 ST GILES’

OXFORD, OX1 3LB

UNITED KINGDOM



# PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
pacific@math.ucla.edu

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
balmer@math.ucla.edu

Don Blasius  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
blasius@math.ucla.edu

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
chari@math.ucr.edu

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
cooper@math.ucsb.edu

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
finn@math.stanford.edu

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
liu@math.ucla.edu

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
jhlu@maths.hku.hk

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
popa@math.ucla.edu

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
qing@cats.ucsc.edu

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
yang@math.princeton.edu

## PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2013 is US \$400/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 261 No. 1 January 2013

---

Hierarchies and compatibility on Courant algebroids	1
PAULO ANTUNES, CAMILLE LAURENT-GENGOUX and JOANA M. NUNES DA COSTA	
A new characterization of complete linear Weingarten hypersurfaces in real space forms	33
CÍCERO P. AQUINO, HENRIQUE F. DE LIMA and MARCO A. L. VELÁSQUEZ	
Calogero–Moser versus Kazhdan–Lusztig cells	45
CÉDRIC BONNAFÉ and RAPHAËL ROUQUIER	
Coarse median spaces and groups	53
BRIAN H. BOWDITCH	
Geometrization of continuous characters of $\mathbb{Z}_p^\times$	95
CLIFTON CUNNINGHAM and MASOUD KAMGARPOUR	
A note on Lagrangian cobordisms between Legendrian submanifolds of $\mathbb{R}^{2n+1}$	101
ROMAN GOLOVKO	
On slope genera of knotted tori in 4-space	117
YI LIU, YI NI, HONGBIN SUN and SHICHENG WANG	
Formal groups of elliptic curves with potential good supersingular reduction	145
ÁLVARO LOZANO-ROBLEDO	
Codimension-one foliations calibrated by nondegenerate closed 2-forms	165
DAVID MARTÍNEZ TORRES	
The trace of Frobenius of elliptic curves and the $p$ -adic gamma function	219
DERMOT MCCARTHY	
$(DN)$ - $(\Omega)$ -type conditions for Fréchet operator spaces	237
KRZYSZTOF PISZCZEK	



0030-8730(201301)261:1;1-D