

*Pacific
Journal of
Mathematics*

ORBIFOLDS WITH SIGNATURE $(0; k, k^{n-1}, k^n, k^n)$

ANGEL CAROCCA, RUBÉN A. HIDALGO AND RUBÍ E. RODRÍGUEZ

ORBIFOLDS WITH SIGNATURE $(0; k, k^{n-1}, k^n, k^n)$

ANGEL CAROCCA, RUBÉN A. HIDALGO AND RUBÍ E. RODRÍGUEZ

Two interesting problems that arise in the theory of closed Riemann surfaces are (i) computing algebraic curves representing the surface and (ii) deciding if the field of moduli is a field of definition.

In this paper we consider pairs (S, H) , where S is a closed Riemann surface and H is a subgroup of $\text{Aut}(S)$, the group of automorphisms of S , so that S/H is an orbifold with signature $(0; k, k^{n-1}, k^n, k^n)$ where $k, n \geq 2$ are integers.

In the case that S is the highest abelian branched cover of S/H we provide explicit algebraic curves representing S . In the case that k is an odd prime, we also describe algebraic curves for some intermediate abelian covers.

For $k = p \geq 3$ a prime and H a p -group, we prove that H is a p -Sylow subgroup of $\text{Aut}(S)$, and if $p \geq 7$ we prove that H is normal in $\text{Aut}(S)$. Also, when $n \neq 3$ we prove that the field of moduli in such cases is a field of definition. If, moreover, S is the highest abelian branched cover of S/H , then we compute explicitly the field of moduli.

1. Introduction

A closed Riemann surface S of genus $g \geq 2$ may be described by many different objects, for instance, by algebraic curves (by the Riemann–Roch theorem [Farkas and Kra 1992]), by torsion-free cocompact Fuchsian groups (by the Koebe–Poincaré uniformization theorem [Koebe 1907a; 1907b; Poincaré 1907]), by Schottky groups (by the retrosection theorem [Bers 1975; Koebe 1907b]), or by certain principally polarized abelian varieties (by the Torelli theorem [Torelli 1913; Weil 1956]). In general, to provide different explicit representations for the same Riemann surface has been a difficult problem, in spite of huge efforts to solve it. It seems that Burnside [1893] and Klein [1878] provided the first examples of algebraic curves and Fuchsian groups, both representing the same Riemann surfaces. In many cases, the group $\text{Aut}(S)$ of automorphisms of S and its subgroups play

Carocca and Rodríguez were supported by Fondecyt grants 1095165 and 1100767. Hidalgo was supported by Fondecyt grant 1110001 and UTFSM grant 12.11.01.

MSC2000: primary 30F10, 30F40; secondary 14H37.

Keywords: algebraic curves, Riemann Surfaces, automorphisms, field of moduli.

a fundamental role in finding algebraic curves representing S . For instance, if $S/\text{Aut}(S)$ has signature of the form $(0; r, s, t)$, then there are known examples having an explicit Fuchsian group and an explicit algebraic curve, both representing S [Burnside 1893; Klein 1878] (we also recommend reading [Karcher and Weber 1999]).

A *field of definition* of S is a subfield \mathbb{K} of \mathbb{C} for which it is possible to find an irreducible nonsingular projective algebraic curve representing S , defined by polynomials whose coefficients belong to \mathbb{K} . If C is an algebraic curve describing S , then the *field of moduli* of S is defined as the fixed field of the group of field automorphisms σ of \mathbb{C} such that C and C^σ are isomorphic, where C^σ is the algebraic curve defined as the zeroes of the polynomials obtained from the ones defining C after σ acts on their coefficients. The field of moduli is always contained in any field of definition, but it may happen that the field of moduli is not a field of definition.

In this article we study closed Riemann surfaces S admitting subgroups $H < \text{Aut}(S)$ so that S/H has signature $(0; k, k^{n-1}, k^n, k^n)$, where $n, k \geq 2$ are integers. For $k = 2$ this type of surface was considered in [Carvacho 2010; González-Diez and Hidalgo 1997] to provide examples of closed Riemann surfaces admitting topologically equivalent but conformally nonequivalent cyclic groups of order 2^n .

In the general case, if S is the homology cover of S/H , then we compute the field of moduli and we give explicit algebraic curves for S . These explicit algebraic curves for homology covers allow us to find algebraic curves for those Riemann surfaces S admitting an abelian group $G < \text{Aut}(S)$ such that S/G has signature $(0; k, k^{n-1}, k^n, k^n)$. We describe such a situation for the case that k is a prime and $G \cong \mathbb{Z}_k \times \mathbb{Z}_{k^n}$. Also, for k an odd prime, we describe the group $\text{Aut}(S)$ and we prove that the field of moduli of S is in fact a field of definition.

In this article we will use letters such as S, R, \tilde{S} to denote (closed) Riemann surfaces, orbifolds will usually be denoted using italic letters such as $\mathbb{O}, \tilde{\mathbb{O}}$ or as S/H (where S is a Riemann surface and $H < \text{Aut}(S)$), groups will be denoted by letters such as H, Γ, G , etc.

2. Preliminaries

2.1. Orbifolds. An orbifold is a tuple $\mathbb{O} = (S, \{(p_1, k_1), \dots, (p_n, k_n), \dots\})$ where (i) S is a Riemann surface, called the *Riemann surface structure* of \mathbb{O} , (ii) $\{p_1, p_2, \dots\} \subset S$ is a collection of different isolated points, called the *cone points* of \mathbb{O} , and (iii) each $k_j \geq 2$ is an integer, called the *cone order* of p_j . An orbifold of signature $(\gamma; k_1, \dots, k_n)$ is given by an orbifold $\mathbb{O} = (S, \{(p_1, k_1), \dots, (p_n, k_n)\})$ where S is a closed Riemann surface of genus γ . An orbifold without cone points is just a Riemann surface.

A *conformal homeomorphism* between two orbifolds, say $\mathbb{O}_1 = (S_1, \{(p_1, k_1), \dots, (p_n, k_n), \dots\})$ and $\mathbb{O}_2 = (S_2, \{(q_1, l_1), \dots, (q_n, l_n), \dots\})$, is a conformal homeomorphism between S_1 and S_2 (the corresponding Riemann surface structures), sending cone points to cone points, and preserving the cone point orders. If $\mathbb{O}_1 = \mathbb{O}_2 = \mathbb{O}$, then we speak about a *conformal automorphism* of the orbifold \mathbb{O} . We use the notation $\mathbb{O}_1 \cong \mathbb{O}_2$ to indicate that \mathbb{O}_1 and \mathbb{O}_2 are conformally equivalent orbifolds.

We denote by $\text{Aut}_{\text{orb}}(\mathbb{O})$ the group of conformal automorphisms of the orbifold \mathbb{O} . If S is the conformal Riemann surface structure of \mathbb{O} , then we denote by $\text{Aut}(S)$ its group of conformal automorphisms. There is a natural inclusion $\text{Aut}_{\text{orb}}(\mathbb{O}) < \text{Aut}(S)$, but in general these two groups are different.

If \mathbb{O} is an orbifold and $H < \text{Aut}_{\text{orb}}(\mathbb{O})$ acts discontinuously on the Riemann surface structure, then the quotient \mathbb{O}/H may be seen again as an orbifold as follows. We denote by $\pi : \mathbb{O} \rightarrow \mathbb{O}/H$ the canonical quotient map. A cone point of \mathbb{O}/H may be obtained in two different ways. In the first case, if $p \in \mathbb{O}$ is not a cone point and it has nontrivial H -stabilizer $H(p)$, then $\pi(p)$ is a cone point with order equal to the order of $H(p)$. In the second case, if $p \in \mathbb{O}$ is a cone point of order n and its H -stabilizer has order m , then $\pi(p)$ is a cone point with order equal to nm .

The orbifolds we consider in this paper are the *good orbifolds* in Thurston's terminology; they are obtained as quotient spaces $\mathbb{O} = \tilde{S}/F$, where \tilde{S} is a (not necessarily closed) Riemann surface and $F < \text{Aut}(\tilde{S})$ is a discontinuous group of conformal automorphisms of \tilde{S} . The cone points are those equivalence classes of points of \tilde{S} with nontrivial F -stabilizer.

2.2. Homology covers. Good orbifolds admit as (branched) universal cover either the Riemann sphere, the complex plane or the hyperbolic plane; this is a consequence of the classical uniformization theorem. Let us consider a good orbifold $\mathbb{O} = (S, \{(p_1, k_1), \dots, (p_n, k_n)\})$ of signature $(\gamma; k_1, \dots, k_n)$. The first (orbifold) fundamental group of \mathbb{O} is

$$(2-1) \quad \pi_1^{\text{orb}}(\mathbb{O}) = \left\langle \alpha_1, \dots, \alpha_\gamma, \beta_1, \dots, \beta_\gamma, \delta_1, \dots, \delta_n : \prod_{j=1}^{\gamma} [\alpha_j, \beta_j] \prod_{k=1}^n \delta_k = \delta_1^{k_1} = \dots = \delta_n^{k_n} = 1 \right\rangle,$$

where $\pi_1(S) = \langle \alpha_1, \dots, \alpha_\gamma, \beta_1, \dots, \beta_\gamma : \prod_{j=1}^{\gamma} [\alpha_j, \beta_j] = 1 \rangle$, with $[a, b] = aba^{-1}b^{-1}$, and the element δ_j represents a simple small loop around p_j in $S - \{p_1, \dots, p_n\}$, for each $j = 1, \dots, n$.

It is clear that to each normal subgroup N of finite index of $\pi_1^{\text{orb}}(\mathbb{O})$ there corresponds an orbifold $\tilde{\mathbb{O}}$ and a finite group $H < \text{Aut}_{\text{orb}}(\tilde{\mathbb{O}})$, so that $\mathbb{O} = \tilde{\mathbb{O}}/H$. Observe that H is isomorphic to $\pi_1^{\text{orb}}(\mathbb{O})/N$. When $N = \pi_1^{\text{orb}}(\mathbb{O})'$ (the derived subgroup of $\pi_1^{\text{orb}}(\mathbb{O})$), the corresponding cover orbifold $\tilde{\mathbb{O}}$ is called the *homology*

orbifold cover of \mathbb{O} . We will be interested only in the particular case when the homology orbifold cover is a closed Riemann surface (i.e., there are no cone points), in which case we call it the *homology cover* of \mathbb{O} , and say that \mathbb{O} is a *homology orbifold*.

Clearly, the homology orbifold cover of \mathbb{O} is the homology cover if and only if $\pi_1^{\text{orb}}(\mathbb{O})'$ has finite index in $\pi_1^{\text{orb}}(\mathbb{O})$ and it acts freely on the universal cover space of \mathbb{O} . The finite index condition is equivalent to the condition that the underlying Riemann surface structure of \mathbb{O} is the Riemann sphere; that is, $\gamma = 0$. The free action condition is equivalent to the following one.

Theorem 1 [Maclachlan 1965]. *Let \mathbb{O} be an orbifold of signature $(\gamma; k_1, \dots, k_n)$. Then $\pi_1^{\text{orb}}(\mathbb{O})'$ is torsion-free if and only if*

$$(2-2) \quad \text{lcm}(k_1, \dots, k_{j-1}, k_{j+1}, \dots, k_n) = \text{lcm}(k_1, \dots, k_n) \quad \text{for all } j = 1, \dots, n.$$

The homology cover (when it exists) is the highest abelian Galois cover of \mathbb{O} .

2.3. Fuchsian groups. The basic theory of Fuchsian groups may be found, for instance, in the classical book [Beardon 1983]. A *cocompact Fuchsian group* acting on the upper half-plane \mathbb{H}^2 is a discrete group $\Gamma < \text{PSL}(2, \mathbb{R})$ such that \mathbb{H}^2 / Γ is an orbifold of some signature; that is, the underlying Riemann surface is a closed Riemann surface. It is known that a cocompact Fuchsian group Γ has a presentation of the form

$$(2-3) \quad \Gamma = \left\langle a_1, b_1, \dots, a_\gamma, b_\gamma, \delta_1, \dots, \delta_n : \prod_{j=1}^{\gamma} [a_j, b_j] \prod_{j=1}^n \delta_j = \delta_1^{k_1} = \dots = \delta_n^{k_n} = 1 \right\rangle,$$

where γ and n are nonnegative integers, the $k_j \geq 2$ are integers, and $2\gamma - 2 + n - \sum_{j=1}^n k_j^{-1} > 0$. The tuple $(\gamma; k_1, \dots, k_n)$ is known as the *signature* of Γ (this is the signature of its quotient orbifold \mathbb{H}^2 / Γ).

An orbifold \mathbb{O} is *of hyperbolic type* if there is a cocompact Fuchsian group Γ so that $\mathbb{O} \cong \mathbb{H}^2 / \Gamma$. By the Poincaré–Koebe uniformization theorem [Koebe 1907a; 1907b; Poincaré 1907], every orbifold with signature $(\gamma; k_1, \dots, k_n)$ is of hyperbolic type if and only if $2\gamma - 2 + n - \sum_{j=1}^n k_j^{-1} > 0$.

By the hyperbolic area of a Fuchsian group Γ (respectively, of a hyperbolic orbifold) of signature $(\gamma; k_1, \dots, k_n)$ we refer to the hyperbolic area of a fundamental polygon domain for it; it is given by

$$(2-4) \quad A(\Gamma) = 2\pi \left(2\gamma - 2 + \sum_{j=1}^n \left(1 - \frac{1}{k_j} \right) \right).$$

We say that a cocompact Fuchsian group Γ , with presentation (2-3), is a *homology Fuchsian group* if $\gamma = 0$ and it satisfies Maclachlan's conditions (2-2). In other

words, homology Fuchsian groups are exactly those cocompact Fuchsian groups providing a Fuchsian uniformization of a hyperbolic homology orbifold of genus zero. If Γ is a homology Fuchsian group of signature $(0; k_1, \dots, k_n)$, then the homology cover of the homology orbifold $\mathbb{O} = \mathbb{H}^2 / \Gamma$ is $S = \mathbb{H}^2 / \Gamma'$, where Γ' denotes the derived subgroup of Γ .

2.4. Fields of moduli and fields of definition. As a consequence of the implicit function theorem, every irreducible nonsingular projective algebraic curve defines a closed Riemann surface; conversely, by the Riemann–Roch theorem, every closed Riemann surface may be described by an irreducible nonsingular projective algebraic curve. It is this equivalence which allows the work in the analytical and in the algebraic settings in a parallel way.

Let C be an irreducible nonsingular projective algebraic curve, say defined by homogeneous polynomials P_1, \dots, P_r , each one with coefficients in a subfield $\mathbb{K} < \mathbb{C}$. Let g denote the genus of the closed Riemann surface corresponding to C . If $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q})$, the group of field automorphisms of \mathbb{C} , then we may consider the new polynomials $P_1^\sigma, \dots, P_r^\sigma$, where the coefficients of P_j^σ are the corresponding images under σ of the coefficients of the original polynomial P_j . The algebraic curve C^σ , defined by these new polynomials, is still an irreducible nonsingular projective algebraic curve, and it defines a new closed Riemann surface of genus g . It is not difficult to see that if \tilde{C} is another irreducible nonsingular projective algebraic curve that is birationally equivalent to C , then C^σ and \tilde{C}^σ are also birationally equivalent. Therefore, a natural action of $\text{Aut}(\mathbb{C}/\mathbb{Q})$ is defined on the moduli space of genus g . The stabilizer of the moduli class of C under such action is the subgroup

$$K_C = \{\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}) : C \cong C^\sigma\} < \text{Aut}(\mathbb{C}/\mathbb{Q}).$$

The fixed field of K_C , denoted by $\mathbb{M}(C)$, is called the *field of moduli* of C .

A subfield \mathbb{K} of \mathbb{C} is called a *field of definition* of C if there is an irreducible nonsingular projective algebraic curve \tilde{C} defined over \mathbb{K} which is birationally equivalent to C . At this point it is important to note that it is not clear that given a field of definition $\mathbb{L} < \mathbb{C}$ of C there is a smaller subfield $\mathbb{F} < \mathbb{L}$ which is again a field of definition of C .

The field of moduli $\mathbb{M}(C)$ is contained in any field of definition of C , and it coincides with the intersection of all fields of definition of C [Koizumi 1972]. Moreover, there is a field of definition of C which is an extension of finite degree of the field of moduli [Dèbes and Emsalem 1999; Hammer and Herrlich 2003].

If $g = 0$, then $C \cong \mathbb{P}^1$, so in this case $\mathbb{M}(C) = \mathbb{Q}$ is a field of definition. If $g = 1$, then C is equivalent to an (affine) elliptic curve $E_\eta = \{y^2 = x(x-1)(x-\eta)\}$, where $\eta \in \mathbb{C} - \{0, 1\}$. If $j(\eta) = (1 - \eta + \eta^2)^3 / \eta^2(\eta - 1)^2$ is its j -invariant and

$$a(\eta) = \frac{27j(\eta)}{j(\eta) - 1},$$

then E_η is also described by $D_\eta = \{y^2 = 4x^3 - a(\eta)x - a(\eta)\}$. It follows that $\mathbb{Q}(j(\eta))$ is a field of definition for E_η . Moreover, if $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q})$ and $E_\eta^\sigma = E_{\sigma(\eta)}$ is conformally equivalent to E_η , then they must have the same j -invariant; that is, $\sigma(j(\eta)) = j(\eta)$. It follows that $\mathbb{M}(C) = \mathbb{M}(E_\eta) = \mathbb{Q}(j(\eta))$ is also a field of definition.

In genus $g \geq 2$, the situation is more difficult. There are examples for which the field of moduli is not a field of definition [Earle 1971; Huggins 2007; Shimura 1972]; all of the examples there are hyperelliptic curves. It is stated in [Earle 1971] that there are examples of nonhyperelliptic Riemann surfaces with the same properties, but no explicit one is given. An explicit example of a nonhyperelliptic Riemann surface of genus $g = 17$ which cannot be defined over \mathbb{R} and whose field of moduli lies inside \mathbb{R} is given in [Hidalgo 2009] (this example is related to the hyperelliptic example in [Earle 1971]).

A. Weil [1956] provided the following sufficient and necessary conditions for the moduli field to be a field of definition.

Theorem 2 [Weil 1956]. *Let C be an irreducible nonsingular projective algebraic curve defined over a finite Galois extension \mathbb{L} of its field of moduli $\mathbb{M}(C)$. If for every $\sigma \in \text{Aut}(\mathbb{L}/\mathbb{M}(C))$ there is a biholomorphism $f_\sigma : C \rightarrow C^\sigma$ defined over \mathbb{L} such that the compatibility condition $f_{\tau\sigma} = f_\sigma^\tau \circ f_\tau$ holds for all $\sigma, \tau \in \text{Aut}(\mathbb{L}/\mathbb{M}(C))$, then there exists an irreducible nonsingular projective algebraic curve E defined over $\mathbb{M}(C)$ and there exists a biregular map $F : C \rightarrow E$, defined over \mathbb{L} , such that $F^\sigma \circ f_\sigma = F$.*

As a consequence of Theorem 2, it follows that if C has no nontrivial automorphism, then it may be defined over its field of moduli. Unfortunately, if C has nontrivial automorphisms, then it is a very difficult task to check whether Weil's conditions hold. But if $C/\text{Aut}(C)$ has signature of the form $(0; a, b, c)$ (quasiplatonic surfaces, or platonic if some cone order is equal to 2), then C may be defined over its field of moduli [Coombes and Harbater 1985; Wolfart 2006].

Consider a (branched) holomorphic covering between closed Riemann surfaces, say $f : S \rightarrow R$. Assume S and R are given by fixed algebraic curves and that R is defined over $\mathbb{M}(S)$. For each $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{M}(S))$ we may consider the (branched) holomorphic covering $f^\sigma : S^\sigma \rightarrow R^\sigma = R$. We say that they are *equivalent coverings*, denoted by $\{f^\sigma : S^\sigma \rightarrow R\} \cong \{f : S \rightarrow R\}$, if there is a holomorphic isomorphism $\phi_\sigma : S \rightarrow S^\sigma$ so that $f^\sigma \circ \phi_\sigma = f$. The *field of moduli* of $f : S \rightarrow R$, denoted by $\mathbb{M}(f : S \rightarrow R)$, is the fixed field of the subgroup

$$K(f : S \rightarrow R) = \{\sigma \in \text{Aut}(\mathbb{C}/\mathbb{M}(S)) : \{f^\sigma : S^\sigma \rightarrow R\} \cong \{f : S \rightarrow R\}\}.$$

It is clear from the definition that $\mathbb{M}(S) < \mathbb{M}(f : S \rightarrow R)$, but in general they may be different fields. For the particular case that $R = S / \text{Aut}(S)$ and S has genus at least two, the following is well known (a direct consequence of [Theorem 2](#)).

Theorem 3 [[Dèbes and Emsalem 1999](#)]. *If C is an irreducible nonsingular projective algebraic curve of genus $g \geq 2$, then there exists an irreducible nonsingular projective algebraic curve C_1 , defined over $\mathbb{M}(C)$, and there exists a Galois cover $f : C \rightarrow C_1$, with $\text{Aut}(C)$ as deck group, so that $\mathbb{M}(f : C \rightarrow C_1) = \mathbb{M}(C)$. Moreover, if $(C_1)_f$ denotes the branch locus of f and if $C_1 - (C_1)_f$ contains at least one $\mathbb{M}(C)$ -rational point, then $\mathbb{M}(C)$ is also a field of definition of C . Such a curve C_1 is called a canonical model of $C / \text{Aut}(C)$.*

3. Main results

Let S be a closed Riemann surface and let $H_1, H_2 < \text{Aut}(S)$. We say that H_1 and H_2 are (weakly) *topologically equivalent* (respectively, *conformally equivalent*) if there is an orientation preserving self-homeomorphism (respectively, conformal automorphism) $h : S \rightarrow S$ so that $H_2 = fH_1f^{-1}$. If $H < \text{Aut}(S)$, then we denote by $\text{Aut}_H(S)$ the normalizer of H in $\text{Aut}(S)$.

3.1. p -groups of automorphisms. A *regular cover* of an orbifold \mathbb{O} is a closed Riemann surface S together with a group H of conformal automorphisms such that the quotient orbifold S/H is isomorphic to \mathbb{O} . In the case that H is an abelian group, we say that the regular cover is an *abelian cover* of the orbifold. In this section we consider regular p^{n+1} -covers of orbifolds of type $(0; p, p^{n-1}, p^n, p^n)$, where $n \geq 2$ and p is an odd prime; that is, H is a p -group of order p^{n+1} . The interest in this type of example is that examples were constructed in [[Carvacho 2010](#); [González-Diez and Hidalgo 1997](#)] of closed Riemann surfaces S admitting topologically equivalent but conformally nonequivalent cyclic groups of order 2^{n+1} , where $n \geq 2$, so the quotient of S by the 2-group generated by these two cyclic subgroups is an orbifold with signature $(0; 2, 2^n, 2^{n+1}, 2^{n+1})$.

Let S be a closed Riemann surface and let $H < \text{Aut}(S)$ be a p -group such that S/H has signature of the form $(0; p, p^{n-1}, p^n, p^n)$, with $n \geq 2$, and consider the regular branched cover $P : S \rightarrow \widehat{\mathbb{C}}$, with H as deck group.

Since $n \geq 3$, then (up to left composition by a suitable Möbius transformation) we may assume that the branch values of P are ∞ of order p , 0 of order p^{n-1} , and 1 and some $\lambda \in \mathbb{C} - \{0, 1\}$ are the ones of order p^n . The choice of λ is not unique, but the only other possible choice is $1/\lambda$.

Theorem 4. *Let $p \geq 3$ be a prime and let $n \geq 2$ be an integer. Consider a closed Riemann surface S with a subgroup $H < \text{Aut}(S)$ such that H is a p -group with S/H of signature $(0; p, p^{n-1}, p^n, p^n)$. Let $\lambda \in \mathbb{C} - \{0, 1\}$ be as defined above. Then:*

- (1) H is a p -Sylow subgroup of $\text{Aut}(S)$. In particular, if $H_1, H_2 < \text{Aut}(S)$ are p -groups with S/H_j of signature $(0; p, p^{n-1}, p^n, p^n)$, then H_1 and H_2 are conformally equivalent.
- (2) If $n \geq 3$, then
- (a) $\text{Aut}_H(S) = H$ for $\lambda \neq -1$,
 - (b) $[\text{Aut}_H(S) : H] \in \{1, 2\}$ for $\lambda = -1$.
- (3) If $n = 2$, then
- (a) $[\text{Aut}_H(S) : H] \in \{1, 2\}$ for $\lambda \neq -1$,
 - (b) $[\text{Aut}_H(S) : H] \in \{1, 2, 4\}$ for $\lambda = -1$.
- (4) If $p \geq p_0$, where
- (a) $p_0 = 7$ for $n = 2$, and
 - (b) $p_0 = 5$ for $n \geq 3$,
- then $\text{Aut}_H(S) = \text{Aut}(S)$.

Remark 5. In the case $\lambda = -1$ and $n \geq 3$, part (2) of [Theorem 4](#) asserts that either $\text{Aut}_H(S) = H$ or $[\text{Aut}_H(S) : H] = 2$. In the last case, $S/\text{Aut}_H(S)$ has signature $(0; 2p, 2p^{n-1}, p^n)$, which is a maximal signature [[Singerman 1972](#)], so $\text{Aut}_H(S) = \text{Aut}(S)$.

3.2. Normality condition. Let S be a closed Riemann surface and $H < \text{Aut}(S)$. Let $\mathcal{M}(S, H)$ denote the locus in the moduli space $\mathcal{M}(S)$ of S consisting of those classes of Riemann surfaces \widehat{S} admitting a group \widehat{H} of conformal automorphisms, which is topologically equivalent to H . In general, one should expect that $\mathcal{M}(S, H)$ is a singular variety. The following shows that this is not the case if H is a p -group and S/H has signature $(0; p, p^{n-1}, p^n, p^n)$.

Corollary 6. Let $p \geq 3$ be a prime and let $n \geq 2$ be an integer. Consider a closed Riemann surface S and let $H < \text{Aut}(S)$ be a p -group such that S/H has signature $(0; p, p^{n-1}, p^n, p^n)$. Then $\mathcal{M}(S, H)$ is a normal subvariety of $\mathcal{M}(S)$.

Proof. The normality condition for $\mathcal{M}(S, H)$ is equivalent to the following property: given any two pairs (S_1, H_1) and (S_2, H_2) , where S_j is a closed Riemann surface (of the same genus as S) and H_j is a p -group of conformal automorphisms of S_j so that S_j/H_j has signature $(0; p, p^{n-1}, p^n, p^n)$, and there is an orientation preserving homeomorphism $f : S_1 \rightarrow S_2$ with $fH_1f^{-1} = H_2$, then f may be replaced by a biholomorphism with the same properties. This property is exactly what part (1) of [Theorem 4](#) states. \square

3.3. Homology rigidity.

Corollary 7. Every orbifold of signature $(0; p, p^{n-1}, p^n, p^n)$, where $p \geq 3$ is a prime and $n \geq 2$ is an integer, is uniquely determined, up to conformal equivalence, by its homology cover Riemann surface.

Proof. A consequence of part (1) of [Theorem 4](#). \square

Remark 8 (Torelli's theorem). Let \mathbb{O} be an orbifold of signature $(0; p, p^{n-1}, p^n, p^n)$, where $p \geq 3$ is a prime and $n \geq 2$ is an integer. As any two homology covers of \mathbb{O} are conformally equivalent Riemann surfaces, we may define the Jacobian of \mathbb{O} , denoted by $J(\mathbb{O})$, as the Jacobian of any of these covers. It follows that $J(\mathbb{O})$ is uniquely determined, up to equivalence of principally polarized abelian varieties, by \mathbb{O} . As a consequence of Torelli's theorem, $J(\mathbb{O})$ determines the conformal class of the homology cover of \mathbb{O} and, by [Corollary 7](#), it also determines the conformal class of \mathbb{O} . In this way, a kind of Torelli's theorem is obtained for this class of orbifolds. We may wonder how to describe the Jacobian of \mathbb{O} in terms of multivalued holomorphic differential forms so that it looks more similar to the construction for the case of Riemann surfaces. In order to do this, we use as homology the orbifold homology group

$$H_1^{\text{orb}}(\mathbb{O}) = \pi_1^{\text{orb}}(\mathbb{O}) / \pi_1^{\text{orb}}(\mathbb{O})',$$

and as holomorphic forms those multivalued holomorphic forms whose liftings to the homology cover define the holomorphic one forms of it.

3.4. Algebraic curves in the abelian case. Curves for the hyperelliptic homology covers and for the homology covers of homology orbifolds with triangular signature have been described in [\[Hidalgo 2012\]](#). Algebraic curves for the homology covers of orbifolds with signature of the form $(0; k, \dots, k)$ have been obtained in [\[González-Diez et al. 2009\]](#). We next provide the algebraic curves for the homology covers of orbifolds with signature $(0; k, k^{n-1}, k^n, k^n)$, where $k, n \geq 2$ are integers. As a consequence of the results in [\[Hidalgo 2012\]](#), the homology covers of such orbifolds cannot be hyperelliptic. Note that if R is the homology cover of such an orbifold \mathbb{O} , then $\mathbb{O} = R/H$, where $H \cong \mathbb{Z}_k \times \mathbb{Z}_{k^{n-1}} \times \mathbb{Z}_{k^n}$.

Theorem 9. *Let $k, n \geq 2$ be integers and let \mathbb{O} be an orbifold with signature $(0; k, k^{n-1}, k^n, k^n)$. Denote by R a homology cover of \mathbb{O} , let $H < \text{Aut}(R)$ be so that $R/H = \mathbb{O}$, and let $P : R \rightarrow \mathbb{O}$ be the Galois cover with H as deck group. We may assume (up to a Möbius transformation) that the cone points of \mathbb{O} (that is, the branch values of P) are given by the points $0, 1, \infty$ and $\lambda \in \mathbb{C} - \{0, 1\}$. We may also assume that ∞ is the cone point of order k , that 0 is the cone point of order k^{n-1} and that 1 and λ are the cone points of order k^n .*

Then R is represented by the (singular) projective algebraic curve

$$C_\lambda : \left\{ \begin{array}{l} z_0^k z_3^{k^n - k} + z_1^{k^{n-1}} z_3^{k^n - k^{n-1}} + z_2^{k^n} = 0 \\ \lambda z_0^k z_3^{k^{n-1} - k} + z_1^{k^{n-1}} + z_3^{k^{n-1}} = 0 \end{array} \right\} \subset \mathbb{P}^3,$$

H is generated by the projective linear transformations

$$\begin{aligned} a_0([z_0 : z_1 : z_2 : z_3]) &= [\rho_1 z_0 : z_1 : z_2 : z_3], \\ b_0([z_0 : z_1 : z_2 : z_3]) &= [z_0 : \rho_{n-1} z_1 : z_2 : z_3], \\ c_0([z_0 : z_1 : z_2 : z_3]) &= [z_0 : z_1 : \rho_n z_2 : z_3], \end{aligned}$$

where $\rho_s = e^{2\pi i/k^s}$, for each positive integer s , and the branched covering map P is represented in this model by

$$P([z_0 : z_1 : z_2 : z_3]) = -\left(\frac{z_1^{k^{n-1}}}{z_0^k z_3^{k^{n-1}-k}}\right).$$

The only singular point of the above curve is $[1 : 0 : 0 : 0]$.

Theorem 9 may be used to find algebraic curves for closed Riemann surfaces S admitting an abelian group $G < \text{Aut}(S)$ whose quotient orbifold S/G has signature of the form $(0; k, k^{n-1}, k^n, k^n)$. In fact, let $Q : S \rightarrow S/G = \mathbb{C}$ be a regular abelian branched cover with G as deck group. Let R be the homology cover of \mathbb{C} , let $P : R \rightarrow \mathbb{C}$ be the regular abelian branched cover, with deck group $H < \text{Aut}(R)$. Then there exists a subgroup $K < H$, acting freely on R and so that $G \cong H/K$, and there exists a regular unbranched cover $F : R \rightarrow S$, with K as deck group, satisfying $P = Q \circ F$. As we have explicit curves for R and an explicit presentation for H , the classical invariant theory permits us to obtain explicit algebraic curves for S and an explicit presentation of G . We show an application in the next section.

3.5. Families with Galois group of order p^{n+1} . As mentioned before, we are interested in regular p^{n+1} -covers of orbifolds of type $(0; p, p^{n-1}, p^n, p^n)$, where $n \geq 2$ and p is an odd prime. In **Section 9** we will see that the algebraic structure of the corresponding groups of order p^{n+1} is restricted to only two algebraic types: a direct or a semidirect product of \mathbb{Z}_{p^n} and \mathbb{Z}_p . The geometric types (classified by either geometric signature or generating vector for the corresponding action) are more varied: four different types are found in each algebraic case.

We study the corresponding families of Riemann surfaces, giving their algebraic curves in the abelian case.

The next result makes the above more explicit for the case when $G \cong \mathbb{Z}_p \times \mathbb{Z}_{p^n}$, where p is a prime. As we will see in its proof, this is a heavy computational procedure, but not a hard one.

Theorem 10. *Let S be a closed Riemann surface admitting a group $G < \text{Aut}(S)$ such that $G = \langle A, B : A^p = B^{p^n} = [A, B] = 1 \rangle \cong \mathbb{Z}_p \times \mathbb{Z}_{p^n}$ and $\mathbb{C} = S/G$ is an orbifold with signature $(0; p, p^{n-1}, p^n, p^n)$, where $n \geq 2$ and p is an odd prime. Let R be a homology cover of \mathbb{C} , let $H < \text{Aut}(R)$ be so that $R/H = \mathbb{C}$. Let $K < H$ be the normal subgroup so that $S = R/K$ and $G = H/K$.*

(1) If $K \cong \mathbb{Z}_{p^{n-1}}$, there exist $\beta \in \{1, 2, \dots, p^{n-1} - 1\}$, $\alpha \in \{0, 1, \dots, p - 1\}$ and $q \in \{1, \dots, [(p^n - 1)/p]\}$, with $(\beta, p) = 1 = (p, q)$, such that a (singular) projective algebraic curve representation of S is given by one of the following two families.

(a) If $\alpha = 0$, there exists λ in \mathbb{C} , with $\lambda \neq 0, 1$, such that

$$S : \left\{ \begin{array}{l} (\lambda - 1)w_0^p - w_1^p + w_3^p = 0 \\ (-1)^{q+1}(w_0^p + w_1^p)^q w_1^{p^{n-1}-\beta} + w_2^{p^{n-1}} w_3^{q_{p-\beta}} = 0 \end{array} \right\} \subset \mathbb{P}^3$$

and the action of G is generated by the projective linear transformations

$$\begin{aligned} A([w_0 : w_1 : w_2 : w_3]) &= [\rho_1 w_0 : w_1 : w_2 : w_3], \\ B([w_0 : w_1 : w_2 : w_3]) &= [w_0 : \rho_1 w_1 : \rho_n^{p^{n-1}-\beta} w_2 : w_3], \end{aligned}$$

where $\rho_k = e^{2\pi i/p^k}$. The regular branched covering map $Q : S \rightarrow S/G$ in this model is represented by

$$Q([w_0 : w_1 : w_2 : w_3]) = \frac{w_0^p + w_1^p}{w_0^p}.$$

The singular points of the above curve are given by the $(p + 1)$ points $[0 : 0 : 1 : 0]$ and $[1 : 0 : 0 : (1 - \lambda)^{1/p}]$.

(b) If $\alpha > 0$, there exists λ in \mathbb{C} , with $\lambda \neq 0, 1$, such that

$$S : \left\{ \begin{array}{l} v_1^{p^{n-1}} + \frac{(-1)^{q+1}}{(\lambda - 1)^q} (\lambda v_1^p - v_3^p)^q v_1^{p^{n-1}-\beta} v_3^{\beta-pq} = 0 \\ v_2^p v_3^{p(p^r-\beta)+\alpha p-p} + \frac{(-1)^{\alpha+1}}{(\lambda - 1)^{\alpha+p^r-\beta}} (v_0^p - v_3^p)^{p^r-\beta} (\lambda v_0^p - v_3^p)^\alpha = 0 \end{array} \right\} \subset \mathbb{P}^3$$

and the group G is generated by the transformations

$$\begin{aligned} A([v_0 : v_1 : v_2 : v_3]) &= [v_0 : v_1 : \rho_1^{p^r-\beta} v_2 : v_3], \\ B([v_0 : v_1 : v_2 : v_3]) &= [\rho_n^{p^{n-1}} v_0 : \rho_n^{p^{n-1}-\beta} v_1 : v_2 : v_3]. \end{aligned}$$

The regular branched covering map $Q : S \rightarrow S/G$ in this model is represented by

$$Q([v_0 : v_1 : v_2 : v_3]) = \frac{\lambda v_0^p - v_3^p}{v_0^p + v_3^p}.$$

(2) If $K \cong \mathbb{Z}_{p^{n-2}} \times \mathbb{Z}_p$, there exist λ in \mathbb{C} , with $\lambda \neq 0, 1$, and integers $\gamma, v \in \{1, \dots, p - 1\}$ such that a (singular) projective algebraic curve representation of S

is provided by the plane projective curve

$$\left\{ \frac{(-1)^{p^{n-1}(p-\gamma)}}{\lambda^{p^{n-1}(p-\gamma)+1}} (u_0^p + u_2^p)^{p^{n-1}(p-\gamma)} u_1^{p^2 v} ((\lambda - 1)u_0^p - u_2^p) + u_1^{p^n} u_2^{p^n(p-\gamma-1)+p+p^2 v} = 0 \right\} \subset \mathbb{P}^2,$$

and the group G is generated by the transformations

$$A([u_0 : u_1 : u_2]) = [\rho_1 u_0 : u_1 : u_2], \quad B([u_0 : u_1 : u_2]) = [u_0 : \rho_n u_1 : u_2].$$

The regular branched covering map $Q : S \rightarrow S/G$ in this model is represented by

$$Q([u_0 : u_1 : u_2]) = \frac{\lambda u_0^p}{u_0^p + u_1^p}.$$

3.6. Field of moduli. If S is a closed Riemann surface, then it follows from the Riemann–Roch theorem that S may be described by an irreducible nonsingular projective algebraic curve C . It is clear from the definition that we may define the field of moduli of S as the field of moduli of C and a field of definition of S as a field of definition of C .

Theorem 11. *Let $p \geq 3$ be a prime, $n \geq 3$ be an integer, S be a closed Riemann surface, and $H < \text{Aut}(S)$ be a p -group with S/H of signature $(0; p, p^{n-1}, p^n, p^n)$. Then S may be defined over its field of moduli.*

Remark 12. Under the hypotheses of [Theorem 11](#), if $\text{Aut}_{\text{orb}}(S/H)$ is nontrivial, then S/H admits an extra conformal involution J such that $(S/H)/\langle J \rangle$ is the orbifold whose underlying Riemann surface is \widehat{C} , with exactly three cone points (of orders $2p$, $2p^{n-1}$ and p^n). It follows that S is a Belyi curve and hence it may be defined over a finite extension of \mathbb{Q} .

Our next result computes the field of moduli for the homology covers of orbifolds with signature $(0; p, p^{n-1}, p^n, p^n)$, where $p \geq 3$ is a prime and $n \geq 2$.

Theorem 13. *Let $p \geq 3$ be a prime and $n \geq 2$ be an integer. For each $\lambda \in \mathbb{C} - \{0, 1\}$, let C_λ be as in [Theorem 9](#) with $k = p$. Then:*

- (1) $C_\lambda \cong C_\mu$ for $\lambda, \mu \in \mathbb{C} - \{0, 1\}$ if and only if $\mu \in \{\lambda, 1/\lambda\}$.
- (2) $\mathbb{M}(C_\lambda) = \mathbb{Q}(\lambda + \lambda^{-1})$.
- (3) $\mathbb{M}(C_\lambda)$ is a field of definition for C_λ .

[Theorem 13](#) will be proved using arguments similar to those given by Dèbes and Emsalem in the proof of [Theorem 3](#). In our case, we do not consider the quotient by the full group of automorphisms, but just the quotient by the abelian group H in [Theorem 9](#).

4. Proof of Theorem 4

Proof of part 4. As previously noted, there is a regular branched cover $P : S \rightarrow \widehat{\mathbb{C}}$, with H as deck group, so that its branch values are ∞ of order p , 0 of order p^{n-1} , 1 of order p^n and λ of order p^n . Let us denote by \mathbb{O}_λ the orbifold whose underlying Riemann surface is $\widehat{\mathbb{C}}$ and whose cone points are ∞ of order p , 0 of order p^{n-1} , 1 of order p^n and λ of order p^n ; that is, $\mathbb{O}_\lambda = S/H$.

If H is not a p -Sylow subgroup, then there is some $H \triangleleft K < \text{Aut}(S)$, where K is a p -group and $[K : H] = p$. It follows that there is an automorphism of order $p \geq 3$ of the orbifold \mathbb{O}_λ . As there are no three cone points with the same order, this is impossible. \square

Proof of parts (2) and (3). If $n \geq 3$, then it is easy to see that

$$\text{Aut}_{\text{orb}}(\mathbb{O}_\lambda) = \begin{cases} \{I\}, & \lambda \in \mathbb{C} - \{0, \pm 1\}, \\ \langle \tau(z) = -z \rangle, & \lambda = -1. \end{cases}$$

Since $\text{Aut}_H(S)/H < \text{Aut}_{\text{orb}}(\mathbb{O}_\lambda)$, it follows that

$$\text{Aut}_H(S) = \begin{cases} H, & \lambda \in \mathbb{C} - \{0, \pm 1\}, \\ K, & \lambda = -1, \end{cases}$$

where $[K : H] \in \{1, 2\}$.

If $n = 2$, then

$$\text{Aut}_{\text{orb}}(\mathbb{O}_\lambda) = \begin{cases} \langle \alpha(z) = \lambda/z \rangle, & \lambda \in \mathbb{C} - \{0, \pm 1\}, \\ \langle \tau(z) = -z, \beta(z) = -1/z \rangle, & \lambda = -1. \end{cases}$$

Again as $\text{Aut}_H(S)/H < \text{Aut}_{\text{orb}}(\mathbb{O}_\lambda)$, it follows that

$$\text{Aut}_H(S) = \begin{cases} \widehat{H}, & \lambda \in \mathbb{C} - \{0, \pm 1\}, \\ \widehat{K}, & \lambda = -1, \end{cases}$$

where $[\widehat{H} : H] \in \{1, 2\}$ and $[\widehat{K} : H] \in \{1, 2, 4\}$. \square

Proof of part (4). As a consequence of the results in [Leyton and Hidalgo 2007], there exists a prime p_0 such that the group H is a normal subgroup in $\text{Aut}(S)$ for $p \geq p_0$; that is, $\text{Aut}(S) = \text{Aut}_H(S)$. Next, we proceed to prove that p_0 may be chosen as desired.

Let $p \geq 3$ be any odd prime. We already know that H is a p -Sylow subgroup of $\text{Aut}(S)$ and that S/H has signature $(0; p, p^{n-1}, p^n, p^n)$. If $S/\text{Aut}(S)$ has signature of the form $(0; a, b, c, d)$, then it follows from Singerman's list [1972] of maximal Fuchsian groups that $(0; a, b, c, d) = (0; p, p^{n-1}, p^n, p^n)$ and, in particular, that $H = \text{Aut}(S)$.

Thus we need only take care of the case when $S/\text{Aut}(S)$ has signature of the form $(0; r, s, t)$. In this case, at least one of the values r, s, t should be a multiple of

p^n . We may assume $t = kp^n$, where k is a positive integer. We may also assume that $2 \leq r \leq s$ and, moreover, that if $r = 2$, then $s \geq 3$. Let

$$D = [\text{Aut}(S) : H].$$

If $D = 2$, then clearly $\text{Aut}_H(S) = \text{Aut}(S)$.

From now on assume that $D \geq 3$. Riemann–Hurwitz (hyperbolic area comparison) asserts that

$$(4-1) \quad D \left(1 - \frac{1}{r} - \frac{1}{s} - \frac{1}{kp^n} \right) = 2 - \frac{1}{p} - \frac{1}{p^{n-1}} - \frac{2}{p^n},$$

where both sides are necessarily positive.

Lemma 14. *If either*

- (1) $p \geq 7$, or
- (2) $p \in \{3, 5\}$ and $n \geq 3$,

then $D \leq 11$.

Proof. Assume $D \geq 12$. As $(r, s) \neq (2, 2)$, it follows from (4-1) that

$$D \left(\frac{1}{6} - \frac{1}{kp^n} \right) \leq 2 - \frac{1}{p} - \frac{1}{p^{n-1}} - \frac{2}{p^n}.$$

Since the quantity in parentheses is positive, the last inequality implies that

$$k \leq \frac{12}{2 + p + p^{n-1}}.$$

Therefore, if $p \geq 7$ then

$$k \leq \frac{12}{2 + p + p^{n-1}} \leq \frac{12}{2 + 2p} \leq \frac{3}{4} < 1,$$

and if $p \in \{3, 5\}$ and $n \geq 3$ then

$$k \leq \frac{12}{2 + p + p^{n-1}} \leq \frac{12}{2 + 3 + 3^2} \leq \frac{6}{7} < 1,$$

obtaining a contradiction in all cases. □

The following proposition gives the desired result.

Proposition 15.

- (1) *If $n \geq 2$, then $p_0 \leq 7$.*
- (2) *If $n \geq 3$, then $p_0 \leq 5$.*

Proof. Let us denote by N_p be the number of p -Sylow subgroups of $\text{Aut}(S)$. We need to prove that $N_p = 1$, if either (i) $p \geq 7$ is prime and $n \geq 2$ or (ii) $p \geq 5$ is a prime and $n \geq 3$.

As $N_p \equiv 1 \pmod{p}$, we may write $N_p = 1 + pL_p$, where L_p is a nonnegative integer.

If we assume that $N_p > 1$, then $N_p \geq 1 + p$. As N_p divides $|\text{Aut}(S)| = D|H|$, it follows that N_p must divide D .

If $p \geq 11$, then $N_p \geq 12$; as $D \leq 11$ by [Lemma 14](#), we obtain a contradiction.

For the remaining cases, we will make use of the following equality, obtained from (4-1):

$$(4-2) \quad \left(D \left(1 - \frac{1}{r} - \frac{1}{s} \right) - 2 \right) p^n + p^{n-1} + p + 2 = \frac{D}{k} \in \{1, \dots, D\}.$$

Note that both sides in this equality are positive integers.

If $p = 7$, since $D \leq 11$ by [Lemma 14](#), we must have that $L_7 = 1$ and $N_7 = D = 8$. If either $r, s \geq 3$ or $r = 2$ and $s \geq 4$, then

$$\left(8 \left(1 - \frac{1}{r} - \frac{1}{s} \right) - 2 \right) \geq 0$$

and the left side of (4-2) is bigger than 8, a contradiction to the fact that the right side should be less than or equal to D .

We are left with the case $r = 2$ and $s = 3$. But in this case the left side of (4-2) equals

$$\left(8 \left(1 - \frac{1}{r} - \frac{1}{s} \right) - 2 \right) 7^n + 7^{n-1} + 9 < 0,$$

again a contradiction.

Now we consider $p = 5$ and $n \geq 3$. In this case either (i) $L_5 = 1$ and $N_5 = D = 6$ or (ii) $L_5 = 2$ and $N_5 = D = 11$.

For $D = 6$, if either (a) $r, s \geq 3$ or (b) $r = 2$ and $s \geq 6$, then

$$\left(6 \left(1 - \frac{1}{r} - \frac{1}{s} \right) - 2 \right) \geq 0$$

and the left side of (4-2) is bigger than D , a contradiction. The remaining cases are $r = 2$ and $3 \leq s \leq 5$. But in these cases we have

$$\left(6 \left(1 - \frac{1}{r} - \frac{1}{s} \right) - 2 \right) 5^n + 5^{n-1} + 7 < 0,$$

again a contradiction.

For $D = 11$, if either (a) $r, s \geq 3$ or (b) $r = 2$ and $s \geq 4$, then

$$\left(11 \left(1 - \frac{1}{r} - \frac{1}{s} \right) - 2 \right) \geq 0$$

and the left side of (4-2) is bigger than D , a contradiction. The remaining cases are $r = 2$ and $s = 3, 4$. But in these cases we have

$$\left(11\left(1 - \frac{1}{r} - \frac{1}{s}\right) - 2\right)5^n + 5^{n-1} + 7 < 0,$$

again a contradiction. \square

5. Proof of Theorem 9

Let R be the homology cover of an orbifold \mathbb{O} with signature $(0; k, k^{n-1}, k^n, k^n)$, where $k, n \geq 2$. The closed Riemann surface R admits a group $H < \text{Aut}(R)$, where $H \cong \mathbb{Z}_k \times \mathbb{Z}_{k^{n-1}} \times \mathbb{Z}_{k^n}$ and such that $R/H = \mathbb{O}$.

First consider the orbifold \mathbb{O}^* obtained from \mathbb{O} , but assuming all cone points are of order k^n . The homology cover of this new orbifold is a closed Riemann surface S admitting a group $H^* < \text{Aut}(S)$, $H^* \cong \mathbb{Z}_{k^n} \times \mathbb{Z}_{k^n} \times \mathbb{Z}_{k^n}$, and such that $\mathbb{O}^* = S/H^*$. It is known (see [González-Diez et al. 2009]) that an algebraic curve representation of S is given by

$$\widehat{C} : \left\{ \begin{array}{l} x_0^{k^n} + x_1^{k^n} + x_2^{k^n} = 0 \\ \lambda x_0^{k^n} + x_1^{k^n} + x_3^{k^n} = 0 \end{array} \right\} \subset \mathbb{P}^3,$$

that H^* is generated by the projective transformations

$$\begin{aligned} a([x_0 : x_1 : x_2 : x_3]) &= [\rho_n x_0 : x_1 : x_2 : x_3], & b([x_0 : x_1 : x_2 : x_3]) &= [x_0 : \rho_n x_1 : x_2 : x_3], \\ c([x_0 : x_1 : x_2 : x_3]) &= [x_0 : x_1 : \rho_n x_2 : x_3] \end{aligned}$$

and that the holomorphic map

$$\pi : \widehat{C} \rightarrow \widehat{\mathbb{C}} : [x_0 : x_1 : x_2 : x_3] \mapsto -\left(\frac{x_1}{x_0}\right)^{k^n}$$

has degree k^{3n} and is a branched regular cover with H^* as deck group. In this case, $\pi(\text{Fix}(a)) = \infty$, $\pi(\text{Fix}(b)) = 0$, $\pi(\text{Fix}(c)) = 1$ and $\pi(\text{Fix}(abc)) = \lambda$.

Now consider the subgroup of H^* given by $K = \langle a^k, b^{k^{n-1}} \rangle \cong \mathbb{Z}_{k^{n-1}} \times \mathbb{Z}_k$, and set $\mathbb{O}_0 = S/K$. The group $H_0 = H^*/K$ is a group of conformal automorphisms of \mathbb{O}_0 , $H_0 \cong H$, and $\mathbb{O}_0/H_0 = \mathbb{O}^*$.

Clearly, if R_0 denotes the underlying Riemann surface structure of the orbifold \mathbb{O}_0 , then R_0/H_0 is the orbifold \mathbb{O} . In this way, since any two homology covers of \mathbb{O} are conformally equivalent, we may assume $R = R_0$.

In order to find an algebraic curve representation for R_0 we proceed as follows. First, we consider the affine curve representation of S defined by $x = x_0/x_3$,

$y = x_1/x_3$ and $z = x_2/x_3$; that is,

$$\widehat{C}_0 = \left\{ \begin{array}{l} x^{k^n} + y^{k^n} + z^{k^n} = 0 \\ \lambda x^{k^n} + y^{k^n} + 1 = 0 \end{array} \right\} \subset \mathbb{C}^3$$

and the action of H^* is generated by the linear transformations

$$a(x, y, z) = (\rho_n x, y, z), \quad b(x, y, z) = (x, \rho_n y, z), \quad c(x, y, z) = (x, y, \rho_n z).$$

The subalgebra of $\langle a^k, b^{k^{n-1}} \rangle$ invariant polynomials, $\mathbb{C}[x, y, z]^{\langle a^k, b^{k^{n-1}} \rangle}$, is generated by the monomials $x^{k^{n-1}}$, y^k and z . It follows that the holomorphic map

$$F : \mathbb{C}^3 \rightarrow \mathbb{C}^3, \\ (x, y, z) \mapsto (x^{k^{n-1}}, y^k, z) = (u, v, w)$$

is a regular branched covering with $\langle a^k, b^{k^{n-1}} \rangle$ as deck group, and therefore $F(\widehat{C}_0)$ provides an affine algebraic curve representation of R , given by

$$F(\widehat{C}_0) = \left\{ \begin{array}{l} u^k + v^{k^{n-1}} + w^{k^n} = 0 \\ \lambda u^k + v^{k^{n-1}} + 1 = 0 \end{array} \right\} \subset \mathbb{C}^3,$$

where the action of $H = H^*/K$ is generated by

$$a_0(u, v, w) = (\rho_1 u, v, w), \quad b_0(u, v, w) = (u, \rho_{n-1} v, w), \\ c_0(u, v, w) = (u, v, \rho_n w).$$

If we consider the projective space \mathbb{P}^3 with coordinates $[z_0 : z_1 : z_2 : z_3]$, and we set

$$u = \frac{z_0}{z_3}, \quad v = \frac{z_1}{z_3}, \quad w = \frac{z_2}{z_3},$$

then we obtain that R is represented by the projective algebraic curve

$$C = \left\{ \begin{array}{l} z_0^k z_3^{k^n-k} + z_1^{k^{n-1}} z_3^{k^n-k^{n-1}} + z_2^{k^n} = 0 \\ \lambda z_0^k z_3^{k^{n-1}-k} + z_1^{k^{n-1}} + z_3^{k^{n-1}} = 0 \end{array} \right\} \subset \mathbb{P}^3.$$

As the branched covering map $P : R \rightarrow R/H$ must satisfy $\pi = P \circ F$ and

$$F([x_0 : x_1 : x_2 : x_3]) = [x_0^{k^{n-1}} : x_1^k x_3^{k^{n-1}-k} : x_2 x_3^{k^{n-1}-1} : x_3^{k^{n-1}}],$$

then

$$P([z_0 : z_1 : z_2 : z_3]) = - \left(\frac{z_1^{k^{n-1}}}{z_0^k z_3^{k^{n-1}-k}} \right).$$

6. Proof of Theorem 10

Proof. Consider a closed Riemann surface S admitting a group $G < \text{Aut}(S)$ such that $G \cong \mathbb{Z}_p \times \mathbb{Z}_{p^n}$ and $\mathbb{O} = S/G$ is an orbifold with signature $(0; p, p^{n-1}, p^n, p^n)$, where $n \geq 2$ and p is an odd prime. Denote by $P : S \rightarrow \mathbb{O}$ the natural holomorphic branched cover with G as deck group.

In this section we will find algebraic curves representing S and the action of G on them.

Let R be the homology cover of \mathbb{O} , and let $Q : R \rightarrow \mathbb{O} = R/H$ be the branched regular covering with H as deck group, where $H = \mathbb{Z}_p \times \mathbb{Z}_{p^{n-1}} \times \mathbb{Z}_{p^n}$.

Since G is abelian, there is a subgroup $K < H$ such that $S = R/K$ (and hence K acts freely on R), $G = H/K$, and there is a regular holomorphic covering $T : R \rightarrow S$ with K as deck group and $Q = P \circ T$.

Consider the affine algebraic curve C_0 representing R , obtained from Theorem 9 by making $z_3 = 1$:

$$C_0 = \left\{ \begin{array}{l} z_0^p + z_1^{p^{n-1}} + z_2^{p^n} = 0 \\ \lambda z_0^p + z_1^{p^{n-1}} + 1 = 0 \end{array} \right\} \subset \mathbb{C}^3,$$

in which case the group H is generated by

$$\begin{aligned} a_0(z_0, z_1, z_2) &= (\rho_1 z_0, z_1, z_2), & b_0(z_0, z_1, z_2) &= (z_0, \rho_{n-1} z_1, z_2), \\ c_0(z_0, z_1, z_2) &= (z_0, z_1, \rho_n z_2). \end{aligned}$$

6.1. Algebraic structure of K . We next describe the algebraic structure of K . At this point we should note that, using the model of R given in Theorem 9, the transformations in H acting with fixed points on S are exactly the ones that belong to $\langle a_0 \rangle \cup \langle b_0 \rangle \cup \langle c_0 \rangle \cup \langle a_0 b_0 c_0 \rangle$.

Proposition 16. *Consider the algebraic model of (R, H) provided by Theorem 9. Let $K < H$ be such that K acts freely on R and $H/K \cong \mathbb{Z}_p \times \mathbb{Z}_{p^n}$. Then, either*

- (1) $\mathbb{Z}_{p^{n-1}} \cong K = \langle a_0^\alpha b_0 c_0^{p^q} \rangle$, where $(p, q) = 1$ and $0 \leq \alpha \leq p-1$; or
- (2) $\mathbb{Z}_{p^{n-2}} \times \mathbb{Z}_p \cong K = \langle b_0^{-p} c_0^{p^2 v} \rangle \times \langle a_0 c_0^{p^{n-1} \gamma} \rangle$, where $(p, v) = 1$ and $1 \leq \gamma \leq p-1$.

Proof. Consider a surjective homomorphism

$$\Phi : H \rightarrow J = \mathbb{Z}_p \times \mathbb{Z}_{p^n}$$

with $K = \ker(\Phi)$ acting freely on R . Note that the order of K is p^{n-1} . Then:

- a) $K \cap \langle a_0 \rangle = \{I\}$, which implies that $\Phi(a_0)$ has order p .
- b) $K \cap \langle b_0 \rangle = \{I\}$, which implies that $\Phi(b_0)$ has order p^{n-1} .

c) $K \cap \langle c_0 \rangle = \{I\}$, which implies that $\Phi(c_0)$ has order p^n .

d) $K \cap \langle a_0 b_0 c_0 \rangle = \{I\}$, which implies that $\Phi(a_0)\Phi(b_0)\Phi(c_0)$ has order p^n .

Hence the subgroups of J given by $\langle \Phi(b_0) \rangle$ and $\langle \Phi(c_0) \rangle$ have respective indices p^2 and p , and there are two cases to be considered, as follows.

Case i). Assume $\langle \Phi(b_0) \rangle \subset \langle \Phi(c_0) \rangle$. Then there exists $1 \leq u \leq p-1$ such that $\Phi(b_0) = \Phi(c_0^{p^u})$, in which case $h = b_0 c_0^{-p^u}$ is an element of K of order p^{n-1} , and therefore $K = \langle h \rangle$ is cyclic of the form given in case (1).

Case ii). Assume $\langle \Phi(b_0) \rangle \not\subset \langle \Phi(c_0) \rangle$. Then we have the following commutative diagram of subgroup inclusions and corresponding indices:

$$\begin{array}{ccc}
 & J & \\
 p \nearrow & & \nwarrow \\
 \langle \Phi(c_0) \rangle & & \langle \Phi(b_0) \rangle_{p^2} \\
 p^2 \nwarrow & & \nearrow \\
 & \langle \Phi(c_0) \rangle \cap \langle \Phi(b_0) \rangle_p &
 \end{array}$$

and it follows that

$$\langle \Phi(c_0) \rangle \cap \langle \Phi(b_0) \rangle = \langle \Phi(c_0^{p^2}) \rangle = \langle \Phi(b_0^p) \rangle.$$

Hence there exists v such that $h_0 = c_0^{p^2 v} b_0^{-p}$ is in K , and h_0 has order p^{n-2} . Also note that $(v, p) = 1$, since otherwise an adequate power of h_0 would be a nontrivial power of b_0 in K . It follows that there are two possibilities for K , either $K \cong \mathbb{Z}_{p^{n-1}}$ or $K = \langle h_0 \rangle \times \langle t \rangle \cong \mathbb{Z}_{p^{n-2}} \times \mathbb{Z}_p$.

Subcase K is not cyclic. As previously noted, in this case $K = \langle h_0 \rangle \times \langle t \rangle \cong \mathbb{Z}_{p^{n-2}} \times \mathbb{Z}_p$, where $h_0 = c_0^{p^2 v} b_0^{-p}$ and $(p, v) = 1$. As $t \in H$ has order p , it has the form

$$t = a_0^\alpha b_0^\beta c_0^{\gamma p^{n-2}},$$

where $\alpha, \beta, \gamma \in \{0, 1, \dots, p-1\}$.

Let us assume $\alpha = 0$. If $\gamma = 0$, then $t \in \langle b_0 \rangle$. As K acts freely on R , necessarily $t = 1$ and we get a contradiction. If $(\gamma, p) = 1$, then we may assume $t = b_0^\beta c_0^{\gamma p^{n-2}}$ (by considering an appropriate power of the original t); hence

$$\tilde{h} = t h_0^{-p^{n-3}} = b_0^{(\beta+v)p^{n-2}} \in K \cap \langle b_0 \rangle.$$

Again, as K acts freely, \tilde{h} must be trivial, and t would belong to $\langle h_0 \rangle$, again a contradiction. Then we have proved that $\alpha > 0$.

Since t has order p , we may replace t by a suitable power of it in order to assume that $t = a_0 b_0^{\beta p^{n-2}} c_0^{\gamma p^{n-1}}$.

We now claim that we may assume $\beta = 0$. Indeed, if $\beta > 0$, then $t h_0^{\beta p^{n-3}} = a_0 c_0^{p^{n-1}(\gamma+\beta)}$ is an element of order p in K that does not belong to $\langle h_0 \rangle$.

Therefore we may write $t = a_0 c_0^{p^{n-1}\gamma}$, and observe that $1 \leq \gamma \leq p-1$ because $K \cap \langle a_0 \rangle = \{I\}$. This is case (2).

Subcase K is cyclic. In this case, $K = \langle h \rangle \cong \mathbb{Z}_{p^{n-1}}$. Let us write

$$h = a_0^\alpha b_0^\beta c_0^\gamma$$

where $\alpha \in \{0, 1, \dots, p-1\}$, $\beta \in \{0, 1, \dots, p^{n-1}-1\}$, $\gamma \in \{0, 1, \dots, p^n-1\}$.

The condition $c_0^{\gamma p^{n-1}} = h^{p^{n-1}} = 1$ ensures that $\gamma \equiv 0 \pmod{p}$. It follows that either $\gamma = 0$ or $\gamma = p^s q$, where $s \in \{1, \dots, n-1\}$ and $(p, q) = 1$.

Next, we need to ensure that, for $\delta \in \{1, 2, \dots, p^{n-1}-1\}$, no power h^δ acts with fixed points in C ; that is, $h^\delta \notin \langle a_0 \rangle \cup \langle b_0 \rangle \cup \langle c_0 \rangle \cup \langle a_0 b_0 c_0 \rangle$.

But if $\gamma = 0$ then $h^p = b_0^{p\beta}$ is a nontrivial element of the group generated by b_0 , a contradiction. Similarly, if $s > 1$ then $h^{p^{n-s}} = b_0^{\beta p^{n-s}}$ is a nontrivial element of the group generated by b_0 , a contradiction.

Therefore $h = a_0^\alpha b_0^\beta c_0^{pq}$, with $(p, q) = 1$, and it follows that h^δ is not in $\langle b_0 \rangle$.

But if $\beta \equiv 0 \pmod{p}$, then $h^{p^{n-2}} = c_0^{q p^{n-2}}$ is a nontrivial element of the group generated by c_0 , a contradiction. Hence $(p, \beta) = 1$, and h^δ is not in $\langle c_0 \rangle$.

We note that $h^\delta \in \langle a_0 \rangle$ implies that $\beta \delta \equiv 0 \pmod{p^{n-1}}$, and since $(\beta, p) = 1$, to have $\delta \equiv 0 \pmod{p^{n-1}}$, which is not possible by our choice for δ .

The condition $h^\delta \in \langle a_0 b_0 c_0 \rangle$ implies that $\beta \delta \equiv pq\delta \pmod{p^{n-1}}$, from which $(\beta - pq)\delta \equiv 0 \pmod{p^{n-1}}$, and then $\delta \equiv 0 \pmod{p^{n-1}}$, which is not possible by our choice for δ .

By taking an appropriate power of h , we may assume that

$$K = \langle a_0^\alpha b_0 c_0^{pq} \rangle,$$

where $(p, q) = 1$.

Now note that in this case $1 \leq \alpha \leq p-1$, since $\alpha = 0$ implies that $\Phi(b_0) = \Phi(c_0)^{-pq}$ is an element of $\langle \Phi(c_0) \rangle$, which is a contradiction, as we are in case ii). This is case (1). \square

6.2. The cyclic case. As a consequence of [Proposition 16](#), we may assume

$$K = \langle a_0^\alpha b_0 c_0^{pq} \rangle,$$

where $(p, q) = 1$ and $\alpha \in \{0, 1, \dots, p-1\}$. Note that

$$a_0^\alpha b_0 c_0^{pq}(z_0, z_1, z_2) = (\rho_1^\alpha z_0, \rho_{n-1} z_1, \rho_{n-1}^q z_2).$$

6.2.1. The case $\alpha = 0$. We next search for polynomials in $\mathbb{C}[z_0, z_1, z_2]^K$. We first note that $z_0 \in \mathbb{C}[z_0, z_1, z_2]^K$. Next, we search for polynomials of the form $z_1^u z_2^v \in \mathbb{C}[z_0, z_1, z_2]^K$, where $u, v \in \{0, 1, \dots, p^{n-1}\}$. The invariance property requires that the values u and v satisfy the relation

$$u + vq \equiv 0 \pmod{p^{n-1}}.$$

As $(p, q) = 1$, we have that some of those polynomials are given by

$$z_1^{p^{n-1}}, \quad z_2^{p^{n-1}}, \quad z_1^q z_2^{p^{n-1}-1}.$$

Let us consider the holomorphic map

$$F : \mathbb{C}^3 \rightarrow \mathbb{C}^4, \\ F(z_0, z_1, z_2) = (z_0, z_1^{p^{n-1}}, z_2^{p^{n-1}}, z_1^q z_2^{p^{n-1}-1}) = (x_1, x_2, x_3, x_4).$$

Let us note that $x_4/x_3 = z_1^q/z_2$. As $(p^{n-1}, q) = 1$, it follows that there exist integers a, b so that $aq + bp^{n-1} = 1$; that is, $z_1 = (z_1^q)^a (z_1^{p^{n-1}})^b = (x_4/x_3)^a x_2^b$. It follows that z_1 is uniquely determined by the tuple (x_1, x_2, x_3, x_4) and a choice for z_2 . In particular, as z_0 is uniquely determined by x_1 , one sees that the map F has degree p^{n-1} and it is K -invariant. In this way, an affine algebraic curve defining $F(C_0)$ is given by

$$F(C_0) = \left\{ \begin{array}{l} x_1^p + x_2 + x_3^p = 0 \\ \lambda x_1^p + x_2 + 1 = 0 \\ x_4^{p^{n-1}} - x_2^q x_3^{p^{n-1}-1} = 0 \end{array} \right\} \subset \mathbb{C}^4$$

and a projective one is provided by taking $x_1 = y_0/y_4$, $x_2 = y_1/y_4$, $x_3 = y_2/y_4$, $x_4 = y_3/y_4$, where $[y_0 : y_1 : y_2 : y_3, y_4] \in \mathbb{P}^4$, as follows:

$$\left\{ \begin{array}{l} y_0^p + y_1 y_4^{p-1} + y_2^p = 0 \\ \lambda y_0^p + y_1 y_4^{p-1} + y_4^p = 0 \\ y_3^{p^{n-1}} - y_1^q y_2^{p^{n-1}-1} y_4^{1-q} = 0 \end{array} \right\} \subset \mathbb{P}^4.$$

The map F is, in projective coordinates, given as

$$F([z_0 : z_1 : z_2 : z_3]) = [z_0 z_3^{p^{n-1}-1} : z_1^{p^{n-1}} : z_2^{p^{n-1}} : z_1^q z_2^{p^{n-1}-1} z_3^{1-q} : z_3^{p^{n-1}}] \\ = [y_0 : y_1 : y_2 : y_3 : y_4].$$

As, by the first equality above,

$$y_1 = -\left(\frac{y_0^p + y_2^p}{y_4^{p-1}} \right),$$

the above also provides the (birational) algebraic curve

$$\left\{ \begin{array}{l} (\lambda - 1)y_0^p - y_2^p + y_4^p = 0 \\ (-1)^{q+1}(y_0^p + y_2^p)^q y_2^{p^{n-1}-1} + y_3^{p^{n-1}} y_4^{qp-1} = 0 \end{array} \right\} \subset \mathbb{P}^3.$$

By making the change of coordinates $w_0 = y_0$, $w_1 = y_2$, $w_2 = y_3$, $w_3 = y_4$, the above is written as

$$\left\{ \begin{array}{l} (\lambda - 1)w_0^p - w_1^p + w_3^p = 0 \\ (-1)^{q+1}(w_0^p + w_1^p)^q w_1^{p^{n-1}-1} + w_2^{p^{n-1}} w_3^{qp-1} = 0 \end{array} \right\} \subset \mathbb{P}^3$$

and the map F is given as

$$F([z_0 : z_1 : z_2 : z_3]) = [z_0 z_3^{p^{n-1}-1} : z_2^{p^{n-1}} : z_1^q z_2^{p^{n-1}-1} z_3^{1-q} : z_3^{p^{n-1}}] = [w_0 : w_1 : w_2 : w_3].$$

In this case, the group $G = H/K$ is generated by the transformations

$$A_1([w_0 : w_1 : w_2 : w_3]) = [\rho_1 w_0 : w_1 : w_2 : w_3],$$

$$B_1([w_0 : w_1 : w_2 : w_3]) = [w_0 : w_1 : \rho_{n-1}^q w_2 : w_3],$$

$$C_1([w_0 : w_1 : w_2 : w_3]) = [w_0 : \rho_1 w_1 : \rho_n^{p^{n-1}-1} w_2 : w_3].$$

Notice that the elements $A = A_1$ and $B = C_1$ also generate G as desired. As the branched covering map $Q : S \rightarrow S/G$ must satisfy $P = Q \circ F$, where $P : R \rightarrow R/H$ is (as in [Theorem 9](#)) given by

$$P([z_0 : z_1 : z_2 : z_3]) = -\left(\frac{z_1^{p^{n-1}}}{z_0^p z_3^{p^{n-1}-p}} \right),$$

and since

$$-\left(\frac{z_1^{p^{n-1}}}{z_0^p z_3^{p^{n-1}-p}} \right) = -\left(\frac{y_1 y_4^{p-1}}{y_0^p} \right) = \frac{y_0^p + y_2^p}{y_0^p} = \frac{w_0^p + w_1^p}{w_0^p},$$

we obtain

$$Q([w_0 : w_1 : w_2 : w_3]) = \frac{w_0^p + w_1^p}{w_0^p}.$$

6.2.2. *The case $\alpha \in \{1, 2, \dots, p-1\}$.* Next, we search for polynomials of the form $z_0^t z_1^u z_2^v \in \mathbb{C}[z_0, z_1, z_2]^K$, where $t \in \{0, 1, \dots, p-1\}$ and $u, v \in \{0, 1, \dots, p^{n-1}\}$. The invariance property requires that the values u and v satisfy the relation

$$t\alpha p^{n-2} + u + vq \equiv 0 \pmod{p^{n-1}}.$$

As $(p, q) = (\alpha, p) = 1$, we have that some of those polynomials are given by

$$z_0^p, \quad z_1^{p^{n-1}}, \quad z_2^{p^{n-1}}, \quad z_1^q z_2^{p^{n-1}-1}, \quad z_0^{p-1} z_1^\alpha p^{n-2}.$$

Let us consider the holomorphic map

$$F: \mathbb{C}^3 \rightarrow \mathbb{C}^5,$$

$$F(z_0, z_1, z_2) = (z_0^p, z_1^{p^{n-1}}, z_2^{p^{n-1}}, z_1^q z_2^{p^{n-1}-1}, z_0^{p-1} z_1^{\alpha p^{n-2}}) = (x_1, x_2, x_3, x_4, x_5).$$

Let us note that $x_4/x_3 = z_1^q/z_2$. As $(p^{n-1}, q) = 1$, it follows that there exist integers a, b so that $aq + bp^{n-1} = 1$, from where $z_1 = (z_1^q)^a (z_1^{p^{n-1}})^b = (x_4/x_3)^a x_2^b z_2$. It follows that z_1 is uniquely determined by the tuple $(x_1, x_2, x_3, x_4, x_5)$ and a choice for z_2 .

As z_0^p is uniquely determined by x_1 , and $z_0^{p-1} z_1^{\alpha p^{n-2}}$ is uniquely determined by x_2, x_3, x_4, x_5 and a choice of z_2 , we have that z_0 is also uniquely determined by the previous data.

All the above permits us to see that the map F has degree p^{n-1} and it is K -invariant. In this way, an affine algebraic curve defining $F(C_0)$ is given by

$$F(C_0) = \left\{ \begin{array}{l} x_1 + x_2 + x_3^p = 0 \\ \lambda x_1 + x_2 + 1 = 0 \\ x_4^{p^{n-1}} - x_2^q x_3^{p^{n-1}-1} = 0 \\ x_5^p - x_1^{p-1} x_2^\alpha = 0 \end{array} \right\} \subset \mathbb{C}^5.$$

We may write $x_2 = -(x_1 + x_3^p)$. In this way, writing $u_1 = x_1, u_2 = x_3, u_3 = x_4$ and $u_4 = x_5$, the above curve is

$$\left\{ \begin{array}{l} (\lambda - 1)u_1 - u_2^p + 1 = 0 \\ u_3^{p^{n-1}} + (-1)^{q+1} (u_1 + u_2^p)^q u_2^{p^{n-1}-1} = 0 \\ u_4^p + (-1)^{\alpha+1} u_1^{p-1} (u_1 + u_2^p)^\alpha = 0 \end{array} \right\} \subset \mathbb{C}^4.$$

Now, we may write

$$u_1 = \frac{1}{\lambda - 1} (u_2^p - 1),$$

and setting $y_1 = u_2, y_2 = u_3$ and $y_3 = u_4$, the above curve is

$$\left\{ \begin{array}{l} y_2^{p^{n-1}} + \frac{(-1)^{q+1}}{(\lambda - 1)^q} (\lambda y_1^p - 1)^q y_1^{p^{n-1}-1} = 0 \\ y_3^p + \frac{(-1)^{\alpha+1}}{(\lambda - 1)^{\alpha+p-1}} (y_1^p - 1)^{p-1} (\lambda y_1^p - 1)^\alpha = 0 \end{array} \right\} \subset \mathbb{C}^3$$

and F is of the form

$$F(z_0, z_1, z_2) = (z_2^{p^{n-1}}, z_1^q z_2^{p^{n-1}-1}, z_0^{p-1} z_1^{\alpha p^{n-2}}) = (y_1, y_2, y_3).$$

Writing $y_1 = v_0/v_3$, $y_2 = v_1/v_3$ and $y_3 = v_2/v_3$, we obtain the projective model

$$\left\{ \begin{array}{l} v_1^{p^{n-1}} v_3^{pq-1} + \frac{(-1)^{q+1}}{(\lambda-1)^q} (\lambda v_0^p - v_3^p)^q v_0^{p^{n-1}-1} = 0 \\ v_2^p v_3^{p^2+p(\alpha-2)} + \frac{(-1)^{\alpha+1}}{(\lambda-1)^{\alpha+p-1}} (v_0^p - v_3^p)^{p-1} (\lambda v_0^p - v_3^p)^\alpha = 0 \end{array} \right\} \subset \mathbb{P}^3$$

and for $n \geq 3$ we have that $\max\{p^{n-1}, p^{n-1}+q-1, \alpha p^{n-2}+p-1\} = p^{n-1}+q-1$ and therefore $F: \mathbb{P}^3 \rightarrow \mathbb{P}^3$ is given as

$$F([z_0 : z_1 : z_2 : z_3]) = [z_2^{p^{n-1}} z_3^{q-1} : z_1^q z_2^{p^{n-1}-1} : z_0^{p-1} z_1^\alpha z_2^{p^{n-2}} z_3^{p^{n-1}+q-p-\alpha p^{n-2}} : z_3^{p^{n-1}+q-1}].$$

In the case $n = 2$ a similar formula may be given for F ; the maximum value above is $p+q-1$ if $q \geq \alpha$ and $p+\alpha-1$ otherwise.

Continuing with $n \geq 3$, the group $G = H/K$ is generated by the transformations

$$\begin{aligned} A_2([v_0 : v_1 : v_2 : v_3]) &= [v_0 : v_1 : \rho_1^{p-1} v_2 : v_3], \\ B_2([v_0 : v_1 : v_2 : v_3]) &= [v_0 : \rho_{n-1}^q v_1 : \rho_1^\alpha v_2 : v_3], \\ C_2([v_0 : v_1 : v_2 : v_3]) &= [\rho_1 v_0 : \rho_n^{p^{n-1}-1} v_1 : v_2 : v_3]. \end{aligned}$$

Notice that the elements $A = A_2$ and $B = C_2$ also generate G as desired. As the branched covering map $Q: S \rightarrow S/G$ must satisfy $P = Q \circ F$, where $P: R \rightarrow R/H$ is (as in [Theorem 9](#)) given by

$$\begin{aligned} P([z_0 : z_1 : z_2 : z_3]) &= -\left(\frac{z_1^{p^{n-1}}}{z_0^p z_3^{p^{n-1}-p}}\right) = -\left(\frac{x_2}{x_1}\right) = \frac{u_1 + u_2^p}{u_1} \\ &= 1 + \frac{(\lambda-1)u_2^p}{(u_2^p-1)} = 1 + \frac{(\lambda-1)y_1^p}{(y_1^p-1)} = 1 + \frac{(\lambda-1)v_0^p}{v_0^p - v_3^p}, \end{aligned}$$

we obtain

$$Q([v_0 : v_1 : v_2 : v_3]) = \frac{\lambda v_0^p - v_3^p}{v_0^p + v_3^p}.$$

6.3. The noncyclic case. In this case,

$$K = \langle b_0^{-p} c_0^{p^2 v}, a_0 c_0^\gamma p^{n-1} \rangle,$$

where $(p, v) = 1$ and $\gamma \in \{1, 2, \dots, p-1\}$.

We have that

$$\begin{aligned} b_0^{-p} c_0^{p^2 v} (z_0, z_1, z_2) &= (z_0, \rho_{n-2}^{-1} z_1, \rho_{n-2}^v z_2), \\ a_0 c_0^\gamma p^{n-1} (z_0, z_1, z_2) &= (\rho_1 z_0, z_1, \rho_1^\gamma z_2). \end{aligned}$$

Clearly, $z_0^A z_1^B z_2^C \in \mathbb{C}[z_0, z_1, z_2]^K$ if and only if

$$\begin{cases} A + C\gamma \equiv 0 \pmod{p}, \\ Cv - B \equiv 0 \pmod{p^{n-2}}. \end{cases}$$

In this way,

$$z_0^p, z_1^{p^{n-2}}, z_0^{p-\gamma} z_1^v z_2 \in \mathbb{C}[z_0, z_1, z_2]^K.$$

Let us consider the map

$$\begin{aligned} F : \mathbb{C}^3 &\rightarrow \mathbb{C}^3, \\ F(z_0, z_1, z_2) &= (z_0^p, z_1^{p^{n-2}}, z_0^{p-\gamma} z_1^v z_2) = (x_1, x_2, x_3). \end{aligned}$$

If we fix (x_1, x_2, x_3) , then we have p choices for z_0 ($z_0^p = x_1$) and p^{n-2} choices for z_1 ($z_1^{p^{n-2}} = x_2$). Once we have made such choices, the value of z_2 is uniquely determined from $z_0^{p-\gamma} z_1^v z_2 = x_3$. It follows that F has degree p^{n-1} and is K -invariant as desired.

The algebraic curve $F(C_0)$ is provided by

$$F(C_0) = \left\{ \begin{array}{l} x_1^{p^{n-1}(p-\gamma)} x_2^{p^2 v} (x_1 + x_2^p) + x_3^{p^n} = 0 \\ \lambda x_1 + x_2^p + 1 = 0 \end{array} \right\} \subset \mathbb{C}^3.$$

As

$$x_1 = -\frac{(1 + x_2^p)}{\lambda},$$

this curve is also represented by, taking $y_1 = x_2$ and $y_2 = x_3$,

$$\left\{ \frac{(-1)^{p^{n-1}(p-\gamma)}}{\lambda^{p^{n-1}(p-\gamma)}} (1 + y_1^p)^{p^{n-1}(p-\gamma)} y_1^{p^2 v} \left(y_1^p - \frac{(1 + y_1^p)}{\lambda} \right) + y_2^{p^n} = 0 \right\} \subset \mathbb{C}^2.$$

A projectivization of this plane curve is given by, using the projective coordinates $[u_0 : u_1 : u_2] \in \mathbb{P}^2$ and taking $y_1 = u_0/u_2$ and $y_2 = u_1/u_2$, the following one:

$$\begin{aligned} \left\{ \frac{(-1)^{p^{n-1}(p-\gamma)}}{\lambda^{p^{n-1}(p-\gamma)+1}} (u_0^p + u_2^p)^{p^{n-1}(p-\gamma)} u_1^{p^2 v} ((\lambda - 1)u_0^p - u_2^p) \right. \\ \left. + u_1^{p^n} u_2^{p^n(p-\gamma-1)+p+p^2 v} = 0 \right\} \subset \mathbb{P}^2. \end{aligned}$$

In this case, the transformations a_0 , b_0 and c_0 define the transformations

$$A_3([u_0 : u_1 : u_2]) = [u_0 : \rho_1^{p-\gamma} u_1 : u_2], \quad B_3([u_0 : u_1 : u_2]) = [\rho_1 u_0 : \rho_{n-1}^v u_1 : u_2], \\ C_3([u_0 : u_1 : u_2]) = [u_0 : \rho_n u_1 : u_2].$$

The elements $A = C_3^{-vp} B_3$ and $B = C_3$ also generate G as desired. And since

$$P(z_0, z_1, z_2) = -\left(\frac{z_1^{p^{n-1}}}{z_0^p}\right) = \frac{\lambda y_1^p}{1 + y_1^p},$$

we obtain

$$Q([u_0 : u_1 : u_2]) = \frac{\lambda u_0^p}{u_0^p + u_1^p}. \quad \square$$

7. Proof of Theorem 11

Proof. Let C be a nonsingular projective algebraic curve admitting a p -group H of conformal automorphisms of C with C/H of signature $(0; p, p^{n-1}, p^n, p^n)$ and let $P : C \rightarrow C/H = \widehat{\mathbb{C}}$ be a holomorphic branched covering with H as deck group. We may assume the branch values of P are given by ∞ or order p , 0 of order p^{n-1} , and 1 and $\lambda \in \mathbb{C} - \{0, 1\}$ are the ones of order p^n . We notice that

$$\text{Aut}_{\text{orb}}(S/H) = \begin{cases} \{I\}, & \lambda \neq -1, \\ \{J(z) = -z\}, & \lambda = -1. \end{cases}$$

Let $K_C = \{\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}) : C^\sigma \cong C\}$. For each $\sigma \in K_C$ there is a biholomorphism $f_\sigma : C \rightarrow C^\sigma$. As H^σ is unique up to conjugation in $\text{Aut}(C^\sigma)$, by Theorem 4, we may assume that $f_\sigma H f_\sigma^{-1} = H^\sigma$. It follows that there is a Möbius transformation M_σ so that $P^\sigma \circ f_\sigma = M_\sigma \circ P$. The transformation M_σ is uniquely determined by f_σ . As M_σ must preserve the cone points and their orders, it follows that $M_\sigma(\infty) = \infty$, $M_\sigma(0) = 0$ and that $\{1, \lambda_\sigma\} = \{M_\sigma(1), M_\sigma(\lambda)\}$, where $\lambda_\sigma \in \mathbb{C} - \{0, 1\}$ is branch value of order p^n of $P^\sigma : C^\sigma \rightarrow \widehat{\mathbb{C}}$ (in fact, $\lambda_\sigma = \sigma(\lambda)$). It follows that either (i) $M_\sigma = I$, in which case $\lambda_\sigma = \lambda$ or (ii) $M_\sigma(z) = z/\lambda$, in which case $\lambda_\sigma = 1/\lambda$.

7.1. Let us assume, from now on, that $\lambda \neq -1$.

Lemma 17. *Let $\lambda \neq -1$ and $\sigma \in K_C$. If there is another biholomorphism $\widehat{f}_\sigma : C \rightarrow C^\sigma$ such that $\widehat{f}_\sigma H \widehat{f}_\sigma^{-1} = H^\sigma$, then $\widehat{f}_\sigma = h \circ f_\sigma$, for some $h \in H$.*

Proof. If there is another biholomorphism $\widehat{f}_\sigma : C \rightarrow C^\sigma$ such that $\widehat{f}_\sigma H \widehat{f}_\sigma^{-1} = H^\sigma$, then $f_\sigma^{-1} \circ \widehat{f}_\sigma \in \text{Aut}(C)$ normalizes H . In this way, $f_\sigma^{-1} \circ \widehat{f}_\sigma$ induces an element of $\text{Aut}_{\text{orb}}(S/H)$. As this last group is trivial, we obtain that $f_\sigma^{-1} \circ \widehat{f}_\sigma \in H$. \square

As a consequence of [Lemma 17](#), M_σ is uniquely determined by σ and, in particular, the collection $\{M_\sigma : \sigma \in K_C\}$ satisfies Weil's conditions in [Theorem 2](#). Hence, there is an isomorphism $F : \widehat{\mathbb{C}} \rightarrow C_1$, where C_1 is defined over $\mathbb{M}(C)$, with the property that $F = F^\sigma \circ M_\sigma$ for every $\sigma \in K_C$.

Let us consider the Galois cover $Q : C \rightarrow B$, where $Q = F \circ P$. We note that, for $\sigma \in K_C$, we have (as $P^\sigma = P$)

$$Q^\sigma \circ f_\sigma = F^\sigma \circ P^\sigma \circ f_\sigma = F \circ M_\sigma^{-1} \circ M_\sigma \circ P \circ f_\sigma^{-1} \circ f_\sigma = R \circ P = Q.$$

Now we follow Dèbes and Emsalem's arguments [[1999](#)]. Assume we are able to find a point $c_1 \in C_1$ which is $\mathbb{M}(C)$ -rational and so that c_1 is not a branch value of the Galois covering Q . Fix a point $c \in C$ so that $Q(c) = c_1$. It follows that the H -stabilizer of c is trivial. We have the points $\sigma(c), f_\sigma(c) \in C^\sigma$. As

$$Q^\sigma(\sigma(c)) = \sigma(Q(c)) = \sigma(c_1) = c_1 \quad \text{and} \quad Q^\sigma(f_\sigma(c)) = Q(c) = c_1,$$

it follows that there is some $h_\sigma \in H$ so that $h_\sigma(f_\sigma(c)) = \sigma(c)$. Moreover, as a consequence of [Lemma 17](#) and the fact that c has trivial stabilizer in H , such $h_\sigma \in H$ is unique. In this way, we may assume that $f_\sigma(c) = \sigma(c)$ and, by the above, such an isomorphism is uniquely determined by σ . Again, by the uniqueness, this new family $\{f_\sigma : \sigma \in K_\lambda\}$ satisfies Weil's conditions and, by [Theorem 2](#), C is definable over its field of moduli.

In this way, in order to finish our proof, we only need find a $\mathbb{M}(C)$ -rational point on C_1 outside the branch set. This is equivalent to finding a point $r \in \widehat{\mathbb{C}} - \{\infty, 0, 1, \lambda\}$ with the property that $F(r) = \sigma(F(r))$, for every $\sigma \in K_C$. As $\sigma(F(r)) = F^\sigma(\sigma(r)) = F(M_\sigma^{-1}(\sigma(r)))$, we need to find a point $r \in \mathbb{C} - \{0, 1, \lambda\}$ such that

$$M_\sigma(r) = \sigma(r).$$

In this way, we need to find a point $r \in \mathbb{C} - \{0, 1, \lambda\}$ so that

- (1) if $\sigma(\lambda) = \lambda$, then $\sigma(r) = r$; and
- (2) if $\sigma(\lambda) = 1/\lambda$, then $\sigma(r) = r/\lambda$.

Condition (1) asserts that we need to find $r \in \mathbb{Q}(\lambda)$. Clearly, any point of the form $r = \alpha(1 + \lambda)$, where $\alpha \in \mathbb{Q}$ satisfies (1) and (2).

7.2. Let us now consider the case $\lambda = -1$. We have (see [Remark 5](#)) that either (i) $\text{Aut}_H(C) = H$ or (ii) $\text{Aut}(C) = \text{Aut}_H(C)$ and $[\text{Aut}(C) : H] = 2$.

In case (i) we may proceed as in the case $\lambda \neq -1$ as [Lemma 17](#) is still valid in this situation (the normalizer of H in $\text{Aut}(C)$ is H).

In case (ii) we have that $C/\text{Aut}(C) = (C/H)/\langle J \rangle$; that is, C is quasiplatonic, so it is defined over its field of moduli. \square

8. Proof of Theorem 13

Proof. Since

$$C_\lambda = \left\{ \begin{array}{l} z_0^p z_3^{p^n-p} + z_1^{p^{n-1}} z_3^{p^n-p^{n-1}} + z_2^{p^n} = 0 \\ \lambda z_0^p z_3^{p^{n-1}-p} + z_1^{p^{n-1}} + z_3^{p^{n-1}} = 0 \end{array} \right\} \subset \mathbb{P}^3$$

and

$$P([z_0 : z_1 : z_2 : z_3]) = -\left(\frac{z_1^{k^{n-1}}}{z_0^k z_3^{k^{n-1}-k}} \right),$$

then, for each $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q})$, one has that $C_\lambda^\sigma = C_{\sigma(\lambda)}$ and $P^\sigma = P$.

Let $K_\lambda = \{\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q}) : C_\lambda \cong C_{\sigma(\lambda)}\}$, so $\mathbb{M}(C_\lambda) = \text{Fix}(K_\lambda)$.

If $\sigma \in K_\lambda$, then there is an isomorphism $f_\sigma : C_\lambda \rightarrow C_{\sigma(\lambda)}$. As a consequence of Theorem 4, we may assume $f_\sigma H f_\sigma^{-1} = H$. So, there is a Möbius transformation M_σ such that $M_\sigma \circ P = P^\sigma \circ f_\sigma$. As M_σ must preserve the cone points and their orders, one has that

$$M_\sigma(\infty) = \infty, \quad M_\sigma(0) = 0, \quad M_\sigma\{1, \lambda\} = \{1, \sigma(\lambda)\}.$$

It follows from the first two equalities above that $M_\sigma(z) = Lz$, for a suitable $L \in \mathbb{C} - \{0\}$. The equality $M_\sigma\{1, \lambda\} = \{1, \sigma(\lambda)\}$ asserts that either (1) $L = 1$ and $\sigma(\lambda) = \lambda$ or (2) $L = \sigma(\lambda)$ and $\sigma(\lambda) = 1/\lambda$. As a consequence, we have proved (1) and (2).

Part (3) is a consequence of Theorem 11. □

9. Galois groups of order p^{n+1}

In this section, we consider those groups G of order $|G| = p^{n+1}$ acting on compact Riemann surfaces with signature $(0; p, p^{n-1}, p^n, p^n)$, for any odd prime p .

The algebraic structure for these groups is determined by the following result.

Proposition 18. *Let p be an odd prime number and let $G < \text{Aut}(S)$ be a group of order $|G| = p^{n+1}$ acting on a compact Riemann surface S with S/G of signature $(0; p, p^{n-1}, p^n, p^n)$.*

Then G is isomorphic to either

- (1) $\mathbb{Z}_{p^n} \times \mathbb{Z}_p$, or
- (2) $\langle x, y : x^{p^n} = y^p = 1, y^{-1}xy = x^{p^{n-1}+1} \rangle$.

Remark 19. in the first case we have provided, in Theorem 10, algebraic curves for S . In the second case explicit algebraic curves are more complicated, but we will study this problem elsewhere.

Proof. First notice that G has a presentation of the form

$$G = \langle x_1, x_2, x_3, x_4 : x_1^{p^n} = x_2^{p^n} = x_3^{p^{n-1}} = x_4^p = x_1 x_2 x_3 x_4 = 1, \mathcal{R} \rangle$$

where \mathcal{R} denotes other relations.

Therefore G cannot be cyclic, since otherwise it could not be generated by elements of the given orders.

Moreover, G has a cyclic subgroup of order p^n , which is normal because it has index p , and therefore G is isomorphic to

$$G \cong \mathbb{Z}_{p^n} \rtimes_{\sigma} \mathbb{Z}_p = \langle x \rangle \rtimes_{\sigma} \langle y \rangle$$

where $\sigma(x) = x^u$ with $u^p = 1 \pmod{p^n}$. The only solutions for u are $u = 1$ and the powers of $u = p^{n-1} + 1$, and the result follows. \square

Remark 20. We will denote the groups appearing in [Proposition 18](#) as follows:

$$(9-1) \quad G_u = \langle x, y : x^{p^n} = y^p = 1, y^{-1}xy = x^u \rangle$$

with $u = 1$ or $u = 1 + p^{n-1}$, and we will study the families of algebraic curves admitting G_u actions with signature $(0; p^n, p^n, p^{n-1}, p)$.

Lemma 21. *Consider the groups G_u given by (9-1) and*

$$(9-2) \quad \Gamma = \langle a_0, b_0, c_0, d_0 : a_0^p = b_0^{p^{n-1}} = c_0^{p^n} = d_0^{p^n} = a_0 b_0 c_0 d_0 = 1 \rangle.$$

Assume $\Phi : \Gamma \rightarrow G_u$ is an epimorphism such that $K = \ker \Phi$ is torsion-free.

Then either

- I) $K = \langle\langle b_0 c_0^{-p^q} a_0^{-\alpha}, a_0^{-1} c_0 a_0 c_0^{-u^s} \rangle\rangle$, with $0 \leq \alpha \leq p-1, 0 < s < p$ and $(q, p) = 1$,
or
- II) $K = \langle\langle a_0 c_0^{-p^{n-1}v}, b_0^p c_0^{-p^2q}, b_0^{-1} c_0 b_0 c_0^{-u^s} \rangle\rangle$, with $1 \leq v \leq p-1, 0 < s < p$
and $(q, p) = 1$,

where $\langle\langle \cdot \rangle\rangle$ denotes normal closure in Γ .

Proof. Since K is torsion-free, we obtain that

- a) $K \cap \langle a_0 \rangle = \{1\}$, and it follows that $y_1 = \Phi(a_0)$ has order p ;
- b) $K \cap \langle b_0 \rangle = \{1\}$, and it follows that $y_2 = \Phi(b_0)$ has order p^{n-1} ;
- c) $K \cap \langle c_0 \rangle = \{1\}$, and it follows that $y_3 = \Phi(c_0)$ has order p^n ;
- d) $K \cap \langle a_0 b_0 c_0 \rangle = \{1\}$, and it follows that $y_4 = \Phi(d_0)$ has order p^n .

Since Φ is an epimorphism, $\{y_1, y_2, y_3, y_4\}$ generate G_u . But clearly $y_4 = (y_1 y_2 y_3)^{-1}$, and therefore $\{y_1, y_2, y_3\}$ generate G_u .

We now examine the following two cases separately.

Case I Suppose $\langle y_1, y_3 \rangle = G_u$. We have $G_u = \langle y_3 \rangle \rtimes_{u^s} \langle y_1 \rangle$ for some $0 < s < p$. Also $y_2 = y_1^\alpha y_3^{pq}$ with $(q, p) = 1$. Hence

$$y_2 y_3^{-pq} y_1^{-\alpha} = \Phi(b_0 c_0^{-pq} a_0^{-\alpha}) = 1$$

and it follows that $b_0 c_0^{-pq} a_0^{-\alpha} \in K$.

Furthermore $\Phi(a_0^{-1} c_0 a_0 c_0^{-u^s}) = y_1^{-1} y_3 y_1 y_3^{-u^s} = 1$; hence $a_0^{-1} c_0 a_0 c_0^{-u^s} \in K$.

Then, checking the order of $\Gamma / \langle\langle b_0 c_0^{-pq} a_0^{-\alpha}, a_0 c_0^{-1} a_0^{-1} c_0^{-u^s} \rangle\rangle$, we obtain, as required,

$$K = \langle\langle b_0 c_0^{-pq} a_0^{-\alpha}, a_0 c_0^{-1} a_0^{-1} c_0^{-u^s} \rangle\rangle.$$

Case II Suppose $\langle y_1, y_3 \rangle < G_u$. Then

$$y_1 = y_3^{p^{n-1}v}$$

with $(v, p) = 1$, since $\langle y_3 \rangle$ is a maximal subgroup of G_u . Hence $a_0 c_0^{-p^{n-1}v} \in K$.

In this case,

$$\langle y_2, y_3 \rangle = G_u = \langle y_3 \rangle \rtimes_{u^s} \langle y_2 \rangle$$

for some $0 < s < p$. Hence $y_2^{-1} y_3 y_2 y_3^{-u^s} = 1$ from where $b_0^{-1} c_0 b_0 c_0^{-u^s} \in K$.

Finally, $y_2^p = y_3^{p^2q}$ with $(q, p) = 1$, from where $b_0^p c_0^{-p^2q} \in K$.

Again, by checking the order of $\Gamma / \langle\langle a_0 c_0^{-p^{n-1}v}, b_0^p c_0^{-p^2q}, b_0^{-1} c_0 b_0 c_0^{-u^s} \rangle\rangle$, we obtain

$$K = \langle\langle a_0 c_0^{-p^{n-1}v}, b_0^p c_0^{-p^2q}, b_0^{-1} c_0 b_0 c_0^{-u^s} \rangle\rangle. \quad \square$$

Considering the above notation for the elements $y_1 = \Phi(a_0)$, $y_2 = \Phi(b_0)$, $y_3 = \Phi(c_0)$ and $y_4 = \Phi(d_0)$ in G_u , we have the following result, which states that examples for both cases of [Proposition 18](#) exist, by the Riemann existence theorem.

Corollary 22. *If the group G_u , with $u = 1$ or $u = 1 + p^{n-1}$, acts on a compact Riemann surface with signature $(0; p, p^{n-1}, p^n, p^n)$, then a generating vector for the action may be chosen to be exactly of one of the following forms:*

- $(y_1, y_1^\alpha y_3^{pq}, y_3, y_3^{-1-pq} y_1^{-1-\alpha})$, with $(q, p) = 1$ and $1 \leq \alpha \leq p-2$;
- $(y_1, y_3^{pq}, y_3, y_3^{-1-pq} y_1^{-1})$, with $(q, p) = 1$;
- $(y_1, y_1^{-1} y_3^{pq}, y_3, y_3^{-1-pq})$, with $(q, p) = 1$;
- $(y_3^{p^{n-1}v}, y_2, y_3, y_3^{-1-p^{n-1}v} y_2^{-1})$.

In the first three cases the order of y_1 is p , the order of y_3 is p^n and $y_1^{-1} y_3 y_1 = y_3^{u^s}$ with $0 < s < p$. In the last case y_2 has order p^{n-1} , y_3 has order p^n , $y_2^p = y_3^{qp^2}$ and $y_2^{-1} y_3 y_2 = y_3^{u^s}$ with $0 < s < p$.

The following table gives the genera of some intermediate curves, where g_L denotes the genus of the quotient of S by the subgroup $L \leq \text{Aut}(S)$ and where V is any cyclic maximal subgroup acting freely:

generating vector	$u = 1 + p^{n-1}$	$u = 1$
$(y_1, y_1^\alpha y_3^{pq}, y_3, y_3^{-1-pq} y_1^{-1-\alpha})$	$g_{\langle y_3 \rangle} = \frac{p-1}{2}$ $g_{\langle y_3^{-1-pq} y_1^{-1-\alpha} \rangle} = \frac{p-1}{2}$ $g_{\langle y_1 \rangle} = \frac{2p^n - p^{n-2}(2p-1) - p}{2}$ $g_{\langle y_3^p \rangle} = p^2 - 2p + 1$ $g_{\langle y_3^p, y_1 \rangle} = 0$ $g_V = p - 1$	$g_{\langle y_3 \rangle} = \frac{p-1}{2}$ $g_{\langle y_3^{-1-pq} y_1^{-1-\alpha} \rangle} = \frac{p-1}{2}$ $g_{\langle y_1 \rangle} = \frac{p^n - p}{2}$ $g_{\langle y_3^p \rangle} = p^2 - 2p + 1$ $g_{\langle y_3^p, y_1 \rangle} = 0$ $g_V = p - 1$
$(y_1, y_3^{pq}, y_3, y_3^{-1-pq} y_1^{-1})$	$g_{\langle y_3 \rangle} = 0$ $g_{\langle y_3^{-1-pq} y_1^{-1} \rangle} = 0$ $g_{\langle y_1 \rangle} = \frac{2p^n - p^{n-2}(2p-1) - p}{2}$ $g_{\langle y_3^p \rangle} = \frac{p^2 - 3p}{2} + 1$ $g_{\langle y_3^p, y_1 \rangle} = 0$ $g_V = \frac{p-1}{2}$	$g_{\langle y_3 \rangle} = 0$ $g_{\langle y_3^{-1-pq} y_1^{-1} \rangle} = 0$ $g_{\langle y_1 \rangle} = \frac{p^n - p}{2}$ $g_{\langle y_3^p \rangle} = \frac{p^2 - 3p}{2} + 1$ $g_{\langle y_3^p, y_1 \rangle} = 0$ $g_V = \frac{p-1}{2}$
$(y_1, y_1^{-1} y_3^{pq}, y_3, y_3^{-1-pq})$	$g_{\langle y_3 \rangle} = 0$ $g_{\langle y_3^{-1-pq} \rangle} = 0$ $g_{\langle y_1 \rangle} = \frac{2p^n - p^{n-2}(2p-1) - p}{2}$ $g_{\langle y_3^p \rangle} = p^2 - 2p + 1$ $g_{\langle y_3^p, y_1 \rangle} = 0$ $g_V = p - 1$	$g_{\langle y_3 \rangle} = 0$ $g_{\langle y_3^{-1-pq} \rangle} = 0$ $g_{\langle y_1 \rangle} = \frac{p^n - p}{2}$ $g_{\langle y_3^p \rangle} = p^2 - 2p + 1$ $g_{\langle y_3^p, y_1 \rangle} = 0$ $g_V = p - 1$
$(y_3^{p^{n-1}v}, y_2, y_3, y_3^{-1-p^{n-1}v} y_2^{-1})$	$g_{\langle y_3 \rangle} = 0$ $g_{\langle y_3^{-1-p^{n-1}v} y_2^{-1} \rangle} = 0$ $g_{\langle y_1 \rangle} = \frac{2p^n - p^{n-1} - p}{2}$ $g_{\langle y_3^p \rangle} = \frac{p^2 - 3p}{2} + 1$ $g_{\langle y_3^p, y_1 \rangle} = 0$ $g_V = \frac{p-1}{2}$	$g_{\langle y_3 \rangle} = 0$ $g_{\langle y_3^{-1-p^{n-1}v} y_2^{-1} \rangle} = 0$ $g_{\langle y_1 \rangle} = \frac{2p^n - p^{n-1} - p}{2}$ $g_{\langle y_3^p \rangle} = \frac{p^2 - 3p}{2} + 1$ $g_{\langle y_3^p, y_1 \rangle} = 0$ $g_V = \frac{p-1}{2}$

References

[Beardon 1983] A. F. Beardon, *The geometry of discrete groups*, Graduate Texts in Mathematics **91**, Springer, New York, 1983. MR 85d:22026 Zbl 0528.30001

- [Bers 1975] L. Bers, “Automorphic forms for Schottky groups”, *Advances in Math.* **16** (1975), 332–361. [MR 51 #13218](#) [Zbl 0327.32011](#)
- [Burnside 1893] W. Burnside, “Note on the equation $y^2 = x(x^4 - 1)$ ”, *Proc. London Math. Soc.* **24**:1 (1893), 17–20. [JFM 25.0795.03](#)
- [Carvacho 2010] M. P. Carvacho, *Equivalence of group actions on Riemann surfaces*, Ph.D. thesis, Universidad de Chile, 2010. [arXiv 1012.4337](#)
- [Coombes and Harbater 1985] K. Coombes and D. Harbater, “Hurwitz families and arithmetic Galois groups”, *Duke Math. J.* **52**:4 (1985), 821–839. [MR 87g:14012](#) [Zbl 0601.14023](#)
- [Dèbes and Emsalem 1999] P. Dèbes and M. Emsalem, “On fields of moduli of curves”, *J. Algebra* **211**:1 (1999), 42–56. [MR 99k:14044](#) [Zbl 0934.14019](#)
- [Earle 1971] C. J. Earle, “On the moduli of closed Riemann surfaces with symmetries”, pp. 119–130 in *Advances in the theory of Riemann surfaces* (Stony Brook, NY, 1969), edited by L. V. Ahlfors, Ann. of Math. Studies **66**, Princeton Univ. Press, 1971. [MR 45 #5343](#) [Zbl 0218.32010](#)
- [Farkas and Kra 1992] H. M. Farkas and I. Kra, *Riemann surfaces*, 2nd ed., Graduate Texts in Mathematics **71**, Springer, New York, 1992. [MR 93a:30047](#) [Zbl 0764.30001](#)
- [González-Diez and Hidalgo 1997] G. González-Diez and R. A. Hidalgo, “Conformal versus topological conjugacy of automorphisms on compact Riemann surfaces”, *Bull. London Math. Soc.* **29**:3 (1997), 280–284. [MR 97k:14031](#) [Zbl 0865.14011](#)
- [González-Diez et al. 2009] G. González-Diez, R. A. Hidalgo, and M. Leyton, “Generalized Fermat curves”, *J. Algebra* **321**:6 (2009), 1643–1660. [MR 2010i:14050](#) [Zbl 1175.14023](#)
- [Hammer and Herrlich 2003] H. Hammer and F. Herrlich, “A remark on the moduli field of a curve”, *Arch. Math. (Basel)* **81**:1 (2003), 5–10. [MR 2004g:14030](#) [Zbl 1048.14013](#)
- [Hidalgo 2009] R. A. Hidalgo, “Non-hyperelliptic Riemann surfaces with real field of moduli but not definable over the reals”, *Arch. Math. (Basel)* **93**:3 (2009), 219–224. [MR 2010j:14063](#) [Zbl 1180.14028](#)
- [Hidalgo 2012] R. A. Hidalgo, “Homology closed Riemann surfaces”, *Q. J. Math.* **63**:4 (2012), 931–952. [Zbl 1256.30025](#)
- [Huggins 2007] B. Huggins, “Fields of moduli of hyperelliptic curves”, *Math. Res. Lett.* **14**:2 (2007), 249–262. [MR 2009a:14040](#) [Zbl 1126.14036](#)
- [Karcher and Weber 1999] H. Karcher and M. Weber, “The geometry of Klein’s Riemann surface”, pp. 9–49 in *The eightfold way*, edited by S. Levy, Math. Sci. Res. Inst. Publ. **35**, Cambridge Univ. Press, 1999. [MR 2001a:14027](#) [Zbl 0964.14029](#)
- [Klein 1878] F. Klein, “Ueber die Transformation siebenter Ordnung der elliptischen Functionen”, *Math. Ann.* **14**:3 (1878), 428–471. [MR 1509988](#) [JFM 11.0297.01](#)
- [Koebe 1907a] P. Koebe, “Über die Uniformisierung beliebiger analytischer Kurven, I”, *Nachr. Acad. Wiss. Göttingen* **1907** (1907), 177–190. [JFM 38.0453.01](#)
- [Koebe 1907b] P. Koebe, “Über die Uniformisierung beliebiger analytischer Kurven, II”, *Nachr. Acad. Wiss. Göttingen* **1907** (1907), 633–669. [JFM 38.0455.02](#)
- [Koizumi 1972] S. Koizumi, “The fields of moduli for polarized abelian varieties and for curves”, *Nagoya Math. J.* **48** (1972), 37–55. [MR 50 #4582](#) [Zbl 0246.14006](#)
- [Leyton and Hidalgo 2007] M. Leyton A. and R. A. Hidalgo, “On uniqueness of automorphisms groups of Riemann surfaces”, *Rev. Mat. Iberoam.* **23**:3 (2007), 793–810. [MR 2009c:30107](#) [MR 1144:30017](#)
- [Maclachlan 1965] C. Maclachlan, “Abelian groups of automorphisms of compact Riemann surfaces”, *Proc. London Math. Soc.* (3) **15** (1965), 699–712. [MR 31 #3596](#) [Zbl 0156.08902](#)

- [Poincaré 1907] H. Poincaré, “Sur l’uniformisation des fonctions analytiques”, *Acta Math.* **31**:1 (1907), 1–63. [MR 1555036](#) [JFM 38.0452.02](#)
- [Shimura 1972] G. Shimura, “On the field of rationality for an abelian variety”, *Nagoya Math. J.* **45** (1972), 167–178. [MR 46 #5342](#) [Zbl 0243.14012](#)
- [Singerman 1972] D. Singerman, “Finitely maximal Fuchsian groups”, *J. London Math. Soc.* (2) **6** (1972), 29–38. [MR 48 #529](#) [Zbl 0251.20052](#)
- [Torelli 1913] R. Torelli, “Sulle varietà di Jacobi”, *Rend. R. Accad. Lincei* **22**:2 (1913), 98–103. [JFM 44.0655.03](#)
- [Weil 1956] A. Weil, “The field of definition of a variety”, *Amer. J. Math.* **78** (1956), 509–524. [MR 18,601a](#) [Zbl 0072.16001](#)
- [Wolfart 2006] J. Wolfart, “*ABC* for polynomials, dessins d’enfants and uniformization: A survey”, pp. 313–345 in *Elementare und analytische Zahlentheorie* (Mainz, Germany, 2004), edited by S. Baier et al., Schr. Wiss. Ges. Johann Wolfgang Goethe Univ. Frankfurt am Main **20**, Steiner, Stuttgart, 2006. [MR 2009d:14031](#) [Zbl 1214.11079](#)

Received April 3, 2012. Revised February 3, 2013.

ANGEL CAROCCA
DEPARTAMENTO DE MATEMÁTICA Y ESTADÍSTICA
UNIVERSIDAD DE LA FRONTERA
CASILLA 54-D
4780000 TEMUCO
CHILE
angelcarocca@ufro.cl

RUBÉN A. HIDALGO
DEPARTAMENTO DE MATEMÁTICA
UNIVERSIDAD TÉCNICA FEDERICO SANTA MARÍA
CASILLA 110-V
2340000 VALPARAÍSO
CHILE
ruben.hidalgo@usm.cl

RUBÍ E. RODRÍGUEZ
FACULTAD DE MATEMÁTICAS
PONTIFICIA UNIVERSIDAD CATÓLICA DE CHILE
CASILLA 306–22
8320000 SANTIAGO
CHILE
rubi@mat.puc.cl

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Don Blasius
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2013 is US \$400/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 263 No. 1 May 2013

Biharmonic hypersurfaces in complete Riemannian manifolds	1
LUIS J. ALÍAS, S. CAROLINA GARCÍA-MARTÍNEZ and MARCO RIGOLI	
Half-commutative orthogonal Hopf algebras	13
JULIEN BICHON and MICHEL DUBOIS-VIOLETTE	
Superdistributions, analytic and algebraic super Harish-Chandra pairs	29
CLAUDIO CARMELI and RITA FIORESI	
Orbifolds with signature $(0; k, k^{n-1}, k^n, k^n)$	53
ANGEL CAROCCA, RUBÉN A. HIDALGO and RUBÍ E. RODRÍGUEZ	
Explicit isogeny theorems for Drinfeld modules	87
IMIN CHEN and YOONJIN LEE	
Topological pressures for ϵ-stable and stable sets	117
XIANFENG MA and ERCAI CHEN	
Lipschitz and bilipschitz maps on Carnot groups	143
WILLIAM MEYERSON	
Geometric inequalities in Carnot groups	171
FRANCESCO PAOLO MONTEFALCONE	
Fixed points of endomorphisms of virtually free groups	207
PEDRO V. SILVA	
The sharp lower bound for the first positive eigenvalue of the Folland–Stein operator on a closed pseudohermitian $(2n + 1)$-manifold	241
CHIN-TUNG WU	
Remark on “Maximal functions on the unit n-sphere” by Peter M. Knopf (1987)	253
HONG-QUAN LI	