

*Pacific
Journal of
Mathematics*

EXPLICIT ISOGENY THEOREMS FOR DRINFELD MODULES

IMIN CHEN AND YOONJIN LEE

EXPLICIT ISOGENY THEOREMS FOR DRINFELD MODULES

IMIN CHEN AND YOONJIN LEE

Let $F = \mathbb{F}_q(T)$ and $A = \mathbb{F}_q[T]$. Given two nonisogenous rank- r Drinfeld A -modules ϕ and ϕ' over K , where K is a finite extension of F , we obtain a partially explicit upper bound (dependent only on ϕ and ϕ') on the degree of primes \wp of K such that $P_\wp(\phi) \neq P_\wp(\phi')$, where $P_\wp(*)$ denotes the characteristic polynomial of Frobenius at \wp on a Tate module of $*$. The bounds are completely explicit in terms of the defining coefficients of ϕ and ϕ' , except for one term, which can be made explicit in the case of $r = 2$. An ingredient in the proof of the partially explicit isogeny theorem for general rank is an explicit bound for the different divisor of torsion fields of Drinfeld modules, which detects primes of potentially good reduction.

Our results are a Drinfeld module analogue of Serre's work (1981), but the results we obtain are unconditional because the generalized Riemann hypothesis holds for function fields.

1. Introduction

Let $A = \mathbb{F}_q[T]$, $F = \mathbb{F}_q(T)$, and let \bar{F} be a fixed algebraic closure of F , K a finite extension of F in \bar{F} , \bar{K} the algebraic closure of K in \bar{F} , \mathbb{O} the ring of integers of K , and \mathbb{F}_q a finite field of order q .

By a prime \wp (or place) of K , we mean a discrete valuation ring R with field of fractions K and maximal ideal \wp , and v denotes the discrete valuation associated to a prime \wp of K . For each place v of K , we fix a choice of \bar{K}_v and extend v to \bar{K}_v , which by abuse of notation we also call v . Also, when we speak of finite extensions of K_v , we assume they are initially given as subfields of \bar{K}_v .

Let ∞ be the infinite prime of F with corresponding discrete valuation

$$v_\infty(f/g) = \deg g - \deg f,$$

where $f, g \in A$. Let S_∞^K be the set of the infinite primes of K lying over ∞ , and

Chen was supported by the NSERC. Lee was supported by Priority Research Centers Program through the NRF funded by the Ministry of Education, Science and Technology (2009-0093827) and by the NRF grant funded by the Korea government (MEST) (2012-0005432). Lee is the corresponding author.
MSC2010: primary 11G09; secondary 11R58.

Keywords: Drinfeld modules, Galois representations, Chebotarev density theorem, isogeny theorem, different, ramification.

let $\bar{\infty} \in S_{\infty}^K$ have corresponding discrete valuation $v_{\bar{\infty}}$.

Let τ be the map that raises an element to its q -th power. A *Drinfeld A -module* ϕ over K is given by an \mathbb{F}_q -algebra homomorphism $i : A \rightarrow K$ and an \mathbb{F}_q -algebra homomorphism

$$\phi : A \rightarrow K\{\tau\}$$

such that ϕ_a has constant term $i(a)$ for any $a \in A$, and the image of ϕ is not contained in K .

A rank- r Drinfeld A -module ϕ over K is completely determined by

$$\phi_T = i(T) + a_1(\phi)\tau + \cdots + a_{r-1}(\phi)\tau^{r-1} + \Delta(\phi)\tau^r,$$

where $a_i(\phi), a_r = \Delta(\phi) \in K$ for $1 \leq i \leq r-1$. We call $\Delta(\phi)$ the *discriminant* of ϕ .

For any *monic* $a \in \mathbb{F}_q[T]$, we have

$$(1) \quad \phi_a = i(a) + \sum_{i=1}^{M-1} a_i(\phi, a)\tau^i + \Delta(\phi)^{(q^M-1)/(q^r-1)}\tau^M$$

for some $a_i(\phi, a) \in K$, where $M = r \deg_K a$.

For any $a \in A, a \neq 0$, we define the A -module of a -torsion points as

$$\phi[a] = \{\lambda \in \bar{K} \mid \phi_a(\lambda) = 0\}.$$

If I is a nonzero ideal of A , we similarly define the A -module of I -torsion points:

$$\phi[I] = \{\lambda \in \bar{K} \mid \phi_a(\lambda) = 0 \text{ for every } a \in I\}.$$

We have $\phi[a] \simeq (A/aA)^r$ if ϕ is of rank r [Rosen 2002, Proposition 12.4]. Let $K_{\phi,a} := K(\phi[a])$ be the field obtained by adjoining a -torsion points of ϕ to K , and let $K_{\phi,I} := K(\phi[I])$.

In the following, we briefly explain the definition of good reduction of a Drinfeld module. For more details, refer to [Goss 1996; Thakur 2004]. Let ϕ be a rank- r Drinfeld A -module over K and let \wp be a prime of K . Let \mathcal{O}_{\wp} be the valuation ring of \wp with the maximal ideal \wp and residue field $\mathbb{F}_{\wp} := \mathcal{O}_{\wp}/\wp$. We say that ϕ has *integral coefficients* at \wp if ϕ_a has coefficients in \mathcal{O}_{\wp} for all $a \in A$ and the reduction modulo \wp of these coefficients defines a Drinfeld module over \wp . The reduced Drinfeld module is denoted by ϕ^{\wp} .

Let ϕ and ϕ' be Drinfeld A -modules over K . Then a *morphism* f from ϕ to ϕ' over K is a polynomial f in $K\{\tau\}$ with the property that $f\phi_a = \phi'_a f$ for all $a \in A$. A nonzero morphism from ϕ to ϕ' over K is called an *isogeny* from ϕ to ϕ' (over K). If there exists an isogeny from ϕ to ϕ' over K , then we say that ϕ and ϕ' are *isogenous* (over K). An isogeny f from ϕ to ϕ' over K is called an *isomorphism* (over K) if there is an isogeny g from ϕ' to ϕ over K such that $fg = I = gf$, where I denotes the identity morphism. We note that ϕ and ϕ' are

isomorphic (over K) if and only if there is a $c \in K^*$ such that $c\phi_a = \phi'_a c$ for all $a \in A$ (for more details, refer to [Rosen 2002]).

We say that ϕ has *good reduction* at \wp if there exists a Drinfeld module ψ over K that is isomorphic to ϕ over K , ψ has integral coefficients at \wp , and ψ^\wp is a Drinfeld module of rank r .

By [Takahashi 1982] (see [Goss 1996, Theorem 4.10.5]; also [Goss 1992, Theorem 3.2.3] for one direction), we have that ϕ has good reduction at \wp if and only if the G_K -module $\phi[\mathcal{L}^\infty] := \bigcup_{m \geq 1} \phi[\mathcal{L}^m]$ is unramified at \wp , where G_K is the absolute Galois group of K and \mathcal{L} is a prime ideal of A different from \wp . This is the analogue for Drinfeld modules of the classical result of Ogg, Néron, and Shafarevich in the theory of abelian varieties.

If ϕ is a Drinfeld A -module defined over K and all its defining coefficients $a_i(\phi)$ lie in \mathbb{C} , then we say that ϕ is *integral over \mathbb{C}* . If ϕ is integral over \mathbb{C} , then it has good reduction outside any set of primes S of K that includes the primes lying over ∞ and the primes dividing the discriminant $\Delta(\phi)$ of ϕ . In particular, the G_K -modules $\phi[\mathcal{L}^\infty]$ and $\phi[\mathcal{L}]$ are unramified outside $S \cup \{\text{primes of } K \text{ lying over } \mathcal{L}\}$.

Let \mathcal{L} be a finite prime of A . The \mathcal{L} -torsion points of ϕ in \bar{K} give rise to a representation

$$\rho_{\phi, \mathcal{L}} : G_K \rightarrow \text{Aut}_{A/\mathcal{L}}(\phi[\mathcal{L}]) \cong \text{GL}_r(A/\mathcal{L}A),$$

where G_K is the absolute Galois group of K . For a prime \wp of K , if ϕ has good reduction at \wp , then $\rho_{\phi, \mathcal{L}}$ is unramified at \wp if \wp does not lie over \mathcal{L} .

For an unramified prime \wp of K , let $\text{Frob}_\wp \in G_K$ denote a Frobenius conjugacy class at \wp . Let $a_\wp(\phi)$ denote the trace of Frob_\wp on the $T_\mathcal{L}(\phi)$, and $P_\wp(\phi)(X)$ the characteristic polynomial of Frob_\wp on the $T_\mathcal{L}(\phi)$. It is known that $a_\wp(\phi)$ and $P_\wp(\phi)(X)$ are independent of \mathcal{L} [Goss 1996, Theorem 4.12.12].

Serre [1972] proved that if E is an elliptic curve over a number field L without complex multiplication, then there are only finitely many primes p such that the Galois representation $\rho_{E, p}$ on the p -torsion points of E is not surjective. The analogue of Serre's result [1972] for rank-2 Drinfeld A -modules was proved by Gardeyn [2001], that is, if ϕ is a rank-2 Drinfeld module over K without complex multiplication, then there are only finitely many primes \mathcal{L} such that $\rho_{\phi, \mathcal{L}}$ is not surjective. The case of general rank is proven in [Pink and Rüttsche 2009a; 2009b].

The following theorem is the Tate conjecture for rank- r Drinfeld A -modules over K , and its generalization to t-motives can be found in [Tamagawa 1994].

Theorem 1.1 [Taguchi 1996]. *Let ϕ, ϕ' be rank- r Drinfeld A -modules over K , and $A_\mathcal{L}$ the \mathcal{L} -adic completion of A . Then the natural homomorphism*

$$\text{Hom}_K(\phi, \phi') \otimes_A A_\mathcal{L} \rightarrow \text{Hom}_{A_\mathcal{L}[G_K]}(T_\mathcal{L}(\phi), T_\mathcal{L}(\phi'))$$

is an isomorphism, where $T_\mathcal{L}(\ast)$ is the \mathcal{L} -adic Tate module of \ast .

A consequence of the Tate conjecture is the isogeny theorem [Taguchi 1992, Proposition 3.1] that states that two Drinfeld A -modules ϕ, ϕ' over K are K -isogenous if and only if $P_{\wp}(\phi)(X) = P_{\wp}(\phi')(X)$ for all but finitely many primes \wp .

We prove the following partially explicit and effective version of the isogeny theorem for rank- r Drinfeld A -modules over K . For a Drinfeld A -module ϕ and a place \wp of K , define

$$\tau_{K,\wp}(\phi) = \inf \left\{ \frac{v_{\wp}(a_i(\phi))}{q^i - 1} : i = 1, \dots, r \right\}.$$

For any extension L/F , let $\gamma_L = [\mathbb{F}_L : \mathbb{F}_q]$. It is known that the constant field of

$$K_{\phi,\text{tor}} := K(\phi[a] : a \in A \text{ nonzero})$$

is finite over \mathbb{F}_q (see [David 2001, Lemma 3.2]), so we may define $\gamma_{\phi} = \gamma_{K_{\phi,\text{tor}}}$. More precisely, let $g_{\phi,\infty} = [K_{\infty}(\Lambda_{\phi,\infty}) : K_{\infty}]$, where $\Lambda_{\phi,\infty}$ is the lattice associated to the uniformization of ϕ over C_{∞} . Then we have

$$\gamma_{\phi} \leq g_{\phi} = \min\{g_{\phi,\infty} : \tilde{\infty} \mid \infty\}.$$

One can bound $g_{\phi,\infty}$ using knowledge of the successive minima of the lattices $\Lambda_{\phi,\infty}$ associated to ϕ [Gardeyn 2002, Proposition 4(i)]. Unfortunately, an explicit bound for these successive minima is not currently known except in the case of rank ≤ 2 [Chen and Lee 2013], so this term is currently inexplicit in general.

Throughout, $\ln x$ denotes the natural logarithm of x , $\log_q x$ denotes the logarithm of x to base q , and $\log_q^* x = \log_q \max\{x, 1\}$.

Theorem 1.2. *Let ϕ_1, ϕ_2 be rank- r Drinfeld A -modules that are integral over \mathbb{O} and not K -isogenous. Let S be the set consisting of the primes of K lying over the prime ∞ and the primes dividing $\Delta(\phi_1)\Delta(\phi_2)$. Suppose $\wp \notin S$ is a prime of K of least degree such that $P_{\wp}(\phi_1) \neq P_{\wp}(\phi_2)$. Then*

$$(2) \quad \deg_K \wp \leq \max \left\{ \frac{4}{m_0} (C_{q,r} + W + c_r s_{q,r} \log_q W), s \max\{1 + 2 \log_q s, 7\} \right\},$$

where

$s =$ the geometric extension degree of K/F ,

$$m_0 = \gamma_K,$$

$$c_r = 2r^2 + r + 1,$$

$$d_r = c_r + \log_q 86rs^2(g+1),$$

$$s_{q,r} = \frac{\ln(qd_r)}{\ln(qd_r) - 1},$$

$$C_{q,r} = \log_q 86rs^2(g+1) + c_r \left(1 + s_{q,r} \log_q \frac{4}{m_0} + \log_q d_r \right) + c_r s_{q,r} \log_q \log_q d_r,$$

$$a_r(\phi_i) = \Delta(\phi_i), \quad i = 1, 2,$$

$$W = \log_q^*(\Lambda_K(\phi_1, \phi_2) + 2D(\phi_1, \phi_2)) + g_{\phi_1} g_{\phi_2} m_0,$$

$$\text{where } D(\phi_1, \phi_2) = \deg_K \text{rad}_K \Delta(\phi_1) + \deg_K \text{rad}_K \Delta(\phi_2),$$

$$\Lambda_K(\phi_1, \phi_2) = - \sum_v \tau_{K,v}(\phi_1) \deg_K v - \sum_v \tau_{K,v}(\phi_2) \deg_K v,$$

$$\deg_K \text{rad}_K x = \sum_{v(x) \neq 0} \deg_K v.$$

(The sums are over every place v of K .)

Note that any Drinfeld A -module defined over K is isomorphic over K to a Drinfeld A -module that is integral over \mathbb{O} . In order to reduce the bounds given by the above theorem, in particular the quantity $\deg_K \text{rad}_K \Delta(\phi_1) \Delta(\phi_2)$, one should use minimal models of ϕ_1 and ϕ_2 (see [Taguchi 1993, Section 2]).

The proof follows the strategy in [Serre 1981] adapted to the Drinfeld module situation, with the notable difference that the effective Chebotarev density theorem we use [Kumar Murty and Scherk 1994] is stronger and unconditional because the general Riemann hypothesis holds for function fields. Also, unlike in the number field case, it is necessary to deal with wild ramification when bounding the different divisor. The bound we obtain on the different divisor is completely explicit in terms of the defining coefficients of the Drinfeld modules involved, unlike the results in [Gardeyn 2002], which are effective but not explicit. Also, the bounds are sensitive to primes of potentially good reduction, unlike the bounds in [Taguchi 1992].

We discuss some of the differences between our method and that of [Gardeyn 2002] in more detail in Section 7. In the rank-2 case, it is possible to make explicit the quantities involved in Gardeyn's bounds for the different divisor of torsion fields by determining the Newton polygons of exponential functions attached to Drinfeld modules [Chen and Lee 2013]. However, the computation of Newton polygons grows in complexity for higher rank, so new techniques using weaker information will likely be required to obtain explicit bounds for successive minima so we can apply the bounds of [Gardeyn 2002] for the different divisor and g_ϕ . Further remarks about this will be made in Section 7.

Cojocaru and David [2008] find upper bounds for the number of primes \wp of degree d such that the field extension of F obtained by adjoining a root of the characteristic polynomial of the Frobenius endomorphism of ϕ over the finite field A/\wp is the fixed field K , where ϕ is a Drinfeld module over K of rank 2 and K is an imaginary quadratic field over F . An ingredient in their proof requires the surjectivity results of Pink [1997] and Gardeyn [2001]. However, they do not require explicit versions of these in order to achieve their results; that is, they use the fact that the Galois representation $\rho_{\phi, \mathcal{L}} : \text{Gal}(F^{\text{sep}}/F) \rightarrow \text{GL}_2(A/\mathcal{L}A)$ and its projection in $\text{PGL}_2(A/\mathcal{L}A)$ are surjective for all but finitely many primes \mathcal{L} in A , assuming

$\text{End}_{\bar{F}}(\phi) = A$. As a method, they also use the effective version of the Chebotarev density theorem in [Kumar Murty and Scherk 1994], but for the different divisor bounds they only require the bounds in [Gardeyn 2002, Proposition 6].

2. Preliminaries

Let L be a finite extension of K and let \mathcal{O}_L be the maximal order of L , that is, the integral closure of \mathcal{O} in L . The *constant field* \mathbb{F}_L of L is the algebraic closure of \mathbb{F}_q in L . The *geometric extension degree* of L/K is the degree of L/K' , where K' is the maximal constant field extension of K in L (that is, $[L : K]/[\mathbb{F}_L : \mathbb{F}_K]$). We say L/K is a *geometric extension* if $K = K'$.

For a prime ideal \mathfrak{B} of \mathcal{O}_L , we let $\deg_L \mathfrak{B}$ be the \mathbb{F}_L -dimension of the residue class field $\mathbb{F}_{L, \mathfrak{B}} := \mathcal{O}_L/\mathfrak{B}$ of \mathfrak{B} , extending this to a general ideal I of \mathcal{O}_L by additivity on products. For a in \mathcal{O}_L , we define the degree of a by $\deg_L a := \deg_L(a)$, where (a) is the principal ideal of \mathcal{O}_L generated by a .

More generally, let \mathfrak{B} be a prime of L , $\mathcal{O}_{L, \mathfrak{B}}$ the valuation ring of \mathfrak{B} , and $\mathbb{F}_{L, \mathfrak{B}} := \mathcal{O}_{L, \mathfrak{B}}/\mathfrak{B}$ the residue class field of \mathfrak{B} . Then the *degree* of \mathfrak{B} is defined to be $\deg_L \mathfrak{B} := [\mathbb{F}_{L, \mathfrak{B}} : \mathbb{F}_L]$, the \mathbb{F}_L -dimension of $\mathbb{F}_{L, \mathfrak{B}}$. We extend the definition by linearity to a divisor $\mathfrak{D} = \sum_{\mathfrak{B}} n_{\mathfrak{B}} \mathfrak{B}$ of L by $\deg_L \mathfrak{D} = \sum_{\mathfrak{B}} n_{\mathfrak{B}} \deg_L \mathfrak{B}$. The *finite part* \mathfrak{D}_0 of a divisor $\mathfrak{D} = \sum_{\mathfrak{B}} n_{\mathfrak{B}} \mathfrak{B}$ is the divisor $\sum_{\mathfrak{B} \neq \infty} n_{\mathfrak{B}} \mathfrak{B}$.

Let $i_{L/K} : \text{Div}(K) \rightarrow \text{Div}(L)$ be the *conorm map* from divisors on K to divisors on L , defined by

$$i_{L/K}(\wp) = \sum_{\mathfrak{B}|\wp} e(\mathfrak{B}/\wp) \mathfrak{B}$$

for every prime \wp of K and then extended by linearity, where $e(\mathfrak{B}/\wp)$ denotes the *ramification index* of \mathfrak{B} over \wp .

For \mathfrak{B} a prime of L lying over the prime \wp of K , denote by $f(\mathfrak{B}/\wp)$ the *inertia degree* of \mathfrak{B} over \wp .

Lemma 2.1 [Rosen 2002, Proposition 7.7]. *Let L/K be a finite extension, \mathfrak{D} a divisor of K , and \mathfrak{B} a prime of L lying over the prime \wp of K . Then*

$$\deg_L i_{L/K} \mathfrak{D} = n' \deg_K \mathfrak{D}, \quad \deg_L \mathfrak{B} = \frac{f(\mathfrak{B}/\wp)}{[\mathbb{F}_L : \mathbb{F}_K]} \deg_K \wp,$$

where n' is the *geometric extension degree* of L/K .

Let L/K be a finite extension. Writing divisors in terms of places instead of primes, the *different divisor* $\mathfrak{D}(L/K)$ of L/K is defined as

$$\mathfrak{D}(L/K) = \sum_w w(D(L_w/K_v))w,$$

and its degree is given by

$$\deg_L \mathfrak{D}(L/K) = \sum_w w(D(L_w/K_v)) \deg_L w,$$

where w ranges through all normalized places of L , and $D(L_w/K_v)$ is the *different ideal* of L_w/K_v .

For convenience, we also define *the degree with respect to K* of $\mathfrak{D}(L/K)$ as

$$\deg_K \mathfrak{D}(L/K) = \sum_v \max\{v(D(L_w/K_v)) : w|v\} \deg_K v,$$

where v ranges through all normalized places of K . Similarly, we define the degree with respect to K of $\mathfrak{D}_0(L/K)$ as

$$\deg_K \mathfrak{D}_0(L/K) = \sum_{v \nmid \infty} \max\{v(D(L_w/K_v)) : w|v\} \deg_K v.$$

Lemma 2.2. *Let L/K be a finite extension. Then*

$$\deg_L \mathfrak{D}(L/K) \leq n' \deg_K \mathfrak{D}(L/K),$$

where n' is the geometric extension degree of L/K .

Proof. By the definition, we have

$$\begin{aligned} \deg_L \mathfrak{D}(L/K) &= \sum_w w(D(L_w/K_v)) \deg_L w \\ &= \sum_v \sum_{w|v} w(D(L_w/K_v)) \deg_L w \\ &= \sum_v \sum_{w|v} v(D(L_w/K_v)) e(w/v) f(w/v) \frac{1}{[\mathbb{F}_L : \mathbb{F}_K]} \deg_K v \\ &\leq \frac{1}{[\mathbb{F}_L : \mathbb{F}_K]} \sum_v \max\{v(D(L_w/K_v)) : w|v\} \sum_{w|v} e(w/v) f(w/v) \deg_K v \\ &= n' \sum_v \max\{v(D(L_w/K_v)) : w|v\} \deg_K v = n' \deg_K \mathfrak{D}(L/K), \end{aligned}$$

where \mathbb{F}_L and \mathbb{F}_K are the constant fields of L and K respectively, $f(w/v)$ denotes the *relative degree* of w over v , and we use the identity

$$[L : K] = \sum_{w|v} e(w/v) f(w/v),$$

which is valid because our constant fields are finite and hence perfect [Rosen 2002, Proposition 7.4]. \square

Lemma 2.3 [Serre 1979, Proposition 8, Chapter III.4]. *Let $M/L/K$ be a tower of finite separable extensions. The different divisor satisfies the transitivity relation*

$$\mathfrak{D}(M/K) = \mathfrak{D}(M/L) + i_{M/L}\mathfrak{D}(L/K).$$

Lemma 2.4. *Let K be a local field with ring of integers \mathbb{O} , and let L/K be a finite extension of K with ring of integers \mathbb{O}_L . Let $\alpha \in \mathbb{O}_L$ be such that $L = K(\alpha)$, and suppose $f(X) \in \mathbb{O}[X]$ is the minimal polynomial of α over K . Then the different ideal $D(\mathbb{O}_L/\mathbb{O})$ divides the ideal $(f'(\alpha))$, with equality holding if and only if $\mathbb{O}_L = \mathbb{O}[\alpha]$. Furthermore, we may replace $f(X)$ by any monic polynomial $g(X)$ in $\mathbb{O}[X]$ that α satisfies.*

Proof. See [Serre 1979, Corollary 2, III.6]. For the final remark, note that $g(X) = f(X)h(X)$ for some $g(X) \in \mathbb{O}[X]$, so $(g'(\alpha)) = (f'(\alpha)h(\alpha)) \subseteq (f'(\alpha))$. \square

Lemma 2.5. *Let E/K and L/K be finite extensions of local fields, with \mathbb{O} the ring of integers of K , \mathbb{O}_E the ring of integers of E , \mathbb{O}_{EL} the ring of integers of EL , and \mathbb{O}_L the ring of integers of L .*

Then the different ideals satisfy $D(EL/L) \mid \mathbb{O}_{EL} \cdot D(E/K)$.

Proof. Suppose that $\mathbb{O}_E = \mathbb{O}_K[x]$ for some $x \in B$, so that $E = K(x)$ (see [Serre 1979, Proposition 12, III.6]). Let $f \in \mathbb{O}_K[X]$ be the minimal polynomial of x over K .

Now $EL = K(x)L = K(x)$ and $x \in \mathbb{O}_{EL}$.

As $f \in \mathbb{O}[X]$ is monic and $x \in \mathbb{O}_{EL}$ is a root of f , we may apply Lemma 2.4 to get that $D(EL/L) \mid \mathbb{O}_{EL} \cdot f'(x)$. But as $\mathbb{O}_E = \mathbb{O}[x]$, we have $D(E/K) = \mathbb{O}_E \cdot f'(x)$. Hence, $\mathbb{O}_{EL} \cdot f'(x) = \mathbb{O}_{EL} \cdot \mathbb{O}_E \cdot f'(x) = \mathbb{O}_{EL} \cdot D(E/K)$. The result thus follows. \square

Lemma 2.6. *Let E/K and L/K be finite extensions of global fields. Then*

$$\mathfrak{D}(EL/K) \leq i_{EL/E}\mathfrak{D}(E/K) + i_{EL/L}\mathfrak{D}(L/K).$$

Proof. This follows by localization and applying Lemma 2.3 and Lemma 2.5. \square

3. Effective Chebotarev density theorem

Lemma 3.1. *Let K be a finite extension of $F = \mathbb{F}_q(T)$ with constant field \mathbb{F}_q , where \mathbb{F}_q is a finite field of order q , and let g be the genus of K . Let $S(N)$ be the number of primes \wp of K with $\deg_K \wp = N$. Then*

$$\left| S(N) - \frac{q^N}{N} \right| \leq \left(2g + 1 + \left(2g + \frac{3}{2} \right) \frac{4}{q} \right) \frac{q^{N/2}}{N}.$$

Proof. From the prime number theorem for L [Rosen 2002, Theorem 5.12], we have that

$$S(N) = \frac{q^N}{N} + O\left(\frac{q^{N/2}}{N}\right).$$

We recall the proof to make the constant explicit.

Let $Z_K(u)$ be the zeta function of K . Using the Euler product decomposition of $Z_K(u)$ and [Rosen 2002, Theorem 5.9], we obtain

$$Z_K(u) = \frac{\prod_{i=1}^{2g} (1 - \pi_i u)}{(1-u)(1-qu)} = \prod_{d=1}^{\infty} (1 - u^d)^{-S(d)}.$$

Taking the logarithmic derivative of both sides, multiplying by u , and equating coefficients of u^N yields the relation

$$q^N + 1 - \sum_{i=1}^{2g} \pi_i^N = \sum_{d|N} dS(d).$$

Using the Möbius inversion formula yields

$$NS(N) = \sum_{d|N} \mu(d)q^{N/d} + 0 - \sum_{d|N} \mu(d) \left(\sum_{i=1}^{2g} \pi_i^{N/d} \right).$$

Following the argument in [Rosen 2002, Theorem 2.2], we obtain

$$\left| \sum_{d|N} \mu(d)q^{N/d} - q^N \right| \leq q^{N/2} + Nq^{N/3}.$$

Similarly, using the Riemann hypothesis [Rosen 2002, Theorem 5.10], we obtain

$$\left| \sum_{d|N} \mu(d) \left(\sum_{i=1}^{2g} \pi_i^{N/d} \right) \right| \leq 2gq^{N/2} + 2gNq^{N/4}.$$

It follows that

$$|NS(N) - q^N| \leq (2g+1)q^{N/2} + Nq^{N/3} + 2gNq^{N/4},$$

so

$$(3) \quad \left| S(N) - \frac{q^N}{N} \right| \leq \frac{2g+1}{N}q^{N/2} + q^{N/3} + 2gq^{N/4} \\ \leq \frac{q^{N/2}}{N} \left(2g+1 + \frac{N}{q^{N/6}} + 2g\frac{N}{q^{N/4}} \right).$$

Since $x/q^x \leq 1/q$ for $x \geq 1$, (3) is less than or equal to

$$\frac{q^{N/2}}{N} \left(2g+1 + (2g+\frac{3}{2})\frac{4}{q} \right). \quad \square$$

The next theorem follows from the effective Chebotarev density theorem in [Kumar Murty and Scherk 1994, Theorem 1].

Theorem 3.2. *Let K be a finite extension of $F = \mathbb{F}_{q_0}(T)$ with constant field \mathbb{F}_q and genus g , where $q = q_0^{m_0}$. Let E be a finite Galois extension of K with Galois group G , \mathbb{F}_{q^m} the algebraic closure of \mathbb{F}_q in E , and $K' = \mathbb{F}_{q^m}K$ the maximal constant field extension of K in E .*

Let $\mathcal{C} \subseteq G = \text{Gal}(E/K)$ be a nonempty conjugacy class in G whose restriction to $\mathbb{F}_{q^m}/\mathbb{F}_q \cong K'/K$ is τ^k , where τ is the Frobenius map $\tau(x) = x^q$, and let \mathcal{D} be the different divisor of E/K' . Let Σ be the divisor of K that is the sum of the primes of K that are ramified in E , and suppose Σ' is a divisor of K such that $\Sigma' \geq \Sigma$. Let $B = \max\{\deg_K \Sigma', \deg_E \mathcal{D}, 2|\text{Gal}(E/K')| - 2, 1\}$.

If

$$N \geq \frac{2}{m_0} \log_{q_0} \frac{4}{3} \left(B^2 + B \left(2g + \frac{g}{m} + 3 \right) + 2 \left(5g + \frac{g}{m} + 3 \right) \right)$$

and $N \equiv k \pmod{m}$, there is a prime $\wp \notin \Sigma'$ of K such that $\deg_K \wp = N$ and $\text{Frob}_\wp = \mathcal{C}$.

Proof. The situation at the outset is that we start with $F = \mathbb{F}_{q_0}(T)$ and K a finite extension of F with possibly larger constant field \mathbb{F}_q , where $q = q_0^n$. Next, we replace $F = \mathbb{F}_{q_0}(T)$ by $F = \mathbb{F}_q(T)$, so that K is a geometric extension of $F = \mathbb{F}_q(T)$. This allows us to use [Lemma 3.1](#) without modification, but now q_0 is replaced by q .

Another remark is that if there exists a prime $\wp \notin \Sigma'$ of K such that $\deg_K \wp = N$ and $\text{Frob}_\wp = \mathcal{C}$, then it follows that \mathcal{C} restricted to $K'/K \cong \mathbb{F}_{q^m}/\mathbb{F}_q$ is τ^N by [\[Kumar Murty and Scherk 1994, Lemma 1\]](#). Since $\text{Gal}(\mathbb{F}_{q^m}/\mathbb{F}_q)$ is cyclic of order m , we have that $\tau^N = \tau^k$ in $\text{Gal}(\mathbb{F}_{q^m}/\mathbb{F}_q)$ if and only if $N \equiv k \pmod{m}$.

Let \mathbb{F}_{q^m} be the algebraic closure of \mathbb{F}_q in E , so $K' := \mathbb{F}_{q^m}K$ and E/K' is a geometric extension. Let $\mathcal{D} := \deg_E \mathcal{D}$ and $\delta' = \deg_K \Sigma'$. Let $\pi(N, \Sigma')$ be the number of primes $\wp \notin \text{Supp } \Sigma'$ of K with $\deg_K \wp = N$, and let $\pi_{\mathcal{C}}(N, \Sigma')$ be the number of primes $\wp \notin \text{Supp } \Sigma'$ of K such that $\deg_K \wp = N$ and $\text{Frob}_\wp = \mathcal{C}$.

It suffices to find a lower bound N_0 for N such that for $N \geq N_0$, $\pi_{\mathcal{C}}(N, \Sigma')$ is positive.

In fact, the genus g of K over \mathbb{F}_q is the same as that of K' over \mathbb{F}_{q^m} (see [\[Rosen 2002, Proposition 8.9\]](#)). We know that the genus of K' over \mathbb{F}_{q^m} and the genus of E over \mathbb{F}_{q^m} are related by the Riemann–Hurwitz theorem [\[Rosen 2002, Theorem 7.16\]](#). Thus, letting g_E be the genus of E , we have

$$(4) \quad g_E = 1 + |\text{Gal}(E/K')|(g - 1) + \frac{1}{2}\mathcal{D}.$$

The effective Chebotarev density theorem in [\[Kumar Murty and Scherk 1994, Theorem 1\]](#) gives

$$\frac{m|\mathcal{C}|}{|G|} \pi(N, \Sigma') - \alpha \leq \pi_{\mathcal{C}}(N, \Sigma') \leq \frac{m|\mathcal{C}|}{|G|} \pi(N, \Sigma') + \alpha,$$

where

$$(5) \quad \alpha = \frac{|\mathcal{C}|}{N} q^{N/2} \left(2g_E \frac{1}{|G|} + 2(2g+1) + \frac{1+N/|\mathcal{C}|}{q^{N/2}} \delta' \right).$$

The condition $N \equiv r \pmod{m}$ ensures \mathcal{C} restricted to $\mathbb{F}_{q^m}/\mathbb{F}_q$ is τ^N .

Remark 3.3. When $\Sigma' = \Sigma$, this is what is proved in [Kumar Murty and Scherk 1994, Theorem 1]. However, the proof carries over with Σ replaced by Σ' . In particular, the key identity (2.1) still holds with $y \in Y_r$ unramified replaced by $y \in Y_r$ not in the support of $\Sigma' \geq \Sigma$.

We have $\pi(N, \Sigma') \geq S(N) - \frac{\deg_K \Sigma'}{N}$. Thus,

$$\frac{m^{|\mathcal{C}|}}{|G|} \left(S(N) - \frac{\deg_K \Sigma'}{N} \right) - \alpha \leq \pi_{\mathcal{C}}(N, \Sigma').$$

It is therefore enough to find a lower bound for N such that

$$(6) \quad \frac{m^{|\mathcal{C}|}}{|G|} \left(S(N) - \frac{\deg_K \Sigma'}{N} \right) - \alpha > 0.$$

From Lemma 3.1, we have

$$(7) \quad \frac{q^N}{N} - \left(2g+1 + \left(2g + \frac{3}{2} \right) \frac{4}{q} \right) \frac{q^{N/2}}{N} \leq S(N) \\ \leq \frac{q^N}{N} + \left(2g+1 + \left(2g + \frac{3}{2} \right) \frac{4}{q} \right) \frac{q^{N/2}}{N}.$$

Since $|G|/m = |\text{Gal}(E/L')|$ and $(1+N/|\mathcal{C}|)/q^{N/2} \leq 2$, from (5) we have

$$(8) \quad \alpha \leq \frac{2m^{|\mathcal{C}|}}{N|G|} q^{N/2} \left(|\text{Gal}(E/K')| (2g+1+\delta') + \frac{g_E}{m} \right).$$

Therefore, combining (6) through (8), we obtain

$$(9) \quad \frac{m^{|\mathcal{C}|}}{|G|} \left(S(N) - \frac{\deg_K \Sigma'}{N} \right) - \alpha \\ \geq \frac{m^{|\mathcal{C}|}}{N|G|} q^{N/2} \left(q^{N/2} - \left(c_0 + \frac{\deg_K \Sigma'}{q^{N/2}} + 2|\text{Gal}(E/K')| (2g+1+\delta') + 2\frac{g_E}{m} \right) \right),$$

where $c_0 = 2g+1 + \left(2g + \frac{3}{2} \right) 4/q$.

We thus need to find a lower bound of N such that the right-hand side of the

inequality in (9) is positive, or equivalently,

$$\begin{aligned}
 q^{N/2} &> c_0 + \frac{\deg_K \Sigma'}{q^{N/2}} + 2|\mathrm{Gal}(E/K')|(2g+1+\delta') + 2\frac{gE}{m} \\
 &= c_0 + \frac{\deg_K \Sigma'}{q^{N/2}} + 2|\mathrm{Gal}(E/K')|(2g+1+\delta') \\
 &\quad + \frac{2}{m}(1+|\mathrm{Gal}(E/K')|(g-1) + \frac{1}{2}\mathfrak{D}) \\
 &= c_0 + \frac{\deg_K \Sigma'}{q^{N/2}} + 2|\mathrm{Gal}(E/K')|\left(2g+1+\delta' + \frac{g-1}{m}\right) + \frac{2}{m}\left(1 + \frac{1}{2}\mathfrak{D}\right),
 \end{aligned}$$

using (4).

Let $1 \leq B$, $\delta' \leq B$, $\mathfrak{D} \leq B$, and $|\mathrm{Gal}(E/K')| \leq \frac{1}{2}B + 1$. Note that if $g = 0$, it suffices to take $\delta' \leq B$ and $\mathfrak{D} \leq B$ only, as it is then automatic that $|\mathrm{Gal}(E/K')| \leq \frac{1}{2}\mathfrak{D} + 1 \leq \frac{1}{2}B + 1$. Therefore, we have

$$\begin{aligned}
 c_0 + \frac{\deg_K \Sigma'}{q^{N/2}} + 2|\mathrm{Gal}(E/K')|\left(2g+1+\delta' + \frac{g-1}{m}\right) + \frac{2}{m}\left(1 + \frac{1}{2}\mathfrak{D}\right) \\
 \leq c_0 + \frac{B}{q^{N/2}} + (B+2)\left(2g+1+B + \frac{g-1}{m}\right) + \frac{2}{m}\left(1 + \frac{1}{2}B\right) \\
 \leq 2g+1 + (2g+\frac{3}{2})\frac{4}{q} + \frac{B}{q^{N/2}} + (B+2)\left(2g+1+B + \frac{g-1}{m}\right) + \frac{2}{m}\left(1 + \frac{1}{2}B\right) \\
 \leq B^2 + B\left(2g+3 + \frac{g}{m}\right) + 6g+3 + \frac{2g}{m} + (2g+\frac{3}{2})\frac{4}{q} + \frac{B}{q^{N/2}} \\
 \leq B^2 + B\left(2g+3 + \frac{g}{m}\right) + 10g+6 + \frac{2g}{m} + \frac{B}{q^{N/2}},
 \end{aligned}$$

where the last inequality uses $4/q \leq 2$. Therefore, it suffices to have

$$q^{N/2} > \left(B^2 + B\left(2g+3 + \frac{g}{m}\right) + 10g+6 + \frac{2g}{m}\right) + \frac{B}{q^{N/2}}.$$

This can be satisfied if the following two inequalities hold:

$$\alpha q^{N/2} \geq B^2 + B\left(2g+3 + \frac{g}{m}\right) + 10g+6 + \frac{2g}{m}, \quad (1-\alpha)q^{N/2} > \frac{B}{q^{N/2}},$$

where $0 < \alpha < 1$; equivalently,

$$N \geq 2 \log_q \frac{1}{\alpha} \left(B^2 + B\left(2g+3 + \frac{g}{m}\right) + 10g+6 + \frac{2g}{m} \right), \quad N > \log_q \frac{1}{1-\alpha} B.$$

Taking $\alpha = \frac{3}{4}$, the required inequalities become

$$N \geq 2 \log_q \frac{4}{3} \left(B^2 + B\left(2g+3 + \frac{g}{m}\right) + 10g+6 + \frac{2g}{m} \right), \quad N > \log_q 4B.$$

So if

$$N \geq \frac{2}{m_0} \log_{q_0} \frac{4}{3} \left(B^2 + B \left(2g + 3 + \frac{g}{m} \right) + 2 \left(5g + 3 + \frac{g}{m} \right) \right)$$

and $N \equiv k \pmod{m}$, then there is a prime $\wp \notin \Sigma'$ of K such that $\deg_K \wp = N$ and $\text{Frob}_\wp = \mathcal{C}$. \square

Corollary 3.4. *Let the notation and hypotheses be as in Theorem 3.2. Then there exists a prime $\wp \notin \Sigma'$ of K such that $\text{Frob}_\wp = \mathcal{C}$ and*

$$(10) \quad \deg_K \wp \leq \frac{4}{m_0} \log_{q_0} \frac{4}{3} (B + 3g + 3) + m.$$

Proof. Let M be the integer such that

$$M = \frac{2}{m_0} \log_{q_0} \frac{4}{3} \left(B^2 + B \left(2g + \frac{g}{m} + 3 \right) + 2 \left(5g + \frac{g}{m} + 3 \right) \right) + \delta,$$

where $0 \leq \delta < 1$. Let $N = M + k'$, where $0 \leq k' \leq m - 1$ is chosen so that $N \equiv k \pmod{m}$. Then

$$N \geq \frac{2}{m_0} \log_{q_0} \frac{4}{3} \left(B^2 + B \left(2g + \frac{g}{m} + 3 \right) + 2 \left(5g + \frac{g}{m} + 3 \right) \right)$$

and $N \equiv k \pmod{m}$. By Theorem 3.2, there exists a prime $\wp \notin \Sigma'$ of K such that $\deg_K \wp = N$ and $\text{Frob}_\wp = \mathcal{C}$. Now,

$$\begin{aligned} \deg_K \wp &= N = M + k' \\ &\leq \frac{2}{m_0} \log_{q_0} \frac{4}{3} \left(B^2 + B \left(2g + 3 + \frac{g}{m} \right) + 10g + 6 + \frac{2g}{m} \right) + m \\ &\leq \frac{2}{m_0} \log_{q_0} \frac{4}{3} \left(B + 2g + 3 + \frac{g}{m} \right)^2 + m \\ &\leq \frac{4}{m_0} \log_{q_0} \frac{4}{3} (B + 3g + 3) + m. \end{aligned} \quad \square$$

4. Bounds for the different divisor

Proposition 4.1. *Let ϕ be a rank- r Drinfeld A -module that is integral over K , and let $\mathfrak{L} = (a)$ be a finite prime of A with a monic. Let $\mathfrak{D}_0(K_{\phi, \mathfrak{L}}/K)$ be the finite part of the different divisor $\mathfrak{D}(K_{\phi, \mathfrak{L}}/K)$. Then we have*

$$(11) \quad \deg_K \mathfrak{D}_0(K_{\phi, \mathfrak{L}}/K) \leq r \left(\deg_K a + \frac{(\ell^r - 2)(\ell^r - 1)}{q^r - 1} \deg_K \Delta(\phi) \right),$$

where $\ell = q^{\deg_F \mathfrak{L}}$. In addition, if $v(a\Delta(\phi)) = 0$ for a finite place v of K , then

$$(12) \quad v(D(K_{\phi, \mathfrak{L}, w}/K_v)) = 0,$$

where $D(K_{\phi, \mathfrak{L}, w}/K_v)$ is the different ideal of $K_{\phi, \mathfrak{L}, w}/K_v$, and $w|v$ is a place of $K_{\phi, \mathfrak{L}, w}$.

Proof. This is a slightly modified version of [David 2001, Lemma 4.2], derived from [Taguchi 1992].

Let $\alpha \in \bar{K}$ be a root of a separable polynomial

$$f(X) = b_0X + b_1X^q + \cdots + b_mX^{q^m}$$

with $b_i \in \mathbb{C}$ and $b_0b_m \neq 0$. Then

$$\begin{aligned} h(X) &= b_m^{q^m-1} f(X/b_m) \\ &= b_0b_m^{q^m-2}X + b_1b_m^{q^m-1-q}X^q + \cdots + b_{m-1}b_m^{q^m-1-q^{m-1}}X^{q^{m-1}} + X^{q^m} \in \mathbb{C}[X] \end{aligned}$$

is monic. Since $h(b_m\alpha) = 0$ and $K(\alpha) = K(b_m\alpha)$, we may apply Lemma 2.4 to $b_m\alpha$ and $h(X)$ to show that the different ideal $D(K(\alpha)/K)$ divides the principal ideal $(b_0b_m^{q^m-2})$.

Let $\mathfrak{L} = (a)$ and $f(X) = \phi_a(X)$. Then $f(X) = aX + \cdots + \Delta(\phi)^{(q^m-1)/(q^r-1)}X^{q^m}$, where $m = r \deg_F a$ (see [Rosen 2002, Proposition 13.8]). There are r roots β_1, \dots, β_r of $\phi_a(X)$ that generate $K_{\phi, \mathfrak{L}}$ over K . Using the transitivity of the different (see Lemma 2.3), it follows that

$$(13) \quad D(K_{\phi, \mathfrak{L}}/K) \mid (b_0b_m^{q^m-2})^r = (a(\Delta(\phi))^{(q^m-2)(q^m-1)/(q^r-1)})^r.$$

This shows that if $v(a\Delta(\phi)) = 0$ for a finite place v , then $v(D(K_{\phi, \mathfrak{L}, w}/K_v)) = 0$. Furthermore, taking the degree with respect to K of (13), we obtain

$$\deg_K \mathfrak{D}_0(K_{\phi, \mathfrak{L}}/K) \leq r \left(\deg_K a + \frac{(\ell^r - 2)(\ell^r - 1)}{q^r - 1} \deg_K \Delta(\phi) \right). \quad \square$$

It is possible to obtain a bound on $\deg_K \mathfrak{D}(K_{\phi, \mathfrak{L}}/K)$ based on Proposition 4.1 and Lemma 4.2, but instead we find a slightly more refined bound in Proposition 4.3, using additional techniques.

Lemma 4.2. *Let $\bar{\infty}$ be an infinite prime of K , $K_{\bar{\infty}}$ the completion of K at $\bar{\infty}$, $\mathbb{C}_{\bar{\infty}}$ the valuation ring of $\bar{\infty}$, $v_{\bar{\infty}}$ the valuation associated to $\bar{\infty}$, and e the ramification index of $\bar{\infty}$ over ∞ .*

Let $\phi_T(X) = TX + a_1X^q + \cdots + a_iX^{q^i} + \cdots + a_rX^{q^r}$ be a rank- r Drinfeld A -module defined over K , and write

$$\phi_{T^n}(X) = T^nX + b_1X^q + \cdots + b_iX^{q^i} + \cdots + b_{rn}X^{q^{rn}},$$

where $n \geq 1$.

Let $\omega_1 = \max \left\{ e, -\frac{v_{\bar{\infty}}(a_i)}{q^i} : i = 1, \dots, r \right\}$ and $\omega_n = n\omega_1$. Then

$$\omega_n \geq \max \left\{ ne, -\frac{v_{\bar{\infty}}(b_i)}{q^i} : i = 1, \dots, rn \right\}.$$

Proof. We use induction on n . First note that

$$\phi_{T^n}(\lambda_n X) = T^n \lambda_n X + b_1 \lambda_n^q X^q + \cdots + b_i \lambda_n^{q^i} X^{q^i} + \cdots + b_{rn} \lambda_n^{q^{rn}} X^{q^{rn}},$$

so taking $\lambda_n \in K$ with

$$v_{\infty}(\lambda_n) = \omega_n \geq \max \left\{ ne, -\frac{v_{\infty}(b_i)}{q^i} : i = 1, \dots, rn \right\}$$

implies that $\phi_{T^n}(\lambda_n X) \in \mathbb{O}_{\infty}[X]$.

The result is true for $n = 1$, as

$$\omega_1 = \max \left\{ e, -\frac{v_{\infty}(a_i)}{q^i} : i = 1, \dots, r \right\}.$$

Assume

$$\omega_n = n\omega_1 \geq \max \left\{ ne, -\frac{v_{\infty}(b_i)}{q^i} : i = 1, \dots, rn \right\}.$$

Now consider the terms in the product

$$\phi_{T^{n+1}} = \phi_{T^n} \circ \phi_T = (T^n + b_1 \tau + \cdots + b_{rn} \tau^{rn}) \circ (T + a_1 \tau + \cdots + a_r \tau^r),$$

where there are $2(r+1)$ types of terms to consider:

$$\begin{aligned} b_i \tau^i T &= b_i T^{q^i} \tau^i, & 1 \leq i \leq rn, \\ b_i \tau^i a_1 \tau &= b_i a_1^{q^i} \tau^{i+1}, & 1 \leq i \leq rn, \\ &\vdots \\ b_i \tau^i a_r \tau^r &= b_i a_r^{q^i} \tau^{i+r}, & 1 \leq i \leq rn, \\ T^{n+1}, T^n a_1 \tau, T^n a_2 \tau^2, \dots, T^n a_r \tau^r. \end{aligned}$$

We need to show that ω_{n+1} is greater than or equal to the valuations of the coefficients of each type of term, that is, for each i with $1 \leq i \leq rn$,

$$(14) \quad \omega_{n+1} \geq -\frac{v_{\infty}(b_i)}{q^i} + e,$$

$$(15) \quad \omega_{n+1} \geq -\frac{v_{\infty}(b_i)}{q^{i+j}} - \frac{v_{\infty}(a_j)}{q^j}, \quad 1 \leq j \leq r,$$

$$(16) \quad \omega_{n+1} \geq ne + 1,$$

$$(17) \quad \omega_{n+1} \geq ne - \frac{v_{\infty}(a_j)}{q^j}, \quad 1 \leq j \leq r.$$

As $\omega_n \geq -v_{\infty}(b_i)/q^i$ for $1 \leq i \leq 2n$, we have

$$\omega_{n+1} = \omega_n + \omega_1 \geq \frac{\omega_n}{q^j} + \omega_1 \geq -\frac{v_{\infty}(b_i)}{q^{i+j}} + \omega_1$$

for $j = 0, 1, \dots, r$ and $i = 1, 2, \dots, rn$, so (14) and (15) are satisfied. Since $\omega_1 = \max\{e, -v_\infty(a_j)/q^j : j = 1, \dots, r\}$,

$$\begin{aligned} \omega_{n+1} &= (n+1)\omega_1 = n\omega_1 + \omega_1 \geq ne + \omega_1 \\ &\geq \max\left\{(n+1)e, ne - \frac{v_\infty(a_j)}{q^j} : j = 1, \dots, r, \right\} \end{aligned}$$

so the last inequalities in (16) and (17) are satisfied. \square

In the following proposition, we obtain an upper bound on the degree of the different divisor of $K_{\phi, \mathfrak{L}}/K$ that uses mild information from the Newton polygons of $\phi_a(X)$ and takes into account primes of potentially good reduction.

Proposition 4.3. *Let ϕ be a rank- r Drinfeld A -module that is integral over K , and let $\mathfrak{L} = (a)$ be a finite prime of A with a monic. Let $\mathfrak{D}(K_{\phi, \mathfrak{L}}/K)$ be the different divisor of $K_{\phi, \mathfrak{L}}/K$. Then we have*

$$\deg_K \mathfrak{D}(K_{\phi, \mathfrak{L}}/K) \leq r \left(\frac{\ell^r - 1}{q - 1} (s \deg_K a + \Lambda(\phi)) + 2 \deg_K a \operatorname{rad}_K \Delta(\phi) \right),$$

where s denotes the geometric extension degree of K/F , $\ell = q^{\deg_F \mathfrak{L}}$, $\Lambda(\phi) = -\sum_v \tau_v(\phi) \deg_K v$, and for $x \in K$ we let $\deg_K \operatorname{rad}_K x := \sum_{v(x) \neq 0} \deg_K v$ (the sums are over every place v of K).

Proof. Let $\phi_T(X) = TX + a_1 X^q + \dots + a_r X^{q^r}$, where $a_i \in \mathbb{O}$. Let

$$f(X) = \phi_a(X) = b_0 X + b_1 X^q + \dots + b_{rn} X^{q^{rn}} = b_{rn} \prod_{i=1}^{q^{rn}} (X - \alpha_i),$$

where $b_0 = a$, $b_{rn} = a_r^{(q^{rn}-1)/(q^r-1)}$, and $n = \deg_K a = \deg_K \mathfrak{L}$. Let α be any one of the α_i .

Let \wp be a finite place of K with corresponding discrete valuation v_\wp , and let

$$\tau_\wp = \inf \left\{ \frac{v_\wp(a_i)}{q^i - 1}, i = 1, \dots, r. \right\}$$

Note that $\tau_\wp \geq 0$. Let K_\wp be the completion of K at \wp , and K'_\wp/K_\wp a totally tamely ramified extension with ramification index $1/(q^{rn} - 1)$ and ring of integers \mathbb{O}'_\wp .

Over K'_\wp , ϕ_T is isomorphic to a Drinfeld A -module

$$\phi'_T(X) = TX + a'_1 X^q + \dots + a'_r X^{q^r},$$

where $a'_i = a_i/\lambda^{q^i-1}$, $v_\wp(a'_i) \geq 0$ for $1 \leq i \leq r$, $v_\wp(\lambda) = \tau_\wp$, and $\lambda \in K'_\wp$.

Let $\phi'_a(X) = b'_0 X + b'_1 X^q + \dots + b'_{rn} X^{q^{rn}}$. As $b'_i = b_i/\lambda^{q^i-1}$, we have

$$v_\wp(b_i) \geq (q^i - 1)v_\wp(\lambda) = (q^i - 1)\tau_\wp.$$

From the Newton polygon of $f(X)$, we have

$$v_{\wp}(\alpha) \geq -\frac{v_{\wp}(a_r) \frac{q^{rn} - 1}{q^r - 1} - (q^{rn-1} - 1)\tau_{\wp}}{q^{rn} - q^{rn-1}} =: -\delta.$$

Pick a $\mu \in K'_{\wp}$ such that $v_{\wp}(\mu) = \delta + \epsilon$, where $0 \leq \epsilon < \frac{1}{q^{rn} - 1}$. Now

$$f(X/\mu) = b_{rn}/\mu^{q^{rn}} \prod_{i=1}^{q^{rn}} (X - \mu\alpha_i),$$

and we know that $g(X) = \prod_i (X - \mu\alpha_i)$ is monic and lies in $\mathbb{O}'_{\wp}[X]$, where \mathbb{O}'_{\wp} is the ring of integers of K'_{\wp} . Thus, $g'(X) = \mu^{q^{rn}-1}a/b_{rn}$. Hence,

$$\begin{aligned} v_{\wp}(g'(\mu\alpha)) &= v_{\wp}(\mu)(q^{rn} - 1) + v_{\wp}(a) - v_{\wp}(b_{rn}) \\ &\leq \delta(q^{rn} - 1) + 1 + v_{\wp}(a) - v_{\wp}(a_r) \frac{q^{rn} - 1}{q^r - 1} \\ &\leq v_{\wp}(a_r) \frac{q^{rn} - 1}{q^r - 1} \left(\frac{q^{rn} - 1}{q^{rn} - q^{rn-1}} - 1 \right) - \frac{(q^{rn-1} - 1)(q^{rn} - 1)}{q^{rn} - q^{rn-1}} \tau_{\wp} + 1 + v_{\wp}(a) \\ &\leq v_{\wp}(a_r) \frac{q^{rn} - 1}{q^r - 1} \cdot \frac{1 - q^{1-rn}}{q - 1} - \frac{q^{2rn-1} - q^{rn} - q^{rn-1} + 1}{q^{rn} - q^{rn-1}} \tau_{\wp} + 1 + v_{\wp}(a) \\ &= v_{\wp}(a_r) \frac{q^{rn} - 1}{(q^r - 1)(q - 1)} - \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \tau_{\wp} + 1 + v_{\wp}(a). \end{aligned}$$

It follows that

$$v_{\wp}(D(K'_{\wp}(\mu\alpha)/K'_{\wp})) \leq v_{\wp}(a_r) \frac{q^{rn} - 1}{(q^r - 1)(q - 1)} - \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \tau_{\wp} + 1 + v_{\wp}(a)$$

and

$$\begin{aligned} v_{\wp}(D(K_{\wp}(\alpha)/K_{\wp})) &\leq v_{\wp}(D(K'_{\wp}(\mu\alpha)/K'_{\wp})) + v_{\wp}(D(K'_{\wp}/K_{\wp})) \\ &\leq v_{\wp}(a_r) \frac{q^{rn} - 1}{(q^r - 1)(q - 1)} - \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \tau_{\wp} + 2 + v_{\wp}(a). \end{aligned}$$

Since $\tau_{\wp} \leq v_{\wp}(a_r)/(q^r - 1)$, we have

$$\begin{aligned} v_{\wp}(a_r) \frac{q^{rn} - 1}{(q^r - 1)(q - 1)} - \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \tau_{\wp} + 2 + v_{\wp}(a) &\geq v_{\wp}(a_r) \frac{q^{rn} - 1}{(q^r - 1)(q - 1)} - \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \frac{v_{\wp}(a_r)}{q^r - 1} + 2 + v_{\wp}(a) \\ &= v_{\wp}(a_r) \frac{q - q^{1-rn}}{(q^r - 1)(q - 1)} + 2 + v_{\wp}(a) \geq 2. \end{aligned}$$

From [Proposition 4.1](#), we know that for a finite place v_\wp of K , $v_\wp(D(K(\alpha)/K)) = 0$ if $v_\wp(aa_r) = 0$. It follows that

$$(18) \quad v_\wp(D(K_\wp(\alpha)/K_\wp)) \\ \leq v_\wp(a_r) \frac{q^{rn} - 1}{(q^r - 1)(q - 1)} - \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \tau_\wp + 2\nu + v_\wp(a),$$

where $\nu = 1$ if $v_\wp(aa_r) > 0$ and $\nu = 0$ if $v_\wp(aa_r) = 0$.

Let $\tilde{\infty} \in S_\infty^K$ be an infinite prime of K with corresponding valuation $v_{\tilde{\infty}}$, and let K'_∞/K_∞ be a totally tamely ramified extension with ramification index $1/(q^{rn} - 1)$ and ring of integers $\mathbb{O}'_{\tilde{\infty}}$.

Let

$$\tau_{\tilde{\infty}}(\phi) = \inf \left\{ \frac{v_{\tilde{\infty}}(a_i)}{q^i - 1}, i = 1, \dots, r \right\}$$

Note that $\tau_{\tilde{\infty}} \leq 0$.

Over K'_∞ , ϕ_T is isomorphic to a Drinfeld A -module

$$\phi'_T(X) = TX + a'_1 X^q + \dots + a'_r X^{q^r},$$

where $a'_i = a_i/\lambda^{q^i - 1}$, $v_{\tilde{\infty}}(a'_i) \geq 0$ for $1 \leq i \leq r$, $v_{\tilde{\infty}}(\lambda) = \tau_{\tilde{\infty}}$, and $\lambda \in K'_\infty$.

Let $\phi'_a(X) = b'_0 X + b'_1 X^q + \dots + b'_{rn} X^{q^{rn}}$. Set

$$\omega_1 = \max \left\{ e, -\frac{v_{\tilde{\infty}}(a'_i)}{q^i} : i = 1, \dots, r \right\} = 1.$$

From [Lemma 4.2](#), we know that

$$\omega_n = n\omega_1 \geq \max \left\{ ne, -\frac{v_{\tilde{\infty}}(b'_i)}{q^i} : i = 1, \dots, rn \right\}.$$

Thus, $v_{\tilde{\infty}}(b'_i) \geq -q^i ne$ for $i = 1, \dots, rn$. As $b'_i = b_i/\lambda^{q^i - 1}$, we have

$$v_{\tilde{\infty}}(b_i) \geq -q^i ne + (q^i - 1)v_{\tilde{\infty}}(\lambda) = -q^i ne + (q^i - 1)\tau_{\tilde{\infty}}.$$

From the Newton polygon of $f(X)$, it follows that

$$v_{\tilde{\infty}}(\alpha) \geq -\frac{v_{\tilde{\infty}}(a_r) \frac{q^{rn} - 1}{q^r - 1} + neq^{rn-1} - (q^{rn-1} - 1)\tau_{\tilde{\infty}}}{q^{rn} - q^{rn-1}} =: -\delta_{\tilde{\infty}}.$$

Let $\mu_{\tilde{\infty}}$ be such that $v_{\tilde{\infty}}(\mu_{\tilde{\infty}}) = \delta_{\tilde{\infty}} + \epsilon_{\tilde{\infty}}$, where $0 \leq \epsilon_{\tilde{\infty}} < 1/(q^{rn} - 1)$. Now

$$f(X/\mu_{\tilde{\infty}}) = b_{rn}/\mu_{\tilde{\infty}}^{q^{rn}} \prod_{i=1}^{q^{rn}} (X - \mu_{\tilde{\infty}} \alpha_i),$$

and we know that $g(X) = \prod_{i=1}^{q^{rn}} (X - \mu_{\infty} \alpha_i)$ is monic and lies in $\mathbb{O}'_{\infty}[X]$, where \mathbb{O}'_{∞} is the ring of integers of K'_{∞} . Thus, $g'(X) = \mu_{\infty}^{q^{rn}-1} a/b_{rn}$. Hence,

$$\begin{aligned}
& v_{\infty}(g'(\mu_{\infty} \alpha)) \\
&= v_{\infty}(\mu_{\infty})(q^{rn} - 1) + v_{\infty}(a) - v_{\infty}(b_{rn}) \\
&\leq \delta_{\infty}(q^{rn} - 1) + 1 + v_{\infty}(a) - v_{\infty}(a_r) \frac{q^{rn} - 1}{q^r - 1} \\
&\leq v_{\infty}(a_r) \frac{q^{rn} - 1}{q^r - 1} \left(\frac{q^{rn} - 1}{q^{rn} - q^{rn-1}} - 1 \right) \\
&\quad + ne \frac{q^{rn} - 1}{q - 1} - \frac{(q^{rn-1} - 1)(q^{rn} - 1)}{q^{rn} - q^{rn-1}} \tau_{\infty} + 1 + v_{\infty}(a) \\
&= v_{\infty}(a_r) \frac{q^{rn} - 1}{q^r - 1} \cdot \frac{1 - q^{1-rn}}{q - 1} + ne \frac{q^{rn} - 1}{q - 1} - \frac{q^{2rn-1} - q^{rn} - q^{rn-1} + 1}{q^{rn} - q^{rn-1}} \tau_{\infty} + 1 + v_{\infty}(a) \\
&= v_{\infty}(a_r) \frac{q^{rn} - 1}{(q^r - 1)(q - 1)} + ne \frac{q^{rn} - 1}{q - 1} - \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \tau_{\infty} + 1 + v_{\infty}(a).
\end{aligned}$$

It follows that

$$\begin{aligned}
(19) \quad v_{\infty}(D(K_{\infty}(\alpha)/K_{\infty})) &\leq v_{\infty}(a_r) \frac{q^{rn} - 1}{(q^r - 1)(q - 1)} + ne \frac{q^{rn} - 1}{q - 1} \\
&\quad - \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \tau_{\infty} + 2 + v_{\infty}(a).
\end{aligned}$$

Let $\mathfrak{D}(K(\alpha)/K)$ be the different divisor of $K(\alpha)$ over K , and Ω_P the set of conjugates of α over K_P . Using (18) and (19), we obtain

$$\begin{aligned}
\deg_K \mathfrak{D}(K(\alpha)/K) &= \sum_P \max\{v_P(D(K_P(\alpha)/K_P)) : \alpha \in \Omega_P\} \deg_K P \\
&\leq n \frac{q^{rn} - 1}{q - 1} \sum_{\infty \in S_{\infty}^K} e(\infty/\infty) \deg_K \infty \\
&\quad - \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \sum_v \tau_P \deg_K P + 2 \deg_K \text{rad}_K aa_r \\
&= n \frac{q^{rn} - 1}{q - 1} \sum_{\infty \in S_{\infty}^K} e(\infty/\infty) \frac{f(\infty/\infty)}{[\mathbb{F}_K : \mathbb{F}_F]} \deg_F \infty \\
&\quad - \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \sum_v \tau_P \deg_K P + 2 \deg_K \text{rad}_K aa_r \\
&\leq n \frac{q^{rn} - 1}{q - 1} s - \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \sum_v \tau_P \deg_K P + 2 \deg_K \text{rad}_K aa_r,
\end{aligned}$$

where the summation runs through all the primes P of K , s is the geometric extension degree of K/F , and we use the fact that $\sum_P v_P(x) \deg_K P = 0$ for $x \in K$. Remark that $\sum_P \tau_P \deg_K P \leq 0$; so we finally get

$$\begin{aligned}
& \deg_K \mathfrak{D}(K(\alpha)/K) \\
& \leq ns \frac{q^{rn} - 1}{q - 1} + \frac{q^{rn} - q - 1 + q^{1-rn}}{q - 1} \left(- \sum_v \tau_P \deg_K P \right) + 2 \deg_K \text{rad}_K aa_r \\
& \leq ns \frac{q^{rn} - 1}{q - 1} + \frac{q^{rn} - 1}{q - 1} \left(- \sum_P \tau_P \deg_K P \right) + 2 \deg_K \text{rad}_K aa_r \\
& \leq \frac{q^{rn} - 1}{q - 1} \left(ns - \sum_v \tau_P \deg_K P \right) + 2 \deg_K \text{rad}_K aa_r \\
& \leq \frac{\ell^r - 1}{q - 1} (ns + \Lambda(\phi)) + 2 \deg_K \text{rad}_K aa_r \\
& \leq \frac{\ell^r - 1}{q - 1} (s \deg_K a + \Lambda(\phi)) + 2 \deg_K \text{rad}_K a \Delta(\phi).
\end{aligned}$$

Using transitivity of the different (see [Lemma 2.3](#)) and the fact that $K_{\phi, \mathfrak{L}}$ is generated by r of the roots α_i , the result follows. \square

Corollary 4.4. *Assume the notation of [Proposition 4.3](#). Let ϕ_1 and ϕ_2 be rank- r Drinfeld A -modules that are integral over \mathbb{O} . Let $\mathfrak{D}(\tilde{K}/K)$ be the different divisor of \tilde{K}/K , where $\tilde{K} = K_{\phi_1, \mathfrak{L}} K_{\phi_2, \mathfrak{L}}$. Then we have*

$$\deg_K \mathfrak{D}(\tilde{K}/K) \leq r \left(\frac{\ell^r - 1}{q - 1} (2s \deg_K a + \Lambda(\phi_1, \phi_2)) + 2D(\phi_1, \phi_2) + 4 \deg_K a \right),$$

where $\Lambda(\phi_1, \phi_2) = \Lambda(\phi_1) + \Lambda(\phi_2)$.

Proof. The result follows from [Lemma 2.6](#) and [Proposition 4.3](#). \square

5. Proof of [Theorem 1.2](#)

We first recall some intermediate results, which are function field analogues of those found in [\[Serre 1981\]](#) (see [\[Gardeyn 2002\]](#)).

Lemma 5.1. *We have*

$$\sum_{1 \leq \deg_F \mathfrak{L} \leq N} \deg_F \mathfrak{L} \geq q^N$$

for all positive integers N , where the sum is over finite primes \mathfrak{L} of F .

Proof. The product of all finite primes \mathfrak{L} of F such that $\deg \mathfrak{L}$ divides N is equal to $T^{q^N} - T$, so the inequality follows. \square

Lemma 5.2. *For any nonzero $n \in A$, there exists a finite prime \mathfrak{L} of F such that $n \not\equiv 0 \pmod{\mathfrak{L}}$ with $\deg_F \mathfrak{L} \leq 1 + \log_q \deg_F n$.*

Proof. Suppose $n \equiv 0 \pmod{\mathfrak{L}}$ for all the primes \mathfrak{L} such that

$$1 \leq \deg_F \mathfrak{L} \leq 1 + \log_q \deg_F n.$$

Choose $k := \lfloor 1 + \log_q \deg_F n \rfloor$, so that $k - 1 \leq \log_q \deg_F n < k$, and hence $q^{k-1} \leq \deg_F n < q^k$.

Then $\prod_{1 \leq \deg_F \mathfrak{L} \leq k} \mathfrak{L}$ divides n , so $q^k \leq \deg_F n$, by [Lemma 5.1](#). But $\deg_F n < q^k$, which is a contradiction. \square

For the proof of [Theorem 1.2](#), we will require an estimate of the form

$$(20) \quad \gamma x^t \leq \frac{x}{1 + \log_q x},$$

for $x \geq C$.

Lemma 5.3. *Let $c^* \geq 1$ be given and set $t^* = 1 - 1/\ln(qc^*)$. Then we have*

$$(21) \quad \gamma x^{t^*} \leq \frac{x}{1 + \log_q x}$$

for $x \geq c^*$, where

$$\gamma = \frac{(c^*)^{1-t^*}}{1 + \log_q c^*} = \frac{(c^*)^1 / \ln(qc^*)}{1 + \log_q c^*}.$$

Proof. The inequality

$$\gamma x^t \leq \frac{x}{1 + \log_q x}$$

is equivalent to

$$f(x, t) = \frac{x^{1-t}}{1 + \log_q x} \geq \gamma.$$

For a fixed t , taking the derivative of f with respect to x ,

$$f'(x, t) = x^{-t} \left((1-t)(1 + \log_q x) - \frac{1}{\ln q} \right) / *^2,$$

where $* = (1 + \log_q x)$. Hence, $f'(x, t) \geq 0$ is equivalent to

$$(1-t)(1 + \log_q x) - \frac{1}{\ln q} \geq 0,$$

or equivalently,

$$(22) \quad (1-t)(\ln q + \ln x) \geq 1.$$

Assuming $t < 1$, (22) is equivalent to

$$x \geq \frac{e^{1/(1-t)}}{q} =: \beta(t).$$

Thus, for a fixed $t < 1$, $f(x, t)$ is increasing with respect to x , when $x \geq \beta(t)$; that is, $f(x, t) \geq f(\beta(t), t)$ if $x \geq \beta(t)$. Now, $\beta(t^*) = c^*$ and $t^* < 1$, so we obtain

$$x^{t^*} f(c^*, t^*) \leq \frac{x}{1 + \log_q x},$$

for $x \geq c^*$. □

Lemma 5.4.

$$(23) \quad \log_q(x + y) \leq \max\{\log_q(2x), \log_q(2y)\},$$

$$(24) \quad \log_q(x + y) \leq \log_q x + \log_q y \quad \text{if } x, y \geq 2.$$

Proof. In order to have $z \geq \log_q(x + y)$, it suffices to have

$$\frac{1}{2}q^z \geq x \quad \text{and} \quad \frac{1}{2}q^z \geq y,$$

which is equivalent to

$$z \geq \log_q(2x) \quad \text{and} \quad z \geq \log_q(2y).$$

Thus, taking $z = \max\{\log_q(2x), \log_q(2y)\}$, we have

$$\log_q(x + y) \leq \max\{\log_q(2x), \log_q(2y)\}. \quad \square$$

Conclusion of the proof of Theorem 1.2. Let $\wp \notin S$ be a prime of K with least degree such that $P_\wp(\phi_1) \neq P_\wp(\phi_2)$, where S is the given finite set of primes of K outside of which both ϕ_1 and ϕ_2 have good reduction. Let α_0 be a nonzero coefficient of $P_\wp(\phi_1) - P_\wp(\phi_2)$.

It is known that a root γ of $P_\wp(\phi_1)$ or $P_\wp(\phi_2)$ satisfies

$$v_\infty(\gamma) = -\frac{1}{r} \deg_K \wp$$

(see [Goss 1992, Theorem 3.2.3(c)(d); Gardeyn 2002, Proposition 9]). This implies that each coefficient β of $P_\wp(\phi_1)$ and $P_\wp(\phi_2)$ satisfies $\deg_F \beta \leq \deg_K \wp$, and hence each coefficient α of $P_\wp(\phi_1) - P_\wp(\phi_2)$ also satisfies $\deg_F \alpha \leq \deg_K \wp$; in particular $\deg_F \alpha_0 \leq \deg_K \wp$.

We choose a finite prime \mathfrak{L} of F by Lemma 5.2 such that

$$(25) \quad \alpha_0 \not\equiv 0 \pmod{\mathfrak{L}} \quad \text{and} \quad \deg_F \mathfrak{L} \leq 1 + \log_q \deg_K \wp,$$

and write $\mathfrak{L} = (a)$, where a is monic in A .

Suppose \wp lies above the prime \mathfrak{p} of F . For $x \geq 7$, we have $\log_q x < \frac{1}{2}(x-1)$ (since if we let $f(x) = \frac{1}{2}(x-1) - \log_q x$, then $f'(x) > 0$ for $x \geq 7$ and $f(7) > 0$). Hence, we obtain that $x < q^{(1/2)(x-1)}$, so $q^{(1/2)(x-1)}/x > 1$; hence, $q^{x-1}/x > q^{(1/2)(x-1)}$ for $x \geq 7$. Thus, noting that

$$s \geq \frac{f(\wp/\mathfrak{p})}{[\mathbb{F}_K : \mathbb{F}_F]},$$

if $x \geq \max\{1 + 2 \log_q s, 7\}$, we get that

$$\frac{q^{x-1}}{x} > q^{(1/2)(x-1)} \geq s \geq \frac{f(\wp/\mathfrak{p})}{[\mathbb{F}_K : \mathbb{F}_F]}.$$

But then if $\mathfrak{L} = \wp$, we would have

$$\deg_F \mathfrak{p} \leq 1 + \log_q \deg_K \wp = 1 + \log_q \frac{f(\wp/\mathfrak{p})}{[\mathbb{F}_K : \mathbb{F}_F]} \deg_F \mathfrak{p};$$

in other words,

$$\frac{q^{x-1}}{x} \leq \frac{f(\wp/\mathfrak{p})}{[\mathbb{F}_K : \mathbb{F}_F]},$$

where $x = \deg_F \mathfrak{p} = \deg_F \mathfrak{L}$. Therefore, we either have that

$$\deg_F \mathfrak{p} \leq \max\{1 + 2 \log_q s, 7\},$$

or $\mathfrak{L} \neq \mathfrak{p}$ by the above inequality. In the former case, it follows that $\deg_K \wp \leq s \max\{1 + 2 \log_q s, 7\}$.

Suppose we are now in the latter case, where $\mathfrak{L} \neq \mathfrak{p}$. Consider the representation

$$\psi_{\mathfrak{L}} : G_K \rightarrow \text{Aut}_{A/\mathfrak{L}}(\phi_1[\mathfrak{L}]) \times \text{Aut}_{A/\mathfrak{L}}(\phi_2[\mathfrak{L}]) \cong \text{GL}_r(A/\mathfrak{L}) \times \text{GL}_r(A/\mathfrak{L}),$$

where $\psi_{\mathfrak{L}} = \rho_{\phi_1, \mathfrak{L}} \times \rho_{\phi_2, \mathfrak{L}}$. Let $G_{\mathfrak{L}}$ be the image of this homomorphism. Let $C_{\mathfrak{L}}$ be the subset of $G_{\mathfrak{L}}$ consisting of pairs (a, b) such that the characteristic polynomials of a and b are not equal. Note that $C_{\mathfrak{L}}$ is invariant under conjugation, so it is a union of conjugacy classes in $G_{\mathfrak{L}}$. Since $\mathfrak{L} \neq \mathfrak{p}$, we have $C_{\mathfrak{L}} \neq \emptyset$; in particular, there is some conjugacy class $\mathcal{C} \subseteq C_{\mathfrak{L}}$ in $G_{\mathfrak{L}}$ with $\mathcal{C} \neq \emptyset$.

Let $S_{\mathfrak{L}} = S \cup \{\text{primes } \mathfrak{l} \text{ of } K \text{ lying over } \mathfrak{L}\}$. Then the Galois representation $\psi_{\mathfrak{L}}$ is unramified outside $S_{\mathfrak{L}}$. We have $A/\mathfrak{L} \cong \mathbb{F}_{\ell}$, where $\ell = q^{\deg_F \mathfrak{L}}$.

Let \tilde{K}/K be the field extension associated to $\psi_{\mathfrak{L}}$, and let n (resp. n') be its degree (resp. geometric extension degree). Applying [Corollary 3.4](#) to \tilde{K}/K , and using [Lemma 2.2](#) together with the bound for the degree with respect to K of $\mathfrak{D} = \mathfrak{D}(\tilde{K}/K)$ given in [Corollary 4.4](#), we deduce that there is a prime $P \notin S_{\mathfrak{L}}$ such that $\text{Frob}_P = \mathcal{C} \subseteq C_{\mathfrak{L}}$ and

$$\deg_K P \leq \frac{4}{m_0} \log_q \frac{4}{3} (B + 3g + 3) + m,$$

where

$$\Sigma' = \sum_{\mathfrak{p} \in \mathcal{S}_\Sigma} \mathfrak{p} \geq \Sigma = \sum_{\mathfrak{p} \in \mathcal{S}} \mathfrak{p}, \quad m = [\mathbb{F}_{\tilde{K}} : \mathbb{F}_K], \quad m_0 = [\mathbb{F}_K : \mathbb{F}_F],$$

$$\deg_K \Sigma' = \deg_K \text{rad}_K \Delta(\phi_1) \Delta(\phi_2) + \deg_K \mathfrak{L},$$

$$B = \max\{\deg_K \Sigma', \deg_{\tilde{K}} \mathfrak{D}, 2|\text{Gal}(E/K')| - 2, 2\},$$

$$\deg_{\tilde{K}} \mathfrak{D} \leq rn' \left(\frac{\ell^r - 1}{q - 1} (2s \deg_K a + \Lambda(\phi_1, \phi_2)) + 2D(\phi_1, \phi_2) + 4 \deg_K a \right).$$

Then

$$(26) \quad \deg_K P \leq \frac{4}{m_0} \log_q \frac{4}{3} (B + 3g + 3) + m \leq \frac{4}{m_0} (\log_q \frac{4}{3} B + \log_q 4(g + 1)) + m,$$

using $B \geq 2$ and [Lemma 5.4](#). Note that regarding B , the terms $\deg_K \Sigma'$ and $2|\text{Gal}(E/K')| - 2$ are less than the bound we use for $\deg_{\tilde{K}} \mathfrak{D}$, so we can ignore them later on when we bound B .

Using [Lemma 5.4](#), we obtain

$$\begin{aligned} \log_q \deg_{\tilde{K}} \mathfrak{D} &= \log_q rn' \left(\frac{\ell^r - 1}{q - 1} \Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2) + \left(2s \frac{\ell^r - 1}{q - 1} + 4 \right) \deg_K a \right) \\ &\leq \log_q rn' + \log_q \left(\frac{\ell^r - 1}{q - 1} (\Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2)) + \left(2s \frac{\ell^r - 1}{q - 1} + 4 \right) \deg_K a \right) \\ &\leq \log_q rn' + \max\{V_1, V_2\}, \end{aligned}$$

where

$$\begin{aligned} V_1 &:= \log_q 2 \frac{\ell^r - 1}{q - 1} (\Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2)) \\ &= \log_q 2 + \log_q \frac{\ell^r - 1}{q - 1} + \log_q (\Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2)), \\ V_2 &:= \log_q 2 \left(2s \frac{\ell^r - 1}{q - 1} + 4 \right) \deg_K a \\ &\leq \log_q 2 + \log_q 8s + \log_q \frac{\ell^r - 1}{q - 1} + \log_q \deg_K a \leq V_1 + \log_q 8s + \log_q \deg_K a. \end{aligned}$$

Thus,

$$\begin{aligned} \log_q B &\leq \log_q rn' + V_1 + \log_q 8s + \log_q \deg_K a \\ &= \log_q rn' + \log_q 16s + \log_q \frac{\ell^r - 1}{q - 1} + \log_q \deg_K a + \log_q (\Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2)). \end{aligned}$$

Since $n' \leq n = |G_\Sigma| < \ell^{2r^2}$, $\log_q \ell = \deg_F \mathfrak{L} = \deg_F a$, and $\deg_K a \leq s \deg_F a = s \log_q \ell$, we finally obtain

$$(27) \quad \log_q B \leq \log_q 16rs^2 + (2r^2 + r) \log_q \ell + \log_q \log_q \ell + \log_q (\Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2)).$$

Note that if $\log_q (\Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2)) = 0$, the derivation of the bound (27) above can be modified so as to obtain

$$(28) \quad \log_q B \leq \log_q 16rs^2 + (2r^2 + r) \log_q \ell + \log_q \log_q \ell.$$

Thus, we have

$$(29) \quad \log_q \frac{4}{3} B \leq \log_q \frac{64}{3} rs^2 + (2r^2 + r + 1) \log_q \ell + \log_q^* (\Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2)).$$

Returning to (26), we obtain

$$(30) \quad \deg_K P \leq \frac{4}{m_0} (\log_q 86rs^2(g+1) + (2r^2 + r + 1) \log_q \ell + \log_q^* (\Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2))) + m.$$

By construction of $C_{\mathcal{L}}$, we have $P_P(\phi_1) \not\equiv P_P(\phi_2) \pmod{\mathcal{L}}$. Thus, $\deg_K \wp \leq \deg_K P$, and from (25), it follows that

$$(31) \quad \begin{aligned} \deg_K \wp &\leq \frac{4}{m_0} (\log_q 86rs^2(g+1) + (2r^2 + r + 1) \log_q \ell + \log_q^* (\Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2))) + m \\ &\leq \frac{4}{m_0} (\log_q 86rs^2(g+1) + (2r^2 + r + 1)(1 + \log_q \deg_K \wp) + \log_q^* (\Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2))) + m. \end{aligned}$$

As $1 + \log_q x \geq 1$, $\frac{\log_q x}{x} \leq 1$, we have

$$\frac{\deg_K \wp}{1 + \log_q (\deg_K \wp)} \leq \frac{4}{m_0} (d_r + W),$$

where $c_r = 2r^2 + r + 1$, $d_r := c_r + \log_q 86rs^2(g+1)$, and

$$W := \log_q^* (\Lambda(\phi_1, \phi_2) + 2D(\phi_1, \phi_2)) + mm_0.$$

If $x \geq d_r$, then using Lemma 5.3 with $c^* = d_r$ and $x = \deg_K \wp$, we obtain

$$\gamma x^{t^*} \leq \frac{x}{1 + \log_q x} \leq \frac{4}{m_0} (d_r + W),$$

where γ is as in Lemma 5.3. This implies that

$$x^{t^*} \leq \frac{4}{m_0} \frac{(d_r + W)}{\gamma},$$

so that

$$\begin{aligned}
 (32) \quad \log_q \deg_K \wp &= \log_q x \leq \frac{1}{t^*} \log_q \frac{4}{m_0} (d_r + W) \cdot \frac{1 + \log_q d_r}{(d_r)^{1/\ln(qd_r)}} \\
 &\leq s^* \left(\log_q \frac{4}{m_0} + \log_q (d_r + W) + \log_q (1 + \log_q d_r) - \frac{1}{\ln(qd_r)} \log_q d_r \right) \\
 &\leq s^* \left(\log_q \frac{4}{m_0} + \log_q d_r + \log_q W + \log_q (1 + \log_q d_r) - \frac{1}{\ln(qd_r)} \log_q d_r \right) \\
 &\leq s^* \left(\log_q \frac{4}{m_0} + \log_q W + \log_q \log_q d_r \right) + \log_q d_r,
 \end{aligned}$$

using $d_r, W \geq 2$, and where

$$t^* = \frac{\ln(qd_r) - 1}{\ln(qd_r)} \quad \text{and} \quad s^* = s_{q,r}^* = \frac{1}{t^*} = \frac{\ln(qd_r)}{\ln(qd_r) - 1}.$$

We note that when q or r is large, $s_{q,r}^*$ tends to 1 from above.

Substitution of (32) into (31) yields

$$\begin{aligned}
 (33) \quad \frac{1}{4} \deg_K \wp &\leq \log_q 86rs^2(g+1) + c_r(1 + \log_q \deg_K \wp) + W \\
 &\leq \log_q 86rs^2(g+1) \\
 &\quad + c_r \left(1 + s^* \left(\log_q \frac{4}{m_0} + \log_q W + \log_q \log_q d_r \right) + \log_q d_r \right) + W \\
 &= \log_q 86rs^2(g+1) + c_r \left(1 + s^* \log_q \frac{4}{m_0} + \log_q d_r \right) \\
 &\quad + c_r s^* \log_q \log_q d_r + W + c_r s^* \log_q W \\
 &= C_{q,r} + W + c_r s_{q,r}^* \log_q W,
 \end{aligned}$$

where

$$C_{q,r} = \log_q 86rs^2(g+1) + c_r \left(1 + s_{q,r}^* \log_q \frac{4}{m_0} + \log_q d_r \right) + c_r s_{q,r}^* \log_q \log_q d_r.$$

Therefore, we either have the above upper bound (33) on $\deg_K \wp$ or $\deg_K \wp \leq d_r \leq C_{q,r}$, so in the end we get

$$(34) \quad \deg_K \wp \leq \frac{4}{m_0} (C_{q,r} + W + c_r s_{q,r} \log_q W).$$

Finally, we note from the discussion in the introduction that $m \leq g_{\phi_1} g_{\phi_2}$.

6. The case of rank 2

In this section, we consider the case of rank 2 and $K = F$, and explain how to make all the terms explicit in our isogeny theorem.

For a Drinfeld A -module ϕ of rank 2 over $K = F = \mathbb{F}_q(T)$, the successive minima of the lattices associated to the uniformizations of ϕ are determined in [Chen and Lee 2013], and this is used to obtain an explicit bound for the valuation $v_\infty(D(K_\infty(\phi[a])/K_\infty))$ of the different of $K_{\phi,a} = K(\phi[a])$ over K at the infinite prime ∞ of K and $v_p(D(K_p(\phi[a])/K_p))$ at a finite prime p of K , following the work of Goss [1996].

The infinite prime case is obtained using the explicit information about the Newton polygon of the exponential map $e_{\phi,\infty}$ attached to ϕ from its uniformization over C_∞ .

Assume the same notation as in the proof and statement of Proposition 4.3, taking $K = F = \mathbb{F}_q(T)$ and $\tilde{\infty} = \infty$; the explicit bounds given in [Chen and Lee 2013] are as follows.

Let $\phi_T = T + a_1\tau + a_2\tau^2$ and $j(\phi) = a_1^{q+1}/a_2$, and let m be the least positive integer such that $-v_\infty(j(\phi)) \leq q^{m+1}$. Then we have

$$v_\infty(D(K_\infty(\phi[a])/K_\infty)) \leq \begin{cases} 1 & \text{if } -v_\infty(j(\phi)) \leq q, \\ 1 + \kappa(q^{\kappa+1} - 1) & \text{if } q < -v_\infty(j(\phi)) \leq q^{m+1}, \end{cases}$$

where

$$\kappa = \frac{-v_\infty(j(\phi)) - q^m}{q^m(q - 1)} + m - 1,$$

and

$$v_p(D(K_p(\phi[a])/K_p)) \leq \begin{cases} 2v_p(a) & \text{if } \phi \text{ has good reduction} \\ & \text{over } K_p, \\ 2v_p(a) + 1 & \text{if } v_p(j(\phi)) \geq 0 \text{ and } \phi \text{ has} \\ & \text{bad reduction over } K_p, \\ 2v_p(a) + 1 - \frac{2}{q-1}v_p(j(\phi)) & \text{if } v_p(j(\phi)) < 0. \end{cases}$$

Putting this together yields the following explicit bound on the different divisor of $F(\phi[a])/F$ when ϕ has rank 2, which can be used in place of the more general bound that we use in this paper. See Section 7 for a comparison of the two bounds in the context of our application.

Theorem 6.1. *Let ϕ be a Drinfeld A -module of rank 2 over F , and $\mathfrak{D}(F(\phi[a])/F)$ the different divisor of $F(\phi[a])/F$. Then*

$$\deg_F \mathfrak{D}(F(\phi[a])/F) \leq 2 \deg_F a + \deg_F \eta + \frac{2}{q-1} \deg_F \delta + v_\infty(D(F_\infty(\phi[a])/F_\infty)),$$

where δ is the (monic) denominator of $j(\phi)$ as represented by a fraction in reduced form, and η is the product of finite primes p such that ϕ has bad reduction over F_p .

Concerning the term g_ϕ , we have from [Gardeyn 2002] that

$$g_\phi = g_{\phi, \infty} \leq (q^2 - 1)(q^2 - q)v_{2, \phi, \infty}/v_{1, \phi, \infty},$$

where $v_{i, \phi, \infty}$ is the i -th successive minimum of ϕ associated to its uniformization over C_∞ . In [Chen and Lee 2013], the $v_{i, \phi, \infty}$ are determined as follows.

Case 1: If $-v(j(\phi)) \leq q$, then $v_{1, \phi, \infty} = v_{2, \phi, \infty} = -s_1$.

Case 2: If $q < -v(j(\phi)) \leq q^{m+1}$, then $v_{1, \phi, \infty} = -s_1$, $v_{2, \phi, \infty} = -s_1 - \kappa$, where $s_1 = (v(a_2) + q^2)/(q^2 - 1)$ in Case 1 and $s_1 = (v(a_1) + q)/(q - 1)$ in Case 2, and m, κ are as above.

7. Comparison with work of Gardeyn

In this section we make some detailed comparisons with the work in [Gardeyn 2002], where an effective isogeny theorem is proven.

For the proof of our Theorem 1.2, an essential ingredient is the bound on the different divisor given in Proposition 4.3,

$$(35) \quad \deg_K \mathcal{D}(K_{\phi, \mathcal{L}}/K) \leq r \left(\frac{\ell^r - 1}{q - 1} (s \deg_K a + \Lambda(\phi)) + 2 \deg_K \text{rad}_K \Delta(\phi) + 2 \deg_K a \right),$$

where we recall that $\Lambda(\phi) = -\sum_v \tau_v(\phi) \deg_K v$. The counterpart of (35) in [Gardeyn 2002] is

$$(36) \quad \deg_K \mathcal{D}(K_{\phi, \mathcal{L}}/K) \leq r \deg_K a + \deg_K \Delta_\phi,$$

where Δ_ϕ is a divisor of K that is determined from the Newton polygons of the exponential functions associated to uniformizations of ϕ over C_\wp , where \wp is a prime of K .

Although there is a larger dependence on ℓ in our different bounds when we take degrees with respect to K , what is required in the application is the degree with respect to $K_{\phi, \mathcal{L}}$, which necessitates multiplying the degree with respect to K by $n' < \ell^{r^2}$. This means both bounds end up being comparable in their dependence on ℓ , as we later take the \log_q of this degree with respect to $K_{\phi, \mathcal{L}}$.

The quantity Δ_ϕ is more difficult to make explicit and compare, as we saw in Section 6, where its determination in the case of rank 2 and $K = F = \mathbb{F}_q(T)$ is recalled from [Chen and Lee 2013]. The method in [Chen and Lee 2013] yields the entire Newton polygon and uses Gekeler's theory of Drinfeld modular forms as well as Rosen's theory of formal Drinfeld modules. It may be possible to obtain weaker information using the more elementary approach of Chen and Lee [2012] in the infinite prime case, and to generalize Rosen's work to higher rank in the finite prime case, in such a way that Gardeyn's bounds can be made explicit.

As for the terms g_ϕ , it would seem that this also requires some knowledge relating to the successive minima of the lattices associated to the uniformization of ϕ over infinite primes.

Finally, two other places of difference are in our use of [Kumar Murty and Scherk 1994] for the Chebotarev density theorem instead of [Geyer and Jarden 1998], and in our analytic estimation methods, which differ slightly from [Gardeyn 2002; Serre 1981] because we have attempted to reduce the size of the constants in the different divisor bound, especially in front of the dominating terms.

Acknowledgements

We would like to thank A. Cojocaru and C. David for useful discussions pertaining to the subject of this paper.

References

- [Chen and Lee 2012] I. Chen and Y. Lee, “Coefficients of exponential functions attached to Drinfeld modules of rank 2”, *Manuscripta Math.* **139**:1-2 (2012), 123–136. MR 2959673 Zbl pre06084034
- [Chen and Lee 2013] I. Chen and Y. Lee, “Newton polygons, successive minima, and different bounds for Drinfeld modules of rank 2”, *Proc. Amer. Math. Soc.* **141**:1 (2013), 83–91. MR 2988712 Zbl 06146598
- [Cojocaru and David 2008] A. C. Cojocaru and C. David, “Frobenius fields for Drinfeld modules of rank 2”, *Compos. Math.* **144**:4 (2008), 827–848. MR 2009g:11071 Zbl 1158.11029
- [David 2001] C. David, “Frobenius distributions of Drinfeld modules of any rank”, *J. Number Theory* **90**:2 (2001), 329–340. MR 2002k:11084 Zbl 1012.11048
- [Gardeyn 2001] F. Gardeyn, *t-motives and Galois representations*, Ph.D. thesis, Universiteit Gent, 2001, <https://biblio.ugent.be/publication/522143>. MR 2715545
- [Gardeyn 2002] F. Gardeyn, “Une borne pour l’action de l’inertie sauvage sur la torsion d’un module de Drinfeld”, *Arch. Math. (Basel)* **79**:4 (2002), 241–251. MR 2003i:11073 Zbl 1019.11015
- [Geyer and Jarden 1998] W.-D. Geyer and M. Jarden, “Bounded realization of l -groups over global fields: the method of Scholz and Reichardt”, *Nagoya Math. J.* **150** (1998), 13–62. MR 99d:12001 Zbl 0906.12002
- [Goss 1992] D. Goss, “ L -series of t -motives and Drinfeld modules”, pp. 313–402 in *The arithmetic of function fields* (Columbus, OH, 1991), edited by D. Goss et al., Ohio State Univ. Math. Res. Inst. Publ. **2**, de Gruyter, Berlin, 1992. MR 94b:11048 Zbl 0806.11028
- [Goss 1996] D. Goss, *Basic structures of function field arithmetic*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **35**, Springer, Berlin, 1996. MR 97i:11062 Zbl 0874.11004
- [Kumar Murty and Scherk 1994] V. Kumar Murty and J. Scherk, “Effective versions of the Chebotarev density theorem for function fields”, *C. R. Acad. Sci. Paris Sér. I Math.* **319**:6 (1994), 523–528. MR 95j:11104 Zbl 0822.11077
- [Pink 1997] R. Pink, “The Mumford–Tate conjecture for Drinfeld-modules”, *Publ. Res. Inst. Math. Sci.* **33**:3 (1997), 393–425. MR 98f:11062 Zbl 0895.11025
- [Pink and Rütscche 2009a] R. Pink and E. Rütscche, “Adelic openness for Drinfeld modules in generic characteristic”, *J. Number Theory* **129**:4 (2009), 882–907. MR 2010f:11092 Zbl 05542339

- [Pink and Rüttsche 2009b] R. Pink and E. Rüttsche, “Image of the group ring of the Galois representation associated to Drinfeld modules”, *J. Number Theory* **129**:4 (2009), 866–881. [MR 2010f:11091](#) [Zbl 05542338](#)
- [Rosen 2002] M. Rosen, *Number theory in function fields*, Graduate Texts in Mathematics **210**, Springer, New York, 2002. [MR 2003d:11171](#) [Zbl 1043.11079](#)
- [Serre 1972] J.-P. Serre, “Propriétés Galoisiennes des points d’ordre fini des courbes elliptiques”, *Invent. Math.* **15**:4 (1972), 259–331. [MR 52 #8126](#) [Zbl 0235.14012](#)
- [Serre 1979] J.-P. Serre, *Local fields*, Graduate Texts in Mathematics **67**, Springer, New York, 1979. [MR 82e:12016](#) [Zbl 0423.12016](#)
- [Serre 1981] J.-P. Serre, “Quelques applications du théorème de densité de Chebotarev”, *Inst. Hautes Études Sci. Publ. Math.* **54** (1981), 323–401 (123–201). [MR 83k:12011](#) [Zbl 0496.12011](#)
- [Taguchi 1992] Y. Taguchi, “Ramifications arising from Drinfeld modules”, pp. 171–187 in *The arithmetic of function fields* (Columbus, OH, 1991), edited by D. Goss et al., Ohio State Univ. Math. Res. Inst. Publ. **2**, de Gruyter, Berlin, 1992. [MR 94b:11049](#) [Zbl 0801.11027](#)
- [Taguchi 1993] Y. Taguchi, “Semi-simplicity of the Galois representations attached to Drinfeld modules over fields of ‘infinite characteristics’”, *J. Number Theory* **44**:3 (1993), 292–314. [MR 94k:11064](#) [Zbl 0781.11024](#)
- [Taguchi 1996] Y. Taguchi, “On ϕ -modules”, *J. Number Theory* **60**:1 (1996), 124–141. [MR 97e:11067](#) [Zbl 0862.11038](#)
- [Takahashi 1982] T. Takahashi, “Good reduction of elliptic modules”, *J. Math. Soc. Japan* **34**:3 (1982), 475–487. [MR 83h:12023](#) [Zbl 0476.14010](#)
- [Tamagawa 1994] A. Tamagawa, “Generalization of Anderson’s t -motives and Tate conjecture”, *Sūrikaiseikikenkyūsho Kōkyūroku* **884** (1994), 154–159. [MR 96g:11061](#) [Zbl 0900.11036](#)
- [Thakur 2004] D. S. Thakur, *Function field arithmetic*, World Scientific, River Edge, NJ, 2004. [MR 2005h:11115](#) [Zbl 1061.11001](#)

Received December 20, 2011.

IMIN CHEN
DEPARTMENT OF MATHEMATICS
SIMON FRASER UNIVERSITY
BURNABY, BRITISH COLUMBIA V5A 1S6
CANADA

ichen@math.sfu.ca

YOONJIN LEE
DEPARTMENT OF MATHEMATICS
EWhA WOMANS UNIVERSITY
11-1 DAEHYUN-DONG, SEODAEMUN-GU
SEOUL 120-750
SOUTH KOREA

yoonjinl@ewha.ac.kr

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Don Blasius
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Vijayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2013 is US \$400/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 263 No. 1 May 2013

Biharmonic hypersurfaces in complete Riemannian manifolds	1
LUIS J. ALÍAS, S. CAROLINA GARCÍA-MARTÍNEZ and MARCO RIGOLI	
Half-commutative orthogonal Hopf algebras	13
JULIEN BICHON and MICHEL DUBOIS-VIOLETTE	
Superdistributions, analytic and algebraic super Harish-Chandra pairs	29
CLAUDIO CARMELI and RITA FIORESI	
Orbifolds with signature $(0; k, k^{n-1}, k^n, k^n)$	53
ANGEL CAROCCA, RUBÉN A. HIDALGO and RUBÍ E. RODRÍGUEZ	
Explicit isogeny theorems for Drinfeld modules	87
IMIN CHEN and YOONJIN LEE	
Topological pressures for ϵ -stable and stable sets	117
XIANFENG MA and ERCAI CHEN	
Lipschitz and bilipschitz maps on Carnot groups	143
WILLIAM MEYERSON	
Geometric inequalities in Carnot groups	171
FRANCESCO PAOLO MONTEFALCONE	
Fixed points of endomorphisms of virtually free groups	207
PEDRO V. SILVA	
The sharp lower bound for the first positive eigenvalue of the Folland–Stein operator on a closed pseudohermitian $(2n + 1)$ -manifold	241
CHIN-TUNG WU	
Remark on “Maximal functions on the unit n -sphere” by Peter M. Knopf (1987)	253
HONG-QUAN LI	