

*Pacific
Journal of
Mathematics*

**QUASISYMMETRIC HOMEOMORPHISMS
ON REDUCIBLE CARNOT GROUPS**

XIANGDONG XIE

Volume 265 No. 1

September 2013

QUASISYMMETRIC HOMEOMORPHISMS ON REDUCIBLE CARNOT GROUPS

XIANGDONG XIE

We show that quasisymmetric homeomorphisms between (most) reducible Carnot groups are bilipschitz. This implies rigidity for quasi-isometries between certain negatively curved homogeneous manifolds. The proof uses Pansu's differentiability theorem for quasisymmetric homeomorphisms between Carnot groups.

1. Introduction

We study quasisymmetric homeomorphisms between reducible Carnot groups. The main result says that in most cases, the quasisymmetric homeomorphism must be bilipschitz.

A Carnot group is *reducible* if it is isomorphic to the direct product of two Carnot groups. Otherwise, a Carnot group is called *irreducible*. A reducible Carnot group G can be written as $G = G_0 \times G_1 \times \cdots \times G_m$, where G_0 is abelian (i.e., isomorphic to some \mathbb{R}^n), and G_j ($1 \leq j \leq m$) is nonabelian irreducible. Such a decomposition is not unique in general; see Example 2.1.

All Carnot groups in this paper are equipped with the Carnot metric (see Section 3).

Theorem 1.1. *Let $F : G \rightarrow G'$ be a quasisymmetric map between two Carnot groups. Suppose G is reducible and admits a direct product decomposition of irreducible Carnot groups where at least two of the factors are not isomorphic. Then F is bilipschitz.*

The same claim remains open in the case when G is isomorphic to a direct product $N \times \cdots \times N$, where N is nonabelian irreducible.

Quasisymmetric homeomorphisms between general metric spaces are quasiconformal. In the case of Carnot groups (and of Loewner spaces more generally), a map is quasisymmetric if and only if it is quasiconformal (see [Heinonen and Koskela 1998]).

Partially supported by NSF grant DMS-1265735.

MSC2010: 53C17, 53C23, 30L10.

Keywords: quasisymmetric homeomorphism, Carnot groups, Pansu differentiability.

Theorem 1.1 has consequences for the rigidity of quasi-isometries between certain negatively curved homogeneous manifolds. Recall that a quasi-isometry between two metric spaces is an almost isometry if it preserves distance up to an additive constant. A quasi-isometry between two negatively curved spaces induces a quasisymmetric homeomorphism between the ideal boundaries (of the negatively curved spaces), where the ideal boundaries are equipped with visual metrics. Conversely, under mild conditions on the negatively curved spaces, each quasisymmetric homeomorphism between the ideal boundaries is the boundary map of a quasi-isometry; see [Bonk and Schramm 2000]. Similarly, almost isometries between negatively curved spaces correspond to bilipschitz maps between the ideal boundaries [ibid.]. On the other hand, Carnot groups arise as the ideal boundary of certain negatively curved homogeneous manifolds (see below for more details). Hence a direct consequence of Theorem 1.1 is that each quasi-isometry between certain negatively curved homogeneous manifolds is an almost isometry.

Heintze [1974] characterized homogeneous manifolds with negative sectional curvature (HMNs): Each HMN is isometric to a simply connected solvable Lie group S equipped with a left invariant Riemannian metric, and furthermore $S = N \rtimes \mathbb{R}$ is a semidirect product of a nilpotent Lie group N with \mathbb{R} , where \mathbb{R} acts on N by expanding (and contracting) automorphisms; conversely, every semidirect product as above admits a left invariant Riemannian metric with negative sectional curvature (hence is an HMN). The ideal boundary of an HMN $S = N \rtimes \mathbb{R}$ can be naturally identified with (the one-point compactification of) N . On the other hand, each Carnot group N is a simply connected nilpotent Lie group having a one-parameter family of dilations (see Section 3 for more details). These dilations induce an action of \mathbb{R} on the Carnot group by expanding (and contracting) automorphisms, so there is an HMN $N \rtimes \mathbb{R}$ associated with each Carnot group N . It follows that each Carnot group can be identified with the ideal boundary of some HMN. Hence Theorem 1.1 implies that each quasi-isometry between these HMNs is an almost isometry.

We next make some comments about the proof of Theorem 1.1. A main step in the proof is to show that the quasisymmetric homeomorphism preserves a certain foliation. Then the arguments in [Shanmugalingam and Xie 2012] show that the quasisymmetric homeomorphism is bilipschitz. To show that the quasisymmetric homeomorphism preserves a foliation, one first proves that infinitesimally it preserves a foliation. The global result then follows by integration. Recall that Pansu's differentiability theorem (see [Pansu 1989] or Section 3) says that a quasisymmetric homeomorphism $F : G \rightarrow G'$ between Carnot groups is Pansu-differentiable a.e., and the Pansu differential is a.e. a graded isomorphism between the two Carnot groups. Under the assumption of Theorem 1.1, we show that there are (proper) connected and simply connected subgroups $N \subset G$ and $N' \subset G'$ such that $\phi(N) = N'$ for every graded isomorphism $\phi : G \rightarrow G'$; see Section 2.

In Section 2 we show that graded isomorphisms between reducible Carnot algebras preserve certain subalgebras, which implies that graded isomorphisms between reducible Carnot groups preserve certain subgroups (as indicated in the preceding paragraph). And in Section 3 we show that quasisymmetric homeomorphisms are bilipschitz.

2. Graded isomorphisms of Carnot algebras

In this section we show that graded isomorphisms between reducible Carnot algebras preserve certain subalgebras. This implies that graded isomorphisms between reducible Carnot groups preserve certain subgroups (see Section 3).

A *Carnot Lie algebra* is a finite-dimensional Lie algebra \mathcal{G} together with a direct sum decomposition $\mathcal{G} = V_1 \oplus V_2 \oplus \cdots \oplus V_r$ of vector subspaces such that $[V_1, V_i] = V_{i+1}$ for all $1 \leq i \leq r$, where we set $V_{r+1} = \{0\}$. The integer r is called the degree of nilpotency of \mathcal{G} . Every Carnot algebra $\mathcal{G} = V_1 \oplus V_2 \oplus \cdots \oplus V_r$ admits a one-parameter family of automorphisms $\lambda_t : \mathcal{G} \rightarrow \mathcal{G}$ for $t \in (0, \infty)$, where $\lambda_t(x) = t^i x$ for $x \in V_i$. Let $\mathcal{G} = V_1 \oplus V_2 \oplus \cdots \oplus V_r$ and $\mathcal{G}' = V'_1 \oplus V'_2 \oplus \cdots \oplus V'_s$ be two Carnot algebras. A Lie algebra homomorphism $\phi : \mathcal{G} \rightarrow \mathcal{G}'$ is graded if ϕ commutes with λ_t for all $t > 0$; that is, if $\phi \circ \lambda_t = \lambda_t \circ \phi$. We observe that $\phi(V_i) \subset V'_i$ for all $1 \leq i \leq r$.

A Carnot algebra \mathcal{G} is called reducible if there exist two nontrivial Carnot algebras \mathcal{G}_1 and \mathcal{G}_2 and a graded isomorphism between \mathcal{G} and $\mathcal{G}_1 \oplus \mathcal{G}_2$. It is called irreducible otherwise. The finite dimensionality implies that every reducible Carnot algebra \mathcal{G} can be written as a direct sum of Carnot algebras $\mathcal{G} = \mathcal{G}_0 \oplus \mathcal{G}_1 \oplus \cdots \oplus \mathcal{G}_m$, where \mathcal{G}_0 is abelian, and each \mathcal{G}_i with $i \geq 1$ is nonabelian and irreducible.

Let $\mathcal{G} = V_1 \oplus V_2 \oplus \cdots \oplus V_r$ be a Carnot algebra. When \mathcal{G} is reducible, it also has a decomposition $\mathcal{G} = \mathcal{G}_0 \oplus \mathcal{G}_1 \oplus \cdots \oplus \mathcal{G}_m$ as a direct sum of an abelian factor and irreducible nonabelian factors. We are interested in the question whether graded isomorphisms preserve such decompositions (after possibly permuting the factors). This question is equivalent to the uniqueness problem of such a decomposition. In general the decomposition is not unique, as the following example shows. The author thanks Bruce Kleiner for suggesting the example.

Example 2.1 (Kleiner). Let \mathcal{G} be a nonabelian irreducible Carnot algebra, and let $f : \mathcal{G} \rightarrow \mathbb{R}^m$ be a nontrivial Lie algebra homomorphism into an abelian group. Let $G(f) \subset \mathcal{G} \oplus \mathbb{R}^m$ be the graph of f . Then $G(f)$ is a Carnot algebra (being isomorphic to \mathcal{G}), and $\mathcal{G} \oplus \mathbb{R}^m$ has two different decompositions $\mathcal{G} \oplus \mathbb{R}^m = G(f) \oplus \mathbb{R}^m$. Alternatively, let $g : \mathcal{G} \oplus \mathbb{R}^m \rightarrow \mathcal{G} \oplus \mathbb{R}^m$ be the map given by $g(x, a) = (x, a + f(x))$. Then g is a graded isomorphism and it does not preserve the factor \mathcal{G} .

Despite this example, we show that graded isomorphisms always preserve the abelian factor (Proposition 2.4) and, in the case of a trivial abelian factor, preserve

the decomposition after possibly permuting the factors (Proposition 2.5).

Definition 2-1. Let \mathcal{G} be a Lie algebra and $x \in \mathcal{G}$. Define $d(x) = \dim(\ker(\text{ad } x))$, where $\text{ad } x : \mathcal{G} \rightarrow \mathcal{G}$ is the linear map given by $\text{ad } x(y) = [x, y]$.

Lemma 2.2. *If $d(x) = \dim \mathcal{G} > 1$ for some $x \in V_1 \setminus \{0\}$, then \mathcal{G} is reducible.*

Proof. Note $[x, y] = 0$ for all $y \in \mathcal{G}$. Let \mathcal{G}_1 be the one-dimensional subspace of V_1 spanned by x , and let W be a complementary subspace of \mathcal{G}_1 in V_1 . Set $\mathcal{G}_2 = W \oplus V_2 \oplus \cdots \oplus V_r$. Then $\mathcal{G} = \mathcal{G}_1 \oplus \mathcal{G}_2$ is a direct sum of vector subspaces. The assumption on x now implies that \mathcal{G}_1 and \mathcal{G}_2 are ideals of \mathcal{G} and that both \mathcal{G}_1 and \mathcal{G}_2 are Carnot algebras. Hence \mathcal{G} is reducible. \square

The next lemma provides an intrinsic characterization of the abelian factor \mathcal{G}_0 .

Lemma 2.3. *Let $\mathcal{G} = V_1 \oplus V_2 \oplus \cdots \oplus V_r$ be a Carnot algebra, and consider a direct sum decomposition $\mathcal{G} = \mathcal{G}_0 \oplus \mathcal{G}_1 \oplus \cdots \oplus \mathcal{G}_m$ of \mathcal{G} into an abelian factor and irreducible nonabelian factors. Let $x \in V_1$. Then $x \in \mathcal{G}_0$ if and only if $d(x) = \dim \mathcal{G}$.*

Proof. It is clear that $x \in \mathcal{G}_0$ implies $d(x) = \dim \mathcal{G}$. We assume $d(x) = \dim \mathcal{G}$ and shall prove that $x \in \mathcal{G}_0$. Note $[x, y] = 0$ for all $y \in \mathcal{G}$. Write $x = x_0 + x_1 + \cdots + x_m$ with $x_i \in \mathcal{G}_i \cap V_1$. Suppose $x \notin \mathcal{G}_0$. Then $x_i \neq 0$ for some $i \geq 1$. Since $[x_i, y] = [x, y] = 0$ for all $y \in \mathcal{G}_i$, Lemma 2.2 implies \mathcal{G}_i is reducible, contradicting the assumption. \square

Recall that the goal of this section is to show that a graded isomorphism of reducible Carnot algebras preserves certain Lie subalgebras. The case when the abelian factor is nontrivial is covered by Proposition 2.4.

Proposition 2.4. *Let $\mathcal{G} = \mathcal{G}_0 \oplus \mathcal{G}_1 \oplus \cdots \oplus \mathcal{G}_m$ and $\mathcal{G}' = \mathcal{G}'_0 \oplus \mathcal{G}'_1 \oplus \cdots \oplus \mathcal{G}'_n$ be two reducible Carnot algebras written as direct sums of an abelian factor and irreducible nonabelian factors. Let $\phi : \mathcal{G} \rightarrow \mathcal{G}'$ be a graded isomorphism. Then $\phi(\mathcal{G}_0) = \mathcal{G}'_0$.*

Proof. By Lemma 2.3, $\phi(\mathcal{G}_0) \subset \mathcal{G}'_0$. Since ϕ is an isomorphism, the conclusion follows by considering ϕ^{-1} . \square

Proposition 2.5 treats the case when the abelian factor is trivial.

Proposition 2.5. *Let $\mathcal{G} = \mathcal{G}_0 \oplus \mathcal{G}_1 \oplus \cdots \oplus \mathcal{G}_m$ and $\mathcal{G}' = \mathcal{G}'_0 \oplus \mathcal{G}'_1 \oplus \cdots \oplus \mathcal{G}'_n$ be two reducible Carnot algebras written as direct sums of an abelian factor and irreducible nonabelian factors. Let $\phi : \mathcal{G} \rightarrow \mathcal{G}'$ be a graded isomorphism. Suppose \mathcal{G} has no abelian factor (that is, $\mathcal{G}_0 = \{0\}$). Then $\mathcal{G}'_0 = \{0\}$, $m = n$ and after possibly permuting the factors $\mathcal{G}'_1, \dots, \mathcal{G}'_m$, there exist graded isomorphisms $\phi_i : \mathcal{G}_i \rightarrow \mathcal{G}'_i$ such that $\phi = \phi_1 \oplus \cdots \oplus \phi_m$.*

Now we start the proof of Proposition 2.5. First observe that Proposition 2.4 implies $\mathcal{G}'_0 = \{0\}$. In the following proofs, we shall use both decompositions $\mathcal{G} = V_1 \oplus \cdots \oplus V_r = \mathcal{G}_1 \oplus \cdots \oplus \mathcal{G}_m$ of \mathcal{G} , as well as those for \mathcal{G}' .

Lemma 2.6. *Let $x \in V_1$. Write $x = x_{i_1} + \cdots + x_{i_k}$ ($1 \leq i_1 < \cdots < i_k \leq m$) with $x_{i_j} \in (\mathcal{G}_{i_j} \cap V_1) \setminus \{0\}$. If $k \geq 2$, then $d(x_{i_j}) > d(x)$.*

Proof. We first show that $\ker(\operatorname{ad} x) \subset \ker(\operatorname{ad} x_{i_j})$ for all $1 \leq j \leq k$. Let $y \in \ker(\operatorname{ad} x)$. Write $y = y_1 + \cdots + y_m$ with $y_i \in \mathcal{G}_i$. Then $0 = [x, y] = [x_{i_1}, y_{i_1}] + \cdots + [x_{i_k}, y_{i_k}]$. Since $[x_{i_j}, y_{i_j}] \in \mathcal{G}_{i_j}$, we have $[x_{i_j}, y_{i_j}] = 0$. Hence $[x_{i_j}, y] = [x_{i_j}, y_{i_j}] = 0$; that is, $y \in \ker(\operatorname{ad} x_{i_j})$.

Next we shall find an element $y \in \ker(\operatorname{ad} x_{i_j}) \setminus \ker(\operatorname{ad} x)$. Since $k \geq 2$, there is some $1 \leq l \leq k$ with $l \neq j$. By Lemma 2.2, since \mathcal{G}_{i_l} is nonabelian and irreducible, there is some $y \in \mathcal{G}_{i_l}$ such that $[x_{i_l}, y] \neq 0$. Now notice that $[x_{i_j}, y] = 0$ and $[x, y] = [x_{i_l}, y] \neq 0$. □

For each $1 \leq i \leq m$, set

$$A_i = \{x \in \mathcal{G}_i \cap V_1 : \phi(x) \in \mathcal{G}'_j \text{ for some } j\}.$$

Let $N_i \subset \mathcal{G}_i \cap V_1$ be the vector subspace spanned by A_i . Similarly, for each $1 \leq j \leq n$ set

$$A'_j = \{y \in \mathcal{G}'_j \cap V'_1 : \phi^{-1}(y) \in \mathcal{G}_i \text{ for some } i\}.$$

Let $N'_j \subset \mathcal{G}'_j \cap V'_1$ be the vector subspace spanned by A'_j .

Lemma 2.7. *We have $N_i = \mathcal{G}_i \cap V_1$ for each i and $N'_j = \mathcal{G}'_j \cap V'_1$ for each j .*

Proof. We prove by contradiction. Suppose $N_i \neq \mathcal{G}_i \cap V_1$ for some i or $N'_j \neq \mathcal{G}'_j \cap V'_1$ for some j . Let $d_1 = 0$ if $N_i = \mathcal{G}_i \cap V_1$ for all i ; otherwise, let

$$d_1 = \max\{d(x) : x \in (\mathcal{G}_i \cap V_1) \setminus N_i \text{ for some } i\}.$$

Similarly, let $d_2 = 0$ if $N'_j = \mathcal{G}'_j \cap V'_1$ for all j ; otherwise, let

$$d_2 = \max\{d(y) : y \in (\mathcal{G}'_j \cap V'_1) \setminus N'_j \text{ for some } j\}.$$

Let $d_0 = \max\{d_1, d_2\}$. After possibly replacing ϕ with ϕ^{-1} , we may assume $d_0 = d_1$. Pick $x \in (\mathcal{G}_i \cap V_1) \setminus N_i$ (for some i) with $d(x) = d_0$. By the definition of N_i we have $x \notin A_i$. Hence $\phi(x)$ can be written as

$$(2-2) \quad \phi(x) = y_1 + \cdots + y_k,$$

where $k \geq 2$ and $y_s \in (\mathcal{G}'_{j_s} \cap V'_1) \setminus \{0\}$ for each $1 \leq s \leq k$, and $1 \leq j_1 < \cdots < j_k \leq n$. By Lemma 2.6, $d(y_s) > d(\phi(x)) = d(x) = d_0$. It follows from the definition of d_0 that $y_s \in N'_{j_s}$. Hence there is an expression

$$(2-3) \quad y_s = z_{s,1} + \cdots + z_{s,u_s} + w_{s,1} + \cdots + w_{s,v_s}$$

with $z_{s,p}, w_{s,q} \in A'_{j_s}$ such that $\phi^{-1}(z_{s,p}) \in \mathcal{G}_i \cap V_1$ and $\phi^{-1}(w_{s,q}) \in \mathcal{G}_t \cap V_1$ for some $t \neq i$ (here t may depend on q). Notice that (2-2) and (2-3) imply

$$x = \sum_{s,p} \phi^{-1}(z_{s,p}) + \sum_{s,q} \phi^{-1}(w_{s,q}).$$

Since $x \in \mathcal{G}_i$ and $\mathcal{G} = \mathcal{G}_1 \oplus \cdots \oplus \mathcal{G}_m$ is a direct sum decomposition, we obtain $x = \sum_{s,p} \phi^{-1}(z_{s,p})$. Notice that each $\phi^{-1}(z_{s,p}) \in A_i$. It follows that $x \in N_i$, contradicting the assumption. \square

Lemma 2.8. *For each i , there is some j such that $\phi(\mathcal{G}_i \cap V_1) \subset \mathcal{G}'_j \cap V'_1$.*

Proof. Fix i . By Lemma 2.7, $\mathcal{G}_i \cap V_1 = N_i$. Hence there is a vector space basis B of $\mathcal{G}_i \cap V_1$ consisting of elements of A_i . Write B as a disjoint union $B = \sqcup B_j$, where B_j consists of those elements of B that are mapped into \mathcal{G}'_j under ϕ . Since ϕ is an isomorphism and \mathcal{G}'_{j_1} and \mathcal{G}'_{j_2} commute for $j_1 \neq j_2$, we see that $[X, Y] = 0$ for $X \in B_{j_1}$ and $Y \in B_{j_2}$. Let $E_j \subset \mathcal{G}_i$ be the subalgebra of \mathcal{G}_i generated by B_j . Observe that E_j is an ideal of \mathcal{G}_i and \mathcal{G}_i admits the direct sum decomposition $\mathcal{G}_i = E_1 \oplus \cdots \oplus E_n$. Since \mathcal{G}_i is irreducible, $E_j = \{0\}$ for all j except exactly one. It follows that for some j , all the elements in B are mapped into \mathcal{G}'_j . Since B is a basis of $\mathcal{G}_i \cap V_1$, we have $\phi(\mathcal{G}_i \cap V_1) \subset \mathcal{G}'_j$. \square

Applying Lemma 2.8 to ϕ^{-1} , we see that for each j , there is some i such that $\phi^{-1}(\mathcal{G}'_j \cap V'_1) \subset \mathcal{G}_i \cap V_1$. From this it is easy to see that $m = n$, and after possibly permuting the factors \mathcal{G}'_j we have $\phi(\mathcal{G}_i) = \mathcal{G}'_i$. Proposition 2.5 follows.

3. Quasisymmetric homeomorphisms are bilipschitz

In this section we show that in most cases quasisymmetric homeomorphisms between reducible Carnot groups are bilipschitz.

A simply connected nilpotent Lie group is a *Carnot group* if its Lie algebra is a Carnot algebra. Let G be a Carnot group with Lie algebra $\mathcal{G} = V_1 \oplus \cdots \oplus V_r$. The subspace V_1 defines a left invariant distribution $HG \subset TG$ on G . We fix a left invariant inner product on HG . An absolutely continuous curve γ in G whose velocity vector $\gamma'(t)$ is contained in $H_{\gamma(t)}G$ for a.e. t is called a horizontal curve. By Chow's theorem ([Bellaïche and Risler 1996], Theorem 2.4), any two points of G can be connected by horizontal curves. Let $p, q \in G$; the *Carnot distance* $d(p, q)$ between them is defined as the infimum of length of horizontal curves that join p and q .

Since the inner product on HG is left invariant, the Carnot metric on G is also left invariant. Different choices of inner product on HG result in Carnot metrics that are bilipschitz equivalent. The Hausdorff dimension of G with respect to a Carnot metric is given by $\sum_{i=1}^r i \cdot \dim V_i$. We use the corresponding Hausdorff measure on G . When $G = G_1 \times G_2$ is a direct product of two Carnot groups (with

a suitable choice of inner product on HG), the Carnot metric on G is the product of the Carnot metrics on G_1 and G_2 , and the Hausdorff measure on G is the product of the Hausdorff measures on G_1 and G_2 .

Recall that, for a simply connected nilpotent Lie group G with Lie algebra \mathcal{G} , the exponential map $\exp : \mathcal{G} \rightarrow G$ is a diffeomorphism. Furthermore, the exponential map induces a one-to-one correspondence between Lie subalgebras of \mathcal{G} and connected Lie subgroups of G .

Let G be a Carnot group with Lie algebra $\mathcal{G} = V_1 \oplus \dots \oplus V_r$. Since $\lambda_t : \mathcal{G} \rightarrow \mathcal{G}$ ($t > 0$) is a Lie algebra automorphism and G is simply connected, there is a unique Lie group automorphism $\Lambda_t : G \rightarrow G$ whose differential at the identity is λ_t . For each $t > 0$, Λ_t is a similarity with respect to the Carnot metric: $d(\Lambda_t(p), \Lambda_t(q)) = t d(p, q)$ for any two points $p, q \in G$. A Lie group homomorphism $f : G \rightarrow G'$ between two Carnot groups is a graded homomorphism if it commutes with Λ_t for all $t > 0$; that is, if $f \circ \Lambda_t = \Lambda_t \circ f$. Notice that a Lie group homomorphism $f : G \rightarrow G'$ between two Carnot groups is graded if and only if the corresponding Lie algebra homomorphism is graded.

A Carnot group is reducible if its Lie algebra is reducible. Equivalently, a Carnot group is reducible if it is isomorphic to the direct product of two Carnot groups. A Carnot group is called irreducible otherwise.

Proposition 2.4 and Proposition 2.5 respectively immediately imply Corollary 3.1 and Corollary 3.2, which say that a graded isomorphism of reducible Carnot groups preserves certain Lie subgroups.

Corollary 3.1. *Let $G = G_0 \times G_1 \times \dots \times G_m$ and $G' = G'_0 \times G'_1 \times \dots \times G'_n$ be two reducible Carnot groups written as direct products of an abelian factor and irreducible nonabelian factors. Let $f : G \rightarrow G'$ be a graded isomorphism. Then $f(G_0) = G'_0$.*

Corollary 3.2. *Let $G = G_0 \times G_1 \times \dots \times G_m$ and $G' = G'_0 \times G'_1 \times \dots \times G'_n$ be two reducible Carnot groups written as direct products of an abelian factor and irreducible nonabelian factors. Let $f : G \rightarrow G'$ be a graded isomorphism. Suppose G has no abelian factor (that is, $G_0 = \{e\}$). Then $G'_0 = \{e\}$, $m = n$ and after possibly permuting the factors G'_1, \dots, G'_m , there exist graded isomorphisms $f_i : G_i \rightarrow G'_i$ such that $f = f_1 \times \dots \times f_m$.*

Definition 3-1. Let G and G' be two Carnot groups endowed with Carnot metrics. A map $F : G \rightarrow G'$ is *Pansu-differentiable* at $x \in G$ if there exists a graded homomorphism $L : G \rightarrow G'$ such that

$$\lim_{y \rightarrow x} \frac{d(F(x)^{-1}F(y), L(x^{-1}y))}{d(x, y)} = 0.$$

In this case, the graded homomorphism $L : G \rightarrow G'$ is called the *Pansu differential* of F at x , and is denoted by $dF(x)$.

Definition 3-2. Let $\eta : [0, \infty) \rightarrow [0, \infty)$ be a homeomorphism. A homeomorphism of metric spaces $F : X \rightarrow Y$ is an η -*quasisymmetric* homeomorphism if for all distinct triples $x, y, z \in X$, we have

$$\frac{d(F(x), F(y))}{d(F(x), F(z))} \leq \eta\left(\frac{d(x, y)}{d(x, z)}\right).$$

A map $F : X \rightarrow Y$ is a *quasisymmetric homeomorphism* if it is an η -quasisymmetric homeomorphism for some η .

The following result (except the terminology) is due to Pansu [1989].

Theorem 3.3. *Let $F : G \rightarrow G'$ be a quasisymmetric homeomorphism between two Carnot groups. Then F is a.e. Pansu-differentiable. Furthermore, at a.e. $x \in G$, the Pansu differential $dF(x) : G \rightarrow G'$ is a graded isomorphism.*

In Theorem 3.3 and the proofs below, ‘‘a.e.’’ is with respect to the Hausdorff measure on G .

For the proof of Theorem 1.1, we need the following:

Proposition 3.4. *Let G and G' be two Carnot groups, $W \subset V_1$, $W' \subset V'_1$ be subspaces. Denote by $\mathcal{G}_W \subset \mathcal{G}$ and $\mathcal{G}'_{W'} \subset \mathcal{G}'$, respectively, the Lie subalgebras generated by W and W' . Let $H \subset G$ and $H' \subset G'$, respectively, be the connected Lie subgroups of G and G' corresponding to \mathcal{G}_W and $\mathcal{G}'_{W'}$. Let $F : G \rightarrow G'$ be a quasisymmetric homeomorphism. If $dF(x)(W) \subset W'$ for a.e. $x \in G$, then F sends left cosets of H into left cosets of H' .*

Proof. For each nonzero vector $u \in W$, the set $\{\exp(tu) : t \in \mathbb{R}\}$ is a subgroup of G . It is a geodesic with respect to the Carnot metric and shall be called a horizontal line. For each nonzero vector $u \in W$, let \mathcal{F}_u be the set of left cosets of $\{\exp(tu) : t \in \mathbb{R}\}$ in G . By the main result in [Balogh et al. 2007], $F : G \rightarrow G'$ is absolutely continuous on almost every curve. It follows that for almost every $L \in \mathcal{F}_u$, the map $F|_L : L \rightarrow G'$ is an absolutely continuous curve in G' . On the other hand, by Pansu’s theorem, F is a.e. Pansu-differentiable and the Pansu differential $dF(x) : G \rightarrow G'$ is a graded isomorphism for a.e. $x \in G$. Also by assumption, $dF(x)(W) \subset W'$ for a.e. $x \in G$. It follows from Fubini’s theorem that, for almost every $L \in \mathcal{F}_u$, the Pansu differential $dF(x) : G \rightarrow G'$ exists, is a graded isomorphism and satisfies $dF(x)(W) \subset W'$ for a.e. $x \in L$. Hence, the tangent vectors of the curve $F|_L$ lie in W' almost everywhere. It follows that for almost every $L \in \mathcal{F}_u$, $F(L)$ lies in a left coset of H' . Now the continuity of F and a limiting argument show that the same is true for all $L \in \mathcal{F}_u$. Conceivably, it might be possible for distinct $L_1, L_2 \in \mathcal{F}_u$ to lie in the same left

coset of H , while their images $F(L_1)$ and $F(L_2)$ lie in distinct cosets of H' . We next show that this cannot happen.

In a Carnot group, every two points can be joined by a piecewise geodesic, where each piece is a left translation of a segment in a horizontal line. The preceding paragraph shows that the image under F of each piece lies in a left coset of H' . It follows that the image of the entire piecewise geodesic lies in a left coset of H' . Hence F sends left cosets of H into left cosets of H' . \square

Now we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. Let $F : G \rightarrow G'$ be a quasymmetric homeomorphism between two Carnot groups. Suppose G is reducible and admits a direct product decomposition of irreducible Carnot groups where at least two of the factors are not isomorphic. We first use Proposition 3.4 to show that F preserves a certain foliation. The arguments in [Shanmugalingam and Xie 2012] then show that F is bilipschitz.

Write $G = G_0 \times G_1 \times \cdots \times G_m$ and $G' = G'_0 \times G'_1 \times \cdots \times G'_n$, where G_0, G'_0 are abelian and G_i, G'_j are irreducible nonabelian factors.

First consider the case when G_0 is nontrivial. Let \mathcal{F} be the foliation of G consisting of the cosets of G_0 , and similarly let \mathcal{F}' be the foliation of G' consisting of the cosets of G'_0 . The leaf space of \mathcal{F} can be naturally identified with $N := G_1 \times \cdots \times G_m$, and that of \mathcal{F}' with $N' := G'_1 \times \cdots \times G'_n$. By Corollary 3.1 and Proposition 3.4, the map F sends the leafs of \mathcal{F} to the leafs of \mathcal{F}' . Hence F induces a map $F_1 : N \rightarrow N'$. Notice that $G = G_0 \times N$ with the Carnot metric is isometric to the product of G_0 and N (also equipped with the Carnot metric); similarly for G' . The arguments in [Shanmugalingam and Xie 2012] go through and imply that F_1 is also quasymmetric. Since both the leafs and the leaf spaces are geodesic metric spaces, the arguments further show that F is bilipschitz.

Next we consider the case when G_0 is trivial. Then $G = G_1 \times \cdots \times G_m$ is a direct product of nonabelian irreducible Carnot groups. We combine all isomorphic factors in the above decomposition to obtain $G = N_1 \times \cdots \times N_s$. Each N_j is a direct product of isomorphic nonabelian irreducible Carnot groups, and the factors in N_i and N_j are not isomorphic for $i \neq j$. Similarly, G' can also be written as such a product $G' = N'_1 \times \cdots \times N'_t$. Notice that the assumption of Theorem 1.1 implies that $s \geq 2$. Corollary 3.2 implies that $s = t$, and after possibly permuting the factors N'_i , the Pansu differential satisfies $dF(x)(N_i) = N'_i$ for all i and a.e. $x \in N$. Now the arguments in the preceding paragraph show that F is bilipschitz. The proof of Theorem 1.1 is now complete. \square

Acknowledgments

I thank Bruce Kleiner for helpful discussions, and the referee for useful remarks.

References

- [Balogh et al. 2007] Z. M. Balogh, P. Koskela, and S. Rogovin, “Absolute continuity of quasi-conformal mappings on curves”, *Geom. Funct. Anal.* **17**:3 (2007), 645–664. MR 2009g:30023 Zbl 1134.30014
- [Bellaïche and Risler 1996] A. Bellaïche and J.-J. Risler (editors), *Sub-Riemannian geometry* (Paris, 1992), Progress in Mathematics **144**, Birkhäuser, Basel, 1996. MR 97f:53002 Zbl 0848.00020
- [Bonk and Schramm 2000] M. Bonk and O. Schramm, “Embeddings of Gromov hyperbolic spaces”, *Geom. Funct. Anal.* **10**:2 (2000), 266–306. MR 2001g:53077 Zbl 0972.53021
- [Heinonen and Koskela 1998] J. Heinonen and P. Koskela, “Quasiconformal maps in metric spaces with controlled geometry”, *Acta Math.* **181**:1 (1998), 1–61. MR 99j:30025 Zbl 0915.30018
- [Heintze 1974] E. Heintze, “On homogeneous manifolds of negative curvature”, *Math. Ann.* **211** (1974), 23–34. MR 50 #5695 Zbl 0273.53042
- [Pansu 1989] P. Pansu, “Métriques de Carnot–Carathéodory et quasiisométries des espaces symétriques de rang un”, *Ann. of Math. (2)* **129**:1 (1989), 1–60. MR 90e:53058 Zbl 0678.53042
- [Shanmugalingam and Xie 2012] N. Shanmugalingam and X. Xie, “A rigidity property of some negatively curved solvable Lie groups”, *Comment. Math. Helv.* **87**:4 (2012), 805–823. MR 2984572 Zbl 1255.22005

Received May 17, 2012. Revised May 5, 2013.

XIANGDONG XIE
DEPARTMENT OF MATHEMATICS AND STATISTICS
BOWLING GREEN STATE UNIVERSITY
BOWLING GREEN, OH 43403
UNITED STATES
xiex@bgsu.edu

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Don Blasius
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2013 is US \$400/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 265 No. 1 September 2013

Genus-two Goeritz groups of lens spaces	1
SANGBUM CHO	
A compact embedding theorem for generalized Sobolev spaces	17
SENG-KEE CHUA, SCOTT RODNEY and RICHARD L. WHEEDEN	
Partial integrability of almost complex structures and the existence of solutions for quasilinear Cauchy–Riemann equations	59
CHONG-KYU HAN and JONG-DO PARK	
An overdetermined problem in potential theory	85
DMITRY KHAVINSON, ERIK LUNDBERG and RAZVAN TEODORESCU	
Quasisymmetric homeomorphisms on reducible Carnot groups	113
XIANGDONG XIE	
Capillarity and Archimedes’ principle	123
JOHN MCCUAN and RAY TREINEN	
Generalized eigenvalue problems of nonhomogeneous elliptic operators and their application	151
DUMITRU MOTREANU and MIEKO TANAKA	
Weighted Ricci curvature estimates for Hilbert and Funk geometries	185
SHIN-ICHI OHTA	
On generalized weighted Hilbert matrices	199
EMMANUEL PREISSMANN and OLIVIER LÉVÊQUE	
Unique prime decomposition results for factors coming from wreath product groups	221
J. OWEN SIZEMORE and ADAM WINCHESTER	
On volume growth of gradient steady Ricci solitons	233
GUOFANG WEI and PENG WU	
Classification of moduli spaces of arrangements of nine projective lines	243
FEI YE	



0030-8730(201309)265:1;1-1