

*Pacific  
Journal of  
Mathematics*

**VARIATION OF COMPLEX STRUCTURES AND  
THE STABILITY OF KÄHLER-RICCI SOLITONS**

STUART J. HALL AND THOMAS MURPHY

# VARIATION OF COMPLEX STRUCTURES AND THE STABILITY OF KÄHLER–RICCI SOLITONS

STUART J. HALL AND THOMAS MURPHY

**We investigate the linear stability of Kähler–Ricci solitons for perturbations induced by varying the complex structure within a fixed Kähler class. We calculate stability for the known examples of Kähler–Ricci solitons.**

## 0. Introduction

We consider a stability problem for shrinking Kähler–Ricci solitons. These are critical points of the  $\nu$ -functional, defined by Perelman on the space of Riemannian metrics on a closed manifold  $M$ . The main result is a formula for the second variation of this functional when restricted to perturbations obtained by varying the complex structure within a fixed Kähler class. Such perturbations were first studied by Tian and Zhu [2008] for Kähler–Einstein manifolds, and our paper attempts to extend their results to Kähler–Ricci solitons. Definitions and notation from the main theorem are explained below.

**Theorem 0.1** (Main Theorem). *Let  $(M, g, f)$  be a normalised Kähler–Ricci soliton and let  $h$  be an  $f$ -essential variation. The second variation of the  $\nu$ -functional at  $g, \langle Nh, h \rangle_f$ , is given as*

$$\langle Nh, h \rangle_f = 2 \int_M f \|h\|^2 e^{-f} dV_g.$$

The main utility of this result is that if one had explicit knowledge of the metric and the function  $f$  then it is possible to calculate the quantity  $\langle Nh, h \rangle_f$  quite easily. In Section 4, we do this for all the known examples of Kähler–Ricci solitons. Notice also that for Kähler–Einstein metrics  $f = 0$  and so  $N(h) = 0$ , recovering a result of Tian and Zhu.

The structure of this paper is as follows: In Section 1, we begin with background on Ricci solitons and the stability problem. In Section 2, the space  $\mathcal{W}(g)$  and the space of  $f$ -essential variations in the above theorem are studied. We obtain several

---

Murphy is supported by an A.R.C. grant. We acknowledge the support of a Dennison research grant from the University of Buckingham, which funded a research visit by Murphy.

MSC2010: primary 53C44; secondary 53C25.

Keywords: Kähler–Ricci soliton, Perelman functional, linear stability.

useful characterizations of elements of these spaces. In [Section 3](#), we give a proof of the main theorem. In [Section 4](#), the stability of the known examples of Ricci solitons is investigated.

After a preliminary version of this work was posted on the arXiv, Yuanqi Wang kindly made us aware that he had independently obtained our [Main Theorem](#) as part of his Ph.D. thesis [[Wang 2011](#)]. His proof is similar to ours but proceeds by direct calculation rather than using the results of Dai, Wang and Wei. His thesis also contains interesting results about convergence of the Kähler–Ricci flow to a Kähler–Einstein metric when the complex structure is allowed to vary.

### 1. Ricci solitons and stability

**Background on solitons.** Throughout this paper,  $(M, g)$  is a smooth closed Riemannian manifold.

**Definition 1.1** (Ricci soliton). Let  $X \in \Gamma(TM)$  be a smooth vector field. The triple  $(M, g, X)$  is called a Ricci soliton if it satisfies the equation

$$(1-1) \quad \text{Ric}(g) + L_X g = cg$$

for a constant  $c \in \mathbb{R}$ . If  $c < 0, c = 0, c > 0$  then the soliton is referred to as expanding, steady and shrinking respectively. When  $c \neq 0$ , set  $c = 1/2\tau$ . If  $X = \nabla f$  for a smooth function  $f$  then the soliton is called a gradient Ricci soliton and (1-1) becomes

$$(1-2) \quad \text{Ric}(g) + \text{Hess}(f) = \frac{1}{2\tau}g.$$

When the vector field  $X$  is Killing, an Einstein metric is recovered; Einstein metrics are therefore referred to as trivial Ricci solitons. We can set  $c = 1$  to factor out homothety, and as one may change the soliton potential  $f$  by a constant, let us also require that

$$\int_M f e^{-f} dV_g = 0.$$

A soliton with these choices will be referred to as a *normalised gradient Ricci soliton*.

As well as being interesting as generalisations of Einstein metrics, Ricci solitons also occur as the fixed points of the Ricci flow

$$(1-3) \quad \frac{\partial g}{\partial t} = -2 \text{Ric}(g)$$

up to diffeomorphism. In this paper we will be considering nontrivial Ricci solitons on compact manifolds. Foundational results due to Hamilton [[1995](#)] and Perelman [[2002](#)] imply that expanding and steady Ricci solitons on compact manifolds must

be trivial. Hence our focus is on shrinking Ricci solitons. Perelman also showed that such solitons are necessarily gradient Ricci solitons. We will henceforth refer to these metrics as nontrivial shrinkers.

Due to the work of many people [Cao 1996; Dancer and Wang 2011; Koiso 1990; Podestà and Spiro 2010; Wang and Zhu 2004] there are now many (infinitely many) examples of nontrivial shrinkers. One striking feature all known nonproduct examples share is that they are Kähler. This means that  $\text{Hess}(f)$  is  $J$ -invariant and so the real vector field  $\nabla f$  is holomorphic (see [Besse 1987, 2.124]). In this case the underlying manifold  $M$  is in fact a smooth Fano variety.

Perelman [2002] showed that gradient Ricci solitons are the critical points of a functional, which is usually denoted by  $\nu(g)$ . Let  $f \in C^\infty(M)$  and  $\tau \in \mathbb{R}$ . We say that  $(f, \tau)$  is compatible if

$$\int_M e^{-f} (4\pi\tau)^{-n/2} = 1.$$

**Definition 1.2.** The  $\nu$ -functional is given by

$$\nu(g) = \inf_{\text{compatible } (f, \tau)} \int_M [(R + |\nabla f|^2)\tau + f - n] e^{-f} (4\pi\tau)^{-n/2} dV_g,$$

where  $R$  is the scalar curvature of  $g$ .

As well as giving a variational characterization of Ricci solitons, Perelman showed that the functional is monotonically increasing under the Ricci flow. Hence, if one could perturb a soliton in a direction that increases  $\nu$  and then continue the flow, one would not flow back to the soliton and the soliton would be regarded as unstable.

**Linear stability.** In order to determine the behaviour of the flow around a soliton one can investigate the second variation of  $\nu(g)$  for an admissible perturbation.

**Definition 1.3.** Let  $h \in s^2(T^*M)$ . Then  $g + th$ ,  $t \in \mathbb{R}^+$  is said to be an admissible perturbation. We have  $\partial g / \partial t|_{t=0} = h$ .

If the second variation is strictly negative then the fixed point is stable and attracting. If the second variation has positive directions then one may perturb the soliton and then flow away. Natasha Sesum [2006] has obtained fundamental results on this topic.

**Proposition 1.4** [Cao et al. 2004; Cao and Zhu 2012]. Let  $h \in s^2(TM^*)$  be an admissible variation of a Ricci soliton  $g$ . The second variation of  $\nu$  is given by

$$D_g^2\nu(h, h) = \frac{\tau}{(4\pi\tau)^{n/2}} \int_M \langle Nh, h \rangle e^{-f} dV_g,$$

where

$$(1-4) \quad Nh = \frac{1}{2} \Delta_f h + \text{Rm}(h, \cdot) + \text{div}^* \text{div}_f h + \frac{1}{2} \text{Hess}(v_h) - C(h, g) \text{Ric}.$$

Here  $\Delta_f(\cdot) = \Delta(\cdot) - \nabla_{\nabla_f}(\cdot)$ ,  $\text{div}_f(\cdot) = \text{div}(\cdot) - \iota_{\nabla_f}$ ,  $v_h$  is the solution of the equation

$$\Delta_f v_h + \frac{v_h}{2\tau} = \text{div}_f \text{div}_f(h),$$

and

$$C(h, g) = \frac{\int_M (\text{Ric}, h) e^{-f} dV_g}{\int_M R e^{-f} dV_g}.$$

This operator allows us to define the concept of linear stability.

**Definition 1.5.** Let  $(M, g, f)$  be a Ricci soliton. The soliton is *linearly stable* if the operator  $N$  is nonpositive definite, and *unstable* otherwise.

We now focus upon Kähler–Ricci solitons. The first result regarding stability is the following:

**Theorem 1.6** [Cao et al. 2004; Hall and Murphy 2011; Tian and Zhu 2008]. *Let  $(M, g, f)$  be a Kähler–Ricci soliton. If  $\dim H^{(1,1)}(M) > 1$  then  $(M, g, f)$  is unstable.*

Kähler–Ricci solitons can be viewed as fixed points of a flow related to the Ricci flow (1-3) called the Kähler–Ricci flow, which in the Fano case can be written as

$$(1-5) \quad \frac{\partial g}{\partial t} = -\text{Ric}(g) + g, \quad g(0) = g_0.$$

One important point about this flow is that it preserves the Kähler class. A foundational result about this flow, due to [Cao 1985], is that it exists for all time. The convergence of it is an extremely subtle issue because the complex structure can jump in the limit at infinity. Hence the type of convergence one expects is rather weak. This is illustrated by the following example:

**Theorem 1.7** [Tian and Zhu 2007]. *Let  $M$  be a compact manifold which admits a Kähler–Ricci soliton  $(g_{KRS}, f)$ . Then any solution of (1-5) will converge to  $g_{KRS}$  in the sense of Cheeger–Gromov if the initial metric  $g_0$  is invariant under the maximal compact subset of the automorphism group of  $M$ .*

The unstable perturbations in Theorem 1.6 do not preserve the canonical class. Therefore, from the point of view of the Kähler–Ricci flow it is natural to consider perturbations which fix the Kähler class but allow the complex structure of the manifold to vary. This was initiated by Tian and Zhu [2008].

**Definition 1.8.** Let  $(M, g_{KRS})$  be a Kähler–Ricci soliton with complex structure  $J_{KRS}$ . The space of perturbations is defined as

$$\begin{aligned} \mathcal{W}(g_{KRS}) = \{h \in s^2(TM^*) \mid & \text{there is a family of Kähler metrics } (g_t, J_t) \\ & \text{with } \partial g_t / \partial t|_{t=0} = h, [g_t(J_t \cdot, \cdot)] = c_1(M, J_{KRS}), \\ & (g_0, J_0) = (g_{KRS}, J_{KRS})\}. \end{aligned}$$

The following result was our main motivation for considering this space of perturbations:

**Theorem 1.9 [Tian and Zhu 2008].** *Let  $(M, g_{KE})$  be a Kähler–Einstein metric and let  $h \in \mathcal{W}(g_{KE})$ . Then*

$$\langle N(h), h \rangle_f \leq 0.$$

Tian and Zhu then conjectured that a similar result should be true for Ricci solitons. Our formula in [Theorem 0.1](#) shows that this might not be true in general. The integral in the main theorem does not seem to have a sign in general. However, the examples we calculate in [Section 4](#) do all have  $\langle N(h), h \rangle_f = 0$ ; this seems to be an artefact of their construction rather than a manifestation of some result in complex differential geometry.

We mention here the related study of stability by Dai, Wang, and Wei [[Dai et al. 2007](#)]. They prove that Kähler–Einstein metrics with negative scalar curvature are stable. There is also the recent work of Nefton Pali [[2012](#)] in this area. He considers a related functional known in the literature as the  $W$ -functional (here one is free to pick a volume form whereas in the definition of the  $\nu$ -functional one is determined by the metric).

**Notation and convention.** We use the curvature convention that  $\text{Rm}(X, Y)Z = \nabla_Y \nabla_X Z - \nabla_X \nabla_Y Z + \nabla_{[X, Y]}Z$ . The convention for divergence that we adopt is  $\text{div}(h) = \text{tr}_{1,2}(\nabla h)$ . The rough Laplacian

$$\Delta h = \text{div}(\nabla h) = -\nabla^* \nabla h$$

is then negative definite. Set

$$\langle \cdot, \cdot \rangle_f = \int_M \langle \cdot, \cdot \rangle e^{-f} dV_g$$

to be the twisted inner product on tensors at a Ricci soliton  $(M, g, f)$ . We will denote pointwise inner products induced on tensor bundles by  $g$  with round brackets  $\langle \cdot, \cdot \rangle$ . The adjoint of a differential operator (such as  $\nabla$ ) with respect to this inner product will be denoted with a subscript  $f$  (for example,  $\text{div}_f$ ) throughout.

## 2. Background on variations of complex structure

**Variations of complex structure.** We recall that an almost complex structure on a manifold  $M$  is a section  $J$  of the endomorphism bundle  $\text{End}(TM)$  satisfying  $J^2 = -\text{id}$ . For  $M$  to be a complex manifold we require that the complex structure is *integrable*. By the Newlander–Nirenberg theorem we may take integrable to mean that the Nijenhuis tensor  $\mathcal{N}(J) = 0$ . We will be concerned with infinitesimal variations of complex structure that are modelled on those coming from a one-parameter family of complex structures  $J_t$ . As we are only working at an infinitesimal level, we don’t actually mind if our variations are induced by such a family.

**Definition 2.1** (Infinitesimal variation of complex structure). Let  $(M, g, J)$  be a Kähler manifold. A tensor  $\zeta \in \text{End}(TM)$  is called an infinitesimal variation of complex structure if it satisfies the two equations

$$(2-1) \quad \zeta J + J\zeta = 0,$$

$$(2-2) \quad \dot{\mathcal{N}}(\zeta) = 0,$$

where  $\dot{\mathcal{N}}(\zeta)$  is the infinitesimal variation in the Nijenhuis tensors  $\mathcal{N}(J + t\zeta)$ .

Equation (2-1) simply says that the  $J_t$  are almost complex structures, and (2-2) comes from requiring that they are integrable. In the above definition we are viewing  $\zeta$  as a section of the bundle  $\text{End}(TM)$  which is defined for any manifold. Switching in the usual manner to the complex viewpoint, (2-1) can be thought of as saying that  $\zeta$  is a section of the bundle  $\Lambda^{(0,1)} \otimes TM^{(1,0)}$ . We will variously view the variation as an element of the real bundle  $\text{End}(TM)$ , a section of the bundle  $\Lambda^{(0,1)} \otimes TM^{(1,0)}$ , and, using the metric to lower indices, as a section of  $TM^* \otimes TM^*$  and  $\Lambda^{(0,1)} \otimes \Lambda^{(0,1)}$ . We note that in complex coordinates Equations (2-1) and (2-2) become

$$\zeta_\alpha^\beta = 0 \quad \text{and} \quad \nabla_\alpha \zeta_{\beta\gamma} = \nabla_\beta \zeta_{\alpha\gamma}.$$

The bundle  $\Lambda^{(0,1)} \otimes TM^{(0,1)}$  is an element of the Dolbeault complex

$$TM^{(1,0)} \xrightarrow{\bar{\partial}} \Lambda^{(0,1)} \otimes TM^{(1,0)} \xrightarrow{\bar{\partial}} \Lambda^{(0,2)} \otimes TM^{(1,0)} \xrightarrow{\bar{\partial}} \dots,$$

where  $\bar{\partial}$  is the usual d-bar operator associated to a holomorphic vector bundle over a complex manifold. Equation (2-2) is equivalent to requiring that  $\bar{\partial}\zeta = 0$ .

Analogous to [Tian and Zhu 2008] and following [Koiso 1983], we will decompose the space of infinitesimal variations into *trivial* variations and *f-essential* variations.

By analogy with the twisted inner product, set

$$\Delta_{\bar{\partial}, f} := \bar{\partial}\bar{\partial}_f^* + \bar{\partial}_f^*\bar{\partial}$$

to be the twisted  $\bar{\partial}$ -Laplacian.

**Definition 2.2** (*f*-essential variation). Let  $\zeta$  be an infinitesimal variation of the complex structure  $J$ . We say  $\zeta$  is trivial if  $\zeta = L_Z J$  for a smooth vector field  $Z \in TM$ . A variation  $\zeta$  is said to be *f*-essential if

$$\int_M \langle \zeta, L_Z J \rangle e^{-f} dV_g = 0$$

for all  $Z \in \Gamma(TM)$ .

The following lemma gives a useful characterisation of *f*-essential variations:

**Lemma 2.3** [Koiso 1983, Lemma 6.4]. *Let  $\zeta$  be an f-essential variation and let  $h(\cdot, \cdot) = \omega(\cdot, \zeta \cdot)$ . If  $h$  is symmetric then*

- (1)  $\bar{\partial}_f^* \zeta = 0$ , and
- (2)  $\text{div}_f h = 0$ .

*In particular, an f-essential variation is  $\Delta_{\bar{\partial}, f}$ -harmonic.*

*Proof.* (1) As  $\zeta$  is *f*-essential,

$$\int_M \langle L_Z J, \zeta \rangle e^{-f} = 0$$

for all  $Z \in \Gamma(TM)$ . The Lie derivative of the complex structure is related to the  $\bar{\partial}$ -operator by

$$\bar{\partial}.Z = -\frac{1}{2} J L_Z J(\cdot).$$

Hence, up to a constant,  $\langle L_Z J, \zeta \rangle_f = \langle \bar{\partial} Z, \zeta \rangle_f$  and  $\bar{\partial}_f^* \zeta = 0$ , as claimed.

(2) We begin by noting that  $\zeta$  being *f*-essential means that

$$\langle L_Z J, \zeta \rangle_f = \langle \omega(\cdot, L_Z J(\cdot)), h \rangle_f = 0.$$

Rewriting and using the Cartan formula we have

$$\omega(\cdot, L_Z J(\cdot)) = L_Z g(\cdot, \cdot) - L_Z \omega(\cdot, \cdot) = 2 \text{div}^* Z^\flat(\cdot, \cdot) - (d \circ \iota_Z \omega)(\cdot, J \cdot).$$

The result follows by noting that

$$\langle (d \circ \iota_Z \omega)(\cdot, J \cdot), h \rangle_f = -\langle (d \circ \iota_Z \omega)(\cdot, \cdot), h(\cdot, J \cdot) \rangle_f,$$

and that  $h(\cdot, J \cdot)$  is symmetric. □

In the previous lemma we have assumed that  $h$  is symmetric. This is not strictly necessary by the following argument: If there existed an antisymmetric,  $\Delta_{\bar{\partial}, f}$ -harmonic section of  $\Lambda^{(0,1)} \otimes TM^{(1,0)}$  then there would have to exist an antisymmetric  $\Delta_{\bar{\partial}}$ -harmonic section of  $\Lambda^{(0,1)} \otimes TM^{(1,0)}$  as

$$\mathcal{H}^{p,q}(E) \equiv H^q(M, E \otimes \Lambda^{(p,0)}) \equiv \mathcal{H}_f^{p,q}(E)$$



for any holomorphic vector bundle  $E$ . The Dai–Wei–Wang Weitzenböck formula (Lemma 3.3) and Lemma 3.4 then imply that the associated  $(0, 2)$ -form is parallel. This would imply that  $h^{0,2}(M) > 0$ . One can then appeal to a classical result of Bochner to show that on a Fano manifold such a holomorphic form cannot exist (see [Besse 1987, 11.24]). Tian and Zhu [2008] give a straightforward proof of this fact in the case one is at a Kähler–Einstein metric.

Tian and Zhu decompose the space  $\mathcal{W}(g)$  modulo the action of the diffeomorphism group. They show that

$$\mathcal{W}(g)/\mathcal{D}(M) = \mathcal{A}^{(1,1)} \oplus H^1(M, TM),$$

where  $\mathcal{A}^{(1,1)}$  is the space of  $\partial\bar{\partial}$ -exact  $(1,1)$ -forms and  $H^1(M, TM)$  is the usual cohomology for the holomorphic vector bundle  $TM$ . Tian and Zhu then show that for a general Kähler–Ricci soliton,  $N|_{\mathcal{A}^{(1,1)}} \leq 0$  so that potentially destabilising elements of  $\mathcal{W}$  actually lie in  $H^1(M, TM)$  (they then show that  $N$  vanishes on this space when  $g$  is an Einstein metric). Hence we will only consider perturbations in  $H^1(M, TM)$  and we will use the special representatives given by  $f$ -essential perturbations. Formally:

**Proposition 2.4** [Tian and Zhu 2008]. *Let  $(M, g_{KRS}, J)$  be a Kähler–Ricci soliton. Then we have the following decomposition:*

$$\mathcal{W}(g_{KRS})/\mathcal{D}(M) \cong \mathcal{A}^{(1,1)}(M, J) \oplus H^1(M, TM),$$

where  $\mathcal{D}(M)$  is the diffeomorphism group of  $M$ . The operator  $N$  is nonpositive when restricted to  $\mathcal{A}^{(1,1)}(M, J)$ .

### 3. Proof of Main Theorem

Consider an  $f$ -essential variation of the complex structure  $h \in H^1(M, TM)$ . Firstly, as  $h$  is  $J$ -anti-invariant it is apparent that  $C(h, g) = 0$ . Thus

$$\langle N(h), h \rangle_f = \langle \frac{1}{2} \Delta_f h + \text{Rm}(h, \cdot), h \rangle_f.$$

In order to evaluate the above we will use a Weitzenböck formula. In order to explain the formula we will digress briefly into the spinorial construction used in [Dai et al. 2007]. This is a powerful generalisation of the techniques used by Koiso [1983].

As  $M$  is Fano it has a canonical  $\text{spin}^c$  structure and parallel spinor  $\sigma_0 \in \Gamma(\mathcal{S}^c)$ , where  $\mathcal{S}^c \rightarrow M$  is the  $\text{spin}^c$  spinor bundle. This induces a map

$$\begin{aligned} \Phi : s^2(TM^*) &\rightarrow \mathcal{S}^c \otimes TM^*, \\ \Phi(h) &= h_{ij} e_i \cdot \sigma_0 \otimes e^j, \end{aligned}$$

where  $\{e_i\}$  is an orthonormal basis of  $TM$  and  $e_i \cdot \sigma_0$  denotes Clifford multiplication in  $\mathcal{S}^c$ .

For  $1 \leq i \leq m$ , following [Dai et al. 2007], choose

$$X_i = \frac{e_i - \sqrt{-1}J e_i}{\sqrt{2}} \quad \text{and} \quad \bar{X}_i = \frac{e_i + \sqrt{-1}J e_i}{\sqrt{2}}.$$

Then  $\{X_1, \dots, X_m\}$  is a local unitary frame for  $T^{1,0}M$ . Set  $\{\theta^1, \dots, \theta^m\}$  to be its dual frame. Then

$$\Phi(h) = h(\bar{X}_i, \bar{X}_j)\bar{\theta}^i \otimes \bar{\theta}^j.$$

This can be identified with

$$\Psi(h) = h(\bar{X}_i, \bar{X}_j)\bar{\theta}^i \otimes X_j \in \wedge^{0,1}(TM^{1,0}),$$

where  $TM^{1,0}$  is the holomorphic tangent bundle.

**Lemma 3.1** [Dai et al. 2005, Lemma 2.3]. *For  $h, \tilde{h} \in s^2(TM^*)$ ,*

$$\text{Re}(\Phi(h), \Phi(\tilde{h})) = (h, \tilde{h}).$$

We will also need the following, which is a result of the calculations on page 680 of [Dai et al. 2007]:

**Lemma 3.2.** *Let  $(M, g)$  be a Fano manifold with canonical spin<sup>c</sup> spinor bundle  $\mathcal{S}^c$  and Dirac operator  $D$ . Let  $\Phi$  and  $\Psi$  be defined as above. Then*

$$D\Phi(h) = \sqrt{2}(\bar{\partial} - \bar{\partial}^*)\Psi(h).$$

The main result we need is the following Weitzenböck formula:

**Lemma 3.3** [Dai et al. 2007, Lemma 2.3]. *Let  $h \in s^2(TM^*)$  and let  $D$  be the Dirac operator. Then*

$$(3-1) \quad D^*D(\Phi(h)) = \Phi(\nabla^*\nabla h - 2\text{Rm}(h, \cdot) + \text{Ric} \circ h - h \circ i\rho),$$

where  $\rho$  is the Ricci form.

In order to deal with the Ricci curvature terms we use the following lemma, which is implicit in the proof of Theorem 2.5 in [Dai et al. 2007]:

**Lemma 3.4.** *Let  $h$  be a skew-hermitian section of  $s^2(TM^*)$ . Then*

$$(\text{Ric} \circ h - h \circ i\rho, h) = 0.$$

*Proof.* This is a pointwise calculation. Choose normal coordinates at  $p \in M$ ,  $\{e_1, \dots, e_{2m}\}$ , where  $e_{m+i} = J e_i$  for  $1 \leq i \leq m$ . We can also choose this basis so that the Ricci tensor is diagonalised; that is,  $\text{Ric}(e_i, e_j) = c_i \delta_{ij}$ , where  $c_{m+i} = c_i$ . We have

$$\begin{aligned} \operatorname{Re}(\Phi(\operatorname{Ric} \circ h), \Phi(h)) &= \sum_{i,j=1}^{2m} c_i h_{ij}^2, \\ -\operatorname{Re}(\Phi(h \circ i\rho), \Phi(h)) &= -2 \sum_{j=1}^m \sum_{i=1}^m c_j (h_{(i+m)j} h_{i(j+m)} - h_{ij} h_{(i+m)(j+m)}). \end{aligned}$$

If  $h$  is skew-Hermitian then

$$h_{ij} = -h_{(i+m)(j+m)} \quad \text{and} \quad h_{i(j+m)} = h_{(i+m)j}.$$

Hence

$$-\operatorname{Re}(\Phi(h \circ i\rho), \Phi(h)) = -2 \sum_{i=1}^m \sum_{j=1}^{2m} c_j (h_{ij}^2) = -\sum_{i,j=1}^{2m} c_i h_{ij}^2,$$

and the result follows.  $\square$

The final lemma we need to prove the main result in this section is a technical lemma to deal with the extra term one obtains by using the rescaled volume form  $e^{-f} dV_g$ .

**Lemma 3.5.** *Let  $A \in \Omega^1(M)$  be a one-form and  $B \in \otimes^k TM^*$ . Then*

- (1)  $\operatorname{div}(A \otimes B) = \operatorname{div}(A) \otimes B + \nabla_{A^\sharp} B$ ,
- (2)  $\operatorname{div}(df \otimes h) = (\Delta f)h + \nabla_{\nabla f} h$ , and
- (3)  $-\langle \nabla_{\nabla f} h, h \rangle_f = \frac{1}{2} \int_M \Delta f f \|h\|^2 e^{-f} dV_g$ .

*Proof.* (1) We calculate using a normal, orthonormal basis  $\{e_i\}$ ,

$$\operatorname{div}(A \otimes B) = \nabla_{e_i} (A \otimes B)(e_i, \cdot) = \operatorname{div}(A) \otimes B + \nabla_{A^\sharp} B.$$

(2) We use  $A = df$ ,  $B = h$  in (1).

(3) We note that

$$\langle \nabla_{\nabla f} h, h \rangle_f = \langle \iota_{\nabla f} \nabla h, h \rangle_f = \langle \nabla h, df \otimes h \rangle_f = -\langle h, \operatorname{div}_f(df \otimes h) \rangle_f.$$

Now using (2) we have

$$\begin{aligned} \langle \nabla_{\nabla f} h, h \rangle_f &= \int_M |\nabla f|^2 \|h\|^2 e^{-f} dV_g - \langle h, \operatorname{div}(df \otimes h) \rangle_f \\ &= -\int_M (\Delta f f) \|h\|^2 e^{-f} dV_g - \langle \nabla_{\nabla f} h, h \rangle_f. \end{aligned} \quad \square$$

As is well known, the soliton potential function of a normalised gradient Ricci soliton solves the equation

$$\Delta f = -2f.$$

*Proof of Main Theorem.* Lemmas 3.1, 3.2, and 3.3 yield that pointwise

$$\begin{aligned} \left(\frac{1}{2}\Delta h + \text{Rm}(h, \cdot), h\right) &= \text{Re}\left(\Phi\left(\frac{1}{2}\Delta h + \text{Rm}(h, \cdot)\right), \Phi(h)\right) \\ &= \text{Re}\left((D^*D\Phi)(h), \Phi(h)\right) \\ &= \text{Re}\left(-2\Delta_{\bar{\partial}}\Psi(h), \Psi(h)\right). \end{aligned}$$

However, as  $h$  is  $f$ -essential then  $\Psi(h)$  is orthogonal to the image of  $\Delta_{\bar{\partial}}$  with respect to the global inner product. Hence

$$\int_M \left(\frac{1}{2}\Delta h + \text{Rm}(h, \cdot), h\right)e^{-f} dV_g = 0. \quad \square$$

#### 4. Examples and applications

**Setup.** As mentioned in the introduction, there are three main sources for concrete examples of Kähler–Ricci solitons: the Dancer–Wang, Podestà–Spiro, and the Wang–Zhu examples. The Wang–Zhu solitons exist on toric–Kähler manifolds and are nontrivial precisely when the Futaki invariant is nonzero. Unfortunately, this class of manifold does not admit any nontrivial deformations of complex structure:

**Theorem 4.1** [Bien and Brion 1996, Theorem 3.2]. *Every Fano toric–Kähler manifold  $M$  has  $H^1(M, TM) = 0$ .*

Similarly, one can see the Podestà–Spiro examples are rigid. The next class of examples to investigate are provided by the Dancer–Wang solitons. These solitons are generalisations of the soliton on  $\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$  constructed by Koiso [1990] and Cao [1996]. We begin by reviewing their construction.

Let  $(V_i, r_i, J_i)$ ,  $1 \leq i \leq r$  be Fano Kähler–Einstein manifolds with first Chern class  $c_1(V_i, J_i) = p_i a_i$ , where  $p_i$  are positive integers and  $a_i \in H^2(V_i; \mathbb{Z})$  are indivisible classes. The Kähler–Einstein metrics  $r_i$  are normalised so that  $\text{Ric}(r_i) = p_i r_i$ . For  $q = (q_1, \dots, q_r)$  with  $q_i \in \mathbb{Z} - \{0\}$ , let  $P_q$  be the total space of the principal  $U(1)$ -bundle over  $B := V_1 \times V_2 \times \dots \times V_r$  with Euler class  $\sum_1^r q_i \pi_i^* a_i$ , where

$$\pi_i : V_1 \times \dots \times V_r \rightarrow V_i$$

is the projection onto the  $i$ -th factor. Denote by  $M_0$  the product  $I \times P_q$  for the unit interval  $I$ . We denote by  $\theta$  the principal  $U(1)$ -connection on  $P_q$  with curvature

$$\Omega := \sum_{i=1}^r q_i \pi_i^* \eta_i,$$

where  $\eta_i$  is the Kähler form of  $r_i$ . There is a one-parameter family of metrics on  $P_q$  given by

$$g_t := f^2(t)\theta \otimes \theta + \sum_{i=1}^r l_i^2(t)\pi_i^* r_i,$$

where  $f$  and  $l_i$  are smooth functions on  $I$  with prescribed boundary behaviour. Finally, consider the metric on  $M_0$  given by

$$g = dt^2 + g_t,$$

with the correct boundary behaviour of  $f$  and the  $l_i$ . This metric then extends to a metric on a compactification of  $M_0$ , which we denote  $M$ .

The complex structure on this manifold can be described explicitly by lifting the complex structure on the base and requiring that  $J(N) = -f(t)^{-1}Z$ , where  $N = \partial_t$  is normal to the hypersurfaces, and  $Z$  is the Killing vector that generates the isometric  $U(1)$  action on  $P_q$ .

**Deformations of Dancer–Wang solitons.** The Ricci soliton equations in this setting reduce to a system of ODEs. We have the following existence theorem:

**Theorem 4.2** [Dancer and Wang 2011, Theorem 4.30]. *Let  $M$  denote the compactification of  $M_0$  as above. Then  $M$  admits a Kähler–Ricci soliton  $(M, g, u)$  which is Einstein if and only if the associated Futaki invariant vanishes.*

We refer to [Dancer and Wang 2011] for details of the constructions. If one chooses the components  $V_i$  to be homogeneous Kähler–Einstein manifolds then the resulting  $M$  is toric. However, by choosing the components  $V_i$  to be nonhomogeneous, Fano and Kähler–Einstein, and calculating the Futaki invariant, they give examples of nontoric Kähler–Ricci solitons. It is these that may admit complex deformations.

Suppose that  $V_i$  is a Fano, Kähler–Einstein manifold admitting deformations of its complex structure  $J_i$ . We consider an essential variation  $h_i$  in the Kähler metric  $r_i$  such that the Kähler form  $\eta_i = r_i(J_i \cdot, \cdot)$  remains in the class  $c_1(V_i, J_0)$ . This induces a variation in the metric on the whole space given by

$$h = l_i^2(t)\pi^*h_i.$$

Clearly the same procedure works for any product of Kähler–Einstein manifolds with some (or all) of the factors admitting complex deformations. Here it is simply stated for one factor for simplicity. Let us state our final result:

**Theorem 4.3.** *For this perturbation  $h$ , one has  $N(h) = 0$ .*

*Proof.* It follows from the construction of  $h$  that the pointwise norm  $\|h\|$  is independent of  $t$ . It also follows that if  $h_i$  is essential then  $h$  is  $u$ -essential. We see now that

$$\langle Nh, h \rangle = \int_M u \|h\|^2 e^{-u} dV_g = \|h\|_{L^2}^2(V_i) \int_I u e^{-u} dt = 0. \quad \square$$

**Remark 4.4.** The significance of this result is that it verifies Tian–Zhu’s conjecture for every obvious example of a complex deformation of the known Kähler–Ricci solitons. We do not know of any explicit deformations beyond these.

It is notable that for all  $f$ -essential perturbations  $h$  known to us, one has  $N(h) = 0$ . Understanding if this is always the case would involve calculating  $H^1(M, TM)$ , which is not easy to calculate in general.

### Acknowledgements

It is a pleasure to thank Joel Fine for useful conversations and Nefton Pali for helpful comments on the paper. We also thank the anonymous referees for corrections and suggestions.

### References

- [Besse 1987] A. L. Besse, *Einstein manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **10**, Springer, Berlin, 1987. [MR 88f:53087](#) [Zbl 0613.53001](#)
- [Bien and Brion 1996] F. Bien and M. Brion, “Automorphisms and local rigidity of regular varieties”, *Compositio Math.* **104**:1 (1996), 1–26. [MR 97h:14034](#) [Zbl 0910.14004](#)
- [Cao 1985] H.-D. Cao, “Deformation of Kähler metrics to Kähler–Einstein metrics on compact Kähler manifolds”, *Invent. Math.* **81**:2 (1985), 359–372. [MR 87d:58051](#) [Zbl 0574.53042](#)
- [Cao 1996] H.-D. Cao, “Existence of gradient Kähler–Ricci solitons”, pp. 1–16 in *Elliptic and parabolic methods in geometry* (Minneapolis, MN, 1994), edited by B. Chow et al., A K Peters, Wellesley, MA, 1996. [MR 98a:53058](#) [Zbl 0868.58047](#) [arXiv 1203.4794](#)
- [Cao and Zhu 2012] H.-D. Cao and M. Zhu, “On second variation of Perelman’s Ricci shrinker entropy”, *Math. Ann.* **353**:3 (2012), 747–763. [MR 2923948](#) [Zbl 1252.53074](#)
- [Cao et al. 2004] H.-D. Cao, R. S. Hamilton, and T. Ilmanen, “Gaussian densities and stability for some Ricci solitons”, preprint, 2004. [arXiv math/0404165](#)
- [Dai et al. 2005] X. Dai, X. Wang, and G. Wei, “On the stability of Riemannian manifold with parallel spinors”, *Invent. Math.* **161**:1 (2005), 151–176. [MR 2006h:53041](#) [Zbl 1075.53042](#)
- [Dai et al. 2007] X. Dai, X. Wang, and G. Wei, “On the variational stability of Kähler–Einstein metrics”, *Comm. Anal. Geom.* **15**:4 (2007), 669–693. [MR 2009a:53078](#) [Zbl 1149.53027](#)
- [Dancer and Wang 2011] A. S. Dancer and M. Y. Wang, “On Ricci solitons of cohomogeneity one”, *Ann. Global Anal. Geom.* **39**:3 (2011), 259–292. [MR 2012a:53124](#) [Zbl 1215.53040](#)
- [Hall and Murphy 2011] S. J. Hall and T. Murphy, “On the linear stability of Kähler–Ricci solitons”, *Proc. Amer. Math. Soc.* **139**:9 (2011), 3327–3337. [MR 2012g:53135](#) [Zbl 1231.53056](#)
- [Hamilton 1995] R. S. Hamilton, “The formation of singularities in the Ricci flow”, pp. 7–136 in *Proceedings of the Conference on Geometry and Topology* (Cambridge, MA, 1993), edited by C.-C. Hsiung and S.-T. Yau, Surveys in Differential Geometry **2**, Int. Press, Cambridge, MA, 1995. [MR 97e:53075](#) [Zbl 0867.53030](#)
- [Koiso 1983] N. Koiso, “Einstein metrics and complex structures”, *Invent. Math.* **73**:1 (1983), 71–106. [MR 85d:58018](#) [Zbl 0515.53040](#)
- [Koiso 1990] N. Koiso, “On rotationally symmetric Hamilton’s equation for Kähler–Einstein metrics”, pp. 327–337 in *Recent topics in differential and analytic geometry*, edited by T. Ochiai, Adv. Stud. Pure Math. **18**, Academic Press, Boston, 1990. [MR 93d:53057](#) [Zbl 0739.53052](#)
- [Pali 2012] N. Pali, “The total second variation of Perelman’s  $\mathcal{W}$ -functional”, preprint, 2012. [arXiv 1201.0969](#)

- [Perelman 2002] G. Perelman, “The entropy formula for the Ricci flow and its geometric applications”, preprint, 2002. [Zbl 1130.53001](#) [arXiv math/0211159](#)
- [Podestà and Spiro 2010] F. Podestà and A. Spiro, “Kähler–Ricci solitons on homogeneous toric bundles”, *J. Reine Angew. Math.* **642** (2010), 109–127. [MR 2011f:32049](#) [Zbl 1210.14057](#)
- [Sesum 2006] N. Sesum, “Linear and dynamical stability of Ricci-flat metrics”, *Duke Math. J.* **133**:1 (2006), 1–26. [MR 2007c:53089](#) [Zbl 1103.53040](#)
- [Tian and Zhu 2007] G. Tian and X. Zhu, “Convergence of Kähler–Ricci flow”, *J. Amer. Math. Soc.* **20**:3 (2007), 675–699. [MR 2007k:53107](#) [Zbl 1185.53078](#)
- [Tian and Zhu 2008] G. Tian and X. Zhu, “Perelman’s  $\mathcal{W}$ -functional and stability of Kähler–Ricci flow”, preprint, 2008. [arXiv 0801.3504](#)
- [Wang 2011] Y. Wang, *On Ricci solitons and Ricci flows*, Ph.D. thesis, University of Wisconsin, Madison, WI, 2011, Available at <http://search.proquest.com/docview/910543286>. [MR 2982322](#)
- [Wang and Zhu 2004] X.-J. Wang and X. Zhu, “Kähler–Ricci solitons on toric manifolds with positive first Chern class”, *Adv. Math.* **188**:1 (2004), 87–103. [MR 2005d:53074](#) [Zbl 1086.53067](#)

Received July 6, 2012. Revised April 16, 2013.

STUART J. HALL  
DEPARTMENT OF APPLIED COMPUTING  
UNIVERSITY OF BUCKINGHAM  
HUNTER STREET  
BUCKINGHAM, MK18 1G  
UNITED KINGDOM  
[stuart.hall@buckingham.ac.uk](mailto:stuart.hall@buckingham.ac.uk)

THOMAS MURPHY  
DÉPARTAMENT DE MATHÉMATIQUE  
UNIVERSITÉ LIBRE DE BRUXELLES  
CP 218, BOULEVARD DU TRIOMPHE  
1050 BRUXELLES  
BELGIUM  
[thomas.murphy@ulb.ac.be](mailto:thomas.murphy@ulb.ac.be)

# PACIFIC JOURNAL OF MATHEMATICS

[msp.org/pjm](http://msp.org/pjm)

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pacific@math.ucla.edu](mailto:pacific@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Don Blasius  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Vijayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2013 is US \$400/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).


---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers



# PACIFIC JOURNAL OF MATHEMATICS

Volume 265    No. 2    October 2013

---

Singularity removability at branch points for Willmore surfaces	257
YANN BERNARD and TRISTAN RIVIÈRE	
On Bach flat warped product Einstein manifolds	313
QIANG CHEN and CHENXU HE	
On plane sextics with double singular points	327
ALEX DEGTYAREV	
A computational approach to the Kostant–Sekiguchi correspondence	349
HEIKO DIETRICH and WILLEM A. DE GRAAF	
Landau–Toeplitz theorems for slice regular functions over quaternions	381
GRAZIANO GENTILI and GIULIA SARFATTI	
On surgery curves for genus-one slice knots	405
PATRICK M. GILMER and CHARLES LIVINGSTON	
Characterizing abelian varieties by the reduction of the Mordell–Weil group	427
CHRIS HALL and ANTONELLA PERUCCA	
Variation of complex structures and the stability of Kähler–Ricci Solitons	441
STUART J. HALL and THOMAS MURPHY	
On crossed homomorphisms of the volume preserving diffeomorphism groups	455
RYOJI KASAGAWA	
Regularity at the boundary and tangential regularity of solutions of the Cauchy–Riemann system	491
TRAN VU KHANH and GIUSEPPE ZAMPIERI	
On the Steinberg character of a semisimple $p$ -adic group	499
JU-LEE KIM and GEORGE LUSZTIG	