

*Pacific
Journal of
Mathematics*

**DEFORMATION OF THREE-DIMENSIONAL HYPERBOLIC
CONE STRUCTURES: THE NONCOLLAPSING CASE**

ALEXANDRE PAIVA BARRETO

DEFORMATION OF THREE-DIMENSIONAL HYPERBOLIC CONE STRUCTURES: THE NONCOLLAPSING CASE

ALEXANDRE PAIVA BARRETO

Dedicated to my wife Cynthia

This work is devoted to the study of deformations of hyperbolic cone structures under the assumption that the length of the singularity remains uniformly bounded over the deformation. Let (M_i, p_i) be a sequence of pointed hyperbolic cone manifolds with cone angles of at most 2π and topological type (M, Σ) , where M is a closed, orientable and irreducible 3-manifold and Σ an embedded link in M . Assuming that the length of the singularity remains uniformly bounded, we prove that either the sequence M_i collapses and M is Seifert fibered or a Sol manifold, or the sequence M_i does not collapse and, in this case, a subsequence of (M_i, p_i) converges to a complete three dimensional Alexandrov space endowed with a hyperbolic metric of finite volume on the complement of a finite union of quasigeodesics. We apply this result to a question proposed by Thurston and to provide universal constants for hyperbolic cone structures when Σ is a small link in M .

1. Introduction

This text focuses on deformations of hyperbolic cone structures on a closed, orientable and irreducible 3-manifold M which are singular along a fixed embedded link $\Sigma = \Sigma_1 \sqcup \cdots \sqcup \Sigma_l$. Unlike complete hyperbolic structures, which are rigid by Mostow's theorem, the hyperbolic cone structures can be deformed (see [Hodgson and Kerckhoff 1998]). The difficulty in understanding these deformations lies in the possibility that the structure degenerates. In other words, the Hausdorff–Gromov limit (see Section 2 for the definition) of the deformation is only an Alexandrov space which may have dimension strictly smaller than 3, although its curvature remains bounded from below by -1 (see [Kojima 1998]).

From [Kojima 1998; Hodgson and Kerckhoff 2005; Fujii 2000] it is known that the degeneration of the hyperbolic cone structures occurs if and only if the

This work is part of Barreto's doctoral thesis, made at Université Paul Sabatier–Toulouse under the supervision of Professor Michel Boileau and supported by CAPES and FAPESP (2009/16234-5).

MSC2010: primary 57M50; secondary 57N16, 53C23.

Keywords: hyperbolic 3-manifolds, cone manifolds, Alexandrov spaces.

singular link of these structures intersects itself during the deformation. When the cone angles vary between 0 and π , the Dirichlet polyhedron of the hyperbolic cone structures is convex and we can use this fact to avoid self intersections of the singular link over deformations (see [Kojima 1998]). In this article we will not use this restrictive assumption and allow the cone angles vary until 2π .

We are interested in studying the following question that was proposed by W. Thurston in the 1980s:

Question 1. *Let M be a closed and orientable hyperbolic 3-manifold and suppose there exists a simple closed geodesic Σ in M . Can the hyperbolic structure of M be deformed to the complete hyperbolic structure on $M - \Sigma$ through a path M_α of hyperbolic cone structures with topological type (M, Σ) and parametrized by the cone angles $\alpha \in [0, 2\pi]$?*

If the deformation proposed by Thurston exists, it is a consequence of his hyperbolic Dehn surgery theorem that the length of the singular link must converge to zero. In particular, we have that its length remains uniformly bounded over the deformation. This conclusion give us a necessary condition for the existence of Thurston's desired deformation. For this reason, we will focus only on deformations of hyperbolic cone structures with this additional hypothesis on the singularity's length. We remark that this assumption is automatically verified when the holonomy representations of the hyperbolic cone structures are convergent.

We started studying this question in [Barreto 2012]. In that paper we obtained the following result (see Section 3 for the definition of collapse):

Theorem 2. *Let M be a closed, orientable and irreducible 3-manifold and let $\Sigma = \Sigma_1 \sqcup \dots \sqcup \Sigma_l$ be an embedded link in M . Suppose there exists a sequence M_i of hyperbolic cone manifolds with topological type (M, Σ) and having cone angles $\alpha_{ij} \in (0, 2\pi]$ along Σ_j for $i \in \mathbb{N}$. Denote by $\mathcal{L}_{M_i}(\Sigma_j)$ the length of the connected component Σ_j of Σ in the hyperbolic cone manifold M_i . If*

$$(1-1) \quad \sup\{\mathcal{L}_{M_i}(\Sigma_j) \mid i \in \mathbb{N} \text{ and } j \in \{1, \dots, l\}\} < \infty$$

and the sequence M_i collapses, then M is Seifert fibered or a Sol manifold.

As a consequence of this theorem, we obtained the following result yielding some information on Thurston's question:

Corollary 3. *Let M be a closed and orientable hyperbolic 3-manifold and suppose there exists a finite union of disjoint simple closed geodesics Σ in M . Let M_α be a (angle decreasing) deformation of this structure along a continuous path of hyperbolic cone structures with topological type (M, Σ) and having cone angles $\alpha \in (L, 2\pi] \subset [0, 2\pi]$ (the same for all components of Σ). If*

$$(1-2) \quad \sup\{\mathcal{L}_{M_\alpha}(\Sigma_j) \mid \alpha \in (L, 2\pi] \text{ and } j \in \{1, \dots, l\}\} < \infty,$$

then every convergent sequence M_{α_i} , with α_i converging to L , does not collapse.

In this article, we will focus on noncollapsing deformations of hyperbolic cone structures. The principal result of this paper is the following one:

Theorem 4. *Let M be a closed, orientable and irreducible 3-manifold and let $\Sigma = \Sigma_1 \sqcup \cdots \sqcup \Sigma_l$ be an embedded link in M . Suppose there exists a sequence M_i of hyperbolic cone manifolds with topological type (M, Σ) and having cone angles $\alpha_{ij} \in (0, 2\pi]$ along Σ_j for $i \in \mathbb{N}$. Denote by $\mathcal{L}_{M_i}(\Sigma_j)$ the length of the connected component Σ_j of Σ in the hyperbolic cone manifold M_i . If*

$$\sup\{\mathcal{L}_{M_i}(\Sigma_j) \mid i \in \mathbb{N} \text{ and } j \in \{1, \dots, l\}\} < \infty,$$

then one of the following statements holds:

- (i) *The sequence M_i collapses and M is Seifert fibered or a Sol manifold.*
- (ii) *The sequence M_i does not collapse and there exists a sequence of points $p_{i_k} \in M - \Sigma$ such that the sequence (M_{i_k}, p_{i_k}) converges to a three-dimensional pointed Alexandrov space (Z, z_0) . The Alexandrov space Z is endowed with a (noncomplete) hyperbolic metric of finite volume on the complement of a finite union Σ_Z of quasigeodesics. Moreover, Z is homeomorphic to M (in particular, Z is compact) if there exists $\varepsilon \in (0, 2\pi)$ such that the cone angles α_{ij} belong to $(\varepsilon, 2\pi]$. Further, the following statements are equivalent:*
 - (a) *Z is compact.*
 - (b) $\inf\{\text{cone angle}_{M_{i_k}}(\Sigma_j) \mid k \in \mathbb{N} \text{ and } \Sigma_j \subset \Sigma\} > 0.$
 - (c) $\inf\{\mathcal{L}_{M_{i_k}}(\Sigma_j) \mid k \in N\} > 0$ for each component Σ_j of Σ .

Remark 5. A byproduct of this theorem is that the length of a connected component Σ_j of Σ shrinks down to zero if and only if the same arises for its cone angles α_{ij} (when i goes to infinity). If the cone angles are supposed to be the same on all connected components of Σ , it follows from this (see [Corollary 23](#)) that the sequence of cone angles converges to zero if and only if the following statements hold:

- (i) $\sup\{\mathcal{L}_{M_i}(\Sigma) \mid i \in \mathbb{N}\} < \infty.$
- (ii) $\lim_{i \rightarrow \infty} \text{diam}(M_i) = \infty.$
- (iii) The sequence M_i does not collapse.

In general, the limiting singular locus Σ_Z need not be a disjoint union of quasigeodesics since the singular link could intersect itself as cone angles are changed. It seems possible that the components of Σ_Z are continuous geodesics and that the limit is a hyperbolic cone manifold in a more general sense allowing singularities along a graph instead of a link. The main problem in understanding the limiting singular locus lies in the possibility that the singularity intersects itself infinitely

many times at the limit. More precisely, Σ_Z may be a graph with infinite degree vertices. A better comprehension of the limiting singular locus is an interesting problem for further investigation.

As an application of [Theorem 4](#), we obtain the following result related to [Question 1](#).

Corollary 6. *Let M be a closed and orientable hyperbolic 3-manifold and suppose there exists a finite union of disjoint simple closed geodesics Σ in M . Let M_α be a deformation of this structure along a continuous path of hyperbolic cone structures with topological type (M, Σ) and having cone angles $\alpha \in (\theta, 2\pi] \subset [0, 2\pi]$ (the same for all components of Σ). The following statements are equivalent:*

- (i) $\theta = 0$ and the path M_α extends continuously to $[0, 2\pi]$, where M_0 denotes $M - \Sigma$ with the complete hyperbolic metric.
- (ii) $\lim_{\alpha \rightarrow \theta} \mathcal{L}_{M_\alpha}(\Sigma) = \lim_{\alpha \rightarrow \theta} \sum_{i=1}^l \mathcal{L}_{M_\alpha}(\Sigma_i) = 0$.
- (iii) *There exists a sequence $\alpha_i \in (\theta, 2\pi]$ converging to θ satisfying*

$$\sup\{\mathcal{L}_{M_{\alpha_i}}(\Sigma_j) \mid \alpha_i \in (\theta, 2\pi] \text{ and } j \in \{1, \dots, l\}\} < \infty$$

and such that the sequence $\text{diam}(M_{\alpha_i})$ goes to infinity with i .

Remark 7. [Corollary 6](#) provides a necessary and sufficient condition for the existence of the deformation proposed by Thurston. Using the notation in [Question 1](#),

$$\theta = 0 \quad \text{if and only if} \quad \lim_{\alpha \rightarrow \theta} \mathcal{L}_{M_\alpha}(\Sigma) = 0.$$

Supposing in addition that M is not Seifert fibered and that Σ is a small link in M , we have also the following theorem (see [Corollaries 25](#) and [26](#)) providing universal constants for hyperbolic cone structures with topological type (M, Σ) .

Theorem 8. *Let M be a closed, orientable, irreducible and non-Seifert fibered 3-manifold and let Σ be a small link in M . There exists a constant $V = V(M, \Sigma) > 0$ and a constant $K = K(M, \varepsilon) > 0$, for each $\varepsilon \in (0, 2\pi)$, such that*

- (i) $\text{Vol}(\mathcal{M}) > V$ for every hyperbolic cone manifold \mathcal{M} with topological type (M, Σ) , and
- (ii) $\text{diam}(\mathcal{M}) < K$ for every hyperbolic cone manifold \mathcal{M} with topological type (M, Σ) and having cone angles in the interval $(\varepsilon, 2\pi]$.

2. Metric geometry

In this section, we recall some definitions about Alexandrov spaces and Hausdorff–Gromov convergence. We refer to [\[Burago et al. 2001; Burago et al. 1992; Gromov 1981; Perelman and Petrunin 1994\]](#) for details.

Given a metric space Z , the metric on Z will always be denoted by $d_Z(\cdot, \cdot)$. The open ball of radius $r > 0$ about a subset A of Z will be denoted by

$$B_Z(A, r) = \bigcup_{a \in A} \{z \in Z \mid d_Z(z, a) < r\}.$$

A metric space Z is called a *length space* (and its metric is called *intrinsic*) when the distance between every pair of points in Z is given by the infimum of the lengths of all rectifiable curves connecting them. When a minimizing geodesic between every pair of points exists, we say that Z is *complete*.

For all $k \in \mathbb{R}$, denote by \mathbb{M}_k^2 the complete and simply connected two-dimensional Riemannian manifold of constant sectional curvature equal to k .

Let $\Delta(x, y, z) \subset Z$ be a geodesic triangle in Z with vertices $x, y, z \in Z$. The angle of $\Delta(x, y, z)$ at vertex x , for example, will be denoted by $\angle_\Delta(x)$. A *comparison triangle* for $\Delta(x, y, z) \subset Z$ in \mathbb{M}_k^2 is a geodesic triangle $\bar{\Delta}_k(\bar{x}, \bar{y}, \bar{z}) \subset \mathbb{M}_k^2$ satisfying

$$d_{\mathbb{M}_k^2}(\bar{x}, \bar{y}) = d_Z(x, y), \quad d_{\mathbb{M}_k^2}(\bar{y}, \bar{z}) = d_Z(y, z), \quad \text{and} \quad d_{\mathbb{M}_k^2}(\bar{z}, \bar{x}) = d_Z(z, x).$$

Definition 9. A length space Z is called an Alexandrov space of curvature not smaller than $k \in \mathbb{R}$ if every point of Z has a neighborhood U such that, the angles of every triangle $\Delta(x, y, z) \subset U$ are well defined and satisfy the inequalities

$$\angle_\Delta(x) \geq \angle_{\bar{\Delta}_k}(\bar{x}), \quad \angle_\Delta(y) \geq \angle_{\bar{\Delta}_k}(\bar{y}), \quad \text{and} \quad \angle_\Delta(z) \geq \angle_{\bar{\Delta}_k}(\bar{z})$$

for every comparison triangle $\bar{\Delta}_k(\bar{x}, \bar{y}, \bar{z}) \subset \mathbb{M}_k^2$ of Δ .

Suppose from now on that Z is an n -dimensional Alexandrov space of curvature not smaller than $k \in \mathbb{R}$ and fix a point $O \in \mathbb{M}_k^2$. We next recall the definition of quasigeodesics on an Alexandrov space (see [Perelman and Petrunin 1994]). Let $\gamma : [a, b] \rightarrow Z$ be a 1-Lipschitz curve and let $z \in Z$ be a point satisfying

$$(2-1) \quad 0 < d_Z(z, \gamma(t)) < \frac{\pi}{\sqrt{k}}$$

for all $t \in [a, b]$. We say that a curve $\tilde{\gamma} : [a, b] \rightarrow \mathbb{M}_k^2$ is a *development* of γ with respect to $z \in Z$ when

$$d_Z(z, \gamma(t)) = d_{\mathbb{M}_k^2}(O, \tilde{\gamma}(t))$$

for all $t \in [a, b]$.

Definition 10. A 1-Lipschitz curve $\gamma : [a, b] \rightarrow Z$ is a quasigeodesic of Z if it is parametrized by arc length and, for every point $z \in Z$ satisfying (2-1) and every development $\tilde{\gamma} : [a, b] \rightarrow \mathbb{M}_k^2$ of γ with respect to $z \in Z$, the curvilinear triangle bounded by the segments $O\tilde{\gamma}(t \pm \delta)$ and the arc $\tilde{\gamma}|_{[t-\delta, t+\delta]}$, where $t \in (a, b)$ and $\delta > 0$ sufficiently small, is convex.

Given three points $x, y, z \in Z$, let $\bar{\Delta}_k(\bar{x}, \bar{y}, \bar{z})$ be a triangle in \mathbb{M}_k^2 satisfying

$$d_{\mathbb{M}_k^2}(\bar{x}, \bar{y}) = d_Z(x, y), \quad d_{\mathbb{M}_k^2}(\bar{y}, \bar{z}) = d_Z(y, z), \quad \text{and} \quad d_{\mathbb{M}_k^2}(\bar{z}, \bar{x}) = d_Z(z, x).$$

We denote by $\angle_k(x; y, z)$ the angle of $\bar{\Delta}_k(\bar{x}, \bar{y}, \bar{z})$ at \bar{x} . Note that this definition does not depend on the choice of the triangle $\bar{\Delta}_k(\bar{x}, \bar{y}, \bar{z})$.

Consider $z \in Z$ and $\lambda \in (0, \pi)$. The point z is said to be λ -strained if there exists a set $\{(a_i, b_i) \in Z \times Z \mid i \in \{1, \dots, n\}\}$, called a λ -strainer at z , such that $\angle_k(z; a_i, b_i) > \pi - \lambda$ and

$$\max \left\{ \left| \angle_k(z; a_i, a_j) - \frac{\pi}{2} \right|, \left| \angle_k(z; b_i, b_j) - \frac{\pi}{2} \right|, \left| \angle_k(z; a_i, b_j) - \frac{\pi}{2} \right| \right\} < \lambda$$

for all $i \neq j \in \{1, \dots, n\}$. The set $R_\lambda(Z)$ of λ -strained points of Z is called the *set of λ -regular points of Z* . It is a remarkable fact that $R_\lambda(Z)$ is an open and dense subset of Z .

We now recall the notion of (pointed) Hausdorff–Gromov convergence:

Definition 11 [Burago et al. 2001]. Let (Z_i, z_i) be a sequence of (pointed) metric spaces. We say that the sequence (Z_i, z_i) converges in the (pointed) Hausdorff–Gromov sense to a (pointed) metric space (Z, z_0) , if the following holds: For every $r > \varepsilon > 0$, there exist $i_0 \in \mathbb{N}$ and a sequence of (maybe noncontinuous) maps $f_i : B_{Z_i}(z_i, r) \rightarrow Z$ ($i > i_0$) such that

- (i) $f_i(z_i) = z_0$,
- (ii) $\sup \{ d_Z(f_i(z_1), f_i(z_2)) - d_Z(z_1, z_2) \mid z_1, z_2 \in Z \} < \varepsilon$,
- (iii) $B_Z(z_0, r - \varepsilon) \subset B_Z(f_i(B_{Z_i}(z_i, r)), \varepsilon)$,
- (iv) $f_i(B_{Z_i}(z_i, r)) \subset B_Z(z_0, r + \varepsilon)$.

For the rest of the paper, the term “converges” will stand for “converges in the (pointed) Hausdorff–Gromov sense”.

Let (Z_i, z_i) be a convergent sequence of Alexandrov spaces with the same lower curvature bound $k \in \mathbb{R}$ and the same dimension $n \in \mathbb{N}$. The limit Alexandrov space must have the same lower curvature bound k , but can have dimension less than or equal to n (see [Burago et al. 2001, Corollary 10.8.25]). When the limit Alexandrov space has dimension n , Perelman’s stability theorem (see [Kapovitch 2007]) assures that it is homeomorphic to Z_i , for sufficiently large indexes.

It is a fundamental fact that the class of Alexandrov spaces of curvature not smaller than $k \in \mathbb{R}$ is precompact with respect to the Hausdorff–Gromov convergence (see [Gromov 1981, Proposition 5.2] and [Burago et al. 2001, Corollary 10.8.25]). More precisely, every sequence of pointed Alexandrov spaces of curvature not smaller than $k \in \mathbb{R}$ admits a convergent subsequence to an Alexandrov space with the same lower bound for the curvature.

Another important fact concerning Alexandrov spaces is that the Hausdorff–Gromov limit of quasigeodesics is a quasigeodesic (see [Perelman and Petrunin 1994]). More precisely, if $\gamma_i : [a, b] \rightarrow Z_i$ is a convergent sequence of quasigeodesics, then the limit curve is a quasigeodesic on the limit space.

3. Sequences of hyperbolic cone manifolds

Let M be a closed, orientable and irreducible differential manifold of dimension 3 and let $\Sigma = \Sigma_1 \sqcup \cdots \sqcup \Sigma_l$ be an embedded link in M . A *hyperbolic cone structure* with topological type (M, Σ) is a complete intrinsic metric on M such that every nonsingular point (i.e., every point in $M - \Sigma$) has a neighborhood isometric to an open set of \mathbb{H}^3 , the hyperbolic space of dimension 3, and that every singular point (i.e., every point in Σ) has a neighborhood isometric to an open neighborhood of a singular point of $\mathbb{H}^3(\alpha)$, the space obtained by identifying the sides of a wedge of angle $\alpha \in (0, 2\pi]$ in \mathbb{H}^3 by a rotation about the axis of the wedge. The angles α are called *cone angles* and they may vary from one connected component of Σ to the other. We emphasize that we only allow cone angles of at most 2π in this paper. By convention, the complete hyperbolic structure M_0 on $M - \Sigma$ (see [Kojima 1996]) is considered as a hyperbolic cone structure with topological type (M, Σ) and cone angles equal to zero.

We point out that every hyperbolic cone manifold is an Alexandrov space of curvature not smaller than -1 . Furthermore, every geodesic on it is a quasigeodesic.

A natural way to study degenerating deformations of hyperbolic cone structures on (M, Σ) is to consider sequences of hyperbolic cone structures converging (in the pointed Hausdorff–Gromov sense) to the limit Alexandrov space. To study these kind of sequences, we need the important notion of collapse which illustrates the intuitive fact that the volume of the sequence may or may not go to zero.

Definition 12. We say that a sequence M_i of hyperbolic cone manifolds with topological type (M, Σ) collapses if, for every sequence of points $p_i \in M - \Sigma$, the sequence $r_{\text{inj}}^{M_i - \Sigma}(p_i)$ consisting of their Riemannian injectivity radii in $M_i - \Sigma$ converges to zero. Otherwise, we say that the sequence M_i does not collapse.

When a convergent sequence of hyperbolic cone manifolds collapses, most of the geometric information can be lost. This happens because the dimension of the limit Alexandrov space is strictly smaller than 3 (see [Barreto 2012]). On the noncollapsing case, however, the limit Alexandrov space must have dimension 3 and, in this case, many kinds of geometric information are preserved and can be used to study the deformation.

Given a sequence M_i of hyperbolic cone manifolds with topological type (M, Σ) , fix indices $i \in \mathbb{N}$ and $j \in \{1, \dots, l\}$. For sufficiently small radius $R > 0$, the metric

neighborhood

$$B_{M_i}(\Sigma_j, R) = \{x \in M_i \mid d_{M_i}(x, \Sigma_j) < R\}$$

of Σ is a solid torus embedded in M_i . The supremum of the radius $R > 0$ satisfying the above property will be called *normal injectivity radius of Σ_j in M_i* and it is going to be denoted by $R_i(\Sigma_j)$. Analogously we can define $R_i(\Sigma)$, the *normal injectivity radius of Σ* . It is a remarkable fact (see [Fujii 2000; Hodgson and Kerckhoff 2005]) that the existence of a uniform lower bound for $R_i(\Sigma)$ ensures the existence of a sequence of points $p_{i_k} \in M$ such that the sequence (M_{i_k}, p_{i_k}) converges to a pointed hyperbolic cone manifold (M_∞, p_∞) with topological type (M, Σ) . Moreover, M_∞ must be compact provided that the cone angles of M_{i_k} are uniformly bounded from below.

Let us also emphasize that the sequence $\text{Vol}(M_i)$ consisting of the Riemannian volumes of the hyperbolic manifolds $M_i - \Sigma$ is always uniformly bounded. More precisely (see [Dunfield 1999; Francaviglia 2004]), we have

$$(3-1) \quad \text{Vol}(M_i) < \text{Vol}(M_0),$$

where M_0 denotes the complete hyperbolic manifold that is homeomorphic to $M - \Sigma$.

The purpose of this section is to prove [Theorem 4](#). It is divided into two parts. The first part contains some preliminary results whereas the remaining part deals with the proof of [Theorem 4](#).

Let us point out that, throughout the rest of the paper, the term ‘‘component’’ is going to stand for ‘‘connected component’’.

3.1. Preliminary results. Let us recall some definitions and elementary results which will be important for the proof of [Theorem 4](#). We will begin with the classification of two-dimensional embedded tori in $M - \Sigma$ (see [Barreto 2012]).

Lemma 13. *Suppose that $M - \Sigma$ is hyperbolic and let T be a two-dimensional torus embedded in $M - \Sigma$. Then T separates M . Moreover, one and only one of the following statements holds:*

- (i) T is parallel to a component of Σ (hence it bounds a solid torus in M).
- (ii) T is not parallel to a component of Σ and it bounds a solid torus in $M - \Sigma$.
- (iii) T is not parallel to a component of Σ and it is contained in a ball B of $M - \Sigma$. Furthermore, T bounds a region in B which is homeomorphic to the exterior of a knot in S^3 .

We turn to the geometric classification of the thin part of a hyperbolic manifold.

Definition 14. Fix $\delta > 0$ and let M be a hyperbolic manifold of dimension 3 (without boundary and perhaps noncomplete). Define the δ -thin part $M_{\text{thin}}(\delta)$ of M by

$$M_{\text{thin}}(\delta) = \{q \in M \mid r_{\text{inj}}^M(q) < \delta \text{ and } \exp_q \text{ is defined on } B_{T_q M}(0, 3\delta)\}.$$

The following result concerning the thin part of hyperbolic manifolds will be needed later.

Proposition 15. *Let M be a hyperbolic manifold of dimension 3 (without boundary and perhaps noncomplete) of finite volume. If $\delta > 0$ is small enough, then each component of $M_{\text{thin}}(\delta)$ contains a maximal region which is isometric to either*

- (i) *the quotient of a metric neighborhood of a geodesic γ in \mathbb{H}^3 by a loxodromic element of $\text{PSL}_2(\mathbb{C})$ leaving γ invariant and whose translation length is not bigger than δ , or*
- (ii) *a parabolic cusp of rank 2.*

In addition, when $\text{Vol}(M) < \infty$, it follows that M has finitely many ends.

This proposition is a consequence of the existence of a Margulis foliation for the thin part of a hyperbolic manifold. A proof for this proposition is given in [Boileau et al. 2005, Theorem 5.3 and Corollary 5.5] where the authors study the thin part of hyperbolic cone manifolds with topological type (M, Σ) and whose cone angles are not bigger than π . Note that the condition imposed on the cone angles is used only in the description of the singular components of the thin part. We summarize below their proof for the first part of the proposition which, indeed, makes unnecessary the angle condition.

Consider a hyperbolic manifold M and denote by $\pi : \tilde{M} \rightarrow M$ the universal cover of M . Let $\delta > 0$ be the constant given by the Margulis lemma (see [Každan and Margulis 1968; Ballmann et al. 1985; Boileau et al. 2005]). Then for every component \mathcal{P} of $M_{\text{thin}}(\delta)$, the stabilizer of a component of $\pi^{-1}(\mathcal{P}) \subset \tilde{M}$ is an elementary subgroup of $\text{PSL}_2(\mathbb{C})$ generated by a loxodromic element or by at most two parabolic elements. Associated to this group we have a canonical foliation of \mathbb{H}^3 . The pull-back of this foliation by a developing map gives a foliation on $\pi^{-1}(\mathcal{P})$ which is equivariant by the action of $\pi_1 M$. The quotient of this foliation is the Margulis foliation on \mathcal{P} .

To finish the proof, it is sufficient to show that the leaves of this foliation are two-dimensional tori.

First, we remark that the leaves are complete. This is a consequence of the fact that injectivity radius is constant on them (see [Barreto 2009]). When the stabilizer of a component of $\pi^{-1}(\mathcal{P})$ is generated by a loxodromic element, the conclusion follows immediately. In the second case, we need to use the fact that the leaves are flat (they were obtained from horospheres) and the Gauss–Bonnet theorem. The hypothesis that the volume of the manifold is finite excludes undesirable euclidean surfaces other than torus.

3.2. Proof of Theorem 4. The purpose of this section is to study a noncollapsing sequence M_i . Without loss of generality, this hypothesis implies the existence of a

sequence $p_i \in M - \Sigma$ satisfying

$$r_0 = \inf\{r_{\text{inj}}^{M_i}(p_i) \mid i \in \mathbb{N}\} > 0,$$

and such that the sequence (M_i, p_i) converges to a pointed Alexandrov space (Z, z_0) . By definition of the pointed Hausdorff–Gromov convergence, the ball $B_Z(z_0, r_0)$ is isometric to a ball of radius r_0 in \mathbb{H}^3 and this implies that Z has dimension equal to 3.

We are interested in the case where the length of the singularity remains uniformly bounded, i.e., where

$$\sup\{\mathcal{L}_{M_i}(\Sigma_j) \mid i \in \mathbb{N} \text{ and } j \in \{1, \dots, l\}\} < \infty.$$

Fix $j \in \{1, \dots, l\}$. We can suppose (passing to a subsequence if necessary) that

$$\sup\{d_{M_i}(p_i, \Sigma_j) \mid i \in \mathbb{N}\} < \infty \quad \text{or} \quad \lim_{i \rightarrow \infty} d_{M_i}(p_i, \Sigma_j) = \infty.$$

In the first case, we can use again the precompactness to suppose that the component $\Sigma_j \subset M_i$, viewed as a sequence of Alexandrov spaces of dimension 1, converges to a closed curve Σ_j^Z in Z . Since Z has dimension 3 and it is the limit of a sequence of Alexandrov spaces with same dimension 3 and same lower curvature bound -1 , we can conclude that Σ_j^Z is a quasigeodesic in Z (see [Perelman and Petrunin 1994]).

Summarizing, each component Σ_j of Σ satisfies one, and only one, of the following statements:

- (1) $\sup\{d_{M_i}(p_i, \Sigma_j) \mid i \in \mathbb{N}\} < \infty$ and Σ_j converges to a quasigeodesic $\Sigma_j^Z \subset Z$.
- (2) $\lim_{i \rightarrow \infty} d_{M_i}(p_i, \Sigma_j) = \infty$.

This dichotomy allows us to write $\Sigma = \Sigma_0 \sqcup \Sigma_\infty$, where Σ_0 contains the components Σ_j of Σ which satisfy item (1) and Σ_∞ those that satisfy item (2).

The following lemma shows that the hypothesis of noncollapsing imposes restrictions on the length and on the cone angles of the singular components of Σ contained in Σ_0 .

Lemma 16. *Suppose that the sequence M_i does not collapse and let $p_i \in M - \Sigma$ be a sequence of points such that $r_0 = \inf\{r_{\text{inj}}^{M_i}(p_i) \mid i \in \mathbb{N}\} > 0$. If*

$$L = \sup\{\mathcal{L}_{M_i}(\Sigma_j) \mid i \in \mathbb{N} \text{ and } j \in \{1, \dots, l\}\} < \infty,$$

the following inequalities hold:

- (i) $\inf\{\mathcal{L}_{M_i}(\Sigma_j) \mid i \in \mathbb{N} \text{ and } \Sigma_j \subset \Sigma_0\} > 0$.
- (ii) $\inf\{\alpha_{ij} \mid i \in \mathbb{N} \text{ and } \Sigma_j \subset \Sigma_0\} > 0$.
- (iii) $\sup\{R_i(\Sigma_j) \mid i \in \mathbb{N} \text{ and } \Sigma_j \subset \Sigma_0\} < \infty$.

Proof. Consider $\mathcal{R} > \sup\{d_{M_i}(p_i, \Sigma_j) \mid i \in \mathbb{N} \text{ and } \Sigma_j \subset \Sigma_0\} + r_0$. By construction, $\mathcal{R} < \infty$ and $B_{M_i}(p_i, r_0) \subset B_{M_i}(\Sigma_j, \mathcal{R})$, for all $i \in \mathbb{N}$ and all components Σ_j of Σ_0 .

Fix $i \in \mathbb{N}$ and fix a component Σ_j of Σ_0 . Let \mathcal{A} be a region of $\mathbb{H}^3(\alpha_{ij})$ which is bounded by two planes orthogonal to the singular geodesic σ of $\mathbb{H}^3(\alpha_{ij})$ and having distance $\mathcal{L}_{M_i}(\Sigma_j)$ between them. Using a developing map for $M_i - \Sigma$ and the minimizing geodesics leaving Σ_j orthogonally, the manifold M_i can be developed in a compact domain $K \subset \mathcal{A}$ such that $\text{Vol}(K) = \text{Vol}(M_i)$.

Since $B_{M_i}(p_i, r_0) \subset B_{M_i}(\Sigma_j, \mathcal{R})$, the development of $B_{M_i}(p_i, r_0)$ in K is contained in $B_{\mathbb{H}^3(\alpha_{ij})}(\sigma, \mathcal{R}) \cap \mathcal{A}$. If V_0 represents the volume of a ball of radius r_0 in \mathbb{H}^3 , we have

$$V_0 = \text{Vol}(B_{M_i}(p_i, r_0)) \leq \text{Vol}(B_{\mathbb{H}^3(\alpha_{ij})}(\sigma, \mathcal{R}) \cap \mathcal{A}) = \frac{\alpha_{ij}}{2} \mathcal{L}_{M_i}(\Sigma_j) \sinh^2(\mathcal{R})$$

and therefore

$$\mathcal{L}_{M_i}(\Sigma_j) \geq \frac{V_0}{\pi \sinh^2(\mathcal{R})} > 0 \quad \text{and} \quad \alpha_{ij} \geq \frac{2V_0}{L \sinh^2(\mathcal{R})} > 0.$$

Finally, item (iii) follows from the fact that the sequence $\text{Vol}(M_i)$ is uniformly bounded from above (see (3-1)). \square

With the preceding notations, set

$$\Sigma_Z = \bigcup_{\Sigma_j \subset \Sigma_0} \Sigma_j^Z \subset Z.$$

We present now the main result for the noncollapsing:

Theorem 17 (noncollapsing). *Suppose that there exists a sequence $p_i \in M - \Sigma$ satisfying*

$$r_0 = \inf\{r_{\text{inj}}^{M_i}(p_i) \mid i \in \mathbb{N}\} > 0$$

and such that the sequence (M_i, p_i) converges to a pointed Alexandrov space (Z, z_0) of dimension 3. If

$$\sup\{\mathcal{L}_{M_i}(\Sigma_j) \mid i \in \mathbb{N} \text{ and } j \in \{1, \dots, l\}\} < \infty.$$

Then:

- (i) $Z - \Sigma_Z$ is a hyperbolic 3-manifold of finite volume whose convex and unbounded ends are finite in number and are parabolic cusps of rank 2.
- (ii) Z is compact (and therefore homeomorphic to M) if and only if $\Sigma_\infty = \emptyset$.
- (iii) If Z is not compact, there is a bijection between the connected components of Σ_∞ and the complete ends of $Z - \Sigma_Z$. In fact, each unbounded end C_j of $Z - \Sigma_Z$ is the Hausdorff-Gromov limit of metric neighborhoods (homeomorphic to solid tori) $B_{M_i}(\Sigma_j, r_i)$ of a component Σ_j of Σ_∞ , where $r_i > 0$ is an

increasing sequence going off to infinity. In addition, the cone angles α_{ij} and the lengths of these components converge to 0.

Proof of (i). According to [Fujii 2000, Lemma 2], every point of $Z - \Sigma_Z$ is the limit of a sequence of points of $M_i - \Sigma$ whose injectivity radius is uniformly bounded from below. This implies that $Z - \Sigma_Z$ is a (without boundary and noncomplete) hyperbolic manifold. Note that the unbounded ends of Z are those of $Z - \Sigma_Z$. In view of Proposition 15, to prove item (i) it is sufficient to show the following:

Claim. $\text{Vol}(Z - \Sigma_Z) < \infty$.

Proof of claim. Suppose for contradiction the statement is false. Let K_∞ be a compact set of $Z - \Sigma_Z$ whose Riemannian volume is strictly greater than $\text{Vol}(M_0)$, where M_0 is $M - \Sigma$ with its complete hyperbolic metric. Since the convergence is bilipschitz on compact subsets (see [Cooper et al. 2000, Theorem 6.20]), there exists an index $i_0 \in \mathbb{N}$ and a compact subset K_{i_0} of $M_{i_0} - \Sigma$ (near K_∞) such that

$$\text{Vol}(M_0) < \text{Vol}_{M_{i_0}}(K_{i_0}) \leq \text{Vol}(M_{i_0}).$$

This is however impossible since $\text{Vol}(M_{i_0}) < \text{Vol}(M_0)$ (see (3-1)). This proves the claim, and thus completes the proof of item (i) of Theorem 17. \square

Proof of (ii) and (iii). If Z is compact then $\Sigma_\infty = \emptyset$. Suppose now that Z is not compact. By Lemma 16 we can choose $R > 0$ such that

$$B_{M_i}(\Sigma_j, R_i(\Sigma_j)) \subset B_{M_i}(p_i, R/2)$$

for all connected components Σ_j of Σ_0 and all $i \in N$. Let K be a compact subset of Z which contains the ball $B_Z(z_0, R)$ (and hence Σ_Z) in its interior and satisfies

$$\mathcal{E} = Z - \text{int}(K) = C_1 \sqcup \cdots \sqcup C_m,$$

where each $C_k \approx T^2 \times [0, \infty)$ is a cuspidal end of Z .

Consider a sequence $C_{1i} = T^2 \times [0, t_i]$ of compact subsets of C_1 , where $t_i > 0$ is an unbounded and strictly increasing sequence.

Let $\varepsilon_i > 0$ be a sequence converging to zero. Without loss of generality, there exists (according to [Cooper et al. 2000, Theorem 6.20]) a sequence of $(1 + \varepsilon_i)$ -bilipschitz embeddings $f_{1i} : C_{1i} \rightarrow M_i - \Sigma$ onto their images. Therefore, the sequence $B_{1i} = f_{1i}(C_{1i})$ converges in the bilipschitz sense to the compact set C_{11} .

Consider now a sequence of holonomy representations $\zeta_{1i} : \mathbb{Z} \times \mathbb{Z} \rightarrow \text{PSL}_2(\mathbb{C})$ for the hyperbolic structures on the interior sets B_{1i} . According to [Cooper et al. 2000, Theorem 6.22], we can assume that

$$(3-2) \quad \zeta_{1i} \circ (f_{1i})_* \longrightarrow \varphi_1,$$

where $\varphi_1 : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathrm{PSL}_2(\mathbb{C})$ is a holonomy representation of the hyperbolic structure in the interior of C_1 and where $(f_{1i})_* : \mathbb{Z} \times \mathbb{Z} \rightarrow \pi_1(M - \Sigma)$ is the canonical homomorphism induced by the map f_{1i} .

Consider the torus $T_{1i} = f_{1i}(T^2 \times \{0\})$ embedded in $M - \Sigma$. Since K contains the ball $B_Z(z_0, R)$, the torus T_{1i} cannot be parallel to a component Σ_j of Σ_0 . For i sufficiently large, the torus T_{1i} cannot be contained in a ball of $M - \Sigma$. To see this, consider a homotopically nontrivial loop γ_1 on $T^2 \times \{0\} \subset C_{11}$. Since C_1 is a parabolic cusp, $\varphi_1(\gamma_1)$ is a nontrivial parabolic element of $\mathrm{PSL}_2(\mathbb{C})$ and therefore the convergence (3-2) implies that $\zeta_{1i} \circ (f_{1i})_*(\gamma_1)$ is not trivial for i very large. The same then holds for the sequence $(f_{1i})_*(\gamma_1)$.

According to Lemma 13, we can suppose that the torus T_{1i} bounds a solid torus W_{1i} in M (with perhaps a singular soul). Note that

$$(3-3) \quad \lim_{i \rightarrow \infty} \mathrm{diam}_{M_i}(W_{1i}) = \infty,$$

because $f_{1i}(C_{1i}) \subset W_{1i}$, for all $i \in \mathbb{N}$.

We can repeat the same construction for each cusp C_k of \mathcal{L} in order to obtain sequences of embedded tori $T_{ki} \subset M - \Sigma$ ($k \in \{1, \dots, m\}$ and $i \in \mathbb{N}$), each of them bounds solid torus W_{ki} in $M - \Sigma_0$. Furthermore whose diameters become infinite with i . This yields a sequence of 3-manifolds with torus boundary

$$\mathcal{M}_i = M_i - \bigcup_{k=1}^m W_{ki}$$

such that M can be obtained by Dehn filling on their boundary components. By construction, the sequence \mathcal{M}_i converges to the compact K and then (by Perelman's stability theorem [Kapovitch 2007]), we can assume that the manifolds \mathcal{M}_i are all homeomorphic to K .

For all $i \in \mathbb{N}$ and all $k \in \{1, \dots, m\}$, fix a homotopically nontrivial loop μ_{ki} in $T^2 \times \{0\} \subset C_k$ satisfying:

- the loop $f_{ki} \circ \mu_{ki}$ bounds a disc in W_{ki} ,
- if, for some index $j \in \mathbb{N}$, a loop μ_{kj} belongs to the same homotopy class of the loop μ_{ki} , then $\mu_{kj} = \mu_{ki}$.

The rest of the proof is going to be divided in two cases depending on whether or not Σ_0 is empty.

First case: $\Sigma_0 = \emptyset$. Since the link Σ was supposed to be nonempty, it follows that $\Sigma_\infty \neq \emptyset$. Since the distance between p_i and Σ_∞ becomes infinite, we can assume that Σ_∞ is contained in the complement of \mathcal{M}_i . More precisely, we can also assume (see Lemma 13) that each solid torus of $M_i - \mathcal{M}_i$ contains at most one component

of Σ_∞ and, in the latter case, this component corresponds to the soul of the solid torus in question.

The singular set Σ_∞ has a finite number of components. Passing to a subsequence if necessary, we obtain an one-to-one map which associates each component Σ_j of Σ_∞ to a component C_{k_j} of \mathcal{L} , that is, the component Σ_j is contained in the component $W_{k_j i}$ of $M_i - \mathcal{M}_i$, for all $i \in \mathbb{N}$.

Recall that $\lim_{i \rightarrow \infty} d_{M_i}(p_i, \Sigma_j) = \infty$ for every connected component Σ_j of Σ_∞ . Since the tori $T_{k_j i}$ remain at a finite distance to the points p_i and they are parallel to the components Σ_j , we must have $\lim_{i \rightarrow \infty} R_i(\Sigma_j) = \infty$.

Since $\Sigma_0 = \emptyset$ and thanks to [Fujii 2000, Theorem 1], the cone angles of Σ converge to zero and Z has a complete hyperbolic structure whose ends are associated with components of Σ_∞ . In other words, the injection defined above between the components of Σ_∞ and the components of \mathcal{L} is, indeed, a bijection.

Second case: $\Sigma_0 \neq \emptyset$. Denote by Λ the subset of $\{1, \dots, m\}$ containing the indices that are not associated with components of Σ_∞ . Denote also by Ω the subset of $\{1, \dots, m\}$ containing the indices that are associated with components of Σ_∞ whose sequence of cone angles does not converge to zero.

Lemma 18. *There exist $i_0 \in \mathbb{N}$ satisfying: for each $k \in \Lambda \cup \Omega$, the homotopy classes of loops μ_{ki} ($i > i_0$) are pairwise distinct.*

Proof. Suppose for a contradiction that the statement of the lemma does not hold. Without loss of generality, there exists $k_0 \in \Lambda \cup \Omega$ such that all loops $\mu_{k_0 i}$ ($i \in \mathbb{N}$) belongs to the same homotopy class. By construction, this implies that the loops $\mu_{k_0 i}$ ($i \in \mathbb{N}$) are the same loop, say μ .

Suppose first that $k_0 \in \Lambda$. By construction,

$$(3-4) \quad \zeta_{k_0 i} \circ (f_{k_0 i})_*(\mu) = \zeta_{k_0 i}(f_{k_0 i} \circ \mu) = 1_{\text{PSL}_2(\mathbb{C})},$$

for all $i \in \mathbb{N}$. Because $\varphi_{k_0}([\mu])$ is a nontrivial parabolic element of $\text{PSL}_2(\mathbb{C})$, we have a contradiction.

Suppose now that $k_0 \in \Omega$. Then $k_0 = k_j$, for some component Σ_j of Σ_∞ whose sequence of cone angles converges to $\alpha_{\infty j} \neq 0$. Since the maps $f_{k_0 i}$ are $(1 + \varepsilon_i)$ -bilipschitz embeddings (with ε_i shrinks down to zero), the loops $f_{k_0 i} \circ \mu$ must have bounded lengths.

As noted in the preceding case, the sequence $R_i(\Sigma_j)$ of the normal injectivity radii of the component Σ_j goes off to infinity. Since $\alpha_{\infty j} \neq 0$, the sequence $\mathcal{L}_{M_i}(f_{k_0 i} \circ \mu)$ formed by the lengths of the loops $f_{k_0 i} \circ \mu$ cannot be bounded. This is a contradiction with the above paragraph. \square

As a consequence of the above lemma, we will show that the set $\Lambda \cup \Omega$ is empty. To do this, the following lemma will be needed:

Lemma 19. *Given $k \in \Lambda$, there exists $i_0 = i_0(k) \in \mathbb{N}$ such that the solid tori W_{ki} contains a simple closed geodesic σ_{ki} , for every $i > i_0$.*

Proof. Fix $k \in \Lambda$ and let

$$\delta = \frac{1}{2} \inf \{ r_{\text{inj}}^{Z-\Sigma_Z}(z) \mid z \in C_{k1} \} > 0.$$

Since the map $f_{ki}|_{C_{k1}} : C_{k1} \rightarrow B_{ki}$ becomes closer and closer to isometries, there exists $i_1 \in \mathbb{N}$ such that

$$r_{\text{inj}}^{M_i}(q) > \delta,$$

for all $i > i_1$ and for all $q \in B_{ki}$ (in particular, for all $q \in T_{ki}$).

Claim. *There is $i_2 \in \mathbb{N}$ such that, for all $i > i_2$, we can find a loop γ_{ki} in W_{ki} which is homotopically nontrivial in the interior $M - \Sigma$ and has length smaller than δ .*

Proof of claim. Consider the loops consisting of two geodesic segments with same ends and equal lengths which, furthermore, are smaller than $\delta/2$. These loops are always homotopically nontrivial; otherwise we would obtain, after development, two distinct geodesic arcs with the same ends and equal lengths in \mathbb{H}^3 , which is not possible.

The fact that W_{ki} does not admit this type of loop in its interior is equivalent to saying that all points of W_{ki} have injectivity radius not smaller than $\delta/2$. This is a contradiction because the sequence $\text{Vol}(M_i)$ is uniformly bounded from above (see (3-1)) and the diameter of components W_{ki} becomes infinite. This proves the claim.

Consider $i_o = \max\{i_1, i_2\}$ and fix $i > i_o$. Let $\gamma_{ki} \subset W_{ki}$ be a loop as above. By [Kojima 1998, Lemma 1.2.4], the loop γ_{ki} is freely homotopic (in $M - \Sigma$) to a closed geodesic $\sigma_{ki} \subset M - \Sigma$. Moreover, the length of σ_{ki} is smaller than δ because the length of loops is strictly decreasing along this homotopy. Because the points of the torus T_{ki} have injectivity radius bigger than δ , all the loops involved in this homotopy must lie entirely in the interior of W_{ki} . In particular, $\sigma_{ki} \subset W_{ki}$.

If σ_{ki} is not simple, then it gives rise to a loop γ'_{ki} consisting of two geodesic segments with same ends and equal lengths which are smaller than $\delta/4$. This implies that the injectivity radius of the ends of γ'_{ki} is smaller than $\delta/4$. We can apply the same construction for the loop γ'_{ki} in order to obtain a new closed geodesic $\sigma_{ki} \subset W_{ki}$ whose length is smaller than $\delta/4$. Since the injectivity radius of points of W_{ki} bounded from below by compactness, this process must end after a finite number of steps and therefore we can suppose that σ_{ki} is simple. This completes the proof of Lemma 19. \square

The following lemma shows that Σ_∞ is not empty and the cone angles of its components goes to zero. Moreover the map between the components of Σ_∞ and the components of \mathcal{L} must be a bijection.

Lemma 20. *The set $\Lambda \cup \Omega$ is empty.*

Proof. According to the above lemma, we can suppose there exists a simple closed geodesic σ_{ki} in the solid torus W_{ki} , for every $i \in \mathbb{N}$ and every $k \in \Lambda$. If the manifolds M_i are regarded as hyperbolic cone manifolds with topological type (M, Σ') , where

$$\Sigma' = \Sigma \cup \bigcup_{k \in \Lambda} \sigma_{ki}$$

and the cone angles on the geodesics σ_{ki} are equal to 2π , it follows from [Lemma 13](#) that the tori T_{ki} are parallel to the geodesics σ_{ki} . In addition, $M - \Sigma'$ admits a complete hyperbolic structure (see [\[Kojima 1996\]](#)) that will be denoted by \mathcal{M}_0 .

For all $i \in \mathbb{N}$ and all $k \in \Lambda$, denote the homotopy class of the loop μ_{ik} by $(p_{ki}, q_{ki}) \in \mathbb{Z} \times \mathbb{Z} \approx \pi_1 C_k$. Without loss of generality, the Thurston's hyperbolic Dehn surgery [\[Cooper et al. 2000, Theorem 1.13\]](#) gives a sequence of complete hyperbolic manifolds $\mathcal{M}(p_{i1}, q_{i1}, \dots, p_{im}, q_{im})$ diffeomorphic to $M - \Sigma$ and such that

$$(3-5) \quad V_i := \text{Vol}(\mathcal{M}(p_{i1}, q_{i1}, \dots, p_{im}, q_{im})) < \text{Vol}(\mathcal{M}_0),$$

where $(p_{ki}, q_{ki}) = \infty$, for all $i \in \mathbb{N}$ and all $k \in \{1, \dots, m\} - \Lambda$.

Since, for each $k \in \Lambda$, the pairs $(p_{ki}, q_{ki})_{i \in \mathbb{N}}$ are pairwise distinct (the homotopy classes of μ_{ik} are pairwise distinct), a subsequence $\mathcal{M}(p_{1i_s}, q_{1i_s}, \dots, p_{mi_s}, q_{mi_s})$ such that

$$\lim_{s \rightarrow \infty} \|(p_{ki_s}, q_{ki_s})\| = \lim_{s \rightarrow \infty} (p_{ki_s})^2 + (q_{ki_s})^2 = \infty \quad \text{for every } k \in \Lambda$$

always exists. Thurston's hyperbolic Dehn surgery then gives

$$(3-6) \quad \lim_{s \rightarrow \infty} V_{i_s} = \text{Vol}(\mathcal{M}_0).$$

Recall that the Riemannian volume of a complete hyperbolic manifold with finite volume is a topological invariant (Mostow's theorem). Since the manifolds $\mathcal{M}(p_{i1}, q_{i1}, \dots, p_{im}, q_{im})$ are diffeomorphic, the sequence V_i must be constant. This contradicts the statements (3-5) and (3-6). Hence $M_i - \mathcal{M}_i$ cannot have nonsingular components. Therefore, $\Sigma_\infty \neq \emptyset$ and the map between the components of Σ_∞ and the components of \mathcal{Z} is a bijection. This proves [Lemma 20](#), and thus completes the proof of items (ii) and (iii) of [Theorem 17](#). \square

Corollary 21. *Suppose that the sequence M_i does not collapse and verifies*

$$\sup\{\mathcal{L}_{M_i}(\Sigma_j) \mid i \in \mathbb{N} \text{ and } j \in \{1, \dots, l\}\} < \infty.$$

If there is $\varepsilon \in (0, 2\pi)$ such that the cone angles α_{ij} belong to $(\varepsilon, 2\pi]$, then there exists a sequence of points $p_{ik} \in M - \Sigma$ such that the sequence (M_{i_k}, p_{i_k}) converges

to a compact and 3-dimensional pointed Alexandrov space (Z, z_0) (in fact homeomorphic to M). Moreover, there exists a finite union of quasigeodesics Σ_Z such that $Z - \Sigma_Z$ is a noncomplete hyperbolic manifold of finite volume.

Remark 22. Suppose that Σ is not connected. If (M_i, p_i) is a sequence as in the statement of [Theorem 17](#), then the inequality

$$\sup\{\text{diam}_{M_i}(\Sigma) \mid i \in \mathbb{N}\} < \infty$$

is a necessary and sufficient condition to ensure that the sequence $\text{diam}(M_i)$ remains bounded.

We have also the following less immediate corollary:

Corollary 23. *Let M be a closed, orientable and irreducible 3-manifold and let Σ be an embedded link in M . Assume that there exists a sequence M_i of hyperbolic cone manifolds with topological type (M, Σ) and having the same cone angles $\alpha_i \in (0, 2\pi]$ for all components of Σ . Then there is a pointed subsequence M_{i_k} converging to M_0 ($M - \Sigma$ with its complete hyperbolic metric) if and only if the following conditions hold:*

- (i) $\sup\{\mathcal{L}_{M_i}(\Sigma) \mid i \in \mathbb{N}\} < \infty$.
- (ii) $\sup\{\text{diam}(M_i) \mid i \in \mathbb{N}\} = \infty$.
- (iii) *The sequence M_i does not collapse.*

Proof. By Kojima's result [\[1998\]](#), the existence of a subsequence M_{i_k} converging to M_0 is equivalent to the convergence of the cone angles α_{i_k} to zero.

Suppose that the sequence α_i converges to zero. Without loss of generality, we can assume that $\alpha_i \in (0, \pi]$, for every $i \in \mathbb{N}$. According to [\[Kojima 1998\]](#), there exists a continuous path (parametrized by cone angles) of hyperbolic cone structures with topological type (M, Σ) which connects the hyperbolic cone structure of M_0 to the complete hyperbolic structure on $M - \Sigma$. Moreover, by uniqueness of the hyperbolic cone structures with cone angles not bigger than π (see [\[Kojima 1998\]](#)), this path contains the hyperbolic cone structures of M_i , for every $i \in \mathbb{N}$. Then for every point $p \in M$, the sequence (M_i, p) converges to $(M - \Sigma, p)$ with the complete hyperbolic structure. This implies items (ii) and (iii). Item (i) is a consequence of Thurston's hyperbolic Dehn surgery theorem which implies that the sequence $\mathcal{L}_{M_i}(\Sigma)$ converges to zero.

Conversely, suppose now that items (i), (ii) and (iii) are true. Then there exists a sequence of points $p_{i_k} \in M - \Sigma$ satisfying

$$\inf\{r_{\text{inj}}^{M_i}(p_{i_k}) \mid k \in \mathbb{N}\} > 0$$

and such that the sequence (M_{i_k}, p_{i_k}) converges to a noncompact and 3-dimensional pointed Alexandrov space (Z, z_0) . [Corollary 21](#) then shows that the sequence α_i must converge to zero. \square

4. Applications

4.1. Small links. An embedded link Σ in a 3-manifold M is called small (in M) if it has an open tubular neighborhood U such that $M - U$ does not contain an embedded essential surface whose boundary is empty or an union of meridians of Σ . An important fact due to W. Thurston and A. Hatcher [[1985](#), Lemma 3] is that every 3-manifold containing a small link does not admit an embedded essential surface.

Given a 3-manifold M , let Σ be an embedded link in M . Suppose there exists a sequence M_i of hyperbolic cone manifolds with topological type (M, Σ) and consider the sequence $\mathcal{L}_{M_i}(\Sigma)$ formed by the lengths of the singular set Σ in M_i . As a consequence of the Culler–Shalen theory [[1983](#)], the holonomy representations of M_i are convergent. Therefore, we have the following proposition:

Proposition 24. *Let M_i be a sequence of hyperbolic cone manifolds with topological type (M, Σ) . If Σ is a small link in M , then*

$$\sup\{\mathcal{L}_{M_i}(\Sigma_j) \mid i \in \mathbb{N} \text{ and } \Sigma_j \text{ component of } \Sigma\} < \infty.$$

When Σ is a small link in M , [Theorem 4](#) yields the following corollaries:

Corollary 25. *Suppose that M is a closed, orientable, irreducible and non-Seifert fibered 3-manifold and let Σ be an embedded small link in M . Then there exists a constant $V = V(M, \Sigma) > 0$ such that $\text{Vol}(\mathcal{M}) > V$, for every hyperbolic cone manifold \mathcal{M} with topological type (M, Σ) and having cone angles of at most 2π .*

Proof. First note that M is not a Sol manifold. In fact every Sol manifold is foliated by essential two-dimensional tori and this is not possible since Σ is small (see [[Hatcher and Thurston 1985](#), Lemma 3]).

Suppose that the lower bound V does not exist. Since Σ is small in M , the nonexistence of V implies the existence of a sequence of hyperbolic cone manifolds \mathcal{M}_i with topological type (M, Σ) satisfying

- $\sup\{\mathcal{L}_{\mathcal{M}_i}(\Sigma_j) \mid i \in \mathbb{N} \text{ and } \Sigma_j \text{ component of } \Sigma\} < \infty$,
- the sequence $\text{Vol}(\mathcal{M}_i - \Sigma)$ formed by the Riemannian volumes of the hyperbolic manifolds $\mathcal{M}_i - \Sigma$ shrinks down to zero (and therefore the sequence \mathcal{M}_i collapses).

According to [Theorem 4](#), M must be Seifert fibered, contradicting our hypothesis. \square

Corollary 26. *Suppose that M is a closed, orientable, irreducible and non-Seifert fibered 3-manifold and let Σ be an embedded small link in M . Given $\varepsilon \in (0, 2\pi)$, there is a constant $K = K(M, \varepsilon) > 0$ such that $\text{diam}(\mathcal{M}) < K$, for every hyperbolic cone manifold \mathcal{M} with topological type (M, Σ) and having cone angles belonging to $(\varepsilon, 2\pi]$.*

Proof. As seen in the previous corollary, M is not a Sol manifold. Fix $\varepsilon \in (0, 2\pi)$ and suppose that the upper bound K does not exist. Since Σ is small in M , the nonexistence of K implies the existence of a sequence of hyperbolic cone manifolds \mathcal{M}_i with topological type (M, Σ) , having cone angles $\alpha_{ji} \in (\varepsilon, 2\pi]$ and satisfying these conditions:

- (i) $\sup\{\mathcal{L}_{\mathcal{M}_i}(\Sigma_j) \mid i \in \mathbb{N} \text{ and } \Sigma_j \text{ component of } \Sigma\} < \infty$.
- (ii) The sequence $\text{diam}(\mathcal{M}_i)$ formed by the diameters of the hyperbolic cone manifolds \mathcal{M}_i go to infinity.

Since M is neither Seifert fibered nor a Sol manifold, it follows from item (i) and [Theorem 4](#) that the sequence \mathcal{M}_i does not collapse. Moreover, since the cone angles α_{ji} belong to $(\varepsilon, 2\pi]$, it follows that the sequence $\text{diam}(\mathcal{M}_i)$ is bounded and this yields a contradiction with item (ii). □

4.2. Proof of Corollary 6. First, we would like to recall that the existence of a deformation M_α as in [Corollary 6](#) is a consequence of the local deformation theorem due to [\[Hodgson and Kerckhoff 1998\]](#).

Proof. The implication (i) \Rightarrow (ii) is immediate (see [\[Kojima 1998\]](#)). Suppose now that the sequence $\mathcal{L}_{M_\alpha}(\Sigma)$ converges to 0 when α converges to θ . Then

$$\sup\{\mathcal{L}_{M_{\alpha_i}}(\Sigma_j) \mid i \in \mathbb{N} \text{ and } \Sigma_j \text{ component of } \Sigma\} < \infty,$$

for every sequence $\alpha_i \in (\theta, 2\pi]$ converging to θ . Consider such a sequence α_i . Since M is hyperbolic (and therefore is neither Seifert fibered nor a Sol manifold), it follows from [Theorem 4](#) that the sequence M_{α_i} does not collapse. Moreover, since the sequence $\mathcal{L}_{M_{\alpha_i}}(\Sigma)$ converges to zero, we must have $\lim_{i \rightarrow \infty} \text{diam}(M_{\alpha_i}) = \infty$. This concludes the proof of the implication (ii) \Rightarrow (iii).

To prove (iii) \Rightarrow (i) take a sequence α_i satisfying item (iii). Again by [Theorem 4](#), it follows that the sequence M_{α_i} does not collapse. Moreover, since the sequence $\text{diam}(M_{\alpha_i})$ is not bounded, we must have $\theta = 0$ because all the components of Σ have the same cone angle. Then, by [\[Kojima 1998\]](#), it follows that M_i converges to M_0 . □

References

[Ballmann et al. 1985] W. Ballmann, M. Gromov, and V. Schroeder, *Manifolds of nonpositive curvature*, Progress in Mathematics **61**, Birkhäuser, Boston, 1985. MR 87h:53050 Zbl 0591.53001

- [Barreto 2009] A. P. Barreto, *Déformations de structures hyperboliques coniques*, thesis, Université de Toulouse, 2009, Available at http://thesesups.ups-tlse.fr/838/1/Paiva-Barreto_Alexandre.pdf.
- [Barreto 2012] A. P. Barreto, “Deformation of hyperbolic cone-structures: study of the collapsing case”, preprint, 2012. [arXiv 1201.2923](https://arxiv.org/abs/1201.2923)
- [Boileau et al. 2005] M. Boileau, B. Leeb, and J. Porti, “Geometrization of 3-dimensional orbifolds”, *Ann. of Math. (2)* **162**:1 (2005), 195–290. [MR 2007f:57028](#) [Zbl 1087.57009](#)
- [Burago et al. 1992] Y. Burago, M. Gromov, and G. Perel’man, “пространства А. Д. Александрова с ограниченными снизу кривизнами”, *Uspekhi Mat. Nauk* **47**:2(284) (1992), 3–51. Translated as “A. D. Alexandrov spaces with curvature bounded below” in *Russ. Math. Surv.* **47**:2 (1992), 1–58. [MR 93m:53035](#) [Zbl 0802.53018](#)
- [Burago et al. 2001] D. Burago, Y. Burago, and S. Ivanov, *A course in metric geometry*, Graduate Studies in Mathematics **33**, Amer. Math. Soc., Providence, RI, 2001. [MR 2002e:53053](#) [Zbl 0981.51016](#)
- [Cooper et al. 2000] D. Cooper, C. D. Hodgson, and S. P. Kerckhoff, *Three-dimensional orbifolds and cone-manifolds*, MSJ Memoirs **5**, Math. Soc. Japan, Tokyo, 2000. [MR 2002c:57027](#) [Zbl 0955.57014](#)
- [Culler and Shalen 1983] M. Culler and P. B. Shalen, “Varieties of group representations and splittings of 3-manifolds”, *Ann. of Math. (2)* **117**:1 (1983), 109–146. [MR 84k:57005](#) [Zbl 0529.57005](#)
- [Dunfield 1999] N. M. Dunfield, “Cyclic surgery, degrees of maps of character curves, and volume rigidity for hyperbolic manifolds”, *Invent. Math.* **136**:3 (1999), 623–657. [MR 2000d:57022](#) [Zbl 0928.57012](#)
- [Francaviglia 2004] S. Francaviglia, “Hyperbolic volume of representations of fundamental groups of cusped 3-manifolds”, *Int. Math. Res. Not.* **2004**:9 (2004), 425–459. [MR 2004m:57032](#) [Zbl 1088.57015](#)
- [Fujii 2000] M. Fujii, “A cone angle condition on strong convergence of hyperbolic 3-cone-manifolds”, *RIMS Kōkyūroku* **1163** (2000), 126–131. [MR 1799375](#) [Zbl 0969.57509](#)
- [Gromov 1981] M. Gromov, *Structures métriques pour les variétés Riemanniennes*, Textes Mathématiques **1**, CEDIC, Paris, 1981. [MR 85e:53051](#) [Zbl 0509.53034](#)
- [Hatcher and Thurston 1985] A. Hatcher and W. Thurston, “Incompressible surfaces in 2-bridge knot complements”, *Invent. Math.* **79**:2 (1985), 225–246. [MR 86g:57003](#) [Zbl 0602.57002](#)
- [Hodgson and Kerckhoff 1998] C. D. Hodgson and S. P. Kerckhoff, “Rigidity of hyperbolic cone-manifolds and hyperbolic Dehn surgery”, *J. Differential Geom.* **48**:1 (1998), 1–59. [MR 99b:57030](#) [Zbl 0919.57009](#)
- [Hodgson and Kerckhoff 2005] C. D. Hodgson and S. P. Kerckhoff, “Universal bounds for hyperbolic Dehn surgery”, *Ann. of Math. (2)* **162**:1 (2005), 367–421. [MR 2006g:57031](#) [Zbl 1087.57011](#)
- [Kapovitch 2007] V. Kapovitch, “Perelman’s stability theorem”, pp. 103–136 in *Metric and comparison geometry*, edited by J. Cheeger and K. Grove, Surv. Differ. Geom. **11**, International Press, Somerville, MA, 2007. [MR 2009g:53057](#) [Zbl 1151.53038](#)
- [Každan and Margulis 1968] D. A. Každan and G. A. Margulis, “Доказательство гипотезы Сельберга”, *Mat. Sb. (N.S.)* **75**(117):1 (1968), 163–168. Translated as “A proof of Selberg’s hypothesis” in *Math. USSR Sb.* **4**:1 (1968), 147–152. [MR 36 #6535](#)
- [Kojima 1996] S. Kojima, “Nonsingular parts of hyperbolic 3-cone-manifolds”, pp. 115–122 in *Topology and Teichmüller spaces* (Katinkulta, 1995), edited by S. Kojima et al., World Scientific, River Edge, NJ, 1996. [MR 99i:57028](#) [Zbl 0928.57010](#)
- [Kojima 1998] S. Kojima, “Deformations of hyperbolic 3-cone-manifolds”, *J. Differential Geom.* **49**:3 (1998), 469–516. [MR 2000d:57023](#) [Zbl 0990.57004](#)

[Perelman and Petrunin 1994] G. Perelman and A. Petrunin, “[Quasigeodesics and gradient curves in Alexandrov spaces](#)”, 1994, Available at http://www.math.psu.edu/petrunin/papers/qg_ams.pdf.

Received January 19, 2012. Revised October 23, 2013.

ALEXANDRE PAIVA BARRETO
DEPARTAMENTO DE MATEMÁTICA
UNIVERSIDADE FEDERAL DE SÃO CARLOS
ROD. WASHINGTON LUÍS, KM 235
C.P. 676
13565-905 SÃO CARLOS
BRAZIL
alexandre@dm.ufscar.br

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Don Blasius
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2014 is US \$410/year for the electronic version, and \$535/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 268 No. 1 March 2014

	1
ALEXANDRE PAIVA BARRETO	
A transport inequality on the sphere obtained by mass transport	23
DARIO CORDERO-ERAUSQUIN	
A cohomological injectivity result for the residual automorphic spectrum of GL_n	33
HARALD GROBNER	
Gradient estimates and entropy formulae of porous medium and fast diffusion equations for the Witten Laplacian	47
GUANGYUE HUANG and HAIZHONG LI	
Controlled connectivity for semidirect products acting on locally finite trees	79
KEITH JONES	
An indispensable classification of monomial curves in $\mathbb{A}^4(\mathbb{k})$	95
ANARGYROS KATSABEKIS and IGNACIO OJEDA	
Contracting an axially symmetric torus by its harmonic mean curvature	117
CHRISTOPHER KIM	
Composition operators on strictly pseudoconvex domains with smooth symbol	135
HYUNGWOON KOO and SONG-YING LI	
The Alexandrov problem in a quotient space of $\mathbb{H}^2 \times \mathbb{R}$	155
ANA MENEZES	
Twisted quantum Drinfeld Hecke algebras	173
DEEPAK NAIDU	
L^p harmonic 1-forms and first eigenvalue of a stable minimal hypersurface	205
KEOMKYO SEO	
Reconstruction from Koszul homology and applications to module and derived categories	231
RYO TAKAHASHI	
A virtual Kawasaki–Riemann–Roch formula	249
VALENTIN TONITA	