

*Pacific
Journal of
Mathematics*

**THE VIRTUAL FIRST BETTI NUMBER
OF SOLUBLE GROUPS**

MARTIN R. BRIDSON AND DESSLAVA H. KOCHLOUKOVA

THE VIRTUAL FIRST BETTI NUMBER OF SOLUBLE GROUPS

MARTIN R. BRIDSON AND DESSISLAVA H. KOCHLOUKOVA

We show that if a group G is finitely presented and nilpotent-by-abelian-by-finite, then there is an upper bound on $\dim_{\mathbb{Q}} H_1(M, \mathbb{Q})$, where M runs through all subgroups of finite index in G .

1. Introduction

The *virtual first betti number* of a finitely generated group G is defined as

$$\text{vb}_1(G) = \sup\{\dim H_1(S, \mathbb{Q}) \mid S \leq G \text{ of finite index}\}.$$

A group is said to be *large* if it has a subgroup of finite index that maps onto a nonabelian free group. If G is large then $\text{vb}_1(G) = \infty$. It is easy to find finitely generated groups G that are not large but have $\text{vb}_1(G) = \infty$. For example, in the metabelian group $\mathbb{Z} \wr \mathbb{Z} = \langle a, t \mid [a, t^{-n}at^n] = 1 \text{ for all } n \rangle$, the subgroup $S_m < \mathbb{Z} \wr \mathbb{Z}$ generated by t^m and the conjugates of a has index m and $H_1(S_m, \mathbb{Z}) = \mathbb{Z}^{m+1}$. In contrast, no example is known of a *finitely presented* group that is not large but has $\text{vb}_1(G) = \infty$ (see [Button 2010; Lackenby 2010]). Since amenable groups do not contain nonabelian free subgroups, one might hope to resolve this issue by finding a finitely presented amenable group with $\text{vb}_1(G) = \infty$, but this seems to be a nontrivial matter.

We shall prove in this paper that for large classes of finitely presented soluble groups $\text{vb}_1(G)$ is always finite. One would like to prove that the same is true for all finitely presented soluble groups, but here one faces the profound difficulty of deciding which soluble groups admit finite presentations; this is unknown even for abelian-by-polycyclic and nilpotent-by-abelian groups.

In the case of metabelian groups, finite presentability is completely understood in terms of the Bieri–Strebel invariant [Bieri and Strebel 1980]. Some sufficient conditions for finite presentability of nilpotent-by-abelian groups were considered by McIsaac [1984] and later Groves [1991]. In the case of S -arithmetic nilpotent-by-abelian groups G one knows more thanks to the work of Abels [1987]: if G is an extension of a nilpotent group N by an abelian group Q then G is finitely

MSC2010: 20F16, 20J05.

Keywords: soluble groups, virtual first betti number, nilpotent-by-abelian-by-finite.

presented if and only if it is of type FP_2 , which it is if and only if $H_2(N, \mathbb{Z})$ is finitely generated as a $\mathbb{Z}Q$ -module (where the Q action is induced by conjugation) and¹ G/N' is finitely presented as a group. The first of these conditions is an easy consequence of the fact that $\mathbb{Z}Q$ is a Noetherian ring, and the second is a corollary of a result in [Bieri and Strebel 1980] that every metabelian quotient of a group of type FP_2 that does not contain noncyclic free subgroups is finitely presented. The case where G is an extension of an abelian normal subgroup A by a polycyclic group Q was approached by Brookes and Groves who studied modules over crossed products of a division ring by a free abelian group; see [Brookes and Groves 1995; 2000; 2002].

Given this background, the natural place to begin our investigation into the virtual first betti number of finitely presented soluble groups is in the setting of metabelian groups. Using methods from commutative algebra, we prove (Theorem 4.3) that if G is finitely presented and metabelian, then $vb_1(G)$ is finite. (The hypothesis that one actually needs to impose on G is somewhat weaker than finite presentability; see Remark 6.5.) The metabelian case is used in the proof of our main theorem, which is the following.

Theorem A. *Let G be a finitely presented group. If G is nilpotent-by-abelian-by-finite, then $vb_1(G)$ is finite.*

Our proof of this theorem relies on the fact that all metabelian quotients of soluble groups of type FP_2 are finitely presented [Bieri and Strebel 1980, Theorem 5.5], as well as a technical result concerning the homology of subgroups of finite index (Proposition 6.2). Groves, Kochloukova and Rodrigues [Groves et al. 2008, Theorem A] proved that if an abelian-by-polycyclic group G is of type FP_3 then it is nilpotent-by-abelian-by-finite, in which case $vb_1(G)$ is finite by Theorem A. The same is true of all soluble groups of type FP_∞ , because they are constructible [Kropholler 1986], hence nilpotent-by-abelian-by-finite, but in this case stronger finiteness results were already known: constructible soluble groups are obtained from the trivial group by finite sequences of ascending HNN extensions and finite extensions, from which it follows that they have finite Prüfer rank (i.e., there is an upper bound on the number of generators for the finitely generated subgroups).

It is natural to wonder if Theorem A might remain true when the field of rationals \mathbb{Q} in the definition of virtual betti number is replaced with other coefficient fields, such as the field with p elements F_p . We shall see in Section 5 that it does not.

Conjecture. If G is finitely presented and soluble, then $vb_1(G)$ is finite.

It is difficult to construct finitely presented soluble groups that are not nilpotent-by-abelian-by-finite. The examples provided by the constructions of Robinson and Strebel [1982] all satisfy the conjecture.

¹Throughout this article, H' denotes the commutator subgroup of a group H .

While editing the final version of this work, we learnt that Andrei Jaikin-Zapirain has, in unpublished work, also proved Theorem A in the metabelian case. Higher dimensional analogues of Theorem A are considered in the forthcoming PhD thesis of Fatemeh Mokari.

2. Preliminary results

2A. Preliminaries on finitely presented metabelian groups. We fix a short exact sequence of groups $A \twoheadrightarrow G \twoheadrightarrow Q$, where A and Q are abelian and G is finitely generated. The action of G on A by conjugation induces an action of Q , which enables us to regard A as a right $\mathbb{Z}Q$ -module. Because G is finitely generated and Q is finitely presented, A is finitely generated as a $\mathbb{Z}Q$ -module.

Associated to a nonzero real character $\chi : Q \rightarrow \mathbb{R}$ one has the monoid

$$Q_\chi = \{g \in Q \mid \chi(g) \geq 0\}.$$

The character sphere $S(Q)$ is the set of equivalence classes in $\text{Hom}(Q, \mathbb{R}) \setminus \{0\}$ under the relation that identifies $\chi_1 \sim \chi_2$ if $\chi_1 = \lambda\chi_2$ for some $\lambda > 0$. We write $[\chi]$ for the class of χ . Following [Bieri and Strebel 1980], let

$$\Sigma_A(Q) = \{[\chi] \mid A \text{ is finitely generated as a } \mathbb{Z}Q_\chi\text{-module}\}.$$

By definition, the $\mathbb{Z}Q$ -module A is *2-tame* if $\Sigma_A(Q)^c = S(Q) \setminus \Sigma_A(Q)$ contains no pair of antipodal points. According to [op. cit., Theorem 5.4], G is finitely presented if and only if A is a 2-tame $\mathbb{Z}Q$ -module, and this happens precisely when G is of homological type FP_2 . We refer the reader to [Bieri 1981] for general results concerning groups of type FP_m . If $A_1 \twoheadrightarrow A_2 \twoheadrightarrow A_3$ is an exact sequence of finitely generated $\mathbb{Z}Q$ -modules, then $\Sigma_{A_2}(Q)^c = \Sigma_{A_1}(Q)^c \cup \Sigma_{A_3}(Q)^c$ (see [Bieri and Strebel 1980, Proposition 2.2]), hence every quotient of a 2-tame $\mathbb{Z}Q$ -module is 2-tame.

2B. Tensor products and finite presentability. Let R be a noetherian commutative ring with unit 1 and let W be a finitely generated RQ -module. As above, we have a Sigma invariant $\Sigma_W(Q) = \{[\chi] \mid W \text{ is finitely generated as an } RQ_\chi\text{-module}\}$, and W is defined to be 2-tame as an RQ -module if $\Sigma_W^c(Q) = S(Q) \setminus \Sigma_W(Q)$ has no pair of antipodal points.

The question of when the tensor square $W \otimes_R W$ is finitely generated as an RQ -module (with Q acting diagonally) is addressed in [Bieri and Groves 1985], where it is shown that $[\chi]$ lies in $\Sigma_W^c(Q)$ if and only if the ring $S = RQ / \text{ann}_{RQ}(W)$ admits a real valuation $v : S \rightarrow \mathbb{R} \cup \{\infty\}$ (in the sense of Bourbaki) that extends χ and is such that the restriction v_0 of v to the image \bar{R} of R in S is nonnegative and discrete. By [loc. cit.], $W \otimes_R W$ is finitely generated as an RQ -module if and only if there is no pair of antipodal elements $[\chi], [-\chi] \in \Sigma_W^c(Q)$ that can be

lifted to valuations of S that have the same restriction v_0 to \bar{R} , with v_0 discrete and nonnegative. (These last conditions on v_0 are automatic if \bar{R} is \mathbb{Z} .)

Returning to the context of [Section 2A](#), we apply these general considerations with $W = A \otimes \mathbb{Q}$ and $R = \mathbb{Q}$, in which case $W \otimes_R W \cong (A \otimes_{\mathbb{Z}} A) \otimes_{\mathbb{Z}} \mathbb{Q}$. We deduce that if there exists a group extension $A \twoheadrightarrow G \twoheadrightarrow Q$, with G finitely presented, then $W = A \otimes \mathbb{Q}$ is 2-tame as a $\mathbb{Q}Q$ -module, and $W \otimes_R W \cong (A \otimes_{\mathbb{Z}} A) \otimes_{\mathbb{Z}} \mathbb{Q}$ is finitely generated as a $\mathbb{Q}Q$ -module via the diagonal Q -action.

We shall also need a refinement of this observation that involves the annihilator $\text{ann}_{\mathbb{Z}Q}(A)$ of A in $\mathbb{Z}Q$, which we denote I . Bieri and Strebel [[1981](#), (1.3)] prove that

$$\Sigma_A(Q) = \Sigma_{\mathbb{Z}Q/I}(Q).$$

Thus if A is 2-tame as a $\mathbb{Z}Q$ -module, then so is $\mathbb{Z}Q/I$.

Lemma 2.1. *If there exists a group extension $A \twoheadrightarrow G \twoheadrightarrow Q$ with A and Q abelian and G finitely presented, and $I = \text{ann}_{\mathbb{Z}Q}(A)$, then $(\mathbb{Z}Q/I) \otimes_{\mathbb{Z}} (\mathbb{Z}Q/I) \otimes_{\mathbb{Z}} \mathbb{Q}$ is finitely generated as a $\mathbb{Q}Q$ -module via the diagonal Q -action.*

2C. Preliminaries on commutative algebra. We will need the following basic facts from commutative algebra; for details see, for example, [[Bourbaki 1961–1965](#); [Atiyah and Macdonald 1969](#); [Eisenbud 1995](#)]. Let Q be a finitely generated abelian group and recall that the *Krull dimension* of a commutative ring is the supremum of the lengths of all chains of prime ideals in the ring.

- (1) The radical \sqrt{J} of each ideal $J \triangleleft \mathbb{Q}Q$ is the intersection of the finitely many prime ideals that contain J and are minimal subject to this condition.
- (2) Finite dimensional \mathbb{Q} -algebras are Artinian and thus have Krull dimension 0.

Throughout, if R is a commutative ring and m a positive integer, then R^m will denote the subring generated by m -th powers, *except* that \mathbb{Z}^n and \mathbb{Q}^n will denote Cartesian powers. Where no ring is specified, tensor products are assumed to be taken over \mathbb{Z} .

3. A finiteness result in commutative algebra

[Lemma 2.1](#) assures us that the following theorem applies to the modules that arise from short exact sequences $N \twoheadrightarrow G \twoheadrightarrow \mathbb{Z}^n$ associated to finitely presented metabelian groups.

Theorem 3.1. *Let $Q \cong \mathbb{Z}^n$ be a group and let $S = \mathbb{Z}Q/I$ be a commutative ring such that $(S \otimes_{\mathbb{Z}} S) \otimes_{\mathbb{Z}} \mathbb{Q}$ is finitely generated as a $\mathbb{Q}Q$ -module via the diagonal Q -action. Then,*

$$\sup_m \dim_{\mathbb{Q}}(S \otimes_{\mathbb{Z}} \mathbb{Q}^m) < \infty.$$

Proof. Let $B = S \otimes \mathbb{Q} = \mathbb{Q}Q/J$ and for each positive integer m define $J_m \triangleleft \mathbb{Q}$ to be $(J, Q^m - 1)$ and

$$B_m := B \otimes_{\mathbb{Q}} \mathbb{Q} = \mathbb{Q}Q/J_m \cong S \otimes_{\mathbb{Z}} \mathbb{Q}.$$

As $\mathbb{Q}Q/(Q^m - 1)$ is finite dimensional over \mathbb{Q} , so is $B_m = \mathbb{Q}Q/J_m$. Hence B_m has Krull dimension 0; i.e., every prime ideal in B_m is a maximal one. Therefore, the finite collection of primes ideals $P_{m,t}$ whose intersection is $\sqrt{B_m}$ are the only prime ideals in $\mathbb{Q}Q$ above J_m , and each of the quotients $\mathbb{Q}Q/P_{m,t}$ is a field.

We shall establish the theorem by proving the following:

Claim 1. *There exist only finitely many fields F such that for some $m \geq 1$ (depending on F) the field F is a quotient of B_m .*

Claim 1 provides an integer m_0 such that if a field F is a quotient of B_m then the natural map $\mathbb{Q}Q \rightarrow F$ factors through $\mathbb{Q}Q/(Q^{m_0} - 1)$.

Claim 2. *If m_0 divides m then $J_m = J_{mr}$ for every $r \in \mathbb{N}$.*

To see that the theorem follows from these claims, note that for an arbitrary positive integer m we have $J_m \supseteq J_{mm_0} = J_{m_0}$, whence

$$\begin{aligned} \dim_{\mathbb{Q}}(\mathbb{Q}Q/J_m) &\leq \dim_{\mathbb{Q}}(\mathbb{Q}Q/J_{m_0}) \leq \dim_{\mathbb{Q}}(\mathbb{Q}Q/(Q^{m_0} - 1)) \\ &= \dim_{\mathbb{Q}} \mathbb{Q}[Q/Q^{m_0}] = m_0^n. \end{aligned}$$

Proof of Claim 1. Our hypothesis on S implies that $B \otimes_{\mathbb{Q}} B$ is finitely generated as $\mathbb{Q}Q$ -module via the diagonal Q -action, by d elements say. Let F be a field quotient of B_m and let $\theta : \mathbb{Q}Q \rightarrow F$ be the canonical projection; so $Q^m - 1 \subseteq \ker(\theta)$. Then, $\theta(Q)$ is a finitely generated multiplicative subgroup of F^* that has finite exponent and F , being finite dimensional over \mathbb{Q} , embeds in \mathbb{C} . Hence $\theta(Q)$ is a finite cyclic group, generated by a root of unity, ϵ of order s , say. Thus we obtain a subgroup $H < Q$ such that Q/H is cyclic of order s and $H - 1 \subseteq \ker(\theta)$. Now, $F \cong \mathbb{Q}[x]/(f)$, where f is the minimal polynomial of ϵ over \mathbb{Q} . And f is an irreducible factor of $x^s - 1$ in $\mathbb{Q}[x]$, whose zeroes are distinct roots of unity with order precisely s . Thus $\dim_{\mathbb{Q}} F = \deg(f) = \varphi(s)$, where φ is Euler's totient function. On the other hand, $F \otimes_{\mathbb{Q}} F$ is an epimorphic image of the $\mathbb{Q}Q$ -module $B \otimes_{\mathbb{Q}} B$ and the action of Q on $F \otimes_{\mathbb{Q}} F$ factors through the action of Q/H , so $F \otimes_{\mathbb{Q}} F$ is generated as a $\mathbb{Q}[Q/H]$ -module by d elements. Hence

$$\varphi(s)^2 = (\dim_{\mathbb{Q}} F)^2 = \dim_{\mathbb{Q}}(F \otimes_{\mathbb{Q}} F) \leq d \dim_{\mathbb{Q}} \mathbb{Q}[Q/H] = ds.$$

An elementary calculation shows that $\varphi(n)/\sqrt{n} \rightarrow \infty$ as $n \rightarrow \infty$, so for fixed d there are only finitely many possible values of s and ϵ . Let b be a natural number such that the order of ϵ is at most b . Then, the order of ϵ is a divisor of $m_0 = b!$ and

$$F \text{ is a quotient of } \mathbb{Q}Q/(Q^{m_0} - 1).$$

Since $\mathbb{Q}Q/(Q^{m_0} - 1)$ is finite dimensional over \mathbb{Q} it has Krull dimension 0, so has only finitely many prime ideals and finitely many field quotients. This completes the proof of Claim 1. \square

Proof of Claim 2. Since m_0 divides m we have $J_m \subseteq J_{m_0}$, so the prime ideals containing J_{m_0} also contain J_m . On the other hand, we saw earlier that for each of the prime ideals $P_{m,i}$ containing J_m , the quotient $F_i := \mathbb{Q}Q/P_{m,i}$ is a field. By definition, m_0 is such that $\mathbb{Q}Q \rightarrow F_i$ factors through $\mathbb{Q}Q/(Q^{m_0} - 1)$, and therefore $P_{m,i}$ (which already contains $J \subset J_m$) contains $J_{m_0} = (J, Q^{m_0} - 1)$. The radical of J_m is the intersection of the prime ideals containing it, so

$$\sqrt{J_m} = \sqrt{J_{m_0}}.$$

Arguing by induction on r , Claim 2 will follow if we can prove that for every prime number p we have $J_m = J_{mp}$, which is equivalent to the assertion that $q^m - 1 \in J_{mp}$ for all $q \in Q$.

We now fix $q \in Q$. From the preceding argument, $\sqrt{J_m} = \sqrt{J_{mp}}$. In particular, $Q^m - 1 \subseteq J_m \subseteq \sqrt{J_m} = \sqrt{J_{mp}}$, so there is a natural number s (over which we have no control) such that

$$(3-1) \quad (q^m - 1)^s \in J_{mp}.$$

As $Q^{mp} - 1 \subseteq J_{mp}$, we also have

$$(3-2) \quad q^{mp} - 1 \in J_{mp}.$$

Let $g(x)$ be the greatest common divisor of $x^{pm} - 1$ and $(x^m - 1)^s$ in $\mathbb{Q}[x]$. In characteristic zero, the polynomial $x^{pm} - 1$ has no repeated roots, so neither does $g(x)$. Since $g(x)$ divides $(x^m - 1)^s$, it must actually divide $x^m - 1$, so in fact $g(x) = x^m - 1$. From (3-1), (3-2) and Bézout's lemma, we have $g(q) \in J_{mp}$. Since $q \in Q$ is arbitrary, this implies that $J_{mp} = J_m$. \square

4. The main theorem for metabelian groups

In this section we prove that all finitely presented metabelian groups have finite virtual first betti number. The proof relies on the finiteness theorem proved in the previous section and two technical lemmas, the first of which is a simple observation about commensurable groups.

Lemma 4.1. *Let G be a group. If $G_0 < G$ is a subgroup of finite index, then $\text{vb}_1(G) = \text{vb}_1(G_0)$.*

Proof. By definition, $\text{vb}_1(G) = \sup_M \dim H_1(M, \mathbb{Q})$, where the supremum is taken over finite-index subgroups of G . If M has finite index in G_0 , then it also has finite index in G , so $\text{vb}_1(G) \geq \text{vb}_1(G_0)$. Conversely, if S has finite index in G , then

$S_0 = G_0 \cap S$ has finite index in G_0 , and since it also has finite index in S , we have $\dim H_1(S_0, \mathbb{Q}) \geq \dim H_1(S, \mathbb{Q})$, so $\text{vb}_1(G_0) \geq \text{vb}_1(G)$. \square

Lemma 4.2. *Let $A \twoheadrightarrow G \twoheadrightarrow Q$ be a short exact sequence of groups with A and Q abelian and let n be the torsion-free rank of Q . Then:*

(a) *Writing $[G, A] = \langle \{[g, a] = g^{-1}a^{-1}ga \mid g \in G, a \in A\} \rangle$, we have*

$$\dim_{\mathbb{Q}} H_1(G, \mathbb{Q}) \leq \dim_{\mathbb{Q}}(A/[G, A] \otimes \mathbb{Q}) + n.$$

In the split case, $G = A \rtimes Q$, we have $H_1(G, \mathbb{Q}) \cong (G/[G, A]) \otimes_{\mathbb{Z}} \mathbb{Q}$, and

$$\dim_{\mathbb{Q}} H_1(G, \mathbb{Q}) = \dim_{\mathbb{Q}}(A/[G, A] \otimes \mathbb{Q}) + n.$$

(b) *If G_m is a subgroup of finite index in G and Q_m is the image of G_m in Q , then*

$$\dim_{\mathbb{Q}} H_1(G_m, \mathbb{Q}) \leq \dim_{\mathbb{Q}}(A \otimes_{\mathbb{Z}Q_m} \mathbb{Q}) + n.$$

In the split case, $G_m = (A \cap G_m) \rtimes Q_m$, equality is attained:

$$\dim_{\mathbb{Q}} H_1(G_m, \mathbb{Q}) = \dim_{\mathbb{Q}}(A \otimes_{\mathbb{Z}Q_m} \mathbb{Q}) + n.$$

(c) *If $G = A \rtimes Q$ and \mathcal{B} denotes the set of subgroups of finite index in Q , then*

$$\text{vb}_1(G) = \sup_{S \in \mathcal{B}} \dim_{\mathbb{Q}}(A \otimes_{\mathbb{Z}S} \mathbb{Q}) + n.$$

Proof. (a) As $[G, A] \subseteq [G, G]$, we see that $H_1(G, \mathbb{Z}) = G/[G, G]$ is a quotient of $G/[G, A]$. So from the central extension $A/[G, A] \twoheadrightarrow G/[G, A] \twoheadrightarrow Q$, we get $\dim_{\mathbb{Q}} H_1(G, \mathbb{Q}) \leq \dim_{\mathbb{Q}}(A/[G, A] \otimes \mathbb{Q}) + \dim_{\mathbb{Q}}(Q \otimes \mathbb{Q}) = \dim_{\mathbb{Q}}(A/[G, A] \otimes \mathbb{Q}) + n$.

If $G = A \rtimes Q$ then, using that A, Q are abelian and A is normal in G , we get $[G, G] = [AQ, AQ] = [Q, A] \subseteq [G, A] \subseteq [G, G]$, hence $[G, G] = [G, A]$ and $A/[G, A] \twoheadrightarrow G/[G, G] \twoheadrightarrow Q$ is an exact sequence of abelian groups.

(b) We consider the short exact sequence $A_m \twoheadrightarrow G_m \twoheadrightarrow Q_m$, where $A_m = A \cap G_m$. From part (a) we have

$$(4-1) \quad \dim_{\mathbb{Q}} H_1(G_m, \mathbb{Q}) \leq \dim_{\mathbb{Q}}(A_m \otimes_{\mathbb{Z}Q_m} \mathbb{Q}) + n,$$

with equality if the sequence splits. Furthermore, since A/A_m is finite we have

$$0 = \text{Tor}_1^{\mathbb{Z}Q_m}(A/A_m, \mathbb{Q}) \quad \text{and} \quad (A/A_m) \otimes_{\mathbb{Z}Q_m} \mathbb{Q} = 0.$$

Thus there is an exact sequence (part of the long exact sequence in Tor associated to $A \cap G_m \twoheadrightarrow A \twoheadrightarrow A/(A \cap G_m)$)

$$0 = \text{Tor}_1^{\mathbb{Z}Q_m}(A/A_m, \mathbb{Q}) \rightarrow A_m \otimes_{\mathbb{Z}Q_m} \mathbb{Q} \rightarrow A \otimes_{\mathbb{Z}Q_m} \mathbb{Q} \rightarrow (A/A_m) \otimes_{\mathbb{Z}Q_m} \mathbb{Q} = 0,$$

whence $A_m \otimes_{\mathbb{Z}Q_m} \mathbb{Q} \cong A \otimes_{\mathbb{Z}Q_m} \mathbb{Q}$. Thus, we may replace $A_m \otimes_{\mathbb{Z}Q_m} \mathbb{Q}$ in (4-1) by $A \otimes_{\mathbb{Z}Q_m} \mathbb{Q}$, and (b) is proved.

(c) From the first part of (b) we have

$$\text{vb}_1(G) \leq \sup_{S \in \mathfrak{B}} \dim_{\mathbb{Q}}(A \otimes_{\mathbb{Z}S} \mathbb{Q}) + n,$$

and to obtain the reverse inequality, we use the second part of (b)

$$\sup_{S \in \mathfrak{B}} \dim_{\mathbb{Q}}(A \otimes_{\mathbb{Z}S} \mathbb{Q}) + n = \sup_{S \in \mathfrak{B}} \dim_{\mathbb{Q}} H_1(A \rtimes S, \mathbb{Q}),$$

noting that $A \rtimes S$ has finite index in G . □

Theorem 4.3. *Let $A \twoheadrightarrow G \twoheadrightarrow Q$ be a short exact sequence of groups with A and Q abelian. If G is finitely presented then its virtual first betti number $\text{vb}_1(G)$ is finite.*

Proof. By passing to a subgroup of finite index in Q and replacing G by the inverse image of this subgroup, we may assume that Q is free abelian. Lemma 4.1 assures us that it is enough to consider this case, and Lemma 4.2(b) tells us that we will be done if we can establish an upper bound on $\dim_{\mathbb{Q}}(A \otimes_{\mathbb{Z}Q_m} \mathbb{Q})$ as Q_m ranges over the subgroups of finite index in Q .

Recall that A is finitely generated as a $\mathbb{Z}Q$ -module, say by d elements. Thus, denoting the annihilator $\text{ann}_{\mathbb{Z}Q}(A) = \{\lambda \in \mathbb{Z}Q \mid A\lambda = 0\}$ by I , we have an epimorphism of $\mathbb{Z}Q$ -modules

$$(\mathbb{Z}Q/I)^{[d]} = \mathbb{Z}Q/I \oplus \cdots \oplus \mathbb{Z}Q/I \rightarrow A$$

that induces an epimorphism of \mathbb{Q} -vector spaces

$$((\mathbb{Z}Q/I) \otimes_{\mathbb{Z}Q_m} \mathbb{Q})^{[d]} = (\mathbb{Z}Q/I)^{[d]} \otimes_{\mathbb{Z}Q_m} \mathbb{Q} \rightarrow A \otimes_{\mathbb{Z}Q_m} \mathbb{Q}.$$

Thus,

$$\dim_{\mathbb{Q}}(A \otimes_{\mathbb{Z}Q_m} \mathbb{Q}) \leq d \dim_{\mathbb{Q}}((\mathbb{Z}Q/I) \otimes_{\mathbb{Z}Q_m} \mathbb{Q})$$

and it suffices to show that

$$\sup_m \dim_{\mathbb{Q}}((\mathbb{Z}Q/I) \otimes_{\mathbb{Z}Q_m} \mathbb{Q}) < \infty.$$

For every m there is a natural number α_m such that $Q^{\alpha_m} \subseteq Q_m$, and $\mathbb{Z}Q/I \otimes_{\mathbb{Z}Q_m} \mathbb{Q}$ is a quotient of $\mathbb{Z}Q/I \otimes_{\mathbb{Z}Q^{\alpha_m}} \mathbb{Q}$. Thus,

$$\dim_{\mathbb{Q}}((\mathbb{Z}Q/I) \otimes_{\mathbb{Z}Q_m} \mathbb{Q}) \leq \dim_{\mathbb{Q}}((\mathbb{Z}Q/I) \otimes_{\mathbb{Z}Q^{\alpha_m}} \mathbb{Q}),$$

and we have reduced to showing that

$$\sup_s \dim_{\mathbb{Q}}((\mathbb{Z}Q/I) \otimes_{\mathbb{Z}Q^s} \mathbb{Q}) < \infty.$$

The theorem now follows from Lemma 2.1 and Theorem 3.1. □

5. Characteristic p case

In this section we shall construct examples which show that the restriction to fields of characteristic 0 in [Theorem A](#) is essential, even in the metabelian case.² To this end, we consider the mod p *virtual first betti number* of a finitely generated group G ,

$$\text{vb}_1^{(p)}(G) = \sup\{\dim H_1(S, F_p) \mid S < G \text{ of finite index}\}.$$

Proposition 5.1. *For every prime p there exist finitely presented metabelian groups Γ such that $\text{vb}_1^{(p)}(\Gamma)$ is infinite.*

Proof. Let Q be a free abelian group with generators x and y and let $A = F_p Q/I$, where I is the ideal of $F_p Q$ generated by $y - x^2 + x - 1$. Then,

$$A \cong F_p \left[x, x^{-1}, \frac{1}{x^2 - x + 1} \right].$$

Consider

$$A_m = A \otimes_{\mathbb{Z} Q^{p^m}} F_p \cong F_p Q / (I, Q^{p^m} - 1).$$

Since $(x^2 - x + 1)^{p^m} - 1 = x^{2p^m} - x^{p^m} + 1 - 1 = x^{p^m}(x^{p^m} - 1)$, we have

$$\begin{aligned} A_m &= F_p \left[x, x^{-1}, \frac{1}{x^2 - x + 1} \right] / (x^{p^m} - 1, (x^2 - x + 1)^{p^m} - 1) \\ &= F_p \left[x, x^{-1}, \frac{1}{x^2 - x + 1} \right] / (x^{p^m} - 1) \end{aligned}$$

is the localisation

$$B_m S^{-1}$$

where $B_m = F_p[x, x^{-1}]/(x^{p^m} - 1)$ and S is the image of $\{(x^2 - x + 1)^j\}_{j \geq 1}$ in B_m . Note that $x^{p^m} - 1$ and $x^2 - x + 1$ do not have a common root in any finite field extension of F_p , for if z were a common root we would have $1 = z^{2p^m} = (z - 1)^{p^m} = z^{p^m} - 1 = 0$, which is a contradiction. Thus the polynomials $x^{p^m} - 1$ and $(x^2 - x + 1)^j$ are coprime in $F_p[x, x^{-1}]$; i.e., they generate the whole ring as an ideal, and so the elements of S are invertible in B_m . Therefore $B_m S^{-1} = B_m$ and

$$\dim_{F_p} A_m = \dim_{F_p} B_m S^{-1} = \dim_{F_p} B_m = p^m.$$

Now define

$$\Gamma = A \rtimes Q \quad \text{and} \quad \Gamma_m = A \rtimes Q^{p^m}.$$

Then, as in the split case of [Lemma 4.2\(b\)](#) (with coefficients in F_p in place of \mathbb{Q}),

$$\dim_{F_p} H_1(\Gamma_m, F_p) = \dim_{F_p} A_m + 2 = p^m + 2,$$

²John Wilson [1998] proved that the dimension of $H_1(S, F_p)$ can grow at most like the square root of the index $[G : S]$. Jack Button [2010] exhibited a finitely presented soluble group that exhibits this growth for all p .

which tends to infinity with m .

By the calculation [Bieri and Strebel 1981, Theorem 5.2] of $\Sigma_A(Q)$ for $A = F_p Q/I$, where the ideal I is 1-generated, or by the link between $\Sigma_A^c(Q)$ and valuation theory (as described in Section 2B), we have

$$\Sigma_A^c(Q) = \{[\chi_1], [\chi_2], [\chi_3]\},$$

with

$$\begin{aligned} \chi_1(x) &= 0, & \chi_2(x) &= 1, & \chi_3(x) &= -1, \\ \chi_1(y) &= 1, & \chi_2(y) &= 0, & \chi_3(y) &= -2. \end{aligned}$$

Thus, A is 2-tame as a $\mathbb{Z}Q$ -module, and by the classification of finitely presented metabelian groups in [Bieri and Strebel 1980], Γ is finitely presented. \square

Corollary 5.2. *There exists a finitely presented metabelian group G such that for the class \mathcal{A} of all subgroups of finite index in G ,*

$$\sup_{M \in \mathcal{A}} d(M) = \infty,$$

where $d(M)$ is the minimal number of generators of M .

Proof. Immediate, since $d(M) \geq \dim_{F_p} H_1(M, F_p)$. \square

It is natural to wonder if the lack of finiteness exhibited in the preceding proposition might be avoided by restricting to subgroups whose index is coprime to p . The following refinement shows that this is not the case.

Proposition 5.3. *Let p be a prime. There exist finitely presented metabelian groups G such that*

$$\sup\{\dim_{F_p} H_1(S, F_p) \mid S \in \mathcal{A}_p\} = \infty,$$

where

$$\mathcal{A}_p = \{S \leq G \mid [G : S] \text{ is finite and coprime to } p\}.$$

Proof. Let $A = F_p[x, x^{-1}, (x+1)^{-1}]$ and let Q be a free abelian group of rank 2 whose generators x_1, x_2 act on A as multiplication by x and $x+1$, respectively. We consider the group $G = A \rtimes Q$. As an $F_p[Q]$ -module, $A \cong F_p[Q]/I$ where I is the ideal generated by $x_2 - x_1 - 1$, and the argument given in the preceding proposition shows that $\Sigma_A(Q)^c$ consists of precisely 3 points, no pair of which is antipodal. Therefore, G is finitely presented.

Let F be a finite field with p^r elements, $r \geq 2$. Let w be a generator of the multiplicative group $F^* = F \setminus \{0\}$. Let Q_r be the kernel of the homomorphism $Q \rightarrow F^*$ defined by $x_1 \mapsto w$ and $x_2 \mapsto w+1$. Let $G_r = A \rtimes Q_r$ and note that $|G/G_r| = |Q/Q_r| = p^r - 1$ is coprime to p .

The ring epimorphism $A \rightarrow F$ sending x to w provides an epimorphism of the underlying additive groups which extends to a group epimorphism $A \rtimes Q_r \rightarrow F \times \mathbb{Z}^2$.

Since $\dim_{F_p} F = r$, it follows that $\dim_{F_p} H_1(G_r, F_p) \geq r + 2$. And, $r \geq 2$ was arbitrary. \square

6. Beyond the metabelian case

In this section we shall prove [Theorem A](#), but first we present a consequence of [Theorem 4.3](#) that describes what one can deduce about towers of finite-index subgroups above the commutator subgroup in amenable and related groups.

Proposition 6.1. *Let G be a group of type FP_2 that does not contain a nonabelian free group and let \mathcal{C} be the set of finite-index subgroups in G that contain the commutator subgroup G' . Then, $\sup_{M \in \mathcal{C}} \dim_{\mathbb{Q}} H_1(M, \mathbb{Q}) < \infty$.*

Proof. By [[Bieri and Strebel 1980](#), Theorem 5.5], G/G'' is finitely presented. Since $M \supseteq G'$, we have $M' \supseteq G''$ and can replace G by G/G'' and M by MG''/G'' without changing $H_1(M, \mathbb{Q})$. Then we can apply [Theorem 4.3](#). \square

Our proof of [Theorem A](#) relies on the following proposition, which is of interest in its own right.

Proposition 6.2. *Let $N \twoheadrightarrow G \twoheadrightarrow Q$ be a short exact sequence of groups, where N is nilpotent, Q is abelian and G is finitely generated. Let G_n be a subgroup of finite index in G and let \bar{G}_n be the image of G_n in the metabelian group G/N' . Then,*

$$\dim_{\mathbb{Q}} H_1(G_n, \mathbb{Q}) = \dim_{\mathbb{Q}} H_1(\bar{G}_n, \mathbb{Q}).$$

Proof. We argue using the Malcev completion $j_N : N \rightarrow N^*$ [[Malcev 1949](#)]. According to [[Quillen 1969](#), Appendix A, Corollary 3.8], for any nilpotent group N , the homomorphism $j_N : N \rightarrow N^*$ is characterized up to isomorphism by the following properties:

- (a) N^* is nilpotent and uniquely divisible.
- (b) $\ker j_N$ is the torsion subgroup of N .
- (c) For every $x \in N^*$, there is a positive integer n such that $x^n \in N$.

In any nilpotent group, the set $\sqrt[S]{S}$ of elements that have powers in a fixed subgroup S is a subgroup. It follows that, for every subgroup $M < N$, the map $M \rightarrow \sqrt{j_N(M)}$ satisfies properties (a) to (c). Thus we may identify M^* with $\sqrt{j_N(M)} < N^*$. If $M < N$ has finite index, then $M^* = \sqrt{j_N(M)} = N^*$. And $(N^*)' = (N')^*$.

With these facts in hand, for all subgroups of finite index $G_n < G$ we have $(G'_n)^* \supseteq ((G_n \cap N)')^* = ((G_n \cap N)^*)' = (N^*)' = (N')^*$. Thus $(G'_n N')^* = (G'_n)^*$, and from (c) we deduce that $G'_n(N' \cap G_n)/G'_n$ is torsion. As $G'_n(N' \cap G_n)/G'_n$ is the kernel of the canonical epimorphism $G_n/G'_n \rightarrow G_n N'/G'_n N'$, we have

$$H_1(G_n, \mathbb{Q}) \cong (G_n/G'_n) \otimes \mathbb{Q} \cong (G_n N'/G'_n N') \otimes \mathbb{Q} \cong H_1(\bar{G}_n, \mathbb{Q}). \quad \square$$

Theorem 6.3. *Let $N \twoheadrightarrow G \twoheadrightarrow Q$ be a short exact sequence of groups. If N is nilpotent, Q is abelian and G is of type FP_2 , then the virtual first betti number $\text{vb}_1(G)$ is finite.*

Proof. In the light of [Proposition 6.2](#), this follows directly from [Theorem 4.3](#) and the fact [[Bieri and Strebel 1980](#), Theorem 5.5] that G/N' is a finitely presented metabelian group. \square

Corollary 6.4 ([Theorem A](#)). *If a group G is nilpotent-by-abelian-by-finite and of type FP_2 , then $\text{vb}_1(G)$ is finite.*

Proof. Let G_0 be a subgroup of finite index in G such that G_0 is nilpotent-by-abelian. Then, G_0 has type FP_2 , so $\text{vb}_1(G_0)$ is finite, by [Theorem 6.3](#), and hence, so is G , by [Lemma 4.1](#). \square

Remark 6.5. We did not use the full force of finite presentability in establishing [Theorem A](#): in fact, it is enough to assume that G has a subgroup of finite index G_0 in which there is a nilpotent subgroup $N \triangleleft G_0$ such that $Q = G_0/N$ is free abelian and, writing $A = N/N'$, the $\mathbb{Q}Q$ -module $A \otimes A \otimes \mathbb{Q}$, with diagonal action, should be finitely generated. These requirements follow from the finite presentability of G_0/N' but are strictly weaker.

Corollary 6.6. *Every soluble group of type FP_∞ has finite virtual first betti number.*

Proof. Soluble groups S of type FP_∞ are constructible and hence nilpotent-by-abelian-by-finite [[Kropholler 1986](#)]. \square

Corollary 6.7. *Every abelian-by-polycyclic group of type FP_3 has finite virtual first betti number.*

Proof. By the main result of [[Groves et al. 2008](#)], abelian-by-polycyclic groups of type FP_3 are nilpotent-by-abelian-by-finite. \square

Acknowledgements

We thank Fatemeh Mokari and an anonymous referee for helpful comments. We particularly thank the referee for simplifying the proofs of [Proposition 6.2](#) and [Claim 2](#) in [Section 3](#). The work of Prof. Bridson was supported by grants from the EPSRC and by a Wolfson Merit Award from the Royal Society; the work of Prof. Kochloukova was supported by “bolsa de produtividade em Pesquisa”, CNPq, Brazil: we thank all of these organizations.

References

- [Abels 1987] H. Abels, *Finite presentability of S -arithmetic groups. Compact presentability of solvable groups*. Lecture Notes in Math. **1261**, Springer, Berlin, 1987. [MR 89b:22017](#) [Zbl 0621.20015](#)
- [Atiyah and Macdonald 1969] M. F. Atiyah and I. G. Macdonald, *Introduction to commutative algebra*, Addison-Wesley, Reading, MA, 1969. [MR 39 #4129](#) [Zbl 0175.03601](#)

- [Bieri 1981] R. Bieri, *Homological dimension of discrete groups*, 2nd ed., Queen Mary College, Department of Pure Mathematics, London, 1981. MR 84h:20047 Zbl 0357.20027
- [Bieri and Groves 1985] R. Bieri and J. R. J. Groves, “Tensor powers of modules over finitely generated abelian groups”, *J. Algebra* **97**:1 (1985), 68–78. MR 87c:20017 Zbl 0705.20050
- [Bieri and Strebel 1980] R. Bieri and R. Strebel, “Valuations and finitely presented metabelian groups”, *Proc. London Math. Soc.* (3) **41**:3 (1980), 439–464. MR 81j:20080 Zbl 0448.20029
- [Bieri and Strebel 1981] R. Bieri and R. Strebel, “A geometric invariant for modules over an abelian group”, *J. Reine Angew. Math.* **322** (1981), 170–189. MR 82f:20017 Zbl 0448.20010
- [Bourbaki 1961–1965] N. Bourbaki, *Algèbre commutative: chapitres 1–7*, *Eléments de mathématique* **27–28, 30–31**, Hermann, Paris, 1961–1965. Translated as *Commutative algebra: chapters 1–7*, Springer, Berlin, 1998. MR 2001g:13001 Zbl 1103.13001
- [Brookes and Groves 1995] C. J. B. Brookes and J. R. J. Groves, “Modules over nilpotent group rings”, *J. London Math. Soc.* (2) **52**:3 (1995), 467–481. MR 97f:20009 Zbl 0857.20002
- [Brookes and Groves 2000] C. J. B. Brookes and J. R. J. Groves, “Modules over crossed products of a division ring by a free abelian group I”, *J. Algebra* **229**:1 (2000), 25–54. MR 2001g:16052 Zbl 0958.16028
- [Brookes and Groves 2002] C. J. B. Brookes and J. R. J. Groves, “Modules over crossed products of a division ring by a free abelian group II”, *J. Algebra* **253**:2 (2002), 417–445. MR 2003g:16034 Zbl 1011.16018
- [Button 2010] J. O. Button, “Largeness of LERF and 1-relator groups”, *Groups Geom. Dyn.* **4**:4 (2010), 709–738. MR 2011j:20078 Zbl 1248.20033
- [Eisenbud 1995] D. Eisenbud, *Commutative algebra: with a view toward algebraic geometry*, Graduate Texts in Mathematics **150**, Springer, New York, 1995. MR 97a:13001 Zbl 0819.13001
- [Groves 1991] J. R. J. Groves, “Some finitely presented nilpotent-by-abelian groups”, *J. Algebra* **144**:1 (1991), 127–166. MR 93b:20059 Zbl 0806.20023
- [Groves et al. 2008] J. R. J. Groves, D. H. Kochloukova, and C. F. Rodrigues, “Abelian-by-polycyclic groups of homological type FP_3 ”, *J. Group Theory* **11**:5 (2008), 609–621. MR 2009j:20074 Zbl 1180.20042
- [Kropholler 1986] P. H. Kropholler, “Cohomological dimension of soluble groups”, *J. Pure Appl. Algebra* **43**:3 (1986), 281–287. MR 88h:20063 Zbl 0603.20033
- [Lackenby 2010] M. Lackenby, “Detecting large groups”, *J. Algebra* **324**:10 (2010), 2636–2657. MR 2011h:20045 Zbl 1231.20026
- [Malcev 1949] A. I. Malcev, “Нильпотентные группы без кручения”, *Izvestiya Akad. Nauk. SSSR. Ser. Mat.* **13**:3 (1949), 201–212. Translated as “Nilpotent torsion-free groups”. MR 10,507e Zbl 0034.01702
- [McIsaac 1984] A. J. McIsaac, “Some conditions for finite presentation of nilpotent-by-abelian groups”, *Math. Proc. Cambridge Philos. Soc.* **95**:3 (1984), 411–415. MR 85k:20104 Zbl 0541.20019
- [Quillen 1969] D. Quillen, “Rational homotopy theory”, *Ann. of Math.* (2) **90** (1969), 205–295. MR 41 #2678 Zbl 0191.53702
- [Robinson and Strebel 1982] D. J. S. Robinson and R. Strebel, “Some finitely presented soluble groups which are not nilpotent by abelian by finite”, *J. London Math. Soc.* (2) **26**:3 (1982), 435–440. MR 84m:20041 Zbl 0465.20027
- [Wilson 1998] J. S. Wilson, “Finitely presented soluble groups”, pp. 296–316 in *Geometry and cohomology in group theory* (Durham, 1994), edited by P. H. Kropholler et al., London Math. Soc. Lecture Note Ser. **252**, Cambridge Univ. Press, 1998. MR 2000m:20052 Zbl 0908.20025

Received July 25, 2014. Revised September 15, 2014.

MARTIN R. BRIDSON
MATHEMATICAL INSTITUTE
UNIVERSITY OF OXFORD
ANDREW WILES BUILDING
WOODSTOCK ROAD
OXFORD
OX2 6GG
UNITED KINGDOM

bridson@maths.ox.ac.uk

DESSLAVA H. KOCHLOUKOVA
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CAMPINAS (UNICAMP)
13083-859 CAMPINAS-SP
BRAZIL

desi@ime.unicamp.br

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 274 No. 2 April 2015

On Demazure and local Weyl modules for affine hyperalgebras	257
ANGELO BIANCHI, TIAGO MACEDO and ADRIANO MOURA	
On curves and polygons with the equiangular chord property	305
TARIK AOUGAB, XIDIAN SUN, SERGE TABACHNIKOV and YUWEN WANG	
The well-posedness of nonlinear Schrödinger equations in Triebel-type spaces	325
SHAOLEI RU and JIECHENG CHEN	
Hypersurfaces with constant curvature quotients in warped product manifolds	355
JIE WU and CHAO XIA	
The first terms in the expansion of the Bergman kernel in higher degrees	373
MARTIN PUCHOL and JIALIN ZHU	
Determinant rank of C^* -algebras	405
GUIHUA GONG, HUAXIN LIN and YIFENG XUE	
Motion by mixed volume preserving curvature functions near spheres	437
DAVID HARTLEY	
Homomorphisms on infinite direct products of groups, rings and monoids	451
GEORGE M. BERGMAN	
The virtual first Betti number of soluble groups	497
MARTIN R. BRIDSON and DESSISLAVA H. KOCHLOUKOVA	