

*Pacific
Journal of
Mathematics*

HARMONIC MAPS FROM \mathbb{C}^n TO KÄHLER MANIFOLDS

JIANMING WAN

Volume 275 No. 1

May 2015

HARMONIC MAPS FROM \mathbb{C}^n TO KÄHLER MANIFOLDS

JIANMING WAN

We prove that a harmonic map from \mathbb{C}^n ($n \geq 2$) to any Kähler manifold must be holomorphic under an assumption on energy density. This can be considered as a complex analogue of the Liouville-type theorem for harmonic maps obtained by Sealey.

1. Introduction

The classical Liouville theorem says that a bounded harmonic function on \mathbb{R}^n (or holomorphic function on \mathbb{C}^n) has to be constant. Sealey [1982] (see also [Xin 1996]) gave an analogue for harmonic maps. He proved that a harmonic map of finite energy from \mathbb{R}^n ($n \geq 2$) to any Riemannian manifold must be a constant map. In this paper we consider the complex analogue of Sealey's result by asking: *Must a harmonic map with finite $\bar{\partial}$ -energy from \mathbb{C}^n ($n \geq 2$) to any Kähler manifold be holomorphic?*

On the other hand, from Siu and Yau's proof of the Frankel conjecture [1980] (the key is to prove that a stable harmonic map from S^2 to $\mathbb{C}P^n$ is holomorphic or conjugate holomorphic), we know that it is very important to study the holomorphicity of harmonic maps. So the above question is obviously interesting. We hope that it is true. But we do not know how to prove it. Our partial result can be stated as follows:

Theorem 1.1. *Let f be a harmonic map from \mathbb{C}^n ($n \geq 2$) to any Kähler manifold. Let $e(f)$ be the energy density and $e''(f)$ the $\bar{\partial}$ -energy density. If*

$$(1-1) \quad e(f)e''(f)(p) = O\left(\frac{1}{R^{4n+\alpha}}\right)^1$$

for some $\alpha > 0$, where R denotes the distance from the origin to p , then f is a holomorphic map.

The research is supported by the National Natural Science Foundation of China 11301416.

MSC2010: 53C55.

Keywords: harmonic maps, holomorphic maps.

¹The notation O means $e(f)e''(f)(p) \leq C/R^{4n+\alpha}$ for some $C > 0$ and sufficiently large R .

The condition (1-1) implies that the $\bar{\partial}$ -energy is finite. Since

$$(e''(f))^2 \leq e(f)e''(f) = O\left(\frac{1}{R^{4n+\alpha}}\right),$$

one has

$$e''(f) = O\left(\frac{1}{R^{2n+\alpha/2}}\right).$$

This leads to

$$\int_{\mathbb{C}^n} e''(f) dv < \infty.$$

Note that we do not have any curvature assumption for the target manifold.

We should mention some other related results on holomorphicity of harmonic maps. For instance, Dong [2013] established many holomorphicity results under the assumption of the target manifolds having strongly seminegative curvature. Xin [1985] obtained some holomorphicity results on harmonic maps from a complete Riemann surface into $\mathbb{C}\mathbb{P}^n$.

If the target manifold is \mathbb{C}^m (in this case every component of the map is a harmonic function), then the answer of above question is positive (see [Wan 2010]).

The main idea of the proof of Theorem 1.1 is to consider a one-parameter family of maps and study the $\bar{\partial}$ -energy variation.

The rest of the paper is organized as follows. Section 2 contains some basic materials on harmonic maps. In Section 3, we study the first variation of the $\bar{\partial}$ -energy. Theorem 1.1 is proved in Section 4.

2. Preliminaries

The materials in this section may be found in [Xin 1996].

2A. Basic concepts of harmonic maps. Let f be a smooth map between two Riemannian manifolds (M, g) and (N, h) . We can define the energy density of f by

$$e(f) = \frac{1}{2} \text{trace} |df|^2 = \frac{1}{2} \sum_{i=1}^m \langle f_* e_i, f_* e_i \rangle,$$

where $\{e_i\}$ ($i = 1, \dots, m = \dim M$) is a local orthonormal frame field of M . The energy integral is defined by

$$E(f) = \int_M e(f) dv.$$

If we choose local coordinates $\{x^i\}$ and $\{y^\alpha\}$ in M and N , respectively, the energy density can be written as

$$(2-1) \quad e(f)(x) = \frac{1}{2} g^{ij}(x) \frac{\partial f^\alpha(x)}{\partial x^i} \frac{\partial f^\beta(x)}{\partial x^j} h_{\alpha\beta}(f(x)).$$

The tension field of f is

$$\tau(f) = (\nabla_{e_i} df)(e_i),$$

where ∇ is the induced connection on the pullback bundle $f^{-1}TN$ over M from those of M and N .

Definition 2.1. We say that f is a harmonic map if $\tau(f) = 0$.

From the variation point of view, a harmonic map can be seen as the critical point of the energy integral functional. Let f_t be a one-parameter family of maps. We can regard it as a smooth map $M \times (-\epsilon, \epsilon) \rightarrow N$. Let $f_0 = f$ and $(df_t/dt)|_{t=0} = v$. Then we have the first-variation formula (see [Xin 1996])

$$(2-2) \quad \frac{d}{dt} E(f_t)|_{t=0} = \int_M \operatorname{div} W \, dv - \int_M \langle v, \tau(f) \rangle \, dv,$$

where $W = \langle v, f_* e_j \rangle e_j$. If M is compact, then $\int_M \operatorname{div} W \, dv = 0$. We know that a harmonic map is the critical point of the energy functional.

2B. $\bar{\partial}$ -energy. Let us consider the complex case. Let f be a smooth map from \mathbb{C}^n to a Kähler manifold N . Let J be the standard complex structure of \mathbb{C}^n and J' the complex structure of N . Let ω and ω^N be the corresponding Kähler forms of \mathbb{C}^n and N (i.e., $\omega(\cdot, \cdot) = \langle J\cdot, \cdot \rangle$ and $\omega^N(\cdot, \cdot) = \langle J'\cdot, \cdot \rangle$). The $\bar{\partial}$ -energy density is defined by

$$\begin{aligned} e''(f) &= |\bar{\partial}f|^2 = |f_*J - J'f_*|^2 \\ &= \frac{1}{4}(|f_*e_i|^2 + |f_*J e_i|^2 - 2\langle J'f_*e_i, f_*J e_i \rangle) \\ &= \frac{1}{2}(e(f) - \langle f^*\omega^N, \omega^M \rangle), \end{aligned}$$

where $\{e_i, J e_i\}$ ($i = 1, \dots, n$) is the Hermitian frame of \mathbb{C}^n and $\langle f^*\omega^N, \omega \rangle$ denotes the induced norm. We say that f is holomorphic if $f_*J = J'f_*$. Obviously, f is holomorphic if and only if $|\bar{\partial}f|^2 \equiv 0$.

It is well known that a holomorphic map between two Kähler manifolds must be harmonic (see [Xin 1996]).

We denote the $\bar{\partial}$ -energy by

$$E_{\bar{\partial}}(f) = \int_{\mathbb{C}^n} |\bar{\partial}f|^2 \, dv.$$

3. $\bar{\partial}$ -energy variation

Let us consider the one-parameter family of maps $f_t(x) = f(tx) : \mathbb{C}^n \rightarrow N$, $t \in (1 - \epsilon, 1 + \epsilon)$ and $f_1 = f$. Let B_R denote the Euclidean ball in \mathbb{C}^n of radius R around 0. We write

$$E(R, t) = \int_{B_R} |\bar{\partial}f_t|^2 \, dv.$$

Lemma 3.1. $E(R, t) = t^{2-2n} E(Rt, 1)$.

Proof. Under the standard Hermitian metric of \mathbb{C}^n , $g^{ij} = \delta_{ij}$, from (2-1) we have

$$e(f_t)(x) = t^2 e(f)(tx).$$

By using the natural coordinates, it is easy to show that

$$\langle f_t^* \omega^N, \omega \rangle(x) = t^2 \langle f^* \omega^N, \omega \rangle(tx).$$

So we get

$$|\bar{\partial} f_t|^2(x) = t^2 |\bar{\partial} f|^2(tx).$$

It is easy to check that

$$\int_{B_R} |\bar{\partial} f_t|^2 dv = t^{2-2n} \int_{B_{Rt}} |\bar{\partial} f|^2 dv.$$

Thus we obtain the lemma. □

We now prove the following variation formula for $\bar{\partial}$ -energy:

Lemma 3.2. $\left. \frac{\partial E(R, t)}{\partial t} \right|_{t=1} = \frac{R}{2} \int_{\partial B_R} \left(\left| f_* \frac{\partial}{\partial r} \right|^2 - \left\langle J' f_* \frac{\partial}{\partial r}, f_* J \frac{\partial}{\partial r} \right\rangle \right) dv.$

Proof. Let $\{e_1, \dots, e_{2n} = \partial/\partial r\}$ be a local orthonormal frame field, where $\partial/\partial r$ denotes the unit radial vector field. By the definition of $f_t(x)$, it is easy to see that the variation vector field of f_t at $t = 1$ is

$$v = \left. \frac{df_t}{dt} \right|_{t=1} = r f_* \frac{\partial}{\partial r}.$$

The proof is separated into two steps.

Step 1. From (2-2), we have

$$\begin{aligned} \left. \frac{d}{dt} \int_{B_R} e(f_t) dv \right|_{t=1} &= \int_{B_R} \operatorname{div}(v, f_* e_j) e_j dv - \int_{B_R} \langle v, \tau(f) \rangle dv \\ &= \int_{\partial B_R} \left\langle v, f_* \frac{\partial}{\partial r} \right\rangle dv = R \int_{\partial B_R} \left| f_* \frac{\partial}{\partial r} \right|^2 dv. \end{aligned}$$

Since f is harmonic, we know that the tension field $\tau(f)$ is 0, and the second equality follows from the divergence theorem.

Step 2. On the other hand, from [Xin 1996] we know that $(d/dt)f_t^*\omega^N = d\theta_t$, where $\theta_t = f_t^*i(f_{t*}\partial/\partial t)\omega^N$. Since $(df_t/dt)|_{t=1} = rf_*\partial/\partial r$, we get $\theta_1 = \theta = rf^*i(f_*\partial/\partial r)\omega^N$. Then

$$\begin{aligned}
\frac{d}{dt} \int_{B_R} \langle f_t^*\omega^N, \omega \rangle dv \Big|_{t=1} &= \int_{B_R} \langle d\theta, \omega \rangle dv \\
&= \int_{B_R} d(\theta \wedge *\omega) + \int_{B_R} \langle \theta, \delta\omega \rangle dv \\
&= \int_{\partial B_R} \theta \wedge *\omega - \int_{B_R} \langle \theta, *d\omega^{n-1} \rangle dv \\
&= \int_{\partial B_R} \theta \wedge *\omega \\
&= - \int_{\partial B_R} \theta(e_i)\omega\left(e_i, \frac{\partial}{\partial r}\right) dv \\
&= -R \int_{\partial B_R} \omega^N\left(f_*\frac{\partial}{\partial r}, f_*e_i\right)\omega\left(e_i, \frac{\partial}{\partial r}\right) dv \\
&= -R \int_{\partial B_R} \left\langle J'f_*\frac{\partial}{\partial r}, f_*e_i \right\rangle \left\langle Je_i, \frac{\partial}{\partial r} \right\rangle dv \\
&= R \int_{\partial B_R} \left\langle J'f_*\frac{\partial}{\partial r}, f_*J\frac{\partial}{\partial r} \right\rangle dv.
\end{aligned}$$

Noting that $\langle d\theta, \omega \rangle dv = d\theta \wedge *\omega$, the second equality follows from the differential rules, where δ and $*$ are the codifferential and star operators. By Stokes' theorem and the definition of δ , the third equality holds. The fifth equality follows from direct computation. Since we may choose $e_1 = J\partial/\partial r$, the last equality holds.

Combining Steps 1 and 2, we obtain

$$\frac{d}{dt} \int_{B_R} |\bar{\partial}f_t|^2 dv \Big|_{t=1} = \frac{R}{2} \int_{\partial B_R} \left(\left| f_*\frac{\partial}{\partial r} \right|^2 - \left\langle J'f_*\frac{\partial}{\partial r}, f_*J\frac{\partial}{\partial r} \right\rangle \right) dv. \quad \square$$

Remark 3.3. If M is a compact manifold, $\int_M \langle f^*\omega^N, \omega^M \rangle dv$ is a homotopy invariant. This was observed first by Lichnerowicz [1970].

4. Proof of Theorem 1.1

We use a similar trick to [Sealey 1982].

By Lemma 3.1, we obtain

$$\frac{\partial E(R, t)}{\partial t} \Big|_{t=1} = (2 - 2n)E(R, 1) + R \frac{\partial E(R, 1)}{\partial R}.$$

On the other hand, from Lemma 3.2 and the condition (1-1), one has

$$\begin{aligned} \frac{\partial E(R, t)}{\partial t} \Big|_{t=1} &= \frac{R}{2} \int_{\partial B_R} \left(\left| f_* \frac{\partial}{\partial r} \right|^2 - \left\langle J' f_* \frac{\partial}{\partial r}, f_* J \frac{\partial}{\partial r} \right\rangle \right) dv \\ &\geq \frac{R}{2} \int_{\partial B_R} \left(\left| f_* \frac{\partial}{\partial r} \right|^2 - \left| f_* \frac{\partial}{\partial r} \right| \left| f_* J \frac{\partial}{\partial r} \right| \right) dv \\ &\geq -\frac{R}{2} \int_{\partial B_R} \left| f_* \frac{\partial}{\partial r} \right| \left| \left| f_* \frac{\partial}{\partial r} \right| - \left| f_* J \frac{\partial}{\partial r} \right| \right) dv \\ &\geq -\frac{R}{2} R^{2n-1} \frac{1}{R^{2n+\alpha/2}} C \\ &= -\frac{C}{2} R^{-\alpha/2}, \end{aligned}$$

where C is a positive constant. Hence for any $\epsilon > 0$, there exists an R_0 such that

$$\frac{\partial E(R, t)}{\partial t} \Big|_{t=1} \geq -\epsilon$$

for all $R \geq R_0$. Therefore

$$R \frac{\partial E(R, 1)}{\partial R} \geq -\epsilon + (2n - 2)E(R, 1)$$

for $R \geq R_0$.

If $E(\infty, 1) = \int_{\mathbb{C}^n} |\bar{\partial}f|^2 dv = E > 0$, then there exists an R_1 such that for all $R \geq R_1$ we have $E(R, 1) \geq E_0 > 0$. Since $n \geq 2$, we can choose a sufficiently small ϵ such that

$$R \frac{\partial E(R, 1)}{\partial R} \geq A = -\epsilon + (2n - 2)E_0 > 0$$

when $R \geq R_2 = \max(R_0, R_1)$. Then

$$E(\infty, 1) = \int_{\mathbb{C}^n} |\bar{\partial}f|^2 dv \geq \int_{R_2}^{\infty} \frac{A}{R} dR = \infty.$$

This is a contradiction. Therefore $\int_{\mathbb{C}^n} |\bar{\partial}f|^2 dv = 0$. Hence f is a holomorphic map.

Remark 4.1. Compared with the real case [Sealey 1982], Lemma 3.2 has the term $\langle J' f_*(\partial/\partial r), f_* J(\partial/\partial r) \rangle$. We need to use condition (1-1) to control it.

Remark 4.2. If we consider the ∂ -energy density $e'(f) = |\partial f|^2 = |f_* J + J' f_*|^2$, the corresponding result of Theorem 1.1 also holds; i.e., if the condition (1-1) is replaced by $e(f)e'(f)(p) = O(1/R^{4n+\alpha})$, then the conclusion is that f is a conjugate holomorphic map ($|\partial f|^2 \equiv 0$).

References

- [Dong 2013] Y. Dong, “Monotonicity formulae and holomorphicity of harmonic maps between Kähler manifolds”, *Proc. Lond. Math. Soc.* (3) **107**:6 (2013), 1221–1260. MR 3149846 Zbl 1295.53070
- [Lichnerowicz 1970] A. Lichnerowicz, “Applications harmoniques et variétés Kähleriennes”, pp. 341–402 in *Symposia Mathematica* (Rome, 1968/69), vol. III, Academic Press, London, 1970. MR 41 #7598 Zbl 0193.50101
- [Sealey 1982] H. C. J. Sealey, “Some conditions ensuring the vanishing of harmonic differential forms with applications to harmonic maps and Yang–Mills theory”, *Math. Proc. Cambridge Philos. Soc.* **91**:3 (1982), 441–452. MR 83i:58038 Zbl 0494.58002
- [Siu and Yau 1980] Y. T. Siu and S. T. Yau, “Compact Kähler manifolds of positive bisectional curvature”, *Invent. Math.* **59**:2 (1980), 189–204. MR 81h:58029 Zbl 0442.53056
- [Wan 2010] J. Wan, *Harmonic maps and harmonic complex structures*, thesis, Zhejiang University, Hangzhou, 2010.
- [Xin 1985] Y. L. Xin, “Holomorphicity of a class of harmonic maps”, *Acta Math. Sinica* **28**:3 (1985), 382–386. In Chinese. MR 87c:58035 Zbl 0598.58018
- [Xin 1996] Y. L. Xin, *Geometry of harmonic maps*, Progress in Nonlinear Differential Equations and their Applications **23**, Birkhäuser, Boston, 1996. MR 97h:58052 Zbl 0848.58014

Received May 6, 2014. Revised August 31, 2014.

JIANMING WAN
DEPARTMENT OF MATHEMATICS
NORTHWEST UNIVERSITY
XI'AN, 710127
CHINA
wanj_m@aliyun.com

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 275 No. 1 May 2015

Constant-speed ramps	1
OSCAR M. PERDOMO	
Surfaces in \mathbb{R}_+^3 with the same Gaussian curvature induced by the Euclidean and hyperbolic metrics	19
NILTON BARROSO and PEDRO ROITMAN	
Cohomology of local systems on the moduli of principally polarized abelian surfaces	39
DAN PETERSEN	
On certain dual q -integral equations	63
OLA A. ASHOUR, MOURAD E. H. ISMAIL and ZEINAB S. MANSOUR	
On a conjecture of Erdős and certain Dirichlet series	103
TAPAS CHATTERJEE and M. RAM MURTY	
Normal forms for CR singular codimension-two Levi-flat submanifolds	115
XIANGHONG GONG and JIŘÍ LEBL	
Measurements of Riemannian two-disks and two-spheres	167
FLORENT BALACHEFF	
Harmonic maps from \mathbb{C}^n to Kähler manifolds	183
JIANMING WAN	
Eigenvarieties and invariant norms	191
CLAUS M. SORENSEN	
The Heegaard distances cover all nonnegative integers	231
RUIFENG QIU, YANQING ZOU and QILONG GUO	



0030-8730(201505)275:1;1-0