

*Pacific
Journal of
Mathematics*

**CONVEX SOLUTIONS TO THE
POWER-OF-MEAN CURVATURE FLOW**

SHIBING CHEN

CONVEX SOLUTIONS TO THE POWER-OF-MEAN CURVATURE FLOW

SHIBING CHEN

We prove some estimates for convex ancient solutions (the existence time for the solution starts at $-\infty$) to the power-of-mean curvature flow, when the power is strictly greater than $\frac{1}{2}$. As an application, we prove that in dimension two, the blow-down of an entire convex translating solution, namely $u_h = \frac{1}{h}u(\frac{1}{h^{1+\alpha}}x)$, locally uniformly converges to $\frac{1}{1+\alpha}|x|^{1+\alpha}$ as $h \rightarrow \infty$. Another application is that for the generalized curve shortening flow (convex curve evolving in its normal direction with speed equal to a power of its curvature), if the convex compact ancient solution sweeps the whole space \mathbb{R}^2 , it must be a shrinking circle. Otherwise the solution must be defined in a strip region.

1. Introduction

Classifying ancient convex solutions to mean curvature flow is very important in studying the singularities of mean curvature flow. Translating solutions arise as a special case of ancient solutions when one uses a proper procedure to blow up the mean convex flow near type II singular points, and general ancient solutions arise at general singularities. Some important progress was made by Wang [2011], and Daskalopoulos, Hamilton and Sesum [Daskalopoulos et al. 2010]. Wang proved that in dimension $n = 2$, an entire convex translating solution to mean curvature flow must be rotationally symmetric in an appropriate coordinate system, which was a conjecture formulated explicitly by White [2000], but for $n \geq 3$ such solutions are not necessarily rotationally symmetric.

Wang also constructed some entire convex translating solutions with level sets neither spherical nor cylindrical in dimension greater or equal to 3. In the same paper, Wang also proved that if a convex ancient solution to the curve shortening flow sweeps the whole space \mathbb{R}^2 , then it must be a shrinking circle—otherwise the convex ancient solution must be defined in a strip region, and he indeed constructed such solutions by a compactness argument. Daskalopoulos et al. [2010] showed that apart from the shrinking circle, the so called *Angenent oval* (a convex ancient

MSC2010: 35J60.

Keywords: mean curvature flow, convexity, translating solution, ancient solution.

solution of the curve shortening flow discovered by Angenent that decomposes into two translating solutions of the flow) is the only other embedded convex compact ancient solution of the curve shortening flow. That means that the corresponding curve shortening solution defined in a strip region constructed by Wang is exactly the “Angenent oval”.

The power-of-mean curvature flow, in which a hypersurface evolves in its normal direction with speed equal to a power α of its mean curvature H , is well-studied [Andrews 1998; 2003; 2002; Schulze 2005; Chou and Zhu 2001; Sheng and Wu 2009]. Schulze [2005] called it H^α -flow. In the following, we will also call the one dimensional power-of-curvature flow the *generalized curve shortening flow*. It would be very interesting if one could classify the ancient convex solutions. In this paper, we use the method developed in [Wang 2011] to study the geometric asymptotic behavior of ancient convex solutions to H^α -flow. The general equation for H^α -flow is

$$\frac{\partial F}{\partial t} = -H^\alpha v,$$

where $F : M \times [0, T) \rightarrow \mathbb{R}^{n+1}$ is a time-dependent embedding of the evolving hypersurface, v is the unit normal vector to the hypersurface $F(M, t)$ in \mathbb{R}^{n+1} , and H is its mean curvature. If the evolving hypersurface can be represented as a graph of a function $u(x, t)$ over some domain in \mathbb{R}^n , then we can project the evolution equation to the $(n + 1)$ -st coordinate direction of \mathbb{R}^{n+1} and the equation becomes

$$u_t = \sqrt{1 + |Du|^2} \left(\operatorname{div} \frac{Du}{\sqrt{1 + |Du|^2}} \right)^\alpha.$$

Then a translating solution to the H^α -flow will satisfy the equation

$$\sqrt{1 + |Du|^2} \left(\operatorname{div} \frac{Du}{\sqrt{1 + |Du|^2}} \right)^\alpha = 1,$$

which is equivalent to the special case $\sigma = 1$ of the following:

$$\begin{aligned} (1) \quad L_\sigma(u) &= (\sqrt{\sigma + |Du|^2})^{\frac{1}{\alpha}} \operatorname{div} \frac{Du}{\sqrt{\sigma + |Du|^2}} \\ (2) \quad &= (\sigma + |Du|^2)^{\frac{1}{2\alpha} - \frac{1}{2}} \sum_{i,j=1}^n \left(\delta_{ij} - \frac{u_i u_j}{\sigma + |Du|^2} \right) u_{ij} \\ (3) \quad &= 1, \end{aligned}$$

where $\sigma \in [0, 1]$, $\alpha \in (\frac{1}{2}, \infty]$ is a constant, $n = 2$ is the dimension of \mathbb{R}^2 . If u is a convex solution of (3), then $u + t$, as a function of $(x, t) \in \mathbb{R}^2 \times \mathbb{R}$, is a translating

solution to the flow

$$(4) \quad u_t = \sqrt{\sigma + |Du|^2} \left(\operatorname{div} \frac{Du}{\sqrt{\sigma + |Du|^2}} \right)^\alpha.$$

When $\sigma = 1$, Equation (4) is the nonparametric power-of-mean curvature flow. When $\sigma = 0$, Equation (3) is the level set flow. That is, if u is a solution of (3) with $\sigma = 0$, then the level set $\{u = -t\}$, where $-\infty < t < -\inf u$, evolves by the power-of-mean curvature.

In the following we will assume $\sigma \in [0, 1]$, $\alpha \in (\frac{1}{2}, \infty]$, and the dimension $n = 2$, although some of the estimates do hold in higher dimension. The main results of this paper are the following theorems.

Theorem 1. *Let u be an entire convex solution of (3). Let*

$$u_h(x) = h^{-1}u(h^{\frac{1}{1+\alpha}}x).$$

Then, u_h locally uniformly converges to

$$\frac{1}{1+\alpha}|x|^{1+\alpha} \quad \text{as } h \rightarrow \infty.$$

Theorem 2. *Let u_σ be an entire convex solution of (3). Then,*

$$u_0(x) = \frac{1}{1+\alpha}|x|^{1+\alpha}$$

up to a translation of the coordinate system. When $\sigma \in (0, 1]$, if

$$|D^2u(x)| = O(|x|^\beta) \quad \text{as } |x| \rightarrow \infty$$

for some fixed constant β satisfying $\beta < 3\alpha - 2$, then u_σ is rotationally symmetric after a proper translation of the coordinate system.

Corollary 3. *A convex compact ancient solution to the generalized curve shortening flow which sweeps the whole space \mathbb{R}^2 must be a shrinking circle.*

Remark 4. The condition $\alpha > \frac{1}{2}$ is necessary for our results. One can consider the translating solution $v(x)$ to (3) with $\sigma = 1$ in one dimension. In fact, when $\alpha \leq \frac{1}{2}$, the translating solution $v(x)$ is a convex function defined on the entire real line [Chou and Zhu 2001, p. 28]. Then one can construct a function $u(x, y) = v(x) - y$, defined on the entire plane, and u will satisfy (3) with $\sigma = 0$; it is obviously not rotationally symmetric. We can also let $u(x, y) = v(x)$, which is an entire solution to (3) with $\sigma = 1$, and it is not rotationally symmetric.

When the dimension is at least two, similar examples can be given: we can take an entire rotationally symmetric solution $v(x)$ to (3) with $n \geq 2$ and $\sigma = 1$, and again let $u(x, y) = v(x) - y$ (here, y is the $(n + 1)$ -st coordinate for \mathbb{R}^{n+1}). It is easy to see that u will satisfy (3) with n replaced by $n + 1$ and $\sigma = 0$, and the level set of u is neither a sphere nor a cylinder.

We would also like to point out that this elementary construction can be used to give a slight simplification of the proof of [Wang 2011, Theorem 2.1] (corresponding to our Corollary 10 for $\alpha = 1$). Let v_σ be an entire convex solution to (3) in dimension n with $\sigma \in (0, 1]$. Then $u(x, y) = v_\sigma(x) - \sqrt{\sigma}y$ will be an entire convex solution to (3) in dimension $n + 1$ with $\sigma = 0$. Hence if one has proved the estimate in Corollary 10 for $\sigma = 0$ in all dimensions, the estimate for $\sigma \in (0, 1]$ follows immediately from the above construction. The remainder of the paper is divided into four sections. Sections 2 and 3 contain the proof of Theorem 1 and the first part of Theorem 2. Section 4 is devoted to the proof of Corollary 10, and the last section completes the proof of Theorem 2.

2. Power growth estimate

In this section, we prove a key estimate, which says that any entire convex solution u to (3) must satisfy

$$u(x) \leq C(1 + |x|^{1+\alpha}),$$

where the constant C depends only on the upper bound of $u(0)$ and $|Du(0)|$. When $\alpha = 1$, the estimate was proved by Wang [2011, Theorem 2.1]. To apply Wang's method, the main difficulty is that now the speed function is nonlinear in the curvature. We overcome this difficulty by further exploiting some elementary convexity properties.

For any constant $h > 0$, we denote

$$\Gamma_h = \{x \in \mathbb{R}^n : u(x) = h\},$$

$$\Omega_h = \{x \in \mathbb{R}^n : u(x) < h\},$$

so that Γ_h is the boundary of Ω_h . Let κ be the curvature of the level curve Γ_h . We have

$$(5) \quad L_\sigma(u) = (\sigma + u_\gamma^2)^{\frac{1}{2\alpha} - \frac{1}{2}} \left(\kappa u_\gamma + \frac{\sigma u_{\gamma\gamma}}{\sigma + u_\gamma^2} \right)$$

$$(6) \quad \geq \kappa u_\gamma^{\frac{1}{\alpha}} = L_0(u),$$

where γ is the unit outward normal to Ω_h , and $u_{\gamma\gamma} = \gamma_i \gamma_j u_{ij}$.

Before starting the proof of our main results, we recall a well known convergence result for the generalized curve shortening flow.

Lemma 5 [Andrews 2003, Theorems 1.3, 1.4, 1.5]. *Let ℓ_t be a time-dependent family of closed curves in \mathbb{R}^2 evolving under the generalized curve shortening flow with $\alpha > \frac{1}{3}$. Suppose the initial curve ℓ_0 is convex. Then the curve converges in finite time T to a round point P in the sense that $((1 + \alpha)(T - t))^{-\frac{1}{1+\alpha}} (\ell_t - P)$ is asymptotic to the unit circle.*

Next, we prove a lemma which will be used to control the shape of the level set of a complete convex solution to (3).

Lemma 6. *Let u be a complete convex solution of (3). Suppose that $u(0) = 0$ and that the infimum $\inf\{|x| : x \in \Gamma_1\}$ is attained at $x_0 = (0, -\delta) \in \Gamma_1$ for some $\delta > 0$ sufficiently small. Let D_1 be the projection of Γ_1 onto the axis $\{x_2 = 0\}$. Then, D_1 contains the interval $(-R, R)$, and when $\alpha \leq 1$, R satisfies*

$$(7) \quad R \geq C_1(-\log \delta - C_2)^{\frac{\alpha}{\alpha+1}},$$

where $C_1, C_2 > 0$ are independent of δ ; when $\alpha > 1$, $R \geq C$ for some positive constant C .

The proof of this lemma follows that of [Wang 2011, Lemma 2.4] with minor modifications; the for reader’s convenience, we give some details here.

Proof. First, we prove the lemma when $\frac{1}{2} < \alpha \leq 1$. Suppose that near x_0 , Γ_1 is given by $x_2 = g(x_1)$. Then, g is a convex function satisfying $g(0) = -\delta$ and $g'(0) = 0$. Let $b > 0$ be a constant such that $g'(b) = 1$. To prove (7), it suffices to prove

$$(8) \quad b \geq C_1(-\log \delta - C_2)^{\frac{\alpha}{\alpha+1}}.$$

For any $y = (y_1, y_2) \in \Gamma_1$, where $y_1 \in [0, b]$, as in the proof of [Wang 2011, Lemma 2.4] we have

$$(9) \quad u_\gamma(y) \geq \frac{\sqrt{1 + g'^2}}{y_1 g' - y_2},$$

where γ is the unit normal of the sublevel set Ω_1 . Since $L_0 u \leq 1$, we have

$$(10) \quad \frac{g''}{(1 + g'^2)^{\frac{3}{2}}} \frac{(1 + g'^2)^{\frac{1}{2\alpha}}}{(y_1 g' - y_2)^{\frac{1}{\alpha}}} \leq \kappa u_\gamma^{\frac{1}{\alpha}} \leq 1,$$

where κ is the curvature of the level curve Γ_1 . Hence,

$$(11) \quad g''(y_1) \leq (1 + g'^2)^{\frac{3}{2} - \frac{1}{2\alpha}} (y_1 g' - y_2)^{\frac{1}{\alpha}}$$

$$(12) \quad \leq 10y_1^{\frac{1}{\alpha}} g' + 10\delta$$

where $y_2 = g(y_1)$ and $g'(y_1) \leq 1$ for $y_1 \in (0, b)$. The inequality from (11) to (12) is trivial when $y_2 \geq 0$. When $y_2 \leq 0$, since $|y_2| \leq \delta$, we have either $y_1 g' \leq \delta$ or $y_1 g' > \delta$. For the former we have

$$(y_1 g' - y_2)^{\frac{1}{\alpha}} \leq (2\delta)^{\frac{1}{\alpha}} \leq 4\delta;$$

for the latter, since $g'(y_1) \leq 1$, we have

$$(y_1 g' - y_2)^{\frac{1}{\alpha}} \leq (2y_1 g')^{\frac{1}{\alpha}} \leq 4y_1^{\frac{1}{\alpha}} g'.$$

We consider the equation

$$(13) \quad \rho''(t) = 10t^{\frac{1}{\alpha}}\rho' + 10\delta$$

with initial conditions $\rho(0) = -\delta$ and $\rho'(0) = 0$. Then for $t \in (0, b)$, we have

$$(14) \quad \rho'(t) = 10\delta e^{\frac{10\alpha}{\alpha+1}t^{\frac{\alpha+1}{\alpha}}} \int_0^t e^{-\frac{10\alpha}{\alpha+1}s^{\frac{\alpha+1}{\alpha}}} ds.$$

Since $\int_0^\infty e^{-\frac{10\alpha}{\alpha+1}s^{\frac{\alpha+1}{\alpha}}} ds$ is bounded above by some constant C , we have

$$(15) \quad 1 \leq \rho'(b) = 10\delta e^{\frac{10\alpha}{\alpha+1}b^{\frac{\alpha+1}{\alpha}}} \int_0^b e^{-\frac{10\alpha}{\alpha+1}s^{\frac{\alpha+1}{\alpha}}} ds$$

$$(16) \quad \leq C_1 \delta e^{\frac{10\alpha}{\alpha+1}b^{\frac{\alpha+1}{\alpha}}},$$

from which (8) follows.

When $\alpha > 1$, the situation is different. First, we introduce a number a such that $g'(a) = \frac{1}{2}$. Then, we can follow the proof above until (11). For (12) the inequality becomes

$$g''(y_1) \leq 10y_1^{\frac{1}{\alpha}}g' + 10\delta^{\frac{1}{\alpha}},$$

for $y_1 \in [a, b]$. Now (16) becomes

$$e^{-\frac{10\alpha}{\alpha+1}b^{\frac{\alpha+1}{\alpha}}}\rho'(b) - e^{-\frac{10\alpha}{\alpha+1}a^{\frac{\alpha+1}{\alpha}}}\rho'(a) \leq C_1\delta^{\frac{1}{\alpha}}.$$

Then, it is easy to see that when δ is small, $b \geq C$, for some fixed constant C . \square

Remark 7. When $\alpha \leq 1$, it follows from Lemma 6 that when δ is sufficiently small, by convexity and in view of Figure 1, we have that Ω_1 contains the shadowed region. Then it is easy to check that Ω_1 contains an ellipse

$$(17) \quad E = \left\{ (x_1, x_2) \mid \frac{x_1^2}{\left(\frac{R}{6}\right)^2} + \frac{\left(x_2 - \frac{7\delta^* - 5\delta}{12}\right)^2}{\left(\frac{\delta^* + \delta}{4}\right)^2} = 1 \right\},$$

where δ^* is a positive constant such that $u(0, \delta^*) = 1$ and R is defined in Lemma 5.

When $\alpha > 1$, if δ^* is very large, in the part $\{x : u(x) \leq 1, x_1 \geq 0\}$, by convexity we can find an ellipse which has the length of short axis bounded from below and the length of long axis very large, and if we let the ellipse evolve under the generalized curve shortening flow, it will take time more than 1 for it to converge to a round point. When δ^* is less than some fixed constant, we need to consider two cases.

Case 1: The set $\{u \leq 1\}$ is not compact. In this case when we project $\{u(x) = 1\}$ to the axis $\{x_2 = 0\}$, and denote the leftmost and rightmost points as $(-l, 0)$ and $(r, 0)$, respectively. Then either l or r is very large, which guarantees that one can




Figure 1. Γ_1 contains the shadow part.

still find an ellipse inside $\{x : u(x) \leq 1, x_1 \leq 0\}$ (or $\{x : u(x) \leq 1, x_1 \geq 0\}$) with the similar property as before.

Case 2: The set $\{u \leq 1\}$ is compact. For this case, we will always assume 0 is the minimum point of u , and $u(0) = 0$. We claim that when δ is very small, for the purpose of the proof of [Corollary 10](#), we can assume one of l or r is very large. Indeed, if the claim is not true, then we have a sequence of functions u_i such that $\{u_i \leq 1\}$ has width bounded by some constant independent of i , and $\text{dist}(0, \{u_i \leq 1\}) \rightarrow 0$ as $i \rightarrow \infty$. In view of the following proof of [Corollary 10](#), we can assume u_i satisfies (3) with $\sigma_i \rightarrow 0$. Then by passing to a subsequence, we can assume that $\{u_i = 1\}$ converges to a convex curve C_0 in hausdorff distance. Let C_0 evolve under the generalized curve shortening flow; by [Lemma 5](#), it will converge to a point P , but by the above discussion we see that P is on C_0 , which is clearly impossible. Once l or r is very large, we can find an ellipse with the similar property as in the case 1.

Remark 8. One can also establish a similar lemma in higher dimensions, which says that D_1 (a convex set with dimension greater than 1) contains a ball centered at the origin with radius

$$R \geq C_n(-\log \delta - C)^{\frac{\alpha}{\alpha+1}},$$

where C_n is a constant depending only on n and C is a positive constant independent of δ . The proof can be reduced to the two dimensional case; for the details, refer to the proof of [[Wang 2011](#), Lemma 2.6].

Lemma 9. *Let u be a complete convex solution of (3). Suppose $u(0) = 0$, δ and δ^* are defined as in [Lemma 6](#) and [Remark 7](#). If δ and δ^* are sufficiently small, then u is defined in a strip region.*

When $\alpha = 1$, this lemma is proved by Wang [2011, Corollary 2.2]. The proof of Lemma 9 is based on a careful study of the shape of the level set of u . Before giving the proof, we will give an important corollary first.

Corollary 10. *Let u be an entire convex solution of (3) in \mathbb{R}^2 , then*

$$(18) \quad u(x) \leq C(1 + |x|^{1+\alpha}),$$

where the constant C depends only on the upper bound for $u(0)$ and $|Du(0)|$.

Proof. The proof of this Corollary follows the proof of [Wang 2011, Theorem 2.1]. We only record some necessary changes here. First, the rescaling $u_h(x) = \frac{1}{h}u(h^{\frac{1}{2}}x)$ used that proof should be replaced by $u_h(x) = \frac{1}{h}u(h^{\frac{1}{1+\alpha}}x)$. Note that u_h solves (3) with $\sigma = \sigma_h \rightarrow 0$ as $h \rightarrow \infty$. Second, the ellipse used in that proof when applying the comparison argument should be replaced by the one discussed in Remark 7. \square

Proof of Lemma 9. By a rotation of coordinates we may assume that the axial directions of E in Remark 7 coincide with those of the coordinate system. Let \mathcal{M}_u be the graph of u , which consists of two parts, $\mathcal{M}_u = \mathcal{M}^+ \cup \mathcal{M}^-$, where

$$\mathcal{M}^+ = \{(x, u(x)) \in \mathbb{R}^3 : \partial_{x_2}u \geq 0\} \quad \text{and} \quad \mathcal{M}^- = \{(x, u(x)) \in \mathbb{R}^3 : \partial_{x_2}u \leq 0\}.$$

Then \mathcal{M}^\pm can be represented as the graphs of functions g^\pm of the form $x_2 = g^\pm(x_1, x_3)$, for $(x_1, x_3) \in D$ where D is the projection of \mathcal{M}_u onto the plane $\{x_2 = 0\}$. The functions g^+ and g^- are concave and convex, respectively, and we have $x_3 = u(x_1, g^\pm(x_1, x_3))$. Set

$$(19) \quad g = g^+ - g^-.$$

Then g is a positive, concave function on D , vanishing on ∂D . For any $h > 0$ let $g_h(x_1) = g(x_1, h)$, $g_h^\pm(x_1) = g^\pm(x_1, h)$, and $D_h = \{x_1 \in \mathbb{R}^1 : (x_1, h) \in D\}$. Then g_h is a positive, concave function in D_h , vanishing on ∂D_h , and $D_h = (-\underline{a}_h, \bar{a}_h)$ is an interval containing the origin. Let $b_h = g_h(0)$. We consider the case $\sigma = 0$ first.

Claim 1: Suppose h is large, $g_1(0) = \delta^* + \delta$ is small, $b_h \leq 4$, and $\underline{a}_h, \bar{a}_h \geq b_h$. Then,

$$\bar{a}_h \geq \frac{1}{1000} \frac{h}{b_h^\alpha} \quad \text{for } \alpha \leq 1 \quad \text{and} \quad \bar{a}_h \geq \frac{1}{1000} \frac{h^{1/(2\alpha-1)}}{b_h^{1/(2\alpha-1)}} \quad \text{for } \alpha > 1.$$

Proof. Without loss of generality, we assume $\bar{a}_h \leq \underline{a}_h$. Let $U_h = \Omega_h \cap \{x_1 > 0\}$. By the convexity of U_h and the assumption $\underline{a}_h, \bar{a}_h \geq b_h$, we have $\underline{a}_s, \bar{a}_s \geq \frac{1}{2}b_h$ for all $s \in (\frac{1}{2}h, h)$. Hence by the concavity of g ,

$$\left| \frac{d}{dx_1} g_s(0) \right| \leq 2 \quad \text{for } s \in \left(\frac{1}{2}h, h\right),$$




Figure 2. $\Gamma_s \cap \{x_1 > 0\}$ is trapped between two lines.

which means the arclength of the image of $\Gamma_s \cap \{x_1 > 0\}$ under the Gauss map is bigger than $\frac{\pi}{6}$. Notice that Ω_1 contains E , which was defined in Remark 7. When δ and δ^* are very small, E is very thin and long. The center of E is very close to the origin; in fact, for our purpose we can just pretend E is centered at the origin. By convexity of Ω_h and in view of Figure 2, we see that $\Gamma_s \cap \{x_1 > 0\}$ is trapped between two lines ℓ_1 and ℓ_2 , and the slopes of ℓ_1 and ℓ_2 are very close to 0 when E is very long and thin. Then it is clear that the largest distance from the points on $\Gamma_s \cap \{x_1 > 0\}$ to the origin can not be bigger than $10\bar{a}_h$. By convexity of u ,

$$u_\gamma(x) \geq \frac{h}{20\bar{a}_h} \quad \text{for } x \in \Gamma_s \cap \{x_1 > 0\}.$$

Since $\Gamma_s \cap \{x_1 > 0\}$ evolves under the generalized curve shortening flow, when $\alpha \leq 1$ we have the estimate

$$(20) \quad \frac{d}{ds} |U_s| = \int_{\Gamma_s \cap \{x_1 > 0\}} \kappa^\alpha d\xi$$

$$(21) \quad = \int_{\Gamma_s \cap \{x_1 > 0\}} u_\gamma^{\frac{1}{\alpha}-1} \kappa d\xi$$

$$(22) \quad \geq \frac{1}{50} \left(\frac{h}{\bar{a}_h} \right)^{\frac{1}{\alpha}-1} \frac{\pi}{6},$$

where from (20) to (21) we used the equation $\kappa u_\gamma^{\frac{1}{\alpha}} = 1$. The claim follows by the simple fact that

$$\frac{3}{2} b_h \bar{a}_h \geq |U_h| \geq \frac{1}{50} \left(\frac{h}{\bar{a}_h} \right)^{\frac{1}{\alpha}-1} \frac{\pi}{6} h.$$

When $\alpha > 1$, let l_s denote the arclength of $\Gamma_s \cap \{x_1 > 0\}$. Then, by the above discussion, it is not hard to see that $l_s \approx C\bar{a}_h$. Then by a simple application of

Jensen's inequality,

$$\begin{aligned} \frac{d}{ds}(|U_s|) &= \int_{\Gamma_s \cap \{x_1 > 0\}} \kappa^\alpha d\xi \\ &= l_s \int_{\Gamma_s \cap \{x_1 > 0\}} \kappa^\alpha \frac{1}{l_s} d\xi \\ &\geq l_s \left(\int_{\Gamma_s \cap \{x_1 > 0\}} \frac{\kappa}{l_s} d\xi \right)^\alpha \geq C l_s^{1-\alpha} \geq C \bar{a}_h^{1-\alpha}. \end{aligned}$$

Again by the simple fact that $\frac{3}{2} b_h \bar{a}_h \geq |U_h|$, we can complete the proof in the same way as in the previous case. \square

From here until (55) we will prove the lemma for the case $\frac{1}{2} < \alpha \leq 1$, and then we will give the details for the case $\alpha > 1$.

Claim 2: Let $h_k = 2^k$, $\bar{a}_k = \bar{a}_{h_k}$, $b_k = b_{h_k}$, $g_k = g_{h_k}$, and $D_k = D_{h_k}$. Then,

$$(23) \quad g_k(0) \leq g_{k-1}(0) + C_0 2^{-k/C} \quad \text{for all } k \text{ large,}$$

where C_0 is a fixed constant and C depends only on α .

Lemma 9 follows from Claims 1 and 2 in the following way. Let the convex set P be the projection of the graph of g onto the plane $\{x_3 = 0\}$. By Claim 2 and the fact that P contains x_1 -axis (it follows from Claim 1), P must equal $I \times \mathbb{R}$ for some interval

$$I \subset \left[0, \lim_{k \rightarrow \infty} g_k(0)\right].$$

Then, by (19), \mathcal{M}_u is also contained in a strip region as stated in Lemma 9.

Proof of Claim 2. To prove (23), observe that since g is positive and concave,

$$g_k(0) \leq h_k g_0(0) \leq 2^k (\delta + \delta^*).$$

Hence, we can start from sufficiently large k_0 , satisfying $g_{k_0}(0) \leq 1$ and

$$(24) \quad g_{k_0} + C_0 \sum_{j=k_0}^{\infty} 2^{-j/C} \leq 2.$$

Suppose (23) holds up to k . Then by (24), we have $g_k(0) \leq 2$. By the concavity of g and the fact that $g \geq 0$, we have $g_{k+1}(0) \leq 2g_k(0) \leq 4$. By Claim 1, we have $\bar{a}_{k+1} \geq \frac{1}{10000} h_k$. To prove (23) at $k+1$,

$$(25) \quad L_k = \left\{ x_1 \in \mathbb{R}^1 : -\frac{C_1}{4} h_k < x_1 < \frac{C_1}{4} h_k \right\}, \quad C_1 = \frac{1}{10000},$$

$$(26) \quad Q_k = L_k \times [h_k, h_{k+1}] \subset D.$$

Since $g > 0$ and g is concave, we have the estimates

$$(27) \quad g(x_1, h) \leq 8,$$

$$(28) \quad |\partial_h g(x_1, h)| \leq \frac{16}{h_k},$$

$$(29) \quad |\partial_{x_1} g(x_1, h)| \leq \frac{16}{h_k} \quad \text{for all } (x_1, h) \in Q_k.$$

Let $\mathcal{X}^\pm = \{(x_1, h) \in Q_k : |\partial_{x_1 x_1} g^\pm(x_1, h)| \geq h_k^{-\beta}\}$, where β is chosen such that $\frac{1}{\alpha} < \beta < 2$. For any $h \in (h_k, h_{k+1})$, by (29),

$$(30) \quad |\{x_1 \in L_k : (x_1, h) \in \mathcal{X}^+\}| h_k^{-\beta} \leq \int_{L_k} \partial_{x_1 x_1} g^+$$

$$(31) \quad \leq \int_{L_k} \partial_{x_1 x_1} g$$

$$(32) \quad \leq 2 \sup_{L_k} |\partial_{x_1} g|$$

$$(33) \quad \leq \frac{C}{h_k}.$$

So, $|\mathcal{X}^+| \leq C h_k^\beta$. Similarly, we have $|\mathcal{X}^-| \leq C h_k^\beta$.

For any given $y_1 \in L_k$, let $\mathcal{X}_{y_1}^\pm = \mathcal{X}^\pm \cap \{x_1 = y_1\}$. Then, by the estimate above, there is a set $\tilde{L}^\pm \subset L_k$ with measure

$$|\tilde{L}^\pm| \leq C h_k^{\beta/2},$$

such that for any $y_1 \in L_k - \tilde{L}^\pm$, we have $|\mathcal{X}_{y_1}^\pm| \leq h_k^{\beta/2}$. When k is large, we can always find $y_1 = C h^{\beta/2} \in L_k - \tilde{L}^\pm$, where the constant C is under control. For such y_1 , we have

$$(34) \quad g(y_1, h_{k+1}) - g(y_1, h_k) \\ = g^+(y_1, h_{k+1}) - g^+(y_1, h_k) + |g^-(y_1, h_{k+1}) - g^-(y_1, h_k)|.$$

In the following, we will estimate $g^+(y_1, h_{k+1}) - g^+(y_1, h_k)$. The estimate for $|g^-(y_1, h_{k+1}) - g^-(y_1, h_k)|$ is analogous. By the same reason as that for [Wang 2011, §2.21], we have

$$(35) \quad \begin{cases} (\partial_h g^+)^{-1} = (1 + \varepsilon_1) u_\gamma, \\ \partial_{x_1 x_1} g^+ = (1 + \varepsilon_2) \kappa. \end{cases}$$

Then, by the equation $u_\gamma^\alpha \kappa = 1$, we have

$$(36) \quad \partial_h g_h^+(y_1, h) \leq C (\partial_{x_1 x_1} g^+)^{\alpha} \leq C h_k^{-\beta \alpha}.$$

Now,

$$(37) \quad g^+(y_1, h_{k+1}) - g^+(y_1, h_k) = \int_{h_k}^{h_{k+1}} \partial_h g^+(y_1, h) dh$$

$$(38) \quad = \int_{\mathcal{X}_{y_1}^+} \partial_h g^+(y_1, h) dh + \int_{[h_k, h_{k+1}] - \mathcal{X}_{y_1}^+} dh$$

$$(39) \quad \leq C_1 h_k^{\frac{\beta}{2}} \frac{1}{h_k} + C_2 h_k^{-\beta\alpha} h_k.$$

Recall that β satisfies $\frac{1}{\alpha} < \beta < 2$, and we have $\eta := \min\{1 - \frac{\beta}{2}, \beta\alpha - 1\} > 0$. From (34) and (39), we have the estimate

$$g(y_1, h_{k+1}) - g(y_1, h_k) \leq \frac{C}{h_k^\eta},$$

for some fixed constant C . Then, we will assume $\partial_{x_1} g(0, h_k) < 0$ (otherwise we can replace x_1 by $-x_1$); therefore, by the above estimate,

$$g(y_1, h_{k+1}) \leq g(y_1, h_k) + \frac{C}{h_k^\eta} \leq g(0, h_k) + \frac{C}{h_k^\eta}.$$

Since g is positive, concave, and defined on $[0, \bar{a}_{k+1}]$, with $\bar{a}_{k+1} \geq Ch_{k+1}$,

$$\frac{g_{k+1}(0)}{g_{k+1}(y_1)} \leq \frac{\bar{a}_{k+1}}{\bar{a}_{k+1} - y_1} \leq 1 + Ch_{k+1}^{\frac{\beta}{2} - 1}.$$

Therefore, by the two estimates above,

$$g_{k+1}(0) \leq g_k(0) + Ch_k^{-\eta},$$

which implies (23) immediately. \square

For the proof of Lemma 6 when $\sigma \in (0, 1]$, we need to use (5) and (6). In fact, by (6) we see that Γ_h is moving at a velocity greater than or equal to its curvature to the power α . Hence, we still have the lower bound of $\frac{d}{ds}(|U_s|)$ as in the proof of Claim 1. Then we can follow the above proof for the case $\sigma = 0$ until (37), replacing the equalities “=” in (20) and (21) with inequalities “ \geq ”. As in [Wang 2011], when $\sigma = 0$, in order to control the second integral in (38) we used the equation $\kappa u_\gamma^{1/\alpha} = 1$. But when $\sigma \neq 0$, by (28) and (35) we have

$$(40) \quad u_\gamma \geq C(\partial_h g^+)^{-1} \geq Ch_k.$$

Hence, we may assume that u_γ is as large as we want, which means that in formula (5), the only important extra term is

$$(\sigma + u_\gamma^2)^{\frac{1}{2\alpha} - \frac{1}{2}} \frac{\sigma u_\gamma \gamma}{\sigma + u_\gamma^2}.$$

To handle this term, we divide the integral (39) into three parts:

$$(41) \quad g^+(y_1, h_{k+1}) - g^+(y_1, h_k) = \int_{h_k}^{h_{k+1}} \partial_h g^+(y_1, h) dh$$

$$(42) \quad = \left(\int_{I_1} + \int_{I_2} + \int_{I_3} \right) \partial_h g^+(y_1, h) dh,$$

where

$$(43) \quad I_1 = \mathcal{X}_{y_1}^+,$$

$$(44) \quad I_2 = \left\{ h \in [h_k, h_{k+1}] - I_1 : (\sigma + u_\gamma^2)^{\frac{1}{2\alpha} - \frac{1}{2}} \frac{\sigma u_{\gamma\gamma}}{\sigma + u_\gamma^2} \leq \frac{1}{2} \right\},$$

$$(45) \quad I_3 = [h_k, h_{k+1}] - I_1 \cup I_2.$$

For the first integral, we can do exactly the same thing as we have done from (38) to (39), namely,

$$\int_{I_1} \partial_h g^+(y_1, h) dh \leq \frac{C}{h_k} h_k^{\frac{\beta}{2}} = C h_k^{\frac{\beta}{2} - 1}.$$

Note that the power $\frac{\beta}{2} - 1$ is a negative number.

Then we estimate the second integral, note that when $(y_1, h) \in I_2$, we have

$$(\sigma + u_\gamma^2)^{\frac{1}{2\alpha}} \kappa u_\gamma \geq \frac{1}{2}.$$

By (40) u_γ is large, so we have $\kappa u_\gamma^{\frac{1}{\alpha}} \geq \frac{1}{4}$, hence by (35) we have

$$(46) \quad \partial_h g^+ \leq C(\partial_{x_1 x_1} g^+)^{\alpha} \leq C h_k^{-\alpha\beta}.$$

Therefore,

$$\int_{I_2} \partial_h g^+(y_1, h) dh \leq C h_k^{-\beta\alpha} h_k = C h_k^{1-\beta\alpha}.$$

Note that $1 - \beta\alpha$ is a negative number. Observe that we can assume I_2 is on the right hand side of I_3 , since by the concavity of g^+ we know that when $h \geq \inf I_2$, $\partial_h g^+(y_1, h)$ will satisfy the estimate (46).

For the third integral, notice that by the same argument as that for [Wang 2011, §2.24],

$$(47) \quad \begin{cases} u_\gamma(y_1, h) = u_{x_2}(1 + \varepsilon_1), \\ u_{\gamma\gamma}(y_1, h) = u_{x_2 x_2}(1 + \varepsilon_2) + \varepsilon_3 u_{x_2}. \end{cases}$$

Hence, by (47),

$$(48) \quad (\sigma + u_{x_2}^2)^{\frac{1}{2\alpha} - \frac{1}{2}} \frac{\sigma u_{x_2 x_2}}{\sigma + u_\gamma^2} \geq \frac{1}{3}.$$

Since $\sigma \in [0, 1]$ and u_γ is large, we have

$$(49) \quad u'' = u_{x_2 x_2} \geq \frac{1}{4}(u')^{3-\frac{1}{\alpha}}.$$

By differentiating the equation $u(x_1, g^+(x_1, h)) = h$ twice with respect to h ,

$$(50) \quad (g^+)'' = -u''(g^+)'^3 \leq -\frac{1}{4}(g^+)'^{\frac{1}{\alpha}-3}(g^+)^3 = -\frac{1}{4}(g^+)'^{\frac{1}{\alpha}}.$$

Note that (50) is for points with corresponding $h \in I_3$. By the discussion after (46) we only need to estimate

$$\int_{[h_k+h_k^{\frac{\beta+2}{4}}, \inf I_2]} (g^+)'' dh.$$

Therefore, by (50) and noticing that $(g^+)'' \geq 0$,

$$(51) \quad \frac{\alpha}{\alpha-1}(g^+)'^{\frac{\alpha-1}{\alpha}}(h) \leq \frac{\alpha}{\alpha-1}(g^+)'^{\frac{\alpha-1}{\alpha}}(h_k) - \frac{1}{4}|I_3 \cap [h_k, h]|.$$

Hence, when $h \in [h_k^{(\beta+2)/4}, \inf I_2]$,

$$(52) \quad (g^+)''(h) \leq ((g^+)'^{\frac{\alpha-1}{\alpha}}(h_k) + C(h-h_k))^{\frac{\alpha}{\alpha-1}}.$$

Finally,

$$(53) \quad \int_{[h_k+h_k^{\frac{\beta+2}{4}}, \inf I_2]} (g^+)'' dh \leq \int_{h_k}^{h_k^{h_k+1}} ((g^+)'^{\frac{\alpha-1}{\alpha}}(h_k) + C(h-h_k))^{\frac{\alpha}{\alpha-1}} dh$$

$$(54) \quad \leq \frac{\alpha-1}{2\alpha-1} ((g^+)'^{\frac{\alpha-1}{\alpha}}(h_k) + C(h-h_k))^{\frac{\alpha}{\alpha-1}+1} \Big|_{h_k}^{2h_k}$$

$$(55) \quad \leq C(g^+)'^{\frac{2\alpha-1}{\alpha}} \leq Ch_k^{\frac{1-2\alpha}{\alpha}}.$$

Note that $\frac{1-2\alpha}{\alpha} < 0$ when $\alpha > \frac{1}{2}$, so we can complete the proof as in the $\sigma = 0$ case.

When $\alpha > 1$, we need to choose the constants and exponents more carefully. First of all, in view of the Lemma 9 for $\alpha > 1$, in order to have properties (35) and (47), we need only to replace the number 2 in (24) with some number much smaller than the constant C in Lemma 9. The definition of L_k in (25) should be modified to

$$L_k = \left\{ x_1 \in \mathbb{R}^1 : -\frac{C_1}{4}h_k^{\frac{1}{2\alpha-1}} < x_1 < \frac{C_1}{4}h_k^{\frac{1}{2\alpha-1}} \right\}, \quad C_1 = \frac{1}{10000},$$

and the definition of Q_k in (26) remains the same. It is easy to see that we still have the estimates (27)–(28), but (29) becomes

$$|\partial_{x_1} g(x_1, h)| \leq 16h_k^{-\frac{1}{2\alpha-1}} \quad \text{for all } (x_1, h) \in Q_k.$$

Then for the definition of

$$\mathcal{Q}^\pm = \{(x_1, h) \in Q_k : |\partial_{x_1 x_1} g^\pm(x_1, h)| \geq h_k^{-\beta}\},$$

we need to choose the exponent β so that $\frac{1}{\alpha} < \beta < \frac{2}{2\alpha-1}$. By doing the same computation as (30)–(32),

$$|\{x_1 \in L_k : (x_1, h) \in \mathcal{X}^+\}| h_k^{-\beta} \leq \int_{L_k} \partial_{x_1 x_1} g^+ \leq C h_k^{-\frac{1}{2\alpha-1}}.$$

Hence,

$$|\mathcal{X}^+| \leq C h_k^{1+\beta-\frac{1}{2\alpha-1}} \quad \text{and} \quad |\mathcal{X}^-| \leq C h_k^{1+\beta-\frac{1}{2\alpha-1}}.$$

Then, by the above estimate there is a set $\tilde{L}^\pm \subset L_k$ with measure

$$|\tilde{L}^\pm| \leq C h_k^{\beta+\varepsilon-\frac{1}{2\alpha-1}}$$

such that for any $y_1 \in L_k - \tilde{L}^\pm$, we have $|\mathcal{X}_{y_1}^\pm| \leq h_k^{1-\varepsilon}$, where ε is chosen such that $\beta + \varepsilon < \frac{2}{2\alpha-1}$. Now, (35)–(38) remain the same, and (39) becomes

$$g^+(y_1, h_{k+1}) - g^+(y_1, h_k) \leq C_1 h_k^{1-\varepsilon} \frac{1}{h_k} + C_2 h_k^{-\beta\alpha} h_k.$$

By the choice of β , all the exponents of h_k are negative. We do not need to change anything from (40) to (49). Finally from (50) we need to replace the computation in the case $\alpha \leq 1$ with the following computation.

First, we have $(g^+)'' \leq -\frac{1}{4}(g^+)'\frac{1}{\alpha} \leq -\frac{1}{4}(g^+)'$, and we only need to bound

$$\int_{[h_k+h_k^{1-\varepsilon/2}, \inf I_2]} (g^+)'\, dh.$$

Note that $(g^+)'\geq 0$. By integrating the above differential inequality, we have

$$(g^+)'(h) \leq (g^+)'(h_k) e^{-\frac{1}{4}|I_3|} \leq (g^+)'(h_k) e^{\frac{1}{8}(h-h_k)}$$

when $h \in [h_k + h_k^{1-\varepsilon/2}, \inf I_2]$. Therefore, we have

$$\begin{aligned} \int_{[h_k+h_k^{1-\varepsilon/2}, \inf I_2]} (g^+)'\, dh &\leq \int_{h_k}^{h_{k+1}} (g^+)'(h_k) e^{\frac{1}{8}(h-h_k)}\, dh \\ &\leq C(g^+)'(h_k) \leq \frac{C}{h_k}. \end{aligned} \quad \square$$

3. Blow-down of an entire convex ancient solutions converges to a power function

In this section we prove that the blow-down of an entire convex solution to (3) converges to a power function.

Proof of Theorem 1 and the first part of Theorem 2. First, we prove that there is a subsequence of u_h converging to $\frac{1}{1+\alpha}|x|^{1+\alpha}$, where $u_h(x) = h^{-1}u(h^{\frac{1}{1+\alpha}}x)$.

By adding a constant we may suppose $u(0) = 0$. Let $x_{n+1} = a \cdot x$ be the equation of the tangent plane of u at 0. By [Corollary 10](#) and the convexity of u we have

$$a \cdot x \leq u(x) \leq C(1 + |x|^{1+\alpha}).$$

Hence,

$$h^{-\frac{\alpha}{1+\alpha}} a \cdot x \leq u_h(x) \leq C\left(\frac{1}{h} + |x|^{1+\alpha}\right).$$

By convexity, Du_h is locally uniformly bounded. Hence, u_h subconverges to a convex function u_0 which satisfies $u_0(0) = 0$, and

$$0 \leq u_0(x) \leq C|x|^{1+\alpha}.$$

It is easy to check that u_0 is an entire convex viscosity solution to [\(3\)](#) with $\sigma = 0$, and the comparison principle holds on any bounded domain.

Now we will prove that $\{u_0(x) = 0\} = \{0\}$. In fact, if $\{x : u_0(x) = 0\}$ is a bounded set, then $\{u_0(x) = h\}$ is a closed, bounded convex curve which evolves under the generalized curve shortening flow; from [\[Andrews 1998\]](#) it follows that $\{u_0(x) = 0\} = \{0\}$. If $\{u_0(x) = 0\}$ contains a straight line, say the line $\{(t, 0) : t \in \mathbb{R}\}$, then by convexity, u is independent of x_1 , which is impossible. So we only need to rule out the possibility that $\{u_0(x) = 0\}$ contains a ray but no straight lines. In this case, for fixed $h > 0$, we can find an ellipse E inside $\{u_0(x) < h\}$, with the short axis bounded from below by a constant depending only on h and with the long axis as long as we want (one only needs to look at the asymptotic cone of $\{u_0(x) = h\}$), but since $\{u_0(x) = h\}$ evolves under the generalized curve shortening flow and $E \subset \{u_0(x) \leq h\}$, this is impossible by comparison principle.

Then since $\{u_0(x) = 0\} = \{0\}$, $\Gamma_{1, u_0} = \{u_0(x) = 1\}$ is a bounded convex curve, and the level set $\{u_0(x) = -t\}$ evolves under the generalized curve shortening flow, with time $t \in (-\infty, 0)$. From [\[Andrews 1998; 2003\]](#) we have the following asymptotic behavior of the convex solution u_0 of $L_0 u = 1$:

$$(56) \quad u_0(x) = \frac{1}{1+\alpha}|x|^{1+\alpha} + \varphi(x),$$

where $\varphi(x) = o(|x|^{\alpha+1})$ for $x \neq 0$ near the origin. In fact, if the initial level curve is in a sufficiently small neighborhood of circle, by [Lemma 13](#), $|\varphi(x)| \leq C|x|^{1+\alpha+\eta}$ for some small positive η , where C is a constant depending only on the initial closeness to the circle. Hence, given any $\epsilon > 0$, for sufficiently small $h' > 0$,

$$B_{(1-\epsilon)r}(0) \subset \Omega_{h', u_0} \subset B_{(1+\epsilon)r}(0),$$

where $r = ((1 + \alpha)h')^{\frac{1}{1+\alpha}}$. Hence, there is a sequence $h_m \rightarrow \infty$ such that

$$B_{(1-\frac{1}{m})r_{m,i}}(0) \subset \Omega_{h_m, u} \subset B_{(1+\frac{1}{m})r_{m,i}}(0),$$

where

$$r_{m,i} = ((1 + \alpha)ih_m)^{\frac{1}{1+\alpha}}, \quad i = 1, \dots, m.$$

Then u_{h_m} subconverges to $\frac{1}{1+\alpha}|x|^{1+\alpha}$.

Since u_0 is an entire convex solution to $L_0u = 1$ (we still use the notation u_0 , but it means an arbitrary entire convex solution), from the above argument, we can find a sequence h_m such that

$$u_{0h_m}(x) = \frac{1}{h_m}u_0(h_m^{\frac{1}{1+\alpha}}x)$$

locally uniformly converges to $\frac{1}{1+\alpha}|x|^{1+\alpha}$. Hence, the sublevel set $\Omega_{\frac{1}{1+\alpha}, u_{0h_m}}$ satisfies

$$B_{1-\epsilon_m}(0) \subset \Omega_{\frac{1}{1+\alpha}, u_{0h_m}} \subset B_{1+\epsilon_m}(0),$$

where $\epsilon_m \rightarrow 0$ as $m \rightarrow \infty$. By the discussion below (56),

$$u_{0h_m}(x) = \frac{1}{1+\alpha}|x|^{1+\alpha} + \varphi(x),$$

where $|\varphi(x)| \leq C|x|^{1+\alpha+\eta}$ for some fixed small positive η , and the constant C is independent of m . Replacing x by $h_m^{-1/(1+\alpha)}x$ in the asymptotic formula above,

$$u_0(x) = \frac{1}{1+\alpha}|x|^{1+\alpha} + h_m\varphi(h_m^{-\frac{1}{1+\alpha}}x),$$

where for any fixed x , $h_m\varphi(h_m^{-\frac{1}{1+\alpha}}x) \rightarrow 0$. Hence $u_0(x) = \frac{1}{1+\alpha}|x|^{1+\alpha}$. So we have proved [Theorem 1](#) and the first part of [Theorem 2](#). \square

4. One-dimensional entire convex ancient solution must be a shrinking circle

This section is devoted to the proof of [Corollary 3](#), which is completed by combining the following lemma (corresponding to [[Wang 2011](#), Lemma 4.1]) and [Theorem 2](#).

Lemma 11. *Let Ω be a smooth, bounded, convex domain in \mathbb{R}^2 . Let u be the solution of (3) with $\sigma = 0$, vanishing on $\partial\Omega$. Then for any constant h satisfying $\inf_{\Omega} u < h < 0$, the level set $\Gamma_{h,u} = \{u = h\}$ is convex. Moreover, $\log(-u)$ is a concave function.*

Proof. Observe that $\varphi := -\log(-u)$ satisfies

$$|D\varphi|^{\frac{1}{\alpha}-1} \sum_{i,j=1}^2 \left(\delta_{ij} - \frac{\varphi_i\varphi_j}{|D\varphi|^2} \right) \varphi_{ij} = e^{\frac{1}{\alpha}\varphi}.$$

Since $\varphi(x) \rightarrow +\infty$ as $x \rightarrow \partial\Omega$, [[Kawohl 1985](#), Theorem 3.13] implies that φ is convex. \square

With the previous lemma and [[Wang 2011](#), Lemma 4.4], we know that any convex compact ancient solution to the generalized curve shortening flow can be represented as a convex solution u to (3) with $\sigma = 0$, and if the solution to the flow

sweeps the whole space, then the corresponding u will be an entire solution. Thus, [Theorem 2](#) implies [Corollary 3](#) immediately.

Remark 12. We can also use the method in [[Wang 2011](#), Section 4] to construct a rotationally nonsymmetric convex compact ancient solution for generalized curve shortening flow with power $\alpha \in (\frac{1}{2}, 1)$. Indeed, for mean curvature flow, a rotationally nonsymmetric convex compact ancient solution is constructed in Lemmas 4.1–4.4 of that reference. By examining the proofs of these lemmas, we can see that they work well for the generalized curve shortening flow considered here.

5. Two-dimensional entire convex translating solution

In this section, by using the previous results and a delicate iteration argument, we prove that under some extra condition on the asymptotic behavior of the solution at infinity the translating solution must be rotationally symmetric.

First of all, we would like to point out that instead of using Gage and Hamilton's exponential convergence of the curve shortening flow [[1986](#)], we need to use the corresponding exponential convergence for the generalized curve shortening flow and we will state it as a lemma, corresponding to [[Wang 2011](#), Lemma 3.2].

Lemma 13. *Let $\{\ell_t\}$ be a convex solution to the generalized curve shortening flow with initial curve $\{\ell_0\}$ uniformly convex. Suppose $\{\ell_0\}$ is in the δ_0 -neighborhood of a unit circle, $\{\ell_t\}$ shrinks to the origin at $t = \frac{1}{1+\alpha}$. Let*

$$\tilde{\ell}_t = (1 - (1 + \alpha)t)^{-\frac{1}{1+\alpha}} \ell_t$$

be the normalization of ℓ_t . Then $\tilde{\ell}_t$ is in the δ_t -neighborhood of the unit circle centered at the origin,

$$\tilde{\ell}_t \subset N_{\delta_t} S^1,$$

with

$$\delta_t \leq C \delta_0 \left(\frac{1}{1+\alpha} - t \right)^t$$

for some small positive constant ι .

Remark 14. Exponential convergence of the standard curve shortening flow (when $\alpha = 1$) was proved by Gage and Hamilton [[1986](#)]. For the general case (when $\alpha > \frac{1}{3}$), as discussed in the following proof, Gage and Hamilton's method combined with Andrews' estimates [[1998](#), Propositions II.1.1 and II.1.2] can still be used to prove the corresponding exponential convergence result.

Proof. The proof of [Lemma 13](#) is similar to the proof of [[Wang 2011](#), Lemma 3.2]. Since the initial curve ℓ_0 is uniformly convex and close to a unit circle, by [[Andrews 1998](#), Propositions II.1.1 and II.1.2], the curvature of $\tilde{\ell}_t$ is bounded from below and from above by some constant depending only on δ_0 , when $t \in (\frac{1}{4\alpha+4}, \frac{1}{2\alpha+2})$.

Hence the evolution equation for $\tilde{\ell}_t$ is uniformly parabolic. Therefore, we can apply Schauder's estimates safely for $\alpha > \frac{1}{2}$, as in [Wang 2011], which says that

$$\|\tilde{\ell}_t - S^1\|_{C^k} \leq C\delta_0 \quad \text{for } t \in \left(\frac{1}{4\alpha+4}, \frac{1}{2\alpha+2}\right).$$

Although the constant C will depend on the lower and upper bounds of the curvature of the initial curve, it is not a problem for our purpose, since when we blow down the solution for $\sigma = 0$, the norm of the gradient Du_h on the curve $\{u_h(x) = 1\}$ approaches 1.

By the equation $\kappa u_h^{\frac{1}{\alpha}} = 1$, we see that the curvature κ is also very close to 1 on that curve. For the exponential decay rate of the derivative of curvature, one can imitate the proof in [Gage and Hamilton 1986, §§5.7.10–5.7.15], and our corresponding estimate will be $|\kappa'(\tau)| \leq C\delta_0 e^{-\iota\tau}$ for some small positive number ι , where $\tau = -\frac{1}{1+\alpha} \log\left(\frac{1}{1+\alpha} - t\right)$. Indeed, in the case $\alpha > 1$, this is done by Chen and Huang [Huang 2011, Corollary 3.2], and it is easy to check that their computation also works for the case $\frac{1}{3} < \alpha < 1$ by taking ℓ small enough. This estimate immediately implies our lemma. \square

In the following we will consider the case when $\sigma = 1$ and $\alpha > 1$. By translating and adding some constant we can assume $u(0) = \inf u$. Let

$$u_h(x) = \frac{1}{h} u(h^{\frac{1}{1+\alpha}} x).$$

Then u_h satisfies the equation $L_\sigma u_h = 1$ with $\sigma = h^{-\frac{2\alpha}{1+\alpha}}$. By Theorem 1, u_h converges to $\frac{1}{1+\alpha} |x|^{1+\alpha}$, and the level set $\Gamma_{\frac{1}{1+\alpha}, u_h}$ converges to the unit circle as $h \rightarrow \infty$.

Lemma 15 [Wang 2011, Lemma 3.3]. *The function u satisfies*

$$(57) \quad u(x) = \frac{1}{1+\alpha} |x|^{1+\alpha} + O(|x|^{1+\alpha-2\alpha\beta}),$$

where β is a constant, chosen such that $\frac{1}{2\alpha} < \beta < \min\{1, \frac{1+\alpha}{2\alpha}\}$.

Proof. For any given small $\delta_0 > 0$, take h sufficiently large such that

$$(58) \quad \Gamma_{\frac{1}{1+\alpha}, u_h} \subset N_{\delta_0}(S^1)$$

for the unit circle S^1 with center p_0 . Note that when h is large, δ_0 is very close to 0. Then we will prove the following claim:

Claim 3: For small fixed τ ,

$$(59) \quad \Gamma_{\tau, u_h} \subset ((1+\alpha)\tau)^{\frac{1}{1+\alpha}} N_{\delta_\tau} \left(\left(1 + \frac{\alpha_0}{\tau}\right)^{\frac{1}{1+\alpha}} S^1 \right)$$

with

$$(60) \quad \delta_\tau \leq C_1(\tau)\sigma^\beta + C_2\delta_0\tau^\eta,$$

where η is a small positive constant, the constants C_1 and C_2 are independent of δ_0 and h , and C_2 is also independent of τ . Let u_0 be the solution of

$$L_0(u) = 1 \quad \text{in } \Omega_{\frac{1}{1+\alpha}, u_h},$$

satisfying

$$u_0 = u_h = \frac{1}{1+\alpha} \quad \text{on } \partial\Omega_{\frac{1}{1+\alpha}, u_h},$$

where $a_0 = |\inf u_0|$ and the center of

$$(1 + \frac{a_0}{\tau})^{\frac{1}{1+\alpha}} S^1$$

is the minimum point of u_0 multiplied by the factor $((1 + \alpha)\tau)^{-\frac{1}{1+\alpha}}$.

Proof of Claim 3. We only need to prove that

$$(61) \quad \text{dist}((1 + \alpha)^{\frac{1}{1+\alpha}} (\tau + a_0)^{\frac{1}{1+\alpha}} S^1, \Gamma_{\tau, u}) \leq C_1(\tau)\sigma^\beta + C_2\delta_0\tau^{\frac{1}{1+\alpha} + \eta},$$

where η is some small positive constant and C_2 is independent of τ . By [Theorem 1](#) we know that u_h converges to $\frac{1}{1+\alpha}|x|^{1+\alpha}$ uniformly on any compact subset of \mathbb{R}^2 . Then by the convexity of u_h , we have that $|Du_h|$ is bounded above and below by some constants depending on τ_0 for large h when

$$x \in \left\{x \in \Omega_{\frac{1}{1+\alpha}, u_h} : \tau_0 \leq u_h < \frac{1}{1+\alpha}\right\}.$$

Hence, by the growth condition for D^2u in [Theorem 2](#), we have $\sigma(u_h)_{\gamma\gamma} \leq C\sigma^\beta$, where C is a constant depending on τ_0 . Therefore, we have

$$\kappa(u_h)_{\gamma}^{1/\alpha} \approx 1 - C\sigma^\beta \quad \text{on } \left\{x \in \Omega_{\frac{1}{1+\alpha}, u_h} : \tau \leq u_h < \frac{1}{1+\alpha}\right\},$$

where C depends on τ_0 . Let

$$\tilde{u}_0 = (1 - C\sigma^\beta)^\alpha (u_0 - \frac{1}{1+\alpha}) + \frac{1}{1+\alpha};$$

then

$$L_0(\tilde{u}_0) = 1 - C\sigma^\beta \quad \text{in } \Omega_{\frac{1}{1+\alpha}, u_h}$$

with

$$\tilde{u}_0 = u_h = \frac{1}{1+\alpha} \quad \text{on } \partial\Omega_{\frac{1}{1+\alpha}, u_h}.$$

Now by the comparison principle, $\Omega_{\tau, u_0} \supset \Omega_{\tau, u_h} \supset \Omega_{\tau, \tilde{u}_0}$, and by the asymptotic behavior of u_0 ,

$$\Gamma_{\tau, u_0} \subset N_\zeta((\tau + a_0)^{\frac{1}{1+\alpha}} S^1) \quad \text{and} \quad \Gamma_{\tau, \tilde{u}_0} \subset N_\zeta((\tau + a_0 - C\sigma^\beta)^{\frac{1}{1+\alpha}} S^1),$$

where $\zeta = C\delta_0(\tau + a_0)^\eta$. Let

$$\ell_1 = (\tau + a_0)^{\frac{1}{1+\alpha}} S^1 \quad \text{and} \quad \ell_2 = (\tau + a_0 - C\sigma^\beta)^{\frac{1}{1+\alpha}} S^1,$$

both centered at p_1 , which is the minimum point of u_0 . Hence

$$(62) \quad \text{dist}((\tau + a_0)^{\frac{1}{1+\alpha}} S^1, \Gamma_{\tau, u_h}) \leq \text{dist}(\ell_1, \ell_2) + C\delta_0(\tau + a_0)^{\frac{1}{1+\alpha} + \eta},$$

where $\text{dist}(\ell_1, \ell_2)$ can be bounded by $C_1(\tau)\sigma^\beta$; hence (60) follows from the above discussion. \square

Now, we will use an iteration argument to prove the following claim, which will enable us to simplify (59) and (60).

Claim 4: $a_0 \leq C\sigma^\beta |\log(\sigma)|$

Proof. We fix a large constant A such that $\{u_{A/\tau} = \frac{1}{1+\alpha}\}$ is very close to a unit circle. Let u_{0, τ^k} solve $L_0 u = 1$ with boundary condition $u = \tau^k$ on $\{u_h = \tau^k\}$. Denote $a_k = |\inf u_{0, \tau^k}|$. From the proof of Claim 3, we see that

$$\{u_0 < \tau\} \supset \{u_{0, \tau} < \tau\} \supset \{\tilde{u}_0 < \tau\},$$

by the comparison principle, we have $\inf u_0 < \inf u_{0, \tau} < \inf \tilde{u}_0$. So by the construction of \tilde{u}_0 and a simple computation, we have $a_0 - a_1 \leq \inf \tilde{u}_0 - \inf u_0 \leq C\sigma^\beta$. When $\tau^k \geq \frac{A}{h}$, we can iterate this argument for u_{0, τ^k} and $u_{0, \tau^{k+1}}$ by rescaling them to

$$\frac{1}{1+\alpha} \tau^{-k} u_{0, \tau^k} \left((1+\alpha)^{\frac{1}{1+\alpha}} \tau^{\frac{k}{1+\alpha}} x \right) \quad \text{and} \quad \frac{1}{1+\alpha} \tau^{-k} u_{0, \tau^{k+1}} \left((1+\alpha)^{\frac{1}{1+\alpha}} \tau^{\frac{k}{1+\alpha}} x \right),$$

respectively. After rescaling back, we have $a_k - a_{k+1} \leq C\sigma^\beta$. Note that the choice of A and the condition $\tau^k \geq \frac{A}{h}$ ensure the uniform gradient bound needed in the above argument. Let k_0 be an integer satisfying $\tau^{k_0} \geq \frac{A}{h} \geq \tau^{k_0+1}$. After k_0 steps we stop the iteration, and notice that

$$\{u_h = \frac{A}{h}\} = h^{-\frac{1}{1+\alpha}} \{u = A\}$$

is contained in a circle with radius $Ch^{-\frac{1}{1+\alpha}}$ for some constant C . Hence it takes at most time $Ch^{-1} = C\sigma^{\frac{1+\alpha}{2\alpha}}$ for $\{u_h = \frac{A}{h}\}$ to shrink to a point. Claim 4 follows from the above discussion. \square

By omitting the lower order term we can rewrite (59) and (60) as

$$\Gamma_{\tau, u_h} \subset ((1+\alpha)\tau)^{\frac{1}{1+\alpha}} N_{\delta_\tau}(S^1)$$

with

$$(63) \quad \delta_\tau \leq C_1(\tau)\sigma^\beta + C_2\delta_0\tau^\eta.$$

If we take τ small such that $C_2\tau^\eta \leq \frac{1}{4}$, then (63) becomes

$$(64) \quad \delta_\tau \leq C_1(\tau)\sigma^\beta + \frac{1}{4}\delta_0.$$

Now we can carry out an iteration argument similar to that in [Wang 2011]. We start at the level $\frac{1}{1+\alpha}\tau^{-k_0}$ for some sufficient large k_0 . Let

$$\Omega_k = \tau^{\frac{k}{1+\alpha}} \Omega_{\frac{1}{1+\alpha}\tau^{-k}, u} \quad \text{and} \quad \Gamma_k = \partial\Omega_k.$$

Note that Γ_k converges to a unit circle as $k \rightarrow \infty$. Suppose that Γ_k is in the δ_k neighborhood of S^1 centered at y_k , where $\delta_k \rightarrow 0$ as $k \rightarrow \infty$ and y_k is the minimum point of the solution of $L_0u = 1$ in Ω_k with $u = \frac{1}{1+\alpha}$ on Γ_{k+1} . By (64) we have

$$(65) \quad \delta_{k-1} \leq C_1(\tau) \tau^{(k-1)\frac{2\alpha\beta}{1+\alpha}} + \frac{1}{4}\delta_k$$

for $k = k_0, k_0 + 1, \dots$. Then we have

$$(66) \quad \Gamma_j \subset N_{\delta_j}(S^1)$$

with

$$(67) \quad \delta_j \leq C\tau^{j\frac{2\alpha\beta}{1+\alpha}}$$

It follows that

$$(68) \quad \Gamma_{\frac{1}{1+\alpha}\tau^{-j}, u} \subset N_{\tilde{\delta}_j}(\tau^{\frac{-j}{1+\alpha}}S^1)$$

with

$$(69) \quad \tilde{\delta}_j \leq C\tau^{\frac{2\alpha\beta-1}{1+\alpha}j},$$

where $\tau^{\frac{-j}{1+\alpha}}S^1$ is centered at $z_j = \tau^{\frac{-j}{1+\alpha}}y_j$. From Lemma 13 and (64), it is not hard to see that

$$(70) \quad |z_j - z_{j-1}| \leq C\tau^{\frac{2\alpha\beta-1}{1+\alpha}j}.$$

Let $z_0 = \lim_{j \rightarrow \infty} z_j$. Then

$$(71) \quad |z_j - z_0| \leq C\tau^{\frac{2\alpha\beta-1}{1+\alpha}j},$$

which means that in (68) we can assume the circle is centered at z_0 by changing the constant C a little bit. In fact when we choose different τ , the corresponding z_0 will not change, so we can assume $z_0 = 0$. Hence, for $h = \frac{1}{1+\alpha}\tau^{-j}$,

$$\Gamma_{h,u} \subset N_\delta((1+\alpha)^{\frac{1}{1+\alpha}}h^{\frac{1}{1+\alpha}}S^1),$$

where

$$(72) \quad \delta \leq Ch^{\frac{1-2\alpha\beta}{1+\alpha}}$$

and S^1 is centered at the origin. By choosing different τ , we see that the estimate holds for all large h . Lemma 15 follows from the above estimates. \square

Remark 16. For the mean curvature flow (when $\alpha = 1$), I learned the proof of [Claim 4](#) from Professor Xu-Jia Wang. Indeed, in this case one can do it as follows. Let v_k solve $L_0 w = 1$ with boundary condition $w = \tau^k$ on $\{u_h = \tau^k\}$, for $k = 1, 2, \dots$. Let $a_k = |\inf v_k|$. Then by [[Wang 2011](#), Lemma 3.1], we have $a_0 - a_1 \leq C\sigma$. By rescaling invariance, we can iterate the argument to show that $a_k - a_{k+1} \leq C\sigma$, provided $\tau^k \geq \frac{1}{h}$. Hence, we stop the iteration at k_0 when $\tau^{k_0} \geq \frac{1}{h} > \tau^{k_0+1}$. Notice that

$$\{u_h \leq \tau^{k_0}\} = h^{-\frac{1}{2}}\{u = h\tau^{k_0}\} \subset h^{-\frac{1}{2}}\{u \leq \frac{1}{h}\}.$$

So, it is easy to see that it takes at most time $C\sigma = \frac{C}{h}$ for $\{u_h \leq \tau^{k_0}\}$ to shrink to a point, namely, $a_{k_0} \leq C\sigma$. Therefore,

$$a_0 = a_{k_0} + \sum_{i=0}^{k_0-1} a_i - a_{i+1} \leq Ck_0\sigma \leq C\sigma |\log \sigma|.$$

In order to finish the proof of [Theorem 2](#) we need to use the following fundamental Liouville theorem by Bernstein [[Simon 1997](#), p. 245].

Lemma 17. *Let u be an entire solution to the elliptic equation*

$$\sum_{i,j=1}^n a_{ij}(x)u_{ij} = 0 \quad \text{in } \mathbb{R}^2.$$

If u satisfies the asymptotic estimate

$$|u(x)| = o(|x|) \quad \text{as } x \rightarrow \infty,$$

then u is a constant.

Proof of the second part of [Theorem 2](#). Let u^* be the Legendre transform of u . Then u^* satisfies equation

$$(73) \quad G(x, D^2 u^*) = \frac{\det D^2 u^*}{(\delta_{ij} - \frac{x_i x_j}{1+|x|^2}) F^{ij}(u^*)} = (1 + |x|^2)^{\frac{1}{2\alpha} - \frac{1}{2}},$$

where $F^{ij}(u^*) = \frac{\partial \det r}{\partial r_{ij}}$ at $r = D^2 u^*$. We have

$$(74) \quad u^*(x) = C(\alpha)|x|^{1+\alpha} + O(|x|^{\frac{1+\alpha-2\alpha\beta}{\alpha}}),$$

where $C(\alpha)$ is a constant depending only on α . In fact, for big h , by [Lemma 15](#),

$$u_h(x) = \frac{1}{1+\alpha}|x|^{1+\alpha} + O(|h|^{\frac{-2\alpha\beta}{1+\alpha}})$$

in $B_1(0)$. Denote by u_h^* the Legendre transform of u_h . Then,

$$u_h^*(x) = C(\alpha)|x|^{1+\frac{1}{\alpha}} + O(|h|^{\frac{-2\alpha\beta}{1+\alpha}}),$$

where $C(\alpha)$ is a constant depending only on α and it comes from the Legendre transform of the function $\frac{1}{1+\alpha}|x|^{1+\alpha}$. Note that $u_h^*(x) = h^{-1}u^*(h^{\frac{\alpha}{1+\alpha}}x)$, we obtain (74).

Let u_0 be the unique radial solution of (3) with $\sigma = 1$, and let u_0^* be the Legendre transform of u_0 . Similar to (74) we have

$$(75) \quad u_0^*(x) = C(\alpha)|x|^{1+\alpha} + O(|x|^{\frac{1+\alpha-2\alpha\beta}{\alpha}}).$$

Since both u^* and u_0^* satisfy (73), $v = u^* - u_0^*$ satisfies the elliptic equation

$$\sum_{i,j=1}^n a_{ij}(x)v_{ij} = 0 \quad \text{in } \mathbb{R}^2,$$

where

$$a_{ij} = \int_0^1 G^{ij}(x, D^2u_0^* + t(D^2u^* - D^2u_0^*)) dt.$$

Here,

$$G^{ij} = \frac{\partial G(x, r)}{\partial r_{ij}}$$

for any symmetric matrix r . Note that by the choice of β , $\frac{1+\alpha-2\alpha\beta}{\alpha} < 1$; hence, by (74) and (75),

$$v = O(|x|^{\frac{1+\alpha-2\alpha\beta}{\alpha}}) = o(|x|) \quad \text{as } |x| \rightarrow \infty.$$

By Lemma 17 we conclude that v is a constant. □

References

- [Andrews 1998] B. Andrews, “Evolving convex curves”, *Calc. Var. PDE* **7**:4 (1998), 315–371. [MR 99k:58038](#) [Zbl 0931.53030](#)
- [Andrews 2002] B. Andrews, “Non-convergence and instability in the asymptotic behaviour of curves evolving by curvature”, *Comm. Anal. Geom.* **10**:2 (2002), 409–449. [MR 2003e:53086](#) [Zbl 1029.53079](#)
- [Andrews 2003] B. Andrews, “Classification of limiting shapes for isotropic curve flows”, *J. Amer. Math. Soc.* **16**:2 (2003), 443–459. [MR 2004a:53083](#) [Zbl 1023.53051](#)
- [Chou and Zhu 2001] K.-S. Chou and X.-P. Zhu, *The curve shortening problem*, Chapman & Hall/CRC, Boca Raton, FL, 2001. [MR 2003e:53088](#) [Zbl 1061.53045](#)
- [Daskalopoulos et al. 2010] P. Daskalopoulos, R. Hamilton, and N. Sesum, “Classification of compact ancient solutions to the curve shortening flow”, *J. Differential Geom.* **84**:3 (2010), 455–464. [MR 2012d:53213](#) [Zbl 1205.53070](#)
- [Gage and Hamilton 1986] M. Gage and R. S. Hamilton, “The heat equation shrinking convex plane curves”, *J. Differential Geom.* **23**:1 (1986), 69–96. [MR 87m:53003](#) [Zbl 0621.53001](#)
- [Huang 2011] R. L. Huang, “Blow-up rates for the general curve shortening flow”, *J. Math. Anal. Appl.* **383**:2 (2011), 482–489. [MR 2012h:53152](#) [Zbl 1220.53082](#)

- [Kawohl 1985] B. Kawohl, *Rearrangements and convexity of level sets in PDE*, Lecture Notes in Mathematics **1150**, Springer, Berlin, 1985. MR 87a:35001 Zbl 0593.35002
- [Schulze 2005] F. Schulze, “Evolution of convex hypersurfaces by powers of the mean curvature”, *Math. Z.* **251**:4 (2005), 721–733. MR 2006h:53065 Zbl 1087.53062
- [Sheng and Wu 2009] W. Sheng and C. Wu, “On asymptotic behavior for singularities of the powers of mean curvature flow”, *Chin. Ann. Math. (B)* **30**:1 (2009), 51–66. MR 2010h:53100 Zbl 1180.53066
- [Simon 1997] L. Simon, “The minimal surface equation”, pp. 239–266 in *Geometry V*, edited by R. Osserman, Encyclopaedia of Mathematical Sciences **90**, Springer, Berlin, 1997. MR 99b:53014 Zbl 0905.53003
- [Wang 2011] X.-J. Wang, “Convex solutions to the mean curvature flow”, *Ann. of Math. (2)* **173**:3 (2011), 1185–1239. MR 2800714 Zbl 1231.53058
- [White 2000] B. White, “The size of the singular set in mean curvature flow of mean-convex sets”, *J. Amer. Math. Soc.* **13**:3 (2000), 665–695. MR 2001j:53098 Zbl 0961.53039

Received February 19, 2013. Revised October 19, 2014.

SHIBING CHEN
DEPARTMENT OF MATHEMATICS
ZHEJIANG UNIVERSITY OF TECHNOLOGY
HANGZHOU, ON 310023
CHINA

Current address:

MATHEMATICAL SCIENCES INSTITUTE
THE AUSTRALIAN NATIONAL UNIVERSITY
JOHN DEDMAN BUILDING 27
CANBERRA ACT 2601
AUSTRALIA

chenshibing1982@hotmail.com

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 276 No. 1 July 2015

On the degree of certain local L -functions	1
U. K. ANANDAVARDHANAN and AMIYA KUMAR MONDAL	
Torus actions and tensor products of intersection cohomology	19
ASILATA BAPAT	
Cyclicity in Dirichlet-type spaces and extremal polynomials II: functions on the bidisk	35
CATHERINE BÉNÉTEAU, ALBERTO A. CONDORI, CONSTANZE LIAW, DANIEL SECO and ALAN A. SOLA	
Compactness results for sequences of approximate biharmonic maps	59
CHRISTINE BREINER and TOBIAS LAMM	
Criteria for vanishing of Tor over complete intersections	93
OLGUR CELIKBAS, SRIKANTH B. IYENGAR, GREG PIEPMEYER and ROGER WIEGAND	
Convex solutions to the power-of-mean curvature flow	117
SHIBING CHEN	
Constructions of periodic minimal surfaces and minimal annuli in Sol_3	143
CHRISTOPHE DESMONTS	
Quasi-exceptional domains	167
ALEXANDRE EREMENKO and ERIK LUNDBERG	
Endoscopic transfer for unitary groups and holomorphy of Asai L -functions	185
NEVEN GRBAC and FREYDOON SHAHIDI	
Quasiconformal harmonic mappings between Dini-smooth Jordan domains	213
DAVID KALAJ	
Semisimple super Tannakian categories with a small tensor generator	229
THOMAS KRÄMER and RAINER WEISSAUER	
On maximal Lindenstrauss spaces	249
PETR PETRÁČEK and JIŘÍ SPURNÝ	