

*Pacific
Journal of
Mathematics*

**FROM QUASIMODES TO RESONANCES:
EXPONENTIALLY DECAYING PERTURBATIONS**

ORAN GANNOT

Volume 277 No. 1

September 2015

FROM QUASIMODES TO RESONANCES: EXPONENTIALLY DECAYING PERTURBATIONS

ORAN GANNOT

We consider self-adjoint operators of black-box type which are exponentially close to the free Laplacian near infinity, and prove an exponential bound for the resolvent in a strip away from resonances. Here the resonances are defined as poles of the meromorphic continuation of the resolvent between appropriate exponentially weighted spaces. We then use a local version of the maximum principle to prove that any cluster of real quasimodes generates at least as many resonances, with multiplicity, rapidly converging to the quasimodes.

1. Introduction

It is expected that for open systems, trapping of classical trajectories produces scattering resonances close to the real axis; this is often referred to as the Lax–Phillips conjecture [1989, Section V.3]. When trapping is weak, for instance in the sense of hyperbolicity, the general conjecture is not true, as shown by Ikawa [1982]. For an account of recent results about resonances near the real axis under weak trapping; see the review by Wunsch [2012]. On the other hand, when the trapping is sufficiently strong so that a construction of real quasimodes is possible, there exist resonances close to the quasimodes [Stefanov and Vodev 1996; Tang and Zworski 1998; Stefanov 1999]. These results were established in the setting of compactly supported perturbations, or more generally for perturbations which are dilation analytic near infinity [Sjöstrand and Zworski 1991; Sjöstrand 1997].

Complementary to the aforementioned results, in this note we prove analogues for “black box” operators which are exponentially close to the free Laplacian at infinity. More precisely, we allow both metric and potential perturbations of the Laplacian outside a compact set (the black box), but require only minimal assumptions on the operator in the black box. Standard techniques give a meromorphic continuation of the exponentially weighted resolvent through the real axis to a strip whose width is of size $O(h)$; the choice of exponential weight and the width of the strip depend on the decay rate of the perturbation. We then apply a complex analytic

MSC2010: primary 35P25; secondary 47F05, 47A40.

Keywords: scattering resonances, quasimodes, exponentially decaying potentials.

framework — summarized, for example, in [Petkov and Zworski 2001] — to deduce an exponential a priori bound on the weighted resolvent away from resonances.

A typical application of such an exponential bound — well-established in [Stefanov and Vodev 1996; Tang and Zworski 1998; Stefanov 1999; 2005] — is to show that any family of sufficiently independent quasimodes generates at least as many resonances, counting multiplicity; these resonances converge rapidly not only to the real axis, but to the quasimodes; see Theorem B for a precise statement. The general assumptions are presented beginning in Section 1B.

One motivation for this work comes from a recent investigation of resonances for Schwarzschild–AdS black holes, where quasimodes have been constructed [Gannot 2014; Holzegel and Smulevici 2014]. Due to the spherical symmetry in that setting, the stationary wave operator P decomposes as a sum of one-dimensional operators P_ℓ on a half-line, which are just restrictions to spaces of spherical harmonics with angular momentum ℓ . Each P_ℓ is a self-adjoint perturbation of the Laplacian by an exponentially decaying potential which is singular near the origin — the results of this paper imply that the resolvent $R_\ell(\sigma)$ of P_ℓ has a meromorphic continuation through the real axis. Although meromorphy of each one-dimensional resolvent does not imply meromorphy for the global resolvent (this requires uniform control as $\ell \rightarrow \infty$ and was recently established in the Schwarzschild–AdS setting; see [Warnick 2015]), the results of this paper do imply the existence of a sequence of poles σ_ℓ for $R_\ell(\sigma)$ satisfying

$$0 < -\operatorname{Im} \sigma_\ell < C e^{-\ell/C} \quad \text{for } \ell \text{ sufficiently large.}$$

We also remark that in the Schwarzschild–AdS case the effective potential is dilation analytic, so the results of [Sjöstrand 1997] indeed apply. One advantage to the approach taken here is that the exponential decay of the potential remains stable under small (radial, static) perturbations of the Schwarzschild–AdS metric.

1A. Free resolvent. We begin by gathering several results about the free resolvent. The Laplacian $-\Delta$ on \mathbb{R}^n with domain $H^2(\mathbb{R}^n)$ is self-adjoint and we denote by $R_0(\sigma)$ the free resolvent

$$R_0(\sigma) = (-\Delta - \sigma^2)^{-1} : L^2(\mathbb{R}^n) \rightarrow H^2(\mathbb{R}^n), \quad \operatorname{Im} \sigma > 0.$$

Choose $\varphi \in C^\infty(\mathbb{R}^n)$ with the property that $\varphi(x) = |x|$ for $|x|$ large enough. If \mathcal{A} denotes some function space, we will use the notation $\mathcal{A}_\gamma = e^{-\gamma\varphi}\mathcal{A}$ for its weighted counterpart. We will also freely move between the equivalent notions

$$T : \mathcal{A}_\alpha \rightarrow \mathcal{B}_\beta \iff e^{\beta\varphi} T e^{-\alpha\varphi} : \mathcal{A} \rightarrow \mathcal{B},$$

depending on convenience.

Our starting point is the well known fact [McLeod 1967] that for each $\gamma > 0$ the weighted resolvent

$$e^{-\gamma\varphi} R_0(\sigma) e^{-\gamma\varphi} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$$

extends holomorphically across $\operatorname{Re} \sigma > 0$ as a bounded operator to the strip $\operatorname{Im} \sigma > -\gamma$, with the usual caveats in even dimensions when winding around the origin. We also have the standard representation,

$$(1-1) \quad e^{-\gamma\varphi} R_0(\sigma) e^{-\gamma\varphi} = e^{-\gamma\varphi} R_0(-\sigma) e^{-\gamma\varphi} + \sigma^{n-2} e^{-\gamma\varphi} M(\sigma) e^{-\gamma\varphi}$$

whenever $\operatorname{Re} \sigma > 0$ and $-\gamma < \operatorname{Im} \sigma < 0$. Here $M(\sigma)$ is the operator with kernel

$$M(\sigma, x, y) = (i/2)(2\pi)^{-n+1} \int_{\mathbb{S}^{n-1}} e^{i\sigma \langle \omega, x-y \rangle} d\omega.$$

We can also write

$$(1-2) \quad M(\sigma) = (i/2)(2\pi)^{-n+1} \Phi^t(\sigma) \Phi(-\sigma),$$

where $\Phi(\sigma) : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{S}^{n-1})$ has kernel $\Phi(\sigma, \omega, x) = e^{i\sigma \langle \omega, x \rangle}$ and $\Phi^t : L^2(\mathbb{S}^{n-1}) \rightarrow L^2(\mathbb{R}^n)$ has the transposed kernel.

The following two lemmas establish standard polynomial bounds for the free resolvent in the case of exponential weights.

Lemma 1.1. *For each $\epsilon > 0$ there exists a constant $C = C(\epsilon) > 0$ such that whenever $|\operatorname{Im} \sigma| < \gamma - \epsilon$ and $\operatorname{Re} \sigma \geq 1$,*

$$\|e^{-\gamma\varphi} M(\sigma) e^{-\gamma\varphi}\|_{L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)} < C |\sigma|^{1-n}.$$

Proof. The proof is adapted from [Burq 2002]. First note that the Fourier transform $\mathcal{F}(e^{-\gamma\varphi})(\xi)$ extends holomorphically to the strip $\{\xi \in \mathbb{C}^n : |\operatorname{Im} \xi| < \gamma - \epsilon\}$ and

$$(1-3) \quad |\mathcal{F}(e^{-\gamma\varphi})(\xi)| < C_N \langle \xi \rangle^{-N}$$

in the strip for each N . In light of (1-1) and (1-2), it suffices to prove that

$$\|\Phi(\sigma) e^{-\gamma\varphi}\|_{L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{S}^{n-1})} < C |\sigma|^{(1-n)/2},$$

which by Plancherel's theorem is equivalent to the same estimate for the composition $(\Phi(\sigma) e^{-\gamma\varphi}) \circ \mathcal{F}$. The operator $(\Phi(\sigma) e^{-\gamma\varphi}) \circ \mathcal{F}$ has kernel $\mathcal{F}(e^{-\gamma\varphi})(\sigma\omega - \xi)$. By Schur's lemma it suffices to obtain an estimate of the form

$$\sup_{\xi \in \mathbb{R}^n} \int_{\mathbb{S}^{n-1}} |\mathcal{F}(e^{-\gamma\varphi})(\sigma\omega - \xi)| d\omega < C |\sigma|^{1-n},$$

since in the other direction we may use (1-3) to obtain the trivial estimate

$$\sup_{\omega \in \mathbb{S}^{n-1}} \int_{\mathbb{R}^n} |\mathcal{F}(e^{-\gamma\varphi})(\sigma\omega - \xi)| d\xi < C.$$

Write ξ as $\xi = \langle \xi, \omega \rangle \omega + \xi^\perp(\omega)$ where $\langle \xi^\perp(\omega), \omega \rangle = 0$. Then, by (1-3), we are left estimating

$$\int_{\mathbb{S}^{n-1}} (1 + |\langle \xi, \omega \rangle - \operatorname{Re} \sigma| + |\xi^\perp(\omega)|)^{-N} d\omega.$$

Fix $\xi \in \mathbb{R}^n$ and $\delta > 0$, and decompose the sphere into two sets,

$$U = \{\omega \in \mathbb{S}^{n-1} : |\langle \xi, \omega \rangle - \operatorname{Re} \sigma| < \delta \operatorname{Re} \sigma, |\xi^\perp(\omega)| < \delta \operatorname{Re} \sigma\}$$

and its complement U^c . The integral over U^c is of the order $O(|\operatorname{Re} \sigma|^{-\infty})$, so it suffices to examine the integral over U .

Observe that unless $\operatorname{Re} \sigma$ is comparable to $|\xi|$, the set U is empty. Indeed, if $\omega \in U$ then $(1 - \delta) \operatorname{Re} \sigma < \langle \xi, \omega \rangle < (1 + \delta) \operatorname{Re} \sigma$. Hence,

$$\frac{\operatorname{Re} \sigma}{2} < (1 - \delta) \operatorname{Re} \sigma < \langle \xi, \omega \rangle \leq |\xi|,$$

while on the other hand,

$$|\xi|^2 = |\langle \xi, \omega \rangle|^2 + |\xi^\perp(\omega)|^2 < 3(\operatorname{Re} \sigma)^2$$

for δ sufficiently small.

Write a typical point of \mathbb{R}^n as (y, y') where $y \in \mathbb{R}^{n-1}$ and $y' \in \mathbb{R}$. By a rotation we may assume that $\xi = (0, |\xi|)$. In that case U is contained in the upper hemisphere, in a cap around $|\xi|^{-1}\xi = (0, 1)$ whose size is independent of ξ . This is true since $\omega \in U$ implies

$$\langle |\xi|^{-1}\xi, \omega \rangle > \frac{1}{2\sqrt{3}} > 0.$$

We then parametrize the upper hemisphere \mathbb{S}_+^{n-1} (which contains ξ) by the diffeomorphism

$$p : \mathbb{R}^{n-1} \rightarrow \mathbb{S}_+^{n-1}, \quad y \mapsto \frac{(y, |\xi|)}{|(y, |\xi|)|}.$$

Whenever $y \in p^{-1}(U)$ we have

$$|\xi^\perp(p(y))| \geq |y|.$$

To see this, compute

$$|\xi^\perp(p(y))|^2 = |\xi|^2 - |\langle \xi, p(y) \rangle|^2 = |\xi|^2 - \frac{|\xi|^4}{|y|^2 + |\xi|^2} = |y|^2 \frac{|\xi|^2}{|y|^2 + |\xi|^2} \geq |y|^2.$$

Furthermore, the Jacobian satisfies

$$|\partial p / \partial y| = O(|\xi|^{1-n}).$$

We can now bound the integral over U by

$$\begin{aligned} \int_U (1 + |\xi^\perp(\omega)|)^{-N} d\omega &= \int_{p^{-1}(U)} (1 + |\xi^\perp(p(y))|)^{-N} |\partial p / \partial y| dy \\ &\leq C_1 |\operatorname{Re} \sigma|^{1-n} \int_{\mathbb{R}^{n-1}} (1 + |y|)^{-N} dy \leq C_2 |\operatorname{Re} \sigma|^{1-n} \end{aligned}$$

for N large enough. In the second step, we used the fact that $|\xi|$ and $\operatorname{Re} \sigma$ were comparable. \square

Lemma 1.2. *For each $\epsilon > 0$ and $|\alpha| \leq 2$ there exists $C_\alpha = C_\alpha(\gamma, \epsilon)$ such that whenever $\operatorname{Im} \sigma > -\gamma + \epsilon$ and $\operatorname{Re} \sigma \geq 1$,*

$$\|D^\alpha (e^{-\gamma\varphi} R_0(\sigma) e^{-\gamma\varphi})\|_{L^2 \rightarrow L^2} \leq C_\alpha |\sigma|^{|\alpha|-1}.$$

Proof. (1) First we handle the case $|\alpha| = 0$ and $n > 1$; see [Rauch 1978; Vodev 1994] for similar arguments. Let $U(t) = \cos(t\sqrt{-\Delta})$ denote the propagator for the Cauchy problem

$$\begin{cases} (\partial_t^2 - \Delta)U(t)f(x) = 0, & (t, x) \in \mathbb{R} \times \mathbb{R}^n, \\ U(0)f(x) = f(x), & \partial_t U(0)f(x) = 0. \end{cases}$$

For $\operatorname{Im} \sigma > 0$, write the resolvent

$$(1-4) \quad e^{-\gamma\varphi} R_0(\sigma) e^{-\gamma\varphi} = \frac{i}{\sigma} \int_0^\infty e^{i\sigma t} e^{-\gamma\varphi} U(t) e^{-\gamma\varphi} dt.$$

Let r_0 be such that $\varphi(x) = |x|$ for $|x| \geq r_0$. Notice that $\|U(t)\|_{L^2 \rightarrow L^2} \leq 1$ and

$$\|1_{\{|x| \geq t/4\}} e^{-\gamma\varphi}\|_{L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)} \leq e^{-\gamma t/4}, \quad t \geq 4r_0.$$

Writing

$$\begin{aligned} U(t) &= 1_{\{|x| < t/4\}} U(t) 1_{\{|x| < t/4\}} + 1_{\{|x| \geq t/4\}} U(t) 1_{\{|x| < t/4\}} \\ &\quad + 1_{\{|x| < t/4\}} U(t) 1_{\{|x| \geq t/4\}} + 1_{\{|x| \geq t/4\}} U(t) 1_{\{|x| \geq t/4\}}, \end{aligned}$$

we see the norms of the latter three terms are of size $O(e^{-\gamma t/4})$ after multiplication by $e^{-\gamma\varphi}$ on the left and right. Hence, we only need to estimate the norm of the operator with kernel

$$1_{\{|x| < t/4\}}(x) e^{-\gamma\varphi(x)} U(t, x, y) e^{-\gamma\varphi(y)} 1_{\{|x| < t/4\}}(y),$$

using explicit knowledge of the kernel $U(t, x, y)$.

In odd dimensions, the kernel vanishes identically by the strong Huygens principle. In even dimensions, the kernel vanishes unless $|x|, |y| < t/4$, which implies that $|x - y| < t/2$ and thus

$$|1_{\{|x| < t/4\}}(x) U(t, x, y) 1_{\{|x| < t/4\}}(y)| \leq C t^{-n},$$

again from explicit formulas for $U(t, x, y)$. Schur's lemma then gives

$$\|1_{\{|x|<t/4\}}e^{-\gamma\varphi}U(t)e^{-\gamma\varphi}1_{\{|x|<t/4\}}\|_{L^2(\mathbb{R}^n)\rightarrow L^2(\mathbb{R}^n)}\leq Ct^{-n}.$$

Therefore we see that the integral in (1-4) actually converges for $\text{Im } \sigma \geq 0$ with the uniform estimate

$$\|e^{-\gamma\varphi}R_0(\sigma)e^{-\gamma\varphi}\|_{L^2\rightarrow L^2}\leq C|\sigma|^{-1}, \quad \text{Im } \sigma \geq 0 \text{ and } \text{Re } \sigma \geq 1.$$

The result for $-\gamma + \epsilon < \text{Im } \sigma < 0$ follows immediately by reflection from (1-1) and Lemma 1.1.

(2) In the case $\alpha = 0$ and $n = 1$, one can simply apply Schur's lemma to the Schwartz kernel

$$e^{-\gamma\varphi}R_0(x, y, \sigma)e^{-\gamma\varphi} = e^{-\gamma\varphi(x)} \frac{ie^{i\sigma|x-y|}}{\sigma} e^{-\gamma\varphi(y)}.$$

The $|\alpha| = 1, 2$ cases follow from the $|\alpha| = 0$ case by interpolation, as in [Zworski 1989, Lemma 3]; we supply a proof for the reader's convenience. Consider first the case $|\alpha| = 2$. By analytic continuation, if $u \in L^2(\mathbb{R}^n)$, then

$$(1-5) \quad \Delta R_0(\sigma)e^{-\gamma\varphi}u = -e^{-\gamma\varphi}u - \sigma^2 R_0(\sigma)e^{-\gamma\varphi}u$$

and hence $R_0(\sigma) : L^2_{-\gamma} \rightarrow H^2_{-\gamma}$ is bounded for $\text{Im } \sigma > -\gamma$. Now, choose $u \in L^2(\mathbb{R}^n)$ and set $f = R_0(\sigma)e^{-\gamma\varphi}u$. Then

$$(1-6) \quad \begin{aligned} \Delta(e^{-\gamma\varphi}R_0(\sigma)e^{-\gamma\varphi}u) \\ = (\gamma^2|\nabla\varphi|^2 - \gamma\Delta\varphi)e^{-\gamma\varphi}f - 2\gamma\nabla\varphi \cdot (e^{-\gamma\varphi}\nabla f) + e^{-\gamma\varphi}\Delta f \end{aligned}$$

In light of (1-5) it suffices to estimate the L^2 norm of $-\gamma\nabla\varphi \cdot (e^{-\gamma\varphi}\nabla f)$. But since φ has uniformly bounded derivatives,

$$\|\nabla\varphi \cdot (e^{-\gamma\varphi}\nabla f)\|_{L^2}^2 \leq C\|e^{-\gamma\varphi}\nabla f\|_{L^2}^2.$$

We now integrate by parts and estimate

$$(1-7) \quad \begin{aligned} \|e^{-\gamma\varphi}\nabla f\|_{L^2}^2 \\ \leq 2\int|\gamma\nabla\varphi||e^{-\gamma\varphi}\nabla f||e^{-\gamma\varphi}f|dx + \int|e^{-\gamma\varphi}\Delta f||e^{-\gamma\varphi}f|dx. \end{aligned}$$

Applying the inequality $2ab \leq 2a^2 + \frac{1}{2}b^2$ to the integrand, the first term on the right hand side is bounded by

$$\int 2|\gamma\nabla\varphi|^2|e^{-\gamma\varphi}f|^2dx + \int \frac{1}{2}|e^{-\gamma\varphi}\nabla f|^2dx,$$

while for the second term we use (1-5). We conclude that

$$\|e^{-\gamma\varphi}\nabla f\|_{L^2}^2 \leq C(1 + |\sigma|^2)\|e^{-\gamma\varphi}f\|_{L^2}^2 + \|e^{-2\gamma\varphi}u\|_{L^2}^2.$$

Returning to (1-6), it follows that

$$\|\Delta(e^{-\gamma\varphi}R_0(\sigma)e^{-\gamma\varphi}u)\|_{L^2} \leq C(1 + |\sigma|^2)\|u\|_{L^2}.$$

Moreover, (1-7) actually shows

$$\|\nabla(e^{-\gamma\varphi}R_0(\sigma)e^{-\gamma\varphi}u)\|_{L^2} \leq C\|u\|_{L^2}. \quad \square$$

We now introduce the semiclassical rescaling by setting $\lambda = h\sigma$. Let $R_0(\sigma, h)$ denote $(-h^2\Delta - \lambda^2)^{-1}$ and its corresponding analytic continuation. We are interested in λ lying in a set of the form

$$(a, b) + i((-\gamma + \epsilon)h, 1),$$

where $0 < a < b$. For the remainder of the paper, equip $H^k(\mathbb{R}^n)$ with the h -dependent norm $\|u\|_{H^k}^2 = \sum_{|\alpha| \leq k} \|(hD)^\alpha u\|_{L^2}^2$. Since $R_0(\lambda, h) = h^{-2}R_0(\lambda/h)$, we have uniform estimates

$$\|R_0(\lambda, h)\|_{L^2_\gamma \rightarrow H^s_{-\gamma}} = O(h^{-1}), \quad s = 0, 1, 2,$$

for $\lambda \in (a, b) + i((-\gamma + \epsilon)h, 1)$.

1B. Black box model. As our scattering problem, we consider exponentially decaying perturbations of the Laplacian outside a compact set, formulated in the black box setting as follows. Suppose \mathcal{H} is a Hilbert space with an orthogonal decomposition

$$\mathcal{H} = \mathcal{H}_{R_0} \oplus L^2(\mathbb{R}^n \setminus B(0, R_0))$$

where $B(x, R) = \{y \in \mathbb{R}^n : |x - y| < R\}$ and R_0 is fixed. The orthogonal projections onto \mathcal{H}_{R_0} and $L^2(\mathbb{R}^n \setminus B(0, R_0))$ will be denoted $1_{B(0, R_0)}u = u|_{B(0, R_0)}$ and $1_{\mathbb{R}^n \setminus B(0, R_0)}u = u|_{\mathbb{R}^n \setminus B(0, R_0)}$ for $u \in \mathcal{H}$. Note that any bounded continuous function $\chi \in C_b(\mathbb{R}^n)$ which is constant near $B(0, R_0)$ acts naturally on \mathcal{H} by

$$\chi u = C_0 u + (\chi - C_0)1_{\mathbb{R}^n \setminus B(0, R_0)}u,$$

where $\chi \equiv C_0$ near $B(0, R_0)$.

Now consider an unbounded self-adjoint operator $P(h)$ on \mathcal{H} with domain $\mathcal{D} \subset \mathcal{H}$ (independent of h for simplicity) with the following properties:

- If $u \in \mathcal{D}$, then $1_{\mathbb{R}^n \setminus B(0, R_0)}u \in H^2(\mathbb{R}^n \setminus B(0, R_0))$.
- If $u \in H^2(\mathbb{R}^n \setminus B(0, R_0))$ vanishes near $B(0, R_0)$, then $u \in \mathcal{D}$.

We assume there exists a real-valued and uniformly positive-definite matrix (a_{ij}) and a real-valued function V (which are allowed to be h -dependent) such that for $u \in \mathcal{D}$,

$$(1-8) \quad (P(h)u)|_{\mathbb{R}^n \setminus B(0, R_0)} = \left(- \sum_{i,j} (h\partial_i) a_{ij} (h\partial_j) + V \right) (u|_{\mathbb{R}^n \setminus B(0, R_0)}).$$

Furthermore, we require that

$$a_{ij}(x; h) \in C_b^\infty(\mathbb{R}^n \setminus B(0, R_0)) \quad \text{and} \quad V(x; h) \in C_b^\infty(\mathbb{R}^n \setminus B(0, R_0)),$$

with all derivatives uniformly bounded in h .

The perturbation is assumed to decay exponentially to the Laplacian in the sense that there exists $\gamma > 0$, $\delta > 0$ so that for $x \in \mathbb{R}^n \setminus B(0, R_0)$,

$$(1-9) \quad |a_{ij}(x; h) - \delta_{ij}| \leq C e^{-(2\gamma+\delta)|x|} \quad \text{and} \quad |V(x; h)| \leq C e^{-(2\gamma+\delta)|x|}.$$

Finally, assume that the mapping

$$(1-10) \quad 1_{B(0, R_0)}(P(h) + i)^{-1} : \mathcal{H} \rightarrow \mathcal{H}_{R_0}$$

is compact.

Under these hypotheses, we show that

$$R(\lambda, h) = (P(h) - \lambda^2)^{-1}, \quad \text{Re } \lambda > 0 \text{ and } \text{Im } \lambda > 0,$$

admits a meromorphic continuation to the strip $\text{Im } \lambda > (-\gamma + \epsilon)h$ as an operator $\mathcal{H}_\gamma \rightarrow \mathcal{H}_{-\gamma}$. In order that the associated weighted space \mathcal{H}_γ makes sense, we choose $\varphi \in C^\infty(\mathbb{R}^n)$, as above, satisfying $\varphi \equiv 0$ near $B(0, R_0)$.

Remark. All of the results in this note also apply to black box operators on the half-line $(0, \infty)$. For the most part this amounts to replacing the Laplacian on \mathbb{R}^n with the Dirichlet Laplacian on $(0, \infty)$, and replacing $H^s(\mathbb{R}^n)$ with $H^s(0, \infty) \cap H_0^1(0, \infty)$. Estimates for the free resolvent on $(0, \infty)$ follow from those on \mathbb{R} by the method of odd reflection; all other necessary modifications should be clear.

1C. Meromorphic continuation. As a preliminary, arbitrarily extend a_{ij} and V to functions defined on all of \mathbb{R}^n with the same properties as their original counterparts. Since the choice of extension has no bearing on the final result, we denote them by the same letters. Now define

$$\begin{aligned} \tilde{P}(h) &= - \sum_{i,j} (h\partial_i) a_{ij} (h\partial_j) + V, \\ \tilde{R}(\lambda, h) &= (\tilde{P}(h) - \lambda)^{-1}, \quad \lambda^2 \notin \sigma(\tilde{P}(h)). \end{aligned}$$

Since $\tilde{P}(h)$ is uniformly elliptic, it is self-adjoint with domain $H^2(\mathbb{R}^n)$. We will also write $A(h)$ for the difference

$$A(h) = \tilde{P}(h) - (-h^2 \Delta).$$

The important fact about $A(h)$ is that it is bounded as a map $H_\alpha^s \rightarrow H_{\alpha+2\gamma+\delta}^{s-2}$ for each $s, \alpha \in \mathbb{R}$.

We will need information about the $L_\gamma^2 \rightarrow H_\gamma^s$ mapping properties of $\tilde{R}(\lambda, h)$ for $\lambda^2 \notin \sigma(\tilde{P}(h))$.

Lemma 1.3. *Fix an interval $(a, b) \in \mathbb{R}_+$. For each $\gamma > 0$ there exists $T_0 > 0$ such that*

$$\|e^{\gamma\varphi} \tilde{R}(\lambda, h)e^{-\gamma\varphi}\|_{L^2 \rightarrow H^s} = O(|\operatorname{Im} \lambda|^{-1}), \quad s = 0, 1, 2$$

uniformly for $\lambda \in (a, b) + i(T_0h, 1)$.

Proof. Conjugating $\tilde{P}(h)$ by $e^{\gamma\varphi}$ yields

$$e^{\gamma\varphi} \tilde{P}(h)e^{-\gamma\varphi} = \tilde{P}(h) + h^2B,$$

where

$$B = \sum_{i,j} (2\gamma a_{ij} \partial_i \varphi) \partial_j - \gamma^2 a_{ij} \partial_i \varphi \partial_j \varphi + \gamma \partial_i (a_{ij} \partial_j \varphi)$$

is a first order operator with uniformly bounded coefficients. It follows that for $\lambda^2 \notin \sigma(\tilde{P}(h))$ (in particular for $\operatorname{Im} \lambda > 0$ and $\operatorname{Re} \lambda > 0$) we can write

$$e^{\gamma\varphi} \tilde{P}(h)e^{-\gamma\varphi} - \lambda^2 = (I + h^2B \tilde{R}(\lambda, h))(\tilde{P}(h) - \lambda^2).$$

Since $\tilde{P}(h) : H^2 \rightarrow L^2$ is self-adjoint,

$$\|u\|_{H^2} < C \|(\tilde{P}(h) + i)u\|_{L^2}.$$

It follows that for $\lambda \in (a, b) + i(0, 1)$,

$$(1-11) \quad \|\tilde{R}(\lambda, h)\|_{L^2 \rightarrow H^s} = O(|\operatorname{Im} \lambda|^{-1}), \quad s = 0, 1, 2.$$

We immediately deduce that

$$\|h^2B \tilde{R}(\lambda, h)\|_{L^2 \rightarrow L^2} = O(h|\operatorname{Im} \lambda|^{-1}) \leq \frac{1}{2}, \quad \lambda \in (a, b) + i[T_0h, 1),$$

for $T_0 > 0$ large enough. In particular, $I + h^2B \tilde{R}(\lambda, h) : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is invertible for $\lambda \in (a, b) + i[T_0h, 1)$ and

$$e^{\gamma\varphi} \tilde{R}(\lambda, h)e^{-\gamma\varphi} = \tilde{R}(\lambda, h)(I + h^2B \tilde{R}(\lambda, h))^{-1}.$$

This also shows that

$$\|e^{\gamma\varphi} \tilde{R}(\lambda, h)e^{-\gamma\varphi}\|_{L^2 \rightarrow H^s} = O(|\operatorname{Im} \lambda|^{-1}), \quad s = 0, 1, 2$$

for $\lambda \in (a, b) + i[T_0h, 1)$. □

The following lemma is useful in the proof of the meromorphic continuation. Equip \mathcal{D} with the h -dependent norm

$$\|u\|_{\mathcal{D}} = \|(P(h) + i)u\|_{\mathcal{H}}.$$

Then it is easy to see that under the uniform boundedness conditions on the derivatives of a_{ij} and V , the analog of [Sjöstrand and Zworski 1991, Proposition 4.1] remains true:

Lemma 1.4. *Suppose $\chi \in C_b^\infty(\mathbb{R}^n)$ has support disjoint from $\overline{B(0, R_0)}$. Then multiplication by χ is bounded $\mathcal{D} \rightarrow H^2(\mathbb{R}^n)$ and $H^2(\mathbb{R}^n) \rightarrow \mathcal{D}$ with a norm bounded independently of h .*

Proof. Consider first the map $\chi : \mathcal{D} \rightarrow H^2(\mathbb{R}^n)$. Since $\tilde{P}(h)$ is elliptic, we have the a priori estimate

$$\begin{aligned} \|\chi u\|_{H^2(\mathbb{R}^n)}^2 &\leq C_1 \left(\|\chi_1 \tilde{P}(h) \mathbf{1}_{\mathbb{R}^n \setminus B(0, R_0)} u\|_{L^2(\mathbb{R}^n \setminus B(0, R_0))}^2 \right. \\ &\quad \left. + \|\chi_1 \mathbf{1}_{\mathbb{R}^n \setminus B(0, R_0)} u\|_{L^2(\mathbb{R}^n \setminus B(0, R_0))}^2 \right) \\ &\leq C_2 \|(P(h) + i)u\|_{\mathcal{H}}^2, \end{aligned}$$

where $\chi_1 \equiv 1$ on $\text{supp } \chi$ and χ_1 also has support disjoint from $\overline{B(0, R_0)}$. All the constants are independent of h . For the case $\chi : H^2(\mathbb{R}^n) \rightarrow \mathcal{D}$ this is equivalent to the uniform boundedness of $\tilde{P}(h)$ on $H^2(\mathbb{R}^n)$, namely

$$\|\chi u\|_{\mathcal{D}} = \|(\tilde{P}(h) + i)(\chi u)\|_{L^2(\mathbb{R}^n)} \leq C \|u\|_{H^2(\mathbb{R}^n)}. \quad \square$$

In what follows, we will always be concerned with λ ranging in a precompact neighborhood of \mathbb{R}_+ . So fix $0 < a_0 < b_0$ and $\epsilon_0 > 0$, and define

$$\Omega(h) = (a_0, b_0) + i((-\gamma + \epsilon_0)h, 1).$$

For each $\epsilon > 0$, we also define a shrunken neighborhood,

$$\Omega_\epsilon(h) = (a_0 + \epsilon, b_0 - \epsilon) + i((-\gamma + \epsilon_0 + \epsilon)h, 1).$$

Proposition 1.5. *The function $R(\lambda, h)$ has a meromorphic continuation in $\Omega(h)$ as a family of bounded operators $\mathcal{H}_\gamma \rightarrow \mathcal{H}_{-\gamma}$.*

Proof. Choose cutoff functions $\chi, \chi_i \in C_c^\infty(\mathbb{R}^n)$, $i = 0, 1, 2$, so that $\chi_0 \equiv 1$ near $B(0, R_0)$ with $\chi_i \equiv 1$ on $\text{supp } \chi_{i-1}$ and $\chi \equiv 1$ on $\text{supp } \chi_2$. We can always choose these so that $\chi\varphi = 0$ and $\chi_i\varphi \equiv 0$. Approximate $R(\lambda, h)$ by a parametrix of the form $Q_0(\lambda, \lambda_0, h) + Q_1(\lambda_0, h)$ where

$$\begin{aligned} Q_0(\lambda, \lambda_0, h) &= (1 - \chi_0)(R_0(\lambda, h) - \tilde{R}(\lambda_0, h)A(h)R_0(\lambda, h))(1 - \chi_1), \\ Q_1(\lambda_0, h) &= \chi_2 R(\lambda_0, h) \chi_1; \end{aligned}$$

see also [Sá Barreto and Zworski 1995]. Here, $\lambda_0 = \lambda_0(h)$ denotes a point in $\Omega(h)$ with $\text{Im } \lambda_0 \geq T_0 h$. We now compute

$$(P(h) - \lambda^2)Q_0(\lambda, \lambda_0, h) = (1 - \chi_1) + K_0(\lambda, \lambda_0, h) + K_1(\lambda, \lambda_0, h)$$

where

$$\begin{aligned} K_0(\lambda, \lambda_0, h) &= -[\tilde{P}(h), \chi_0](R_0(\lambda, h) - \tilde{R}(\lambda_0, h)A(h)R_0(\lambda, h))(1 - \chi_1), \\ K_1(\lambda, \lambda_0, h) &= (1 - \chi_0)(\lambda^2 - \lambda_0^2)\tilde{R}(\lambda_0, h)A(h)R_0(\lambda, h)(1 - \chi_1), \end{aligned}$$

and

$$(P(h) - \lambda^2)Q_1(\lambda_0, h) = \chi_1 + K_2(\lambda_0, h) + K_3(\lambda, \lambda_0, h),$$

where

$$\begin{aligned} K_2(\lambda_0, h) &= -[\tilde{P}(h), \chi_2]R(\lambda_0, h)\chi_1, \\ K_3(\lambda, \lambda_0, h) &= \chi_2(\lambda_0^2 - \lambda^2)R(\lambda_0, h)\chi_1. \end{aligned}$$

If we let $K = K_0 + K_1 + K_2 + K_3$, then

$$(P(h) - \lambda^2)(Q_0(\lambda, \lambda_0, h) + Q_1(\lambda_0, h)) = I + K(\lambda, \lambda_0, h).$$

Note that if $\psi \in C_c^\infty(\mathbb{R}^n)$ then $[\tilde{P}(h), \psi]$ is a first order operator with compactly supported coefficients and $\|[\tilde{P}(h), \psi]\|_{H^1(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)} = O(h)$.

It is easy to see that $Q_0 + Q_1 : \mathcal{H}_\gamma \rightarrow \mathcal{H}_{-\gamma}$. For Q_0 this follows from the mapping properties of $R_0(\lambda, h)$, $A(h)$, and $\tilde{R}(\lambda_0, h)$. For Q_1 , this fact is trivial since Q_1 contains compactly supported cutoffs. We also remark that by the resolvent identity,

$$K_0(\lambda_0, \lambda_0, h) = -[\tilde{P}(h), \chi_0]\tilde{R}(\lambda_0, h)(1 - \chi_1).$$

To apply the Fredholm theory, we begin by showing that $K : \mathcal{H}_\gamma \rightarrow \mathcal{H}_\gamma$ is compact. First note that

$$K_0(\lambda, \lambda_0, h) = -[\tilde{P}(h), \chi_0](R_0(\lambda, h) - \tilde{R}(\lambda_0, h)A(h)R_0(\lambda, h))(1 - \chi_1) : \mathcal{H}_\gamma \rightarrow \mathcal{H}_\gamma$$

is compact: we see that $R_0(\lambda, h) : L_\gamma^2(\mathbb{R}^n) \rightarrow H_{-\gamma}^2(\mathbb{R}^n)$ and $\tilde{R}(\lambda_0, h)A(h)R_0(\lambda, h) : L_\gamma^2(\mathbb{R}^n) \rightarrow H_{\gamma+\delta}^2(\mathbb{R}^n)$. On the other hand $[\tilde{P}(h), \chi_0]$ is compactly supported and hence maps $H_\alpha^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n \setminus B(0, R_0))$ compactly for each $\alpha \in \mathbb{R}$.

Similarly, we can write

$$K_2(\lambda_0, h) = [\tilde{P}(h), \chi_2](1 - \chi_0)R(\lambda_0, h)\chi_1$$

which is compact since $(1 - \chi_0)R(\lambda_0, h)\chi_1 : \mathcal{H}_\gamma \rightarrow H^2(\mathbb{R}^n \setminus B(0, R_0))$ and $[\tilde{P}(h), \chi_2]$ is compactly supported. To see that K_1 is compact, again use that $\tilde{R}(\lambda_0, h)A(h)R_0(\lambda, h) : L_\gamma^2(\mathbb{R}^n) \rightarrow H_{\gamma+\delta}^2(\mathbb{R}^n)$ and now appeal to the fact that the inclusion

$$H_{\gamma+\delta}^2(\mathbb{R}^n) \hookrightarrow L_\gamma^2(\mathbb{R}^n)$$

is compact. Finally, the compactness of $K_3(\lambda, \lambda_0, h)$ follows from (1-10).

Next, we need to verify the invertibility of $I + K(\lambda, \lambda_0, h)$ for at least one value of $\lambda \in \Omega(h)$. Recall that multiplication by $(1 - \chi_0) : H^2(\mathbb{R}^n) \rightarrow \mathcal{D}$ is uniformly bounded in h . It follows that for $\lambda_0 \in \Omega(h)$ in the upper half-plane, for $u \in \mathcal{H}$,

$$\|(1 - \chi_0)R(\lambda_0, h)u\|_{H^2(\mathbb{R}^n)} \leq C_1\|(P(h) + i)R(\lambda_0, h)u\|_{\mathcal{H}} \leq C_2|\operatorname{Im} \lambda_0|^{-1}\|u\|_{\mathcal{H}},$$

and hence

$$\|(1 - \chi)R(\lambda_0, h)\|_{\mathcal{H} \rightarrow H^2(\mathbb{R}^n)} = O(|\operatorname{Im} \lambda_0|^{-1}), \quad \lambda_0 \in \Omega(h), \operatorname{Im} \lambda_0 > 0.$$

Here, we used

$$(P(h) + i)R(\lambda_0, h) = I + (\lambda_0^2 + i)R(\lambda_0, h)$$

and $R(\lambda_0, h) = O_{\mathcal{H} \rightarrow \mathcal{H}}(|\operatorname{Im} \lambda_0|^{-1})$. Combining this with (1-11), we see there exists $T_1 > T_0$ such that if $\lambda_0 \in \Omega(h)$ satisfies $\operatorname{Im} \lambda_0 \geq T_1 h$, then

$$\|K(\lambda_0, \lambda_0, h)\|_{\mathcal{H}_\gamma \rightarrow \mathcal{H}_\gamma} = O(h|\operatorname{Im} \lambda_0|^{-1}) \leq \frac{1}{2},$$

and hence $I + K(\lambda_0, \lambda_0, h)$ will be invertible. \square

Remark. The poles and their multiplicities of the extension obtained above do not depend on the particular choice of φ . Indeed, if φ_1 and φ_2 both vanish near $\mathbb{R}^n \setminus B(0, R_0)$ and equal $|x|$ for large $|x|$, then

$$e^{-\gamma\varphi_1} R(\lambda, h) e^{-\gamma\varphi_1} = e^{-\gamma(\varphi_1 - \varphi_2)} e^{-\gamma\varphi_2} R(\lambda, h) e^{-\gamma\varphi_2} e^{-\gamma(\varphi_1 - \varphi_2)}$$

and vice versa. Hence the poles and multiplicities of one such extension agree with those of any other.

Remark. As pointed out by the anonymous referee, an interesting question is whether $R(\lambda, h)$ can be continued to a larger region in the lower half plane when the perturbations are smooth functions of $\exp((-2\gamma - \delta)|x|)$ for large $|x|$ (and also whether the corresponding resolvent estimates hold). Such hypotheses are satisfied for stationary wave operators arising from black hole metrics with nondegenerate event horizons; see [Dyatlov 2011; Gannot 2014] for two examples.

At this point we need to introduce a new assumption on a reference operator $P^\sharp(h)$, defined as follows: choose $R_1 > R_0$ and $R_2 > 2R_1$ and let \mathbb{T} denote the torus $\mathbb{T} = (\mathbb{R}/R_2\mathbb{Z})^n$. Let

$$\mathcal{H}^\sharp = \mathcal{H}_{R_0} \oplus L^2(\mathbb{T} \setminus B(0, R_0)),$$

where $B(0, R_1)$ is considered a subset of \mathbb{T} . Define the dense subspace

$$\mathcal{D}^\sharp = \{u \in \mathcal{H}^\sharp : \psi u \in \mathcal{D}, (1 - \psi)u \in H^2(\mathbb{T})\},$$

where $\psi \in C_c^\infty(B(0, R_1))$ satisfies $\psi \equiv 1$ near $B(0, R_0)$. Now set

$$P^\sharp(h)u = P(h)\psi u + \left(- \sum_{i,j} (h\partial_i) a_{ij} (h\partial_j) + V \right) (1 - \psi)u, \quad u \in \mathcal{D}^\sharp.$$

Then $P^\sharp(h)$ is self-adjoint on \mathcal{D}^\sharp with discrete spectrum. We require that

$$(1-12) \quad \#\{z \in \sigma(P^\sharp(h)) : z \in [-L, L]\} \leq C(L/h^2)^{n^\sharp/2}$$

for some $n^\sharp \geq n$ and each $L \geq 1$. Here the eigenvalues are counted with multiplicity. If z_1, z_2, z_3, \dots are the eigenvalues of $P^\sharp(h)$ ordered so $|z_1| \leq |z_2| \leq |z_3| \leq \dots$, then the singular values of $(P^\sharp(h) - \lambda_0^2)^{-1}$ are $\mu_j((P^\sharp(h) - \lambda_0^2)^{-1}) = |z_j - \lambda_0^2|^{-1}$. If $\text{Im } \lambda_0 = T_1 h$, then (1-12) implies that there exists a constant $C > 0$ so that

$$\mu_j((P^\sharp(h) - \lambda_0^2)^{-1}) \leq Ch^{-2} j^{-2/n^\sharp}, \quad j > Ch^{-n^\sharp}.$$

2. Resolvent estimates

To estimate $R(\lambda, h)$, we make use of the following general fact [Gohberg and Kreĭn 1969, Chapter V, Theorem 5.1]: Suppose A is a compact operator lying in some p -class. If $(I + A)$ is invertible, then

$$\|(I + A)^{-1}\| \leq \frac{\det(I + |A|^p)}{|\det(I + A^p)|}.$$

We wish to apply this inequality to $(I + K)$, but first we need to verify that a suitable power of K is of trace class. Under our hypotheses we cannot estimate the singular values of K_2 ; nevertheless, the proof of Proposition 1.5 shows that $I + K_2(\lambda_0, h)$ is invertible on \mathcal{H}_γ for $\text{Im } \lambda_0 > T_1 h$, so we use the decomposition

$$(I + K(\lambda, \lambda_0, h)) = (I + K_2(\lambda_0, h))(I + \tilde{K}(\lambda, \lambda_0, h)),$$

where $\tilde{K} = (I + K_2)^{-1}(K_0 + K_1 + K_3)$. Note that $I + K$ and $I + \tilde{K}$ have the same poles.

2A. Singular values. From now on we will always choose $\lambda_0 \in \Omega(h)$ with fixed imaginary part $\text{Im } \lambda_0 = T_1 h$. Throughout, it will be clear that whenever an estimate depends on $\lambda_0 \in \Omega(h)$, it really only depends on $\text{Im } \lambda_0$.

Proposition 2.1. *The operator $\tilde{K}(\lambda, \lambda_0, h)^{n^\sharp+1} : \mathcal{H}_\gamma \rightarrow \mathcal{H}_\gamma$ is of trace class for $\lambda \in \Omega(h)$.*

Proof. We estimate the singular values of each summand in \tilde{K} . Since the weighted resolvent only continues to a narrow strip in the lower half-plane, in such a region it is particularly simple to estimate $\mu_j(K_0)$: choose an open ball $B \subseteq \mathbb{R}^n$ containing $\text{supp } \nabla \chi_0$ and let $-\Delta_B$ denote the Dirichlet Laplacian on B . Again using that the inclusion $1_{\mathbb{R}^n \setminus B(0, R_0)} : \mathcal{D}_\gamma \rightarrow H_\gamma^2$ is uniformly bounded in h , we consider K_0 as a map $\mathcal{H}_\gamma \rightarrow H^1(B)$. By Weyl asymptotics,

$$\mu_j((-h^2 \Delta_B)^{-1}) \leq Ch^{-2} j^{-2/n}, \quad j = 1, 2, 3, \dots$$

Thus, we estimate

$$\begin{aligned} \mu_j(K_0(\lambda, \lambda_0, h)) &\leq C \mu_j((-h^2 \Delta_B)^{-1/2}) \|(-h^2 \Delta_B)^{1/2} K_0(\lambda, \lambda_0, h)\|_{\mathcal{H}_\gamma \rightarrow L^2(B)} \\ &\leq Ch^{-3} j^{-1/n}, \quad \lambda \in \Omega(h). \end{aligned}$$

By the same reasoning we estimate $\mu_j(K_1)$, writing

$$\begin{aligned} \mu_j(K_1(\lambda, \lambda_0, h)) &\leq C \mu_j(e^{\gamma\varphi} \tilde{R}(\lambda_0, h) e^{-(\gamma+\delta)\varphi}) \|e^{(\gamma+\delta)\varphi} A(h) e^{\gamma\varphi} e^{-\gamma\varphi} R_0(\lambda)\|_{L^2_\gamma(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)}. \end{aligned}$$

In order to bound $\mu_j(e^{\gamma\varphi} \tilde{R}(\lambda_0, h) e^{-(\gamma+\delta)\varphi})$, let $P_0(h) = -h^2\Delta + x^2$ denote the harmonic oscillator. The inequality $\mu_j(P_0(h)^{-1}) \leq Ch^{-1} j^{-1/n}$ follows, in this case by explicit knowledge of the spectrum. Since $P_0(h)e^{-\delta\varphi} : H^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is bounded,

$$\begin{aligned} \mu_j(e^{\gamma\varphi} \tilde{R}(\lambda_0, h) e^{-(\gamma+\delta)\varphi}) &\leq \mu_j(P_0(h)^{-1}) \|P_0(h) e^{-\delta\varphi} e^{(\gamma+\delta)\varphi} \tilde{R}(\lambda_0, h) e^{-(\gamma+\delta)\varphi}\|_{L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)} \\ &\leq Ch^{-2} j^{-1/n}. \end{aligned}$$

Combined with the previous estimate we obtain

$$\mu_j(K_1) \leq Ch^{-3} j^{-1/n}, \quad \lambda \in \Omega(h).$$

Next we estimate the singular values of K_3 using (1-12). Recall that $(P(h) - \lambda^2)\chi = (P^\sharp(h) - \lambda^2)\chi$, which implies that

$$(P(h) - \lambda_0^2)^{-1}\chi_1 = \chi(P^\sharp(h) - \lambda_0^2)^{-1}\chi_1 - (P(h) - \lambda_0^2)^{-1}[P^\sharp(h), \chi](P^\sharp(h) - \lambda_0^2)^{-1}\chi_1.$$

Multiply this equation on the left by χ_2 and apply Fan's inequality, $\mu_{2k-1}(A+B) \leq \mu_k(A) + \mu_k(B)$. Using the fact that $(P(h) - \lambda_0^2)^{-1}[P^\sharp(h), \chi]$ has norm $O(1)$,

$$\mu_j(K_3(\lambda, \lambda_0, h)) \leq Ch^{-2} j^{-2/n^\sharp}, \quad j > Fh^{-n^\sharp}$$

for some constant $F > 0$. For $j \leq Fh^{-n^\sharp}$, we simply bound $\mu_j(K_3) \leq Ch^{-1}$ using the trivial norm estimate.

It is now clear that $\mu_j(K_j)^{n^\sharp}$ is summable for $j = 0, 1, 3$. \square

Applying the resolvent estimate as above, we obtain

$$\begin{aligned} (2-1) \quad \|R(\lambda, h)\|_{\mathcal{H}_\gamma \rightarrow \mathcal{H}_{-\gamma}} &\leq \|Q_0 + Q_1\|_{\mathcal{H}_\gamma \rightarrow \mathcal{H}_{-\gamma}} \|(I + K_2)^{-1}\|_{\mathcal{H}_\gamma \rightarrow \mathcal{H}_\gamma} \|(I + \tilde{K})^{-1}\|_{\mathcal{H}_\gamma \rightarrow \mathcal{H}_\gamma} \\ &\leq C \|Q_0 + Q_1\|_{\mathcal{H}_\gamma \rightarrow \mathcal{H}_{-\gamma}} \frac{\det(I + (\tilde{K}^* \tilde{K})^{\frac{n^\sharp+1}{2}})}{|\det(I + \tilde{K}^{n^\sharp+1})|}. \end{aligned}$$

Since

$$\|Q_0 + Q_1\|_{\mathcal{H}_\gamma \rightarrow \mathcal{H}_{-\gamma}} = O(h^{-2}), \quad \lambda \in \Omega(h),$$

it remains only to estimate the determinants. Define

$$f(\lambda, h) = \det(I + \tilde{K}^{n^\sharp+1}(\lambda, \lambda_0, h))$$

in $\Omega(h)$. By Weyl convexity inequalities, it follows that $|f(\lambda, h)| \leq M(h)$, $\lambda \in \Omega(h)$, where

$$M(h) = \sup_{\lambda \in \Omega(h)} \det(I + (\tilde{K}^* \tilde{K})^{\frac{n^\sharp+1}{2}}).$$

We therefore need to bound $M(h)$ from above and $|f(\lambda, h)|$ from below.

2B. Estimating the determinant from above. Here we obtain an upper bound for $M(h)$ of the form $M(h) \leq e^{Ch^{-p}}$. For the application in mind, the value of p is unimportant and we do not attempt to optimize the exponent. In fact h^{-p} also represents a polynomial upper bound for the number of resonances in a disk of radius h , but again obtaining an optimal value is unimportant in this context.

Proposition 2.2. *There exists $C > 0$ depending only on $\text{Im } \lambda_0$, and $p > 0$ such that*

$$M(h) \leq e^{Ch^{-p}}.$$

Proof. We estimate $M(h)$ using Fan's inequalities:

$$\begin{aligned} \prod_{j \geq 1} (1 + \mu_j(\tilde{K}^{n^\sharp+1})) &= \prod_{j \geq 1} (1 + \mu_j(\tilde{K})^{n^\sharp+1}) \leq \prod_{j \geq 1} (1 + \mu_{3j-2}(\tilde{K})^{n^\sharp+1})^3 \\ &\leq \prod_{j \geq 1} (1 + C_0(\mu_j(K_0)^{n^\sharp+1} + \mu_j(K_1)^{n^\sharp+1} + \mu_j(K_3)^{n^\sharp+1}))^3 \\ &\leq \prod_{i=0,1,3} \prod_{j \geq 1} (1 + C_0 \mu_j(K_i)^{n^\sharp+1})^3. \end{aligned}$$

For $i = 0, 1$, the singular values occurring in this product are bounded above by $\mu_j(K_i) \leq Ch^{-3} j^{-1/n^\sharp}$, and so we bound the product by the trace,

$$\prod_{j \geq 1} (1 + C_0 \mu_j(K_i)^{n^\sharp+1}) \leq \exp\left(C_1 h^{-3n^\sharp-3} \sum_{j \geq 1} j^{-1+1/n^\sharp}\right) \leq e^{Ch^{-3n^\sharp-3}}.$$

On the other hand for K_3 ,

$$\begin{aligned} \prod_{j \geq 1} (1 + C_0 \mu_j(K_3)^{n^\sharp+1}) &\leq \prod_{1 \leq j \leq Fh^{-n^\sharp}} (1 + C_0 \mu_j(K_3)^{n^\sharp+1}) \prod_{j > Fh^{-n^\sharp}} (1 + C_0 \mu_j(K_3)^{n^\sharp+1}) \\ &\leq \left(e^{Ch^{-n^\sharp} \log(1/h)}\right) \left(e^{Ch^{-n^\sharp}}\right). \end{aligned}$$

Thus,

$$M(h) \leq e^{Ch^{-p}}$$

for some $p > 0$, where the constant C only depends on $\text{Im } \lambda_0$. \square

2C. Estimating the determinant from below. Next we need to estimate $|f(\lambda, h)|$ from below. Note that λ_0 is not a zero of $f(\lambda, h)$ and that we have

$$(I + \tilde{K}(\lambda_0, \lambda_0, h)^{n^\sharp+1})^{-1} = I - \tilde{K}(\lambda_0, \lambda_0, h)^{n^\sharp+1}(I + \tilde{K}(\lambda_0, \lambda_0, h)^{n^\sharp+1})^{-1}.$$

By taking determinants and arguing as in the previous section, we obtain a lower bound at λ_0 ,

$$|f(\lambda_0, h)| \geq e^{-Ch^{-p}},$$

where the constant again depends only on $\text{Im } \lambda_0$. Since we can bound $|f(\lambda, h)|$ from above by $M(h)$ and from below at a chosen point, we are in a position to employ Cartan's principle [Levin 1972, Theorem 11] to obtain a lower bound away from resonances.

Proposition 2.3. *For each $\epsilon > 0$ there exists $C = C(\epsilon)$ such that*

$$|f(\lambda, h)| \geq e^{-Ah^{-p} \log(1/S(h))}, \quad \lambda \in \Omega_\epsilon(h) \setminus \bigcup_j D(r_j(h), S(h)),$$

where $S(h) \ll 1$ and $\{r_j(h)\}$ denote the resonances of $P(h)$ in $\Omega_\epsilon(h)$.

Proof. Rather than applying [Levin 1972, Theorem 11] directly, we prefer to control the set where the lower bound holds at the expense of the quality of the lower bound, just as in [Petkov and Zworski 2001]. For the reader's convenience we reproduce the proof, making the necessary adjustments.

Choose λ_0 with fixed real part. Define radii and disks

$$\rho_s(h) = T_1 + \gamma - \epsilon_0 - s\epsilon, \quad D_s(h) = D(\lambda_0, \rho_s(h)), \quad s = 1, 2, 3.$$

We see that $f(\lambda, h)$ is analytic in the disk $D_1(h)$. Let $r_j(h)$, $j = 1, \dots, N(h)$ denote the zeros of $f(\lambda, h)$ in $D_2(h)$, including multiplicity, and define the Blaschke product

$$\phi(\lambda, h) = \frac{(-\rho_2(h))^{N(h)}}{(r_1(h) - \lambda_0) \cdots (r_{N(h)}(h) - \lambda_0)} \prod_j \frac{\rho_2(h)(\lambda - r_j(h))}{\rho_2(h)^2 - (\overline{r_j(h)} - \lambda_0)(\lambda - \lambda_0)}.$$

Then ϕ has the same zeros as $f(\lambda, h)$, no poles in $D_2(h)$, and satisfies $\phi(\lambda_0, h) = 1$. Moreover, on the boundary of $D_2(h)$,

$$(2-2) \quad |\phi(\lambda, h)| = \frac{\rho_2^{N(h)}(h)}{|(\lambda_0 - r_1(h)) \cdots (\lambda_0 - r_{N(h)}(h))|} \geq 1.$$

Since the function defined by

$$\psi(\lambda, h) = \frac{f(\lambda, h)}{\phi(\lambda, h)}$$

has no zeros in $D_2(h)$, we may apply (2-2) and Carathéodory's estimate [Levin 1972, Theorem 8] to conclude that in $D_3(h)$ we have the lower bound

$$\begin{aligned} \log |\psi(\lambda, h)| &\geq -\frac{2\rho_3(h)}{\epsilon} \log \sup_{\lambda \in D_1(h)} |\psi(\lambda, h)| + \frac{\rho_2(h) + \rho_3(h)}{\epsilon} \log |\psi(\lambda_0, h)| \\ &\geq -\frac{2\rho_3(h)}{\epsilon} \log \sup_{\lambda \in D_2(h)} |f(\lambda, h)| + \frac{\rho_2(h) + \rho_3(h)}{\epsilon} \log |f(\lambda_0, h)|. \end{aligned}$$

It therefore suffices to bound $|\phi(\lambda, h)|$ from below in $D_3(h)$.

Outside the set $\bigcup_j D(r_j(h), S(h))$, the polynomial appearing in the numerator of $\phi(\lambda, h)$ is bounded below by $S(h)^{N(h)}$. On the other hand, the polynomial in the denominator of $\phi(\lambda, h)$ is bounded above in $D_3(h)$ by $\rho_2(h)^{N(h)}(\rho_2(h) + \rho_3(h))^{N(h)}$. Therefore

$$|\phi(\lambda, h)| \geq \left(\frac{S(h)}{\rho_2(h)(\rho_2(h) + \rho_3(h))} \right)^{N(h)}, \quad \lambda \in D_3(h) \setminus \bigcup_j D(r_j(h), S(h)).$$

Moreover, we can apply Jensen's formula to estimate the number of zeros $N(h)$ in $D_2(h)$ by

$$\begin{aligned} N(h) &\leq \frac{1}{\log \frac{\rho_1(h)}{\rho_2(h)}} (\log \sup_{\lambda \in D_1} |f(\lambda, h)| - \log |f(\lambda_0, h)|) \\ &\leq \frac{1}{\log \frac{\rho_1(h)}{\rho_2(h)}} (\log M(h) - \log |f(\lambda_0, h)|) \\ &= O(h^{-p}). \end{aligned}$$

Combining all the contributions, we obtain

$$|f(\lambda, h)| \geq e^{-Ch^{-p} \log(1/S(h))}, \quad \lambda \in D_3(h) \setminus \bigcup_j D(r_j(h), S(h)).$$

Since all the constants appearing are uniform in $\operatorname{Re} \lambda_0$, we can vary the real part in $\Omega_\epsilon(h)$ and obtain the necessary lower bound. Of course, ϵ is arbitrary and the result follows. \square

We can now establish our main theorem on resolvent estimates.

Theorem A. *For each $\epsilon > 0$, there exists $A = A(\epsilon)$ such that*

$$\|R(\lambda, h)\|_{\mathcal{H}_\gamma \rightarrow \mathcal{H}_{-\gamma}} < e^{Ah^{-p} \log(1/S(h))}, \quad \lambda \in \Omega_\epsilon(h) \setminus \bigcup_j D(r_j(h), S(h)),$$

where $S(h) \ll 1$ and $\{r_j(h)\}$ denote the resonances of $P(h)$ in $\Omega_\epsilon(h)$.

Proof. Apply Propositions 2.2 and 2.3 to (2-1). \square

3. From quasimodes to resonances

The passage from quasimodes to resonances is essentially an argument by contradiction. In the absence of resonances, the exponential bound appearing in Theorem A would hold throughout $\Omega_\epsilon(h)$; combined with the self-adjoint bound in the upper half-plane, an application of the “semiclassical maximum principle” implies a resolvent estimate on the real axis that contradicts the existence of a real quasimode. First results in this direction are due to Stefanov and Vodev [1996] who used the Phragmén–Lindelöf principle to show that having high energy real quasimodes implies the existence of resonances converging to the real axis. Bounds on the resolvent play a central role in that argument which go back to the work of Carleman [1936] on the completeness of sets of eigenfunctions. Tang and Zworski [1998] replaced the Phragmén–Lindelöf principle with a local version of the maximum principle which showed that there exists a resonance close to each quasimode. Stefanov further refined these method by dealing with multiplicities [1999], and modifying the maximum principle [2005] to allow the localization of resonances exponentially close to the real axis.

3A. Quasimodes. Suppose that $u(h) \in \mathcal{D}$ satisfies $\|u(h)\| = 1$ and

$$\text{supp } u(h) \subset K \quad \text{for a compact set } K \text{ independent of } h.$$

Suppose further that there exists $\lambda(h)^2 \in (a_0, b_0)$ such that

$$\|(P(h) - \lambda(h)^2)u(h)\| \leq R(h)$$

for a function $R(h) \geq 0$. We refer to such functions as quasimodes with accuracy $R(h)$. For the resolvent, choose a weight φ so that $\varphi \equiv 0$ on K . Also choose χ_1 with $\varphi \equiv 0$ on $\text{supp } \chi_1$ and $\chi_1 \equiv 1$ on K . Notice that for λ in the upper half-plane,

$$e^{-\gamma\varphi} R(\lambda, h) e^{-\gamma\varphi} (P(h) - \lambda^2)u = e^{-\gamma\varphi} R(\lambda, h) e^{-\gamma\varphi} (P(h) - \lambda^2) \chi_1 u = u,$$

and hence this equation holds away from poles by analytic continuation. We also recall the following standard fact: consider the Laurent expansion of $e^{-\gamma\varphi} R(\lambda, h) e^{-\gamma\varphi}$ near a resonance $r(h)$:

$$e^{-\gamma\varphi} R(\lambda, h) e^{-\gamma\varphi} = \text{holomorphic}(\lambda) + \sum_{j=1}^N A_j (\lambda^2 - r(h)^2)^{-j}.$$

Then, $\text{range}(A_j) \subseteq \text{range}(A_1)$ for $j = 1, \dots, N$. For a very general discussion of these types of results, see [Agmon 1998]. Consider the resonances $r_i(h)$ for $i = 1, \dots, N(h)$ contained in the set $\Omega_\epsilon(h)$, each with the associated residue $A_1^{(i)}$. If Π denotes the projection onto $\bigoplus_i \text{range}(A_1^{(i)})$, then $(I - \Pi)A_j^{(i)} = 0$ for each i, j .

Hence,

$$(I - \Pi) e^{-\gamma\varphi} R(\lambda, h) e^{-\gamma\varphi}$$

is holomorphic in $\Omega_\epsilon(h)$. By the maximum principle, this operator satisfies the bound given by Theorem A in a set slightly smaller than $\Omega_\epsilon(h)$ — see the proof of [Stefanov 1999, Theorem 1] or [Stefanov 2005, Theorem 3] for a precise statement.

3B. Semiclassical maximum principle. We now review the semiclassical maximum principle, as presented in [Stefanov 2005].

Lemma 3.1. *Let $a(h) < b(h)$ and suppose that $S_\pm(h)$, $\alpha(h)$, $w(h)$ are functions satisfying*

$$0 < S_+(h) \leq S_-(h), \quad 1 \leq \alpha(h), \quad S_-(h)\alpha(h) \log \alpha(h) \leq w(h).$$

Also, suppose $F(\lambda, h)$ is a holomorphic function defined in a neighborhood of

$$[a(h) - w(h), b(h) + w(h)] + i[-\alpha(h)S_-(h), S_+(h)].$$

If

$$\begin{cases} |F(\lambda, h)| \leq e^{\alpha(h)}, & \lambda \in [a(h) - w(h), b(h) + w(h)] + i[-\alpha(h)S_-(h), S_+(h)], \\ |F(\lambda, h)| \leq M(h), & \lambda \in [a(h) - w(h), b(h) + w(h)] + iS_+(h), \end{cases}$$

with $M(h) \geq 1$, then there exists $h_1 = h_1(S_-, S_+, \alpha) > 0$ such that

$$|F(\lambda, h)| \leq e^3 M(h), \quad \lambda \in [a(h), b(h)] + i[S_-(h), S_+(h)]$$

for $h \leq h_1$.

For our application, we will apply this lemma with

- $S_-(h) = S_+(h) = S(h)$,
- $F(\lambda, h) = (I - \Pi) e^{-\gamma\varphi} R(\lambda, h) e^{-\gamma\varphi}$,
- $\alpha(h) = Ch^{-p} \log(1/S(h))$,
- $M(h) = 1/S(h)$.

The choice of $S(h)$ and $w(h)$ is made as in [Stefanov 2005] according to the accuracy $R(h)$ of the quasimodes.

3C. Lower bounds on the number of resonances. Here we state the main theorem on the existence of resonances rapidly converging to the real axis. We refer to [Stefanov 2005, Theorem 3] for the proof; the only modification is that instead of a compactly truncated resolvent $(I - \Pi)\chi R(\lambda, h)\chi$, we use $(I - \Pi)e^{-\gamma\varphi} R(\lambda, h)e^{-\gamma\varphi}$.

Theorem B. *Let $P(h)$ satisfy the black box hypotheses. Let $0 < a_0 < a(h) < b(h) < b_0 < \infty$. Assume there is an h_0 such that for $h < h_0$ there exists $m(h) \in \{1, 2, \dots\}$, $\lambda_n(h)^2 \in [a(h), b(h)]$, and $u_n(h) \in \mathcal{D}$ with $\|u_n(h)\| = 1$ for $1 \leq n \leq m(h)$ such that $\text{supp } u_n(h) \subset K$ for a compact set K , independent of h . Suppose further that*

- (1) $\|(P(h) - \lambda_n(h)^2)u_n(h)\| \leq R(h)$,
- (2) whenever a collection $\{v_n(h)\}_{n=1}^{m(h)} \subset \mathcal{H}$ satisfies $\|u_n(h) - v_n(h)\| < h^N/M$, $\{v_n(h)\}_{n=1}^{m(h)}$ is linearly independent,

where $R(h) \leq h^{p+N+1}/C \log(1/h)$ and $C \gg 1$, $N \geq 0$, $M > 0$. Then there exists $C_0 > 0$ depending on a_0, b_0 and the operator $P(h)$ such that for $B > 0$ there exists $h_1 < h_0$ depending on A, B, M, N so that the following holds: whenever $h \in (0, h_1)$, the operator $P(h)$ has at least $m(h)$ resonances in the strip

$$\left[a(h) - c(h) \log \frac{1}{h}, b(h) + c(h) \log \frac{1}{h} \right] - i[0, c(h)]$$

where $c(h) = \max(C_0 B M R(h) h^{-p-N-1}, e^{-B/h})$.

Acknowledgments

I would like to thank Maciej Zworski for suggesting the problem, along with many valuable conversations. Thanks to the anonymous referee for suggesting improvements in the exposition. I am also grateful to Jeffrey Galkowski for his interest in the problem and some helpful discussions.

References

- [Agmon 1998] S. Agmon, “A perturbation theory of resonances”, *Comm. Pure Appl. Math.* **51**:11-12 (1998), 1255–1309. MR 99i:47023 Zbl 0941.47008
- [Burq 2002] N. Burq, “Semi-classical estimates for the resolvent in nontrapping geometries”, *Int. Math. Res. Not.* **2002**:5 (2002), 221–241. MR 2002k:81069 Zbl 1161.81368
- [Carleman 1936] T. Carleman, “Über die asymptotische Verteilung der Eigenwerte partieller Differentialgleichungen”, *Ber. Verh. Sächs. Akad. Wiss. Leipzig. Math.-Phys. Kl.* **88** (1936), 119–134. Zbl 0017.11402
- [Dyatlov 2011] S. Dyatlov, “Quasi-normal modes and exponential energy decay for the Kerr–de Sitter black hole”, *Comm. Math. Phys.* **306**:1 (2011), 119–163. MR 2012g:58055 Zbl 1223.83029
- [Gannot 2014] O. Gannot, “Quasinormal modes for Schwarzschild–AdS black holes: exponential convergence to the real axis”, *Comm. Math. Phys.* **330**:2 (2014), 771–799. MR 3223487 Zbl 1295.85001
- [Gohberg and Kreĭn 1969] I. C. Gohberg and M. G. Kreĭn, *Introduction to the theory of linear nonselfadjoint operators*, Translations of Mathematical Monographs **18**, American Mathematical Society, Providence, RI, 1969. MR 39 #7447 Zbl 0181.13504
- [Holzegel and Smulevici 2014] G. Holzegel and J. Smulevici, “Quasimodes and a lower bound on the uniform energy decay rate for Kerr–AdS spacetimes”, *Anal. PDE* **7**:5 (2014), 1057–1090. MR 3265959 Zbl 1300.83030 arXiv 1303.5944
- [Ikawa 1982] M. Ikawa, “Decay of solutions of the wave equation in the exterior of two convex obstacles”, *Osaka J. Math.* **19**:3 (1982), 459–509. MR 84e:35018 Zbl 0498.35008
- [Lax and Phillips 1989] P. D. Lax and R. S. Phillips, *Scattering theory*, 2nd ed., Pure and Applied Mathematics **26**, Academic Press, Boston, 1989. MR 90k:35005 Zbl 0697.35004

- [Levin 1972] B. Y. Levin, *Distribution of zeros of entire functions*, American Mathematical Society, Providence, RI, 1972. MR 28 #217 Zbl 0152.06703
- [McLeod 1967] J. B. McLeod, “The analytic continuation to the unphysical sheet of the resolvent kernel associated with the Schroedinger operator”, *Quart. J. Math. Oxford Ser. (2)* **18** (1967), 219–231. MR 37 #608a Zbl 0152.11401
- [Petkov and Zworski 2001] V. Petkov and M. Zworski, “Semi-classical estimates on the scattering determinant”, *Ann. Henri Poincaré* **2**:4 (2001), 675–711. MR 2002h:35222 Zbl 1041.81041
- [Rauch 1978] J. Rauch, “Local decay of scattering solutions to Schrödinger’s equation”, *Comm. Math. Phys.* **61**:2 (1978), 149–168. MR 58 #14590 Zbl 0381.35023
- [Sá Barreto and Zworski 1995] A. Sá Barreto and M. Zworski, “Existence of resonances in three dimensions”, *Comm. Math. Phys.* **173**:2 (1995), 401–415. MR 96j:35183 Zbl 0835.35099
- [Sjöstrand 1997] J. Sjöstrand, “A trace formula and review of some estimates for resonances”, pp. 377–437 in *Microlocal analysis and spectral theory* (Lucca, 1996), edited by L. Rodino, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci. **490**, Kluwer Academic, Dordrecht, 1997. MR 99e:47064 Zbl 0877.35090
- [Sjöstrand and Zworski 1991] J. Sjöstrand and M. Zworski, “Complex scaling and the distribution of scattering poles”, *J. Amer. Math. Soc.* **4**:4 (1991), 729–769. MR 92g:35166 Zbl 0752.35046
- [Stefanov 1999] P. Stefanov, “Quasimodes and resonances: sharp lower bounds”, *Duke Math. J.* **99**:1 (1999), 75–92. MR 2000k:35230 Zbl 0952.47013
- [Stefanov 2005] P. Stefanov, “Approximating resonances with the complex absorbing potential method”, *Comm. Partial Differential Equations* **30**:10-12 (2005), 1843–1862. MR 2006i:35266 Zbl 1095.35017
- [Stefanov and Vodev 1996] P. Stefanov and G. Vodev, “Neumann resonances in linear elasticity for an arbitrary body”, *Comm. Math. Phys.* **176**:3 (1996), 645–659. MR 96k:35122 Zbl 0851.35032
- [Tang and Zworski 1998] S.-H. Tang and M. Zworski, “From quasimodes to resonances”, *Math. Res. Lett.* **5**:3 (1998), 261–272. MR 99i:47088 Zbl 0913.35101
- [Vodev 1994] G. Vodev, “Sharp bounds on the number of scattering poles in the two-dimensional case”, *Math. Nachr.* **170** (1994), 287–297. MR 95i:35209 Zbl 0829.35091
- [Warnick 2015] C. M. Warnick, “On quasinormal modes of asymptotically anti-de Sitter black holes”, *Comm. Math. Phys.* **333**:2 (2015), 959–1035. MR 3296168 Zbl 06394860
- [Wunsch 2012] J. Wunsch, “Resolvent estimates with mild trapping”, preprint, 2012. arXiv 1209.0843
- [Zworski 1989] M. Zworski, “Sharp polynomial bounds on the number of scattering poles”, *Duke Math. J.* **59**:2 (1989), 311–323. MR 90h:35190 Zbl 0705.35099

Received August 6, 2013. Revised December 10, 2014.

ORAN GANNOT
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CALIFORNIA, BERKELEY
EVANS HALL
BERKELEY, CA 94720
UNITED STATES
ogannot@math.berkeley.edu

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 277 No. 1 September 2015

Real positivity and approximate identities in Banach algebras	1
DAVID P. BLECHER and NARUTAKA OZAWA	
On shrinking gradient Ricci solitons with nonnegative sectional curvature	61
MINGLIANG CAI	
From quasimodes to resonances: exponentially decaying perturbations	77
ORAN GANNOT	
A general simple relative trace formula	99
JAYCE R. GETZ and HEEKYOUNG HAHN	
Chern-Simons functions on toric Calabi-Yau threefolds and Donaldson-Thomas theory	119
ZHENG HUA	
On the flag curvature of a class of Finsler metrics produced by the navigation problem	149
LIBING HUANG and XIAOHUAN MO	
Angular distribution of diameters for spheres and rays for planes	169
NOBUHIRO INNAMI and YUYA UNEME	
A note on an L^p -Brunn–Minkowski inequality for convex measures in the unconditional case	187
ARNAUD MARSIGLIETTI	
Structure of seeds in generalized cluster algebras	201
TOMOKI NAKANISHI	
Inequalities of Alexandrov–Fenchel type for convex hypersurfaces in hyperbolic space and in the sphere	219
YONG WEI and CHANGWEI XIONG	
Upper bounds of root discriminant lower bounds	241
SIMAN WONG	



0030-8730(201509)277:1;1-V