

*Pacific
Journal of
Mathematics*

**A NOTE ON AN L^p -BRUNN-MINKOWSKI INEQUALITY
FOR CONVEX MEASURES IN THE UNCONDITIONAL CASE**

ARNAUD MARSIGLIETTI

Volume 277 No. 1

September 2015

A NOTE ON AN L^p -BRUNN–MINKOWSKI INEQUALITY FOR CONVEX MEASURES IN THE UNCONDITIONAL CASE

ARNAUD MARSIGLIETTI

We consider a different L^p -Minkowski combination of compact sets in \mathbb{R}^n than the one introduced by Firey and we prove an L^p -Brunn–Minkowski inequality, $p \in [0, 1]$, for a general class of measures called convex measures that includes log-concave measures, under unconditional assumptions. As a consequence, we derive concavity properties of the function $t \mapsto \mu(t^{1/p}A)$, $p \in (0, 1]$, for unconditional convex measures μ and unconditional convex body A in \mathbb{R}^n . We also prove that the (B)-conjecture for all uniform measures is equivalent to the (B)-conjecture for all log-concave measures, completing recent works by Saroglou.

1. Introduction

The Brunn–Minkowski inequality is a fundamental inequality which states that, for every convex subset $A, B \subset \mathbb{R}^n$ and for every $\lambda \in [0, 1]$, one has

$$(1) \quad |(1-\lambda)A + \lambda B|^{\frac{1}{n}} \geq (1-\lambda)|A|^{\frac{1}{n}} + \lambda|B|^{\frac{1}{n}},$$

where

$$A + B = \{a + b : a \in A, b \in B\}$$

denotes the *Minkowski sum* of A and B and where $|\cdot|$ denotes Lebesgue measure. The inequality and its consequences are well covered in the book [Schneider 1993] and the survey [Gardner 2002].

Several extensions of the Brunn–Minkowski inequality have been developed during the last decades by establishing functional versions (see, e.g., [Henstock and Macbeath 1953; Dubuc 1977; Dancs and Uhrin 1980; Uhrin 1994]), by considering different measures (see, e.g., [Borell 1974; 1975]), by generalizing the Minkowski sum (see, e.g., [Firey 1961; 1962; 1964; Lutwak 1993; 1996]), among others.

This research was supported in part by the Institute for Mathematics and its Applications with funds provided by the National Science Foundation.

MSC2010: 28A75, 52A40, 60B11.

Keywords: Brunn–Minkowski–Firey theory, L^p -Minkowski combination, convex body, convex measure, (B)-conjecture.

In this paper, we will combine these extensions to prove an L^p -Brunn–Minkowski inequality for a large class of measures, including the log-concave measures.

Firstly, let us consider measures other than Lebesgue measure. Following Borell [1974; 1975], we say that a Borel measure μ in \mathbb{R}^n is s -concave, $s \in [-\infty, +\infty]$, if the inequality

$$\mu((1 - \lambda)A + \lambda B) \geq M_s^\lambda(\mu(A), \mu(B))$$

holds for every $\lambda \in [0, 1]$ and for every compact subset $A, B \subset \mathbb{R}^n$ such that $\mu(A)\mu(B) > 0$. Here $M_s^\lambda(a, b)$ denotes the s -mean of the nonnegative real numbers a, b with weight λ , defined as

$$M_s^\lambda(a, b) = ((1 - \lambda)a^s + \lambda b^s)^{\frac{1}{s}} \quad \text{if } s \notin \{-\infty, 0, +\infty\},$$

$M_{-\infty}^\lambda(a, b) = \min(a, b)$, $M_0^\lambda(a, b) = a^{1-\lambda}b^\lambda$, $M_{+\infty}^\lambda(a, b) = \max(a, b)$. Hence the Brunn–Minkowski inequality tells us that Lebesgue measure in \mathbb{R}^n is $\frac{1}{n}$ -concave.

As a consequence of the Hölder inequality, one has $M_p^\lambda(a, b) \leq M_q^\lambda(a, b)$ for every $p \leq q$. Thus every s -concave measure is $-\infty$ -concave. The $-\infty$ -concave measures are also called *convex measures*.

For $s \leq \frac{1}{n}$, Borell showed that every measure μ which is absolutely continuous with respect to n -dimensional Lebesgue measure is s -concave if and only if its density is an α -concave function, with

$$(2) \quad \alpha = \frac{s}{1 - sn} \in \left[-\frac{1}{n}, +\infty\right].$$

A function $f : \mathbb{R}^n \rightarrow [0, +\infty)$ is said to be α -concave, with $\alpha \in [-\infty, +\infty]$, if the inequality

$$f((1 - \lambda)x + \lambda y) \geq M_\alpha^\lambda(f(x), f(y))$$

holds for every $x, y \in \mathbb{R}^n$ such that $f(x)f(y) > 0$ and for every $\lambda \in [0, 1]$.

Secondly, let us consider a generalization of the notion of the Minkowski sum introduced by Firey, which leads to an L^p -Brunn–Minkowski theory. For *convex bodies* A and B in \mathbb{R}^n (i.e., compact convex sets containing the origin in the interior), the L^p -Minkowski combination, $p \in [-\infty, +\infty]$, of A and B with weight $\lambda \in [0, 1]$ is defined by

$$(1 - \lambda) \cdot A \oplus_p \lambda \cdot B = \{x \in \mathbb{R}^n : \langle x, u \rangle \leq M_p^\lambda(h_A(u), h_B(u)) \text{ for all } u \in S^{n-1}\},$$

where h_A denotes the *support function* of A defined by

$$h_A(u) = \max_{x \in A} \langle x, u \rangle, \quad u \in S^{n-1}.$$

Notice that, for every $p \leq q$, one has

$$(1 - \lambda) \cdot A \oplus_p \lambda \cdot B \subset (1 - \lambda) \cdot A \oplus_q \lambda \cdot B.$$

The support function is an important tool in convex geometry: it has the property of determining the convex body, since

$$A = \{x \in \mathbb{R}^n : \langle x, u \rangle \leq h_A(u) \text{ for all } u \in S^{n-1}\},$$

and it is linear with respect to Minkowski sum and dilation:

$$h_{A+B} = h_A + h_B, \quad h_{\mu A} = \mu h_A$$

($A, B \subset \mathbb{R}^n$ and $\mu \geq 0$). Thus,

$$(1 - \lambda) \cdot A \oplus_1 \lambda \cdot B = (1 - \lambda)A + \lambda B.$$

In this paper, we consider a different L^p -Minkowski combination. We denote by \mathbb{R}_+ the set of nonnegative real numbers. Recall that a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is *unconditional* if there exists a basis (a_1, \dots, a_n) of \mathbb{R}^n (the canonical basis in the sequel) such that, for every $x = \sum_{i=1}^n x_i a_i \in \mathbb{R}^n$ and for every $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n) \in \{-1, 1\}^n$, one has $f(\sum_{i=1}^n \varepsilon_i x_i a_i) = f(x)$. A measure which is absolutely continuous with respect to n -dimensional Lebesgue measure is *unconditional* if its density function is unconditional. For $\mathbf{p} = (p_1, \dots, p_n) \in [-\infty, +\infty]^n$, $a = (a_1, \dots, a_n) \in (\mathbb{R}_+)^n$, $b = (b_1, \dots, b_n) \in (\mathbb{R}_+)^n$ and $\lambda \in [0, 1]$, let us denote

$$(1 - \lambda)a +_{\mathbf{p}} \lambda b = (M_{p_1}^\lambda(a_1, b_1), \dots, M_{p_n}^\lambda(a_n, b_n)) \in (\mathbb{R}_+)^n.$$

For subsets $A, B \subset \mathbb{R}^n$ such that $A \cap (\mathbb{R}_+)^n$ and $B \cap (\mathbb{R}_+)^n$ are nonempty, for $\mathbf{p} \in [-\infty, +\infty]^n$ and for $\lambda \in [0, 1]$, we define the L^p -Minkowski combination of A and B with weight λ , denoted by $(1 - \lambda) \cdot A +_{\mathbf{p}} \lambda \cdot B$, to be the unconditional subset (i.e., the indicator function is unconditional) such that

$$((1 - \lambda) \cdot A +_{\mathbf{p}} \lambda \cdot B) \cap (\mathbb{R}_+)^n = \{(1 - \lambda)a +_{\mathbf{p}} \lambda b : a \in A \cap (\mathbb{R}_+)^n, b \in B \cap (\mathbb{R}_+)^n\}.$$

This definition is consistent with the well known fact that an unconditional set (or function) is determined by its restriction to the positive octant $(\mathbb{R}_+)^n$. Moreover, this L^p -Minkowski combination coincides with the classical Minkowski sum when $\mathbf{p} = (1, \dots, 1)$ and A, B are unconditional convex subsets of \mathbb{R}^n (see Proposition 2.1).

Using an extension of the Brunn-Minkowski inequality discovered by Uhrin [1994], we prove the following result:

Theorem 1.1. *Let $\mathbf{p} = (p_1, \dots, p_n) \in [0, 1]^n$ and $\alpha \in \mathbb{R}$ with $\alpha \geq -(\sum_{i=1}^n p_i^{-1})^{-1}$. Let μ be an unconditional measure in \mathbb{R}^n that has an α -concave density function with respect to Lebesgue measure. Then, for every unconditional convex body A, B in \mathbb{R}^n and for every $\lambda \in [0, 1]$,*

$$(3) \quad \mu((1 - \lambda) \cdot A +_{\mathbf{p}} \lambda \cdot B) \geq M_\gamma^\lambda(\mu(A), \mu(B)),$$

where $\gamma = (\sum_{i=1}^n p_i^{-1} + \alpha^{-1})^{-1}$.

In Theorem 1.1, if α or one of the p_i is equal to 0, then $(\sum_{i=1}^n p_i^{-1})^{-1}$ and γ are defined by continuity and are equal to 0.

The case of Lebesgue measure and $\mathbf{p} = (0, \dots, 0)$ is treated by Saroglou [2015], answering a conjecture by Böröczky, Lutwak, Yang and Zhang [Böröczky et al. 2012] in the unconditional case.

Conjecture 1.2 (log-Brunn–Minkowski inequality [Böröczky et al. 2012]). *Let A, B be symmetric convex bodies in \mathbb{R}^n and let $\lambda \in [0, 1]$. Then*

$$(4) \quad |(1 - \lambda) \cdot A \oplus_0 \lambda \cdot B| \geq |A|^{1-\lambda} |B|^\lambda.$$

Useful links between Conjecture 1.2 and the (B)-conjecture have been discovered by Saroglou [2014; 2015].

Conjecture 1.3 ((B)-conjecture [Latała 2002; Cordero-Erausquin et al. 2004]). *Let μ be a symmetric log-concave measure in \mathbb{R}^n and let A be a symmetric convex subset of \mathbb{R}^n . Then the function $t \mapsto \mu(e^t A)$ is log-concave on \mathbb{R} .*

The (B)-conjecture was solved by Cordero-Erausquin, Fradelizi and Maurey [Cordero-Erausquin et al. 2004] for the Gaussian measure and for the unconditional case. As a variant of the (B)-conjecture, one may study concavity properties of the function $t \mapsto \mu(V(t)A)$ where $V : \mathbb{R} \rightarrow \mathbb{R}_+$ is a convex function. As a consequence of Theorem 1.1, we deduce concavity properties of the function $t \mapsto \mu(t^{\frac{1}{p}} A)$, $p \in (0, 1]$, for every unconditional s -concave measure μ and every unconditional convex body A in \mathbb{R}^n (see Proposition 2.4).

Saroglou [2014] has also proved that the log-Brunn–Minkowski inequality for Lebesgue measure — which is to say, inequality (4) — is equivalent to the log-Brunn–Minkowski inequality for all log-concave measures. We continue these kinds of equivalences by proving that the (B)-conjecture for all uniform measures is equivalent to the (B)-conjecture for all log-concave measures (see Proposition 3.1).

We also investigate functional versions of the (B)-conjecture, which may be read as follows:

Conjecture 1.4 (functional version of the (B)-conjecture). *Let $f, g : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be even log-concave functions. Then the function*

$$t \mapsto \int_{\mathbb{R}^n} f(e^{-t}x)g(x) dx$$

is log-concave on \mathbb{R} .

We prove that Conjecture 1.4 is equivalent to Conjecture 1.3 (see Proposition 3.2).

Let us note that other developments in the use of the earlier mentioned extensions of the Brunn–Minkowski inequality have been recently made as well. See, e.g., [Bobkov et al. 2014; Caglar and Werner 2014; Caglar et al. 2015; Gardner et al. 2014].

The rest of the paper is organized as follows: in the next section, we prove Theorem 1.1 and we extend it to m sets, $m \geq 2$. We also compare our L^p -Minkowski combination to the Firey combination and derive an L^p -Brunn–Minkowski inequality for the Firey combination. We then discuss the consequences of a variant of the (B)-conjecture, namely we deduce concavity properties of the function $t \mapsto \mu(t^{\frac{1}{p}} A)$, $p \in (0, 1]$. In Section 3, we prove that the (B)-conjecture for all uniform measures is equivalent to the (B)-conjecture for all log-concave measures, and we also prove that the (B)-conjecture is equivalent to its functional version, Conjecture 1.4.

2. Proof of Theorem 1.1 and consequences

Before proving Theorem 1.1, let us show that our L^p -Minkowski combination coincides with the classical Minkowski sum when $\mathbf{p} = (1, \dots, 1)$, for unconditional convex sets.

Proposition 2.1. *Let A, B be unconditional convex subsets of \mathbb{R}^n and let $\lambda \in [0, 1]$. Then*

$$(1 - \lambda) \cdot A +_{\mathbf{1}} \lambda \cdot B = (1 - \lambda)A + \lambda B,$$

where $\mathbf{1} = (1, \dots, 1)$.

Proof. Since the sets $(1 - \lambda) \cdot A +_{\mathbf{1}} \lambda \cdot B$ and $(1 - \lambda)A + \lambda B$ are unconditional, it is sufficient to prove that

$$((1 - \lambda) \cdot A +_{\mathbf{1}} \lambda \cdot B) \cap (\mathbb{R}_+)^n = ((1 - \lambda)A + \lambda B) \cap (\mathbb{R}_+)^n.$$

Let $x \in ((1 - \lambda)A + \lambda B) \cap (\mathbb{R}_+)^n$. There exists $a = (a_1, \dots, a_n) \in A$ and $b = (b_1, \dots, b_n) \in B$ such that $x = (1 - \lambda)a + \lambda b$ and, for every $i \in \{1, \dots, n\}$, $(1 - \lambda)a_i + \lambda b_i \in \mathbb{R}_+$. Let $\varepsilon, \eta \in \{-1, 1\}^n$ such that $(\varepsilon_1 a_1, \dots, \varepsilon_n a_n) \in (\mathbb{R}_+)^n$ and $(\eta_1 b_1, \dots, \eta_n b_n) \in (\mathbb{R}_+)^n$. Notice that, for every $i \in \{1, \dots, n\}$, $0 \leq (1 - \lambda)a_i + \lambda b_i \leq (1 - \lambda)\varepsilon_i a_i + \lambda \eta_i b_i$. Since the sets A and B are convex and unconditional, it follows that $x \in (1 - \lambda)(A \cap (\mathbb{R}_+)^n) + \lambda(B \cap (\mathbb{R}_+)^n) = ((1 - \lambda) \cdot A +_{\mathbf{1}} \lambda \cdot B) \cap (\mathbb{R}_+)^n$.

The other inclusion is clear due to the definition of the set $(1 - \lambda) \cdot A +_{\mathbf{1}} \lambda \cdot B$. \square

Proof of Theorem 1.1. Let $\lambda \in [0, 1]$ and let A, B be unconditional convex bodies in \mathbb{R}^n .

It has been shown by Uhrin [1994] that if $f, g, h : (\mathbb{R}_+)^n \rightarrow \mathbb{R}_+$ are bounded measurable functions such that, for every $x, y \in (\mathbb{R}_+)^n$, $h((1 - \lambda)x +_{\mathbf{p}} \lambda y) \geq M_\alpha^\lambda(f(x), g(y))$, then

$$\int_{(\mathbb{R}_+)^n} h(x) dx \geq M_\gamma^\lambda \left(\int_{(\mathbb{R}_+)^n} f(x) dx, \int_{(\mathbb{R}_+)^n} g(x) dx \right),$$

where $\gamma = (\sum_{i=1}^n p_i^{-1} + \alpha^{-1})^{-1}$.

Let us denote by ϕ the density function of μ and let us set $h = 1_{(1-\lambda) \cdot A +_p \lambda \cdot B} \phi$, $f = 1_A \phi$ and $g = 1_B \phi$. By assumption, the function ϕ is unconditional and α -concave, hence ϕ is nonincreasing in each coordinate on the octant $(\mathbb{R}_+)^n$. Then for every $x, y \in (\mathbb{R}_+)^n$ one has

$$\phi((1-\lambda)x +_p \lambda y) \geq \phi((1-\lambda)x + \lambda y) \geq M_\alpha^\lambda(\phi(x), \phi(y)).$$

Hence,

$$h((1-\lambda)x +_p \lambda y) \geq M_\alpha^\lambda(f(x), g(y)).$$

Thus we may apply the result mentioned at the beginning of the proof to obtain that

$$\int_{(\mathbb{R}_+)^n} h(x) dx \geq M_\gamma^\lambda \left(\int_{(\mathbb{R}_+)^n} f(x) dx, \int_{(\mathbb{R}_+)^n} g(x) dx \right),$$

where $\gamma = (\sum_{i=1}^n p_i^{-1} + \alpha^{-1})^{-1}$. In other words, one has

$$\mu(((1-\lambda) \cdot A +_p \lambda \cdot B) \cap (\mathbb{R}_+)^n) \geq M_\gamma^\lambda(\mu(A \cap (\mathbb{R}_+)^n), \mu(B \cap (\mathbb{R}_+)^n)).$$

Since the sets $(1-\lambda) \cdot A +_p \lambda \cdot B$, A and B are unconditional, it follows that

$$\mu((1-\lambda) \cdot A +_p \lambda \cdot B) \geq M_\gamma^\lambda(\mu(A), \mu(B)). \quad \square$$

Remark. One may similarly define the L^p -Minkowski combination

$$\lambda_1 \cdot A_1 +_p \cdots +_p \lambda_m \cdot A_m$$

for m convex bodies $A_1, \dots, A_m \subset \mathbb{R}^n$, $m \geq 2$, where $\lambda_1, \dots, \lambda_m \in [0, 1]$ are such that $\sum_{i=1}^m \lambda_i = 1$, by extending the definition of the p -mean M_p^λ to m nonnegative numbers. By induction, one has under the same assumptions of Theorem 1.1 that

$$(5) \quad \mu(\lambda_1 \cdot A_1 +_p \cdots +_p \lambda_m \cdot A_m) \geq M_\gamma^\lambda(\mu(A_1), \dots, \mu(A_m)),$$

where $\gamma = (\sum_{i=1}^n p_i^{-1} + \alpha^{-1})^{-1}$. Indeed, let $m \geq 2$ and let us assume that inequality (5) holds. Notice that

$$\lambda_1 \cdot A_1 +_p \cdots +_p \lambda_m \cdot A_m +_p \lambda_{m+1} \cdot A_{m+1} = \left(\sum_{i=1}^m \lambda_i \right) \cdot \tilde{A} +_p \lambda_{m+1} \cdot A_{m+1},$$

where

$$\tilde{A} := \left(\frac{\lambda_1}{\sum_{i=1}^m \lambda_i} \cdot A_1 +_p \cdots +_p \frac{\lambda_m}{\sum_{i=1}^m \lambda_i} \cdot A_m \right).$$

Thus,

$$\begin{aligned} \mu\left(\left(\sum_{i=1}^m \lambda_i\right) \cdot \tilde{A} +_p \lambda_{m+1} \cdot A_{m+1}\right) &\geq \left(\left(\sum_{i=1}^m \lambda_i\right) \mu(\tilde{A})^\gamma + \lambda_{m+1} \mu(A_{m+1})^\gamma\right)^{\frac{1}{\gamma}} \\ &\geq \left(\sum_{i=1}^{m+1} \lambda_i \mu(A_i)^\gamma\right)^{\frac{1}{\gamma}}. \end{aligned}$$

Consequences. The following result compares the L^p -Minkowski combinations \oplus_p and $+_p$.

Lemma 2.2. *Let $p \in [0, 1]$ and set $\mathbf{p} = (p, \dots, p) \in [0, 1]^n$. For every unconditional convex body A, B in \mathbb{R}^n and for every $\lambda \in [0, 1]$, one has*

$$(1 - \lambda) \cdot A \oplus_p \lambda \cdot B \supset (1 - \lambda) \cdot A +_p \lambda \cdot B.$$

Proof. The case $p = 0$ is proved in [Saroglou 2015]. Let $p \neq 0$. Since the sets $(1 - \lambda) \cdot A \oplus_p \lambda \cdot B$ and $(1 - \lambda) \cdot A +_p \lambda \cdot B$ are unconditional, it is sufficient to prove that

$$((1 - \lambda) \cdot A \oplus_p \lambda \cdot B) \cap (\mathbb{R}_+)^n \supset ((1 - \lambda) \cdot A +_p \lambda \cdot B) \cap (\mathbb{R}_+)^n.$$

Let $u \in S^{n-1} \cap (\mathbb{R}_+)^n$ and let $x \in ((1 - \lambda) \cdot A +_p \lambda \cdot B) \cap (\mathbb{R}_+)^n$. One has

$$\begin{aligned} \langle x, u \rangle &= \sum_{i=1}^n ((1 - \lambda)a_i^p + \lambda b_i^p)^{\frac{1}{p}} u_i = \sum_{i=1}^n ((1 - \lambda)(a_i u_i)^p + \lambda (b_i u_i)^p)^{\frac{1}{p}} \\ &= \|(1 - \lambda)X + \lambda Y\|_{\frac{1}{p}}, \end{aligned}$$

where $X = ((a_1 u_1)^p, \dots, (a_n u_n)^p)$ and $Y = ((b_1 u_1)^p, \dots, (b_n u_n)^p)$. Notice that $\|X\|_{\frac{1}{p}} \leq h_A(u)^p$, $\|Y\|_{\frac{1}{p}} \leq h_B(u)^p$ and that $\|\cdot\|_{\frac{1}{p}}$ is a norm. It follows that

$$\langle x, u \rangle \leq ((1 - \lambda)\|X\|_{\frac{1}{p}} + \lambda\|Y\|_{\frac{1}{p}})^{\frac{1}{p}} \leq ((1 - \lambda)h_A(u)^p + \lambda h_B(u)^p)^{\frac{1}{p}}.$$

Hence, $x \in ((1 - \lambda) \cdot A \oplus_p \lambda \cdot B) \cap (\mathbb{R}_+)^n$. □

From Lemma 2.2 and Theorem 1.1, one obtains the following result:

Corollary 2.3. *Let $p \in [0, 1]$. Let μ be an unconditional measure in \mathbb{R}^n that has an α -concave density function, with $\alpha \geq -\frac{p}{n}$. Then, for every unconditional convex body A, B in \mathbb{R}^n and for every $\lambda \in [0, 1]$,*

$$(6) \quad \mu((1 - \lambda) \cdot A \oplus_p \lambda \cdot B) \geq M_\gamma^\lambda(\mu(A), \mu(B)),$$

where $\gamma = \left(\frac{n}{p} + \frac{1}{\alpha}\right)^{-1}$.

In Corollary 2.3, if α or p is equal to 0, then γ is defined by continuity and is equal to 0.

Remarks. (1) By taking $\alpha = 0$ in Corollary 2.3 (corresponding to log-concave measures), one obtains

$$\mu((1 - \lambda) \cdot A \oplus_0 \lambda \cdot B) \geq \mu(A)^{1-\lambda} \mu(B)^\lambda.$$

(2) By taking $\alpha = +\infty$ in Corollary 2.3 (corresponding to $\frac{1}{n}$ -concave measures), one obtains that, for every $p \in [0, 1]$,

$$\mu((1 - \lambda) \cdot A \oplus_p \lambda \cdot B)^{\frac{p}{n}} \geq (1 - \lambda)\mu(A)^{\frac{p}{n}} + \lambda\mu(B)^{\frac{p}{n}}.$$

Equivalently, for every $p \in [0, 1]$, for every unconditional convex body A, B in \mathbb{R}^n and for every unconditional convex set $K \subset \mathbb{R}^n$,

$$|((1 - \lambda) \cdot A \oplus_p \lambda \cdot B) \cap K|^{\frac{p}{n}} \geq (1 - \lambda)|A \cap K|^{\frac{p}{n}} + \lambda|B \cap K|^{\frac{p}{n}}.$$

Let us recall that the function $t \mapsto \mu(e^t A)$ is log-concave on \mathbb{R} for every unconditional log-concave measure μ and every unconditional convex body A in \mathbb{R}^n (see [Cordero-Erausquin et al. 2004]). By adapting the argument of [Marsiglietti 2015], Proof of Proposition 3.1 (see Proof of Corollary 2.5), it follows that the function $t \mapsto \mu(t^{\frac{1}{p}} A)$ is $\frac{p}{n}$ -concave on \mathbb{R}_+ , for every $p \in (0, 1]$, for every unconditional s -concave measure μ , with $s \geq 0$, and for every unconditional convex body A in \mathbb{R}^n . However, no concavity properties are known for the function $t \mapsto \mu(e^t A)$ when μ is an s -concave measure with $s < 0$. Instead, for these measures we prove concavity properties of the function $t \mapsto \mu(t^{\frac{1}{p}} A)$.

Proposition 2.4. *Let $p \in (0, 1]$ and $\alpha \in [-\frac{p}{n}, 0)$, let μ be an unconditional measure that has an α -concave density function, and let A be an unconditional convex body in \mathbb{R}^n . Then the function $t \mapsto \mu(t^{\frac{1}{p}} A)$ is $(\frac{n}{p} + \frac{1}{\alpha})^{-1}$ -concave on \mathbb{R}_+ .*

Proof. Let $t_1, t_2 \in \mathbb{R}_+$. By applying Corollary 2.3 to the sets $t_1^{\frac{1}{p}} A$ and $t_2^{\frac{1}{p}} A$, one obtains

$$\begin{aligned} \mu(((1 - \lambda)t_1 + \lambda t_2)^{\frac{1}{p}} A) &= \mu((1 - \lambda) \cdot t_1^{\frac{1}{p}} A \oplus_p \lambda \cdot t_2^{\frac{1}{p}} A) \\ &\geq M_\gamma^\lambda(\mu(t_1^{\frac{1}{p}} A), \mu(t_2^{\frac{1}{p}} A)), \end{aligned}$$

where $\gamma = (\frac{n}{p} + \frac{1}{\alpha})^{-1}$. Hence the function $t \mapsto \mu(t^{\frac{1}{p}} A)$ is γ -concave on \mathbb{R}_+ . \square

As a consequence, we derive concavity properties for the function $t \mapsto \mu(tA)$.

Corollary 2.5. *Let $p \in (0, 1]$, let μ be an unconditional measure that has an α -concave density function, with $\alpha \in [-\frac{p}{n}, 0)$, and let A be an unconditional convex body in \mathbb{R}^n . Then the function $t \mapsto \mu(tA)$ is $(\frac{1-p}{n} + \gamma)$ -concave on \mathbb{R}_+ , where $\gamma = (\frac{n}{p} + \frac{1}{\alpha})^{-1}$.*

Proof. We adapt [Marsiglietti 2015], Proof of Proposition 3.1. Let us denote by ϕ the density function of the measure μ and let us denote by F the function $t \mapsto \mu(tA)$. From Proposition 2.4, the function $t \mapsto F(t^{\frac{1}{p}})$ is γ -concave, hence the right derivative of F , denoted by F'_+ , exists everywhere and the function $t \mapsto \frac{1}{p}t^{\frac{1}{p}-1}F'_+(t^{\frac{1}{p}})F(t^{\frac{1}{p}})^{\gamma-1}$ is nonincreasing. Notice that

$$F(t) = t^n \int_A \phi(tx) dx$$

and that $t \mapsto \phi(tx)$ is nonincreasing; thus the function $t \mapsto \frac{1}{t^{1-p}}F(t)^{\frac{1-p}{n}}$ is nonincreasing. Since

$$F'_+(t)F(t)^{\frac{1-p}{n}+\gamma-1} = t^{1-p}F'_+(t)F(t)^{\gamma-1} \cdot \frac{1}{t^{1-p}}F(t)^{\frac{1-p}{n}},$$

it follows that $F'_+(t)F(t)^{\frac{1-p}{n}+\gamma-1}$ is nonincreasing as the product of two nonnegative nonincreasing functions. Hence F is $(\frac{1-p}{n} + \gamma)$ -concave. \square

Remark. For every s -concave measure μ and for every convex subset $A \subset \mathbb{R}^n$, the function $t \mapsto \mu(tA)$ is s -concave. Hence Corollary 2.5 is of value only if $\frac{1-p}{n} + \gamma \geq \alpha/(1 + \alpha n)$ (see relation (2)). Notice that this condition is satisfied if $\alpha \geq -p/(n(1 + p))$. We thus obtain:

Corollary 2.6. *Let $p \in (0, 1]$, let μ be an unconditional measure that has an α -concave density function, with $-p/(n(1 + p)) \leq \alpha < 0$, and let K be an unconditional convex body in \mathbb{R}^n . Then, for all subsets $A, B \in \{\mu K : \mu > 0\}$ and all $\lambda \in [0, 1]$, one has*

$$\mu((1 - \lambda)A + \lambda B) \geq M_{\frac{1-p}{n}+\gamma}^\lambda(\mu(A), \mu(B)),$$

where $\gamma = (\frac{n}{p} + \frac{1}{\alpha})^{-1}$.

In [Marsiglietti 2015] we investigated improvements of concavity properties of convex measures under additional assumptions, such as symmetries. Corollary 2.6 follows the same path and completes the results found there.

We conclude this section with a remark on the question of improving the concavity properties of convex measures.

Remark. Let μ be a Borel measure that has a density function with respect to Lebesgue measure in \mathbb{R}^n . One may write the density function of μ in the form e^{-V} , where $V : \mathbb{R}^n \rightarrow \mathbb{R}$ is a measurable function. Let us assume that V is C^2 . Let $\gamma > 0$. The function e^{-V} is γ -concave if $\text{Hess}(\gamma e^{-\gamma V})$, the Hessian of $\gamma e^{-\gamma V}$, is nonpositive (in the sense of symmetric matrices). One has

$$\text{Hess}(\gamma e^{-\gamma V}) = -\gamma^2 \nabla \cdot (\nabla V e^{-\gamma V}) = \gamma^2 e^{-\gamma V} (\gamma \nabla V \otimes \nabla V - \text{Hess } V),$$

where

$$\nabla V \otimes \nabla V = \left(\frac{\partial V}{\partial x_i} \frac{\partial V}{\partial x_j} \right)_{1 \leq i, j \leq n}.$$

It follows that the matrix $\text{Hess}(\gamma e^{-\gamma V})$ is nonpositive if and only if the matrix $\gamma \nabla V \otimes \nabla V - \text{Hess } V$ is nonpositive.

Let us apply this remark to the Gaussian measure

$$d\gamma_n(x) = \frac{1}{(2\pi)^{\frac{n}{2}}} e^{-\frac{|x|^2}{2}} dx, \quad x \in \mathbb{R}^n.$$

Here $V(x) = \frac{|x|^2}{2} + c_n$, where $c_n = \frac{n}{2} \log(2\pi)$. Thus $\nabla V \otimes \nabla V = (x_i x_j)_{1 \leq i, j \leq n}$ and $\text{Hess } V = \text{Id}$, the identity matrix. The eigenvalues of $\gamma \nabla V \otimes \nabla V - \text{Hess } V$ are -1 (with multiplicity $(n-1)$) and $\gamma|x|^2 - 1$. Hence, if $\gamma|x|^2 - 1 \leq 0$, then $\gamma \nabla V \otimes \nabla V - \text{Hess } V$ is nonpositive. One deduces that, for every $\gamma > 0$, for all compact sets $A, B \subset \frac{1}{\sqrt{\gamma}} B_2^n$ and for every $\lambda \in [0, 1]$, one has

$$(7) \quad \gamma_n((1-\lambda)A + \lambda B) \geq M_{\frac{\lambda}{1+\gamma n}}^\lambda(\gamma_n(A), \gamma_n(B)),$$

where B_2^n denotes the Euclidean closed unit ball in \mathbb{R}^n .

Since the Gaussian measure is a log-concave measure, inequality (7) is an improvement of the concavity of the Gaussian measure when restricted to compact sets $A, B \subset \frac{1}{\sqrt{\gamma}} B_2^n$.

3. Equivalence between (B)-conjecture-type problems

The next proposition reduces the proof of the (B)-conjecture for all uniform measures in \mathbb{R}^n , for every $n \in \mathbb{N}^*$, to proving the (B)-conjecture for all symmetric log-concave measures in \mathbb{R}^n , for every $n \in \mathbb{N}^*$. This completes recent work by Saroglou [2014; 2015].

We will say that a measure μ satisfies the (B)-property if the function $t \mapsto \mu(e^t A)$ is log-concave on \mathbb{R} for every symmetric convex set $A \subset \mathbb{R}^n$.

Proposition 3.1. *If every symmetric uniform measure in \mathbb{R}^n , for every $n \in \mathbb{N}^*$, satisfies the (B)-property, then every symmetric log-concave measure in \mathbb{R}^n , for every $n \in \mathbb{N}^*$, satisfies the (B)-property.*

Proof. The proof is inspired by [Artstein-Avidan et al. 2004, beginning of Section 3].

Step 1: Stability under orthogonal projection. Let us show that the (B)-property is stable under orthogonal projection onto an arbitrary subspace.

Let F be a k -dimensional subspace of \mathbb{R}^n . Let us define, for every compactly supported measure μ in \mathbb{R}^n and every measurable subset $A \subset F$,

$$\Pi_F \mu(A) := \mu(\Pi_F^{-1}(A)),$$

where Π_F denotes the orthogonal projection onto F and

$$\Pi_F^{-1}(A) := \{x \in \mathbb{R}^n : \Pi_F(x) \in A\}.$$

We have $\Pi_F^{-1}(e^t A) = e^t(A \times F^\perp)$, where F^\perp denotes the orthogonal complement of F . Hence if μ satisfies the (B)-property, so does $\Pi_F \mu$.

Step 2: Approximation of log-concave measures. Let us show that for every compactly supported log-concave measure μ in \mathbb{R}^n there exists a sequence $(K_p)_{p \in \mathbb{N}^*}$ of convex subsets of \mathbb{R}^{n+p} such that $\lim_{p \rightarrow +\infty} \Pi_{\mathbb{R}^n} \mu_{K_p} = \mu$ in the sense that the density function of μ is the pointwise limit of the density functions of $(\mu_{K_p})_{p \in \mathbb{N}^*}$, where μ_{K_p} denotes the uniform measure on K_p (up to a constant).

Let μ be a compactly supported log-concave measure in \mathbb{R}^n with density function $f = e^{-V}$, where $V : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ is a convex function. To simplify notation, define

$$(8) \quad W(x) = \left(1 - \frac{V(x)}{p}\right)_+,$$

where $a_+ = \max(a, 0)$ for every $a \in \mathbb{R}$. Notice that $e^{-V(x)} = \lim_{p \rightarrow +\infty} W(x)^p$ for every $x \in \mathbb{R}^n$. Let us define for every $p \in \mathbb{N}^*$

$$K_p = \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^p : |y| \leq W(x)\}.$$

One has, for every $x \in \mathbb{R}^n$,

$$W(x)^p = \int_0^{W(x)} pr^{p-1} dr = p \int_0^{+\infty} 1_{[0, W(x)]}(r)r^{p-1} dr = \frac{1}{v_p} \int_{\mathbb{R}^p} 1_{K_p}(x, y) dy.$$

The last equality follows from an integration in polar coordinates, where v_p denotes the volume of the Euclidean closed unit ball in \mathbb{R}^p . By denoting μ_{K_p} the measure in \mathbb{R}^{n+p} with density function

$$\frac{1}{v_p} 1_{K_p}(x, y), \quad (x, y) \in \mathbb{R}^n \times \mathbb{R}^p,$$

it follows that, for every $p \in \mathbb{N}^*$, the measure $\Pi_{\mathbb{R}^n} \mu_{K_p}$ has density function $W(x)^p, x \in \mathbb{R}^n$. We conclude that $\lim_{p \rightarrow +\infty} \Pi_{\mathbb{R}^n} \mu_{K_p} = \mu$.

Step 3: Conclusion. Let $n \in \mathbb{N}^*$ and let μ be a symmetric log-concave measure in \mathbb{R}^n . By approximation, one can assume that μ is compactly supported. Since μ is symmetric, the sequence $(K_p)_{p \in \mathbb{N}^*}$ defined in Step 2 is a sequence of symmetric convex subsets of \mathbb{R}^{n+p} . If we assume that the (B)-property holds for all uniform measures in \mathbb{R}^m , for every $m \in \mathbb{N}^*$, then, for every $p \in \mathbb{N}^*$, μ_{K_p} satisfies the (B)-property. It follows from Step 1 that, for every $p \in \mathbb{N}^*$, $\Pi_{\mathbb{R}^n} \mu_{K_p}$ satisfies the (B)-property. Since $\lim_{p \rightarrow +\infty} \Pi_{\mathbb{R}^n} \mu_{K_p} = \mu$ (see Step 2) and since a pointwise

limit of log-concave functions is log-concave, we conclude that μ satisfies the (B)-property. \square

Similarly, let us now prove that the functional form of the (B)-conjecture (Conjecture 1.4) is equivalent to the classical (B)-conjecture (Conjecture 1.3).

Proposition 3.2. *One has equivalence between the following properties:*

- (1) *For every $n \in \mathbb{N}^*$, for every symmetric log-concave measure μ in \mathbb{R}^n and for every symmetric convex subset A of \mathbb{R}^n , the function $t \mapsto \mu(e^t A)$ is log-concave on \mathbb{R} .*
- (2) *For every $n \in \mathbb{N}^*$ and for all even log-concave functions $f, g : \mathbb{R}^n \rightarrow \mathbb{R}_+$, the function $t \mapsto \int_{\mathbb{R}^n} f(e^{-t}x)g(x) dx$ is log-concave on \mathbb{R} .*

Proof. (2) \implies (1) This is clear by taking f to be 1_A , the indicator function of a symmetric convex set A , and by taking g to be the density function of a log-concave measure μ .

(1) \implies (2) Let $f, g : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be even log-concave functions. By approximation, one may assume that f and g are compactly supported. Let us write $g = e^{-V}$, where $V : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ is an even convex function. One has

$$G(t) := \int_{\mathbb{R}^n} f(e^{-t}x)e^{-V(x)} dx = \lim_{p \rightarrow +\infty} \int_{\mathbb{R}^n} f(e^{-t}x)W(x)^p dx,$$

where $W(x)$ is as in (8). Let us denote, for $t \in \mathbb{R}$,

$$G_p(t) = \int_{\mathbb{R}^n} f(e^{-t}x)W(x)^p dx.$$

We have seen in the proof of Proposition 3.1 that

$$W(x)^p = \frac{1}{v_p} \int_{\mathbb{R}^p} 1_{K_p}(x, y) dy,$$

where $K_p := \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^p : |y| \leq W(x)\}$ and where v_p denotes the volume of the Euclidean closed unit ball in \mathbb{R}^p . Hence,

$$G_p(t) = \frac{1}{v_p} \int_{K_p} f(e^{-t}x)1_{\mathbb{R}^p}(y) dx dy.$$

Notice that K_p is a symmetric convex subset of \mathbb{R}^{n+p} . The change of variable $\tilde{x} = e^{-t}x$ and $\tilde{y} = e^{-t}y$ leads to

$$G_p(t) = \frac{e^{t(n+p)}}{v_p} \mu_p(e^{-t}K_p),$$

where μ_p is the measure with density function

$$h(x, y) = f(x)1_{\mathbb{R}^p}(y), \quad (x, y) \in \mathbb{R}^n \times \mathbb{R}^p.$$

Since a pointwise limit of log-concave functions is log-concave, we conclude that the function G is log-concave on \mathbb{R} as the pointwise limit of the log-concave functions G_p , $p \in \mathbb{N}^*$. \square

Recall that the (B)-conjecture holds true for the Gaussian measure and for the unconditional case (see [Cordero-Erausquin et al. 2004]). From the techniques of the proof of Proposition 3.2, it follows that Conjecture 1.4 holds true if one function is the density function of the Gaussian measure or if both functions are unconditional.

References

- [Artstein-Avidan et al. 2004] S. Artstein-Avidan, B. Klartag, and V. Milman, “The Santaló point of a function, and a functional form of the Santaló inequality”, *Mathematika* **51**:1-2 (2004), 33–48. MR 2007a:52008 Zbl 1121.52021
- [Bobkov et al. 2014] S. G. Bobkov, A. Colesanti, and I. Fragalà, “Quermassintegrals of quasi-concave functions and generalized Prékopa–Leindler inequalities”, *Manuscripta Math.* **143**:1-2 (2014), 131–169. MR 3147446 Zbl 1290.26019
- [Borell 1974] C. Borell, “Convex measures on locally convex spaces”, *Ark. Mat.* **12** (1974), 239–252. MR 52 #9311 Zbl 0297.60004
- [Borell 1975] C. Borell, “Convex set functions in d -space”, *Period. Math. Hungar.* **6**:2 (1975), 111–136. MR 53 #8359 Zbl 0307.28009
- [Böröczky et al. 2012] K. J. Böröczky, E. Lutwak, D. Yang, and G. Zhang, “The log–Brunn–Minkowski inequality”, *Adv. Math.* **231**:3-4 (2012), 1974–1997. MR 2964630 Zbl 1258.52005
- [Caglar and Werner 2014] U. Caglar and E. M. Werner, “Divergence for s -concave and log concave functions”, *Adv. Math.* **257** (2014), 219–247. MR 3187648 Zbl 1310.26016
- [Caglar et al. 2015] U. Caglar, M. Fradelizi, O. Guédon, J. Lehec, C. Schütt, and E. M. Werner, “Functional versions of L_p -affine surface area and entropy inequalities”, *International Mathematics Research Notices* (2015), 1–28.
- [Cordero-Erausquin et al. 2004] D. Cordero-Erausquin, M. Fradelizi, and B. Maurey, “The (B) conjecture for the Gaussian measure of dilates of symmetric convex sets and related problems”, *J. Funct. Anal.* **214**:2 (2004), 410–427. MR 2005g:60064 Zbl 1073.60042
- [Dancs and Uhrin 1980] S. Dancs and B. Uhrin, “On a class of integral inequalities and their measure-theoretic consequences”, *J. Math. Anal. Appl.* **74**:2 (1980), 388–400. MR 81g:26009 Zbl 0442.26011
- [Dubuc 1977] S. Dubuc, “Critères de convexité et inégalités intégrales”, *Ann. Inst. Fourier (Grenoble)* **27**:1 (1977), 135–165. MR 56 #3210 Zbl 0331.26008
- [Firey 1961] W. J. Firey, “Mean cross-section measures of harmonic means of convex bodies”, *Pacific J. Math.* **11** (1961), 1263–1266. MR 25 #3427 Zbl 0122.41101
- [Firey 1962] W. J. Firey, “ p -means of convex bodies”, *Math. Scand.* **10** (1962), 17–24. MR 25 #4416 Zbl 0188.27303
- [Firey 1964] W. J. Firey, “Some applications of means of convex bodies”, *Pacific J. Math.* **14** (1964), 53–60. MR 28 #4428 Zbl 0126.38405
- [Gardner 2002] R. J. Gardner, “The Brunn–Minkowski inequality”, *Bull. Amer. Math. Soc. (N.S.)* **39**:3 (2002), 355–405. MR 2003f:26035 Zbl 1019.26008

- [Gardner et al. 2014] R. J. Gardner, D. Hug, and W. Weil, “The Orlicz–Brunn–Minkowski theory: A general framework, additions, and inequalities”, *J. Differential Geom.* **97**:3 (2014), 427–476. MR 3263511 Zbl 1303.52002
- [Henstock and Macbeath 1953] R. Henstock and A. M. Macbeath, “On the measure of sum-sets, I: The theorems of Brunn, Minkowski, and Lusternik”, *Proc. London Math. Soc.* (3) **3** (1953), 182–194. MR 15,109g Zbl 0052.18302
- [Latała 2002] R. Latała, “On some inequalities for Gaussian measures”, pp. 813–822 in *Proceedings of the International Congress of Mathematicians, II* (Beijing, 2002), edited by T. Li, Higher Ed. Press, Beijing, 2002. MR 2004b:60055 Zbl 1015.60011
- [Lutwak 1993] E. Lutwak, “The Brunn–Minkowski–Firey theory, I: Mixed volumes and the Minkowski problem”, *J. Differential Geom.* **38**:1 (1993), 131–150. MR 94g:52008 Zbl 0788.52007
- [Lutwak 1996] E. Lutwak, “The Brunn–Minkowski–Firey theory, II: Affine and geominimal surface areas”, *Adv. Math.* **118**:2 (1996), 244–294. MR 97f:52014 Zbl 0853.52005
- [Marsiglietti 2015] A. Marsiglietti, “On the improvement of concavity of convex measures”, *Proceedings of the American Mathematical Society* (2015), 1–12.
- [Saroglou 2014] C. Saroglou, “More on logarithmic sums of convex bodies”, preprint, 2014. arXiv 1409.4346
- [Saroglou 2015] C. Saroglou, “Remarks on the conjectured log-Brunn–Minkowski inequality”, *Geom. Dedicata* **177**:1 (2015), 353–365. MR 3370038
- [Schneider 1993] R. Schneider, *Convex bodies: The Brunn–Minkowski theory*, Encyclopedia of Mathematics and its Applications **44**, Cambridge Univ. Press, 1993. MR 94d:52007 Zbl 0798.52001
- [Uhrin 1994] B. Uhrin, “Curvilinear extensions of the Brunn–Minkowski–Lusternik inequality”, *Adv. Math.* **109**:2 (1994), 288–312. MR 95j:52017 Zbl 0847.52007

Received November 25, 2014. Revised March 13, 2015.

ARNAUD MARSIGLIETTI
INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS
UNIVERSITY OF MINNESOTA
207 CHURCH STREET SE
306 LIND HALL
MINNEAPOLIS, MN 55455
UNITED STATES
arnaud.marsiglietti@ima.umn.edu

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 277 No. 1 September 2015

Real positivity and approximate identities in Banach algebras	1
DAVID P. BLECHER and NARUTAKA OZAWA	
On shrinking gradient Ricci solitons with nonnegative sectional curvature	61
MINGLIANG CAI	
From quasimodes to resonances: exponentially decaying perturbations	77
ORAN GANNOT	
A general simple relative trace formula	99
JAYCE R. GETZ and HEEKYOUNG HAHN	
Chern-Simons functions on toric Calabi-Yau threefolds and Donaldson-Thomas theory	119
ZHENG HUA	
On the flag curvature of a class of Finsler metrics produced by the navigation problem	149
LIBING HUANG and XIAOHUAN MO	
Angular distribution of diameters for spheres and rays for planes	169
NOBUHIRO INNAMI and YUYA UNEME	
A note on an L^p -Brunn–Minkowski inequality for convex measures in the unconditional case	187
ARNAUD MARSIGLIETTI	
Structure of seeds in generalized cluster algebras	201
TOMOKI NAKANISHI	
Inequalities of Alexandrov–Fenchel type for convex hypersurfaces in hyperbolic space and in the sphere	219
YONG WEI and CHANGWEI XIONG	
Upper bounds of root discriminant lower bounds	241
SIMAN WONG	



0030-8730(201509)277:1;1-V