

*Pacific
Journal of
Mathematics*

**STRUCTURE OF SEEDS IN GENERALIZED
CLUSTER ALGEBRAS**

TOMOKI NAKANISHI

Volume 277 No. 1

September 2015

STRUCTURE OF SEEDS IN GENERALIZED CLUSTER ALGEBRAS

TOMOKI NAKANISHI

We study generalized cluster algebras, introduced by Chekhov and Shapiro. When the coefficients satisfy the normalization and quasireciprocity conditions, one can naturally extend the structure theory of seeds in the ordinary cluster algebras by Fomin and Zelevinsky to generalized cluster algebras. As the main result, we obtain formulas expressing cluster variables and coefficients in terms of c -vectors, g -vectors, and F -polynomials.

1. Introduction

Chekhov and Shapiro [2014] introduced *generalized cluster algebras*, which naturally generalize the ordinary cluster algebras by Fomin and Zelevinsky [2002]. In generalized cluster algebras, the celebrated *binomial* exchange relation for cluster variables of ordinary cluster algebras

$$(1-1) \quad \begin{aligned} x'_k x_k &= p_k^- \prod_{j=1}^n x_j^{[-b_{jk}]_+} + p_k^+ \prod_{j=1}^n x_j^{[b_{jk}]_+} \\ &= \left(\prod_{j=1}^n x_j^{[-b_{jk}]_+} \right) (p_k^- + p_k^+ w_k), \quad w_k = \prod_{j=1}^n x_j^{b_{jk}}, \end{aligned}$$

is replaced by the *polynomial* one of arbitrary degree $d_k \geq 1$,

$$(1-2) \quad x'_k x_k = \left(\prod_{j=1}^n x_j^{[-\beta_{jk}]_+} \right)^{d_k} \sum_{s=0}^{d_k} p_{k,s} w_k^s, \quad w_k = \prod_{j=1}^n x_j^{\beta_{jk}},$$

where $\beta_{jk} = b_{jk}/d_k$ are assumed to be integers and the coefficients $p_{k,s}$ should also be mutated appropriately. This generalization is expected to be natural, since it originates in the transformations preserving the associated Poisson bracket [Gekhtman et al. 2005]. In fact, it was shown in [Chekhov and Shapiro 2014] that the generalized cluster algebras have the *Laurent property*, which is regarded as the most characteristic feature of the ordinary cluster algebras. It was also shown in

MSC2010: 13F60.

Keywords: cluster algebra.

the same paper that the *finite-type classification* of the generalized cluster algebras reduces to the one for the ordinary case. These results already imply that, despite the apparent complexity of their exchange relations (1-2), generalized cluster algebras may be well controlled like the ordinary ones. See also [Rupel 2013] for the result on greedy bases in rank 2 generalized cluster algebras.

Besides the above cluster-algebra-theoretic interest, the generalized cluster algebra structure naturally appears for the Teichmüller spaces of Riemann surfaces with orbifold points [Chekhov and Shapiro 2014]. More recently, it also appears in representation theory of quantum affine algebras [Gleitz 2014] and also in the study of WKB analysis [Iwaki and Nakanishi 2014]. In view of these developments, and also for potentially more versatility of polynomial exchange relations than the binomial one, it is not only natural but also necessary to develop a structure theory of seeds in generalized cluster algebras which is parallel to the one for the ordinary cluster algebras by [Fomin and Zelevinsky 2007]. The core notion of the theory of that paper is a cluster pattern with *principal coefficients*, from which other important notions such as *c-vectors*, *g-vectors*, and *F-polynomials* are also induced. Then, the main result of [Fomin and Zelevinsky 2007] is the formulas expressing cluster variables and coefficients in terms of *c-vectors*, *g-vectors*, and *F-polynomials*. These formulas are especially important in view of the categorification of cluster algebras by (generalized) cluster categories (see [Plamondon 2011] and references therein).

The purpose of this paper is to provide results parallel to the above ones for generalized cluster algebras. To be more precise, we consider a class of generalized cluster algebras whose coefficients satisfy the *normalization condition* and what we call the *quasireciprocity condition*. For this class of generalized cluster algebras, we introduce the notions of a cluster pattern with principal coefficients, *c-vectors*, *g-vectors*, and *F-polynomials*. Then, as a main result, we obtain the formulas expressing cluster variables and coefficients in terms of *c-vectors*, *g-vectors*, and *F-polynomials*, which are parallel to the ones in [Fomin and Zelevinsky 2007]. To summarize, *generalized cluster algebras preserve essentially every feature of the ordinary ones*, and this is the main message of the paper.

2. Generalized cluster algebras

In this section we recall basic notions of generalized cluster algebras following [Chekhov and Shapiro 2014]. However, we slightly modify the setting of Chekhov and Shapiro to match the setting of (ordinary) cluster algebras in [Fomin and Zelevinsky 2007].

2A. Generalized seed mutations. Throughout the paper we always assume that any matrix is an *integer* matrix.

Recall that a matrix $B = (b_{ij})_{i,j=1}^n$ is said to be *skew-symmetrizable* if there is an n -tuple of positive integers $\mathbf{d} = (d_1, \dots, d_n)$ such that $d_i b_{ij} = -d_j b_{ji}$.

We start by fixing a semifield \mathbb{P} , whose addition is denoted by \oplus . Let $\mathbb{Z}\mathbb{P}$ be the group ring of \mathbb{P} , and let $\mathbb{Q}\mathbb{P}$ be the field of fractions of $\mathbb{Z}\mathbb{P}$. Let w_1, \dots, w_n be any algebraic independent variables, and let $\mathcal{F} = \mathbb{Q}\mathbb{P}(w)$ be the field of rational functions in $w = (w_1, \dots, w_n)$ with coefficients in $\mathbb{Q}\mathbb{P}$.

The following definition is the usual one [Fomin and Zelevinsky 2007].

Definition 2.1. A (labeled) seed in \mathbb{P} is a triplet $(\mathbf{x}, \mathbf{y}, B)$ such that

- B is a skew-symmetrizable matrix, called an *exchange matrix*,
- $\mathbf{x} = (x_1, \dots, x_n)$ is an n -tuple of elements in \mathcal{F} , called *cluster variables* or *x-variables*,
- $\mathbf{y} = (y_1, \dots, y_n)$ is an n -tuple of elements in \mathbb{P} , called *coefficients* or *y-variables*.

Next we introduce a pair (\mathbf{d}, \mathbf{z}) of data for generalized seed mutations. Firstly, $\mathbf{d} = (d_1, \dots, d_n)$ is an n -tuple of positive integers, and we call these integers the *mutation degrees*. We stress that we do *not* impose the skew-symmetric condition $d_i b_{ij} = -d_j b_{ji}$. Secondly, \mathbf{z} is a family of elements in \mathbb{P} ,

$$(2-1) \quad \mathbf{z} = (z_{i,s})_{i=1,\dots,n; s=1,\dots,d_i-1},$$

satisfying the *reciprocity condition*

$$(2-2) \quad z_{i,s} = z_{i,d_i-s} \quad (s = 1, \dots, d_i - 1).$$

We call them the *frozen coefficients*, since they are not “mutated”, or simply the *z-variables*. We also set

$$(2-3) \quad z_{i,0} = z_{i,d_i} = 1.$$

For $\mathbf{d} = (1, \dots, 1)$, \mathbf{z} is empty, and it reduces to the ordinary case. (Here and below, “ordinary” means the case of ordinary cluster algebras.)

Definition 2.2. Let (\mathbf{d}, \mathbf{z}) be given as above. For any seed $(\mathbf{x}, \mathbf{y}, B)$ in \mathbb{P} and $k = 1, \dots, n$, the (\mathbf{d}, \mathbf{z}) -*mutation of $(\mathbf{x}, \mathbf{y}, B)$ at k* is another seed $(\mathbf{x}', \mathbf{y}', B')$ = $\mu_k(\mathbf{x}, \mathbf{y}, B)$ in \mathbb{P} defined by the following rule:

$$(2-4) \quad b'_{ij} = \begin{cases} -b_{ij} & \text{if } i = k \text{ or } j = k, \\ b_{ij} + d_k([-b_{ik}]_+ b_{kj} + b_{ik}[b_{kj}]_+) & \text{if } i, j \neq k, \end{cases}$$

$$(2-5) \quad y'_i = \begin{cases} y_k^{-1} & \text{if } i = k, \\ y_i (y_k^{[\varepsilon b_{ki}]_+})^{d_k} \left(\bigoplus_{s=0}^{d_k} z_{k,s} y_k^{\varepsilon s} \right)^{-b_{ki}} & \text{if } i \neq k, \end{cases}$$

$$(2-6) \quad x'_i = \begin{cases} x_k^{-1} \left(\prod_{j=1}^n x_j^{[-\varepsilon b_{jk}]_+} \right)^{d_k} \frac{\sum_{s=0}^{d_k} z_{k,s} \hat{y}_k^{\varepsilon s}}{\bigoplus_{s=0}^{d_k} z_{k,s} y_k^{\varepsilon s}} & \text{if } i = k, \\ x_i & \text{if } i \neq k, \end{cases}$$

where $\varepsilon = \pm 1$, $[a]_+ = \max(a, 0)$, and we set

$$(2-7) \quad \hat{y}_i = y_i \prod_{j=1}^n x_j^{b_{ji}}.$$

When the data (\mathbf{d}, \mathbf{z}) is clearly assumed, we may drop the prefix and simply call it the (*generalized*) *mutation*.

Let $D = (d_i \delta_{ij})_{i,j=1}^n$ be the diagonal matrix with diagonal entries \mathbf{d} . It is important to note that the mutation (2-4) is equivalent to the *ordinary* mutation of exchange matrices between DB and DB' , and also between BD and $B'D$ in [Fomin and Zelevinsky 2007].

The following properties are easy to confirm:

- The formulas (2-5) and (2-6) are *independent* of the choice of the sign ε due to (2-2).
- The mutation μ_k is involutive, i.e., $\mu_k(\mu_k(\mathbf{x}, \mathbf{y}, B)) = (\mathbf{x}, \mathbf{y}, B)$.

Remark 2.3. Here we transposed every matrix in [Chekhov and Shapiro 2014]. Also, the matrix B therein is the matrix DB^T here, and β_{ij} therein is b_{ji} here.

Remark 2.4. In this paper we do not use the freedom of the choice of sign ε in (2-5) and (2-6), and it can be safely set as $\varepsilon = 1$ throughout. Nevertheless, we keep it in all formulas involved since it is useful for several purposes, for example, to consider *signed mutations*, which appeared in [Iwaki and Nakanishi 2014].

Proposition 2.5. *Under the mutation μ_k , the \hat{y} -variables (2-7) mutate in the same way as the y -variables, namely,*

$$(2-8) \quad \hat{y}'_i = \begin{cases} \hat{y}_k^{-1} & \text{if } i = k, \\ \hat{y}_i (\hat{y}_k^{[\varepsilon b_{ki}]_+})^{d_k} \left(\sum_{s=0}^{d_k} z_{k,s} \hat{y}_k^{\varepsilon s} \right)^{-b_{ki}} & \text{if } i \neq k. \end{cases}$$

Proof. This is proved using the technique in [Fomin and Zelevinsky 2007, Proposition 3.9]. \square

Next let us explain how our setting is regarded as a specialization of the setting of [Chekhov and Shapiro 2014]. In that paper a seed in \mathbb{P} is defined as a triplet

$(\mathbf{x}, \mathbf{p}, B)$, where \mathbf{x} and B are the same as in this paper (up to the identification of B as in Remark 2.3), but \mathbf{p} is a family of elements in \mathbb{P} ,

$$(2-9) \quad \mathbf{p} = (p_{i,s})_{i=1,\dots,n;s=0,\dots,d_i}.$$

Then, for the mutation $(\mathbf{x}', \mathbf{p}', B') = \mu_k(\mathbf{x}, \mathbf{p}, B)$, the following formulas replace (2-5) and (2-6):

$$(2-10) \quad \begin{aligned} p'_{k,s} &= p_{k,d_k-s}, \\ \frac{p'_{i,s}}{p'_{i,0}} &= \begin{cases} \frac{p_{i,s}}{p_{i,0}} (p_{k,d_k})^s & \text{if } i \neq k, b_{ki} \geq 0, \\ \frac{p_{i,s}}{p_{i,0}} (p_{k,0})^s & \text{if } i \neq k, b_{ki} \leq 0, \end{cases} \end{aligned}$$

$$(2-11) \quad x'_i = \begin{cases} x_k^{-1} \left(\prod_{j=1}^n x_j^{[-b_{jk}]_+} \right)^{d_k} \left(\sum_{s=0}^{d_k} p_{k,s} u_k^s \right) & \text{if } i = k, \\ x_i & \text{if } i \neq k, \end{cases}$$

where

$$(2-12) \quad u_i = \prod_{j=1}^n x_j^{b_{ji}}.$$

Now, let us start from a seed $(\mathbf{x}, \mathbf{y}, B)$ in our setting. Comparing (2-6) and (2-11), we naturally identify

$$(2-13) \quad p_{i,s} = \frac{z_{i,s} y_i^s}{\bigoplus_{r=0}^{d_i} z_{i,r} y_i^r}.$$

Then, it is easy to check that the mutation (2-10) follows from (2-2) and (2-5). Moreover, the specialization (2-13) satisfies the *normalization* property

$$(2-14) \quad \bigoplus_{s=0}^{d_i} p_{i,s} = 1$$

and the *quasireciprocity* property that for each $i = 1, \dots, n$ there is some $y_i \in \mathbb{P}$ such that

$$(2-15) \quad \frac{p_{i,s}}{p_{i,0}} \frac{p_{i,d_i}}{p_{i,d_i-s}} = y_i^{2s}, \quad s = 1, \dots, d_i.$$

Conversely, suppose that a family \mathbf{p} in (2-9) satisfies properties (2-14) and (2-15). First we note that such a y_i is unique, since any semifield \mathbb{P} is torsion-free [Fomin and Zelevinsky 2002, Section 5]. Next we define $z_{i,s} \in \mathbb{P}$ ($i = 1, \dots, n$; $s = 0, \dots, d_i$) by

$$(2-16) \quad \frac{p_{i,s}}{p_{i,0}} = y_i^s z_{i,s}.$$

In particular, we have $z_{i,0} = 1$. Then, substituting (2-16) in (2-15), we obtain

$$(2-17) \quad z_{i,s} z_{i,d_i} z_{i,d_i-s}^{-1} = 1, \quad s = 1, \dots, d_i.$$

In particular, by setting $s = d_i$, we have $z_{i,d_i}^2 = 1$. Once again, since \mathbb{P} is torsion-free, we have $z_{i,d_i} = 1$. Then, again by (2-17), we have the reciprocity $z_{i,s} = z_{i,d_i-s}$ ($s = 1, \dots, d_i - 1$). Meanwhile, by (2-14) and (2-16), we have

$$(2-18) \quad p_{i,0} = \frac{1}{\bigoplus_{s=0}^{d_i} z_{i,s} y_i^s}.$$

Then, by (2-16) again, we recover the specialization (2-13). Finally, it is straightforward to recover the mutation (2-5) from (2-10) and (2-15). Furthermore, by (2-16), one can also confirm that the coefficients $z_{i,s}$ do not mutate.

2B. Generalized cluster algebras and Laurent property. Let \mathbb{T}_n be the n -regular tree whose edges are labeled by the numbers $1, \dots, n$. Following [Fomin and Zelevinsky 2002], let us write $t \stackrel{k}{\sim} t'$ if the vertices t and t' of \mathbb{T}_n are connected by the edge labeled by k .

Definition 2.6. A (\mathbf{d}, \mathbf{z}) -cluster pattern Σ in \mathbb{P} is an assignment of a seed Σ_t in \mathbb{P} to each vertex t of \mathbb{T} such that if $t \stackrel{k}{\sim} t'$ then the assigned seeds Σ_t and $\Sigma_{t'}$ are obtained from each other by the (\mathbf{d}, \mathbf{z}) -mutation at k .

We fix a vertex t_0 of \mathbb{T}_n and call it the *initial vertex*. Accordingly, the assigned seed $\Sigma_{t_0} = (\mathbf{x}_{t_0}, \mathbf{y}_{t_0}, B_{t_0})$ at t_0 is called the *initial seed*. Let us write, for simplicity,

$$(2-19) \quad \mathbf{x}_{t_0} = \mathbf{x} = (x_1, \dots, x_n), \quad \mathbf{y}_{t_0} = \mathbf{y} = (y_1, \dots, y_n), \quad B_{t_0} = B = (b_{ij})_{i,j=1}^n.$$

On the other hand, for the seed $\Sigma_t = (\mathbf{x}_t, \mathbf{y}_t, B_t)$ assigned to a general vertex t of \mathbb{T}_n , we write

$$(2-20) \quad \mathbf{x}_t = (x_1^t, \dots, x_n^t), \quad \mathbf{y}_t = (y_1^t, \dots, y_n^t), \quad B_t = (b_{ij}^t)_{i,j=1}^n.$$

Definition 2.7. The *generalized cluster algebra* \mathcal{A} associated with a (\mathbf{d}, \mathbf{z}) -cluster pattern Σ in \mathbb{P} is a $\mathbb{Z}^{\mathbb{P}}$ -subalgebra of \mathcal{F} generated by all x -variables x_i^t ($t \in \mathbb{T}$, $i = 1, \dots, n$) occurring in Σ . It is denoted by $\mathcal{A} = \mathcal{A}(\mathbf{x}, \mathbf{y}, B; \mathbf{d}, \mathbf{z})$, where $(\mathbf{x}, \mathbf{y}, B)$ is the initial seed of Σ .

For any (\mathbf{d}, \mathbf{z}) -cluster pattern in \mathbb{P} , each x -variable x_i^t is expressed as a subtraction-free rational function of \mathbf{x} with coefficients in $\mathbb{Q}^{\mathbb{P}}$. The following stronger property due to [Chekhov and Shapiro 2014] is of fundamental importance.

Theorem 2.8 (Laurent property [Chekhov and Shapiro 2014, Theorem 2.5]). *For any (\mathbf{d}, \mathbf{z}) -cluster pattern in \mathbb{P} , each x -variable x_i^t is expressed as a Laurent polynomial of \mathbf{x} with coefficients in $\mathbb{Z}^{\mathbb{P}}$.*

2C. Example. As the simplest nontrivial example, we consider $\mathbf{d} = (2, 1)$, $\mathbf{z} = (z_{1,1})$, and an initial seed $(\mathbf{x}, \mathbf{y}, B)$ in \mathbb{P} such that

$$(2-21) \quad B = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

(This example also appears in [Chekhov and Shapiro 2014, proof of Theorem 2.7].) Accordingly,

$$(2-22) \quad \hat{y}_1 = y_1 x_2, \quad \hat{y}_2 = y_2 x_1^{-1}.$$

We note that

$$(2-23) \quad DB = \begin{pmatrix} 0 & -2 \\ 1 & 0 \end{pmatrix}, \quad BD = \begin{pmatrix} 0 & -1 \\ 2 & 0 \end{pmatrix},$$

which are the initial exchange matrices for ordinary cluster algebras of type $B_2 = C_2$. Set $\Sigma(1) = (\mathbf{x}(1), \mathbf{y}(1), B(1))$ to be the initial seed $(\mathbf{x}, \mathbf{y}, B)$, and consider the seeds $\Sigma(t) = (\mathbf{x}(t), \mathbf{y}(t), B(t))$ ($t = 2, \dots, 7$) obtained by the following sequence of alternative mutations of μ_1 and μ_2 .

$$(2-24) \quad \Sigma(1) \xleftrightarrow{\mu_1} \Sigma(2) \xleftrightarrow{\mu_2} \Sigma(3) \xleftrightarrow{\mu_1} \Sigma(4) \xleftrightarrow{\mu_2} \Sigma(5) \xleftrightarrow{\mu_1} \Sigma(6) \xleftrightarrow{\mu_2} \Sigma(7).$$

By (2-4), we have

$$(2-25) \quad B(t) = (-1)^{t+1} B.$$

Then, using the exchange relations (2-5) and (2-6), we obtain the explicit expressions of x - and y -variables in Table 1, where we set $z_{1,1} = z$ for simplicity. We observe the same periodicity of mutations of seeds for the ordinary cluster algebras of type $B_2 = C_2$.

3. Structure of seeds in generalized cluster patterns

The goal of this section is to establish some basic structural results on seeds in a (\mathbf{d}, \mathbf{z}) -cluster pattern which are parallel to the ones in [Fomin and Zelevinsky 2007].

3A. X -functions and Y -functions. Let us temporarily regard $\mathbf{y} = (y_i)_{i=1}^n$ and $\mathbf{z} = (z_{i,s})_{i=1,\dots,n; s=1,\dots,d_i-1}$ with $z_{i,s} = z_{i,d_i-s}$ as formal variables. Let $\mathbb{Q}_{\text{sf}}(\mathbf{y}, \mathbf{z})$ be the *universal semifield* of \mathbf{y} and \mathbf{z} , which consists of the rational functions in \mathbf{y} and \mathbf{z} with subtraction-free expressions [Fomin and Zelevinsky 2007]. Let $\text{Trop}(\mathbf{y}, \mathbf{z})$ be the *tropical semifield* of \mathbf{y} and \mathbf{z} , which is the multiplicative abelian group freely generated by \mathbf{y} and \mathbf{z} with *tropical sum* \oplus defined by

$$(3-1) \quad \left(\prod_i y_i^{a_i} \prod_{i,s} z_{i,s}^{a_{i,s}} \right) \oplus \left(\prod_i y_i^{b_i} \prod_{i,s} z_{i,s}^{b_{i,s}} \right) = \prod_i y_i^{\min(a_i, b_i)} \prod_{i,s} z_{i,s}^{\min(a_{i,s}, b_{i,s})}.$$

$$\begin{array}{l}
\begin{cases} x_1(1) = x_1 \\ x_2(1) = x_2 \end{cases} \\
\begin{cases} x_1(2) = x_1^{-1} \frac{1+z\hat{y}_1+\hat{y}_1^2}{1\oplus zy_1\oplus y_1^2} \\ x_2(2) = x_2 \end{cases} \\
\begin{cases} x_1(3) = x_1^{-1} \frac{1+z\hat{y}_1+\hat{y}_1^2}{1\oplus zy_1\oplus y_1^2} \\ x_2(3) = x_2^{-1} \frac{1+\hat{y}_2+z\hat{y}_1\hat{y}_2+\hat{y}_1^2\hat{y}_2}{1\oplus y_2\oplus zy_1y_2\oplus y_1^2y_2} \end{cases} \\
\begin{cases} x_1(4) = x_1x_2^{-2} \frac{1+2\hat{y}_2+\hat{y}_2^2+z\hat{y}_1\hat{y}_2+z\hat{y}_1\hat{y}_2^2+\hat{y}_1^2\hat{y}_2^2}{1\oplus 2y_2\oplus y_2^2\oplus zy_1y_2\oplus zy_1y_2^2\oplus y_1^2y_2^2} \\ x_2(4) = x_2^{-1} \frac{1+\hat{y}_2+z\hat{y}_1\hat{y}_2+\hat{y}_1^2\hat{y}_2}{1\oplus y_2\oplus zy_1y_2\oplus y_1^2y_2} \end{cases} \\
\begin{cases} x_1(5) = x_1x_2^{-2} \frac{1+2\hat{y}_2+\hat{y}_2^2+z\hat{y}_1\hat{y}_2+z\hat{y}_1\hat{y}_2^2+\hat{y}_1^2\hat{y}_2^2}{1\oplus 2y_2\oplus y_2^2\oplus zy_1y_2\oplus zy_1y_2^2\oplus y_1^2y_2^2} \\ x_2(5) = x_1x_2^{-1} \frac{1+\hat{y}_2}{1\oplus y_2} \end{cases} \\
\begin{cases} x_1(6) = x_1 \\ x_2(6) = x_1x_2^{-1} \frac{1+\hat{y}_2}{1\oplus y_2} \end{cases} \\
\begin{cases} x_1(7) = x_1 \\ x_2(7) = x_2 \end{cases}
\end{array}
\quad
\begin{array}{l}
\begin{cases} y_1(1) = y_1 \\ y_2(1) = y_2 \end{cases} \\
\begin{cases} y_1(2) = y_1^{-1} \\ y_2(2) = y_2(1\oplus zy_1\oplus y_1^2) \end{cases} \\
\begin{cases} y_1(3) = y_1^{-1}(1\oplus y_2\oplus zy_1y_2\oplus y_1^2y_2) \\ y_2(3) = y_2^{-1}(1\oplus zy_1\oplus y_1^2)^{-1} \end{cases} \\
\begin{cases} y_1(4) = y_1(1\oplus y_2\oplus zy_1y_2\oplus y_1^2y_2)^{-1} \\ y_2(4) = y_1^{-2}y_2^{-1}(1\oplus 2y_2\oplus y_2^2 \\ \oplus zy_1y_2\oplus zy_1y_2^2\oplus y_1^2y_2^2)^{-1} \end{cases} \\
\begin{cases} y_1(5) = y_1^{-1}y_2^{-1}(1\oplus y_2) \\ y_2(5) = y_1^2y_2(1\oplus 2y_2\oplus y_2^2 \\ \oplus zy_1y_2\oplus zy_1y_2^2\oplus y_1^2y_2^2)^{-1} \end{cases} \\
\begin{cases} y_1(6) = y_1y_2(1\oplus y_2)^{-1} \\ y_2(6) = y_2^{-1} \end{cases} \\
\begin{cases} y_1(7) = y_1 \\ y_2(7) = y_2 \end{cases}
\end{array}$$

Table 1. x - and y -variables for sequence (2-24).

Definition 3.1. A (\mathbf{d}, z) -cluster pattern with *principal coefficients* is a (\mathbf{d}, z) -cluster pattern in $\mathbb{P} = \text{Trop}(\mathbf{y}, z)$ with initial seed $(\mathbf{x}, \mathbf{y}, B)$, where \mathbf{x} and B are arbitrary.

Definition 3.2. Let Σ be the (\mathbf{d}, z) -cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, B)$. By the Laurent property in Theorem 2.8, each x -variable x_i^t in Σ is expressed as $X_i^t(\mathbf{x}, \mathbf{y}, z) \in \mathbb{Z}\mathbb{P}[\mathbf{x}^{\pm 1}]$ with $\mathbb{P} = \text{Trop}(\mathbf{y}, z)$. We call them the X -functions of Σ .

For principal coefficients, we actually have the following result, which is stronger than Theorem 2.8 and which is parallel to [Fomin and Zelevinsky 2003, Proposition 11.2; 2007, Proposition 3.6].

Proposition 3.3. *We have*

$$(3-2) \quad X_i^t(\mathbf{x}, \mathbf{y}, \mathbf{z}) \in \mathbb{Z}[\mathbf{x}^{\pm 1}, \mathbf{y}, \mathbf{z}].$$

Proof. We follow the argument in the proof of [Fomin and Zelevinsky 2003, Proposition 11.2]. Let p be any variable in \mathbf{y} or \mathbf{z} . Let us view $X_i^t(\mathbf{x}, \mathbf{y}, \mathbf{z})$ as a Laurent polynomial in p , say $h(p)$, whose coefficients are Laurent polynomials in the rest of the variables in \mathbf{x} , \mathbf{y} , and \mathbf{z} . We show that $h(p)$ is a polynomial in p with nonzero constant term having subtraction-free rational expression by induction on the distance between t and t_0 in \mathbb{T}_n . The crucial point is that the coefficients $p_{k,s} = z_{k,s}y_k^s / \bigoplus_{r=0}^{d_k} z_{k,r}y_k^r$ in the mutation (2-6) are normalized as (2-14). Since $\mathbb{P} = \text{Trop}(\mathbf{y}, \mathbf{z})$, this means that $p_{k,s}$ ($s = 0, \dots, d_r$) are polynomials in p , and there is no common factor in p . Thus, the right-hand side of (2-6) is a polynomial in p with nonzero constant term having subtraction-free rational expression by the induction hypothesis and the “trivial lemma” (Lemma 5.2) in [Fomin and Zelevinsky 2003]. □

Definition 3.4. We denote by Σ the (\mathbf{d}, \mathbf{z}) -cluster pattern in the universal semifield $\mathbb{Q}_{\text{sf}}(\mathbf{y}, \mathbf{z})$ with initial seed $(\mathbf{x}, \mathbf{y}, B)$. Each y -variable y_i^t in Σ is expressed as a subtraction-free rational function $Y_i^t(\mathbf{y}, \mathbf{z}) \in \mathbb{Q}_{\text{sf}}(\mathbf{y}, \mathbf{z})$. We call them the Y -functions of Σ .

Due to the universal property of the semifield $\mathbb{Q}_{\text{sf}}(\mathbf{y}, \mathbf{z})$ [Fomin and Zelevinsky 2007, Definition 2.1], the following fact holds.

Lemma 3.5. *For any (\mathbf{d}, \mathbf{z}) -cluster pattern in \mathbb{P} with the same initial exchange matrix B as above, we have*

$$(3-3) \quad y_i^t = Y_i^t|_{\mathbb{P}}(\mathbf{y}, \mathbf{z}),$$

where the right-hand side stands for the evaluation of $Y_i^t(\mathbf{y}, \mathbf{z})$ in \mathbb{P} .

3B. c -vectors, F -polynomials, and g -vectors. Let us extend the notions of c -vectors, F -polynomials, and g -vectors in [Fomin and Zelevinsky 2007] to a (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients.

3B.1. C -matrices and c -vectors. For a (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients, each y -variable $y_i^t \in \text{Trop}(\mathbf{y}, \mathbf{z})$ is, by definition, a Laurent monomial of \mathbf{y} and \mathbf{z} with coefficient 1. The following simple fact was observed in [Iwaki and Nakanishi 2014] in the special case.

Lemma 3.6. *Each y -variable y_i^t is actually a Laurent monomial of \mathbf{y} with coefficient 1.*

Proof. This is equivalent to saying that the frozen coefficients \mathbf{z} never enter in y_i^t . This is true for the initial y -variables. Then, the claim can be shown by induction

on the distance between t and t_0 in \mathbb{T}_n , by inspecting the mutation (2-5) and the definition of the tropical sum (3-1). □

Definition 3.7. Let Σ be a (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients. Let us express each y -variable y_j^t in Σ as

$$(3-4) \quad y_j^t = Y_i^t |_{\text{Trop}(\mathbf{y}, \mathbf{z})}(\mathbf{y}, \mathbf{z}) = \prod_{i=1}^n y_i^{c_{ij}^t}.$$

The resulting matrices $C^t = (c_{ij}^t)_{i,j=1}^n$ and their column vectors $c_j^t = (c_{ij}^t)_{i=1}^n$ are called the C -matrices and the c -vectors of Σ , respectively.

The following mutation/recurrence formula provides a combinatorial description of c -vectors.

Proposition 3.8. *The c -vectors of a (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients satisfy the following recurrence relation for $t \stackrel{k}{\leftarrow} t'$:*

$$(3-5) \quad c_{ij}^{t_0} = \delta_{ij},$$

$$(3-6) \quad c'_{ij} = \begin{cases} -c_{ik}^t & \text{if } j = k, \\ c_{ij}^t + c_{ik}^t[\varepsilon d_k b_{kj}^t]_+ + [-\varepsilon c_{ik}^t]_+ d_k b_{kj}^t & \text{if } j \neq k, \end{cases}$$

where $\varepsilon = \pm 1$ and it is independent of the choice of the sign ε .

Proof. As already remarked in the proof of Lemma 3.6, for a (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients, the mutation (2-5) is simplified as

$$(3-7) \quad y_i^{t'} = \begin{cases} y_k^{t-1} & \text{if } i = k, \\ y_i^t (y_k^{t[\varepsilon b_{ki}^t]_+})^{d_k} \left(\bigoplus_{s=0}^{d_k} y_k^{t \varepsilon s} \right)^{-b_{ki}^t} & \text{if } i \neq k. \end{cases}$$

This is equivalent to (3-6) due to the following formula in $\text{Trop}(\mathbf{y}, \mathbf{z})$:

$$(3-8) \quad \frac{1}{\bigoplus_{s=0}^{d_k} \left(\prod_{j=1}^n y_j^{\varepsilon c_{jk}^t} \right)^s} = \left(\prod_{j=1}^n y_j^{[-\varepsilon c_{jk}^t]_+} \right)^{d_k}. \quad \square$$

We observe that the above relation coincides with the one for the c -vectors of the ordinary cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, DB)$ in [Fomin and Zelevinsky 2002, Proposition 5.8]. Therefore, we have the following result.

Proposition 3.9. *The c -vectors of the (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, B)$ coincide with the c -vectors of the ordinary cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, DB)$.*

Alternatively, one can relate these c -vectors with the c -vectors of the ordinary cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, BD)$ as follows. Let us introduce

$$(3-9) \quad \tilde{c}_{ij}^t = d_i^{-1} c_{ij}^t d_j.$$

Then, $\tilde{c}_{ij}^{t_0} = \delta_{ij}$, and (3-6) is rewritten as

$$(3-10) \quad \tilde{c}'_{ij} = \begin{cases} -\tilde{c}_{ik}^t & \text{if } j = k, \\ \tilde{c}_{ij}^t + \tilde{c}_{ik}^t [\varepsilon b_{kj}^t d_j]_+ + [-\varepsilon \tilde{c}_{ik}^t]_+ b_{kj}^t d_j & \text{if } j \neq k. \end{cases}$$

Therefore, we have the following result.

Proposition 3.10. *The \tilde{c} -vectors, which are the column vectors in (3-9), of the (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, B)$ coincide with the c -vectors of the ordinary cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, BD)$.*

We need this alternative description for the description of the g -vectors below.

3B.2. F -polynomials. Thanks to Proposition 3.3, the following definition makes sense.

Definition 3.11. Let Σ be a (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients. For each $t \in \mathbb{T}_n$ and $i = 1, \dots, n$, a polynomial $F_i^t(\mathbf{y}, \mathbf{z}) \in \mathbb{Z}[\mathbf{y}, \mathbf{z}]$ is defined by the specialization of the X -function $X_i^t(\mathbf{x}, \mathbf{y}, \mathbf{z})$ of Σ with $x_1 = \dots = x_n = 1$. They are called the F -polynomials of Σ .

The following mutation/recurrence formula provides a combinatorial description of F -polynomials.

Proposition 3.12 (cf. [Fomin and Zelevinsky 2007, Proposition 5.1]). *The F -polynomials for a (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients satisfy the following recurrence relation for $t \xrightarrow{k} t'$:*

$$(3-11) \quad F_i^{t_0} = 1,$$

$$(3-12) \quad F_i^{t'} = \begin{cases} F_k^{t-1} \left(\prod_{j=1}^n y_j^{[-\varepsilon c_{jk}^t]_+} F_j^{t[-\varepsilon b_{jk}^t]_+} \right)^{d_k} \sum_{s=0}^{d_k} z_{k,s} \left(\prod_{j=1}^n y_j^{\varepsilon c_{jk}^t} F_j^{t \varepsilon b_{jk}^t} \right)^s & \text{if } i = k, \\ F_i^t & \text{if } i \neq k, \end{cases}$$

where $\varepsilon = \pm 1$ and it is independent of the choice of the sign ε .

Proof. By specializing the mutation (2-6) with $\mathbb{P} = \text{Trop}(\mathbf{y}, \mathbf{z})$, we obtain

$$(3-13) \quad X_i^{t'} = \begin{cases} X_k^{t-1} \left(\prod_{j=1}^n X_j^{t[-\varepsilon b'_{jk}]_+} \right)^{d_k} \frac{\sum_{s=0}^{d_k} z_{k,s} \left(\prod_{j=1}^n y_j^{\varepsilon c'_{jk}} X_j^{t \varepsilon b'_{jk}} \right)^s}{\bigoplus_{s=0}^{d_k} \left(\prod_{j=1}^n y_j^{\varepsilon c'_{jk}} \right)^s} & \text{if } i = k, \\ X_i^t & \text{if } i \neq k. \end{cases}$$

Then, specializing it with $x_1 = \dots = x_n = 1$, and using (3-8), we obtain (3-12). \square

3B.3. G -matrices and g -vectors. Let Σ be the (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, B)$. Let $\mathbb{Z}[\mathbf{x}^{\pm 1}, \mathbf{y}, \mathbf{z}]$ be the one in Proposition 3.3. Following [Fomin and Zelevinsky 2007], we introduce a \mathbb{Z}^n -grading in $\mathbb{Z}[\mathbf{x}^{\pm 1}, \mathbf{y}, \mathbf{z}]$ as follows:

$$(3-14) \quad \deg(x_i) = \mathbf{e}_i, \quad \deg(y_i) = -\mathbf{b}_j, \quad \deg(z_{i,r}) = 0.$$

Here, \mathbf{e}_i is the i -th unit vector of \mathbb{Z}^n , and $\mathbf{b}_j = \sum_{i=1}^n b_{ij} \mathbf{e}_i$ is the j -th column of the initial matrix $B = (b_{ij})_{i,j=1}^n$. Note that $\deg(\hat{y}_i) = 0$ by (2-7).

Proposition 3.13 (cf. [Fomin and Zelevinsky 2007, Proposition 6.1]). *The X -functions are homogeneous with respect to the \mathbb{Z}^n -grading.*

Proof. We repeat the original argument of Fomin and Zelevinsky, by induction on the distance between t and t_0 in \mathbb{T}_n . Using (2-6) and Lemma 3.5 specialized to a (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients, we have

$$(3-15) \quad X_i^{t'} = \begin{cases} X_k^{t-1} \left(\prod_{j=1}^n X_j^{t[-\varepsilon b'_{jk}]_+} \right)^{d_k} \frac{\sum_{s=0}^{d_k} z_{k,s} Y_k^{t \varepsilon s} |_{\mathcal{F}(\hat{\mathbf{y}}, \mathbf{z})}}{\bigoplus_{s=0}^{d_k} z_{k,s} Y_k^{t \varepsilon s} |_{\text{Trop}(\mathbf{y}, \mathbf{z})}(\mathbf{y}, \mathbf{z})} & \text{if } i = k, \\ X_i^t & \text{if } i \neq k. \end{cases}$$

Then, the right-hand side is homogeneous due to the induction hypothesis. \square

Definition 3.14. Let Σ be the (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients and initial matrix $(\mathbf{x}, \mathbf{y}, B)$. Thanks to Proposition 3.13, the degree vector $\deg(X_i^t)$ of each X -function X_i^t of Σ is defined. Let us express it as

$$(3-16) \quad \deg(X_j^t) = \sum_{i=1}^n g_{ij}^t \mathbf{e}_i.$$

The resulting matrices $G^t = (g_{ij}^t)_{i,j=1}^n$ and their column vectors $g_j^t = (g_{ij}^t)_{i=1}^n$ are called the G -matrices and the g -vectors of Σ , respectively.

The following mutation/recurrence formula provides a combinatorial description of g -vectors.

Proposition 3.15. *The g -vectors of the (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, B)$ satisfy the following recurrence relation for $t \stackrel{k}{\leftarrow} t'$:*

$$(3-17) \quad g_{ij}^{t_0} = \delta_{ij},$$

$$(3-18) \quad g_{ij}^{t'} = \begin{cases} -g_{ik}^t + \sum_{\ell=1}^n g_{i\ell}^t [-\varepsilon b_{\ell k}^t d_k]_+ - \sum_{\ell=1}^n b_{i\ell} [-\varepsilon c_{\ell k}^t d_k]_+ & \text{if } j = k, \\ g_{ij}^t & \text{if } j \neq k, \end{cases}$$

where $\varepsilon = \pm 1$ and it is independent of the choice of the sign ε .

Proof. This is obtained by comparing the degrees of both sides of (3-13). □

By using the \tilde{c} -vectors in (3-9), the relation (3-18) is rewritten as follows.

$$(3-19) \quad g_{ij}^{t'} = \begin{cases} -g_{ik}^t + \sum_{\ell=1}^n g_{i\ell}^t [-\varepsilon b_{\ell k}^t d_k]_+ - \sum_{\ell=1}^n b_{i\ell} d_{\ell} [-\varepsilon \tilde{c}_{\ell k}^t]_+ & \text{if } j = k, \\ g_{ij}^t & \text{if } j \neq k. \end{cases}$$

Having Proposition 3.10 in mind, we observe that this relation coincides with the one for the g -vectors of the ordinary cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, BD)$ in [Fomin and Zelevinsky 2007, Proposition 6.6]. Therefore, we have the following result.

Proposition 3.16. *The g -vectors of the (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, B)$ coincide with the g -vectors of the ordinary cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, BD)$.*

For the sake of completeness, we also present the counterpart of Proposition 3.10. Let us introduce

$$(3-20) \quad \tilde{g}_{ij}^t = d_i g_{ij}^t d_j^{-1}.$$

Then, the relation (3-18) is also rewritten as

$$(3-21) \quad \tilde{g}_{ij}^{t'} = \begin{cases} -\tilde{g}_{ik}^t + \sum_{\ell=1}^n \tilde{g}_{i\ell}^t [-\varepsilon d_{\ell} b_{\ell k}^t]_+ - \sum_{\ell=1}^n d_i b_{i\ell} [-\varepsilon c_{\ell k}^t]_+ & \text{if } j = k, \\ \tilde{g}_{ij}^t & \text{if } j \neq k. \end{cases}$$

Having Proposition 3.9 in mind, we observe that this relation coincides with the one for the g -vectors of the ordinary cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, DB)$. Therefore, we have the following result.

Proposition 3.17. *The \tilde{g} -vectors, which are the column vectors in (3-20), of the (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, B)$ coincide with the g -vectors of the ordinary cluster pattern with principal coefficients and initial seed $(\mathbf{x}, \mathbf{y}, DB)$.*

We see a duality between the c -vectors and the g -vectors in Propositions 3.9, 3.10, 3.16, and 3.17. In particular, the c -vectors are associated with the matrix DB , while the g -vectors are associated with the matrix BD . This is somewhat suggested from the beginning in the monomial parts in the relations (2-5) and (2-6).

3B.4. Sign-coherence.

Definition 3.18. Let Σ be a (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients. A c -vector c_j^t of Σ is said to be *sign-coherent* if it is nonzero and all components are either nonnegative or nonpositive.

Proposition 3.19 (cf. [Fomin and Zelevinsky 2007, Proposition 5.6]). *For any (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients, the following two conditions are equivalent.*

- (i) Any F -polynomial $F_i^t(\mathbf{y}, \mathbf{z})$ has constant term 1.
- (ii) Any c -vector c_i^t is sign-coherent.

Proof. This is proved by an argument parallel to the one in [Fomin and Zelevinsky 2007, Proposition 5.6] by using the recursion relation (3-12) for the F -polynomials. We omit the details. \square

In the ordinary case it was conjectured in [Fomin and Zelevinsky 2007, Conjecture 5.6] that the sign-coherence holds for any c -vector of any cluster pattern with principal coefficients. This was proved by Derksen et al. [2010, Theorem 1.7] when the initial exchange matrix B is skew-symmetric, and very recently it was proved in full generality by Gross et al. [2014, Corollary 5.5]. Since our c -vectors are identified with the c -vectors of some ordinary cluster pattern with principal coefficients by Proposition 3.9, we obtain the following theorem as a corollary of [Gross et al. 2014, Corollary 5.5].

Theorem 3.20. *Any c -vector of any (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients is sign-coherent.*

As a consequence of the sign-coherence, we also obtain the following duality between the C - and G -matrices by applying [Nakanishi and Zelevinsky 2012, Equation (3.11)] (see also [Nakanishi 2012, Proposition 3.2]), which is valid under the sign-coherence property. Recall that for a skew-symmetrizable matrix B the matrix DB is still skew-symmetrizable.

Proposition 3.21 (cf. [Nakanishi and Zelevinsky 2012, Equation (3.11)]). *Let C^t and G^t be the C - and G -matrices at $t \in \mathbb{T}_n$ of any (\mathbf{d}, \mathbf{z}) -cluster pattern Σ with principal coefficients. Let $R = (r_i \delta_{ij})_{i,j=1}^n$ be a diagonal matrix with positive diagonal entries such that RDB is skew-symmetric. Then*

$$(3-22) \quad R^{-1} D^{-1} (G^t)^T D R C^t = I.$$

Proof. This is obtained by combining [Nakanishi and Zelevinsky 2012, Equation (3.11)] with Propositions 3.9 and 3.17. \square

3C. Main formulas. Finally, we present the main formulas expressing the x - and y -variables of any (\mathbf{d}, \mathbf{z}) -cluster pattern Σ in any semifield \mathbb{P} in terms of F -polynomials, c -vectors, and g -vectors defined for the same initial exchange matrix of Σ .

Theorem 3.22 (cf. [Fomin and Zelevinsky 2007, Proposition 3.13]). *For any (\mathbf{d}, \mathbf{z}) -cluster pattern in \mathbb{P} ,*

$$(3-23) \quad y_i^t = \prod_{j=1}^n y_j^{c_{ji}^t} \prod_{j=1}^n F_j^t|_{\mathbb{P}}(\mathbf{y}, \mathbf{z})^{b_{ji}^t}.$$

Proof. We apply Lemma 3.5 to a (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients, and we obtain

$$(3-24) \quad \hat{y}_i^t = Y_i^t(\hat{\mathbf{y}}, \mathbf{z}).$$

On the other hand, specializing (2-7) to the same (\mathbf{d}, \mathbf{z}) -cluster pattern with principal coefficients, we have

$$(3-25) \quad \hat{y}_i^t = Y_i^t|_{\text{Trop}(\mathbf{y}, \mathbf{z})}(\mathbf{y}, \mathbf{z}) \prod_{j=1}^n X_j^t(\mathbf{x}, \mathbf{y}, \mathbf{z})^{b_{ji}^t} = \prod_{j=1}^n y_j^{c_{ji}^t} \prod_{j=1}^n X_j^t(\mathbf{x}, \mathbf{y}, \mathbf{z})^{b_{ji}^t},$$

where we used (3-4) in the second equality. Thus, we have

$$(3-26) \quad Y_i^t(\hat{\mathbf{y}}, \mathbf{z}) = \prod_{j=1}^n y_j^{c_{ji}^t} \prod_{j=1}^n X_j^t(\mathbf{x}, \mathbf{y}, \mathbf{z})^{b_{ji}^t}.$$

Now, we set $x_1 = \cdots = x_n = 1$. Then, $\hat{\mathbf{y}} = \mathbf{y}$, and we obtain

$$(3-27) \quad Y_i^t(\mathbf{y}, \mathbf{z}) = \prod_{j=1}^n y_j^{c_{ji}^t} \prod_{j=1}^n F_j^t(\mathbf{y}, \mathbf{z})^{b_{ji}^t}.$$

Finally, evaluating it in \mathbb{P} , we obtain (3-23). \square

Theorem 3.23 (cf. [Fomin and Zelevinsky 2007, Corollary 6.3]). *For any (\mathbf{d}, \mathbf{z}) -cluster pattern in \mathbb{P} ,*

$$(3-28) \quad x_i^t = \left(\prod_{j=1}^n x_j^{g_{ji}^t} \right) \frac{F_i^t|_{\mathcal{F}}(\hat{\mathbf{y}}, \mathbf{z})}{F_i^t|_{\mathbb{P}}(\mathbf{y}, \mathbf{z})}.$$

Proof. First, we obtain the following equality in exactly the same way as [Fomin and Zelevinsky 2007, Theorem 3.7], and we skip its derivation:

$$(3-29) \quad x_i^t = \frac{X_i^t|_{\mathcal{F}}(\mathbf{x}, \mathbf{y}, \mathbf{z})}{F_i^t|_{\mathbb{P}}(\mathbf{y}, \mathbf{z})}.$$

On the other hand, by the definition of the g -vectors, we have

$$(3-30) \quad X_i^t \left(\dots, \gamma_i x_i, \dots; \dots, \prod_{j=1}^n \gamma_k^{-b_{ki}} y_i, \dots; \dots, z_{i,r}, \dots \right) = \left(\prod_{j=1}^n \gamma_j^{g_{ji}^t} \right) X_i^t(\mathbf{x}, \mathbf{y}, \mathbf{z}).$$

By setting $\gamma_i = x_i^{-1}$, we have

$$(3-31) \quad F_i^t(\hat{\mathbf{y}}, \mathbf{z}) = \left(\prod_{j=1}^n x_j^{-g_{ji}^t} \right) X_i^t(\mathbf{x}, \mathbf{y}, \mathbf{z}).$$

Combining it with (3-29), we obtain (3-28). \square

3D. Example. Let us consider the example in Section 2C again. From the data in Table 1, one can read off the following data for the C -matrix $C(t)$, the G -matrix $G(t)$, and the F -polynomials $F_i(t)$ for the seed $\Sigma(t)$ with principal coefficients therein.

$$\begin{aligned} C(1) &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & G(1) &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & \begin{cases} F_1(1) = 1, \\ F_2(1) = 1, \end{cases} \\ C(2) &= \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, & G(2) &= \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, & \begin{cases} F_1(2) = 1 + zy_1 + y_1^2, \\ F_2(2) = 1, \end{cases} \\ C(3) &= \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, & G(3) &= \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, & \begin{cases} F_1(3) = 1 + zy_1 + y_1^2, \\ F_2(3) = 1 + y_2 + zy_1y_2 + y_1^2y_2, \end{cases} \\ C(4) &= \begin{pmatrix} 1 & -2 \\ 0 & -1 \end{pmatrix}, & G(4) &= \begin{pmatrix} 1 & 0 \\ -2 & -1 \end{pmatrix}, & \begin{cases} F_1(4) = 1 + 2y_2 + y_2^2 \\ \quad + zy_1y_2 + zy_1y_2^2 + y_1^2y_2^2, \\ F_2(4) = 1 + y_2 + zy_1y_2 + y_1^2y_2, \end{cases} \\ C(5) &= \begin{pmatrix} -1 & 2 \\ -1 & 1 \end{pmatrix}, & G(5) &= \begin{pmatrix} 1 & 1 \\ -2 & -1 \end{pmatrix}, & \begin{cases} F_1(5) = 1 + 2y_2 + y_2^2 \\ \quad + zy_1y_2 + zy_1y_2^2 + y_1^2y_2^2, \\ F_2(5) = 1 + y_2, \end{cases} \\ C(6) &= \begin{pmatrix} 1 & 0 \\ 1 & -1 \end{pmatrix}, & G(6) &= \begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}, & \begin{cases} F_1(6) = 1, \\ F_2(6) = 1 + y_2, \end{cases} \\ C(7) &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & G(7) &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & \begin{cases} F_1(7) = 1, \\ F_2(7) = 1. \end{cases} \end{aligned}$$

Acknowledgements

We thank Anne-Sophie Gleitz, Kohei Iwaki, and Michael Shapiro for useful discussions and communications.

References

- [Chekhov and Shapiro 2014] L. Chekhov and M. Shapiro, “Teichmüller spaces of Riemann surfaces with orbifold points of arbitrary order and cluster variables”, *Int. Math. Res. Not.* **2014**:10 (2014), 2746–2772. MR 3214284 Zbl 1301.30042
- [Derksen et al. 2010] H. Derksen, J. Weyman, and A. Zelevinsky, “Quivers with potentials and their representations, II: Applications to cluster algebras”, *J. Amer. Math. Soc.* **23**:3 (2010), 749–790. MR 2012c:16044 Zbl 1208.16017
- [Fomin and Zelevinsky 2002] S. Fomin and A. Zelevinsky, “Cluster algebras, I: Foundations”, *J. Amer. Math. Soc.* **15**:2 (2002), 497–529. MR 2003f:16050 Zbl 1021.16017
- [Fomin and Zelevinsky 2003] S. Fomin and A. Zelevinsky, “Cluster algebras, II: Finite type classification”, *Invent. Math.* **154**:1 (2003), 63–121. MR 2004m:17011 Zbl 1054.17024
- [Fomin and Zelevinsky 2007] S. Fomin and A. Zelevinsky, “Cluster algebras, IV: Coefficients”, *Compos. Math.* **143**:1 (2007), 112–164. MR 2008d:16049 Zbl 1127.16023
- [Gekhtman et al. 2005] M. Gekhtman, M. Shapiro, and A. Vainshtein, “Cluster algebras and Weil–Petersson forms”, *Duke Math. J.* **127**:2 (2005), 291–311. Correction at **139**:2 (2007), 407–409. MR 2006d:53103 Zbl 1079.53124
- [Gleitz 2014] A.-S. Gleitz, “Quantum affine algebras at roots of unity and generalised cluster algebras”, preprint, 2014. arXiv 1410.2446
- [Gross et al. 2014] M. Gross, P. Hacking, S. Keel, and M. Kontsevich, “Canonical bases for cluster algebras”, preprint, 2014. arXiv 1411.1394
- [Iwaki and Nakanishi 2014] K. Iwaki and T. Nakanishi, “Exact WKB analysis and cluster algebras, II: Simple poles, orbifold points, and generalized cluster algebras”, preprint, 2014. arXiv 1409.4641
- [Nakanishi 2012] T. Nakanishi, “Tropicalization method in cluster algebras”, pp. 95–115 in *Tropical geometry and integrable systems* (Glasgow, 2011), edited by C. Athorne et al., *Contemp. Math.* **580**, Amer. Math. Soc., Providence, RI, 2012. MR 2985390
- [Nakanishi and Zelevinsky 2012] T. Nakanishi and A. Zelevinsky, “On tropical dualities in cluster algebras”, pp. 217–226 in *Algebraic groups and quantum groups* (Nagoya, 2010), edited by S. Ariki et al., *Contemp. Math.* **565**, Amer. Math. Soc., Providence, RI, 2012. MR 2932428 Zbl 06296873
- [Plamondon 2011] P.-G. Plamondon, “Cluster algebras via cluster categories with infinite-dimensional morphism spaces”, *Compos. Math.* **147**:6 (2011), 1921–1954. MR 2862067 Zbl 1244.13017
- [Rupel 2013] D. Rupel, “Greedy bases in rank 2 generalized cluster algebras”, preprint, 2013. arXiv 1309.2567

Received November 27, 2014. Revised January 21, 2015.

TOMOKI NAKANISHI
GRADUATE SCHOOL OF MATHEMATICS
NAGOYA UNIVERISTY
CHIKUSA-KU
NAGOYA 464-8602
JAPAN
nakanisi@math.nagoya-u.ac.jp

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 277 No. 1 September 2015

Real positivity and approximate identities in Banach algebras	1
DAVID P. BLECHER and NARUTAKA OZAWA	
On shrinking gradient Ricci solitons with nonnegative sectional curvature	61
MINGLIANG CAI	
From quasimodes to resonances: exponentially decaying perturbations	77
ORAN GANNOT	
A general simple relative trace formula	99
JAYCE R. GETZ and HEEKYOUNG HAHN	
Chern-Simons functions on toric Calabi-Yau threefolds and Donaldson-Thomas theory	119
ZHENG HUA	
On the flag curvature of a class of Finsler metrics produced by the navigation problem	149
LIBING HUANG and XIAOHUAN MO	
Angular distribution of diameters for spheres and rays for planes	169
NOBUHIRO INNAMI and YUYA UNEME	
A note on an L^p -Brunn–Minkowski inequality for convex measures in the unconditional case	187
ARNAUD MARSIGLIETTI	
Structure of seeds in generalized cluster algebras	201
TOMOKI NAKANISHI	
Inequalities of Alexandrov–Fenchel type for convex hypersurfaces in hyperbolic space and in the sphere	219
YONG WEI and CHANGWEI XIONG	
Upper bounds of root discriminant lower bounds	241
SIMAN WONG	



0030-8730(201509)277:1;1-V