

*Pacific
Journal of
Mathematics*

GAMES AND ELEMENTARY EQUIVALENCE OF II_1 -FACTORS

ISAAC GOLDBRING AND THOMAS SINCLAIR

GAMES AND ELEMENTARY EQUIVALENCE OF II_1 -FACTORS

ISAAC GOLDBRING AND THOMAS SINCLAIR

We use Ehrenfeucht–Fraïssé games to give a local geometric criterion for elementary equivalence of II_1 -factors. We obtain as a corollary that two II_1 -factors are elementarily equivalent if and only if their unitary groups are elementarily equivalent as \mathbb{Z}_4 -metric spaces.

Introduction

While most mathematicians are concerned with determining when two objects in their field are isomorphic, logicians tend to be concerned with the coarser notion of *elementary equivalence*. Two (classical) structures M and N are said to be elementarily equivalent if and only if for any first-order sentence σ (in the language appropriate to the study of M and N), we have σ is true in M if and only if σ is true in N . For structures appearing in analysis, a continuous logic is used in which sentences can now take a continuum of “truth” values; the appropriate notion of elementary equivalence is that the truth values of all sentences are the same in both structures.

The model-theoretic study of tracial von Neumann algebras began in earnest in [Farah et al. 2013; 2014a; 2014b]. At the moment, there are only three distinct elementary equivalence classes of II_1 -factors known. (This should not be so surprising as it took a while for many isomorphism classes of II_1 -factors to be discovered and elementary equivalence is a much coarser notion.) Indeed, it was observed in [Farah et al. 2014b] that Property (Γ) and the property of being McDuff are both elementary properties (for separable II_1 -factors). Thus, if we let M_{DL} be a separable II_1 -factor that has Property (Γ) but is not McDuff (see [Dixmier and Lance 1969]), then M_{DL} , the hyperfinite II_1 -factor \mathcal{R} and the free group factor $L(\mathbb{F}_2)$ are mutually nonelementarily equivalent. Amongst those studying II_1 -factors from a model-theoretic point of view, it is widely agreed that there should be more than three elementary equivalence classes of II_1 -factors; in fact, there should probably be continuum many elementary equivalence classes. At the moment, we cannot even answer the question: is $\mathcal{R} \otimes L(\mathbb{F}_2)$ elementarily equivalent to \mathcal{R} ? In

Goldbring’s work was partially supported by NSF grant DMS-1007144. Sinclair was supported by an NSF RTG Assistant Adjunct Professorship.

MSC2010: primary 46L10; secondary 03C20.

Keywords: II_1 -factors, model theory, Ehrenfeucht–Fraïssé games.

order to accomplish these goals, we need more tools for understanding elementary equivalence of II_1 -factors.

Ehrenfeucht–Fraïssé games have long been a tool in model theory for establishing that structures are elementarily equivalent. In [Heinrich and Henson 1986], the authors exhibit an Ehrenfeucht–Fraïssé-type game used to establish elementary equivalence for Banach spaces. In this note, we adapt the game from [loc. cit.] and combine it with an argument of Kirchberg [1993] in order to characterize elementary equivalence for II_1 -factors belonging to the class \mathcal{K}_{op} (to be defined below). We should note that, currently, we do not know of a II_1 -factor that does not belong to the class \mathcal{K}_{op} and the existence of such a factor would already lead to two new theories of II_1 -factors!

Recall Dye’s theorem [1955], which states that any two factors not of type I_2^n (e.g., any two II_1 -factors) are isomorphic if and only if their unitary groups are isomorphic (even as discrete groups). Combining Dye’s theorem with the Keisler–Shelah theorem (which states that two structures are elementarily equivalent if and only if they have isomorphic ultrapowers) and the fact that the functors of taking ultrapowers and taking unitary groups commute, we see that two II_1 -factors are elementarily equivalent if and only if their unitary groups are elementarily equivalent as metric groups (with respect to the ℓ_2 metric). Using the aforementioned Ehrenfeucht–Fraïssé games and some further arguments, our main result is that we can improve upon the previous sentence, essentially removing the group structure:

Theorem 0.1. *Suppose that M and N are II_1 -factors belonging to the class \mathcal{K}_{op} . Then M and N are elementarily equivalent if and only if $U(M)$ and $U(N)$ are elementarily equivalent as \mathbb{Z}_4 -metric spaces.*

Here, by a \mathbb{Z}_4 -metric space, we mean a metric space X equipped with an action of \mathbb{Z}_4 on X by isometries. Unitary groups of von Neumann algebras will always be considered as \mathbb{Z}_4 -metric spaces by having the generator of \mathbb{Z}_4 act by multiplication by i .

In this paper, we assume that the reader is familiar with some basic model theory and von Neumann algebra theory. Good references for continuous model theory are [Ben Yaacov et al. 2008] and [Farah et al. 2014a]; the latter is geared towards the model-theoretic study of operator algebras.

All normed spaces are assumed to be over the complex numbers, \mathbb{C} . For a normed space X , we denote the closed unit ball by $(X)_1 := \{x \in X : \|x\| \leq 1\}$.

For the reader’s convenience, we now recall the original notion of Ehrenfeucht–Fraïssé games in the context of continuous logic. This has not appeared in the literature but has appeared in some online lecture notes of Bradd Hart [2012]. Fix an arbitrary language \mathcal{L} and atomic formulae $\varphi_1(\vec{x}), \dots, \varphi_k(\vec{x})$ in the variables $\vec{x} = (x_1, \dots, x_n)$ and $\varepsilon > 0$. The Ehrenfeucht–Fraïssé game $\mathfrak{G}(\varphi_1, \dots, \varphi_k, \varepsilon)$ is played with \mathcal{L} -structures M and N as follows: First Player I chooses $a_1 \in M$ or

$b_1 \in N$ respecting the sort of x_1 . Player II chooses $b_2 \in N$ or $a_2 \in M$ respectively. The players alternate in this manner until they have produced sequences $a_1, \dots, a_n \in M$ and $b_1, \dots, b_n \in N$. Player II then wins the game if and only if for each $i = 1, \dots, k$, we have $|\varphi_i(\vec{a})^M - \varphi_i(\vec{b})^N| \leq \varepsilon$. It is then a theorem that $M \equiv N$ if and only if Player II has a winning strategy in each $\mathfrak{G}(\varphi_1, \dots, \varphi_k, \varepsilon)$.

1. The class \mathcal{K}_{op}

Given a C^* algebra A , recall that its opposite algebra A^{op} is the algebra obtained from A by multiplying elements in the opposite order; that is, for $a, b \in A$, we have $a \cdot_{op} b := b \cdot a$. It is immediate that A^{op} is once again a C^* algebra. Furthermore, if A is a von Neumann algebra, then A^{op} is also a von Neumann algebra. Note also that if $(A_i : i \in I)$ is a family of C^* algebras (resp. tracial von Neumann algebras) and \mathcal{U} is an ultrafilter on I , then $(\prod_{\mathcal{U}} A_i)^{op} \cong \prod_{\mathcal{U}} A_i^{op}$ via the identity map, where the ultraproduct is understood to be the usual C^* algebra ultraproduct (resp. tracial ultraproduct).

Many of the naturally occurring tracial von Neumann algebras are isomorphic to their opposites, e.g., \mathcal{R} and $L(G)$ (G any group). There are examples of tracial von Neumann algebras that are not isomorphic to their opposites (see [Connes 1975]). During a seminar talk given by the first author at Vanderbilt University, Jesse Peterson asked whether or not the class of all tracial von Neumann algebras isomorphic to their opposites is an axiomatizable class. While we do not know the answer to this question (although we suspect the answer is negative), the answer is positive if one replaces the word “isomorphism” by “elementary equivalence” as we show in the following:

Proposition 1.1. *The class of all tracial von Neumann algebras that are elementarily equivalent to their opposites is an elementary class.*

Definition 1.2. We let \mathcal{K}_{op} denote the class of all tracial von Neumann algebras elementarily equivalent to their opposites.

Proof of Proposition 1.1. We present a proof suggested to us by Todor Tsankov as well as independently by the anonymous referee. There is a collection of axioms for the class \mathcal{K}_{op} : for every term t , recursively define the term t^{op} by defining $(t_1 \cdot t_2)^{op} := t_2^{op} \cdot t_1^{op}$. Then one can recursively define, for any formula φ , the formula φ^{op} , the key clause being the atomic formulae, where one replaces every occurrence of a term t by the term t^{op} . Then the conditions $|\sigma - \sigma^{op}| = 0$, as σ ranges over all sentences, axiomatizes the class \mathcal{K}_{op} . \square

We remark in passing that alternately by [Ben Yaacov et al. 2008, Proposition 5.14], it suffices to show that \mathcal{K}_{op} is closed under isomorphisms, ultraproducts, and ultraroots. We leave it as an exercise to verify these properties for \mathcal{K}_{op} .

Since \mathcal{R} and $L(\mathbb{F}_2)$ are isomorphic to their opposites, they belong to \mathcal{K}_{op} . Moreover, the example M_{DL} of a II_1 -factor with Property (Γ) that is not McDuff given by Lance and Dixmier [1969] is also isomorphic to its opposite. Thus, we have this:

Corollary 1.3. *If there is a II_1 -factor that does not belong to \mathcal{K}_{op} , then there are at least five theories of II_1 -factors.*

Proof. If N is a II_1 -factor that does not belong to \mathcal{K}_{op} , then the theories of N and N^{op} differ from each other and from the three known theories of II_1 -factors. \square

Question 1.4. Are there more “explicit” axioms for the class \mathcal{K}_{op} ? Can one use typical model-theoretic preservation theorems to show that \mathcal{K}_{op} is universally axiomatizable or $\forall\exists$ -axiomatizable?

Question 1.5. Is there a single sentence σ such that adding the condition “ $\sigma = 0$ ” to the axioms for II_1 -factors gives an axiomatization of \mathcal{K}_{op} ?

A negative answer to the last question implies that there must be infinitely many elementary equivalence classes of II_1 -factors not belonging to \mathcal{K}_{op} . Indeed, if there are only finitely many elementary equivalence classes of II_1 -factors not belonging to \mathcal{K}_{op} , then the class of II_1 -factors not belonging to \mathcal{K}_{op} is readily verified to be elementary as well, whence a typical compactness argument is used to show that the last question has a positive answer.

2. Model theory of Banach pairs

In order to frame the main results of the paper in the next section on the model theory of II_1 -factors, we introduce a class of linear (unbounded) metric structures (“Banach pairs”) for which II_1 -factors will be the primary set of examples. The important fact which we will see is that the theory of a II_1 -factor regarded as a Banach pair will determine its theory as II_1 -factor. For this reason we feel it is justified to introduce this treatment, despite several existing approaches in the literature for dealing with linear metric structures, e.g., [Ben Yaacov 2008; Ben Yaacov et al. 2008; Henson and Moore 1983], with at least one treatment [Farah et al. 2014a] being devoted to C^* -algebras and tracial von Neumann algebras.

Definition 2.1. A *Banach pair* (X, \mathcal{C}) consist of a normed space X and a distinguished subset $\mathcal{C} \subset (X)_1$ which is

- complete;
- roundly convex, i.e., $\lambda x + \mu y \in \mathcal{C}$ for all $x, y \in \mathcal{C}$ and $\lambda, \mu \in \mathbb{C}$ with $|\lambda| + |\mu| \leq 1$;
- generating, i.e., $\bigcup_n n \cdot \mathcal{C} = X$.

The main examples of Banach pairs we will be interested in are where $X = M$, a tracial von Neumann algebra equipped with the 2-norm $\|x\|_2 := \text{tr}(x^*x)^{1/2}$, and $\mathcal{C} = (M)_1$, the (norm) closed unit ball.

A Banach pair (X, \mathcal{C}) can be interpreted as a structure for the language \mathcal{L}_{BP} below:

- There is one sort each for \mathbb{C} and X .
- There is a sequence of domains of quantification \mathcal{C}_n for X .
- There are function symbols $\iota_{m,n} : \mathcal{C}_m \rightarrow \mathcal{C}_n$ for $m \leq n$ to be interpreted as the usual inclusion maps.
- X is given the usual complex normed space axioms.
- There are axioms which show $0_X \in \mathcal{C}_1 \subset (X)_1$.
- There are axioms to show each \mathcal{C}_n is roundly convex.

For a Banach pair (X, \mathcal{C}) and $x \in X$, we define $\|x\|_{\mathcal{C}} := \inf\{t > 0 : x \in t \cdot \mathcal{C}\}$, which can be checked to be a Banach norm on X . However, note that $\|\cdot\|_{\mathcal{C}}$ is a definable predicate if and only if it is uniformly continuous with respect to the usual norm. (In the case that X is a tracial von Neumann algebra, this will be the case if and only if X is finite-dimensional.)

As an \mathcal{L}_{BP} -structure, the ultrapower $(X, \mathcal{C})^{\mathcal{U}}$ can be identified with the Banach pair $(X^{\mathcal{U}}, \mathcal{C}^{\mathcal{U}})$, where $X^{\mathcal{U}}$ is the quotient space of $\{(x_i) : \lim_{\mathcal{U}} \|x_i\|_{\mathcal{C}} < \infty\}$ modulo the subspace $\{(z_i) : \lim_{\mathcal{U}} \|z_i\|_{\mathcal{C}} < \infty, \lim_{\mathcal{U}} \|z_i\| = 0\}$ and $\mathcal{C}^{\mathcal{U}} \subset X^{\mathcal{U}}$ is defined in the obvious way.

We say that two Banach pairs (X, \mathcal{C}) and (Y, \mathcal{D}) are isomorphic (written $(X, \mathcal{C}) \cong (Y, \mathcal{D})$) if they are isomorphic as \mathcal{L}_{BP} -structures, that is, if there is an isometry $T : X \rightarrow Y$ so that $T(\mathcal{C}) = \mathcal{D}$. By definition, the aforementioned Banach pairs are elementarily equivalent (written $(X, \mathcal{C}) \equiv (Y, \mathcal{D})$) if $\text{Th}(X, \mathcal{C}) = \text{Th}(Y, \mathcal{D})$. As a consequence of the Keisler–Shelah theorem in continuous logic, we have that $(X, \mathcal{C}) \equiv (Y, \mathcal{D})$ if and only if there is an ultrafilter so that $(X, \mathcal{C})^{\mathcal{U}} \cong (Y, \mathcal{D})^{\mathcal{U}}$. See [Henson and Iovino 2002, §10] for a proof of this fact in the context of normed spaces or [Heinrich and Henson 1986, §3] for a more explicit construction for Banach spaces.

Our main observation in this section is that for Banach pairs (X, \mathcal{C}) and (Y, \mathcal{D}) elementary equivalence can be characterized in terms of the pairs “having the same local geometric structure” by the use of Ehrenfeucht–Fraïssé games. For the very similar case of Banach spaces, this was done by Heinrich and Henson [1986, Theorem 4] and the case of normed spaces is largely similar (see [Henson and Iovino 2002, Remark 10.10]).

We now describe precisely what we mean when we say that two Banach pairs (X, \mathcal{C}) and (Y, \mathcal{D}) have the same local geometric structure. For E a subspace of X and F a subspace of Y , we say that a linear bijection $T : E \rightarrow F$ is an ε -almost isometry if $\|T\|, \|T^{-1}\| \leq 1 + \varepsilon$ and $T(E \cap \mathcal{C}) \subset_{\varepsilon} F \cap \mathcal{D}$ and $T^{-1}(F \cap \mathcal{D}) \subset_{\varepsilon} E \cap \mathcal{C}$. (We write $A \subset_{\varepsilon} B$ if $\sup_{x \in A} \inf_{y \in B} \|x - y\| \leq \varepsilon$.)

The following is adapted from [Heinrich and Henson 1986, §2]; see also [Henson and Moore 1983, §8]. We describe a game $\mathfrak{G}(n, \varepsilon)$ played by two players with Banach pairs (X, \mathcal{C}) and (Y, \mathcal{D}) , where $\varepsilon > 0$ and n are fixed parameters.

Step 1. Player I chooses a one-dimensional subspace, either $E_1 \subset X$ or $F_1 \subset Y$. Player II then chooses a subspace, respectively $F_1 \subset Y$ or $E_1 \subset X$ and a linear bijection $T_1 : E_1 \rightarrow F_1$.

Step i . Player I chooses an at most one-dimensional extension, either $E_i \supset E_{i-1}$ or $F_i \supset F_{i-1}$. Player II then chooses a subspace, respectively $F_i \subset Y$ or $E_i \subset X$, and a linear bijection $T_i : E_i \rightarrow F_i$ which extends T_{i-1} .

Step n . The players make their choices, and the game terminates. Player II wins if $T_n : E_n \rightarrow F_n$ is an ε -almost isometry; otherwise, Player I wins.

During the course of proofs, we may speak of Player I playing $x_i \in X$, in which case we mean that Player I plays $\text{span}(E_{i-1} \cup \{x_i\})$. We may then also say that Player II responds with $y_i \in Y$, in which case we mean that Player II plays the linear bijection T_i extending T_{i-1} that sends x_i to y_i .

Definition 2.2. We say that Banach pairs (X, \mathcal{C}) and (Y, \mathcal{D}) are *locally equivalent* (written $(X, \mathcal{C}) \cong_{\text{loc}} (Y, \mathcal{D})$) if for every $\varepsilon > 0$ and every n , Player II has a winning strategy for the game $\mathfrak{G}(n, \varepsilon)$.

Remark 2.3. Since ε is arbitrary, and we need only deal with at most one-dimensional extensions, we see that local isomorphism remains the same under an alternate version of ε -almost isometry, namely, the existence of linear bijections $T : E \rightarrow F$, $S : F \rightarrow E$ with strict containment $T(E \cap \mathcal{C}) \subseteq F \cap \mathcal{D}$ and $S(F \cap \mathcal{D}) \subseteq E \cap \mathcal{C}$ so that $\|ST - \text{id}_E\|$, $\|TS - \text{id}_F\| < \varepsilon$ and $\|T\|, \|S\| < 1 + \varepsilon$.

Proposition 2.4. *The following statements are equivalent:*

- (1) $(X, \mathcal{C}) \equiv (Y, \mathcal{D})$.
- (2) *There exists an ultrafilter so that $(X, \mathcal{C})^{\mathcal{U}} \cong (Y, \mathcal{D})^{\mathcal{U}}$ as Banach pairs.*
- (3) $(X, \mathcal{C}) \cong_{\text{loc}} (Y, \mathcal{D})$.

As noted above, (1) \iff (2) is the Keisler–Shelah theorem applied to the language of Banach pairs. The proof of (2) \implies (3) is straightforward using representing sequences. Therefore we only need to prove (3) \implies (1). The proof is more or less identical to the Banach space version as in [Heinrich and Henson 1986]. However, since we are working in a different logic, we sketch a (nearly complete) proof here for the convenience of the reader.

Sketch of (3) \implies (1). First, we work with the notion of ε -almost isometry as described in Remark 2.3. Let σ be a sentence of the form $\inf_{v_1} \sup_{v_2} \cdots Q_{v_n} \rho(v_1, \dots, v_n)$, where Q is \inf if n is odd and \sup if n is even and where ρ is quantifier-free. (We

suppress mention of the sorts C_i corresponding to each v_i .) Fix $\varepsilon > 0$. It suffices to show that $\sigma^{(Y, \mathcal{D})} \leq \sigma^{(X, \mathcal{C})} + \varepsilon$ for all $\varepsilon > 0$. Indeed, by symmetry of the relation of local equivalence, this shows that all sentences of the above form have the same truth values in (X, \mathcal{C}) and (Y, \mathcal{D}) . Since any sentence in prenex normal form is equivalent to one of the above form (by adding dummy variables) and since the set of sentences in prenex normal form is dense in the set of all sentences (see [Ben Yaacov et al. 2008, §6]), we obtain that $(X, \mathcal{C}) \equiv (Y, \mathcal{D})$.

Fix a sufficiently small $\delta > 0$. (We will see exactly how small δ needs to be in a moment.) Fix a winning strategy \mathcal{S} for Player II in $\mathfrak{G}(n, \delta)$. Call a play of the game $\mathfrak{G}(n, \delta)$ *regular* if

- for odd i , Player I plays $x_i \in X$, while for even i , Player I plays $y_i \in Y$;
- for each i , Player I's move at Round i is always in the sort corresponding to the variable v_i ;
- Player II always plays according to \mathcal{S} .

We say that sequences $x_1, \dots, x_k \in X$ and $y_1, \dots, y_k \in Y$ *correspond* if they are the results of the first k rounds of a regular play of $\mathfrak{G}(n, \delta)$.

For $0 \leq l \leq n$, let $\sigma_l(v_1, \dots, v_{n-l})$ denote the formula obtained from σ by removing the first $n-l$ quantifiers. One now proves, by induction on l ($0 \leq l \leq n$), that if $x_1, \dots, x_{n-l} \in X$ and $y_1, \dots, y_{n-l} \in Y$ correspond, then

$$\sigma_l(y_1, \dots, y_{n-l})^{(Y, \mathcal{D})} \leq \sigma_l(x_1, \dots, x_{n-l})^{(X, \mathcal{C})} + \varepsilon.$$

The base case $l=0$ follows from the fact that $T_n : \text{span}(x_1, \dots, x_n) \rightarrow \text{span}(y_1, \dots, y_n)$ is a δ -almost isometry if δ is chosen sufficiently small. We now prove the induction step. Suppose that the claim holds for l and that $x_1, \dots, x_{n-l-1} \in X$ and $y_1, \dots, y_{n-l-1} \in Y$ correspond. Let $r := \sigma_{l+1}(x_1, \dots, x_{n-l-1})^{(X, \mathcal{C})}$. First suppose that $n-l$ is odd, so that $\sigma_{l+1}(v_1, \dots, v_{n-l-1}) = \inf_{v_{n-l}} \sigma_l(v_1, \dots, v_{n-l})$. Fix $\eta > 0$ and let $x_{n-l} \in X$ be of the same sort as v_{n-l} so that $\sigma_l(x_1, \dots, x_{n-l})^{(X, \mathcal{C})} \leq r + \eta$. Let $y_{n-l} \in Y$ be Player II's response to x_{n-l} according to the strategy \mathcal{S} . Then, by induction,

$$\sigma_l(y_1, \dots, y_{n-l})^{(Y, \mathcal{D})} \leq \sigma_l(x_1, \dots, x_{n-l})^{(X, \mathcal{C})} + \varepsilon \leq r + \varepsilon + \eta.$$

Letting η go to 0 yields the desired result. The case that $n-l$ is even is similar and is left to the reader. \square

3. Elementary equivalence of II_1 -factors

We say that two tracial von Neumann algebras M and N are locally equivalent if the associated Banach pairs $(M, (M)_1)$ and $(N, (N)_1)$ are locally equivalent. Somewhat miraculously, it turns out that for II_1 -factors belonging to \mathcal{K}_{op} , local

equivalence is the same as elementary equivalence. This essentially follows from an argument of Kirchberg [1993]. First, we need to recall a fact about *Jordan morphisms* between von Neumann algebras.

Given a C^* algebra A , the *special Jordan product* on A is the operation \circ defined by $a \circ b := \frac{1}{2}(ab + ba)$ for all $a, b \in A$. If B is also a C^* algebra, then a linear map $T : A \rightarrow B$ is a *Jordan morphism* if it preserves the special Jordan product and the involution. We need the following:

Fact 3.1 (See [Hanche-Olsen and Størmer 1984, Corollary 7.4.9]). If M and N are von Neumann algebras and $T : M \rightarrow N$ is a normal Jordan homomorphism, then T is the sum of a $*$ -homomorphism and a $*$ -antihomomorphism.

Recall that a map $A \rightarrow B$ between C^* algebras is a $*$ -antihomomorphism if and only if it is a $*$ -homomorphism $A \rightarrow B^{op}$.

Suppose that M and N are von Neumann algebras and $T : M \rightarrow N$ is a unital, bijective, normal Jordan homomorphism. Write $T = T_1 + T_2$, where $T_1 : M \rightarrow N$ and $T_2 : M \rightarrow N^{op}$ are $*$ -homomorphisms. Since $T_i(1)$ is a projection for $i = 1, 2$ and $T_1(1) + T_2(1) = 1$, we have that $T_1(1)$ and $T_2(1)$ are orthogonal projections. Since $T(M) = N$, it follows that each $T_i(1)$ is a central projection. Thus, if N is a factor, it follows that $\{T_1(1), T_2(1)\} = \{0, 1\}$, whence T is either an isomorphism or an anti-isomorphism.

The following is basically Proposition 4.6 in [Kirchberg 1993].

Proposition 3.2 (Kirchberg). *Suppose that M and N are II_1 -factors. If there is an isometry $T : L^2(M, \text{tr}_M) \rightarrow L^2(N, \text{tr}_N)$ so that T maps M onto N contractively, then $M \cong N$ or $M \cong N^{op}$.*

Proof. We first show that T maps unitaries to unitaries. If $u \in M$ is a unitary, we have

$$1 = \|u\|_2^2 = \|T(u)\|_2^2 = \langle T(u), T(u) \rangle = \langle T(u)^*T(u), 1 \rangle.$$

On the other hand,

$$\|T(u)^*T(u)\|_2 \leq \|T(u)\| \cdot \|T(u)\|_2 \leq 1.$$

It follows that $T(u)^*T(u) = 1$. We thus have that $T'(x) := T(1)^*T(x)$ is unital, contractive, trace-preserving, and takes unitaries to unitaries. By the same reasoning as in the proof of [Kirchberg 1993, Proposition 4.6], T' is a weakly continuous Jordan morphism and the result follows from the discussion preceding this proposition. \square

Corollary 3.3. *Suppose that M and N are II_1 -factors. Then M is locally equivalent to N if and only if M is elementarily equivalent to N or to N^{op} . In particular, if M and N are II_1 -factors belonging to the class \mathcal{K}_{op} , then M is locally equivalent to N if and only if M is elementarily equivalent to N .*

Proof. By the downward Löwenheim–Skolem theorem (see [Farah et al. 2014a, Section 4.2]), we may suppose that M and N are separable. Suppose that M is locally equivalent to N . Then by Proposition 2.4, there is an isometry $L^2(M^{\mathcal{U}}) \rightarrow L^2(N^{\mathcal{U}})$ that maps $M^{\mathcal{U}}$ into $N^{\mathcal{U}}$ contractively. By Proposition 3.2, $M^{\mathcal{U}}$ is isomorphic to either $N^{\mathcal{U}}$ or $(N^{\mathcal{U}})^{op}$. It follows that M is elementarily equivalent to either N or N^{op} . The converse is trivial. \square

We now introduce a more useful test for determining elementary equivalence which works in the more specific case of Banach pairs $(M, (M)_1)$, where M is a II_1 -factor (or more generally a tracial von Neumann algebra) equipped with the 2-norm, and $(M)_1$ is the (operator norm) unit ball of M .

We define the game $\mathfrak{G}_{\text{vN}}(n, \varepsilon)$ in parameters n and $\varepsilon > 0$ which is played by two players with II_1 -factors M and N as follows.

Step i . Player I chooses a unitary, either $u_i \in U(M)$ or $v_i \in U(N)$. Player II then chooses a unitary, respectively $v_i \in U(N)$ or $u_i \in U(M)$, in the same manner.

Step n . The players make their choices, and the game terminates. Player II wins if $|\langle u_i, u_j \rangle - \langle v_i, v_j \rangle| < \varepsilon$ for all $1 \leq i, j \leq n$; otherwise, Player I wins.

Theorem 3.4. *The II_1 -factors M and N are locally equivalent if and only if Player II has a winning strategy for the game $\mathfrak{G}_{\text{vN}}(n, \varepsilon)$ for all parameters (n, ε) .*

In order to prove this result we will first need one lemma.

Lemma 3.5. *Let M and N be II_1 -factors, $E \subset M$ and $F \subset N$ be subspaces, and $T : (E, E \cap (M)_1) \rightarrow (F, F \cap (N)_1)$ be an ε -almost isometry. If $u \in E$ is a unitary, then there exists a unitary $v \in N$ so that $\|T(u) - v\|_2 \leq 4\sqrt{\varepsilon}$.*

Proof. In a II_1 -factor, a u is a unitary if and only if it is a contraction with $\|u\|_2 = 1$. By definition, we see that there exists a contraction $y \in N$ with $\|y - T(u)\|_2 \leq \varepsilon$. In particular, $\|y\|_2 \geq 1 - 2\varepsilon$. By a standard estimate we have that

$$\|1 - |y|\|_2^2 \leq 1 + \| |y| \|_2^2 - 2 \operatorname{tr}|y| = 1 + \operatorname{tr}(|y|^2) - 2 \operatorname{tr}|y| \leq 1 - \operatorname{tr}|y| \leq 1 - \|y\|_2 \leq 2\varepsilon,$$

whence writing $y = v|y|$ for $v \in U(N)$ we have that $\|T(u) - v\|_2 \leq 4\sqrt{\varepsilon}$. \square

Proof of Theorem 3.4. First suppose that M and N are locally equivalent. Fix n and $\varepsilon > 0$; we describe a winning strategy for Player II in the game $\mathfrak{G}_{\text{vN}}(n, \varepsilon)$. For simplicity, we suppose that $n = 2$ and describe a winning strategy for Player II; the general case is no more difficult, only the notation is more cumbersome. Fix δ sufficiently small (to be specified later) and fix a winning strategy \mathcal{S} for Player II in the game $\mathfrak{G}(2, \delta)$. Suppose that Player I first plays $u_1 \in U(M)$. (The case that Player I first plays a unitary in N is similar.) Let $y_1 \in N$ be Player II's response to u_1 in the game $\mathfrak{G}(2, \delta)$ according to \mathcal{S} . Since $u_1 \mapsto y_1$ determines a δ -almost isometry, by Lemma 3.5, there is $v_1 \in U(N)$ such that $\|y_1 - v_1\|_2 \leq 4\sqrt{\delta}$. Now

suppose that Player II responds with $v_2 \in U(N)$. (The case that Player II responds with a unitary in M is similar.) Let $x_2 \in M$ be Player II's response to (u_1, y_1, v_2) in the game $\mathfrak{G}(2, \delta)$ according to \mathcal{S} . Since $u_1 \mapsto y_1, x_2 \mapsto v_2$ determines a δ -almost isometry, we once again have $u_2 \in U(M)$ such that $\|x_2 - u_2\|_2 \leq 4\sqrt{\delta}$.

We need to verify that $|\langle u_i, u_j \rangle - \langle v_i, v_j \rangle| < \varepsilon$ for $i, j = 1, 2$. If δ is chosen small enough so that a δ -almost isometry preserves inner products within an error of $\varepsilon/3$ (use, for example, the polarization identity) and such that perturbing entries of an inner product by a distance of no more than $4\sqrt{\delta}$ changes the inner product by an amount not exceeding $\varepsilon/3$, then the desired estimates hold. For example,

$$\langle u_1, u_2 \rangle \sim_{\varepsilon/3} \langle u_1, x_2 \rangle \sim_{\varepsilon/3} \langle y_1, v_2 \rangle \sim_{\varepsilon/3} \langle v_1, v_2 \rangle.$$

We now prove the converse. Suppose that Player II has a winning strategy in all of the games $\mathfrak{G}_{v_N}(n, \varepsilon)$; we show that M and N are elementarily equivalent as Banach pairs. By symmetry, it is enough to show that $\sigma^{(M, (M)_1)} \leq r$ implies that $\sigma^{(N, (N)_1)} \leq r$ for any positive real number r and any prenex normal form sentence σ . Since $\sigma \div r$ is equivalent to a prenex normal form sentence, it is enough to prove that $\sigma^{(M, (M)_1)} = 0$ implies $\sigma^{(N, (N)_1)} = 0$ for any prenex normal form sentence σ .

Towards this end, we introduce the ‘‘unitary transform’’ of a sentence in prenex normal form. Suppose that σ is a sentence in prenex normal form, say

$$\sigma = Q_1 x_1 \cdots Q_n x_n \varphi(\vec{x}),$$

where $\varphi(\vec{x})$ is quantifier-free. We form the new sentence σ^u as follows:

- If $Q_i = \inf$ and x_i is of sort n_i , replace each occurrence of the variable x_i by the term $t_i(u_i, v_i) := n_i \cdot ((u_i + v_i)/2)$, where u_i and v_i are variables of sort \mathcal{C}_1 , and replace the quantifier $Q_i x_i$ by the quantifiers $Q_i u_i Q_i v_i$.
- The quantifier-free part of σ^u should now be

$$\max(\varphi, \max_i (\max(1 \div \|u_i\|_2, 1 \div \|v_i\|_2))).$$

For example, if $\sigma = \sup_{x_1} \inf_{x_2} \varphi(x_1, x_2)$, where x_2 is of sort \mathcal{C}_1 (for simplicity), then $\sigma^u = \sup_{x_1} \inf_{u_2} \inf_{v_2} \varphi(x_1, (u_2 + v_2)/2)$.

Also, we let σ^{uu} be the ‘‘formula’’ defined in the exact same way as σ^u except that we only allow quantifiers over the unitary groups rather than the entire unit ball. (Formally, σ^{uu} is not a formula in the sense of continuous logic, but it will be useful in the remainder of the proof.)

Claim 1. We have $\sigma^{(M, (M)_1)} = 0$ if and only if $(\sigma^u)^{(M, (M)_1)} = 0$ (and the corresponding statement for $(N, (N)_1)$).

Claim 1 follows from the fact that, in a *finite* von Neumann algebra, any contraction is an average of two unitaries. Indeed if x is a contraction in a finite von Neumann algebra, then it has polar decomposition $x = u|x|$, where u is a unitary.

As $|x|$ is a self-adjoint contraction, by functional calculus it may be written as the average of two unitaries.

Claim 2. We have $(\sigma^u)^{(M, (M)_1)} = 0$ if and only if $(\sigma^{uu})^{(M, (M)_1)} = 0$ (and the corresponding statement for $(N, (N)_1)$).

The backwards direction of Claim 2 is trivial; the forwards direction follows from the fact that if x is a contraction in a finite factor and $\|x\|_2 \geq 1 - \varepsilon$, then there is a unitary u so that $\|u - x\|_2 \leq 2\sqrt{\varepsilon}$.

Finally, suppose that σ is a sentence in prenex normal form and $\sigma^{(M, (M)_1)} = 0$. Then by Claims 1 and 2, we have $(\sigma^{uu})^{(M, (M)_1)} = 0$. Since atomic formulae are of the form $\|\lambda_1 x_1 + \cdots + \lambda_n x_n\|_2$ and arbitrary quantifier-free formulae are continuous combinations of atomic formulae, it follows from a winning strategy for Player II in $\mathfrak{G}_{\text{vN}}(n, \varepsilon)$ (for suitably small ε) that $(\sigma^{uu})^{(N, (N)_1)} = 0$, whence $\sigma^{(N, (N)_1)} = 0$ by Claims 1 and 2 again. \square

Suppose now that $\mathcal{L}_i = \{\Phi\}$, where Φ is a unary function symbol with modulus of uniform continuity $\Delta_\Phi(\varepsilon) = \varepsilon$. If M is a tracial von Neumann algebra, we view $U(M)$ as an \mathcal{L}_i -structure by interpreting Φ as multiplication by i . We then have this:

Corollary 3.6. *Let M and N be II_1 -factors. Then M and N are locally equivalent if and only if $U(M)$ and $U(N)$ are elementarily equivalent as \mathcal{L}_i -structures.*

Proof. If M and N are locally equivalent, then M is elementarily equivalent to either N or N^{op} . It follows that there is an ultrafilter \mathcal{U} such that $M^\mathcal{U}$ is isomorphic to $N^\mathcal{U}$ or $(N^{op})^\mathcal{U}$. In either case, $(U(M))^\mathcal{U} = U(M^\mathcal{U})$ is isomorphic to $(U(N))^\mathcal{U} = U(N^\mathcal{U})$ as \mathcal{L}_i -structures, whence $U(M)$ and $U(N)$ are elementarily equivalent as \mathcal{L}_i -structures.

Conversely, assume that $U(M)$ and $U(N)$ are elementarily equivalent as \mathcal{L}_i -structures. Then Player II has a winning strategy for the Ehrenfeucht–Fraïssé games for $U(M)$ and $U(N)$ as \mathcal{L}_i -structures. It then follows that Player II has a winning strategy in the games \mathfrak{G}_{vN} for M and N . Indeed, this follows from the fact that

$$\Re\langle u_i, u_j \rangle = 1 - \frac{1}{2}d(u_i, u_j)^2, \quad \Im\langle u_i, u_j \rangle = 1 - \frac{1}{2}d(u_i, i \cdot u_j)^2. \quad \square$$

Remark 3.7. Notice that the proof of the previous corollary gives an alternative proof of the forward direction of [Theorem 3.4](#).

Corollary 3.8. *Let M and N be II_1 -factors in the class \mathcal{K}_{op} . Then M and N are elementarily equivalent if and only if $U(M)$ and $U(N)$ are elementarily equivalent as \mathcal{L}_i -structures.*

Corollary 3.9. *Let M and N be II_1 -factors. Suppose that for every ε there is a $(1 + \varepsilon)$ -Lipschitz homeomorphism $f : U(M) \rightarrow U(N)$; that is, f is bijective with*

$$(1 + \varepsilon)^{-1} \|u - v\|_2 \leq \|f(u) - f(v)\|_2 \leq (1 + \varepsilon) \|u - v\|_2$$

that is further assumed to preserve the action by \mathbb{Z}_4 . Then M and N are locally isomorphic.

We will say that M and N are *approximately Lipschitz isometric* if the condition of the previous corollary is satisfied. Although this relation ought to be in principle much stronger than elementary equivalence, to the best of our knowledge the results of [Farah et al. 2014b] heretofore furnish the only known examples of properties invariant under this relation namely, the McDuff property and property (Γ) . It is, however, tempting to speculate that approximate Lipschitz isometry ought to be equivalent to isomorphism (up to opposites).

In lieu of this, it would be highly interesting to determine whether hyperfiniteness is an invariant of approximate Lipschitz isometry. If true, this would be in contrast with [Farah et al. 2014b, Theorem 4.3] which shows in particular that hyperfiniteness is not an invariant of elementary equivalence. Though one can show, essentially by Fact 3.1 and Proposition 3.2 (see also [Takesaki 2003, Chapter XIV.2]), that for every n , there exists $\varepsilon > 0$ so that for any ε -approximate Lipschitz embedding θ of M_n into a II_1 -factor N , there is a $*$ -homomorphism $\theta' : M_n \rightarrow N$ so that the image of the unit ball under θ is ε -contained in 2-norm in the image unit ball under θ' of M_n , this still does not seem sufficient, unless ε could be taken independent of n .

4. Further remarks and open problems

Of course, Corollary 3.8 raises the question: which \mathbb{Z}_4 -metric spaces arise as unitary groups of II_1 -factors? Even more importantly, what are the theories of such \mathbb{Z}_4 -metric spaces? Ignoring the extra structure for a moment, an important example of a complete theory of (noncompact) metric spaces is the theory of the *Urysohn metric space*. (See, for example, [Ealy and Goldbring 2012].) Recall that the Urysohn metric space is the unique (up to isometry) complete, separable metric space that is universal (that is, every separable metric space isometrically embeds) and ultrahomogeneous (every isometry between finite — even compact — subspaces extends to an isometry of the entire space). However, the Urysohn space (or rather, its bounded counterpart, the Urysohn sphere) could never be isometric to the unitary group of a II_1 -factor as the latter's metric is always negative definite.

Note that for M with separable predual, $U(M)$ isometrically embeds naturally in \mathbb{S}^∞ , the Hilbert sphere in ℓ^2 . The space \mathbb{S}^∞ is the “Hilbertian Urysohn sphere” in the sense described in [Nguyen Van Thé 2010, Section 1.4.2].

It is well worth pointing out the following proposition, which is an immediate consequence of Ozawa's fundamental result [2004] on the nonexistence of a universal, separable II_1 -factor.

Proposition 4.1. *For any separable II_1 -factor M , $U(M)$ is not universal among all \mathbb{Z}_4 -metric spaces which embed (as \mathbb{Z}_4 -metric spaces) in \mathbb{S}^∞ .*

Proof. Suppose, towards a contradiction, that there is a II_1 -factor M for which $U(M)$ is universal among all \mathbb{Z}_4 -metric spaces which embed in \mathcal{S}^∞ . In particular, for any II_1 -factor N with separable predual, $U(N)$ isometrically embeds in $U(M)$ in a way which commutes with the action of i . Since this embedding respects the inner product, it is not hard to see it must extend to an isometric embedding $L^2(N) \rightarrow L^2(M)$ which takes N into M contractively. Thus, as above, there is a unital injective $*$ -homomorphism $N \hookrightarrow pMp \oplus ((1-p)M(1-p))^{op}$, whence N embeds in either M or M^{op} since N is a factor. However, this would contradict the fact that there is no separable universal II_1 -factor [Ozawa 2004] (pick $M \star M^{op}$). \square

Question 4.2. Can $U(M)$ ever be universal among all metric spaces which embed in \mathcal{S}^∞ ?

Proposition 4.1 is good evidence that the answer to the previous question is no. We remark that a positive answer to the previous question would be equivalent to demonstrating the existence of a separable II_1 -factor for which there is an isometric embedding $\mathcal{S}^\infty \hookrightarrow U(M)$. We currently do not know whether \mathcal{S}^∞ embeds isometrically in the unitary group of *any* II_1 -factor. The existence of such an embedding ought to have striking consequences as the following proposition, which is similar in spirit, demonstrates.

Proposition 4.3. *Suppose M is a separable II_1 -factor belonging to the class \mathcal{K}_{op} . Further suppose that, for each n , the n -dimensional complex spheres \mathbb{S}^n isometrically embed in $U(M)$ with respect to the natural \mathbb{Z}_4 -actions. Then M is a locally universal II_1 -factor; that is, every separable II_1 -factor embeds into an ultrapower of M . In particular, if, for each n , the n -dimensional complex spheres \mathbb{S}^n isometrically embed in $U(\mathcal{R})$ with respect to the natural \mathbb{Z}_4 -actions, then Connes' embedding problem has a positive answer.*

Proof. Suppose that M satisfies the assumption of the proposition and let N be a II_1 -factor. Let F be any finite subset of $U(N)$. Then choosing an orthogonal projection P onto a suitably large finite-dimensional subspace so that $\|P(u)\| > 1 - \varepsilon$ for all $u \in F \cup iF$, we can correct to an (effective in) ε -almost \mathbb{Z}_4 -embedding of F into some \mathbb{S}^n , and therefore also in $U(M)$. But \mathbb{Z}_4 -embeddings preserve inner products, whence pairs of inner products in F can be modeled arbitrarily well in $U(M)$. As above, Kirchberg's argument shows that N embeds in $M^{\mathcal{U}}$. \square

We now remark how our main result recasts Kirchberg's characterization of \mathcal{R}^ω -embeddability in a game-theoretical light. Let (A, tr) be an arbitrary tracial C^* -algebra which we view as a normed space with respect to the 2-norm. To introduce a bit of terminology, we say that a subspace $E \subset A$ is ε -almost representable in \mathcal{R} if there exists a subspace $F \subset \mathcal{R}$ and a linear bijection $T : E \rightarrow F$ so that $\|T\|, \|T^{-1}\| \leq 1 + \varepsilon$ and $T(E \cap (A)_1) \subset_\varepsilon F \cap (\mathcal{R})_1$. Then by [Kirchberg

1993, Proposition 4.6], A is \mathcal{R}^ω -embeddable if and only if for every $\varepsilon > 0$, every finite-dimensional subspace of A is ε -representable in \mathcal{R} .

Let us introduce the following “one-sided, one-round game” $\mathfrak{G}_{\mathcal{R}}(n, \varepsilon)$ for which the winning condition is that, for all $u_1, \dots, u_n \in U(A)$ which are linearly independent, there exist n unitaries $v_1, \dots, v_n \in U(\mathcal{R})$ so that the map

$$T : \text{span}\{u_1, \dots, u_n\} \rightarrow \text{span}\{v_1, \dots, v_n\}$$

defined by $T(u_i) = v_i$ satisfies $\|T\|, \|T^{-1}\| \leq 1 + \varepsilon$.

Proposition 4.4. *There is a constant $N = N(n, \varepsilon)$ so that every n -dimensional subspace E of any tracial C^* -algebra (A, tr) is ε -almost representable in \mathcal{R} if $\mathfrak{G}_{\mathcal{R}}(N, \varepsilon/4)$ is winnable.*

Proof. We first claim that there is a uniform constant $K(n, \varepsilon)$ so that for every n -dimensional subspace $E \subset A$ of any tracial C^* -algebra (A, tr) there exists a set of unitaries $\bar{u} = \{u_1, \dots, u_l\} \subset U(A)$ with $l \leq K$ so that every element of $E \cap (A)_1$ is ε -approximated in 2-norm by a convex combination of elements of \bar{u} .

Indeed, choose an $(\varepsilon/2)$ -net $x_1, \dots, x_m \in E \cap (A)_1$. The cardinality of such a net is bounded in particular by the $(\varepsilon/4)$ -covering number of the unit ball in ℓ_n^2 . We may perturb each x_i so that $\|x_i\| < 1 - \varepsilon/4$ and still have an ε -net for $E \cap (A)_1$. By the main result of [Popa 1981], there is a constant C depending only on ε so that each x_i is a convex combination of at most C unitaries in $U(A)$, whence the claim follows.

We next claim that if A is infinite-dimensional and if $E \subset A$ is a finite-dimensional subspace, then for every $\varepsilon > 0$ and $u \in U(A)$, there exists $u' \in U(A)$ with $\|u - u'\|_2 < \varepsilon$ and so that u' is linearly independent from E . To see this, let $P_E : L^2(A) \rightarrow E$ be the orthogonal projection onto E . By the Kaplansky density theorem, we have that $U(A)$ is 2-norm dense in $U(A'')$. Since $M := A'' \subset \mathcal{B}(L^2(A, \text{tr}))$ is infinite-dimensional, it contains a diffuse abelian subalgebra. Therefore, there is a projection $p \in M$ with trace $\text{tr}(p) = 1 - \varepsilon^2/2$ and a sequence of unitaries $v_n \in U(M)$ so that $v_n \rightarrow p$ weakly. Since P_E is a finite-rank operator, we thus have that $P_E(uv_n) \rightarrow P_E(up)$ strongly, whence

$$\|P_E(uv_n)\|_2 \rightarrow \|P_E(up)\|_2 \leq \|p\|_2 = \sqrt{1 - \varepsilon^2/2}.$$

It is now easy to see that choosing n sufficiently large and $u' \in U(A)$ sufficiently close to uv_n works.

We now can proceed with the proof of the proposition. Let

$$E = \text{span}\{u_1, \dots, u_n\} \subset A.$$

(Every n -dimensional subspace of a C^* -algebra is a subspace of a space spanned by at most $4n$ unitaries, so we may assume this is the case without loss of generality.) By the previous claims, we can extend u_1, \dots, u_n to $u_1, \dots, u_n, u_{n+1}, \dots, u_s$

($s \leq n + K(n, \varepsilon)$) to a complete collection of linearly independent unitaries so that all elements in $E \cap (A)_1$ are 2ε -approximated in 2-norm by a convex combination of unitaries in the collection. If $\mathfrak{G}_R(s, \varepsilon/4)$ is winnable, then it is easy to check that for $S = T|_E$, we have that $S(E \cap (A)_1) \subseteq_\varepsilon S(E) \cap (\mathcal{R})_1$, and we are done. \square

Problem 4.5. Let $C \subset \ell_n^2$ be a convex subset of the unit ball in n -dimensional Hilbert space. For every $\varepsilon > 0$, does there exist a II_1 -factor M so that (ℓ_n^2, C) is ε -represented in M ? Can one always choose a locally universal II_1 -factor (in the sense of [Farah et al. 2014b]) or even \mathcal{R} ?

References

- [Ben Yaacov 2008] I. Ben Yaacov, “Continuous first order logic for unbounded metric structures”, *J. Math. Log.* **8**:2 (2008), 197–223. [MR 2011m:03065](#) [Zbl 1191.03026](#)
- [Ben Yaacov et al. 2008] I. Ben Yaacov, A. Berenstein, C. W. Henson, and A. Usvyatsov, “Model theory for metric structures”, pp. 315–427 in *Model theory with applications to algebra and analysis, II*, edited by Z. Chatzidakis et al., London Math. Soc. Lecture Note Ser. **350**, Cambridge Univ. Press, Cambridge, 2008. [MR 2009j:03061](#) [Zbl 1233.03045](#)
- [Connes 1975] A. Connes, “Sur la classification des facteurs de type II”, *C. R. Acad. Sci. Paris Sér. A-B* **281**:1 (1975), Aii, A13–A15. [MR 51 #13706](#) [Zbl 0304.46042](#)
- [Dixmier and Lance 1969] J. Dixmier and E. C. Lance, “Deux nouveaux facteurs de type II_1 ”, *Invent. Math.* **7** (1969), 226–234. [MR 40 #1787](#) [Zbl 0174.18701](#)
- [Dye 1955] H. A. Dye, “On the geometry of projections in certain operator algebras”, *Ann. of Math.* (2) **61** (1955), 73–89. [MR 16,598a](#) [Zbl 0064.11002](#)
- [Ealy and Goldbring 2012] C. Ealy and I. Goldbring, “Thorn-forking in continuous logic”, *J. Symbolic Logic* **77**:1 (2012), 63–93. [MR 2951630](#) [Zbl 1257.03059](#)
- [Farah et al. 2013] I. Farah, B. Hart, and D. Sherman, “Model theory of operator algebras I: stability”, *Bull. Lond. Math. Soc.* **45**:4 (2013), 825–838. [MR 3081550](#) [Zbl 1295.03019](#)
- [Farah et al. 2014a] I. Farah, B. Hart, and D. Sherman, “Model theory of operator algebras II: model theory”, *Israel J. Math.* **201**:1 (2014), 477–505. [MR 3265292](#) [Zbl 1301.03037](#)
- [Farah et al. 2014b] I. Farah, B. Hart, and D. Sherman, “Model theory of operator algebras III: elementary equivalence and II_1 factors”, *Bull. Lond. Math. Soc.* **46**:3 (2014), 609–628. [MR 3210717](#) [Zbl 1303.46049](#)
- [Hanche-Olsen and Størmer 1984] H. Hanche-Olsen and E. Størmer, *Jordan operator algebras*, Monographs and Studies in Mathematics **21**, Pitman (Advanced Publishing Program), Boston, 1984. [MR 86a:46092](#) [Zbl 0561.46031](#)
- [Hart 2012] B. Hart, “Continuous model theory and its applications”, lecture notes, 2012, available at <http://ms.mcmaster.ca/~bradd/courses/math712/index.html>.
- [Heinrich and Henson 1986] S. Heinrich and C. W. Henson, “Banach space model theory, II: Isomorphic equivalence”, *Math. Nachr.* **125** (1986), 301–317. [MR 87j:46139](#) [Zbl 0633.46076](#)
- [Henson and Iovino 2002] C. W. Henson and J. Iovino, “Ultraproducts in analysis”, pp. 1–110 in *Analysis and logic* (Mons, 1997), edited by C. Finet and C. Michaux, London Math. Soc. Lecture Note Ser. **262**, Cambridge Univ. Press, 2002. [MR 2004d:03001](#) [Zbl 1026.46007](#)
- [Henson and Moore 1983] C. W. Henson and L. C. Moore, Jr., “Nonstandard analysis and the theory of Banach spaces”, pp. 27–112 in *Nonstandard analysis — recent developments* (Victoria, B.C., 1980), Lecture Notes in Math. **983**, Springer, Berlin, 1983. [MR 85f:46033](#) [Zbl 0511.46070](#)

- [Kirchberg 1993] E. Kirchberg, “On nonseparable extensions, tensor products and exactness of group C^* -algebras”, *Invent. Math.* **112**:3 (1993), 449–489. MR 94d:46058 Zbl 0803.46071
- [Nguyen Van Thé 2010] L. Nguyen Van Thé, “Structural Ramsey theory of metric spaces and topological dynamics of isometry groups”, *Mem. Amer. Math. Soc.* **206**:968 (2010), x+140. MR 2011k:03096 Zbl 1203.05159
- [Ozawa 2004] N. Ozawa, “There is no separable universal II_1 -factor”, *Proc. Amer. Math. Soc.* **132**:2 (2004), 487–490. MR 2004h:46074 Zbl 1041.46045
- [Popa 1981] S. Popa, “On the Russo–Dye theorem”, *Michigan Math. J.* **28**:3 (1981), 311–315. MR 82j:46074 Zbl 0499.46037
- [Takesaki 2003] M. Takesaki, *Operator Algebras and Non-commutative Geometry, 8: Theory of operator algebras, III*, Encyclopaedia of Mathematical Sciences **127**, Springer, Berlin, 2003. MR 2004g:46080 Zbl 1059.46032

Received August 19, 2014. Revised March 2, 2015.

ISAAC GOLDBRING
DEPARTMENT OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE
UNIVERSITY OF ILLINOIS AT CHICAGO
SCIENCE AND ENGINEERING OFFICES M/C 249
851 SOUTH MORGAN STREET
CHICAGO, IL 60607-7045
UNITED STATES
isaac@math.uic.edu

THOMAS SINCLAIR
DEPARTMENT OF MATHEMATICS
PURDUE UNIVERSITY
150 N UNIVERSITY ST.
WEST LAFAYETTE, IN 47907-2067
UNITED STATES
tsincla@purdue.edu

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 278 No. 1 November 2015

Growth tight actions	1
GOULNARA N. ARZHANTSEVA, CHRISTOPHER H. CASHEN and JING TAO	
A flag structure on a cusped hyperbolic 3-manifold	51
ELISHA FALBEL and RAFAEL SANTOS THEBALDI	
A new upper bound for the Dirac operators on hypersurfaces	79
NICOLAS GINOUX, GEORGES HABIB and SIMON RAULOT	
Games and elementary equivalence of II_1 -factors	103
ISAAC GOLDBRING and THOMAS SINCLAIR	
Grossberg–Karshon twisted cubes and hesitant walk avoidance	119
MEGUMI HARADA and EUNJEONG LEE	
Gamma factors of distinguished representations of $\text{GL}_n(\mathbb{C})$	137
ALEXANDER KEMARSKY	
The W -entropy formula for the Witten Laplacian on manifolds with time dependent metrics and potentials	173
SONGZI LI and XIANG-DONG LI	
A diagrammatic categorification of the affine q -Schur algebra $\hat{S}(n, n)$ for $n \geq 3$	201
MARCO MACKAAY and ANNE-LAURE THIEL	
Showing distinctness of surface links by taking 2-dimensional braids	235
INASA NAKAMURA	
Correction to Modular L -values of cubic level	253
ANDREW KNIGHTLY and CHARLES LI	