

*Pacific
Journal of
Mathematics*

**CORRECTION TO
MODULAR L -VALUES OF CUBIC LEVEL**

ANDREW KNIGHTLY AND CHARLES LI

CORRECTION TO MODULAR L -VALUES OF CUBIC LEVEL

ANDREW KNIGHTLY AND CHARLES LI

Volume **260:2** (2012), 527–563

In the paper in question, equation (2-23) is incorrect, and hence so are equations (3-6) and (3-9). We correct the statements in this note. All other statements in the paper, including the main theorems, are unaffected.

In [Knightly and Li 2012], equation (2-23) was quoted from an early draft of [Knightly and Li 2015], which at the time contained an error. It should read

$$(2-23) \quad f_2^\sigma(zg) = \frac{(p+1)\zeta}{2\omega_p(zd)} \sum_{w \in (\mathbb{Z}/p\mathbb{Z})^*} \overline{\omega_p(w)} \theta_p \left(\frac{-\frac{c}{a}w - \frac{tb}{d}w^{-1}}{p} \right)$$

for $g = \begin{pmatrix} c & dp^{-2} \\ ap & b \end{pmatrix} \in \begin{pmatrix} \mathbb{Z}_p & (1/p^2)\mathbb{Z}_p^* \\ p\mathbb{Z}_p^* & \mathbb{Z}_p \end{pmatrix}$. It is a twisted Kloosterman sum.

Proposition 0.1 (Corrected Proposition 3.4). *Let $\delta = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, so that*

$$J_\delta(s, f^\sigma) = \int_{\mathbb{Q}_p^*} \int_{\mathbb{Q}_p} f^\sigma \left(\begin{pmatrix} 0 & -y \\ 1 & x \end{pmatrix} \right) \overline{\theta_p(rx)} dx \chi_p(y) |y|_p^{k/2-s} d^*y.$$

Then

$$(3-6) \quad J_\delta(s, f^\sigma) = J_\delta(s, f_2^\sigma) = \begin{cases} \frac{(p^3)^{k/2-s} p(p+1)\omega_p(-rp^2)}{2\zeta \chi_p(p^3)} & \text{if } p \nmid r, \\ 0 & \text{otherwise,} \end{cases}$$

and

$$(3-7) \quad I_\delta(s)_p = 0.$$

Proof. By (2-22), the matrix $\begin{pmatrix} 0 & -y \\ 1 & x \end{pmatrix}$ never belongs to $\text{Supp}(f_1^\sigma)$, so $J_\delta(s, f^\sigma) = J_\delta(s, f_2^\sigma)$. Note that $\begin{pmatrix} 0 & -y \\ 1 & x \end{pmatrix} \in \text{Supp}(f_2^\sigma)$ if and only if

$$\begin{pmatrix} 0 & -py \\ p & px \end{pmatrix} \in \begin{pmatrix} \mathbb{Z}_p & p^{-2}\mathbb{Z}_p^* \\ p\mathbb{Z}_p^* & \mathbb{Z}_p \end{pmatrix}.$$

MSC2010: primary 11F41; secondary 11F70.

Keywords: L-functions, relative trace formula, supercuspidal representations, Maass forms.

In this case, we may write $y = -p^{-3}u$ for $u \in \mathbb{Z}_p^*$, and $x' = px \in \mathbb{Z}_p$. Then $dx' = p^{-1}dx$, and dropping the $'$ from the notation, we have

$$\begin{aligned} & J_\delta(s, f_2^\sigma) \\ &= p \int_{\mathbb{Z}_p^*} \int_{\mathbb{Z}_p} f_2^\sigma \left(\begin{pmatrix} p^{-1} & \\ & p^{-1} \end{pmatrix} \begin{pmatrix} 0 & p^{-2}u \\ p & x \end{pmatrix} \right) \theta_p \left(\frac{-rx}{p} \right) dx \chi_p(p^{-3})(p^3)^{k/2-s} d^*u \\ &= \frac{(p^3)^{k/2-s} p(p+1)\zeta \omega_p(p)}{2\chi_p(p^3)} \sum_{w \in (\mathbb{Z}/p\mathbb{Z})^*} \overline{\omega_p(w)} \int_{\mathbb{Z}_p^*} \overline{\omega_p(u)} \int_{\mathbb{Z}_p} \theta_p \left(\frac{-tx}{p} \right) \theta_p \left(\frac{-rx}{p} \right) dx d^*u \end{aligned}$$

by (2-23). Replacing u by $(-uw)^{-1}$, the above is equal to

$$\frac{(p^3)^{k/2-s} p(p+1)\zeta \omega_p(p)}{2\chi_p(p^3)} \sum_{w \in (\mathbb{Z}/p\mathbb{Z})^*} \overline{\omega_p(w)} \int_{\mathbb{Z}_p^*} \omega_p(-uw) \int_{\mathbb{Z}_p} \theta_p \left(\frac{(tu-r)x}{p} \right) dx d^*u.$$

Observe that w is eliminated, and the sum over w contributes $p-1$. Furthermore,

$$\int_{\mathbb{Z}_p} \theta_p \left(\frac{(tu-r)x}{p} \right) dx = \begin{cases} 1 & \text{if } u \in t^{-1}r + p\mathbb{Z}_p, \\ 0 & \text{otherwise.} \end{cases}$$

In particular, it vanishes if $p \mid r$. Assuming $p \nmid r$,

$$\begin{aligned} J_\delta(s, f^\sigma) &= \frac{(p^3)^{k/2-s} p(p+1)\zeta \omega_p(p)}{2\chi_p(p^3)} (p-1) \int_{t^{-1}r+p\mathbb{Z}_p} \omega_p(-u) d^*u \\ &= \frac{(p^3)^{k/2-s} p(p+1)\zeta \omega_p(p)}{2\chi_p(p^3)} \omega_p(-t^{-1}r) \end{aligned}$$

since the coset has multiplicative measure $1/(p-1)$. Equality (3-6) now follows, using the fact that

$$\frac{\zeta \omega_p(p)}{\omega_p(t)} = \frac{\zeta \omega_p(p)^2}{\omega_p(pt)} = \frac{\omega_p(p)^2}{\zeta}.$$

For fixed t , summing (3-6) over $\pm\zeta$ gives 0. Thus $I_\delta(s)_p = \sum_\sigma J_\delta(s, f^\sigma) = 0$. \square

Proposition 0.2 (Corrected Proposition 3.5). *For $a \in \mathbb{Q}^*$, let $\delta_a = \begin{pmatrix} a & -1 \\ 1 & 0 \end{pmatrix}$, so that*

$$J_{\delta_a}(s, f^\sigma) = \int_{\mathbb{Q}_p^*} \int_{\mathbb{Q}_p} f^\sigma \left(\begin{pmatrix} ya & y(xa-1) \\ 1 & x \end{pmatrix} \right) \overline{\theta_p(rx)} dx \chi_p(y) |y|_p^{k/2-s} d^*y.$$

Then $J_{\delta_a}(s, f_1^\sigma)$ vanishes unless $a \in p^2\mathbb{Z}_p$ and $p \nmid r$. In this case, writing $a = p^a a_0$ for $a_0 \in \mathbb{Z}_p^* \cap \mathbb{Q}^*$, we have

$$(3-8) \quad J_{\delta_a}(s, f_1^\sigma) = \begin{cases} \frac{|a|_p^{2s-k} p(p+1) \omega_p(p^{a_p})}{2\chi_p(a^2)} \theta_p\left(\frac{ta}{rp^3} - \frac{r}{a}\right) & \text{if } a_0 \equiv 1 \pmod{p}, \\ 0 & \text{otherwise.} \end{cases}$$

The integral $J_{\delta_a}(s, f_2^\sigma)$ vanishes unless $a \in p^2\mathbb{Z}_p$. For such a ,

$$(3-9) \quad J_{\delta_a}(s, f_2^\sigma) = \begin{cases} \frac{(p^3)^{k/2-s} p(p+1) \omega_p(-p^2r)}{2\chi_p(p^3)\zeta} \theta_p\left(-\frac{ta}{rp^3}\right) & \text{if } p \nmid r, \\ 0 & \text{otherwise.} \end{cases}$$

Finally, $I_{\delta_a}(s)_p$ vanishes unless $p \nmid r$ and $a = p^{a_p} a_0$ for $a_p \geq 2$ and $a_0 \equiv 1 \pmod{p\mathbb{Z}_p}$. If these conditions are satisfied, then

$$(3-10) \quad I_{\delta_a}(s)_p = \frac{|a|_p^{2s-k} p(p+1) \omega_p(p^{a_p}) \theta_p\left(-\frac{t}{a}\right)}{\chi_p(a^2)} \Delta_p(a) \quad \text{for } \Delta_p(a) = \begin{cases} p-1 & \text{if } a_p > 2, \\ -1 & \text{if } a_p = 2. \end{cases}$$

Remark. Only (3-9) differs from the original statement.

Proof of (3-9). Consider

$$J_{\delta_a}(s, f_2^\sigma) = \int_{\mathbb{Q}_p^*} \int_{\mathbb{Q}_p} f_2^\sigma \left(\begin{pmatrix} ya & y(xa-1) \\ 1 & x \end{pmatrix} \right) \overline{\theta_p(rx)} dx \chi_p(y) |y|_p^{k/2-s} d^*y.$$

By (2-23), the integrand is nonzero precisely when

$$\begin{pmatrix} p & \\ & p \end{pmatrix} \begin{pmatrix} ya & y(xa-1) \\ 1 & x \end{pmatrix} = \begin{pmatrix} pya & py(xa-1) \\ p & px \end{pmatrix} \in \begin{pmatrix} \mathbb{Z}_p & p^{-2}\mathbb{Z}_p^* \\ p\mathbb{Z}_p^* & \mathbb{Z}_p \end{pmatrix}.$$

Taking the determinant, this says, in particular, that $p^2y \in p^{-1}\mathbb{Z}_p^*$, so we may write $y = u/p^3$ for $u \in \mathbb{Z}_p^*$. Replacing px by x , we have

$$J_{\delta_a}(s, f_2^\sigma) = \frac{(p^3)^{k/2-s} p \omega_p(p)}{\chi_p(p^3)} \int_{\mathbb{Z}_p^*} \int_{\mathbb{Z}_p} f_2^\sigma \left(\begin{pmatrix} \frac{ua}{p^2} & \frac{u}{p^2} \left(\frac{xa}{p} - 1\right) \\ p & x \end{pmatrix} \right) \theta_p\left(\frac{-rx}{p}\right) dx d^*u.$$

From the upper left entry, the integrand is nonzero only if $a_p \geq 2$. Assuming the latter, we also have $xa/p - 1 \in \mathbb{Z}_p^*$, so the upper right entry belongs to $p^{-2}\mathbb{Z}_p^*$ as required. Hence by (2-23), the above is equal to

$$\frac{(p^3)^{k/2-s} p(p+1)\zeta \omega_p(p)}{2\chi_p(p^3)} \sum_{w \in (\mathbb{Z}/p\mathbb{Z})^*} \overline{\omega_p(w)} \int_{\mathbb{Z}_p^*} \overline{\omega_p(u)} \theta_p\left(-\frac{uaw}{p^3}\right) \int_{\mathbb{Z}_p} \overline{\omega_p\left(\frac{xa}{p} - 1\right)} \theta_p\left(\frac{-txu^{-1}\left(\frac{xa}{p} - 1\right)^{-1} w^{-1}}{p}\right) \theta_p\left(\frac{-rx}{p}\right) dx d^*u.$$

Note that $\omega_p(xa/p - 1) = \omega_p(-1)$ since $p^2 \mid a$. For the same reason,

$$\theta_p\left(\frac{-txu^{-1}\left(\frac{xa}{p} - 1\right)^{-1}w^{-1}}{p}\right) = \theta_p\left(\frac{tu^{-1}w^{-1}x}{p}\right).$$

Therefore the above integral over \mathbb{Z}_p equals

$$\omega_p(-1) \int_{\mathbb{Z}_p} \theta_p\left(\frac{(-r + tu^{-1}w^{-1})x}{p}\right) dx = \begin{cases} \omega_p(-1) & \text{if } u \in tr^{-1}w^{-1} + p\mathbb{Z}_p, \\ 0 & \text{otherwise.} \end{cases}$$

In particular, $J_{\delta_a}(s, f_2^\sigma) = 0$ if $p \mid r$. Assuming $p \nmid r$, $J_{\delta_a}(s, f_2^\sigma)$ equals

$$\begin{aligned} & \frac{(p^3)^{k/2-s} p(p+1)\zeta \omega_p(-p)}{2\chi_p(p^3)} \sum_{w \in (\mathbb{Z}/p\mathbb{Z})^*} \overline{\omega_p(w)} \int_{tr^{-1}w^{-1} + p\mathbb{Z}_p} \overline{\omega_p(u)} \theta_p\left(-\frac{uaw}{p^3}\right) d^*u \\ &= \frac{(p^3)^{k/2-s} p(p+1)\zeta \omega_p(-p)}{2\chi_p(p^3)} \sum_{w \in (\mathbb{Z}/p\mathbb{Z})^*} \overline{\omega_p(w)} \overline{\omega_p(tr^{-1}w^{-1})} \theta_p\left(-\frac{tr^{-1}a}{p^3}\right) \frac{1}{p-1} \\ &= \frac{(p^3)^{k/2-s} p(p+1)\zeta \omega_p(-p)}{2\chi_p(p^3)} \overline{\omega_p(tr^{-1})} \theta_p\left(-\frac{tr^{-1}a}{p^3}\right). \end{aligned}$$

Equation (3-9) follows upon using $\zeta \omega_p(p)/\omega_p(t) = \omega_p(p)^2/\zeta$. □

References

[Knightly and Li 2012] A. Knightly and C. Li, “Modular L -values of cubic level”, *Pacific J. Math.* **260**:2 (2012), 527–563. MR 3001804 Zbl 1286.11069

[Knightly and Li 2015] A. Knightly and C. Li, “Simple supercuspidal representations of $GL(n)$ ”, *Taiwanese J. Math.* **19**:4 (2015), 995–1029.

Received March 1, 2015.

ANDREW KNIGHTLY
 DEPARTMENT OF MATHEMATICS AND STATISTICS
 UNIVERSITY OF MAINE
 NEVILLE HALL
 ORONO, ME 04469-5752
 UNITED STATES
knightly@math.umaine.edu

CHARLES LI
 DEPARTMENT OF MATHEMATICS
 THE CHINESE UNIVERSITY OF HONG KONG
 SHATIN
 HONG KONG
charles@charlesli.org

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 278 No. 1 November 2015

Growth tight actions	1
GOULNARA N. ARZHANTSEVA, CHRISTOPHER H. CASHEN and JING TAO	
A flag structure on a cusped hyperbolic 3-manifold	51
ELISHA FALBEL and RAFAEL SANTOS THEBALDI	
A new upper bound for the Dirac operators on hypersurfaces	79
NICOLAS GINOUX, GEORGES HABIB and SIMON RAULOT	
Games and elementary equivalence of II_1 -factors	103
ISAAC GOLDBRING and THOMAS SINCLAIR	
Grossberg–Karshon twisted cubes and hesitant walk avoidance	119
MEGUMI HARADA and EUNJEONG LEE	
Gamma factors of distinguished representations of $\text{GL}_n(\mathbb{C})$	137
ALEXANDER KEMARSKY	
The W -entropy formula for the Witten Laplacian on manifolds with time dependent metrics and potentials	173
SONGZI LI and XIANG-DONG LI	
A diagrammatic categorification of the affine q -Schur algebra $\hat{S}(n, n)$ for $n \geq 3$	201
MARCO MACKAAY and ANNE-LAURE THIEL	
Showing distinctness of surface links by taking 2-dimensional braids	235
INASA NAKAMURA	
Correction to Modular L -values of cubic level	253
ANDREW KNIGHTLY and CHARLES LI	