

*Pacific
Journal of
Mathematics*

ON EXTENSIONS OF ALGEBRAIC GROUPS
WITH FINITE QUOTIENT

MICHEL BRION

ON EXTENSIONS OF ALGEBRAIC GROUPS WITH FINITE QUOTIENT

MICHEL BRION

Consider an exact sequence of group schemes of finite type over a field k ,

$$1 \longrightarrow N \longrightarrow G \longrightarrow Q \longrightarrow 1,$$

where Q is finite. We show that Q lifts to a finite subgroup scheme F of G ; if Q is étale and k is perfect, then F may be chosen étale as well. As applications, we obtain generalizations of classical results of Arima, Chevalley, and Rosenlicht to possibly nonconnected algebraic groups. We also show that every homogeneous space under such a group has a projective equivariant compactification.

1. Introduction

Consider an extension of algebraic groups, that is, an exact sequence of group schemes of finite type over a field,

$$(1) \quad 1 \longrightarrow N \longrightarrow G \xrightarrow{f} Q \longrightarrow 1.$$

Such an extension is generally not split, i.e., f admits no section which is a morphism of group schemes. In this note, we obtain the existence of a splitting in a weaker sense, for extensions with finite quotient group:

Theorem 1.1. *Let G be an algebraic group over a field k , and N a normal subgroup of G with G/N finite. Then there exists a finite subgroup F of G such that $G = N \cdot F$.*

Here $N \cdot F$ denotes, as in [SGA 3₁ 1970, VIA.5.3.3], the quotient of the semidirect product $N \rtimes F$ by $N \cap F$ embedded as a normal subgroup via $x \mapsto (x, x^{-1})$. If G/N is étale and k is perfect, then the subgroup F may be chosen étale as well. But this fails over any imperfect field k , see Remark 3.3 for details.

In the case where G is smooth and k is perfect, Theorem 1.1 was known to Borel and Serre, and they presented a proof over an algebraically closed field of characteristic 0 (see [Borel and Serre 1964, Lemma 5.11 and footnote on p. 152]). That result was also obtained by Platonov [1966, Lemma 4.14] for smooth linear algebraic groups over perfect fields. In the latter setting, an effective version of

MSC2010: primary 14L15; secondary 14L30, 20G15.

Keywords: algebraic groups, finite quotients, extensions, equivariant compactifications.

[Theorem 1.1](#) has been obtained recently by Lucchini Arteche [[2015a](#), [Theorem 1.1](#)]; see [[Lucchini Arteche 2015b](#), [Proposition 1.1](#)]; [Chernousov et al. 2008](#), p. 473; [Lötscher et al. 2013](#), [Lemma 5.3](#)] for earlier results in this direction.

Returning to an extension (1) with an arbitrary quotient Q , one may ask whether there exists a subgroup H of G such that $G = N \cdot H$ and $N \cap H$ is finite (when Q is finite, the latter condition is equivalent to the finiteness of H). We then say that (1) is *quasisplit*, and H is a *quasicomplement* of N in G , with *defect group* $N \cap H$.

When Q is smooth and N is an abelian variety, every extension (1) is quasisplit (as shown by Rosenlicht [[1956](#), [Theorem 14](#)]; see [[Milne 2013](#), [Section 2](#)] for a modern proof). The same holds when Q is reductive (i.e., Q is smooth and affine, and the radical of $Q_{\bar{k}}$ is a torus), N is arbitrary and $\text{char}(k) = 0$, as we will show in [Corollary 4.8](#). On the other hand, the group G of unipotent 3×3 matrices sits in a central extension

$$1 \longrightarrow \mathbb{G}_a \longrightarrow G \longrightarrow \mathbb{G}_a^2 \longrightarrow 1,$$

which is not quasisplit. It would be interesting to determine which classes of groups N , Q yield quasisplit extensions. Another natural problem is to bound the defect group in terms of N and Q . The proof of [Theorem 1.1](#) yields some information in that direction; see [Remark 3.4](#), and [[Lucchini Arteche 2015a](#)] for an alternative approach via nonabelian Galois cohomology.

This article is organized as follows. In [Section 2](#), we begin the proof of [Theorem 1.1](#) with a succession of reductions to the case where $Q = G/N$ is étale and N is a smooth connected unipotent group, a torus, or an abelian variety. In [Section 3](#), we show that every class of extensions (1) is torsion in that setting ([Lemma 3.1](#)); this quickly implies [Theorem 1.1](#). [Section 4](#) presents some applications of [Theorem 1.1](#) to the structure of algebraic groups: we obtain analogues of classical results of Chevalley, Rosenlicht and Arima on smooth connected algebraic groups (see [[Rosenlicht 1956; 1961; Arima 1960](#)]) and of Mostow [[1956](#)] on linear algebraic groups in characteristic 0. Finally, we show that every homogeneous space under an algebraic group admits a projective equivariant compactification; this result seems to have been unrecorded so far. It is well known that any such homogeneous space is quasiprojective (see [[Raynaud 1970](#), [Corollary VI.2.6](#)]); also, the existence of equivariant compactifications of certain homogeneous spaces having no separable point at infinity has attracted recent interest (see, e.g., [[Gabber 2012; Gabber et al. 2014](#)]).

2. Proof of [Theorem 1.1](#): some reductions

We first fix notation and conventions which will be used throughout this article. We consider schemes and their morphisms over a field k , and choose an algebraic

closure \bar{k} . Given a scheme X and an extension K/k of fields, we denote by X_K the K -scheme obtained from X by the base change $\text{Spec}(K) \rightarrow \text{Spec}(k)$.

We use mostly [SGA 3_I 1970; SGA 3_{II} 1970; SGA 3_{III} 1970], and occasionally [Demazure and Gabriel 1970], as references for group schemes. Given such a group scheme G , we denote by $e_G \in G(k)$ the neutral element, and by G^0 the neutral component of G , with quotient map $\pi : G \rightarrow G/G^0 = \pi_0(G)$. The group law of G is denoted by $\mu : G \times G \rightarrow G$, $(x, y) \mapsto xy$.

Throughout this section, we consider an extension (1) and a subgroup F of G . Then the map

$$\nu : N \rtimes F \longrightarrow G, \quad (x, y) \longmapsto xy$$

is a morphism of group schemes with kernel $N \cap F$, embedded in $N \times F$ via $x \mapsto (x, x^{-1})$. Thus, ν factors through a morphism of group schemes

$$\iota : N \cdot F \longrightarrow G.$$

Also, the composition $F \rightarrow G \rightarrow G/N$ factors through a morphism of group schemes

$$i : F/(N \cap F) \longrightarrow G/N.$$

By [SGA 3_I 1970, VIA.5.4], ι and i are closed immersions of group schemes.

Lemma 2.1. *The following conditions are equivalent:*

- (i) ι is an isomorphism.
- (ii) i is an isomorphism.
- (iii) ν is faithfully flat.
- (iv) For any scheme S and $g \in G(S)$, there exists a faithfully flat morphism of finite presentation $f : S' \rightarrow S$ and $x \in N(S')$, $y \in F(S')$ such that $g = xy$ in $G(S')$.

When G/N is smooth, these conditions are equivalent to:

- (v) $G(\bar{k}) = N(\bar{k})F(\bar{k})$.

Proof. (i) \Leftrightarrow (ii): Recall from [SGA 3_I 1970, VIA.5.5.3] that i factors through an isomorphism $F/(N \cap F) \rightarrow (N \cdot F)/N$. Thus, we obtain a commutative diagram

$$\begin{array}{ccc} N \cdot F & \xrightarrow{\iota} & G \\ \varphi \downarrow & & f \downarrow \\ F/(N \cap F) & \xrightarrow{i} & G/N \end{array}$$

where both vertical arrows are N -torsors for the action of N by right multiplication. As a consequence, this diagram is cartesian. In particular, i is an isomorphism if and only if so is ι ; this yields the desired equivalence.

(i) \Rightarrow (iii): Since ν is identified with the quotient map of $N \rtimes F$ by $N \cap F$, the assertion follows from [SGA 3₁ 1970, VIA.3.2].

(iii) \Rightarrow (iv): This follows by forming the cartesian square

$$\begin{array}{ccc} S' & \longrightarrow & S \\ \downarrow & & \downarrow g \\ N \rtimes F & \xrightarrow{\nu} & G \end{array}$$

and observing that ν is of finite presentation, since the schemes G , N and F are of finite type.

(iv) \Rightarrow (i): By our assumption applied to the identity map $G \rightarrow G$, there exists a scheme S' and morphisms $x : S' \rightarrow N$, $y : S' \rightarrow F$ such that the morphism $\nu \circ (x \times y) : N \rtimes F \rightarrow G$ is faithfully flat of finite presentation. As a consequence, the morphism of structure sheaves $\mathbb{O}_G \rightarrow \nu_*(x \times y)_*(\mathbb{O}_{S'})$ is injective. Thus, so are $\mathbb{O}_G \rightarrow \nu_*(\mathbb{O}_{N \rtimes F})$, and hence $\mathbb{O}_G \rightarrow i_*(\mathbb{O}_{N \cdot F})$. Since i is a closed immersion, it must be an isomorphism.

(ii) \Leftrightarrow (v): When G/N is smooth, i is an isomorphism if and only if it is surjective on \bar{k} -rational points. Since $(G/N)(\bar{k}) = G(\bar{k})/N(\bar{k})$ and likewise for $F/(N \cap F)$, this yields the desired equivalence. \square

We assume from now on that the quotient group Q in the extension (1) is finite.

Lemma 2.2. *If the exact sequence $1 \rightarrow H^0 \rightarrow H \rightarrow \pi_0(H) \rightarrow 1$ is quasisplit for any smooth algebraic group H such that $\dim(H) = \dim(G)$, then (1) is quasisplit as well.*

Proof. Consider first the case where G is smooth. Then Q is étale, and hence N contains G^0 . By our assumption, there exists a finite subgroup $F \subset G$ such that $G = G^0 \cdot F$. In view of Lemma 2.1 (iv), it follows that $G = N \cdot F$.

If $\text{char}(k) = 0$, then the proof is completed as every algebraic group is smooth (see, e.g., [SGA 3₁ 1970, VIB.1.6.1]). So we may assume that $\text{char}(k) = p > 0$. Consider the n -fold relative Frobenius morphism

$$F_G^n : G \longrightarrow G^{(p^n)}$$

and its kernel G_n . Then F_G^n is finite and bijective, so that G_n is an infinitesimal normal subgroup of G . Moreover, the quotient G/G_n is smooth for $n \gg 0$ (see [SGA 3₁ 1970, VIIA.8.3]). We may thus choose n so that G/G_n and N/N_n are smooth. The composition $N \rightarrow G \rightarrow G/G_n$ factors through a closed immersion of group schemes $N/N_n \rightarrow G/G_n$ by [SGA 3₁ 1970, VIA.5.4] again. Moreover, the image of N/N_n is a normal subgroup of G/G_n , as follows, e.g., from our

smoothness assumption and [SGA 3_I 1970, VIB.7.3]. This yields an exact sequence

$$1 \longrightarrow N/N_n \longrightarrow G/G_n \longrightarrow Q' \longrightarrow 1,$$

where Q' is a quotient of Q and hence is finite; moreover, $\dim(G/G_n) = \dim(G)$. By our assumption and the first step, there exists a finite subgroup F' of G/G_n such that $G/G_n = (N/N_n) \cdot F'$. In view of [SGA 3_I 1970, VIA.5.3.1], there exists a unique subgroup F of G containing G_n such that $F/G_n = F'$; then F is finite as well.

We check that $G = N \cdot F$ by using Lemma 2.1 (iv) again. Let S be a scheme, and $g \in G(S)$. Then there exists a faithfully flat morphism of finite presentation $S' \rightarrow S$ and $x' \in (N/N_n)(S')$, $y' \in F'(S')$ such that $F_G^n(g) = x' y'$ in $(G/G_n)(S')$. Moreover, there exists a faithfully flat morphism of finite presentation $S'' \rightarrow S'$ and $x'' \in N(S'')$, $y'' \in F(S'')$ such that $F_G^n(x'') = x'$ and $F_G^n(y'') = y'$. Then $y''^{-1} x''^{-1} g \in G_n(S'')$, and hence $g \in N(S'')F(S'')$, since F contains G_n . \square

Remark 2.3. With the notation of the proof of Lemma 2.2, there is an exact sequence of quasicomplements

$$1 \longrightarrow G_n \longrightarrow F \longrightarrow F' \longrightarrow 1.$$

When $N = G^0$, so that $G_n \subset N$, we also have an exact sequence of defect groups

$$1 \longrightarrow G_n \longrightarrow N \cap F \longrightarrow (N/G_n) \cap F' \longrightarrow 1.$$

By Lemma 2.2, it suffices to prove Theorem 1.1 when G is smooth and $N = G^0$, so that $Q = \pi_0(G)$. We may thus choose a maximal torus T of G (see [SGA 3_{II} 1970, XIV.1.1]). Then the normalizer $N_G(T)$ and the centralizer $Z_G(T)$ are (represented by) subgroups of G (see [SGA 3_I 1970, VIB.6.2.5]). Moreover, $N_G(T)$ is smooth by [SGA 3_{II} 1970, XI.2.4]. We now gather further properties of $N_G(T)$:

Lemma 2.4. (i) $G = G^0 \cdot N_G(T)$.

(ii) $N_G(T)^0 = Z_{G^0}(T)$.

(iii) We have an exact sequence $1 \rightarrow W(G^0, T) \rightarrow \pi_0(N_G(T)) \rightarrow \pi_0(G) \rightarrow 1$, where $W(G^0, T) := N_{G^0}(T)/Z_{G^0}(T) = \pi_0(N_{G^0}(T))$ denotes the Weyl group.

Proof. (i) By Lemma 2.1 (v), it suffices to show that $G(\bar{k}) = G^0(\bar{k})N_G(T)(\bar{k})$. Let $x \in G(\bar{k})$, then xTx^{-1} is a maximal torus of $G^0(\bar{k})$, and hence $xTx^{-1} = yTy^{-1}$ for some $y \in G^0(\bar{k})$. Thus, $x \in yN_G(T)(\bar{k})$, which yields the assertion.

(ii) We may assume that k is algebraically closed and G is connected (since $N_G(T)^0 = N_{G^0}(T)^0$). Then $Z_G(T)$ is a Cartan subgroup of G , and hence equals its connected normalizer by [SGA 3_{II} 1970, XII.6.6].

(iii) By (i), the natural map $N_G(T)/N_{G^0}(T) \rightarrow \pi_0(G)$ is an isomorphism. Combined with (ii), this yields the statement. \square

Remark 2.5. If $N_G(T) = N_G(T)^0 \cdot F$ for some subgroup $F \subset N_G(T)$, then by Lemmas 2.1 and 2.4, $G = G^0 \cdot F$. Moreover, the commutative diagram of exact sequences

$$\begin{array}{ccccccc} 1 & \longrightarrow & N_G(T)^0 \cap F & \longrightarrow & F & \longrightarrow & \pi_0(N_G(T)) \longrightarrow 1 \\ & & \downarrow & & \text{id} \downarrow & & \downarrow \\ 1 & \longrightarrow & G^0 \cap F & \longrightarrow & F & \longrightarrow & \pi_0(G) \longrightarrow 1 \end{array}$$

together with Lemma 2.4 yields the exact sequence

$$1 \longrightarrow Z_{G^0}(T) \cap F \longrightarrow G^0 \cap F \longrightarrow W(G^0, T) \longrightarrow 1.$$

In view of Lemma 2.1 (iv) and Lemma 2.4 (i), it suffices to prove Theorem 1.1 under the additional assumption that T is normal in G . Then T is central in G^0 , and hence G_k^0 is nilpotent by [SGA 3_{II} 1970, XII.6.7]. It follows that G^0 is nilpotent, in view of [SGA 3_I 1970, VIB.8.3]. To obtain further reductions, we will use the following:

Lemma 2.6. *Let N' be a normal subgroup of G contained in N . Assume that the resulting exact sequence $1 \rightarrow N/N' \rightarrow G/N' \rightarrow Q \rightarrow 1$ is quasisplit, and that any exact sequence of algebraic groups $1 \rightarrow N' \rightarrow G' \rightarrow Q' \rightarrow 1$, where Q' is finite, is quasisplit as well. Then (1) is quasisplit.*

Proof. By assumption, there exists a finite subgroup F' of G/N' for which $G/N' = (N/N') \cdot F'$. Denote by G' the subgroup of G containing N' such that $G'/N' = F'$. By assumption again, there is a finite subgroup F of G' containing N' such that $G' = N' \cdot F$. We check that $G = N \cdot F$ using Lemma 2.1 (iv). Let S be a scheme, and $g \in G(S)$; denote by $f' : G \rightarrow G/N'$ the quotient map. Then there exists a faithfully flat morphism of finite presentation $S' \rightarrow S$ and $x \in (N/N')(S')$, $y \in F'(S')$ such that $f'(g) = xy$ in $(G/N')(S')$. Moreover, there exists a faithfully flat morphism of finite presentation $S'' \rightarrow S'$ and $z \in N(S'')$, $w \in G'(S'')$ such that $f'(z) = x$ and $f'(w) = y$. Then $w^{-1}z^{-1}g \in N'(S'')$, and hence $g \in N(S'')G'(S'')$, as G' contains N' . This shows that $G = N \cdot G' = N \cdot (N' \cdot F)$. We conclude by observing that $N \cdot (N' \cdot F) = N \cdot F$, in view of Lemma 2.1 (iv) again. \square

Remark 2.7. With the notation of the proof of Lemma 2.6, we have an exact sequence $1 \rightarrow N' \rightarrow G' = N' \cdot F \rightarrow F' \rightarrow 1$, and hence an exact sequence of quasicomplements

$$1 \longrightarrow N' \cap F \longrightarrow F \longrightarrow F' \longrightarrow 1.$$

Moreover, we obtain an exact sequence $1 \rightarrow N' \rightarrow N \cap G' \rightarrow (N/N') \cap F' \rightarrow 1$, by using [SGA 3_I 1970, VIA.5.3.1]. Since $N \cap G' = N \cap (N' \cdot F) = N' \cdot (N \cap F)$, where the latter equality follows from Lemma 2.1 (iv), this yields an exact sequence

of defect groups

$$1 \longrightarrow N' \cap F \longrightarrow N \cap F \longrightarrow (N/N') \cap F' \longrightarrow 1.$$

Next, we show that it suffices to prove [Theorem 1.1](#) when G^0 is assumed in addition to be *commutative*.

We argue by induction on the dimension of G (assumed to be smooth, with G^0 nilpotent). If $\dim(G) = 1$, then G^0 is either a k -form of \mathbb{G}_a or \mathbb{G}_m , or an elliptic curve; in particular, G^0 is commutative. In higher dimensions, the derived subgroup $D(G^0)$ is a smooth, connected normal subgroup of G contained in G^0 , and the quotient $G^0/D(G^0)$ is commutative of positive dimension (see [\[SGA 3_I 1970, VIB.7.8, 8.3\]](#)). Moreover, $G/D(G^0)$ is smooth, and $\pi_0(G/D(G^0)) = \pi_0(G)$. By the induction assumption, it follows that the exact sequence

$$1 \longrightarrow G^0/D(G^0) \longrightarrow G/D(G^0) \longrightarrow \pi_0(G) \longrightarrow 1$$

is quasisplit. Also, every exact sequence $1 \rightarrow D(G^0) \rightarrow G' \rightarrow Q' \rightarrow 1$, where Q' is finite, is quasisplit, by the induction assumption again together with [Lemma 2.2](#). Thus, [Lemma 2.6](#) yields the desired reduction.

We now show that we may further assume G^0 to be *a torus, a smooth connected commutative unipotent group, or an abelian variety*.

Indeed, we have an exact sequence of commutative algebraic groups

$$1 \longrightarrow T \longrightarrow G^0 \longrightarrow H \longrightarrow 1,$$

where T is the maximal torus of G^0 , and H is smooth and connected. Moreover, we have an exact sequence

$$1 \longrightarrow H_1 \longrightarrow H \longrightarrow H_2 \longrightarrow 1,$$

where H_1 is a smooth connected affine algebraic group, and H_2 is a pseudoabelian variety in the sense of [\[Totaro 2013\]](#), i.e., H_2 has no nontrivial smooth connected affine normal subgroup. Since H_1 contains no nontrivial torus, it is unipotent; also, H_2 is an extension of a smooth connected unipotent group by an abelian variety A , in view of [\[Totaro 2013, Theorem 2.1\]](#). Note that T is a normal subgroup of G (the largest subtorus). Also, H_1 is a normal subgroup of G/T (the largest smooth connected affine normal subgroup of the neutral component), and A is a normal subgroup of $(G/T)/H_1$ as well (the largest abelian subvariety). Thus, arguing by induction on the dimension as in the preceding step, with $D(G^0)$ replaced successively by T , H_1 and A , yields our reduction.

When G^0 is unipotent and $\text{char}(k) = p > 0$, we may further assume that G^0 is *killed by p* . Indeed, by [\[SGA 3_{II} 1970, XVII.3.9\]](#), there exists a composition series $\{e_G\} = G_0 \subset G_1 \subset \cdots \subset G_n = G^0$ such that each G_i is normal in G , and each

quotient G_i/G_{i-1} is a k -form of some $(\mathbb{G}_a)^{r_i}$; in particular, G_i/G_{i-1} is killed by p . Our final reduction follows by induction on n .

3. Proof of [Theorem 1.1](#): extensions by commutative groups

In this section, we consider smooth algebraic groups Q, N such that Q is finite and N is commutative. Given an extension [\(1\)](#), the action of G on N by conjugation factors through an action of Q by group automorphisms, which we denote by $(x, y) \mapsto y^x$, where $x \in Q$ and $y \in N$. Recall that the isomorphism classes of such extensions with a prescribed Q -action on N form a commutative (abstract) group, which we denote by $\text{Ext}^1(Q, N)$; see [\[SGA 3_{II} 1970, XVII.App. I\]](#) (and [\[Demazure and Gabriel 1970, III.6.1\]](#) for the setting of extensions of group sheaves).

Lemma 3.1. *With the above notation and assumptions, the group $\text{Ext}^1(Q, N)$ is torsion.*

Proof. Any extension [\(1\)](#) yields an N -torsor over Q for the étale topology, since Q is finite and étale. This defines a map $\tau : \text{Ext}^1(Q, N) \rightarrow H_{\text{ét}}^1(Q, N)$, which is a group homomorphism (indeed, the sum of any two extensions is obtained by taking their direct product, pulling back under the diagonal map $Q \rightarrow Q \times Q$, and pushing forward under the multiplication $N \times N \rightarrow N$; and the sum of any two torsors is obtained by the analogous operations). The kernel of τ consists of those classes of extensions that admit a section (which is a morphism of schemes). In view of [\[SGA 3_{II} 1970, XVII.App. I.3.1\]](#), this yields an exact sequence

$$0 \longrightarrow HH^2(Q, N) \longrightarrow \text{Ext}^1(Q, N) \xrightarrow{\tau} H_{\text{ét}}^1(Q, N),$$

where HH^i stands for Hochschild cohomology (denoted by H^i in [\[SGA 3_I 1970; SGA 3_{II} 1970; SGA 3_{III} 1970\]](#), and by H_0^i in [\[Demazure and Gabriel 1970\]](#)). Moreover, the group $H_{\text{ét}}^1(Q, N)$ is torsion (as follows, e.g., from [\[Rosenlicht 1956, Theorem 14\]](#)), and $HH^2(Q, N)$ is killed by the order of Q , as a special case of [\[SGA 3_{II} 1970, XVII.5.2.4\]](#). \square

Remark 3.2. The above argument yields that $\text{Ext}^1(Q, N)$ is killed by md if Q is finite étale of order m , and N is a torus split by an extension of k of degree d . Indeed, we just saw that $HH^2(Q, N)$ is killed by m ; also, $H_{\text{ét}}^1(Q, N)$ is a direct sum of groups of the form $H_{\text{ét}}^1(\text{Spec}(k'), N)$ for finite separable extensions k' of k , and these groups are killed by d . This yields a slight generalization of [\[Lucchini Arteché 2015b, Proposition 1.1\]](#), via a different approach.

End of the proof of [Theorem 1.1](#). Recall from our reductions in [Section 2](#) that we may assume G^0 to be a smooth commutative unipotent group, a torus, or an abelian variety. We will rather denote G^0 by N , and $\pi_0(G)$ by Q .

We first assume in addition that $\text{char}(k) = 0$ if N is unipotent. Then the n -th power map

$$n_N : N \longrightarrow N, \quad x \longmapsto x^n$$

is an isogeny for any positive integer n . Consider an extension (1) and denote by γ its class in $\text{Ext}^1(Q, N)$. By Lemma 3.1, we may choose n so that $n\gamma = 0$. Also, $n\gamma = (n_N)_*(\gamma)$ (the pushout of γ by n_N); moreover, the exact sequence of commutative algebraic groups

$$1 \longrightarrow N[n] \longrightarrow N \xrightarrow{n_N} N \longrightarrow 1$$

yields an exact sequence

$$\text{Ext}^1(Q, N[n]) \longrightarrow \text{Ext}^1(Q, N) \xrightarrow{(n_N)_*} \text{Ext}^1(Q, N)$$

due to [SGA 3_{II} 1970, XVII.App. I.2.1]. Thus, there is a class $\gamma' \in \text{Ext}^1(Q, N[n])$ with pushout γ , i.e., we have a commutative diagram of extensions

$$\begin{array}{ccccccc} 1 & \longrightarrow & N[n] & \longrightarrow & G' & \longrightarrow & Q \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow \\ 1 & \longrightarrow & N & \longrightarrow & G & \longrightarrow & Q \longrightarrow 1, \end{array}$$

where the square on the left is cartesian. It follows that G' is a finite subgroup of G , and $G = N \cdot G'$.

Next, we consider the remaining case, where N is unipotent and $\text{char}(k) = p > 0$. In view of our final reduction at the end of Section 2, we may further assume that N is killed by p . Then there exists an étale isogeny $N \rightarrow N_1$, where N_1 is a vector group (see [Conrad et al. 2015, Lemma B.1.10]). This yields another commutative diagram of extensions

$$\begin{array}{ccccccc} 1 & \longrightarrow & N & \longrightarrow & G & \longrightarrow & Q \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow \\ 1 & \longrightarrow & N_1 & \longrightarrow & G_1 & \longrightarrow & Q \longrightarrow 1. \end{array}$$

Assume that there exists a finite subgroup F_1 of G_1 such that $G_1 = N_1 \cdot F_1$. Let F be the pullback of F_1 to G ; then F is a finite subgroup, and one checks that $G = N \cdot F$ using Lemma 2.1 (iv). Thus, we may finally assume that N is a vector group.

Under that assumption, the N -torsor $G \rightarrow Q$ is trivial, since Q is affine. Thus, we may choose a section $s : Q \rightarrow G$. Also, we may choose a finite Galois extension of fields K/k such that Q_K is constant. Then s yields a section $s_K : Q_K \rightarrow G_K$, equivariant under the Galois group $\Gamma_K := \text{Gal}(K/k)$. So we may view $G(K)$ as the set of the $y s(x)$, where $y \in N(K)$ and $x \in Q(K)$, with multiplication

$$y s(x) y' s(x') = y y'^x c(x, x') s(xx'),$$

where $c \in Z^2(Q(K), N(K))^{\Gamma_K}$. Consider the (abstract) subgroup $H \subset N(K)$ generated by the $c(x', x'')^x$, where $x, x', x'' \in Q(K)$. Then H is finite, since $N(K)$ is killed by p and $Q(K)$ is finite. Moreover, $Hs(Q(K))$ is a subgroup of $G(K)$, in view of the above formula for the multiplication. Clearly, $Hs(Q(K))$ is finite and stable under Γ_K ; thus, it corresponds to a finite (algebraic) subgroup G' of G . Also, we obtain as above that $G = N \cdot G'$. This completes the proof of [Theorem 1.1](#). \square

Remark 3.3. If k is perfect, then the subgroup F as in [Theorem 1.1](#) may be chosen étale. Indeed, the reduced subscheme F_{red} is then a subgroup by [[SGA 3_I 1970](#), VIA.0.2]. Moreover, $G(\bar{k}) = G^0(\bar{k})F_{\text{red}}(\bar{k})$, and hence $G = G^0 \cdot F_{\text{red}}$ in view of [Lemma 2.1 \(v\)](#).

In contrast, when k is imperfect, there exists a finite group G admitting no étale subgroup F such that $G = G^0 \cdot F$. Consider for example (as in [[SGA 3_I 1970](#), VIA.1.3.2]) the subgroup G of $\mathbb{G}_{a,k}$ defined by the additive polynomial $X^{p^2} - tX^p$, where $p := \text{char}(k)$ and $t \in k \setminus k^p$. Then G has order p^2 and G^0 has order p . If $G = G^0 \cdot F$ with F étale, then $G^0 \cap F$ is trivial. Thus, $G \cong G^0 \rtimes F$ and F has order p . Let $K := k(t^{1/p})$. Then F_K is contained in $(G_K)_{\text{red}}$, which is the subgroup of $\mathbb{G}_{a,K}$ defined by the additive polynomial $X^p - t^{1/p}X$. By counting dimensions, it follows that $F_K = (G_K)_{\text{red}}$, which yields a contradiction as $(G_K)_{\text{red}}$ is not defined over k .

Remark 3.4. One may obtain information on the defect group $N \cap F$ by examining the steps in the proof of [Theorem 1.1](#) and combining [Remarks 2.3](#), [2.5](#) and [2.7](#). For instance, if G is smooth, then $N \cap F$ is an extension of the Weyl group $W(G^0, T)$ by the nilpotent group $Z_{G^0}(T) \cap F$, where T is a maximal torus of G . If $\text{char}(k) = 0$ (so that G is smooth), then $Z_{G^0}(T) \cap F$ is commutative. Indeed, $Z_{G^0}(T)$ is a connected nilpotent algebraic group, and hence an extension of a semiabelian variety S by a connected unipotent algebraic group U . Thus, $U \cap F$ is trivial, and hence $Z_{G^0}(T) \cap F$ is isomorphic to a subgroup of S .

Remark 3.5. When k is finite, [Theorem 1.1](#) follows readily from our first reduction step ([Lemma 2.2](#)) together with a theorem of Lang [[1956](#), [Theorem 2](#)]. More specifically, let H be a smooth algebraic group and choose representatives x_1, \dots, x_m of the orbits of the Galois group $\Gamma := \text{Gal}(\bar{k}/k)$ in $\pi_0(H)(\bar{k})$. Denote by $\Gamma_i \subset \Gamma$ the isotropy group of x_i and set $k_i := \bar{k}^{\Gamma_i}$ for $i = 1, \dots, m$. Then $x_i \in \pi_0(H)(k_i)$, and hence the fiber $\pi_{k_i}^{-1}(x_i)$ (a torsor under $H_{k_i}^0$) contains a k_i -rational point. Consider the subfield

$$K := \prod_{i=1}^n k_i \subset \bar{k}.$$

Then the finite étale group scheme $\pi_0(H)_K$ is constant, and π is surjective on K -rational points. Thus, $\pi_0(H)$ has a quasicomplement in H : the finite étale group scheme corresponding to the constant, Γ -stable subgroup scheme $H(K)$ of H_K .

4. Some applications

We first recall two classical results on the structure of algebraic groups. The first one is the affinization theorem (see [Demazure and Gabriel 1970, III.3.8] and also [SGA 3_I 1970, VIB.12.2]): any algebraic group G has a smallest normal subgroup H such that G/H is affine. Moreover, H is smooth, connected and contained in the center of G^0 ; we have $\mathbb{C}(H) = k$ (such an algebraic group is called *antiaffine*) and $\mathbb{C}(G/H) = \mathbb{C}(G)$.

Consequently, H is the fiber at e_G of the affinization morphism $G \rightarrow \text{Spec } \mathbb{C}(G)$; moreover, the formation of H commutes with arbitrary field extensions. Also, note that H is the largest antiaffine subgroup of G ; we will denote H by G_{ant} . The structure of antiaffine groups is described in [Brion 2009] and [Sancho and Sancho 2009].

The second structure result is a version of a theorem of Chevalley, due to Raynaud [1970, Lemma IX.2.7] (see also [Bosch et al. 1990, 9.2 Theorem 1]): any connected algebraic group G has a smallest affine normal subgroup N such that G/N is an abelian variety. Moreover, N is connected; if G is smooth and k is perfect, then N is smooth as well. We will denote N by G_{aff} .

We will also need the following observation:

Lemma 4.1. *Let G be an algebraic group, and N a normal subgroup. Then the quotient map $f : G \rightarrow G/N$ yields an isomorphism $G_{\text{ant}}/(G_{\text{ant}} \cap N) \cong (G/N)_{\text{ant}}$.*

Proof. We have a closed immersion of group schemes $G_{\text{ant}}/(G_{\text{ant}} \cap N) \rightarrow G/N$; moreover, $G_{\text{ant}}/(G_{\text{ant}} \cap N)$ is antiaffine. So we obtain a closed immersion of commutative group schemes $i : G_{\text{ant}}/(G_{\text{ant}} \cap N) \rightarrow (G/N)_{\text{ant}}$. The cokernel of i is antiaffine, as a quotient of $(G/N)_{\text{ant}}$. Also, this cokernel is a subgroup of $(G/N)/(G_{\text{ant}}/(G_{\text{ant}} \cap N))$, which is a quotient of G/G_{ant} . Since the latter is affine, it follows that $\text{Coker}(i)$ is affine as well, by using [SGA 3_I 1970, VIB.11.17]. Thus, $\text{Coker}(i)$ is trivial, i.e., i is an isomorphism. \square

We now obtain a further version of Chevalley's structure theorem, for possibly nonconnected algebraic groups:

Theorem 4.2. *Any algebraic group G has a smallest affine normal subgroup N such that G/N is proper. Moreover, N is connected.*

Proof. It suffices to show that G admits an affine normal subgroup N such that G/N is proper. Indeed, given another such subgroup N' , the natural map

$$G/(N \cap N') \longrightarrow G/N \times G/N'$$

is a closed immersion, and hence $G/(N \cap N')$ is proper. Taking for N a minimal such subgroup, it follows that N is the smallest one. Moreover, the natural morphism $G/N^0 \rightarrow G/N$ is finite, since it is a torsor under the finite group N/N^0 (see

[SGA 3_I 1970, VIA.5.3.2]). As a consequence, G/N^0 is proper; hence $N = N^0$ by the minimality assumption. Thus, N is connected.

Also, we may reduce to the case where G is smooth by using the relative Frobenius morphism as in the proof of Lemma 2.2.

If in addition G is connected, then we just take $N = G_{\text{aff}}$. In the general case, we consider the (possibly nonnormal) subgroup $H := (G^0)_{\text{aff}}$; then the homogeneous space G/H is proper, since G/G^0 is finite and G^0/H is proper. As a consequence, the automorphism functor of G/H is represented by a group scheme $\text{Aut}_{G/H}$, locally of finite type; in particular, the neutral component $\text{Aut}_{G/H}^0$ is an algebraic group (see [Matsumura and Oort 1967, Theorem 3.7]). The action of G by left multiplication on G/H yields a morphism of group schemes

$$\varphi : G \longrightarrow \text{Aut}_{G/H}.$$

The kernel N of φ is a closed subscheme of H , and hence is affine. To complete the proof, it suffices to show that G/N is proper. In turn, it suffices to check that $(G/N)^0$ is proper. Since $(G/N)^0 \cong G^0/(G^0 \cap N)$, and $G^0 \cap N$ is the kernel of the restriction $G^0 \rightarrow \text{Aut}_{G/H}^0$, we are reduced to showing that $\text{Aut}_{G/H}^0$ is proper (by using [SGA 3_I 1970, VIA.5.4.1] again).

We claim that $\text{Aut}_{G/H}^0$ is an abelian variety. Indeed, $(G/H)_{\bar{k}}$ is a finite disjoint union of copies of $(G^0/H)_{\bar{k}}$, which is an abelian variety. Also, the natural morphism $A \rightarrow \text{Aut}_A^0$ is an isomorphism for any abelian variety A . Thus, $(\text{Aut}_{G/H}^0)_{\bar{k}}$ is an abelian variety (a product of copies of $(G^0/H)_{\bar{k}}$); this yields our claim, and completes the proof. \square

Remark 4.3. The formation of G_{aff} (for a connected group scheme G) commutes with separable algebraic field extensions, as follows from a standard argument of Galois descent. But this formation does not commute with purely inseparable field extensions, in view of [SGA 3_{II} 1970, XVII.C.5].

Likewise, the formation of N as in Theorem 4.2 commutes with separable algebraic field extensions. As a consequence, $N = (G^0)_{\text{aff}}$ for any smooth group scheme G (since $(G^0)_{\text{aff}}$ is invariant under any automorphism of G , and hence is a normal subgroup of G when k is separably closed). In particular, if k is perfect and G is smooth, then N is smooth as well.

For an arbitrary group scheme G , we may have $N \neq (G^0)_{\text{aff}}$, e.g., when G is infinitesimal: then N is trivial, while $(G^0)_{\text{aff}} = G$.

We do not know if the formations of G_{aff} and N commute with arbitrary separable field extensions.

The structure of proper algebraic groups is easily described as follows:

Proposition 4.4. *Given a proper algebraic group G , there exists an abelian variety A , a finite group F equipped with an action $F \rightarrow \text{Aut}_A$ and a normal subgroup*

$D \subset F$ such that D acts faithfully on A by translations and $G \cong (A \rtimes F)/D$, where D is embedded in $A \rtimes F$ via $x \mapsto (x, x^{-1})$. Moreover, $A = G_{\text{ant}}$ and $F/D \cong G/G_{\text{ant}}$ are uniquely determined by G . Finally, G is smooth if and only if F/D is étale.

Proof. Note that G_{ant} is a smooth connected proper algebraic group, and hence an abelian variety. Moreover, the quotient group G/G_{ant} is affine and proper, hence finite. By [Theorem 1.1](#), there exists a finite subgroup $F \subset G$ such that $G = G_{\text{ant}} \cdot F$. In particular, $G \cong (F \rtimes G_{\text{ant}})/(F \cap G_{\text{ant}})$; this implies the existence assertion. For the uniqueness, just note that $\mathcal{O}(G) \cong \mathcal{O}(G/A) \cong \mathcal{O}(F/D)$, and this identifies the affinization morphism to the natural homomorphism $G \rightarrow F/D$, with kernel A .

If G is smooth, then so is $G/A \cong F/D$; as F/D is finite, it must be étale. Since the homomorphism $G \rightarrow F/D$ is smooth, the converse holds as well. \square

Remark 4.5. The simplest examples of proper algebraic groups are the semidirect products $G = A \rtimes F$, where F is a finite group acting on the abelian variety A . If this action is nontrivial (for example, if A is nontrivial and F is the constant group $\mathbb{Z}/2\mathbb{Z}$ acting via $x \mapsto x^{\pm 1}$), then every morphism of algebraic groups $f : G \rightarrow H$, where H is connected, has a nontrivial kernel. (Otherwise, A is contained in the center of G by the affinization theorem.) This yields examples of algebraic groups which admit no faithful representation in a connected algebraic group.

Remark 4.6. With the notation and assumptions of [Proposition 4.4](#), consider a subgroup $H \subset G$ and the homogeneous space $X := G/H$. Then there exists an abelian variety B quotient of A , a subgroup $I \subset F$ containing D , and a faithful homomorphism $I \rightarrow \text{Aut}_B$ such that the scheme X is isomorphic to the associated fiber bundle $F \times^I B$. Moreover, the schemes F/I and B are uniquely determined by X , and X is smooth if and only if F/I is étale.

Indeed, let $K := A \cdot H$, then $X \cong G \times^K K/H \cong F \times^I K/H$, where $I := F \cap K$. Moreover, $K/H \cong A/(A \cap H)$ is an abelian variety. This shows the existence assertion; those on uniqueness and smoothness are checked as in the proof of [Proposition 4.4](#).

Conversely, given a finite group F and a subgroup $I \subset F$ acting on an abelian variety B , the associated fiber bundle $F \times^I B$ exists (since it is the quotient of the projective scheme $F \times B$ by the finite group I), and is homogeneous whenever F/I is étale (since $(F \times^I B)_{\bar{k}}$ is just a disjoint union of copies of $B_{\bar{k}}$). We do not know how to characterize the homogeneity of $F \times^I B$ when the quotient F/I is arbitrary.

Returning to an arbitrary algebraic group G , we have the ‘‘Rosenlicht decomposition’’ $G = G_{\text{ant}} \cdot G_{\text{aff}}$ when G is smooth and connected (see, e.g., [\[Brion 2009\]](#)). We now extend this result to possibly nonconnected groups:

Theorem 4.7. *Let G be an algebraic group. Then there exists an affine subgroup H of G such that $G = G_{\text{ant}} \cdot H$. If G is smooth and k is perfect, then H may be chosen smooth.*

Proof. By [Theorem 4.2](#), we may choose an affine normal subgroup $N \subset G$ such that G/N is proper. In view of [Proposition 4.4](#), there exists a finite subgroup F of G/N such that $G/N = (G/N)_{\text{ant}} \cdot F$, and $(G/N)_{\text{ant}}$ is an abelian variety. Let H be the subgroup of G containing N such that $G/H = F$. Then H is affine, since it sits in an extension $1 \rightarrow N \rightarrow H \rightarrow F \rightarrow 1$. We check that $G = G_{\text{ant}} \cdot H$ by using [Lemma 2.1 \(iv\)](#). Let S be a scheme, and $g \in G(S)$. Denote by g' the image of g in $(G/N)(S)$. Then there exist a faithfully flat morphism of finite presentation $S' \rightarrow S$ and $x' \in (G/N)_{\text{ant}}(S')$, $y' \in F(S')$ such that $g' = x'y'$ in $(G/N)(S')$. Moreover, in view of [Lemma 4.1](#), x' lifts to some $x'' \in G_{\text{ant}}(S'')$, where $S'' \rightarrow S'$ is faithfully flat of finite presentation. So $gx''^{-1} \in G(S'')$ lifts y' , and hence $g \in G_{\text{ant}}(S'')H(S'')$.

If G is smooth and k is perfect, then N may be chosen smooth by [Remark 4.3](#); also, F may be chosen smooth by [Remark 3.3](#). Then H is smooth as well. \square

We now derive from [Theorem 4.7](#) a generalization of our main [Theorem 1.1](#), under the additional assumption of characteristic 0 (then reductivity is equivalent to linear reductivity, also known as full reducibility):

Corollary 4.8. *Every extension (1) with reductive quotient group Q is quasisplit when $\text{char}(k) = 0$.*

Proof. Choose an affine subgroup $H \subset G$ such that $G = G_{\text{ant}} \cdot H$ and denote by $R_u(H)$ its unipotent radical. By a result of Mostow [[1956](#), Theorem 6.1], H has a Levi subgroup, i.e., a fully reducible algebraic subgroup L such that $H = R_u(H) \rtimes L$. Note that $R_u(H)$ is normal in G , since it is normalized by H and centralized by G_{ant} . It follows that $G_{\text{ant}} \cdot R_u(H)$ is normal in G , and $G = (G_{\text{ant}} \cdot R_u(H)) \cdot L$. Also, note that the quotient map $f : G \rightarrow Q$ sends G_{ant} to e_Q (since Q is affine), and $R_u(H)$ to e_Q as well (since Q is reductive). It follows that the sequence

$$1 \longrightarrow N \cap L \longrightarrow L \xrightarrow{f} Q \longrightarrow 1$$

is exact, where $N = \text{Ker}(f)$. If $N \cap L$ has a quasicomplement H in L , then H is a quasicomplement to N in G (as follows, e.g., from [Lemma 2.1 \(v\)](#)). Thus, we may assume that G is *reductive*. Since every quasicomplement to N^0 in G is a quasicomplement to N , we may further assume that N is *connected*.

We have a canonical decomposition

$$G^0 = D(G^0) \cdot R(G^0),$$

where the derived subgroup $D(G^0)$ is semisimple, the radical $R(G^0)$ is a central torus, and $D(G^0) \cap R(G^0)$ is finite (see, e.g., [[SGA 3_{III} 1970](#), XXII.6.2.4]). Thus, $G = D(G^0) \cdot (R(G^0) \cdot F)$, where $F \subset G$ is a quasicomplement to G^0 . Likewise, $N = D(N) \cdot R(N)$, where $D(N) \subset D(G^0)$, $R(N) \subset R(G^0)$ and both are normal in G . Denote by S the neutral component of the centralizer of $D(N)$ in $D(G^0)$. Then S is a normal semisimple subgroup of G , and a quasicomplement to $D(N)$

in $D(G^0)$. If $R(N)$ admits a quasicomplement T in $R(G^0) \cdot F$, then one readily checks that $S \cdot T$ is a quasicomplement to N in G . As a consequence, we may replace G with $R(G^0) \cdot F$, and hence assume that G^0 is a *torus*.

Denote by $X^*(G^0)$ the character group of G_k^0 ; this is a free abelian group of finite rank, equipped with a continuous action of $F(\bar{k}) \rtimes \Gamma$, where Γ denotes the absolute Galois group of k . Moreover, we have a surjective homomorphism $\rho : X^*(G^0) \rightarrow X^*(N)$, equivariant for $F(\bar{k}) \rtimes \Gamma$. Thus, ρ splits over the rationals, and hence there exists a subgroup $\Lambda \subset X^*(G^0)$, stable by $F(\bar{k}) \rtimes \Gamma$, which is mapped isomorphically by ρ to a subgroup of finite index of $X^*(N)$. The quotient $X^*(G^0)/\Lambda$ corresponds to a subtorus $H \subset G^0$, normalized by G , which is a quasicomplement to N in G^0 . So $H \cdot F$ is the desired quasicomplement to N in G . \square

Remark 4.9. Corollary 4.8 does not extend to positive characteristics, due to the existence of groups without Levi subgroups (see [Conrad et al. 2015, Appendix A.6; McNinch 2010, Section 3.2]). As a specific example, when k is perfect of characteristic $p > 0$, there exists a nonsplit extension of algebraic groups

$$1 \longrightarrow V \longrightarrow G \xrightarrow{f} \mathrm{SL}_2 \longrightarrow 1,$$

where V is a vector group on which SL_2 acts linearly via the Frobenius twist of its adjoint representation. We show that this extension is not quasisplit. Otherwise, let H be a quasicomplement to N in G . Then so is the reduced neutral component of H , and hence we may assume that H is smooth and connected. The quotient map f restricts to an isogeny $H \rightarrow \mathrm{SL}_2$, and hence to an isomorphism. Thus, the above extension is split, a contradiction.

Next, we obtain an analogue of the Levi decomposition (see [Mostow 1956] again) for possibly nonlinear algebraic groups:

Corollary 4.10. *Let G be an algebraic group over a field of characteristic 0. Then $G = R \cdot S$, where $R \subset G$ is the largest connected solvable normal subgroup, and $S \subset G$ is an algebraic subgroup such that S^0 is semisimple; also, $R \cap S$ is finite.*

Proof. By a standard argument, G has a largest connected solvable normal subgroup R . The quotient G/R is affine, since $R \supset G_{\mathrm{ant}}$. Moreover, R/G_{ant} contains the radical of G/G_{ant} , and hence $(G/R)^0$ is semisimple. In particular, G/R is reductive. So Corollary 4.8 yields the existence of the quasicomplement S . \square

Remark 4.11. One may ask for a version of Corollary 4.10 in which the normal subgroup R is replaced with an analogue of the unipotent radical of a linear algebraic group, and the quasicomplement S is assumed to be reductive. But such a version would make little sense when G is an antiaffine semiabelian variety (for example, when G is the extension of an abelian variety A by \mathbb{G}_m , associated with an algebraically trivial line bundle of infinite order on A). Indeed, such a group G

has a largest connected reductive subgroup: its maximal torus, which admits no quasicomplement.

Also, recall that the radical R may admit no complement in G , e.g., when $G = \mathrm{GL}_n$ with $n \geq 2$.

Finally, one may also ask for the uniqueness of a minimal quasicomplement in [Corollary 4.10](#), up to conjugacy in $R(k)$ (as for Levi complements, see [\[Mostow 1956, Theorem 6.2\]](#)). But this fails when k is algebraically closed and G is the semidirect product of an abelian variety A with a group F of order 2. Denote by σ the involution of A induced by the nontrivial element of F ; then $R = A$, and the complements to R in G are exactly the subgroups generated by the involutions $x\sigma$ where $x \in A^{-\sigma}(k)$, i.e., $\sigma(x) = x^{-1}$. The action of $R(k)$ on complements is given by $yx\sigma y^{-1} = xy\sigma(y)^{-1}\sigma$; moreover, the homomorphism $A \rightarrow A^{-\sigma}$, $y \mapsto y\sigma(y)^{-1}$ is generally not surjective. This holds for example when $A = (B \times B)/C$, where B is a nontrivial abelian variety, C is the subgroup of $B \times B$ generated by (x_0, x_0) for some $x_0 \in B(k)$ of order 2, and σ arises from the involution $(x, y) \mapsto (y^{-1}, x^{-1})$ of $B \times B$; then $A^{-\sigma}$ has 2 connected components.

Another consequence of [Theorem 4.7](#) concerns the case where k is finite; then every antiaffine algebraic group is an abelian variety (see [\[Brion 2009, Proposition 2.2\]](#)). This yields readily:

Corollary 4.12. *Let G be an algebraic group over a finite field. Then G sits in an extension of algebraic groups*

$$1 \longrightarrow F \longrightarrow A \times H \longrightarrow G \longrightarrow 1,$$

where F is finite, A is an abelian variety, and H is affine. If G is smooth, then H may be chosen smooth as well.

Returning to an arbitrary base field, we finally obtain the existence of equivariant compactifications of homogeneous spaces:

Theorem 4.13. *Let G be an algebraic group, and H a closed subgroup. Then there exists a projective scheme X equipped with an action of G , and an open G -equivariant immersion $G/H \hookrightarrow X$ with schematically dense image.*

Proof. When G is affine, this follows from a theorem of Chevalley asserting that H is the stabilizer of a line L in a finite-dimensional G -module V (see [\[SGA 3, 1970, VIB.11.16\]](#)). Indeed, one may take for X the closure of the G -orbit of L in the projective space of lines of V ; then X satisfies the required properties in view of [\[Demazure and Gabriel 1970, III.3.5.2\]](#). Note that X is equipped with an ample G -linearized invertible sheaf.

When G is proper, the homogeneous space G/H is proper as well, and hence is projective by [\[Raynaud 1970, Corollary VI.2.6\]](#) (alternatively, this follows from the structure of X described in [Remark 4.5](#)).

In the general case, [Theorem 4.2](#) yields an affine normal subgroup N of G such that G/N is proper. Then $N \cdot H$ is a subgroup of G , and $G/(N \cdot H)$ is proper as well, hence projective. It suffices to show the existence of a projective scheme Y equipped with an action of $N \cdot H$, an open immersion $(N \cdot H)/H \rightarrow Y$ with schematically dense image, and a $N \cdot H$ -linearized ample line bundle: indeed, by [\[Mumford et al. 1994, Proposition 7.1\]](#) applied to the projection $G \times Y \rightarrow Y$ and the $N \cdot H$ -torsor $G \rightarrow G/(N \cdot H)$, the associated fiber bundle $G \times^{N \cdot H} Y$ yields the desired equivariant compactification. In view of Chevalley's theorem used in the first step, it suffices in turn to check that $N \cdot H$ acts on $(N \cdot H)/H$ via an affine quotient group; equivalently, $(N \cdot H)_{\text{ant}} \subset H$.

By [Lemma 4.1](#), $(N \cdot H)_{\text{ant}}$ is a quotient of $(N \rtimes H)_{\text{ant}}$. The latter is the fiber at the neutral element of the affinization morphism $N \rtimes H \rightarrow \text{Spec } \mathbb{C}(N \rtimes H)$. Also, $N \rtimes H \cong N \times H$ as schemes, N is affine and the affinization morphism commutes with products; thus, $(N \rtimes H)_{\text{ant}} = H_{\text{ant}}$. As a consequence, $(N \cdot H)_{\text{ant}} = H_{\text{ant}}$; this completes the proof. \square

Remark 4.14. If $\text{char}(k) = 0$, the equivariant compactification X of [Theorem 4.13](#) may be taken smooth, as follows from the existence of an equivariant desingularization (see [\[Kollár 2007, Proposition 3.9.1, Theorem 3.36\]](#)).

In arbitrary characteristics, X may be taken normal if G is smooth. Indeed, the G -action on any equivariant compactification X stabilizes the reduced subscheme X_{red} (since $G \times X_{\text{red}}$ is reduced), and lifts to an action on its normalization \tilde{X} (since $G \times \tilde{X}$ is normal). But the existence of regular compactifications (equivariant or not) is an open question.

Over any imperfect field k , there exist smooth connected algebraic groups G having no smooth compactification. Indeed, we may take for G the subgroup of $\mathbb{G}_a \times \mathbb{G}_a$ defined by $y^p - y - tx^p = 0$, where $p := \text{char}(k)$ and $t \in k \setminus k^p$. This is a smooth affine curve, and hence has a unique regular compactification X . One checks that X is the curve $(y^p - yz^{p-1} - tx^p = 0) \subset \mathbb{P}^2$, which is not smooth at its point at infinity.

Acknowledgements

I warmly thank Giancarlo Lucchini Arteché for very helpful e-mail discussions, and Zinovy Reichstein for his interest and for pointing out several references. Many thanks also to the referee for a careful reading of this note, and useful comments.

References

- [Arima 1960] S. Arima, “Commutative group varieties”, *J. Math. Soc. Japan* **12** (1960), 227–237. [MR 25 #79](#) [Zbl 0113.36307](#)
- [Borel and Serre 1964] A. Borel and J.-P. Serre, “Théorèmes de finitude en cohomologie galoisienne”, *Comment. Math. Helv.* **39** (1964), 111–164. [MR 31 #5870](#) [Zbl 0143.05901](#)

- [Bosch et al. 1990] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **21**, Springer, Berlin, 1990. MR 91i:14034 Zbl 0705.14001
- [Brion 2009] M. Brion, “Anti-affine algebraic groups”, *J. Algebra* **321**:3 (2009), 934–952. MR 2009j:14059 Zbl 1166.14029
- [Chernousov et al. 2008] V. Chernousov, P. Gille, and Z. Reichstein, “Reduction of structure for torsors over semilocal rings”, *Manuscripta Math.* **126**:4 (2008), 465–480. MR 2009f:14090 Zbl 1157.14028
- [Conrad et al. 2015] B. Conrad, O. Gabber, and G. Prasad, *Pseudo-reductive groups*, 2nd ed., New Mathematical Monographs **17**, Cambridge University Press, 2015. MR 2011k:20093 Zbl 1314.20037
- [Demazure and Gabriel 1970] M. Demazure and P. Gabriel, *Groupes algébriques, I: Géométrie algébrique, généralités, groupes commutatifs*, Masson, Paris, 1970. MR 46 #1800 Zbl 0203.23401
- [Gabber 2012] O. Gabber, “On pseudo-reductive groups and compactification theorems”, *Oberwolfach Reports* **9** (2012), 2371–2374. MR 3156730
- [Gabber et al. 2014] O. Gabber, P. Gille, and L. Moret-Bailly, “Fibrés principaux sur les corps valués Henséliens”, *Algebr. Geom.* **1**:5 (2014), 573–612. MR 3296806
- [Kollár 2007] J. Kollár, *Lectures on resolution of singularities*, Annals of Mathematics Studies **166**, Princeton University Press, 2007. MR 2008f:14026 Zbl 1113.14013
- [Lang 1956] S. Lang, “Algebraic groups over finite fields”, *Amer. J. Math.* **78** (1956), 555–563. MR 19,174a Zbl 0073.37901
- [Lötscher et al. 2013] R. Lötscher, M. MacDonald, A. Meyer, and Z. Reichstein, “Essential p -dimension of algebraic groups whose connected component is a torus”, *Algebra Number Theory* **7**:8 (2013), 1817–1840. MR 3134035 Zbl 1288.20061
- [Lucchini Arteché 2015a] G. Lucchini Arteché, “Extensions of algebraic groups with finite quotient and non-abelian 2-cohomology”, preprint, 2015. arXiv 1503.06582
- [Lucchini Arteché 2015b] G. Lucchini Arteché, “Groupe de Brauer non ramifié algébrique des espaces homogènes”, *Transform. Groups* **20**:2 (2015), 463–493. MR 3348564 Zbl 06458683
- [Matsumura and Oort 1967] H. Matsumura and F. Oort, “Representability of group functors, and automorphisms of algebraic schemes”, *Invent. Math.* **4** (1967), 1–25. MR 36 #181 Zbl 0173.22504
- [McNinch 2010] G. J. McNinch, “Levi decompositions of a linear algebraic group”, *Transform. Groups* **15**:4 (2010), 937–964. MR 2012b:20110 Zbl 1221.20034
- [Milne 2013] J. S. Milne, “A proof of the Barsotti–Chevalley theorem on algebraic groups”, preprint, 2013. arXiv 1311.6060v2
- [Mostow 1956] G. D. Mostow, “Fully reducible subgroups of algebraic groups”, *Amer. J. Math.* **78** (1956), 200–221. MR 19,1181f Zbl 0073.01603
- [Mumford et al. 1994] D. Mumford, J. Fogarty, and F. Kirwan, *Geometric invariant theory*, 3rd ed., Ergebnisse der Mathematik und ihrer Grenzgebiete (2) **34**, Springer, Berlin, 1994. MR 95m:14012 Zbl 0797.14004
- [Platonov 1966] V. P. Platonov, “Теория алгебраических линейных групп и периодические группы”, *Izv. Akad. Nauk SSSR Ser. Mat.* **30**:3 (1966), 573–620. Translated as “Theory of algebraic linear groups and periodic groups”, pp. 61–110 in *Seven papers on algebra*, Transl. Amer. Math. Soc. (2) **69**, Amer. Math. Soc., Providence, RI, 1968. MR 33 #7428 Zbl 0146.04405
- [Raynaud 1970] M. Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, Lecture Notes in Math. **119**, Springer, Berlin, 1970. MR 41 #5381 Zbl 0195.22701
- [Rosenlicht 1956] M. Rosenlicht, “Some basic theorems on algebraic groups”, *Amer. J. Math.* **78** (1956), 401–443. MR 18,514a Zbl 0073.37601

- [Rosenlicht 1961] M. Rosenlicht, “Toroidal algebraic groups”, *Proc. Amer. Math. Soc.* **12** (1961), 984–988. [MR 24 #A3162](#) [Zbl 0107.14703](#)
- [Sancho and Sancho 2009] C. Sancho de Salas and F. Sancho de Salas, “Principal bundles, quasi-abelian varieties and structure of algebraic groups”, *J. Algebra* **322**:8 (2009), 2751–2772. [MR 2010m:14058](#) [Zbl 1191.14055](#)
- [SGA 3_I 1970] M. Demazure and A. Grothendieck, *Schémas en groupes, Tome I: Propriétés générales des schémas en groupes, Exposés I–VII* (Séminaire de Géométrie Algébrique du Bois Marie 1962–1964), Lecture Notes in Math. **151**, Springer, Berlin, 1970. [MR 43 #223a](#) [Zbl 0207.51401](#)
- [SGA 3_{II} 1970] M. Demazure and A. Grothendieck, *Schémas en groupes, Tome II: Groupes de type multiplicatif, et structure des schémas en groupes généraux, Exposés VIII–XVIII* (Séminaire de Géométrie Algébrique du Bois Marie 1962–1964), Lecture Notes in Math. **152**, Springer, Berlin, 1970. [MR 43 #223b](#) [Zbl 0209.24201](#)
- [SGA 3_{III} 1970] M. Demazure and A. Grothendieck, *Schémas en groupes, Tome III: Structure des schémas en groupes réductifs, Exposés XIX–XXVI* (Séminaire de Géométrie Algébrique du Bois Marie 1962–1964), Lecture Notes in Math. **153**, Springer, Berlin, 1970. [MR 43 #223c](#) [Zbl 0212.52810](#)
- [Totaro 2013] B. Totaro, “Pseudo-abelian varieties”, *Ann. Sci. Éc. Norm. Supér. (4)* **46**:5 (2013), 693–721. [MR 3185350](#) [Zbl 1286.14061](#)

Received May 27, 2015. Revised July 12, 2015.

MICHEL BRION
INSTITUT FOURIER
UNIVERSITÉ DE GRENOBLE I (JOSEPH FOURIER)
100 RUE DES MATHÉMATIQUES
38402 SAINT-MARTIN D’HÈRES CEDEX
FRANCE
michel.brion@ujf-grenoble.fr

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 279 No. 1-2 December 2015

In memoriam: Robert Steinberg

Robert Steinberg (1922–2014): In memoriam V. S. VARADARAJAN	1
Cellularity of certain quantum endomorphism algebras HENNING H. ANDERSEN, GUSTAV I. LEHRER and RUIBIN ZHANG	11
Lower bounds for essential dimensions in characteristic 2 via orthogonal representations ANTONIO BABIC and VLADIMIR CHERNOUSOV	37
Cocharacter-closure and spherical buildings MICHAEL BATE, SEBASTIAN HERPEL, BENJAMIN MARTIN and GERHARD RÖHRLE	65
Embedding functor for classical groups and Brauer–Manin obstruction EVA BAYER-FLUCKIGER, TING-YU LEE and RAMAN PARIMALA	87
On maximal tori of algebraic groups of type G_2 CONSTANTIN BELI, PHILIPPE GILLE and TING-YU LEE	101
On extensions of algebraic groups with finite quotient MICHEL BRION	135
Essential dimension and error-correcting codes SHANE CERNELE and ZINOVY REICHSTEIN	155
Notes on the structure constants of Hecke algebras of induced representations of finite Chevalley groups CHARLES W. CURTIS	181
Complements on disconnected reductive groups FRANÇOIS DIGNE and JEAN MICHEL	203
Extending Hecke endomorphism algebras JIE DU, BRIAN J. PARSHALL and LEONARD L. SCOTT	229
Products of partial normal subgroups ELLEN HENKE	255
Lusztig induction and ℓ -blocks of finite reductive groups RADHA KESSAR and GUNTER MALLE	269
Free resolutions of some Schubert singularities MANOJ KUMMINI, VENKATRAMANI LAKSHMIBAI, PRAMATHANATH SASTRY and C. S. SESHADRI	299
Free resolutions of some Schubert singularities in the Lagrangian Grassmannian VENKATRAMANI LAKSHMIBAI and REUVEN HODGES	329
Distinguished unipotent elements and multiplicity-free subgroups of simple algebraic groups MARTIN W. LIEBECK, GARY M. SEITZ and DONNA M. TESTERMAN	357
Action of longest element on a Hecke algebra cell module GEORGE LUSZTIG	383
Generic stabilisers for actions of reductive groups BENJAMIN MARTIN	397
On the equations defining affine algebraic groups VLADIMIR L. POPOV	423
Smooth representations and Hecke modules in characteristic p PETER SCHNEIDER	447
On CRDAHA and finite general linear and unitary groups BHAMA SRINIVASAN	465
Weil representations of finite general linear groups and finite special linear groups PHAM HUU TIEP	481
The pro- p Iwahori Hecke algebra of a reductive p -adic group, V (parabolic induction) MARIE-FRANCE VIGNÉRAS	499
Acknowledgement	531