

*Pacific
Journal of
Mathematics*

PRODUCTS OF PARTIAL NORMAL SUBGROUPS

ELLEN HENKE

Volume 279 No. 1-2

December 2015

PRODUCTS OF PARTIAL NORMAL SUBGROUPS

ELLEN HENKE

In memory of Robert Steinberg.

We show that the product of two partial normal subgroups of a locality (in the sense of Chermak) is again a partial normal subgroup. This generalizes a theorem of Chermak and fits into the context of building a local theory of localities.

1. Introduction

Localities were introduced by Andrew Chermak [2013], in the context of his proof of the existence and uniqueness of centric linking systems. Roughly speaking, localities are group-like structures which are essentially the “same” as the transporter systems of Oliver and Ventura [2007]; see the appendix to [Chermak 2013]. As centric linking systems are special cases of transporter systems, the existence of centric linking systems implies that there is a locality attached to every fusion system. It is work in progress of Chermak to build a local theory of localities similar to the local theory of fusion systems as developed by Aschbacher [2008; 2011]. In fact, it seems often an advantage to work inside of localities, where some group theoretical concepts and constructions can be expressed more naturally than in fusion systems. Thus, one can hope to improve the local theory of fusion systems, once a way of translating between fusion systems and localities is established. The results of this paper can be considered as first evidence that some constructions are easier in the world of localities. We prove that the product of partial normal subgroups of a locality is itself a partial normal subgroup, whereas in fusion systems the product of normal subsystems has only been defined in special cases; see [Aschbacher 2011, Theorem 3]. It is work in progress of Chermak to show that there is a one-to-one correspondence between the normal subsystems of a saturated fusion system \mathcal{F} and the partial normal subgroups of a linking locality attached to \mathcal{F} in the sense of [Henke 2015, Definition 2]. This is one reason why our result seems particularly important in the case of linking localities. Another reason is that the concept of a linking locality generalizes properties of localities corresponding to centric linking

MSC2010: 20D20, 20N99.

Keywords: fusion systems, localities, transporter systems.

systems and is thus interesting for studying the homotopy theory of fusion systems; see [Broto et al. 2003; 2005; 2007; Henke 2015]. It is however crucial for our proof that we work with arbitrary localities, since our arguments rely heavily on the theory of quotient localities introduced by Chermak [2015], and a quotient of a linking locality is not necessarily a linking locality again. Thus, we feel that the method of our proof gives evidence for the value of studying localities in general rather than restricting attention only to the special case of linking localities.

To describe the results of this paper in more detail, let \mathcal{L} be a partial group as defined in [Chermak 2013, Definition 2.1; 2015, Definition 1.1]. Thus, there is an involutory bijection $\mathcal{L} \rightarrow \mathcal{L}$, $f \mapsto f^{-1}$, called an “inversion”, and a multivariable product Π which is only defined on certain words in \mathcal{L} . Let \mathbf{D} be the domain of the product; i.e., \mathbf{D} is a set of words in \mathcal{L} and Π is a map $\mathbf{D} \rightarrow \mathcal{L}$. Following Chermak, we call a nonempty subset \mathcal{H} of \mathcal{L} a *partial subgroup* of \mathcal{L} if $h^{-1} \in \mathcal{H}$ for all $h \in \mathcal{H}$ and $\Pi(v) \in \mathcal{H}$ for all words v in the alphabet \mathcal{H} with $v \in \mathbf{D}$. A partial subgroup \mathcal{N} is called a *partial normal subgroup* if $x^f := \Pi(f^{-1}, x, f) \in \mathcal{N}$ for all $x \in \mathcal{N}$ and $f \in \mathcal{L}$ for which $(f^{-1}, x, f) \in \mathbf{D}$. Given two subsets \mathcal{M} and \mathcal{N} of \mathcal{L} , the product \mathcal{MN} is naturally defined by

$$\mathcal{MN} = \{\Pi(m, n) : m \in \mathcal{M}, n \in \mathcal{N}, (m, n) \in \mathbf{D}\}.$$

The problem is however to show that this is again a partial normal subgroup if \mathcal{M} and \mathcal{N} are partial normal subgroups. Indeed, as we show in Example 2.3, this is not true in general if \mathcal{L} is an arbitrary partial group. It is true however in the important case that (\mathcal{L}, Δ, S) is a locality. Chermak [2015, Theorem 5.1] proved this in a special case and we build on his result to prove the general case. More precisely, we prove the following theorem:

Theorem 1. *Let (\mathcal{L}, Δ, S) be a locality and let \mathcal{M}, \mathcal{N} be partial normal subgroups of \mathcal{L} . Then $\mathcal{MN} = \mathcal{NM}$ is a partial normal subgroup of \mathcal{L} and $(\mathcal{MN}) \cap S = (\mathcal{M} \cap S)(\mathcal{N} \cap S)$. Moreover, for every $g \in \mathcal{MN}$ there exists $m \in \mathcal{M}$ and $n \in \mathcal{N}$ such that $(m, n) \in \mathbf{D}$, $g = \Pi(m, n)$, and $S_g = S_{(m,n)}$.*

To understand the technical conditions stated in the last sentence of the theorem, we recall from [Chermak 2013; 2015] that

$$S_g = \{s \in S : (g^{-1}, s, g) \in \mathbf{D} \text{ and } s^g \in S\}$$

for any $g \in \mathcal{L}$. Moreover, for a word $v = (g_1, \dots, g_n)$ in \mathcal{L} , S_v is the set of all $s \in S$ such that there exist $x_0, \dots, x_n \in S$ with $s = x_0$, $(g_i^{-1}, x_{i-1}, g_i) \in \mathbf{D}$ and $x_{i-1}^{g_i} = x_i$ for $i = 1, \dots, n$. By [op. cit, Proposition 2.6 and Corollary 2.7], the sets S_g and S_v are subgroups of S , $S_g \in \Delta$ for any $g \in \mathcal{L}$, and $S_v \in \Delta$ if and only if $v \in \mathbf{D}$. Therefore, the condition $S_g = S_{(m,n)}$ stated in the theorem is crucial for proving that certain products are defined in \mathcal{L} . This is particularly important for the proof of our

next theorem which concerns products of more than two partial normal subgroups. Given subsets $\mathcal{N}_1, \mathcal{N}_2, \dots, \mathcal{N}_l$ of \mathcal{L} define their product via

$$\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l := \{\Pi(n_1, n_2, \dots, n_l) : (n_1, n_2, \dots, n_l) \in \mathbf{D}, n_i \in \mathcal{N}_i \text{ for } 1 \leq i \leq l\}.$$

We prove:

Theorem 2. *Let $\mathcal{N}_1, \mathcal{N}_2, \dots, \mathcal{N}_l$ be partial normal subgroups of a locality (\mathcal{L}, Δ, S) . Then $\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l$ is a partial normal subgroup of \mathcal{L} . Moreover, the following hold:*

- (1) $\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l = (\mathcal{N}_1 \cdots \mathcal{N}_k)(\mathcal{N}_{k+1} \cdots \mathcal{N}_l)$ for every $1 \leq k < l$.
- (2) $\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l = \mathcal{N}_{1\sigma}\mathcal{N}_{2\sigma} \cdots \mathcal{N}_{l\sigma}$ for every permutation $\sigma \in S_l$.
- (3) For every $g \in \mathcal{N}_1 \cdots \mathcal{N}_l$ there exists $(n_1, \dots, n_l) \in \mathbf{D}$ with $n_i \in \mathcal{N}_i$ for every $i = 1, \dots, l$, $g = \Pi(n_1, \dots, n_l)$, and $S_g = S_{(n_1, \dots, n_l)}$.

As already mentioned above, it is work in progress of Andrew Chermak to show that for every fusion system \mathcal{F} and a linking locality (\mathcal{L}, Δ, S) attached to \mathcal{F} there is a one-to-one correspondence between the normal subsystems of \mathcal{F} and the partial normal subgroups of \mathcal{L} . When this work is complete, our results will imply the existence of a product of an arbitrary finite number of normal subsystems of \mathcal{F} .

In this text only relatively few demands will be made on understanding the concepts introduced in [Chermak 2013; 2015]. In Section 2, we point the reader to the few general results needed about partial groups, give a concise definition of a locality and review some basic facts about localities. In Section 3, we summarize what is needed about partial normal subgroups and quotient localities.

2. Partial groups and localities

We refer the reader to [Chermak 2013, Definition 2.1] or [Chermak 2015, Definition 1.1] for the precise definition of a partial group, and to the elementary properties of partial groups stated in [2013, Lemma 2.2] or [2015, Lemma 1.4]. Adapting Chermak’s notation we write $\mathbf{W}(\mathcal{L})$ for the set of words in a set \mathcal{L} , \emptyset for the empty word, and $v_1 \circ v_2 \circ \dots \circ v_n$ for the concatenation of words $v_1, \dots, v_n \in \mathbf{W}(\mathcal{L})$.

For the remainder of this text let \mathcal{L} be a partial group with product $\Pi : \mathbf{D} \rightarrow \mathcal{L}$ defined on the domain $\mathbf{D} \subseteq \mathbf{W}(\mathcal{L})$.

Again following Chermak’s notation, we set $\mathbf{1} = \Pi(\emptyset)$. Moreover, given a word $v = (f_1, \dots, f_n) \in \mathbf{D}$, we write $f_1 f_2 \dots f_n$ for the product $\Pi(v)$. Recall the definitions of partial subgroups and partial normal subgroups from the introduction. Note that a partial subgroup of \mathcal{L} is always a partial group itself whose product is the restriction of the product Π to $\mathbf{W}(\mathcal{H}) \cap \mathbf{D}$. Observe furthermore that \mathcal{L} forms a group in the usual sense if $\mathbf{W}(\mathcal{L}) = \mathbf{D}$; see [op. cit., Lemma 1.3]. So it makes sense

to call a partial subgroup \mathcal{H} of \mathcal{L} a *subgroup of \mathcal{L}* if $\mathbf{W}(\mathcal{H}) \subseteq \mathbf{D}$. In particular, we can talk about *p -subgroups of \mathcal{L}* meaning subgroups of \mathcal{L} whose order is a power of p .

We will need the Dedekind lemma [Chermak 2015, Lemma 1.10] in the following slightly more general form:

2.1 (Dedekind lemma). *Let $\mathcal{H}, \mathcal{K}, \mathcal{A}$ be subsets of \mathcal{L} such that \mathcal{A} is a partial subgroup of \mathcal{L} and $\mathcal{K} \subseteq \mathcal{A}$. Then $\mathcal{A} \cap (\mathcal{H}\mathcal{K}) = (\mathcal{A} \cap \mathcal{H})\mathcal{K}$ and $\mathcal{A} \cap (\mathcal{K}\mathcal{H}) = \mathcal{K}(\mathcal{A} \cap \mathcal{H})$.*

Proof. Clearly, $(\mathcal{A} \cap \mathcal{H})\mathcal{K} \subseteq \mathcal{A} \cap (\mathcal{H}\mathcal{K})$. Taking $h \in \mathcal{H}$ and $k \in \mathcal{K}$ with $(h, k) \in \mathbf{D}$ and $hk \in \mathcal{A}$, we have $(h, k, k^{-1}) \in \mathbf{D}$ by [op. cit, Lemma 1.4(d)] and then $h = h(kk^{-1}) = (hk)k^{-1} \in \mathcal{A}$ as $\mathcal{K} \subseteq \mathcal{A}$ and \mathcal{A} is a partial subgroup. Hence, $h \in \mathcal{A} \cap \mathcal{H}$ and $hk \in (\mathcal{A} \cap \mathcal{H})\mathcal{K}$. The second equation follows similarly. \square

Before we continue with more definitions, we illustrate the concepts we mentioned so far with examples. For this purpose we say that two groups G_1 and G_2 form an *amalgam*, if the set-theoretic intersection $G_1 \cap G_2$ is a subgroup of both G_1 and G_2 , and the restriction of the multiplication on G_1 to a multiplication on $G_1 \cap G_2$ is the same as the restriction of the multiplication on G_2 to a multiplication on $G_1 \cap G_2$.

Example 2.2. Let G_1 and G_2 be groups which form an amalgam. Set $\mathcal{L} = G_1 \cup G_2$ and $\mathbf{D} = \mathbf{W}(G_1) \cup \mathbf{W}(G_2)$. Define a partial product $\Pi : \mathbf{D} \rightarrow \mathcal{L}$ by sending $v = (f_1, \dots, f_n) \in \mathbf{W}(G_i)$ to the product $f_1 \dots f_n$ in the group G_i for $i = 1, 2$. Define an inversion $\mathcal{L} \rightarrow \mathcal{L}$ by sending $f \in G_i$ to the inverse of f in the group G_i for $i = 1, 2$. Then \mathcal{L} with these structures forms a partial group. (For readers familiar with the concept of an objective partial group as introduced in [Chermak 2013, Definition 2.6] or [Chermak 2015, Definition 2.1] we mention that, setting $\Delta := \{G_1, G_2\}$, (\mathcal{L}, Δ) is an objective partial group if $G_1 \cap G_2$ is properly contained in G_1 and G_2 .)

Let \mathcal{K} be a subset of \mathcal{L} . Then \mathcal{K} is a partial subgroup of \mathcal{L} if and only if $\mathcal{K} \cap G_i$ is a subgroup of G_i for each $i = 1, 2$. The subset \mathcal{K} is a subgroup of \mathcal{L} if and only if \mathcal{K} is a subgroup of G_i for some $i = 1, 2$. Moreover, \mathcal{K} is a partial normal subgroup of \mathcal{L} if and only if $(\mathcal{K} \cap G_i) \trianglelefteq G_i$ for $i = 1, 2$.

We use the construction method introduced in the previous example to show that the product of two partial normal subgroups of a partial group is not in general itself a partial normal subgroup.

Example 2.3. Let $G_1 \cong C_2 \times C_4$ and let G_2 be a dihedral group of order 16. Choose G_1 and G_2 such that G_1 and G_2 form an amalgam with $G_1 \cap G_2 \cong C_2 \times C_2$ and $\Phi(G_1) = Z(G_2)$. Let \mathcal{M} and \mathcal{N} be the two cyclic subgroups of G_1 of order 4. Form the locality \mathcal{L} as in Example 2.2. As G_1 is abelian, a subgroup \mathcal{K} of G_1 is normal in G_1 and thus a partial normal subgroup of \mathcal{L} if and only if $\mathcal{K} \cap G_2 \trianglelefteq G_2$. As $G_1 \cap G_2 \cong C_2 \times C_2$ and \mathcal{M} and \mathcal{N} are cyclic of order 4, we have $\mathcal{M} \cap G_2 =$

$\mathcal{M} \cap G_2 = \Phi(G_1) = Z(G_2) \trianglelefteq G_2$. Thus \mathcal{M} and \mathcal{N} are partial normal subgroups of \mathcal{L} . The product $\mathcal{M}\mathcal{N}$ in \mathcal{L} is the same as the product $\mathcal{M}\mathcal{N}$ in G_1 and thus equal to G_1 . However, as G_2 does not have a normal fours subgroup, $G_1 \cap G_2$ is not normal in G_2 and thus $\mathcal{M}\mathcal{N} = G_1$ is not a partial normal subgroup of \mathcal{L} .

The previous example shows that the concept of a partial group (and even the concept of an objective partial group) is too general for our purposes. Therefore, we will focus on localities. We give a definition of a locality which, in contrast to the definition given by Chermak [2013; 2015], does not require the reader to be familiar with the definition of an objective partial group and can easily be seen to be equivalent to Chermak’s definition. For any $g \in \mathcal{L}$, $\mathbf{D}(g)$ denotes the set of $x \in \mathcal{L}$ with $(g^{-1}, x, g) \in \mathbf{D}$. Thus, $\mathbf{D}(g)$ denotes the set of elements $x \in \mathcal{L}$ for which the conjugation $x^g := \Pi(g^{-1}, x, g)$ is defined. If $g \in \mathcal{L}$ and $X \subseteq \mathbf{D}(g)$ we set $X^g := \{x^g : x \in X\}$. If we write X^g for some $g \in \mathcal{L}$ and some subset $X \subseteq \mathcal{L}$, we will always implicitly mean that $X \subseteq \mathbf{D}(g)$.

Definition 2.4. We say that (\mathcal{L}, Δ, S) is a *locality* if the partial group \mathcal{L} is finite as a set, S is a p -subgroup of \mathcal{L} , Δ is a nonempty set of subgroups of S , and the following conditions hold:

- (L1) S is maximal with respect to inclusion among the p -subgroups of \mathcal{L} .
- (L2) A word $(f_1, \dots, f_n) \in \mathbf{W}(\mathcal{L})$ is an element of \mathbf{D} if and only if there exist $P_0, \dots, P_n \in \Delta$ such that
 - (*) $P_{i-1} \subseteq \mathbf{D}(f_i)$ and $P_{i-1}^{f_i} = P_i$.
- (L3) For any subgroup Q of S , for which there exist $P \in \Delta$ and $g \in \mathcal{L}$ with $P \subseteq \mathbf{D}(g)$ and $P^g \leq Q$, we have $Q \in \Delta$.

If (\mathcal{L}, Δ, S) is a locality and $v = (f_1, \dots, f_n) \in \mathbf{W}(\mathcal{L})$, then we say that $v \in \mathbf{D}$ via P_0, \dots, P_n (or $v \in \mathbf{D}$ via P_0), if $P_0, \dots, P_n \in \Delta$ and (*) holds.

From now on let (\mathcal{L}, Δ, S) be a locality.

Note that $P = P^1 \leq S$ for all $P \in \Delta$. As $\Delta \neq \emptyset$, property (L3) implies thus $S \in \Delta$. For any $g \in \mathcal{L}$, write c_g for the conjugation map

$$c_g : \mathbf{D}(g) \rightarrow \mathcal{L}, x \mapsto x^g.$$

Recall the definitions of S_g and S_v from the introduction. Note that $S_g \subseteq \mathbf{D}(g)$. For any subgroup X of \mathcal{L} set

$$N_{\mathcal{L}}(X) := \{f \in \mathcal{L} : X \subseteq \mathbf{D}(f), X^f = X\}.$$

2.5 (Important properties of localities). *The following hold:*

- (a) $N_{\mathcal{L}}(P)$ is a subgroup of \mathcal{L} for each $P \in \Delta$.

- (b) Let $P \in \Delta$ and $g \in \mathcal{L}$ with $P \subseteq S_g$. Then $Q := P^g \in \Delta$ (so in particular Q is a subgroup of S). Moreover, $N_{\mathcal{L}}(P) \subseteq \mathbf{D}(g)$ and

$$c_g : N_{\mathcal{L}}(P) \rightarrow N_{\mathcal{L}}(Q)$$

is an isomorphism of groups.

- (c) Let $w = (g_1, \dots, g_n) \in \mathbf{D}$ via (X_0, \dots, X_n) . Then

$$c_{g_1} \circ \dots \circ c_{g_n} = c_{\Pi(w)}$$

is a group isomorphism $N_{\mathcal{L}}(X_0) \rightarrow N_{\mathcal{L}}(X_n)$.

- (d) For every $g \in \mathcal{L}$, $S_g \in \Delta$. In particular, S_g is a subgroup of S .

- (e) For any $w \in \mathbf{W}(\mathcal{L})$, S_w is a subgroup of $S_{\Pi(w)}$, and $S_w \in \Delta$ if and only if $w \in \mathbf{D}$.¹

Proof. Properties (a)–(c) correspond to statements in [Chermak 2015, Lemma 2.3] except for the fact stated in (b) that $Q \in \Delta$. This is however true by [op. cit., Proposition 2.6(c)]. Property (d) is true by [op. cit., Proposition 2.6(a)] and property (e) is stated in [op. cit., Corollary 2.7]. □

3. Partial normal subgroups and quotient localities

In this section we continue to assume that (\mathcal{L}, Δ, S) is a locality. The following theorem is a special case of Theorem 1 and will be used to prove the more general theorem.

Theorem 3.1. *Let \mathcal{M}, \mathcal{N} be partial normal subgroups of \mathcal{L} such that $\mathcal{M} \cap \mathcal{N} = 1$. Then $\mathcal{M}\mathcal{N} = \mathcal{N}\mathcal{M}$ is a partial normal subgroup of \mathcal{L} . Moreover, for any $f \in \mathcal{M}\mathcal{N}$ there exists $m \in \mathcal{M}$ and $n \in \mathcal{N}$ such that $(m, n) \in \mathbf{D}$, $f = mn$, and $S_f = S_{(m,n)}$.*

Proof. As $\mathcal{M} \cap \mathcal{N} \subseteq S$, it follows from [Chermak 2015, Lemma 5.3] that \mathcal{M} normalizes $\mathcal{N} \cap S$ and \mathcal{N} normalizes $\mathcal{M} \cap S$. So by [op. cit., Theorem 5.1], $\mathcal{M}\mathcal{N} = \mathcal{N}\mathcal{M}$ is a partial normal subgroup of \mathcal{L} . Moreover, by [op. cit., Lemma 5.2], for any $f \in \mathcal{M}\mathcal{N}$ there exist $m \in \mathcal{M}$ and $n \in \mathcal{N}$ such that $(m, n) \in \mathbf{D}$, $f = mn$, and $S_f = S_{(m,n)}$. □

To deduce Theorem 1 from Theorem 3.1, we need the theory of quotient localities developed in [Chermak 2015]; see also [Chermak 2013, Sections 3 and 4]. For the convenience of the reader we quickly summarize this theory here. After that we state some more specialized lemmas needed in our proof.

Throughout let \mathcal{K} be a partial normal subgroup of \mathcal{L} and $T = S \cap \mathcal{K}$.

- 3.2.** (a) T is strongly closed in (\mathcal{L}, Δ, S) ; that is, $t^g \in T$ for every $g \in \mathcal{L}$ and every $t \in T \cap S_g$. In particular, $T^g = T$ for any $g \in \mathcal{L}$ with $T \subseteq S_g$.

- (b) T is maximal in the poset of p -subgroups of \mathcal{N} .

¹Recall the definition of $S_{(g_1, \dots, g_n)}$ from the introduction.

Proof. Let $g \in \mathcal{L}$ and $t \in T \cap S_g$. Then $t^s \in S$ and, as \mathcal{N} is a partial normal subgroup, $t^s \in \mathcal{N}$. Hence, $t^s \in S \cap \mathcal{N} = T$. This proves (a). Property (b) is proved in [Chermak 2015, Lemma 3.1(c)]. \square

We write $\uparrow_{\mathcal{K}}$ for the relation \uparrow introduced in [op. cit., Definition 3.6], but with the partial normal subgroup \mathcal{N} replaced by \mathcal{K} . Thus $\uparrow_{\mathcal{K}}$ is a relation on the set $\mathcal{L} \circ \Delta$ of pairs $(f, P) \in \mathcal{L} \times \Delta$ with $P \leq S_f$. For $(f, P), (g, Q) \in \mathcal{L} \circ \Delta$, we have $(f, P) \uparrow_{\mathcal{K}} (g, Q)$ if there exist $x \in N_{\mathcal{K}}(P, Q)$ and $y \in N_{\mathcal{K}}(P^f, Q^g)$ such that $xg = fy$. We say then $(f, P) \uparrow_{\mathcal{K}} (g, Q)$ via (x, y) . One easily sees that $\uparrow_{\mathcal{K}}$ is reflexive and transitive. Moreover, $(f, P) \uparrow_{\mathcal{K}} (f, S_f)$ via $(\mathbf{1}, \mathbf{1})$. An element $f \in \mathcal{L}$ is called $\uparrow_{\mathcal{K}}$ -maximal if (f, S_f) is maximal with respect to the relation $\uparrow_{\mathcal{K}}$ (i.e., if $(f, S_f) \uparrow_{\mathcal{K}} (g, Q)$ implies $(g, Q) \uparrow_{\mathcal{K}} (f, S_f)$ for any $(g, Q) \in \mathcal{L} \circ \Delta$). We summarize some important technical properties of the relation $\uparrow_{\mathcal{K}}$ in the following lemma.

3.3. *The following hold:*

- (a) *Every element of $N_{\mathcal{L}}(S)$ is $\uparrow_{\mathcal{K}}$ -maximal. In particular, every element of S is $\uparrow_{\mathcal{K}}$ -maximal.*
- (b) *If $f \in \mathcal{L}$ is $\uparrow_{\mathcal{K}}$ -maximal, then $T \leq S_f$.*
- (c) *(Stellmacher’s splitting lemma) Let $(x, f) \in \mathbf{D}$ such that $x \in \mathcal{K}$ and f is $\uparrow_{\mathcal{K}}$ -maximal. Then $S_{(x, f)} = S_{xf}$.*

Proof. Property (a) is [Chermak 2015, Lemma 3.7(a)], (b) is [op. cit., Proposition 3.9], and (c) is [op. cit., Lemma 3.12]. \square

The relation $\uparrow_{\mathcal{K}}$ is crucial for defining a quotient locality \mathcal{L}/\mathcal{K} somewhat analogously to quotients of groups. A coset of \mathcal{K} in \mathcal{L} is of the form

$$\mathcal{K}f = \{kf : k \in \mathcal{K}, (k, f) \in \mathbf{D}\}$$

for some $f \in \mathcal{L}$. A maximal coset of \mathcal{K} is a coset which is maximal with respect to inclusion among the cosets of \mathcal{K} in \mathcal{L} . The set of these maximal cosets is denoted by \mathcal{L}/\mathcal{K} .

3.4. *The following hold:*

- (a) *$f \in \mathcal{L}$ is $\uparrow_{\mathcal{K}}$ -maximal if and only if $\mathcal{K}f$ is a maximal coset.*
- (b) *The maximal cosets of \mathcal{K} form a partition of \mathcal{L} .*

Proof. This is [op. cit., Proposition 3.14(b),(c),(d)]. \square

The reader might note that what we call a coset would be more precisely called a right coset. The distinction does however not matter very much, since we are mostly interested in the maximal cosets and, by [op. cit., Proposition 3.14(a)], we

have $\mathcal{K}f = f\mathcal{K}$ for any $\uparrow_{\mathcal{K}}$ -maximal element $f \in \mathcal{L}$. By 3.4(b), we can define a map

$$\rho : \mathcal{L} \rightarrow \bar{\mathcal{L}} := \mathcal{L}/\mathcal{K}$$

sending $f \in \mathcal{L}$ to the unique maximal coset of \mathcal{K} containing f . This should be thought of as a “quotient map”. We adopt the bar notation similarly as used for groups. Thus, if X is an element or a subset of \mathcal{L} , then \bar{X} denotes the image of X under ρ . Furthermore, if X is an element or a subset of $\mathbf{W}(\mathcal{L})$ then \bar{X} denotes the image of X under ρ^* , where ρ^* denotes the map $\mathbf{W}(\mathcal{L}) \rightarrow \mathbf{W}(\mathcal{L})$ with $(f_1, \dots, f_n)\rho^* = (f_1\rho, \dots, f_n\rho)$. In particular,

$$\bar{\mathbf{D}} = \mathbf{D}\rho^*.$$

We note:

3.5. *Let $f, g \in \mathcal{L}$ such that $\bar{g} = \bar{f}$ and f is $\uparrow_{\mathcal{K}}$ -maximal. Then $g \in \mathcal{K}f$.*

Proof. By 3.4(a), $\mathcal{K}f$ is a maximal coset, so $\bar{g} = \bar{f} = \mathcal{K}f$ by the definition of ρ . Hence, again by the definition of ρ , $g \in \mathcal{K}f$. □

Recall the definition of a homomorphism of a partial groups from [Chermak 2013, Definition 3.1] and [Chermak 2015, Definition 1.11]. By [op. cit., Lemma 3.16], there is a unique mapping $\bar{\Pi} : \bar{\mathbf{D}} \rightarrow \bar{\mathcal{L}}$ and a unique involutory bijection $\bar{f} \mapsto \bar{f}^{-1}$ such that $\bar{\mathcal{L}}$ with these structures is a partial group and ρ is a homomorphism of partial groups. Since ρ is a homomorphism, we have $\bar{\Pi}(\bar{v}) = \bar{\Pi}(v\rho^*) = \bar{\Pi}(v)\rho = \bar{\Pi}(v)$ for $v \in \mathbf{D}$ and $\bar{f}^{-1} = \bar{f}^{-1}$ by the definition of a homomorphism of partial groups and by [op. cit., Lemma 1.13]. In particular, $\bar{\mathbf{1}} = \bar{\Pi}(\emptyset)$ is the identity element in $\bar{\mathcal{L}}$. So ρ has kernel $\ker(\rho) = \{f \in \mathcal{L} : \bar{f} = \bar{\mathbf{1}}\} = \mathcal{K}\mathbf{1} = \mathcal{K}$. By [op. cit., Proposition 4.2], $(\bar{\mathcal{L}}, \bar{\Delta}, \bar{S})$ is a locality for $\bar{\Delta} := \{\bar{P} : P \in \Delta\}$. We will use this important fact throughout without further reference. We remark:

3.6. *Let $v = (f_1, \dots, f_n) \in \mathbf{W}(\mathcal{L})$ such that each f_i is $\uparrow_{\mathcal{K}}$ -maximal and $\bar{v} \in \bar{\mathbf{D}}$. Then $v \in \mathbf{D}$ and $\bar{\Pi}(\bar{v}) = \bar{\Pi}(v)$.*

Proof. As $\bar{v} \in \bar{\mathbf{D}}$, there is $u = (g_1, \dots, g_n) \in \mathbf{D}$ such that $\bar{u} = \bar{v}$. Then $\bar{g}_i = \bar{f}_i$ for $i = 1, \dots, n$, i.e., $g_i \in \mathcal{K}f_i$ by 3.5. Now by [op. cit., Proposition 3.14(e)], $v \in \mathbf{D}$. As seen above, since ρ is a homomorphism of partial groups, $\bar{\Pi}(\bar{v}) = \bar{\Pi}(v)$. □

There is a nice correspondence between the partial subgroups of \mathcal{L} containing \mathcal{K} and the partial subgroups of $\bar{\mathcal{L}}$.

3.7. *Let \mathfrak{H} be the set of partial subgroups of \mathcal{L} containing \mathcal{K} .*

- (a) *Let $\mathcal{H} \in \mathfrak{H}$. Then the maximal cosets of \mathcal{K} contained in \mathcal{H} form a partition of \mathcal{H} .*
- (b) *Write $\bar{\mathfrak{H}}$ for the set of partial subgroups of $\bar{\mathcal{L}}$. Then the map $\mathfrak{H} \rightarrow \bar{\mathfrak{H}}$ with $\mathcal{H} \mapsto \bar{\mathcal{H}}$ is well defined and a bijection. Moreover, for any $\mathcal{H} \in \mathfrak{H}$, we have $\bar{\mathcal{H}} \trianglelefteq \bar{\mathcal{L}}$ if and only if $\mathcal{H} \trianglelefteq \mathcal{L}$.*

Proof. Property (a) is [Chermak 2015, Lemma 3.15]. The map ρ is a homomorphism of partial groups and $(\bar{\mathcal{L}}, \bar{\Delta}, \bar{S})$ is a locality. From the way \bar{D} and $\bar{\Delta}$ are defined, it follows that ρ is a projection in the sense of [op. cit., Definition 4.5]. Hence, property (b) is a reformulation of [op. cit., Proposition 4.8]. \square

3.8. *For any subset X of \mathcal{L} and for any partial subgroup \mathcal{H} of \mathcal{L} containing \mathcal{K} , $\overline{X \cap \mathcal{H}} = \overline{X} \cap \overline{\mathcal{H}}$.*

Proof. Clearly, $\overline{X \cap \mathcal{H}} \subseteq \overline{X} \cap \overline{\mathcal{H}}$. Let now $x \in X$ such that $\bar{x} \in \overline{\mathcal{H}}$. Then there exists $h \in \mathcal{H}$ such that $\bar{x} = \bar{h}$ and, by 3.7(a), we may choose h such that $\mathcal{K}h$ is a maximal coset. By the definition of ρ , this means $x \in \mathcal{K}h \subseteq \mathcal{H}$ and hence $x \in X \cap \mathcal{H}$. Thus $\bar{x} \in \overline{X \cap \mathcal{H}}$, proving $\overline{X \cap \mathcal{H}} \subseteq \overline{X} \cap \overline{\mathcal{H}}$. \square

3.9. *Let $R \leq S$. Then $\{f \in \mathcal{L} : \bar{f} \in \overline{R}\} = \mathcal{K}R$.*

Proof. Clearly, $\bar{f} \in \overline{R}$ for any $f \in \mathcal{K}R$, as \mathcal{K} is the kernel of ρ . Let now $f \in \mathcal{L}$ and $r \in R$ with $\bar{f} = \bar{r}$. As every element of S is $\uparrow_{\mathcal{K}}$ -maximal by 3.3(a), it follows from 3.5 that $f \in \mathcal{K}r \subseteq \mathcal{K}R$. This proves the assertion. \square

3.10. *Let $T \leq R \leq S$. Then $R = \{s \in S : \bar{s} \in \overline{R}\}$ and $N_{\bar{S}}(\overline{R}) = \overline{N_S(R)}$.*

Proof. By 3.9 and the Dedekind lemma (2.1), we have $\{s \in S : \bar{s} \in \overline{R}\} = S \cap (\mathcal{K}R) = (S \cap \mathcal{K})R = TR = R$. Moreover, for any element $t \in S$ with $\bar{t} \in N_{\bar{S}}(\overline{R})$ and any $r \in R$, we have $\bar{r}^t = \bar{r}^{\bar{t}} \in \overline{R}$, so $r^t \in \{s \in S : \bar{s} \in \overline{R}\} = R$. Hence, $N_{\bar{S}}(\overline{R}) \leq \overline{N_S(R)}$. As ρ is a homomorphism of partial groups, $\overline{N_S(R)} \subseteq N_{\bar{S}}(\overline{R})$, so the assertion holds. \square

3.11. *For every $f \in \mathcal{L}$ such that f is $\uparrow_{\mathcal{K}}$ -maximal, we have $\overline{S_f} = \overline{S_{\bar{f}}}$.*

Proof. Set $P = S_f$ and $Q = P^f$. As ρ is a homomorphism of partial groups, one easily observes that $\overline{P} \subseteq \overline{S_{\bar{f}}}$. As $(\bar{\mathcal{L}}, \bar{\Delta}, \bar{S})$ is a locality, $\overline{S_{\bar{f}}}$ is a p -group. So assuming the assertion is wrong, there exists $a \in S$ such that $\bar{a} \in N_{\overline{S_{\bar{f}}}}(\overline{P}) \setminus \overline{P}$. As f is \uparrow -maximal, $T \leq P = S_f$ by 3.3(b). Hence, by 3.10 applied with P in the role of R , $\bar{a} \in \overline{N_S(P)}$. So by 3.10 now applied with $N_S(P)$ in the role of R , $a \in N_S(P)$. Using 2.5(a),(b), we conclude that $A := P\langle a \rangle$ is a p -subgroup of the group $N_{\mathcal{L}}(P)$ and that A^f is a p -subgroup of the group $N_{\mathcal{L}}(Q)$. As $\overline{A^f} \subseteq \overline{S}$, we have $\overline{A^f} \subseteq N_{\bar{S}}(\overline{Q})$. By 3.2(a), $T = T^f \leq Q$. Thus, by 3.10, $\overline{A^f} \subseteq \overline{N_S(Q)}$. Now 3.9 yields $A^f \subseteq (\mathcal{K}N_S(Q)) \cap N_{\mathcal{L}}(Q) = N_{\mathcal{K}}(Q)N_S(Q)$, where the last equality uses the Dedekind lemma (2.1). Recall that $N_{\mathcal{L}}(Q)$ is a finite group. Clearly, $N_{\mathcal{K}}(Q)$ is a normal subgroup of $N_{\mathcal{L}}(Q)$. It follows from 3.2(b) that $T \in \text{Syl}_p(N_{\mathcal{K}}(Q))$. So $N_S(Q) \in \text{Syl}_p(N_{\mathcal{K}}(Q)N_S(Q))$ and by Sylow's theorem, there exists $c \in N_{\mathcal{K}}(Q)$ such that $A^{f^c} \leq N_S(Q)$. Then $(f, P) \uparrow_{\mathcal{K}} (fc, A)$ via $(\mathbf{1}, c)$ contrary to f being $\uparrow_{\mathcal{K}}$ -maximal. \square

3.12. *Suppose that $f, g \in \mathcal{L}$ such that $\bar{f} = \bar{g}$, $S_f = S_g$, and f is $\uparrow_{\mathcal{K}}$ -maximal. Then g is $\uparrow_{\mathcal{K}}$ -maximal and $\mathcal{K}f = \mathcal{K}g$.*

Proof. As f is $\uparrow_{\mathcal{K}}$ -maximal and $\bar{f} = \bar{g}$, we have $g \in \mathcal{K}f$ by 3.5, i.e., there exists $k \in \mathcal{K}$ with $(k, f) \in \mathbf{D}$ and $g = kf$. By Stellmacher’s splitting lemma 3.3(c), we have $S_g = S_{kf} = S_{(k,f)}$. Hence, $S_f = S_g = S_{(k,f)}$ and thus $k \in N_{\mathcal{L}}(S_f)$. By 2.5(c), $k^{-1} \in N_{\mathcal{L}}(S_f)$ and $(k^{-1}, k, f) \in \mathbf{D}$ as via S_f . Hence, $(k^{-1}, g) = (k^{-1}, kf) \in \mathbf{D}$, $k^{-1}g = k^{-1}(kf) = k^{-1}kf = (k^{-1}k)f = f$ and $S_f^g = S_f^f$. This shows that $(f, S_f) \uparrow_{\mathcal{K}} (g, S_f)$ via $(k^{-1}, \mathbf{1})$. We conclude that g is $\uparrow_{\mathcal{K}}$ -maximal as f is $\uparrow_{\mathcal{K}}$ -maximal and $\uparrow_{\mathcal{K}}$ is transitive. By 3.4, $\mathcal{K}g$ and $\mathcal{K}f$ are both maximal cosets, and the maximal cosets of \mathcal{K} form a partition of \mathcal{L} . So it follows that $\mathcal{K}f = \mathcal{K}g$. \square

4. Proof of Theorem 1

Throughout this section assume the hypothesis of Theorem 1. Set

$$\mathcal{K} := \mathcal{M} \cap \mathcal{N}.$$

Observe that \mathcal{K} is a partial normal subgroup of \mathcal{L} . As in Section 3, let

$$\rho : \mathcal{L} \rightarrow \bar{\mathcal{L}} := \mathcal{L}/\mathcal{K}$$

be the quotient map sending $f \in \mathcal{L}$ to the unique maximal coset of \mathcal{K} containing f , and use the bar notation as introduced there. Set

$$T := \mathcal{K} \cap S.$$

4.1. $\bar{\mathcal{M}} \cap \bar{\mathcal{N}} = 1$.

Proof. As \mathcal{K} is contained in \mathcal{M} and \mathcal{N} , this is a special case of 3.8. \square

4.2. We have $\bar{\mathcal{M}}\bar{\mathcal{N}} = \bar{\mathcal{N}}\bar{\mathcal{M}}$, and $\bar{\mathcal{M}}\bar{\mathcal{N}}$ is a partial normal subgroup of $\bar{\mathcal{L}}$. Moreover, for any $x \in \bar{\mathcal{M}}\bar{\mathcal{N}}$, there exist $\bar{m} \in \bar{\mathcal{M}}$ and $\bar{n} \in \bar{\mathcal{N}}$ such that $(\bar{m}, \bar{n}) \in \bar{\mathbf{D}}$, $x = \bar{m}\bar{n}$ and $\bar{S}_x = \bar{S}_{(\bar{m}, \bar{n})}$.

Proof. By 3.7(b), $\bar{\mathcal{M}}$ and $\bar{\mathcal{N}}$ are partial normal subgroups of $\bar{\mathcal{L}}$. By 4.1, $\bar{\mathcal{M}} \cap \bar{\mathcal{N}} = 1$. Hence, the assertion follows from Theorem 3.1. \square

4.3. Let $x \in \bar{\mathcal{M}}\bar{\mathcal{N}}$. Then there exist $m \in \mathcal{M}$ and $n \in \mathcal{N}$ with $(m, n) \in \mathbf{D}$ such that m, n , and mn are $\uparrow_{\mathcal{K}}$ -maximal, $x = \bar{m}\bar{n} = \overline{mn}$ and $S_{mn} = S_{(m,n)}$.

Proof. By 4.2, there exist $\bar{m} \in \bar{\mathcal{M}}$, $\bar{n} \in \bar{\mathcal{N}}$ such that $(\bar{m}, \bar{n}) \in \bar{\mathbf{D}}$, $x = \bar{m}\bar{n}$, and $\bar{S}_x = \bar{S}_{(\bar{m}, \bar{n})}$. By 3.7(a), we may furthermore choose preimages $m \in \mathcal{M}$ and $n \in \mathcal{N}$ of \bar{m} and \bar{n} such that m and n are $\uparrow_{\mathcal{K}}$ -maximal. Then, by 3.2(a) and 3.3(b), $m, n \in N_{\mathcal{L}}(T)$. By 3.6, $(m, n) \in \mathbf{D}$ and $\bar{m}\bar{n} = \overline{mn}$. It remains to prove that $S_{mn} = S_{(m,n)}$ and that mn is $\uparrow_{\mathcal{K}}$ -maximal. As an intermediate step we prove the following two properties:

$$(4-1) \quad S_f \subseteq S_{(m,n)} \text{ for every } f \in \mathcal{L} \text{ with } \bar{f} = x.$$

$$(4-2) \quad S_f = S_{(m,n)} \text{ for every } \uparrow_{\mathcal{K}}\text{-maximal element } f \in \mathcal{L} \text{ with } \bar{f} = x.$$

For the proof of (4-1) and (4-2) note first that, by 3.11, $\bar{S}_m = \overline{S_m}$ and $\bar{S}_n = \overline{S_n}$ as m and n are $\uparrow_{\mathcal{K}}$ -maximal. Hence,

$$\bar{S}_x = \bar{S}_{(\bar{m}, \bar{n})} = \{\bar{s} : \bar{s} \in \bar{S}_m, \bar{s}^{\bar{m}} \in \bar{S}_n\} = \{\bar{s} : s \in S_m, \bar{s}^{\bar{m}} \in \bar{S}_n\}.$$

If $s \in S_m$ then, by definition of S_m , $(m^{-1}, s, m) \in \mathbf{D}$ and $s^m \in S$. Moreover, as ρ is a homomorphism of partial groups, $\bar{s}^{\bar{m}} = \overline{s^m}$. So $\bar{s}^{\bar{m}} \in \bar{S}_n$ is equivalent to $s^m \in S_n$ by 3.10 since $T \leq S_n$. Hence,

$$\bar{S}_x = \{\bar{s} : s \in S_m, s^m \in S_n\} = \overline{S_{(m,n)}}.$$

As $m, n \in N_{\mathcal{L}}(T)$, $T \leq S_{(m,n)}$. Clearly, $\bar{S}_f \subseteq \bar{S}_x$ for every $f \in \mathcal{L}$ with $\bar{f} = x$. If such f is in addition $\uparrow_{\mathcal{K}}$ -maximal, then $\bar{S}_f = \bar{S}_x$ and $T \leq S_f$ by 3.11 and 3.3(b). Now (4-1) and (4-2) follow from 3.10. As $\overline{mn} = \overline{m} \overline{n} = x$, (4-1) yields in particular $S_{mn} \subseteq S_{(m,n)}$ and thus $S_{mn} = S_{(m,n)}$ by 2.5(e). Choosing $f \in \mathcal{L}$ to be $\uparrow_{\mathcal{K}}$ -maximal with $\bar{f} = x$, we obtain from (4-2) that $S_f = S_{(m,n)} = S_{mn}$. So mn is $\uparrow_{\mathcal{K}}$ -maximal by 3.12 completing the proof. \square

4.4. *Let $f \in \mathcal{L}$ with $\bar{f} \in \overline{\mathcal{M}\mathcal{N}}$. Then $f \in \mathcal{MN}$ and there exist $m \in \mathcal{M}$, $n \in \mathcal{N}$ with $(m, n) \in \mathbf{D}$, $f = mn$, and $S_f = S_{(m,n)}$.*

Proof. By 4.3, we can choose $m \in \mathcal{M}$ and $n \in \mathcal{N}$ with $(m, n) \in \mathbf{D}$ such that mn is $\uparrow_{\mathcal{K}}$ -maximal, $\bar{f} = \overline{mn}$ and $S_{mn} = S_{(m,n)}$. Then there exists $k \in \mathcal{K}$ with $(k, mn) \in \mathbf{D}$ and $f = k(mn)$. As $S_{mn} = S_{(m,n)}$, it follows that $S_{(k,mn)} = S_{(k,m,n)}$ and $(k, m, n) \in \mathbf{D}$ by 2.5(e). Hence, $(km, n) \in \mathbf{D}$ and $f = (km)n$ by the axioms of a partial group. As $\mathcal{K} \subseteq \mathcal{M}$, we have $km \in \mathcal{M}$ and so $f = (km)n \in \mathcal{MN}$. It is now sufficient to show that $S_{(km,n)} = S_f$. As mn is $\uparrow_{\mathcal{K}}$ -maximal, it follows from Stellmacher's splitting lemma 3.3(c) that $S_f = S_{k(mn)} = S_{(k,mn)} = S_{(k,m,n)} \subseteq S_{(km,n)}$. By 2.5(e), $S_{(km,n)} \subseteq S_{(km)n} = S_f$. So $S_f = S_{(km,n)}$, proving the assertion. \square

Proof of Theorem 1. By 4.2 and 3.7(b), there exists a partial normal subgroup \mathcal{H} of \mathcal{L} containing \mathcal{K} such that $\overline{\mathcal{H}} = \overline{\mathcal{M}\mathcal{N}} = \overline{\mathcal{N}\mathcal{M}}$. Then for any $f \in \mathcal{L}$ with $\bar{f} \in \overline{\mathcal{M}\mathcal{N}}$, there exists $h \in \mathcal{H}$ with $\bar{f} = \bar{h}$. By 3.7(a), we can choose h to be $\uparrow_{\mathcal{K}}$ -maximal. So by 3.5, $f \in \mathcal{K}h \subseteq \mathcal{H}$. This shows $\mathcal{H} = \{f \in \mathcal{L} : \bar{f} \in \overline{\mathcal{M}\mathcal{N}}\}$.

We need to prove that $\mathcal{H} = \mathcal{MN} = \mathcal{NM}$. As the situation is symmetric in \mathcal{M} and \mathcal{N} , it is enough to prove that $\mathcal{H} = \mathcal{MN}$. Since ρ is a homomorphism, for any $m \in \mathcal{M}$ and $n \in \mathcal{N}$ with $(m, n) \in \mathbf{D}$, we have $\overline{mn} = \overline{m} \overline{n} \in \overline{\mathcal{M}\mathcal{N}}$ and thus $mn \in \mathcal{H}$. Hence, $\mathcal{MN} \subseteq \mathcal{H}$. By 4.4, we have $\mathcal{H} \subseteq \mathcal{MN}$, so $\mathcal{H} = \mathcal{MN}$. Moreover, 4.4 shows that for every $f \in \mathcal{MN}$, there exists $m \in \mathcal{M}$ and $n \in \mathcal{N}$ such that $(m, n) \in \mathbf{D}$, $f = mn$, and $S_f = S_{(m,n)}$. So it only remains to prove that $S \cap (\mathcal{MN}) = (S \cap \mathcal{M})(S \cap \mathcal{N})$. Clearly, $(S \cap \mathcal{M})(S \cap \mathcal{N}) \subseteq S \cap (\mathcal{MN})$. Let now $s \in S \cap (\mathcal{MN})$. By what we just said, there exists $m \in \mathcal{M}$ and $n \in \mathcal{N}$ with $(m, n) \in \mathbf{D}$, $s = mn$, and $S_s = S_{(m,n)}$. As $S_s = S$, it follows that $m, n \in G := N_{\mathcal{L}}(S)$. By 2.5(a), G is a subgroup of \mathcal{L} . Furthermore, property (L1) in the definition of a locality implies that S is a Sylow

p -subgroup of G . Note that $X := G \cap \mathcal{M}$ and $Y := G \cap \mathcal{N}$ are normal subgroups of G . Hence, $s = mn \in (XY) \cap S = (X \cap S)(Y \cap S) = (\mathcal{M} \cap S)(\mathcal{N} \cap S)$ completing the proof. \square

5. The Proof of Theorem 2

Throughout, let (\mathcal{L}, Δ, S) be a locality with partial normal subgroups $\mathcal{N}_1, \dots, \mathcal{N}_l$. We prove Theorem 2 in a series of lemmas.

5.1. *Let $1 \leq k < l$ such that the products $\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_k$ and $\mathcal{N}_{k+1}\mathcal{N}_{k+2} \cdots \mathcal{N}_l$ are partial normal subgroups. Suppose furthermore that for any $f \in \mathcal{N}_1 \cdots \mathcal{N}_k$ and any $g \in \mathcal{N}_{k+1} \cdots \mathcal{N}_l$ there exist $u = (n_1, \dots, n_k), v = (n_{k+1}, \dots, n_l) \in \mathbf{D}$ such that $n_i \in \mathcal{N}_i$ for $i = 1, \dots, l, f = \Pi(u), g = \Pi(v), S_f = S_u$ and $S_g = S_v$. Then*

$$\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l = (\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_k)(\mathcal{N}_{k+1}\mathcal{N}_{k+2} \cdots \mathcal{N}_l)$$

is a partial normal subgroup of \mathcal{L} , and for every $h \in \mathcal{N}_1, \dots, \mathcal{N}_l$ there exists $w = (n_1, \dots, n_l) \in \mathbf{D}$ such that $n_i \in \mathcal{N}_i$ for $i = 1, \dots, l, h = \Pi(w)$, and $S_h = S_w$.

Proof. By Theorem 1, $(\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_k)(\mathcal{N}_{k+1}\mathcal{N}_{k+2} \cdots \mathcal{N}_l)$ is a partial normal subgroup of \mathcal{L} . If $w = (n_1, \dots, n_l) \in \mathbf{D}$ with $n_i \in \mathcal{N}_i$ for $i = 1, \dots, l$, then $u = (n_1, \dots, n_k), v = (n_{k+1}, \dots, n_l) \in \mathbf{D}$, and

$$\Pi(w) = \Pi(\Pi(u), \Pi(v)) \in (\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_k)(\mathcal{N}_{k+1}\mathcal{N}_{k+2} \cdots \mathcal{N}_l).$$

This proves that $\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l \subseteq (\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_k)(\mathcal{N}_{k+1}\mathcal{N}_{k+2} \cdots \mathcal{N}_l)$. To prove the converse inclusion, let

$$h \in (\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_k)(\mathcal{N}_{k+1}\mathcal{N}_{k+2} \cdots \mathcal{N}_l).$$

Then by Theorem 1, there exist $f \in \mathcal{N}_1 \cdots \mathcal{N}_k$ and $g \in \mathcal{N}_{k+1} \cdots \mathcal{N}_l$ such that $(f, g) \in \mathbf{D}, h = fg$, and $S_h = S_{(f,g)}$. By assumption, there exist $u = (n_1, \dots, n_k)$ and $v = (n_{k+1}, \dots, n_l) \in \mathbf{D}$ such that $n_i \in \mathcal{N}_i$ for $i = 1, \dots, l, f = \Pi(u), g = \Pi(v), S_f = S_u$, and $S_g = S_v$. Then $S_h = S_{(f,g)} = S_{u \circ v}, u \circ v \in \mathbf{D}$ via S_h , and

$$h = fg = \Pi(\Pi(u), \Pi(v)) = \Pi(u \circ v) \in \mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l,$$

proving the assertion. \square

5.2. (a) *The product $\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l$ is a partial normal subgroup, and for every $f \in \mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l$ there exists $w = (n_1, \dots, n_l) \in \mathbf{D}$ such that $n_i \in \mathcal{N}_i$ for $i = 1, \dots, l, f = \Pi(w)$, and $S_f = S_w$.*

(b) *For every $1 \leq k < l$, we have*

$$\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l = (\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_k)(\mathcal{N}_{k+1}\mathcal{N}_{k+2} \cdots \mathcal{N}_l).$$

Proof. We prove this by induction on l . Clearly, the claim is true for $l = 1$. Assume now $l > 1$. Then there exists always $1 \leq k < l$. For any such k , it follows by induction (one time applied with $\mathcal{N}_1, \dots, \mathcal{N}_k$ and one time applied with $\mathcal{N}_{k+1}, \dots, \mathcal{N}_l$ in place of $\mathcal{N}_1, \dots, \mathcal{N}_l$) that the hypothesis of 5.1 is fulfilled, so the assertion follows. \square

5.3. *Let $\sigma \in S_l$ be a permutation. Then $\mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l = \mathcal{N}_{1\sigma}\mathcal{N}_{2\sigma} \cdots \mathcal{N}_{l\sigma}$.*

Proof. We may assume that $\sigma = (i, i + 1)$ for some $1 \leq i < l$, as S_l is generated by transpositions of this form. Note that $\mathcal{N}_1 \cdots \mathcal{N}_{i-1}, \mathcal{N}_i\mathcal{N}_{i+1}$ and $\mathcal{N}_{i+2} \cdots \mathcal{N}_l$ are partial normal subgroups by 5.2(a), where it is understood that $\mathcal{N}_r \cdots \mathcal{N}_s = \{\mathbf{1}\}$ if $r > s$. By Theorem 1, we have $\mathcal{M}\mathcal{N} = \mathcal{N}\mathcal{M}$ for any two partial normal subgroups. Using this fact and 5.2(b) repeatedly, we obtain

$$\begin{aligned} \mathcal{N}_1\mathcal{N}_2 \cdots \mathcal{N}_l &= (\mathcal{N}_1 \cdots \mathcal{N}_{i-1})(\mathcal{N}_i\mathcal{N}_{i+1})(\mathcal{N}_{i+2} \cdots \mathcal{N}_l) \\ &= (\mathcal{N}_1 \cdots \mathcal{N}_{i-1})(\mathcal{N}_{i+1}\mathcal{N}_i)(\mathcal{N}_{i+2} \cdots \mathcal{N}_l) \\ &= \mathcal{N}_{1\sigma}\mathcal{N}_{2\sigma} \cdots \mathcal{N}_{l\sigma}. \end{aligned} \quad \square$$

Proof of Theorem 2. It follows from 5.2(a) that $\mathcal{N}_1 \cdots \mathcal{N}_l$ is a partial normal subgroup and that (c) holds. Property (a) is 5.2(b), and property (b) is 5.3. \square

References

[Aschbacher 2008] M. Aschbacher, “Normal subsystems of fusion systems”, *Proc. Lond. Math. Soc.* (3) **97**:1 (2008), 239–271. MR 2009e:20044 Zbl 1241.20026

[Aschbacher 2011] M. Aschbacher, *The generalized Fitting subsystem of a fusion system*, *Memoirs of the American Mathematical Society* **209**:986, American Mathematical Society, Providence, RI, 2011. MR 2012f:20063 Zbl 1278.20020

[Broto et al. 2003] C. Broto, R. Levi, and B. Oliver, “Homotopy equivalences of p -completed classifying spaces of finite groups”, *Invent. Math.* **151**:3 (2003), 611–664. MR 2004c:55031 Zbl 1042.55008

[Broto et al. 2005] C. Broto, N. Castellana, J. Grodal, R. Levi, and B. Oliver, “Subgroup families controlling p -local finite groups”, *Proc. London Math. Soc.* (3) **91**:2 (2005), 325–354. MR 2007e:20111 Zbl 1090.20026

[Broto et al. 2007] C. Broto, N. Castellana, J. Grodal, R. Levi, and B. Oliver, “Extensions of p -local finite groups”, *Trans. Amer. Math. Soc.* **359**:8 (2007), 3791–3858. MR 2008i:55013 Zbl 1145.55013

[Chermak 2013] A. Chermak, “Fusion systems and localities”, *Acta Math.* **211**:1 (2013), 47–139. MR 3118305 Zbl 1295.20021

[Chermak 2015] A. Chermak, “Localities and projections”, preprint, 2015. arXiv 1505.07786v2

[Henke 2015] E. Henke, “Subcentric linking systems”, preprint, 2015. arXiv 1506.01458

[Oliver and Ventura 2007] B. Oliver and J. Ventura, “Extensions of linking systems with p -group kernel”, *Math. Ann.* **338**:4 (2007), 983–1043. MR 2008k:55029 Zbl 1134.55011

Received June 2, 2015. Revised August 30, 2015.

ELLEN HENKE
INSTITUTE OF MATHEMATICS
UNIVERSITY OF ABERDEEN
FRASER NOBLE BUILDING
ABERDEEN
AB24 3UE
UNITED KINGDOM
ellen.henke@abdn.ac.uk

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 279 No. 1-2 December 2015

In memoriam: Robert Steinberg

Robert Steinberg (1922–2014): In memoriam V. S. VARADARAJAN	1
Cellularity of certain quantum endomorphism algebras HENNING H. ANDERSEN, GUSTAV I. LEHRER and RUIBIN ZHANG	11
Lower bounds for essential dimensions in characteristic 2 via orthogonal representations ANTONIO BABIC and VLADIMIR CHERNOUSOV	37
Cocharacter-closure and spherical buildings MICHAEL BATE, SEBASTIAN HERPEL, BENJAMIN MARTIN and GERHARD RÖHRLE	65
Embedding functor for classical groups and Brauer–Manin obstruction EVA BAYER-FLUCKIGER, TING-YU LEE and RAMAN PARIMALA	87
On maximal tori of algebraic groups of type G_2 CONSTANTIN BELI, PHILIPPE GILLE and TING-YU LEE	101
On extensions of algebraic groups with finite quotient MICHEL BRION	135
Essential dimension and error-correcting codes SHANE CERNELE and ZINOVY REICHSTEIN	155
Notes on the structure constants of Hecke algebras of induced representations of finite Chevalley groups CHARLES W. CURTIS	181
Complements on disconnected reductive groups FRANÇOIS DIGNE and JEAN MICHEL	203
Extending Hecke endomorphism algebras JIE DU, BRIAN J. PARSHALL and LEONARD L. SCOTT	229
Products of partial normal subgroups ELLEN HENKE	255
Lusztig induction and ℓ -blocks of finite reductive groups RADHA KESSAR and GUNTER MALLE	269
Free resolutions of some Schubert singularities MANOJ KUMMINI, VENKATRAMANI LAKSHMIBAI, PRAMATHANATH SASTRY and C. S. SESHADRI	299
Free resolutions of some Schubert singularities in the Lagrangian Grassmannian VENKATRAMANI LAKSHMIBAI and REUVEN HODGES	329
Distinguished unipotent elements and multiplicity-free subgroups of simple algebraic groups MARTIN W. LIEBECK, GARY M. SEITZ and DONNA M. TESTERMAN	357
Action of longest element on a Hecke algebra cell module GEORGE LUSZTIG	383
Generic stabilisers for actions of reductive groups BENJAMIN MARTIN	397
On the equations defining affine algebraic groups VLADIMIR L. POPOV	423
Smooth representations and Hecke modules in characteristic p PETER SCHNEIDER	447
On CRDAHA and finite general linear and unitary groups BHAMA SRINIVASAN	465
Weil representations of finite general linear groups and finite special linear groups PHAM HUU TIEP	481
The pro- p Iwahori Hecke algebra of a reductive p -adic group, V (parabolic induction) MARIE-FRANCE VIGNÉRAS	499
Acknowledgement	531



0030-8730(2015)279:1;1-1