

Pacific Journal of Mathematics

**SOLUTIONS WITH LARGE NUMBER OF PEAKS
FOR THE SUPERCRITICAL HÉNON EQUATION**

ZHONGYUAN LIU AND SHUANGJIE PENG

SOLUTIONS WITH LARGE NUMBER OF PEAKS FOR THE SUPERCRITICAL HÉNON EQUATION

ZHONGYUAN LIU AND SHUANGJIE PENG

This paper is concerned with the Hénon equation

$$\begin{cases} -\Delta u = |y|^\alpha u^{p+\varepsilon}, & u > 0, & \text{in } B_1(0), \\ u = 0 & & \text{on } \partial B_1(0), \end{cases}$$

where $B_1(0)$ is the unit ball in \mathbb{R}^N ($N \geq 4$), $p = (N+2)/(N-2)$ is the critical Sobolev exponent, $\alpha > 0$ and $\varepsilon > 0$. We show that if ε is small enough, this problem has a positive peak solution which presents a new phenomenon: the number of its peaks varies with the parameter ε at the order $\varepsilon^{-1/(N-1)}$ when $\varepsilon \rightarrow 0^+$. Moreover, all peaks of the solutions approach the boundary of $B_1(0)$ as ε goes to 0^+ .

1. Introduction and main results

We study the existence of positive solutions to a type of nonlinear elliptic problem whose typical form is the supercritical problem

$$(1-1) \quad \begin{cases} -\Delta u = |y|^\alpha u^{p+\varepsilon}, & u > 0, & \text{in } B_1(0), \\ u = 0 & & \text{on } \partial B_1(0), \end{cases}$$

where $p = (N+2)/(N-2)$, $\alpha > 0$, $\varepsilon > 0$ and $B_1(0)$ is the unit ball in \mathbb{R}^N ($N \geq 4$).

It is well known that the problem

$$(1-2) \quad \begin{cases} -\Delta u = |y|^\alpha u^q, & u > 0, & \text{in } B_1(0), \\ u = 0 & & \text{on } \partial B_1(0), \end{cases}$$

was proposed by M. Hénon [1973] when he studied rotating stellar structures and is hence called the Hénon equation, and it has attracted a lot of interest in recent years. Ni [1982] first considered (1-2) and proved that it possesses a positive radial solution when $q \in (1, (N+2+2\alpha)/(N-2))$. Due to the appearance of the weighted term $|y|^\alpha$, the classical moving plane method in [Gidas et al. 1979] cannot be applied to problem (1-2). It is natural to ask whether problem (1-2) has nonradial solutions. The existence of a nonradial solution for $1 < q < p$ was obtained by

MSC2010: primary 35J60; secondary 35J65, 58E05.

Keywords: peak solutions, supercritical Hénon equation, reduction method.

Smets, Willem and Su [2002] provided α is large enough. When $q = p - \varepsilon$, Cao and Peng [2003] showed that the ground state solution is nonradial and blows up near the boundary of $B_1(0)$ as $\varepsilon \rightarrow 0$. Later on, Peng [2006] constructed multiple boundary peak solutions for problem (1-2). When $q = p$, Serra [2005] proved that problem (1-2) has a nonradial solution provided α is large enough. More recently, Wei and Yan [2013] showed that there are infinitely many nonradial solutions for problem (1-2) with $\alpha > 0$. For other results related to Hénon type problems, see [Byeon and Wang 2006; 2005; Cao et al. 2009; Hirano 2009; Pistoia and Serra 2007] and the references therein.

On the other hand, using the Pohozaev identity [1965], we know that for $q \geq \frac{N+2+2\alpha}{N-2}$ there are no solutions to problem (1-2) in star-shaped domains with respect to the origin. So it seems more interesting to consider whether there are solutions for q in the range $(\frac{N+2}{N-2}, \frac{N+2+2\alpha}{N-2})$. However, much less is known about that case. When $q = \frac{N+2+2\alpha}{N-2} - \varepsilon$, Gladiali and Grossi [2012] showed that there exists one solution concentrating at $y = 0$ provided $0 < \alpha \leq 1$. By the results in [Gladiali et al. 2013], the same results still hold when α is not an even integer. In [Li and Peng 2009], the asymptotic behavior of the radial solutions obtained by Ni [1982] was analyzed as $\varepsilon \rightarrow 0^+$.

The purpose of this paper is to study the supercritical problem (1-1) and try to construct solutions whose number of peaks varies with ε as $\varepsilon \rightarrow 0^+$. In fact, we will consider the more general problem

$$(1-3) \quad \begin{cases} -\Delta u = K(|y|)u^{p+\varepsilon}, & u > 0, & \text{in } B_1(0), \\ u = 0 & & \text{on } \partial B_1(0), \end{cases}$$

where $K(r) \in C^1[0, 1]$ and $K(1) > 0$.

Without loss of generality, we can assume that

$$K(1) = 1.$$

The main result of this paper is as follows.

Theorem 1.1. *Assume that $N \geq 4$. If $K(r)$ satisfies $K(1) > 0$ and $K'(1) > 0$, then there exists $\varepsilon_0 > 0$ such that for $\varepsilon \in (0, \varepsilon_0)$, problem (1-3) has a solution u_ε whose number of local maximal points is of the order $\varepsilon^{-1/(N-1)}$ as $\varepsilon \rightarrow 0^+$. In particular, problem (1-1) has solutions with a large number of peaks for small $\varepsilon > 0$.*

Remark 1.2. For the case $\alpha = 0$, the well-known Pohozaev identity [1965] implies that (1-1) has no solutions for $\varepsilon > 0$. It was also shown in [Ben Ayed et al. 2003] that problem (1-1) has no single-peak solutions for ε small enough. Our results mean that the weight $|y|^\alpha$ has a great influence on the existence of peak solutions for problem (1-1).

Let us outline the main idea in the proof of [Theorem 1.1](#). We introduce some notation first. For $x \in \mathbb{R}^N$ and $\Lambda > 0$, set

$$U_{x,\Lambda}(y) = C_N \left(\frac{\Lambda}{1 + \Lambda^2 |y - x|^2} \right)^{(N-2)/2}, \quad C_N = (N(N-2))^{(N-2)/4}.$$

It's well known that $U_{x,\Lambda}(y)$ are the only solutions of

$$-\Delta u = u^{(N+2)/(N-2)}, \quad u > 0, \quad \text{in } \mathbb{R}^N.$$

Let

$$k = [\varepsilon^{-1/(N-1)}],$$

where $[a]$ denotes the integer part of a real number a . By the transformation $u(y) \mapsto \varepsilon^{2/(4+(N-2)\varepsilon)} u(\varepsilon^{1/(N-2)} y)$ and setting $B_* = B_{\varepsilon^{-1/(N-2)}}$, we see that (1-3) becomes

$$(1-4) \quad \begin{cases} -\Delta u = K(\varepsilon^{1/(N-2)} |y|) u^{p+\varepsilon}, & u > 0, & \text{in } B_*(0), \\ u = 0 & & \text{on } \partial B_*(0). \end{cases}$$

We denote by $PU_{x,\Lambda}$, the projection of $U_{x,\Lambda}$, the solution of the problem

$$(1-5) \quad \begin{cases} \Delta PU_{x,\Lambda} = \Delta U_{x,\Lambda} & \text{in } B_*(0), \\ PU_{x,\Lambda} = 0 & \text{on } \partial B_*(0). \end{cases}$$

Set $y = (y', y'')$, $y'' \in \mathbb{R}^{N-2}$. Define

$$\mathcal{H}_s = \left\{ u : u \in H_0^1(B_*(0)), \quad u \text{ is even in } y_h, \quad h = 2, 3, \dots, N, \right. \\ \left. u(r \cos \theta, r \sin \theta, y'') = u \left(r \cos \left(\theta + \frac{2\pi j}{k} \right), r \sin \left(\theta + \frac{2\pi j}{k} \right), y'' \right) \right\}.$$

Let

$$x_j = \left(r \cos \frac{2(j-1)\pi}{k}, r \sin \frac{2(j-1)\pi}{k}, 0 \right), \quad j = 1, \dots, k,$$

where 0 is the zero vector in \mathbb{R}^{N-2} , and let

$$W_{r,\Lambda}(y) = \sum_{j=1}^k PU_{x_j,\Lambda}.$$

In what follows, we always assume that

$$r \in [\varepsilon^{-1/(N-2)}(1 - r_0 \varepsilon^{1/(N-1)}), \varepsilon^{-1/(N-2)}(1 - r_1 \varepsilon^{1/(N-1)})]$$

for some constants $r_1 > r_0 > 0$, and that

$$L_0 \leq \Lambda \leq L_1$$

for some constants $L_1 > L_0 > 0$.

We will prove [Theorem 1.1](#) by verifying the following result.

Theorem 1.3. *Under the same assumptions as [Theorem 1.1](#), there exists $\varepsilon_0 > 0$ such that for $\varepsilon \in (0, \varepsilon_0)$, problem (1-4) has a solution u_ε of the form*

$$u_\varepsilon = W_{r_\varepsilon, \Lambda_\varepsilon} + \phi_\varepsilon,$$

where $\phi_\varepsilon \in \mathcal{H}_s$, $\|\phi_\varepsilon\|_{L^\infty} \rightarrow 0$ as $\varepsilon \rightarrow 0^+$, $L_0 \leq \Lambda_\varepsilon \leq L_1$ and

$$r_\varepsilon \in [\varepsilon^{-1/(N-2)}(1 - r_0\varepsilon^{1/(N-1)}), \varepsilon^{-1/(N-2)}(1 - r_1\varepsilon^{1/(N-1)})].$$

Remark 1.4. In our result, the number of peaks k of the solution u_ε varies with the parameter ε at the order $\varepsilon^{-1/(N-1)}$ when $\varepsilon \rightarrow 0^+$. This is a new phenomenon for the Hénon equation and is in contrast to the subcritical or critical case. For example, in [\[Peng 2006\]](#), where $\varepsilon < 0$, it was proved that for any prescribed integer $k > 0$, there exists $\varepsilon_0 > 0$ such that for any $\varepsilon \in (-\varepsilon_0, 0)$, problem (1-4) has a solution which has exactly k peaks.

Remark 1.5. The results of this paper can be considered as a perturbation of those in [\[Wei and Yan 2013\]](#). In fact the number of bubbles k can be taken to be

$$k = [\delta^{-1/(N-2)}]$$

for any $|\varepsilon| < \delta \ll 1$. When $\varepsilon = 0$, we recover Wei and Yan's result.

We use a reduction argument to prove [Theorem 1.3](#). More precisely, we follow the method in [\[Wei and Yan 2010b; 2013\]](#) to construct peak solutions for problem (1-4). In those papers, where no parameter appears in the considered problem, Wei and Yan used k , the number of peaks of the solutions, as the parameter to construct infinitely many positive peak solutions. This idea is very novel and effective for obtaining infinitely many solutions to several types of problems; see [\[Peng and Wang 2013; Wei and Yan 2010a; 2011\]](#). Unlike the situation in [\[Wei and Yan 2010b; 2013\]](#), here we deal with the supercritical case; we cannot use the variational argument. Instead, we will use the Fredholm theory of compact operators in a suitable Banach space and will employ a direct technique to eliminate the Lagrange multipliers caused from the reduction procedure. Another aspect that differs from [\[Wei and Yan 2010b; 2013\]](#) is that, as we mentioned before, in our proof, we use ε as the parameter in the construction of peak solutions, but in this paper the number of peaks depends on the parameter ε . As a final remark, we point out that for $\alpha = 0$, del Pino, Felmer and Musso [\[2003\]](#) have constructed two-peaked solutions for problem (1-1) in a special domain. Hence, we believe that the effect of the weight $|y|^\alpha$ on the existence of solutions is something like that of the domain.

This paper has the following structure. In [Section 2](#), we carry out the finite-dimensional reduction procedure. The main results will be proved in [Section 3](#). We put the energy expansion and some basic estimates used in [Sections 2 and 3](#) in [Appendices A and B](#).

2. Finite-dimensional reduction

In this section, we perform a finite-dimensional reduction. Let

$$(2-1) \quad \|u\|_* = \sup_{y \in B_*(0)} \left(\sum_{i=1}^k \frac{1}{(1 + |y - x_i|)^{\frac{1}{2}(N-2)+\tau}} \right)^{-1} |u(y)|$$

and

$$(2-2) \quad \|v\|_{**} = \sup_{y \in B_*(0)} \left(\sum_{i=1}^k \frac{1}{(1 + |y - x_i|)^{\frac{1}{2}(N+2)+\tau}} \right)^{-1} |v(y)|,$$

where $\tau = (N-2)/(N-1)$. We denote by L_*^∞ and L_{**}^∞ the function spaces defined on $B_*(0)$ with finite $\|\cdot\|_*$ and $\|\cdot\|_{**}$ norm, respectively.

Let

$$Z_{i,1} = \frac{\partial P U_{x_i, \Lambda}}{\partial r}, \quad Z_{i,2} = \frac{\partial P U_{x_i, \Lambda}}{\partial \Lambda}.$$

First, we consider the linear problem

$$(2-3) \quad \begin{cases} -\Delta \phi - (p+\varepsilon) K(\varepsilon^{\frac{1}{N-2}} |y|) W_{r, \Lambda}^{p-1+\varepsilon} \phi = h + \sum_{j=1}^2 c_j \sum_{i=1}^k U_{x_i, \Lambda}^{p-1} Z_{i,j} & \text{in } B_*(0), \\ \phi \in \mathcal{H}_s, \quad \left\langle \sum_{i=1}^k U_{x_i, \Lambda}^{p-1} Z_{i,l}, \phi \right\rangle = 0, & l = 1, 2, \end{cases}$$

for some numbers c_i , where

$$\langle u, v \rangle = \int_{B_*(0)} uv.$$

Lemma 2.1. *Assume there is a sequence $\varepsilon = \varepsilon_n \rightarrow 0$ such that ϕ_ε solves (2-3) for $h = h_\varepsilon$. If $\|h_\varepsilon\|_{**}$ goes to zero as ε goes to zero, so does $\|\phi_\varepsilon\|_*$.*

Proof. The proof of this lemma is very similar to the proof of Lemma 2.1 in [Wei and Yan 2013].

We argue by contradiction. Suppose that there are $\varepsilon \rightarrow 0$, $h = h_\varepsilon$, $\Lambda_\varepsilon \in [L_0, L_1]$ and $r_\varepsilon \in [\varepsilon^{-1/(N-2)}(1 - r_0 \varepsilon^{1/(N-1)}), \varepsilon^{-1/(N-2)}(1 - r_1 \varepsilon^{1/(N-1)})]$ such that ϕ_ε solves (2-3) for $h = h_\varepsilon$, $\Lambda = \Lambda_\varepsilon$, $r = r_\varepsilon$ with $\|h_\varepsilon\|_{**} \rightarrow 0$ and $\|\phi_\varepsilon\|_* \geq c > 0$. Without loss of generality, we may assume that $\|\phi_\varepsilon\|_* = 1$.

Now rewrite (2-3) in the following integral form:

$$\begin{aligned} \phi_\varepsilon(y) = & (p + \varepsilon) \int_{B_*(0)} G_\varepsilon(y, z) K(\varepsilon^{-1/(N-2)} |z|) W_{r, \Lambda}^{p-1+\varepsilon}(z) \phi_\varepsilon(z) dz \\ & + \int_{B_*(0)} G_\varepsilon(y, z) \left(h_\varepsilon(z) + \sum_{j=1}^2 c_j \sum_{i=1}^k Z_{i,j}(z) U_{x_i, \Lambda}^{p-1}(z) \right) dz. \end{aligned}$$

By [Lemma B.3](#), we find

$$\begin{aligned}
& \left| (p + \varepsilon) \int_{B_*(0)} G_\varepsilon(y, z) K(\varepsilon^{-1/(N-2)}|z|) W_{r,\Lambda}^{p-1+\varepsilon}(z) \phi_\varepsilon(z) dz \right| \\
& \leq (p + \varepsilon) \int_{B_*(0)} \frac{1}{|y - z|^{N-2}} K(\varepsilon^{-1/(N-2)}|z|) W_{r,\Lambda}^{p-1+\varepsilon}(z) |\phi_\varepsilon(z)| dz \\
& \leq C \|\phi_\varepsilon\|_* \int_{B_*(0)} \frac{1}{|y - z|^{N-2}} W_{r,\Lambda}^{p-1+\varepsilon} \sum_{j=1}^k \frac{1}{(1 + |z - x_j|)^{\frac{1}{2}(N-2)+\tau}} dz \\
& \leq C \|\phi_\varepsilon\|_* \sum_{j=1}^k \frac{1}{(1 + |z - x_j|)^{\frac{1}{2}(N-2)+\tau+\vartheta}}.
\end{aligned}$$

It follows from [Lemma B.2](#) that

$$\begin{aligned}
\left| \int_{B_*(0)} G_\varepsilon(y, z) h_\varepsilon(z) dz \right| & \leq \int_{B_*(0)} \frac{1}{|y - z|^{N-2}} |h_\varepsilon(z)| dz \\
& \leq C \|h_\varepsilon\|_{**} \sum_{j=1}^k \frac{1}{(1 + |z - x_j|)^{\frac{1}{2}(N-2)+\tau}}
\end{aligned}$$

and

$$\left| \int_{B_*(0)} G_\varepsilon(y, z) \sum_{j=1}^k Z_{i,l}(z) U_{x_i,\Lambda}^{p-1}(z) dz \right| \leq C \sum_{j=1}^k \frac{1}{(1 + |z - x_j|)^{\frac{1}{2}(N-2)+\tau}}.$$

Next, we estimate c_ℓ , $\ell = 1, 2$. Multiplying (2-3) by $Z_{1,t}$ and integrating, we obtain that c_ℓ satisfies

$$\begin{aligned}
(2-4) \quad & \sum_{\ell=1}^2 \sum_{i=1}^k \langle U_{x_i,\Lambda}^{p-1} Z_{i,\ell}, Z_{1,t} \rangle c_\ell \\
& = \langle -\Delta \phi_\varepsilon - (p + \varepsilon) K(\varepsilon^{-1/(N-2)}|y|) W_{r,\Lambda}^{p-1+\varepsilon} \phi_\varepsilon, Z_{1,t} \rangle - \langle h_\varepsilon, Z_{1,t} \rangle.
\end{aligned}$$

It follows from [Lemma B.1](#) that

$$\begin{aligned}
|\langle h_\varepsilon, Z_{1,\ell} \rangle| & \leq C \|h_\varepsilon\|_{**} \int_{\mathbb{R}^N} \frac{1}{(1 + |z - x_1|)^{N-2}} \sum_{j=1}^k \frac{1}{(1 + |z - x_j|)^{\frac{1}{2}(N+2)+\tau}} dz \\
& \leq C \|h_\varepsilon\|_{**}.
\end{aligned}$$

On the other hand, using [Lemma B.3](#), we obtain

$$\begin{aligned}
& \langle -\Delta \phi_\varepsilon - (p + \varepsilon) K(\varepsilon^{1/(N-2)}|y|) W_{r,\Lambda}^{p-1+\varepsilon} \phi_\varepsilon, Z_{1,\ell} \rangle \\
& = \langle -\Delta Z_{1,\ell} - (p + \varepsilon) K(\varepsilon^{1/(N-2)}|y|) W_{r,\Lambda}^{p-1+\varepsilon} Z_{1,\ell}, \phi_\varepsilon \rangle
\end{aligned}$$

$$\begin{aligned}
&= \langle -\Delta Z_{1,\ell} - pK(\varepsilon^{1/(N-2)}|y|)W_{r,\Lambda}^{p-1}Z_{1,\ell}, \phi_\varepsilon \rangle \\
&\quad + p \langle K(\varepsilon^{1/(N-2)}|y|)(W_{r,\Lambda}^{p-1} - W_{r,\Lambda}^{p-1+\varepsilon})Z_{1,\ell}, \phi_\varepsilon \rangle \\
&\quad - \varepsilon \langle K(\varepsilon^{1/(N-2)}|y|)W_{r,\Lambda}^{p-1+\varepsilon}Z_{1,\ell}, \phi_\varepsilon \rangle \\
&= o(\|\phi_\varepsilon\|_*).
\end{aligned}$$

However, there is a constant $c' > 0$ such that

$$\sum_{i=1}^k \langle U_{x_i,\Lambda}^{p-1}Z_{i,t}, Z_{1,\ell} \rangle = (c' + o(1))\delta_{t\ell}.$$

Hence we find from (2-4) that

$$c_\ell = o(\|\phi_\varepsilon\|_*) + O(\|h_\varepsilon\|_{**}).$$

Therefore,

$$\begin{aligned}
(2-5) \quad \|\phi_\varepsilon\|_* &\leq o(1) + \|h_\varepsilon\|_{**} + \frac{\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N-2)+\tau+\vartheta}}}{\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N-2)+\tau}}}.
\end{aligned}$$

Noting that $\|\phi_\varepsilon\|_* = 1$, we obtain from (2-5) that there is $R > 0$ such that

$$(2-6) \quad \|\phi_\varepsilon(y)\|_{L^\infty(B_R(x_i))} \geq a > 0 \quad \text{for some } i.$$

Furthermore, for this particular i , the translated version $\bar{\phi}_\varepsilon(y) = \phi_\varepsilon(y - x_i)$ converges uniformly on any compact set to a solution u of

$$(2-7) \quad -\Delta u - pU_{0,\Lambda}^{p-1}u = 0 \quad \text{in } \mathbb{R}^N$$

for some $\Lambda \in [L_0, L_1]$. Since u is perpendicular to the kernel of (2-7), we have $u \equiv 0$, which contradicts $\|u(y)\|_{L^\infty(B_R(x_i))} \geq a > 0$. \square

The following proposition is a direct consequence of combining Proposition 4.1 in [del Pino et al. 2003] with Lemma 2.1.

Proposition 2.2. *There exists $\varepsilon_0 > 0$ and a constant $C > 0$ such that for all $\varepsilon \leq \varepsilon_0$ and all $h_\varepsilon \in L_{**}^\infty$, problem (2-3) has a unique solution $\phi_\varepsilon \equiv \mathcal{L}_\varepsilon(h_\varepsilon) \in L_*^\infty$. Moreover,*

$$(2-8) \quad \|\mathcal{L}_\varepsilon(h_\varepsilon)\|_* \leq C\|h_\varepsilon\|_{**}, \quad |c_\ell| \leq C\|h_\varepsilon\|_{**}.$$

In order to prove the main theorem, we will prove that problem (1-4) admits a solution of the form $u = W_{r,\Lambda} + \phi$, where $W_{r,\Lambda} = \sum_{j=1}^k P U_{x_j,\Lambda}$ and $\phi \in \mathcal{H}_s$ is small and satisfies $\langle U_{x_i,\Lambda}^{p-1} Z_{i,\ell}, \phi \rangle = 0, i = 1, 2, \dots, k, \ell = 1, 2$.

We consider the perturbation problem

$$(2-9) \quad \begin{cases} -\Delta(W_{r,\Lambda} + \phi) = K(\varepsilon^{\frac{1}{N-2}} |y|)(W_{r,\Lambda} + \phi)^{p+\varepsilon} + \sum_{\ell=1}^2 c_\ell \sum_{i=1}^k U_{x_i,\Lambda}^{p-1} Z_{i,\ell} & \text{in } B_*(0), \\ \phi \in \mathcal{H}_s, \quad \left\langle \sum_{i=1}^k U_{x_i,\Lambda}^{p-1} Z_{i,\ell}, \phi \right\rangle = 0, & \ell = 1, 2. \end{cases}$$

Proposition 2.3. *There is $\varepsilon_0 > 0$ such that for any $\varepsilon \leq \varepsilon_0$, any $\Lambda \in [L_0, L_1]$, and*

$$r \in \left[\varepsilon^{-\frac{1}{N-2}} (1 - r_0 \varepsilon^{\frac{1}{N-1}}), \varepsilon^{-\frac{1}{N-2}} (1 - r_1 \varepsilon^{\frac{1}{N-1}}) \right],$$

problem (2-9) has a unique solution $\phi = \phi_{r,\Lambda}$ satisfying

$$\|\phi\|_* \leq C \varepsilon^{(\frac{1}{2} + \sigma)/(N-2)}, \quad |c_\ell| \leq C \varepsilon^{(\frac{1}{2} + \sigma)/(N-2)},$$

where $\sigma > 0$ is a small constant.

Let

$$\begin{aligned} N_\varepsilon(\phi) &= K(\varepsilon^{\frac{1}{N-2}} |y|) \left((W_{r,\Lambda} + \phi)^{p+\varepsilon} - W_{r,\Lambda}^{p+\varepsilon} - (p + \varepsilon) W_{r,\Lambda}^{p-1+\varepsilon} \phi \right), \\ l_\varepsilon &= K(\varepsilon^{\frac{1}{N-2}} |y|) W_{r,\Lambda}^{p+\varepsilon} - \sum_{j=1}^k U_{x_j,\Lambda}^p. \end{aligned}$$

Then problem (2-9) can be written as

$$(2-10) \quad \begin{cases} -\Delta\phi - (p + \varepsilon) K(\varepsilon^{\frac{1}{N-2}} |y|) W_{r,\Lambda}^{p-1+\varepsilon} \phi \\ \quad \quad \quad = N_\varepsilon(\phi) + l_\varepsilon + \sum_{\ell=1}^2 c_\ell \sum_{i=1}^k U_{x_i,\Lambda}^{p-1} Z_{i,\ell} & \text{in } B_*(0), \\ \phi \in \mathcal{H}_s, \quad \left\langle \sum_{i=1}^k U_{x_i,\Lambda}^{p-1} Z_{i,\ell}, \phi \right\rangle = 0, & \ell = 1, 2. \end{cases}$$

We will use the contraction mapping theorem to prove that problem (2-9) is uniquely solvable under the condition that $\|\phi\|_*$ is small enough. So we need to estimate $N_\varepsilon(\phi)$ and l_ε .

Lemma 2.4. *If $N \geq 4$, then*

$$\|N_\varepsilon(\phi)\|_{**} \leq C \|\phi\|_*^{\min\{p+\varepsilon, 2\}}.$$

Proof. We have

$$|N_\varepsilon(\phi)| \leq \begin{cases} C |\phi|^{p+\varepsilon}, & N \geq 7, \\ C (W_{r,\Lambda}^{p-2+\varepsilon} \phi^2 + |\phi|^{p+\varepsilon}), & N = 4, 5, 6. \end{cases}$$

Firstly, we consider $N \geq 7$. By the Hölder inequality, we have

$$\begin{aligned}
 |N_\varepsilon(\phi)| &\leq C \|\phi\|_*^{p+\varepsilon} \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N-2)+\tau}} \right)^{p+\varepsilon} \\
 &\leq C \|\phi\|_*^{p+\varepsilon} \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}} \right) \\
 &\quad \times \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{p+\varepsilon}{p+\varepsilon-1}(\frac{1}{2}(N-2)+\tau) - \frac{1}{p+\varepsilon-1}(\frac{1}{2}(N+2)+\tau)}} \right)^{p+\varepsilon-1} \\
 &\leq C \|\phi\|_*^{p+\varepsilon} \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}} \right).
 \end{aligned}$$

Thus, the result follows.

Suppose that $N = 4, 5, 6$. Using the fact that $N-2 > \frac{1}{2}(N-2) + \tau$, we get

$$\begin{aligned}
 |N_\varepsilon(\phi)| &\leq C \|\phi\|_*^2 \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{N-2}} \right)^{p-2+\varepsilon} \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N-2)+\tau}} \right)^2 \\
 &\quad + C \|\phi\|_*^{p+\varepsilon} \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}} \right) \\
 &\leq C \|\phi\|_*^2 \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N-2)+\tau}} \right)^{p+\varepsilon} \\
 &\quad + C \|\phi\|_*^{p+\varepsilon} \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}} \right) \\
 &\leq C \|\phi\|_*^2 \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}} \right).
 \end{aligned}$$

So we have proved that for $N \geq 4$,

$$\|N_\varepsilon(\phi)\|_{**} \leq C \|\phi\|_*^{\min\{p+\varepsilon, 2\}}.$$

□

Lemma 2.5. Assume that $r \in [\varepsilon^{-\frac{1}{N-2}}(1-r_0\varepsilon^{\frac{1}{N-1}}), \varepsilon^{-\frac{1}{N-2}}(1-r_1\varepsilon^{\frac{1}{N-1}})]$. If $N \geq 4$, then

$$\|I_\varepsilon\|_{**} \leq C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)}.$$

Proof. Define

$$\Omega_j = \left\{ y : y = (y', y'') \in B_{\varepsilon^{-1/(N-2)}}(0), \left\langle \frac{y'}{|y'|}, \frac{x_j}{|x_j|} \right\rangle \geq \cos \frac{\pi}{k} \right\}.$$

We have

$$\begin{aligned}
 l_\varepsilon &= K(\varepsilon^{1/(N-2)}|y|)(W_{r,\Lambda}^{p+\varepsilon} - W_{r,\Lambda}^p) + K(\varepsilon^{1/(N-2)}|y|)\left(W_{r,\Lambda}^p - \sum_{j=1}^k P U_{x_j,\Lambda}^p\right) \\
 &\quad + K(\varepsilon^{1/(N-2)}|y|)\left(\sum_{j=1}^k P U_{x_j,\Lambda}^p - \sum_{j=1}^k U_{x_j,\Lambda}^p\right) + \sum_{j=1}^k U_{x_j,\Lambda}^p (K(\varepsilon^{1/(N-2)}|y|) - 1) \\
 &=: J_0 + J_1 + J_2 + J_3.
 \end{aligned}$$

Estimate of J_0 .

$$\begin{aligned}
 |J_0| &\leq C\varepsilon W_{r,\Lambda}^p |\ln W_{r,\Lambda}| \\
 &\leq C\varepsilon \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{N-2}} \right)^p \ln \sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{N-2}} \\
 &\leq C\varepsilon \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}} \right) \left(\sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{N+2}{4}(\frac{N-2}{2}-\frac{N-2}{N+2}\tau)}} \right)^{\frac{4}{N-2}} \\
 &\leq C\varepsilon \sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}}.
 \end{aligned}$$

Estimate of J_1 . From the symmetry, we can assume that $y \in \Omega_1$. Then,

$$|y-x_j| \geq |y-x_1|, \quad y \in \Omega_1, \quad j \neq 1.$$

Firstly, we claim

$$(2-11) \quad \frac{1}{1+|y-x_j|} \leq \frac{C}{|x_j-x_1|}, \quad y \in \Omega_1, \quad j \neq 1.$$

In fact, if $|y-x_1| \leq \frac{1}{2}|x_1-x_j|$, then $|y-x_j| \geq \frac{1}{2}|x_j-x_1|$. If $|y-x_1| \geq \frac{1}{2}|x_j-x_1|$, then $|y-x_j| \geq |y-x_1| \geq \frac{1}{2}|x_j-x_1|$.

It's easy to verify that

$$|J_1| \leq C \frac{1}{(1+|y-x_1|)^4} \sum_{j=2}^k \frac{1}{(1+|y-x_j|)^{N-2}} + C \left(\sum_{j=2}^k \frac{1}{(1+|y-x_j|)^{N-2}} \right)^p.$$

Using (2-11), taking $1 < \varrho \leq N-2$, we obtain for $y \in \Omega_1$,

$$\frac{1}{(1+|y-x_1|)^4} \frac{1}{(1+|y-x_j|)^{N-2}} \leq C \frac{1}{(1+|y-x_1|)^{N+2-\varrho}} \frac{1}{|x_j-x_1|^\varrho}, \quad j \neq 1.$$

Take $\varrho > \max\{\frac{1}{2}(N-1), 1\}$ satisfying $N+2-\varrho \geq \frac{1}{2}(N+2)+\tau$. Then

$$\begin{aligned} \frac{1}{(1+|y-x_1|)^4} \sum_{j=2}^k \frac{1}{(1+|y-x_j|)^{N-2}} &\leq \frac{C}{(1+|y-x_1|)^{N+2-\varrho}} (k\varepsilon^{1/(N-2)})^\varrho \\ &= \frac{C}{(1+|y-x_1|)^{N+2-\varrho}} \varepsilon^{\varrho/((N-1)(N-2))} \leq \frac{C}{(1+|y-x_1|)^{\frac{1}{2}(N+2)+\tau}} \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)}. \end{aligned}$$

By the Hölder inequality, we find

$$\begin{aligned} \left(\sum_{j=2}^k \frac{1}{(1+|y-x_j|)^{N-2}} \right)^p &\leq \sum_{j=2}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}} \left(\sum_{j=2}^k \frac{1}{(1+|y-x_j|)^{\frac{N+2}{4}(\frac{N-2}{2}-\tau\frac{N-2}{N+2})}} \right)^{\frac{4}{N-2}}. \end{aligned}$$

Noticing that $\frac{N+2}{N-2} \left(\frac{N-2}{2} - \tau \frac{N-2}{N+2} \right) > \frac{N-1}{2}$ if $N \geq 4$, we deduce that

$$\begin{aligned} \left(\sum_{j=2}^k \frac{1}{(1+|y-x_j|)^{N-2}} \right)^p &\leq C \left(\sum_{j=2}^k \frac{1}{|x_j-x_1|^{\frac{N+2}{4}(\frac{N-2}{2}-\tau\frac{N-2}{N+2})}} \right)^{\frac{4}{N-2}} \sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}} \\ &\leq C (k\varepsilon^{1/(N-2)})^{\frac{N+2}{N-2}(\frac{N-2}{2}-\tau\frac{N-2}{N+2})} \sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}} \\ &\leq C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)} \sum_{j=1}^k \frac{1}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}}. \end{aligned}$$

Hence, we conclude that if $N \geq 4$,

$$\|J_1\|_{**} \leq C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)}.$$

Estimate of J_2 . Let $H(y, x)$ be the regular part of the Green function for $-\Delta$ in $B_1(0)$ with the zero boundary condition. Let \bar{x}_j^* be the reflection point of \bar{x}_j with respect to $\partial B_1(0)$. Then

$$\varepsilon H(\bar{y}, \bar{x}_j) = \frac{C\varepsilon}{|\bar{y} - \bar{x}_j^*|^{N-2}} \leq \frac{C}{(1+|y-x_j|)^{N-2}}.$$

By direct calculation, we have

$$\begin{aligned}
 |J_2| &\leq C \sum_{j=1}^k \frac{C\varepsilon}{(1+|y-x_j|)^4} H(\bar{y}, \bar{x}_j) \\
 &\leq \sum_{j=1}^k \frac{C}{(1+|y-x_j|)^{4+(1-\gamma)(N-2)}} (\varepsilon H(\bar{y}, \bar{x}_j))^\gamma \\
 &\leq C\varepsilon^{\gamma/(N-1)} \sum_{j=1}^k \frac{C}{(1+|y-x_j|)^{4+(1-\gamma)(N-2)}} \\
 &\leq C\varepsilon^{(\frac{1}{2}+\sigma)/(N-2)} \sum_{j=1}^k \frac{C}{(1+|y-x_j|)^{\frac{1}{2}(N+2)+\tau}},
 \end{aligned}$$

where $\gamma > 0$ satisfies $\gamma(N-2)/(N-1) > \frac{1}{2}$ and $4+(1-\gamma)(N-2) \geq \frac{1}{2}(N+2)+\tau$.

Estimate of J_3 . For $y \in \Omega_1$ and $j > 1$, using (2-11), we find

$$U_{x_j, \Lambda}^p \leq \frac{C}{(1+|y-x_1|)^{\frac{1}{2}(N+2)+\tau}} \frac{1}{|x_j-x_1|^{\frac{1}{2}(N+2)-\tau}}.$$

Thus, we have

$$\begin{aligned}
 \left| \sum_{j=2}^k (K(\varepsilon^{1/(N-2)}|y|) - 1) U_{x_j, \Lambda}^p \right| &\leq \frac{C}{(1+|y-x_1|)^{\frac{1}{2}(N+2)+\tau}} \sum_{j=2}^k \frac{1}{|x_j-x_1|^{\frac{1}{2}(N+2)-\tau}} \\
 &\leq \frac{C}{(1+|y-x_1|)^{\frac{1}{2}(N+2)+\tau}} (k\varepsilon^{1/(N-2)})^{\frac{1}{2}(N+2)-\tau} \\
 &\leq \frac{C}{(1+|y-x_1|)^{\frac{1}{2}(N+2)+\tau}} \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)}.
 \end{aligned}$$

If $y \in \Omega_1$ and $||y| - \varepsilon^{-1/(N-2)}| \geq \delta \varepsilon^{-1/(N-2)}$, where $\delta > 0$ is a fixed constant, then

$$||y| - |x_1|| \geq ||y| - \varepsilon^{-1/(N-2)}| - ||x_1| - \varepsilon^{-1/(N-2)}| \geq \frac{1}{2} \delta \varepsilon^{-1/(N-2)}.$$

So, we obtain

$$\left| U_{x_1, \Lambda}^p (K(\varepsilon^{1/(N-2)}|y|) - 1) \right| \leq \frac{C}{(1+|y-x_1|)^{\frac{1}{2}(N+2)+\tau}} \varepsilon^{(\frac{1}{2}(N+2)-\tau)/(N-2)}.$$

If $y \in \Omega_1$ and $\left| |y| - \varepsilon^{-1/(N-2)} \right| \leq \delta \varepsilon^{-1/(N-2)}$, then

$$\begin{aligned} \left| K(\varepsilon^{1/(N-2)}|y|) - 1 \right| &\leq C \left| \varepsilon^{1/(N-2)}|y| - 1 \right| \\ &\leq C \varepsilon^{1/(N-2)} \left(\left| |y| - |x_1| \right| + \left| |x_1| - \varepsilon^{-1/(N-2)} \right| \right) \\ &\leq C \varepsilon^{1/(N-2)} \left| |y| - |x_1| \right| + C \varepsilon^{1/(N-1)} \\ &\leq C \varepsilon^{1/(N-2)} \left| |y| - |x_1| \right| + C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)}, \end{aligned}$$

and

$$\left| |y| - |x_1| \right| \leq \left| |y| - \varepsilon^{-1/(N-2)} \right| + \left| |x_1| - \varepsilon^{-1/(N-2)} \right| \leq 2\delta \varepsilon^{-1/(N-2)}.$$

Since

$$\begin{aligned} \frac{\varepsilon^{1/(N-2)} \left| |y| - |x_1| \right|}{(1 + |y - x_1|)^{N+2}} &\leq C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)} \frac{\left| |y| - |x_1| \right|^{\frac{1}{2}+\sigma}}{(1 + |y - x_1|)^{N+2}} \\ &\leq \frac{C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)}}{(1 + |y - x_1|)^{N+\frac{3}{2}-\sigma}} \leq \frac{C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)}}{(1 + |y - x_1|)^{\frac{1}{2}(N+2)+\tau}}, \end{aligned}$$

we get

$$\left| U_{x_1, \Lambda}^P (K(\varepsilon^{1/(N-2)}|y|) - 1) \right| \leq \frac{C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)}}{(1 + |y - x_1|)^{\frac{1}{2}(N+2)+\tau}}.$$

As a result, we deduce

$$\|J_3\|_{**} \leq C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)}.$$

□

Proof of Proposition 2.3. Recall that

$$k = \lceil \varepsilon^{-1/(N-1)} \rceil, \quad N \geq 4.$$

Let

$$E = \left\{ u \in \mathcal{H}_s \cap L_*^\infty : \|u\|_* \leq \varepsilon^{1/(2(N-2))} \text{ and } \int_{B_*(0)} U_{x_i, \Lambda}^{p-1} Z_{i, \ell} u = 0, \quad i = 1, \dots, k, \quad \ell = 1, 2 \right\}.$$

Then, (2-10) is equivalent to

$$\phi = A_\varepsilon(\phi) =: \mathcal{L}_\varepsilon(N_\varepsilon(\phi)) + \mathcal{L}_\varepsilon(l_\varepsilon),$$

where \mathcal{L}_ε is defined in Proposition 2.2. We will prove that A_ε is a contraction map from E to E . First, $A_\varepsilon(E) \subset E$ because

$$\begin{aligned} \|A_\varepsilon(\phi)\|_* &\leq C \|N_\varepsilon(\phi)\|_{**} + C \|l_\varepsilon\|_{**} \\ &\leq C \|\phi\|_*^{\min\{p+\varepsilon, 2\}} + C \|l_\varepsilon\|_{**} \leq C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)} \leq \varepsilon^{1/(2(N-2))}. \end{aligned}$$

Next we write

$$\|A_\varepsilon(\phi_1) - A_\varepsilon(\phi_2)\|_* = \|\mathcal{L}_\varepsilon(N_\varepsilon(\phi_1)) - \mathcal{L}_\varepsilon(N_\varepsilon(\phi_2))\|_* \leq C\|N_\varepsilon(\phi_1) - N_\varepsilon(\phi_2)\|_{**}.$$

If $N \geq 7$, then

$$|N'_\varepsilon(t)| \leq C|t|^{p-1+\varepsilon}.$$

Thus, we have

$$\begin{aligned} & |N_\varepsilon(\phi_1) - N_\varepsilon(\phi_2)| \\ & \leq C(|\phi_1|^{p-1+\varepsilon} + |\phi_2|^{p-1+\varepsilon})|\phi_1 - \phi_2| \\ & \leq C(\|\phi_1\|_*^{p-1+\varepsilon} + \|\phi_2\|_*^{p-1+\varepsilon})\|\phi_1 - \phi_2\|_* \left(\sum_{j=1}^k \frac{1}{(1 + |y - x_j|)^{\frac{1}{2}(N-2)+\tau}} \right)^{p+\varepsilon} \\ & \leq C(\|\phi_1\|_*^{p-1+\varepsilon} + \|\phi_2\|_*^{p-1+\varepsilon})\|\phi_1 - \phi_2\|_* \sum_{j=1}^k \frac{1}{(1 + |y - x_j|)^{\frac{1}{2}(N+2)+\tau}}. \end{aligned}$$

As a consequence,

$$\begin{aligned} \|A_\varepsilon(\phi_1) - A_\varepsilon(\phi_2)\|_* & \leq C\|N_\varepsilon(\phi_1) - N_\varepsilon(\phi_2)\|_{**} \\ & \leq C(\|\phi_1\|_*^{p-1+\varepsilon} + \|\phi_2\|_*^{p-1+\varepsilon})\|\phi_1 - \phi_2\|_* \leq \frac{1}{2}\|\phi_1 - \phi_2\|_*. \end{aligned}$$

For $N = 4, 5, 6$,

$$|N'_\varepsilon(t)| \leq CW_{r,\Lambda}^{p-2+\varepsilon}|t| + C|t|^{p-1+\varepsilon}.$$

So we have

$$\begin{aligned} & |N_\varepsilon(\phi_1) - N_\varepsilon(\phi_2)| \\ & \leq C(|\phi_1|^{p-1+\varepsilon} + |\phi_2|^{p-1+\varepsilon})|\phi_1 - \phi_2| + C(|\phi_1| + |\phi_2|)W_{r,\Lambda}^{p-2+\varepsilon}|\phi_1 - \phi_2| \\ & \leq C(\|\phi_1\|_*^{p-1+\varepsilon} + \|\phi_2\|_*^{p-1+\varepsilon})\|\phi_1 - \phi_2\|_* \left(\sum_{j=1}^k \frac{1}{(1 + |y - x_j|)^{\frac{1}{2}(N-2)+\tau}} \right)^{p+\varepsilon} \\ & \quad + C(\|\phi_1\|_* + \|\phi_2\|_*)\|\phi_1 - \phi_2\|_* W_{r,\Lambda}^{p-2+\varepsilon} \left(\sum_{j=1}^k \frac{1}{(1 + |y - x_j|)^{\frac{1}{2}(N-2)+\tau}} \right)^2 \\ & \leq C(\|\phi_1\|_* + \|\phi_2\|_*)\|\phi_1 - \phi_2\|_* \sum_{j=1}^k \frac{1}{(1 + |y - x_j|)^{\frac{1}{2}(N+2)+\tau}}. \end{aligned}$$

In either case, we see that A_ε is a contraction map. By the contraction mapping theorem, there is a unique $\phi \in E$ such that

$$\phi = A_\varepsilon(\phi).$$

Moreover, it follows from [Proposition 2.2](#) that

$$\|\phi\|_* \leq C \|l_\varepsilon\|_{**} + C \|N_\varepsilon(\phi)\|_{**} \leq C \|l_\varepsilon\|_{**} + C \|\phi\|_*^{\min\{p+\varepsilon, 2\}},$$

which implies

$$\|\phi\|_* \leq C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)}, \quad |c_\ell| \leq C \varepsilon^{(\frac{1}{2}+\sigma)/(N-2)}. \quad \square$$

3. Proof of the main results

In this section, we will choose (r, Λ) such that $W_{r,\Lambda} + \phi_{r,\Lambda}$ is a solution of [\(1-4\)](#), where $\phi_{r,\Lambda}$ is the map obtained in [Proposition 2.3](#).

Lemma 3.1. *If (r, Λ) satisfies*

$$(3-1) \quad \int_{B_*(0)} \left(\nabla(W_{r,\Lambda} + \phi_{r,\Lambda}) \nabla \frac{\partial W_{r,\Lambda}}{\partial r} - K(\varepsilon^{1/(N-2)}|y|)(W_{r,\Lambda} + \phi_{r,\Lambda})^{p+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial r} \right) = 0,$$

$$(3-2) \quad \int_{B_*(0)} \left(\nabla(W_{r,\Lambda} + \phi_{r,\Lambda}) \nabla \frac{\partial W_{r,\Lambda}}{\partial \Lambda} - K(\varepsilon^{1/(N-2)}|y|)(W_{r,\Lambda} + \phi_{r,\Lambda})^{p+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} \right) = 0,$$

then $W_{r,\Lambda} + \phi_{r,\Lambda}$ is a solution of [\(1-4\)](#).

Proof. It follows from [Proposition 2.3](#) that if [\(3-1\)](#) and [\(3-2\)](#) hold, then by symmetry,

$$(3-3) \quad c_1 \left\langle U_{x_1,\Lambda}^{p-1} \frac{\partial P U_{x_1,\Lambda}}{\partial r}, \frac{\partial W_{r,\Lambda}}{\partial r} \right\rangle = 0 = c_2 \left\langle U_{x_1,\Lambda}^{p-1} \frac{\partial P U_{x_1,\Lambda}}{\partial \Lambda}, \frac{\partial W_{r,\Lambda}}{\partial \Lambda} \right\rangle,$$

which implies that $c_1 = c_2 = 0$. Hence $W_{r,\Lambda} + \phi_{r,\Lambda}$ is a solution of [\(1-4\)](#). \square

In the rest of this section, we need to solve [\(3-1\)](#) and [\(3-2\)](#).

Proposition 3.2. *Equations [\(3-1\)](#) and [\(3-2\)](#) are equivalent to*

$$(3-4) \quad -\frac{\varepsilon H(\bar{x}_1, \bar{x}_1)}{\Lambda^{N-1}} + \sum_{i=2}^k \frac{\varepsilon G(\bar{x}_i, \bar{x}_1)}{\Lambda^{N-1}} + O(\varepsilon^{(1+\sigma)/(N-2)}) = 0$$

and

$$(3-5) \quad \frac{B_2 \varepsilon}{\Lambda^{N-2}} \frac{\partial H(\bar{x}_1, \bar{x}_1)}{\partial d} + B_3 K'(1) + \sum_{i=2}^k \frac{B_2 \varepsilon}{\Lambda^{N-2}} \frac{\partial G(\bar{x}_i, \bar{x}_1)}{\partial d} + O(\varepsilon^{\sigma/(N-2)}) = 0,$$

respectively, where $d = 1 - \varepsilon^{1/(N-2)}r$, B_1 , B_2 and B_3 are the same positive constants as in [Proposition A.1](#) and $\sigma > 0$ is a small constant.

Proof. Here we prove only the first one. The second can be proved similarly by noting that $\partial/\partial d = -\varepsilon^{-1/(N-2)}\partial/\partial r$.

First, we see that

$$\int_{B_*(0)} \nabla(W_{r,\Lambda} + \phi_{r,\Lambda}) \nabla \frac{\partial W_{r,\Lambda}}{\partial \Lambda} = \int_{B_*(0)} \nabla W_{r,\Lambda} \nabla \frac{\partial W_{r,\Lambda}}{\partial \Lambda},$$

and

$$\begin{aligned} & \int_{B_*(0)} K(\varepsilon^{1/(N-2)}|y|)(W_{r,\Lambda} + \phi_{r,\Lambda})^{p+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} \\ &= \int_{B_*(0)} K(\varepsilon^{1/(N-2)}|y|) W_{r,\Lambda}^{p+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} \\ &\quad + (p+\varepsilon) \int_{B_*(0)} K(\varepsilon^{1/(N-2)}|y|) W_{r,\Lambda}^{p-1+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} \phi_{r,\Lambda} \\ &\quad + O\left(\int_{B_*(0)} W_{r,\Lambda}^{p-1+\varepsilon} |\phi_{r,\Lambda}|^2\right). \end{aligned}$$

On the other hand, noticing that $\phi_{r,\Lambda} \in E$, we have

$$\begin{aligned} & \int_{B_*(0)} K(\varepsilon^{1/(N-2)}|y|) W_{r,\Lambda}^{p-1+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} \phi_{r,\Lambda} \\ &= \int_{B_*(0)} K(\varepsilon^{1/(N-2)}|y|) \left(W_{r,\Lambda}^{p-1+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} - \sum_{j=1}^k U_{x_j,\Lambda}^{p-1} \frac{\partial U_{x_j,\Lambda}}{\partial \Lambda} \right) \phi_{r,\Lambda} \\ &\quad + \sum_{j=1}^k \int_{B_*(0)} (K(\varepsilon^{1/(N-2)}|y|) - 1) U_{x_j,\Lambda}^{p-1} \frac{\partial U_{x_j,\Lambda}}{\partial \Lambda} \phi_{r,\Lambda} \\ &= k \int_{\Omega_1} K(\varepsilon^{1/(N-2)}|y|) \left(W_{r,\Lambda}^{p-1+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} - \sum_{j=1}^k U_{x_j,\Lambda}^{p-1} \frac{\partial U_{x_j,\Lambda}}{\partial \Lambda} \right) \phi_{r,\Lambda} \\ &\quad + k \int_{\Omega_1} (K(\varepsilon^{1/(N-2)}|y|) - 1) U_{x_1,\Lambda}^{p-1} \frac{\partial U_{x_1,\Lambda}}{\partial \Lambda} \phi_{r,\Lambda}, \end{aligned}$$

$$\begin{aligned} & \left| \int_{\Omega_1} K(\varepsilon^{1/(N-2)}|y|) \left(W_{r,\Lambda}^{p-1+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} - \sum_{j=1}^k U_{x_j,\Lambda}^{p-1} \frac{\partial U_{x_j,\Lambda}}{\partial \Lambda} \right) \phi_{r,\Lambda} \right| \\ &\leq C \int_{\Omega_1} \left(U_{x_1,\Lambda}^{p-1} (U_{x_1,\Lambda} - P U_{x_1,\Lambda}) + U_{x_1,\Lambda}^{p-1} \sum_{j=2}^k U_{x_j,\Lambda} + \sum_{j=2}^k U_{x_j,\Lambda}^p \right) |\phi_{r,\Lambda}| \\ &\quad + O\left(\varepsilon \int_{\Omega_1} W_{r,\Lambda}^{p-1} \ln W_{r,\Lambda} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} \phi_{r,\Lambda}\right) \\ &\leq C \varepsilon^{1/(N-2)(1+\sigma)}, \end{aligned}$$

and

$$\begin{aligned}
& \left| \int_{\Omega_1} (K(\varepsilon^{1/(N-2)}|y|) - 1) U_{x_1, \Lambda}^{p-1} \frac{\partial U_{x_1, \Lambda}}{\partial \Lambda} \phi_{r, \Lambda} \right| \\
& \leq \left| \int_{|y| - \varepsilon^{-1/(N-2)} \leq \varepsilon^{-1/(2(N-2))}} (K(\varepsilon^{1/(N-2)}|y|) - 1) U_{x_1, \Lambda}^{p-1} \frac{\partial U_{x_1, \Lambda}}{\partial \Lambda} \phi_{r, \Lambda} \right| \\
& \quad + \left| \int_{|y| - \varepsilon^{-1/(N-2)} \geq \varepsilon^{-1/(2(N-2))}} (K(\varepsilon^{1/(N-2)}|y|) - 1) U_{x_1, \Lambda}^{p-1} \frac{\partial U_{x_1, \Lambda}}{\partial \Lambda} \phi_{r, \Lambda} \right| \\
& \leq C \varepsilon^{1/(N-2)(1+\sigma)}.
\end{aligned}$$

So, we have proved

$$\begin{aligned}
& \int_{B_*(0)} \left(\nabla(W_{r, \Lambda} + \phi_{r, \Lambda}) \nabla \frac{\partial W_{r, \Lambda}}{\partial \Lambda} - K(\varepsilon^{1/(N-2)}|y|) (W_{r, \Lambda} + \phi_{r, \Lambda})^{p+\varepsilon} \frac{\partial W_{r, \Lambda}}{\partial \Lambda} \right) \\
& = \int_{B_*(0)} \left(\nabla W_{r, \Lambda} \nabla \frac{\partial W_{r, \Lambda}}{\partial \Lambda} - K(\varepsilon^{1/(N-2)}|y|) W_{r, \Lambda}^{p+\varepsilon} \frac{\partial W_{r, \Lambda}}{\partial \Lambda} \right) + O(k \varepsilon^{(1+\sigma)/(N-2)}),
\end{aligned}$$

and the result follows from [Proposition A.1](#). \square

Proof of Theorem 1.3. Note (see [\[Wei and Yan 2013\]](#)) that

$$H(\bar{x}_1, \bar{x}_1) = \frac{1}{2^{N-2} d^{N-2}} (1 + O(d))$$

and

$$\frac{a_0}{j^N} + O\left(\frac{d}{j^{N-2}}\right) \leq \frac{1}{k^{N-2}} G(\bar{x}_j, \bar{x}_1) \leq \frac{a_1}{j^N} + O\left(\frac{d}{j^{N-2}}\right),$$

where $a_1 \geq a_0 > 0$. Hence, we find that there is a constant $B_4 > 0$ such that

$$\sum_{j=2}^k G(\bar{x}_j, \bar{x}_1) = k^{N-2} \left(\frac{B_4}{|\bar{x}_1|^{N-2}} + O\left(\frac{1}{k^{N-1}}\right) + O(d) \right) = B_4 k^{N-2} + O(k^{N-2} d).$$

Consequently, (3-4) and (3-5) are equivalent to

$$(3-6) \quad -\frac{A_1 \varepsilon}{\Lambda^{N-1} d^{N-2}} + \frac{A_2 k^{N-2} \varepsilon}{\Lambda^{N-1}} + O(\varepsilon^{(1+\sigma)/(N-2)}) = 0$$

and

$$(3-7) \quad -\frac{A_3 \varepsilon}{\Lambda^{N-2} d^{N-1}} + A_4 + O(\varepsilon^{\sigma/(N-2)}) = 0,$$

respectively, for positive constants A_i , $i = 1, 2, 3, 4$. Recall that $d = 1 - \varepsilon^{1/(N-2)} r$. Define $\eta = dk$. Thus, (3-6) and (3-7) read

$$(3-8) \quad -\frac{A_1}{\Lambda^{N-1} \eta^{N-2}} + \frac{A_2}{\Lambda^{N-1}} + O(\varepsilon^{\sigma/(N-2)}) = 0$$

and

$$(3-9) \quad -\frac{A_3}{\Lambda^{N-2}\eta^{N-1}} + A_4 + O(\varepsilon^{\sigma/(N-2)}) = 0.$$

Let

$$f_1(\eta, \Lambda) = -\frac{A_1}{\Lambda^{N-1}\eta^{N-2}} + \frac{A_2}{\Lambda^{N-1}}$$

and

$$f_2(\eta, \Lambda) = -\frac{A_3}{\Lambda^{N-2}\eta^{N-1}} + A_4.$$

It is easy to check that $f_1 = 0$ and $f_2 = 0$ have a unique solution

$$\eta_0 = \left(\frac{A_1}{A_2}\right)^{\frac{1}{N-2}}, \quad \Lambda_0 = \left(\frac{A_3}{A_4\eta_0^{N-1}}\right)^{\frac{1}{N-2}}.$$

On the other hand, we have

$$\frac{\partial f_1(\eta_0, \Lambda_0)}{\partial \Lambda} = 0, \quad \frac{\partial f_2(\eta_0, \Lambda_0)}{\partial \eta} > 0,$$

and

$$\frac{\partial f_1(\eta_0, \Lambda_0)}{\partial \eta} > 0, \quad \frac{\partial f_2(\eta_0, \Lambda_0)}{\partial \Lambda} > 0.$$

Hence the linear operator of $f_1 = 0$ and $f_2 = 0$ at (η_0, Λ_0) is invertible. Therefore, (3-8) and (3-9) have a solution near (η_0, Λ_0) . \square

Appendix A: Energy expansion

Here and in Appendix B, we assume that

$$x_j = \left(r \cos \frac{2(j-1)\pi}{k}, r \sin \frac{2(j-1)\pi}{k}, 0 \right), \quad j = 1, \dots, k,$$

where $r \in [\varepsilon^{-\frac{1}{N-2}}(1 - r_0\varepsilon^{\frac{1}{N-1}}), \varepsilon^{-\frac{1}{N-2}}(1 - r_1\varepsilon^{\frac{1}{N-1}})]$ and 0 is the zero vector in \mathbb{R}^{N-2} .

Let

$$\bar{x}_j = \varepsilon^{\frac{1}{N-2}} x_j.$$

Let $G(x, y)$ be the Green function of $-\Delta$ in $B_1(0)$ with the Dirichlet boundary and let $H(x, y)$ be the regular part of the Green function.

Recall that

$$k = \left\lceil \varepsilon^{-\frac{1}{N-1}} \right\rceil$$

and

$$W_{r,\Lambda}(y) = \sum_{j=1}^k P U_{x_j, \Lambda}(y),$$

where $PU_{x,\Lambda}$ is the solution of (1-5). Moreover,

$$(A-1) \quad \phi_{x_j,\Lambda}(y) = U_{x_j,\Lambda}(y) - PU_{x_j,\Lambda}(y) = \frac{\varepsilon H(\bar{x}_j, \bar{y})}{\Lambda^{\frac{1}{2}(N-2)}} + O\left(\frac{\varepsilon^{N/(N-2)}}{d^N}\right),$$

where $d = 1 - |\bar{x}_j| = 1 - \varepsilon^{1/(N-2)}|x_j|$.

Proposition A.1. *We have*

$$\begin{aligned} & \int_{B_*(0)} \left(\nabla W_{r,\Lambda} \nabla \frac{\partial W_{r,\Lambda}}{\partial \Lambda} - K(\varepsilon^{1/(N-2)}|y|) W_{r,\Lambda}^{p+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} \right) \\ &= k B_1 \left(-\frac{\varepsilon H(\bar{x}_1, \bar{x}_1)}{\Lambda^{N-1}} + \sum_{i=2}^k \frac{\varepsilon G(\bar{x}_i, \bar{x}_1)}{\Lambda^{N-1}} + O(\varepsilon^{(1+\sigma)/(N-2)}) \right), \end{aligned}$$

and

$$\begin{aligned} & \int_{B_*(0)} \left(\nabla W_{r,\Lambda} \nabla \frac{\partial W_{r,\Lambda}}{\partial r} - K(\varepsilon^{1/(N-2)}|y|) W_{r,\Lambda}^{p+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial r} \right) \\ &= k \left(\frac{B_2 \varepsilon}{\Lambda^{N-2}} \frac{\partial H(\bar{x}_1, \bar{x}_1)}{\partial r} - B_3 K'(1) \varepsilon^{1/(N-2)} \right. \\ & \quad \left. + \sum_{i=2}^k \frac{B_2 \varepsilon}{\Lambda^{N-2}} \frac{\partial G(\bar{x}_i, \bar{x}_1)}{\partial r} + O(\varepsilon^{(1+\sigma)/(N-2)}) \right), \end{aligned}$$

where B_1 , B_2 and B_3 are some positive constants.

Proof. The proof is quite standard now. Here we only prove the first equation. The other one can be obtained similarly.

Using symmetry, we find

$$\begin{aligned} I &:= \int_{B_*(0)} \left(\nabla W_{r,\Lambda} \nabla \frac{\partial W_{r,\Lambda}}{\partial \Lambda} - K(\varepsilon^{1/(N-2)}|y|) W_{r,\Lambda}^{p+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} \right) \\ &= k \left(p \sum_{i=1}^k \int_{B_*(0)} PU_{x_1,\Lambda}^{p-1} \frac{\partial PU_{x_1,\Lambda}}{\partial \Lambda} PU_{x_i,\Lambda} - \int_{\Omega_1} K(\varepsilon^{1/(N-2)}|y|) W_{r,\Lambda}^{p+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} \right). \end{aligned}$$

It is easy to check that for $y \in \Omega_1$,

$$\begin{aligned} \frac{\partial}{\partial \Lambda} W_{r,\Lambda}^{p+1} &= \frac{\partial}{\partial \Lambda} PU_{x_1,\Lambda}^{p+1} + (p+1) \frac{\partial}{\partial \Lambda} \left(PU_{x_1,\Lambda}^p \sum_{i=2}^k PU_{x_i,\Lambda} \right) \\ &\quad + O\left(U_{x_1,\Lambda}^{\frac{1}{2}(p+1)} \left(\sum_{i=2}^k U_{x_i,\Lambda} \right)^{\frac{1}{2}(p+1)} \right). \end{aligned}$$

Thus, we have

$$\begin{aligned}
& (p+1) \int_{\Omega_1} K(\varepsilon^{1/(N-2)}|y|) W_{r,\Lambda}^{p+\varepsilon} \frac{\partial W_{r,\Lambda}}{\partial \Lambda} \\
&= \int_{\Omega_1} K(\varepsilon^{1/(N-2)}|y|) \frac{\partial W_{r,\Lambda}^{p+1}}{\partial \Lambda} + O\left(\varepsilon \int_{\Omega_1} W_{r,\Lambda}^{p+1} \ln W_{r,\Lambda}\right) \\
&= \int_{\Omega_1} K(\varepsilon^{1/(N-2)}|y|) \frac{\partial}{\partial \Lambda} P U_{x_1,\Lambda}^{p+1} \\
&\quad + (p+1) \int_{\Omega_1} K(\varepsilon^{1/(N-2)}|y|) \frac{\partial}{\partial \Lambda} \left(P U_{x_1,\Lambda}^p \sum_{i=2}^k P U_{x_i,\Lambda} \right) \\
&\quad + O\left(\int_{\Omega_1} U_{x_1,\Lambda}^{\frac{1}{2}(p+1)} \left(\sum_{i=2}^k U_{x_i,\Lambda} \right)^{\frac{1}{2}(p+1)} \right) + O\left(\varepsilon \int_{\Omega_1} W_{r,\Lambda}^{p+1} \ln W_{r,\Lambda} \right).
\end{aligned}$$

Note that for $y \in \Omega_1$, $|y - x_i| \geq |y - x_1|$. Using (2-11), we see that for $t \in (1, N-2)$,

$$\sum_{i=2}^k U_{x_i,\Lambda} \leq \frac{C}{(1 + |y - x_1|)^{N-2-t}} \sum_{i=2}^k \frac{1}{|x_i - x_1|^t}.$$

If we take t close to $N-2$, then

$$\int_{\Omega_1} U_{x_1,\Lambda}^{\frac{1}{2}(p+1)} \left(\sum_{i=2}^k U_{x_i,\Lambda} \right)^{\frac{1}{2}(p+1)} = O((k\varepsilon^{1/(N-2)})^{Nt/(N-2)}) = O(\varepsilon^{(1+\sigma)/(N-2)}).$$

Moreover, it is easy to show that

$$\varepsilon \int_{\Omega_1} W_{r,\Lambda}^{p+1} \ln W_{r,\Lambda} = O(\varepsilon).$$

As a result, we obtain

$$\begin{aligned}
I &= k \left(- \int_{\Omega_1} K(\varepsilon^{1/(N-2)}|y|) P U_{x_1,\Lambda}^p \frac{\partial P U_{x_1,\Lambda}}{\partial \Lambda} \right. \\
&\quad - \sum_{i=2}^k \int_{\Omega_1} K(\varepsilon^{1/(N-2)}|y|) P U_{x_1,\Lambda}^p \frac{\partial P U_{x_i,\Lambda}}{\partial \Lambda} \\
&\quad + p \sum_{i=2}^k \int_{\Omega_1} (1 - K(\varepsilon^{1/(N-2)}|y|)) P U_{x_1,\Lambda}^{p-1} \frac{\partial P U_{x_1,\Lambda}}{\partial \Lambda} P U_{x_i,\Lambda} \\
&\quad \left. + p \sum_{i=2}^k \int_{B_*(0) \setminus \Omega_1} P U_{x_1,\Lambda}^{p-1} \frac{\partial P U_{x_1,\Lambda}}{\partial \Lambda} P U_{x_i,\Lambda} + O(\varepsilon^{(1+\sigma)/(N-2)}) \right).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
& \int_{\Omega_1} K(\varepsilon^{1/(N-2)}|y|) P U_{x_1, \Lambda}^p \frac{\partial P U_{x_1, \Lambda}}{\partial \Lambda} \\
&= \int_{\Omega_1} P U_{x_1, \Lambda}^p \frac{\partial P U_{x_1, \Lambda}}{\partial \Lambda} + \int_{\Omega_1} (K(\varepsilon^{1/(N-2)}|y|) - 1) P U_{x_1, \Lambda}^p \frac{\partial P U_{x_1, \Lambda}}{\partial \Lambda} \\
&= \int_{\Omega_1} P U_{x_1, \Lambda}^p \frac{\partial P U_{x_1, \Lambda}}{\partial \Lambda} \\
&\quad + \int_{\Omega_1} (K(|\bar{x}_1|) - 1) P U_{x_1, \Lambda}^p \frac{\partial P U_{x_1, \Lambda}}{\partial \Lambda} + O(\varepsilon^{(1+\sigma)/(N-2)}) \\
&= \int_{\Omega_1} P U_{x_1, \Lambda}^p \frac{\partial P U_{x_1, \Lambda}}{\partial \Lambda} - K'(1)d \int_{\Omega_1} P U_{x_1, \Lambda}^p \frac{\partial P U_{x_1, \Lambda}}{\partial \Lambda} + O(\varepsilon^{(1+\sigma)/(N-2)}) \\
&= - \int_{\Omega_1} U_{x_1, \Lambda}^p \frac{\partial \phi_{x_1, \Lambda}}{\partial \Lambda} - p \int_{\Omega_1} U_{x_1, \Lambda}^p \frac{\partial U_{x_1, \Lambda}}{\partial \Lambda} \phi_{x_1, \Lambda} + O(\varepsilon^{(1+\sigma)/(N-2)}) \\
&= \frac{B_1 \varepsilon H(\bar{x}_1, \bar{x}_1)}{\Lambda^{N-1}} + O(\varepsilon^{(1+\sigma)/(N-2)})
\end{aligned}$$

and

$$\begin{aligned}
& \int_{\Omega_1} K(\varepsilon^{1/(N-2)}|y|) P U_{x_1, \Lambda}^p \frac{\partial P U_{x_i, \Lambda}}{\partial \Lambda} \\
&= \int_{\Omega_1} P U_{x_1, \Lambda}^p \frac{\partial P U_{x_i, \Lambda}}{\partial \Lambda} + \int_{\Omega_1} (K(\varepsilon^{1/(N-2)}|y|) - 1) P U_{x_1, \Lambda}^p \frac{\partial P U_{x_i, \Lambda}}{\partial \Lambda} \\
&= \int_{\Omega_1} U_{x_1, \Lambda}^p \frac{\partial U_{x_i, \Lambda}}{\partial \Lambda} - \int_{\Omega_1} U_{x_1, \Lambda}^p \frac{\partial \phi_{x_i, \Lambda}}{\partial \Lambda} + O(\varepsilon^{(1+\sigma)/(N-2)}) \\
&= - \frac{B_1 \varepsilon G(\bar{x}_i, \bar{x}_1)}{\Lambda^{N-1}} + O(\varepsilon^{(1+\sigma)/(N-2)}).
\end{aligned}$$

Other terms can be estimated similarly. Thus, the result follows. \square

Appendix B: Basic estimates

In this section, we will give some basic estimates used in the reduction procedure. We will use the same constant $C > 0$ to denote the different constants.

Lemma B.1. *Let $g_{ij} = 1/((1 + |y - x_i|)^\alpha (1 + |y - x_j|)^\beta)$ for each fixed i and j , $i \neq j$, where $\alpha \geq 1$ and $\beta \geq 1$ are two constants. Then for any $0 < \sigma \leq \min(\alpha, \beta)$, there is a constant $C > 0$ such that*

$$g_{ij}(y) \leq \frac{C}{|x_i - x_j|^\sigma} \left(\frac{1}{(1 + |y - x_i|)^{\alpha+\beta-\sigma}} + \frac{1}{(1 + |y - x_j|)^{\alpha+\beta-\sigma}} \right).$$

Lemma B.2. *For any constant $0 < \sigma < N - 2$, there is a constant $C > 0$ such that*

$$\int_{\mathbb{R}^N} \frac{1}{|y - z|^{N-2}} \frac{1}{(1 + |z|)^{2+\sigma}} dz \leq \frac{C}{(1 + |y|)^\sigma}.$$

The proofs of the above two lemmas can be found in [Wei and Yan 2010b].

Lemma B.3. *Suppose that $\varepsilon > 0$ and $N \geq 4$. Then there is a small $\vartheta > 0$ such that*

$$\begin{aligned} \int_{\mathbb{R}^N} \frac{1}{|y - z|^{N-2}} W_{r,\Lambda}^{p-1+\varepsilon}(z) \sum_{j=1}^k \frac{1}{(1 + |z - x_j|)^{\frac{1}{2}(N-2)+\tau}} dz \\ \leq C \sum_{j=1}^k \frac{1}{(1 + |z - x_j|)^{\frac{1}{2}(N-2)+\tau+\vartheta}}. \end{aligned}$$

Proof. This is similar to the proof of Lemma B.3 in [Wei and Yan 2010b]. So we just sketch it. Note that

$$W_{r,\Lambda}(z) \leq C \sum_{j=1}^k \frac{1}{(1 + |z - x_j|)^{N-2}}.$$

As in [Wei and Yan 2010b], for $y \in \Omega_1$ we have $W_{r,\Lambda}(z) \leq \frac{C}{(1 + |z - x_1|)^{N-2-\tau_1}}$, where $0 < \tau_1 \leq \frac{1}{2}(N - 2)$. Thus,

$$W_{r,\Lambda}^{p-1+\varepsilon}(z) \leq \frac{C}{(1 + |z - x_1|)^{4 - \frac{4\tau_1}{N-2} + (N-2-\tau_1)\varepsilon}}.$$

By virtue of Lemma B.1, for $y \in \Omega_1$ we get

$$\begin{aligned} W_{r,\Lambda}^{p-1+\varepsilon}(z) \sum_{j=1}^k \frac{1}{(1 + |z - x_j|)^{\frac{1}{2}(N-2)+\tau}} \\ \leq \frac{C}{(1 + |z - x_1|)^{\frac{1}{2}(N+6)+\tau - \frac{4\tau_1}{N-2} + (N-2-\tau_1)\varepsilon}} \\ + \sum_{j=2}^k \frac{C}{(1 + |z - x_1|)^{4 - \frac{4\tau_1}{N-2} + (N-2-\tau_1)\varepsilon}} \frac{1}{(1 + |z - x_j|)^{\frac{1}{2}(N-2)+\tau}} \\ \leq \frac{C}{(1 + |z - x_1|)^{\frac{1}{2}(N+6)+\tau - \frac{4\tau_1}{N-2} + (N-2-\tau_1)\varepsilon}} \\ + \frac{C}{(1 + |z - x_1|)^{\frac{1}{2}(N+6)+\tau - \frac{N+2}{N-2}\tau_1 + (N-2-\tau_1)\varepsilon}} \sum_{j=2}^k \frac{1}{|x_j - x_1|^{\tau_1}} \\ \leq \frac{C}{(1 + |z - x_1|)^{\frac{1}{2}(N+6)+\tau - \frac{N+2}{N-2}\tau_1 + (N-2-\tau_1)\varepsilon}}. \end{aligned}$$

Thus, we can obtain

$$\begin{aligned}
 & \int_{\Omega_1} \frac{1}{|y-z|^{N-2}} W_{r,\Lambda}^{p-1+\varepsilon}(z) \sum_{j=1}^k \frac{1}{(1+|z-x_j|)^{\frac{1}{2}(N-2)+\tau}} dz \\
 & \leq \int_{\Omega_1} \frac{1}{|y-z|^{N-2}} \frac{C}{(1+|z-x_1|)^{\frac{1}{2}(N+6)+\tau-\frac{N+2}{N-2}\tau_1+(N-2-\tau_1)\varepsilon}} dz \\
 & \leq \frac{C}{(1+|z-x_1|)^{\frac{1}{2}(N+2)+\tau-\frac{N+2}{N-2}\tau_1+(N-2-\tau_1)\varepsilon}}.
 \end{aligned}$$

As a result, for τ_1 satisfying $2 - (N+2)/(N-2)\tau_1 > 0$, we find that

$$\begin{aligned}
 & \int_{\mathbb{R}^N} \frac{1}{|y-z|^{N-2}} W_{r,\Lambda}^{p-1+\varepsilon}(z) \sum_{j=1}^k \frac{1}{(1+|z-x_j|)^{\frac{1}{2}(N-2)+\tau}} dz \\
 & = \sum_{i=1}^k \int_{\Omega_i} \frac{1}{|y-z|^{N-2}} W_{r,\Lambda}^{p-1+\varepsilon}(z) \sum_{j=1}^k \frac{1}{(1+|z-x_j|)^{\frac{1}{2}(N-2)+\tau}} dz \\
 & \leq \sum_{i=1}^k \frac{C}{(1+|z-x_1|)^{\frac{1}{2}(N+2)+\tau-\frac{N+2}{N-2}\tau_1+(N-2-\tau_1)\varepsilon}} \\
 & \leq C \sum_{j=1}^k \frac{1}{(1+|z-x_j|)^{\frac{1}{2}(N-2)+\tau+\vartheta}}.
 \end{aligned}$$

□

Acknowledgements

The authors would like to thank the referee very much for the careful reading of the paper and valuable comments, including pointing out [Remark 1.5](#). Z. Liu was supported by funds from the NSFC (No. 11426088, No. 11501166); S. Peng was partially supported by funds from the NSFC (No. 11125101, No. 11571130).

References

- [Ben Ayed et al. 2003] M. Ben Ayed, K. El Mehdi, O. Rey, and M. Grossi, “[A nonexistence result of single peaked solutions to a supercritical nonlinear problem](#)”, *Commun. Contemp. Math.* **5**:2 (2003), 179–195. [MR 2004k:35140](#) [Zbl 1066.35035](#)
- [Byeon and Wang 2005] J. Byeon and Z.-Q. Wang, “[On the Hénon equation: asymptotic profile of ground states, II](#)”, *J. Differential Equations* **216**:1 (2005), 78–108. [MR 2006j:35068](#) [Zbl 1114.35070](#)
- [Byeon and Wang 2006] J. Byeon and Z.-Q. Wang, “[On the Hénon equation: asymptotic profile of ground states, I](#)”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **23**:6 (2006), 803–828. [MR 2007g:35046](#) [Zbl 1114.35071](#)
- [Cao and Peng 2003] D. Cao and S. Peng, “[The asymptotic behaviour of the ground state solutions for Hénon equation](#)”, *J. Math. Anal. Appl.* **278**:1 (2003), 1–17. [MR 2003m:35062](#) [Zbl 1086.35036](#)

- [Cao et al. 2009] D. Cao, S. Peng, and S. Yan, “Asymptotic behaviour of ground state solutions for the Hénon equation”, *IMA J. Appl. Math.* **74**:3 (2009), 468–480. MR 2010d:35088 Zbl 1169.35377
- [Gidas et al. 1979] B. Gidas, W. M. Ni, and L. Nirenberg, “Symmetry and related properties via the maximum principle”, *Comm. Math. Phys.* **68**:3 (1979), 209–243. MR 80h:35043 Zbl 0425.35020
- [Gladiali and Grossi 2012] F. Gladiali and M. Grossi, “Supercritical elliptic problem with non-autonomous nonlinearities”, *J. Differential Equations* **253**:9 (2012), 2616–2645. MR 2959382 Zbl 1266.35105
- [Gladiali et al. 2013] F. Gladiali, M. Grossi, and S. L. N. Neves, “Nonradial solutions for the Hénon equation in \mathbb{R}^N ”, *Adv. Math.* **249** (2013), 1–36. MR 3116566 Zbl 06296090
- [Hénon 1973] M. Hénon, “Numerical experiments on the stability of spherical stellar systems”, *Astron. Astrophys.* **24** (1973), 229–238.
- [Hirano 2009] N. Hirano, “Existence of positive solutions for the Hénon equation involving critical Sobolev terms”, *J. Differential Equations* **247**:5 (2009), 1311–1333. MR 2010f:35121 Zbl 1176.35083
- [Li and Peng 2009] S. Li and S. Peng, “Asymptotic behavior on the Hénon equation with supercritical exponent”, *Sci. China Ser. A* **52**:10 (2009), 2185–2194. MR 2011b:35159 Zbl 1184.35136
- [Ni 1982] W. M. Ni, “A nonlinear Dirichlet problem on the unit ball and its applications”, *Indiana Univ. Math. J.* **31**:6 (1982), 801–807. MR 84b:35051 Zbl 0515.35033
- [Peng 2006] S. Peng, “Multiple boundary concentrating solutions to Dirichlet problem of Hénon equation”, *Acta Math. Appl. Sin. Engl. Ser.* **22**:1 (2006), 137–162. MR 2006k:35099 Zbl 1153.35325
- [Peng and Wang 2013] S. Peng and Z.-Q. Wang, “Segregated and synchronized vector solutions for nonlinear Schrödinger systems”, *Arch. Ration. Mech. Anal.* **208**:1 (2013), 305–339. MR 3021550 Zbl 1260.35211
- [del Pino et al. 2003] M. del Pino, P. Felmer, and M. Musso, “Two-bubble solutions in the supercritical Bahri–Coron’s problem”, *Calc. Var. Partial Differential Equations* **16**:2 (2003), 113–145. MR 2004a:35079 Zbl 1142.35421
- [Pistoia and Serra 2007] A. Pistoia and E. Serra, “Multi-peak solutions for the Hénon equation with slightly subcritical growth”, *Math. Z.* **256**:1 (2007), 75–97. MR 2008b:35091 Zbl 1134.35047
- [Pohožaev 1965] S. I. Pohožaev, “On the eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$ ”, *Dokl. Akad. Nauk SSSR* **165** (1965), 36–39. In Russian; translated in *Sov. Math. Dokl.* **6** (1965), 1408–1411. MR 33 #411 Zbl 0141.30202
- [Serra 2005] E. Serra, “Non radial positive solutions for the Hénon equation with critical growth”, *Calc. Var. Partial Differential Equations* **23**:3 (2005), 301–326. MR 2006f:35100 Zbl 1207.35147
- [Smets et al. 2002] D. Smets, M. Willem, and J. Su, “Non-radial ground states for the Hénon equation”, *Commun. Contemp. Math.* **4**:3 (2002), 467–480. MR 2003g:35086 Zbl 1160.35415
- [Wei and Yan 2010a] J. Wei and S. Yan, “Infinitely many positive solutions for the nonlinear Schrödinger equations in \mathbb{R}^N ”, *Calc. Var. Partial Differential Equations* **37**:3–4 (2010), 423–439. MR 2011c:35065 Zbl 1189.35106
- [Wei and Yan 2010b] J. Wei and S. Yan, “Infinitely many solutions for the prescribed scalar curvature problem on \mathbb{S}^N ”, *J. Funct. Anal.* **258**:9 (2010), 3048–3081. MR 2011g:35119 Zbl 1209.53028
- [Wei and Yan 2011] J. Wei and S. Yan, “Infinitely many positive solutions for an elliptic problem with critical or supercritical growth”, *J. Math. Pures Appl.* (9) **96**:4 (2011), 307–333. MR 2832637 Zbl 1253.31008
- [Wei and Yan 2013] J. Wei and S. Yan, “Infinitely many nonradial solutions for the Hénon equation with critical growth”, *Rev. Mat. Iberoam.* **29**:3 (2013), 997–1020. MR 3090144 Zbl 1277.35330

Received November 6, 2014. Revised May 10, 2015.

ZHONGYUAN LIU
SCHOOL OF MATHEMATICS AND STATISTICS
HENAN UNIVERSITY
KAIFENG, HENAN 475004
CHINA

liuzy@henu.edu.cn

SHUANGJIE PENG
SCHOOL OF MATHEMATICS AND STATISTICS
HUBEI KEY LABORATORY OF MATHEMATICAL PHYSICS
CENTRAL CHINA NORMAL UNIVERSITY
WUHAN, HUBEI 430079
CHINA

sjpeng@mail.ccnu.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2016 is US \$440/year for the electronic version, and \$600/year for print and electronic.

Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), PASCAL CNRS Index, [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY



mathematical sciences publishers

nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 280 No. 1 January 2016

Stable capillary hypersurfaces in a wedge	1
JAIGYOUNG CHOE and MIYUKI KOISO	
The Chern–Simons invariants for the double of a compression body	17
DAVID L. DUNCAN	
Compactness and the Palais–Smale property for critical Kirchhoff equations in closed manifolds	41
EMMANUEL HEBEY	
On the equivalence of the definitions of volume of representations	51
SUNGWOON KIM	
Strongly positive representations of even GSpin groups	69
YEANSU KIM	
An Orlik–Raymond type classification of simply connected 6-dimensional torus manifolds with vanishing odd-degree cohomology	89
SHINTARÔ KUROKI	
Solutions with large number of peaks for the supercritical Hénon equation	115
ZHONGYUAN LIU and SHUANGJIE PENG	
Effective divisors on the projective line having small diagonals and small heights and their application to adelic dynamics	141
YÛSUKE OKUYAMA	
Computing higher Frobenius–Schur indicators in fusion categories constructed from inclusions of finite groups	177
PETER SCHAUENBURG	
Chordal generators and the hydrodynamic normalization for the unit ball	203
SEBASTIAN SCHLEISSINGER	
On a question of A. Balog	227
ILYA D. SHKREDOV	
Uniqueness result on nonnegative solutions of a large class of differential inequalities on Riemannian manifolds	241
YUHUA SUN	
Correction to “Closed orbits of a charge in a weakly exact magnetic field”	255
WILL J. MERRY	