

*Pacific  
Journal of  
Mathematics*

**COMPLEX HYPERBOLIC  $(3, 3, n)$  TRIANGLE GROUPS**

JOHN R. PARKER, JIEYAN WANG AND BAOHUA XIE

# COMPLEX HYPERBOLIC $(3, 3, n)$ TRIANGLE GROUPS

JOHN R. PARKER, JIEYAN WANG AND BAOHUA XIE

Let  $p, q, r$  be positive integers. Complex hyperbolic  $(p, q, r)$  triangle groups are representations of the hyperbolic  $(p, q, r)$  reflection triangle group to the holomorphic isometry group of complex hyperbolic space  $H_{\mathbb{C}}^2$ , where the generators fix complex lines. In this paper, we obtain all the discrete and faithful complex hyperbolic  $(3, 3, n)$  triangle groups for  $n \geq 4$ . Our result solves a conjecture of Schwartz in the case when  $p = q = 3$ .

## 1. Introduction

An abstract  $(p, q, r)$  reflection triangle group for positive integers  $p, q, r$  is the group

$$\Delta_{p,q,r} = \langle \sigma_1, \sigma_2, \sigma_3 \mid \sigma_1^2 = \sigma_2^2 = \sigma_3^2 = (\sigma_2\sigma_3)^p = (\sigma_3\sigma_1)^q = (\sigma_1\sigma_2)^r = \text{id} \rangle.$$

We sometimes take (at least) one of  $p, q, r$  to be  $\infty$ , in which case the corresponding relation does not appear.

It is interesting to seek geometrical representations of  $\Delta_{p,q,r}$ . An extremely well-known fact is that  $\Delta_{p,q,r}$  may be realised geometrically as the reflections in the side of a geodesic triangle with internal angles  $\pi/p, \pi/q, \pi/r$ . Furthermore, if  $1/p + 1/q + 1/r > 1, = 1$  or  $< 1$  then this triangle is spherical, Euclidean or hyperbolic respectively. Moreover, up to isometries (or similarities in the Euclidean case) there is a unique such triangle and the representation is rigid. In the case where (at least) one of  $p, q, r$  is  $\infty$ , we omit the relevant term from  $1/p + 1/q + 1/r$  and we insist that the sides of the triangle are asymptotic. Thus the  $(\infty, \infty, \infty)$  triangle is a triangle in the hyperbolic plane with all three vertices on the boundary.

In contrast, if we choose a geometrical representation of  $\Delta_{p,q,r}$  in a space of nonconstant curvature then more interesting things can happen; see, for example, [Brehm 1990]. In this paper, we consider representations of  $\Delta_{p,q,r}$  to  $\text{SU}(2, 1)$ , which is (a triple cover of) the group of holomorphic isometries of complex hyperbolic space  $H_{\mathbb{C}}^2$ . A convenient model of  $H_{\mathbb{C}}^2$  is the unit ball in  $\mathbb{C}^2$  with the

---

Wang and Xie were supported by NSFC (grant numbers 11071059 and 11371126). Xie was also supported by NSFC (grant number 11201134) and the Young Teachers Support program of Hunan University.

MSC2010: primary 20H10; secondary 22E40, 51M10.

Keywords: complex hyperbolic geometry, complex hyperbolic triangle groups.

Bergman metric, having constant holomorphic sectional curvature and  $1/4$ -pinched real sectional curvatures.

A *complex hyperbolic triangle group* will be a representation of  $\Delta_{p,q,r}$  to  $SU(2, 1)$  where the generators fix complex lines. Note we could have made other choices. For example, we could choose the generators to be antiholomorphic isometries, or we could choose reflections in three complex lines but with higher order. These choices lead to interesting results, but we will not consider them here. A crucial observation is that when  $\min\{p, q, r\} \geq 3$ , there is a one (real) dimensional representation space of complex hyperbolic triangle groups with  $1/p + 1/q + 1/r < 1$  (either make a simple dimension count or see [Brehm 1990] for example). This means that the representation is determined up to conjugacy by  $p, q, r$  and one extra variable. This variable is determined by certain traces; see, for example, [Pratoussevitch 2005].

In order to state our main results, we need a little terminology. Elements of  $SU(2, 1)$  act on complex hyperbolic space  $\mathbf{H}_{\mathbb{C}}^2$  and its boundary (see below). An element  $A \in SU(2, 1)$  is called *loxodromic* if it fixes two points, both of which lie on  $\partial\mathbf{H}_{\mathbb{C}}^2$ ; *parabolic* if it fixes exactly one point, and this point lies on  $\partial\mathbf{H}_{\mathbb{C}}^2$ ; *elliptic* if it fixes at least one point of  $\mathbf{H}_{\mathbb{C}}^2$ . Discrete groups cannot contain elliptic elements of infinite order. Therefore in a representation of an abstract group to  $SU(2, 1)$ , if an element of infinite order in the abstract group is represented by an elliptic map then the representation is not discrete or not faithful (or both); compare with [Goldman and Parker 1992].

Complex hyperbolic triangle groups have a rich history; see Schwartz's ICM survey [2002] for an overview. In particular, he presented the following conjectural picture:

**Conjecture 1.1** [Schwartz 2002]. *Let  $\Delta_{p,q,r}$  be a triangle group with  $p \leq q \leq r$ . Then any complex hyperbolic representation  $\Gamma$  of  $\Delta_{p,q,r}$  is discrete and faithful if and only if  $W_A = I_1 I_3 I_2 I_3$  and  $W_B = I_1 I_2 I_3$  are not elliptic. Furthermore:*

- (i) *If  $p < 10$  then  $\Gamma$  is discrete and faithful if and only if  $W_A = I_1 I_3 I_2 I_3$  is nonelliptic.*
- (ii) *If  $p > 13$  then  $\Gamma$  is discrete and faithful if and only if  $W_B = I_1 I_2 I_3$  is nonelliptic.*

The initial step towards solving this conjecture is the following result of Grossi.

**Proposition 1.2** [Grossi 2007]. *Let  $\Delta_{p,q,r}$  be a triangle group with  $p \leq q \leq r$ . Define  $W_A = I_1 I_3 I_2 I_3$  and  $W_B = I_1 I_2 I_3$ . Then for complex hyperbolic representations of  $\Delta_{p,q,r}$ :*

- (i) *If  $p < 10$  and  $W_A = I_1 I_3 I_2 I_3$  is nonelliptic then  $W_B$  is nonelliptic.*
- (ii) *If  $p > 13$  and  $W_B = I_1 I_2 I_3$  is nonelliptic then  $W_A$  is nonelliptic.*

A motivating example, initially considered by Goldman and Parker [1992] and completed by Schwartz [2001b; 2005], concerns complex hyperbolic ideal triangle groups, that is, representations of  $\Delta_{\infty, \infty, \infty}$ . This result may be summarised as follows:

**Theorem 1.3** [Goldman and Parker 1992; Schwartz 2001b; 2005]. *Let  $\Gamma = \langle I_1, I_2, I_3 \rangle$  be a complex hyperbolic  $(\infty, \infty, \infty)$  triangle group. Then  $\Gamma$  is a discrete and faithful representation of  $\Delta_{\infty, \infty, \infty}$  if and only if  $I_1 I_2 I_3$  is nonelliptic.*

Note that this gives a complete solution to Schwartz’s conjecture in the case  $p = q = r = \infty$ . Furthermore, Schwartz [2001a] gives an elegant description of the group where  $I_1 I_2 I_3$  is parabolic.

**Theorem 1.4** [Schwartz 2001a]. *Let  $\Gamma = \langle I_1, I_2, I_3 \rangle$  be the  $(\infty, \infty, \infty)$  complex hyperbolic triangle group for which  $I_1 I_2 I_3$  is parabolic. Let  $\Gamma_2 = \langle I_1 I_2, I_1 I_3 \rangle$  be the index-2 subgroup of  $\Gamma$  with no complex reflections. Then  $\mathbf{H}_{\mathbb{C}}^2 / \Gamma_2$  is a complex hyperbolic orbifold whose boundary is a triple cover of the Whitehead link complement.*

Schwartz [2007] proves his conjecture for  $\min\{p, q, r\}$  sufficiently large (but unfortunately with no effective bound on this minimum).

**Theorem 1.5** [Schwartz 2007]. *Let  $\Gamma = \langle I_1, I_2, I_3 \rangle$  be a complex hyperbolic  $(p, q, r)$  triangle group with  $p \leq q \leq r$ . If  $p$  is sufficiently large, then  $\Gamma$  is a discrete and faithful representation of  $\Delta_{p, q, r}$  if and only if  $I_1 I_2 I_3$  is nonelliptic.*

Our main result solves Schwartz’s conjecture in the case when  $p = q = 3$ .

**Theorem 1.6.** *Let  $n$  be an integer at least 4. Let  $\Gamma = \langle I_1, I_2, I_3 \rangle$  be a complex hyperbolic  $(3, 3, n)$  triangle group. Then  $\Gamma$  is a discrete and faithful representation of  $\Delta_{3, 3, n}$  if and only if  $I_1 I_3 I_2 I_3$  is nonelliptic.*

Note that the “only if” part is a consequence of our observation about elliptic elements above. The “if” part will follow from Corollary 4.4 below.

For the representation where  $I_1 I_3 I_2 I_3$  is parabolic, when  $n = 4$  and 5 we have the following description of the quotient orbifold from the census of Falbel, Koseleff and Rouillier [Falbel et al. 2015]. The case  $n = 4$  combines work of Deraux, Falbel and Wang [Deraux and Falbel 2015; Falbel and Wang 2014]. The cleanest statement may be found in [Deraux 2015, Theorem 4.2], which also treats the case  $n = 5$ .

**Theorem 1.7** [Deraux 2015, Theorem 4.2]. (i) *Let  $\Gamma = \langle I_1, I_2, I_3 \rangle$  be the complex hyperbolic  $(3, 3, 4)$  triangle group for which  $I_1 I_3 I_2 I_3$  is parabolic. Let  $\Gamma_2 = \langle I_1 I_2, I_1 I_3 \rangle$  be the index-2 subgroup of  $\Gamma$  with no complex reflections. Then  $\Gamma_2$  is conjugate to both  $\rho_{1-1}(\pi_1(M_4))$  and  $\rho_{4-1}(\pi_1(M_4))$  from [Falbel et al. 2015]. In particular,  $\mathbf{H}_{\mathbb{C}}^2 / \Gamma_2$  is a complex hyperbolic orbifold whose boundary is the figure eight knot complement.*

(ii) Let  $\Gamma = \langle I_1, I_2, I_3 \rangle$  be the complex hyperbolic  $(3, 3, 5)$  triangle group for which  $I_1 I_3 I_2 I_3$  is parabolic. Let  $\Gamma_2 = \langle I_1 I_2, I_1 I_3 \rangle$  be the index-2 subgroup of  $\Gamma$  with no complex reflections. Then  $\Gamma_2$  is conjugate to both  $\rho_{4-3}(\pi_1(M_9))$  and  $\rho_{3-3}(\pi_1(M_{15}))$  from [Falbel et al. 2015].

It should be possible to give a similar description of the other complex hyperbolic  $(3, 3, n)$  triangle groups for which  $I_1 I_3 I_2 I_3$  is parabolic.

Note that Theorem 1.6 holds in the case  $n = \infty$ . This follows from recent work of Parker and Will [2015b] (see also [Parker and Will 2015a]). Furthermore, if as above  $\Gamma_2 = \langle I_1 I_2, I_1 I_3 \rangle$  is the index-2 subgroup of representation of the  $(3, 3, \infty)$  triangle group for which  $I_1 I_3 I_2 I_3$  is parabolic, then  $\mathbb{H}_{\mathbb{C}}^2 / \Gamma_2$  is a complex hyperbolic orbifold whose boundary is the Whitehead link complement. This is one of the representations in [Falbel et al. 2015].

Finally, we note some further interesting groups in this family.

**Theorem 1.8 [Thompson 2010].** *The complex hyperbolic  $(3, 3, 4)$  triangle group with  $I_1 I_3 I_2 I_3$  of order 7 and the complex hyperbolic  $(3, 3, 5)$  triangle group with  $I_1 I_3 I_2 I_3$  of order 5 are both lattices.*

Our method of proof will be to construct a Dirichlet domain based at the fixed point of the order- $n$  elliptic map  $I_1 I_2$ . Since this point has nontrivial stabiliser, this domain is not a fundamental domain for  $\Gamma$ , but it is a fundamental domain for the coset space of the stabiliser of this point in  $\Gamma$ . Of course, in order to prove directly that this is a Dirichlet domain, we would have to check infinitely many inequalities. Instead, we construct a candidate Dirichlet domain and then use the Poincaré polyhedron theorem for coset decompositions (see [Mostow 1980, Theorem 6.3.2] or [Deraux et al. 2015, Theorem 3.2], for example).

In the case of a Fuchsian  $(3, 3, n)$  triangle group acting on the hyperbolic plane, a fundamental domain is a hyperbolic triangle with internal angles  $\pi/3, \pi/3$  and  $\pi/n$ . The Dirichlet domain with centre the fixed point of an order- $n$  elliptic map is a regular hyperbolic  $2n$ -gon with internal angles  $2\pi/3$ . This  $2n$ -gon is made up of  $2n$  copies of the triangular fundamental domain for the  $(3, 3, n)$  group; see Figure 1. The stabiliser of the order- $n$  fixed point, which is a dihedral group of order  $2n$ , fixes the  $2n$ -gon and permutes the triangles.

For the complex hyperbolic  $(3, 3, n)$  triangle groups, we will see that the combinatorial structure of the Dirichlet domain  $D$  is the same as that in the Fuchsian case. Namely,  $D$  has  $2n$  sides, each of which is contained in a bisector. Each side meets exactly two other sides (in the case where  $I_1 I_3 I_2 I_3$  is parabolic, there are some additional tangencies between sides on the ideal boundary). The sides are permuted by the dihedral group  $\langle I_1, I_2 \rangle$ .

In Section 2 we give the necessary background on complex hyperbolic geometry and the Poincaré polyhedron theorem. In Section 3 we normalise the generators

of  $\Gamma$  and discuss the parameters this involves. Finally, in [Section 4](#) we consider the bisectors and their intersection properties. This is the heart of the paper.

## 2. Background

**Complex hyperbolic space.** Let  $\mathbb{C}^{2,1}$  be the three-dimensional complex vector space equipped with a Hermitian form  $H$  of signature (2, 1). In this paper we consider the diagonal Hermitian form  $H = \text{diag}(1, 1, -1)$ . Thus if  $\mathbf{u} = (u_1, u_2, u_3)^t$  and  $\mathbf{v} = (v_1, v_2, v_3)^t$  then the Hermitian form is given by

$$\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{v}^* H \mathbf{u} = u_1 \bar{v}_1 + u_2 \bar{v}_2 - u_3 \bar{v}_3.$$

Define

$$V_- = \{\mathbf{v} \in \mathbb{C}^{2,1} : \langle \mathbf{v}, \mathbf{v} \rangle < 0\}, \quad V_0 = \{\mathbf{v} \in \mathbb{C}^{2,1} - \{0\} : \langle \mathbf{v}, \mathbf{v} \rangle = 0\}.$$

There is a natural projection map  $\mathbb{P}$  from  $\mathbb{C}^{2,1} - \{0\}$  to  $\mathbb{C}\mathbb{P}^2$  that identifies all nonzero (complex) scalar multiples of a vector in  $\mathbb{C}^{2,1}$ . *Complex hyperbolic space* is defined to be  $\mathbf{H}_{\mathbb{C}}^2 = \mathbb{P}V_-$  and its boundary is  $\partial\mathbf{H}_{\mathbb{C}}^2 = \mathbb{P}V_0$ . Clearly, if  $\mathbf{v}$  lies in  $V_-$  or  $V_0$  then  $v_3 \neq 0$  and so  $\mathbf{H}_{\mathbb{C}}^2 \cup \partial\mathbf{H}_{\mathbb{C}}^2$  is contained in the affine chart of  $\mathbb{C}\mathbb{P}^2$  with  $v_3 \neq 0$ . We canonically identify this chart with  $\mathbb{C}^2$  by setting  $z = v_1/v_3$  and  $w = v_2/v_3$ . Thus a vector  $(z, w) \in \mathbb{C}^2$  corresponds to  $[z : w : 1]^t$  in  $\mathbb{C}\mathbb{P}^2$ . Evaluating the Hermitian form at this point gives  $|z|^2 + |w|^2 - 1 = (|v_1|^2 + |v_2|^2 - |v_3|^2)/|v_3|^2$ . Therefore

$$\mathbf{H}_{\mathbb{C}}^2 = \{(z, w) \in \mathbb{C}^2 : |z|^2 + |w|^2 < 1\}, \quad \partial\mathbf{H}_{\mathbb{C}}^2 = \{(z, w) \in \mathbb{C}^2 : |z|^2 + |w|^2 = 1\}.$$

In other words,  $\mathbf{H}_{\mathbb{C}}^2$  is the unit ball in  $\mathbb{C}^2$  and its boundary is the unit sphere  $S^3$ .

The Bergman metric on  $\mathbf{H}_{\mathbb{C}}^2$  is given in terms of the Hermitian form. Let  $u$  and  $v$  be points in  $\mathbf{H}_{\mathbb{C}}^2$  and let  $\mathbf{u}$  and  $\mathbf{v}$  be vectors in  $V_-$  so that  $\mathbb{P}\mathbf{u} = u$  and  $\mathbb{P}\mathbf{v} = v$ . The Bergman metric is given as a Riemannian metric  $ds^2$  or a distance function  $\rho(u, v)$  by the formulae

$$ds^2 = \frac{-4}{\langle \mathbf{u}, \mathbf{u} \rangle^2} \det \begin{pmatrix} \langle \mathbf{u}, \mathbf{u} \rangle & \langle d\mathbf{u}, \mathbf{u} \rangle \\ \langle \mathbf{u}, d\mathbf{u} \rangle & \langle d\mathbf{u}, d\mathbf{u} \rangle \end{pmatrix}, \quad \cosh^2 \left( \frac{\rho(u, v)}{2} \right) = \frac{\langle \mathbf{u}, \mathbf{v} \rangle \langle \mathbf{v}, \mathbf{u} \rangle}{\langle \mathbf{u}, \mathbf{u} \rangle \langle \mathbf{v}, \mathbf{v} \rangle}.$$

The formulae for the Bergman metric are homogeneous and so the ambiguity in the choice of  $\mathbf{u}$  and  $\mathbf{v}$  does not matter.

Let  $\text{SU}(2, 1)$  be the group of unimodular matrices preserving the Hermitian form  $H$ . An element  $A$  of  $\text{SU}(2, 1)$  acts on  $\mathbf{H}_{\mathbb{C}}^2$  as  $A(u) = \mathbb{P}(A\mathbf{u})$ , where  $\mathbf{u}$  is any vector in  $V_-$  with  $\mathbb{P}\mathbf{u} = u$ . It is clear that scalar multiples of the identity act trivially. Since the determinant of  $A$  is 1, such a scalar multiple must be a cube root of unity. Therefore, we define  $\text{PU}(2, 1) = \text{SU}(2, 1)/\{\omega I : \omega^3 = 1\}$ . Since the Bergman metric is given in terms of the Hermitian form, it is clear that elements of  $\text{SU}(2, 1)$  or  $\text{PU}(2, 1)$ , act as isometries of  $\mathbf{H}_{\mathbb{C}}^2$ . Indeed,  $\text{PU}(2, 1)$  is the full group of holomorphic isometries of  $\mathbf{H}_{\mathbb{C}}^2$ . In what follows, we choose to work with matrices in  $\text{SU}(2, 1)$ .

There are two kinds of totally geodesic two-dimensional submanifolds in  $\mathbf{H}_{\mathbb{C}}^2$ : complex lines and totally real totally geodesic subspaces. Let  $\mathbf{c} \in \mathbb{C}^{2,1}$  be a vector with  $\langle \mathbf{c}, \mathbf{c} \rangle > 0$ . Then a *complex line* is the projection of the set  $\{z \in \mathbb{C}^{2,1} : \langle z, \mathbf{c} \rangle = 0\}$ . The vector  $\mathbf{c}$  is then called a *polar vector* of the complex line. The *complex reflection* with polar vector  $\mathbf{c}$  is defined to be

$$I_{\mathbf{c}}(z) = -z + \frac{2\langle z, \mathbf{c} \rangle}{\langle \mathbf{c}, \mathbf{c} \rangle} \mathbf{c}.$$

**Bisectors and Dirichlet domains.** We will consider subgroups of  $SU(2, 1)$  acting on  $\mathbf{H}_{\mathbb{C}}^2$  and we want to show they are discrete. We will do this by constructing a fundamental polyhedron and using the Poincaré polyhedron theorem. There are no totally geodesic real hypersurfaces in  $\mathbf{H}_{\mathbb{C}}^2$  and so we must choose hypersurfaces for the sides of our polyhedra. We choose to work with bisectors. A *bisector* in  $\mathbf{H}_{\mathbb{C}}^2$  is the locus of points equidistant (with respect to the Bergman metric) from a given pair of points in  $\mathbf{H}_{\mathbb{C}}^2$ . Suppose that these points are  $u$  and  $v$ . Choose lifts  $\mathbf{u} = (u_1, u_2, u_3)^t$  and  $\mathbf{v} = (v_1, v_2, v_3)^t$  to  $V_-$  so that  $\langle \mathbf{u}, \mathbf{u} \rangle = \langle \mathbf{v}, \mathbf{v} \rangle$ . Then the bisector equidistant from  $u$  and  $v$  is

$$\begin{aligned} \mathcal{B} &= \mathcal{B}(u, v) = \{(z, w) \in \mathbf{H}_{\mathbb{C}}^2 : \rho((z, w), u) = \rho((z, w), v)\} \\ &= \{(z, w) \in \mathbf{H}_{\mathbb{C}}^2 : |z\bar{u}_1 + w\bar{u}_2 - \bar{u}_3| = |z\bar{v}_1 + w\bar{v}_2 - \bar{v}_3|\}. \end{aligned}$$

Suppose that we are given three points  $u, v_1$  and  $v_2$  in  $\mathbf{H}_{\mathbb{C}}^2$ . If the three corresponding vectors  $\mathbf{u}, \mathbf{v}_1$  and  $\mathbf{v}_2$  in  $V_-$  form a basis for  $\mathbb{C}^{2,1}$  then the intersection  $\mathcal{B}(u, v_1) \cap \mathcal{B}(u, v_2)$  is called a Giraud disc. This is a particularly nice type of bisector intersection (see [Deraux et al. 2015, Section 2.5]).

Suppose that  $\Gamma$  is a discrete subgroup of  $PU(2, 1)$ . Let  $u$  be a point of  $\mathbf{H}_{\mathbb{C}}^2$  and write  $\Gamma_u$  for the stabiliser of  $u$  in  $\Gamma$  (that is, the subgroup of  $\Gamma$  comprising all elements fixing  $u$ ). Then the *Dirichlet domain*  $D_u(\Gamma)$  for  $\Gamma$  with centre  $u$  is defined to be

$$D_u(\Gamma) = \{v \in \mathbf{H}_{\mathbb{C}}^2 : \rho(v, u) < \rho(v, A(u)) \text{ for all } A \in \Gamma - \Gamma_u\}.$$

Dirichlet domains for certain cyclic groups are particularly simple.

**Proposition 2.1.** *Let  $A$  be a regular elliptic element of  $PU(2, 1)$  of order 3. Then for any point  $u$  not fixed by  $A$ , the Dirichlet domain  $D_u(\langle A \rangle)$  for the cyclic group  $\langle A \rangle$  with centre  $u$  has exactly two sides.*

*Proof.* Since there are only two nontrivial elements in  $\langle A \rangle$ , neither of which fix  $u$ , the Dirichlet domain  $D_u(\langle A \rangle)$  is

$$D_u(\langle A \rangle) = \{v \in \mathbf{H}_{\mathbb{C}}^2 : \rho(v, u) < \rho(v, A(u)), \rho(v, u) < \rho(v, A^{-1}(u))\}.$$

Its images under  $A$  and  $A^{-1}$  are

$$A(D_u(\langle A \rangle)) = \{v : \rho(v, A(u)) < \rho(v, u), \rho(v, A(u)) < \rho(v, A^{-1}(u))\},$$

$$A^{-1}(D_u(\langle A \rangle)) = \{v : \rho(v, A^{-1}(u)) < \rho(v, u), \rho(v, A^{-1}(u)) < \rho(v, A(u))\}.$$

By considering the minimum of  $\rho(v, u)$ ,  $\rho(v, A(u))$ ,  $\rho(v, A^{-1}(u))$  as  $v$  varies over  $\mathbf{H}_{\mathbb{C}}^2$ , it is clear these three domains are disjoint and their closures cover  $\mathbf{H}_{\mathbb{C}}^2$ .  $\square$

**Proposition 2.2** [Phillips 1992]. *Let  $A \in \text{SU}(2, 1)$  have real trace which is at least 3. Then for any  $u \in \mathbf{H}_{\mathbb{C}}^2$ , the bisectors  $\mathcal{B}(u, A(u))$  and  $\mathcal{B}(u, A^{-1}(u))$  are disjoint. Thus, the Dirichlet domain  $D_u(\langle A \rangle)$  has exactly two sides.*

**The Poincaré polyhedron theorem.** Our goal is to construct the Dirichlet domain for a complex hyperbolic representation  $\Gamma$  of the  $(3, 3, n)$  triangle group with centre the fixed point of an order- $n$  elliptic map. If we use the definition of Dirichlet domain, then we need to check infinitely many inequalities. Thus, we need to use another method. This method is to construct a candidate Dirichlet domain and then use the Poincaré polyhedron theorem.

The main tool we use to show discreteness is the Poincaré polyhedron theorem. The version of this theorem that we use is for polyhedra  $D$  with a finite stabiliser; see [Mostow 1980, Theorem 6.3.2] or [Deraux et al. 2015, Theorem 3.2]. Rather than give a general statement of this theorem, we will state it in the particular case we are interested in, namely Dirichlet polyhedra for reflection groups.

Let  $u$  be a point in  $\mathbf{H}_{\mathbb{C}}^2$  and let  $\Upsilon$  be a finite subgroup of  $\text{PU}(2, 1)$  fixing  $u$ . Let  $A_1, \dots, A_n$  be a finite collection of involutions in  $\text{PU}(2, 1)$  (so  $A_i^2$  is the identity for each  $i$ ). Suppose that no  $A_i$  fixes  $u$ . Suppose that the group  $\Upsilon$  preserves this collection of involutions under conjugation. That is, for each  $A_i$  with  $1 \leq i \leq n$  and each  $P \in \Upsilon$ , we suppose that  $PA_iP^{-1} = A_j$  for some  $1 \leq j \leq n$ . Let  $\mathcal{B}_i = \mathcal{B}(u, A_i(u))$  be the bisector equidistant from  $u$  and  $A_i(u)$ . If  $P \in \Upsilon$  satisfies  $PA_iP^{-1} = A_j$  then  $PA_i(u) = A_j(u)$  (since  $P(u) = u$ ) and so  $P$  maps  $\mathcal{B}_i$  to  $\mathcal{B}_j$ . We define  $D$  to be the component of  $\mathbf{H}_{\mathbb{C}}^2 - \bigcup_{i=1}^n \mathcal{B}_i$  containing  $u$ , and we suppose that there are points from each of the  $\mathcal{B}_i$  on the boundary of  $D$  (that is, the  $\mathcal{B}_i$  are not nested). This construction makes  $D$  open. Note that, by construction,  $\Upsilon$  maps  $D$  to itself.

For each  $1 \leq i \leq n$ , let  $s_i = \mathcal{B}_i \cap \bar{D}$ . We call  $s_i$  a *side* of  $D$ . Such a side can be given a cell structure based on how it intersects other sides. We suppose that the involutions  $A_i$  for  $1 \leq i \leq n$  satisfy the following conditions, and so form a *side pairing* of  $D$ :

- (1) For each  $1 \leq i \leq n$ , the involution  $A_i$  sends  $s_i$  to itself, preserving the cell structure. The relation  $A_i^2 = \text{id}$  is called a *reflection relation*.
- (2) For each  $1 \leq i \leq n$ , we have  $\bar{D} \cap A_i(\bar{D}) = s_i$  and  $D \cap A(D) = \emptyset$ .
- (3) If  $v$  is a point in  $s_i$  and in no other side (that is,  $v$  lies in the relative interior of  $s_i$ ) then there is an open neighbourhood  $U_v$  of  $v$  lying in  $\bar{D} \cup A_i(\bar{D})$ .



Note that, unlike the case of reflection groups in constant curvature,  $A_i$  does not fix  $s_i$  pointwise. Therefore, we could have subdivided  $s_i$  into two sets (each of dimension 3) that are interchanged by  $A_i$ . In practice this would cause unnecessary complication.

Suppose that  $s_i$  and  $s_j$  are two sides with nonempty intersection. Their intersection  $r = s_i \cap s_j$  is called a *ridge* of  $D$ . Since  $A_i$  preserves the cell structure of  $s_i$ , we see that  $A_i(r) = s_i \cap s_k$  is another ridge of  $D$ . Applying  $A_k$  gives another ridge in  $s_k$ . Continuing in this way gives a *ridge cycle*

$$(r_1, s_{i_0}, s_{i_1}) \xrightarrow{A_{i_1}} (r_2, s_{i_1}, s_{i_2}) \xrightarrow{A_{i_2}} (r_3, s_{i_2}, s_{i_3}) \cdots$$

Here  $(r_j, s_{i_{j-1}}, s_{i_j})$  is an ordered triple with  $r_j = s_{i_{j-1}} \cap s_{i_j}$ . Since there are finitely many  $\Upsilon$  orbits of  $r_1$ , eventually we find a ridge  $r_{m+1} = s_{i_m} \cap s_{i_{m+1}}$  so that the corresponding ordered triple satisfies

$$(r_{m+1}, s_{i_m}, s_{i_{m+1}}) \xrightarrow{P} (r_1, s_{i_0}, s_{i_1})$$

for some  $P \in \Upsilon$ . We call  $T_1 = PA_{i_m} \cdots A_{i_1}$  the *cycle transformation* associated to  $r_1$ . It means that the ridge cycle starts at  $(r_1, s_{i_0}, s_{i_1})$  and ends to itself by  $T_1$ . Clearly  $T_1$  maps  $r_1$  to itself. Of course,  $T_1$  may not act as the identity on  $r_1$  and even if it does, it may not act as the identity on  $\mathbf{H}_{\mathbb{C}}^2$ . Nevertheless, we suppose  $T_1$  has finite order  $\ell$ . The relation  $T_1^\ell = \text{id}$  is called a *cycle relation*.

In the example we are interested in, the ridge cycle is

$$(r_1, s_{i_0}, s_{i_1}) \xrightarrow{A_{i_1}} (r_2, s_{i_1}, s_{i_2}) \xrightarrow{P} (r_1, s_{i_0}, s_{i_1})$$

and, in fact,  $s_{i_2} = s_{i_0}$  and so  $r_2 = r_1$ . Moreover,  $P$  is an involution with  $P(r_1) = r_1$  and  $P(s_{i_1}) = s_{i_0}$ . Hence the cycle transformation is  $T_1 = PA_{i_1}$ , which happens to have order 3. Thus, the cycle relation is  $T_1^3 = (PA_{i_1})^3 = \text{id}$ .

We suppose that  $D$  satisfies the *cycle condition* which means that copies of  $D$  tessellate a neighbourhood for each ridge  $r$ . Furthermore, the relevant copies of  $D$  are its preimages under suffix subwords of  $T^\ell$ . The full statement is explained in [Deraux et al. 2015]. For brevity, we state this condition only in the special case we are interested in. Let  $r$  be a ridge and let  $T = PA_i$  be its cycle transformation with cycle relation  $(PA_i)^3 = \text{id}$ . Let  $\mathcal{C} = \{\text{id}, PA_i, (PA_i)^2\}$ . Then the cycle condition states that

- (1) 
$$r = \bigcap_{C \in \mathcal{C}} C^{-1}(\bar{D}).$$
- (2) If  $C_1, C_2 \in \mathcal{C}$  with  $C_1 \neq C_2$  then  $C_1^{-1}(D) \cap C_2^{-1}(D) = \emptyset$ .
- (3) If  $v$  is a point in  $r$  and in no other ridge (that is,  $v$  lies in the relative interior of  $r$ ) then there is an open neighbourhood  $U_v$  of  $v$  with

$$U_v \subset \bigcup_{C \in \mathcal{C}} C^{-1}(\bar{D}).$$

It means that there are exactly three copies of  $D$  along each ridge  $r$ , which are  $D$ ,  $T(D)$  and  $T^2(D)$ . Observe that the stabiliser of  $r$  is generated by  $A_i$  and  $P$ . Hence it is a dihedral group of order 6. Since  $A_i$ ,  $P$  and  $PA_iP^{-1}$  preserve one of the three copies and interchange the other two, the stabiliser preserves the three copies of  $D$ .

Finally, if two sides of  $D$  are asymptotic at a point  $v$  of  $\partial\mathbf{H}_{\mathbb{C}}^2$  then there is a horoball  $H_v$  so that  $H_v$  intersects  $\bar{D}$  only in facets of  $D$  containing  $v$  and  $H_v$  is preserved by the stabiliser of  $v$  in  $\Gamma$ . We say that  $H_v$  is a *consistent horoball* at  $v$ . In particular, if  $v$  is a fixed point of a parabolic element of  $\Gamma$  then there exists a consistent horoball at  $v$ .

The Poincaré polyhedron theorem states:

**Theorem 2.3** [Mostow 1980, Theorem 6.3.2; Deraux et al. 2015, Theorem 3.2]. *Suppose that  $D$  is a polyhedron on  $\mathbf{H}_{\mathbb{C}}^2$  with sides contained in bisectors together with a side pairing. Let  $\Upsilon < \text{PU}(2, 1)$  be a discrete group of automorphisms of  $D$ . Let  $\Gamma$  be the group generated by  $\Upsilon$  and the side pairing maps. Suppose that the cycle condition holds at all ridges of  $D$  and that there is a consistent horoball at all points (if any) where sides of  $D$  are asymptotic. Then:*

- (1)  $\Gamma$  is discrete.
- (2) The images of  $D$  under the cosets of  $\Upsilon$  in  $\Gamma$  tessellate  $\mathbf{H}_{\mathbb{C}}^2$ .
- (3) A fundamental domain for  $\Gamma$  may be obtained by intersecting  $D$  with a fundamental domain for  $\Upsilon$ .
- (4) A presentation for  $\Gamma$  is given as follows. The generators are a generating set for  $\Upsilon$  together with all side pairing maps. The relations are generated by all relations in  $\Upsilon$ , all reflection relations and all cycle relations.

### 3. The generators


Consider complex reflections  $I_1$  and  $I_2$  in  $\text{SU}(2, 1)$  so that  $I_1I_2$  has order  $n$  and fixes the origin  $o$ . Writing  $c = \cos(\pi/n)$  and  $s = \sin(\pi/n)$ , we may choose  $I_1$  and  $I_2$  to be

$$(3-1) \quad I_1 = \begin{bmatrix} -c & s & 0 \\ s & c & 0 \\ 0 & 0 & -1 \end{bmatrix}, \quad I_2 = \begin{bmatrix} -c & -s & 0 \\ -s & c & 0 \\ 0 & 0 & -1 \end{bmatrix}.$$

Note that polar vectors of  $I_1$  and  $I_2$  are

$$\mathbf{n}_1 = \begin{bmatrix} s \\ 1+c \\ 0 \end{bmatrix}, \quad \mathbf{n}_2 = \begin{bmatrix} -s \\ 1+c \\ 0 \end{bmatrix}.$$

We want to find  $I_3$  so that  $I_1I_3$  and  $I_2I_3$  both have order 3. Conjugating by a diagonal map  $\text{diag}(e^{i\psi}, e^{i\psi}, e^{-2i\psi})$  if necessary, we may suppose that the polar



**Figure 1.** The  $2n$ -gon in the hyperbolic plane made up of  $2n$  copies of a  $(3, 3, n)$  triangle.

vector of  $I_3$  is

$$n_3 = \begin{bmatrix} a \\ be^{i\theta} \\ d - 1 \end{bmatrix},$$

where  $a, b, d$  are nonnegative real numbers satisfying  $a^2 + b^2 - (d - 1)^2 = 2(d - 1)$ , that is,  $a^2 + b^2 - d^2 = -1$ . Furthermore, complex conjugating if necessary, we may always assume  $\theta \in [0, \pi]$ . Then

$$(3-2) \quad I_3 = \begin{bmatrix} -1 + a^2/(d - 1) & abe^{-i\theta}/(d - 1) & -a \\ abe^{i\theta}/(d - 1) & -1 + b^2/(d - 1) & -be^{i\theta} \\ a & be^{-i\theta} & -d \end{bmatrix}.$$

It is easy to check that  $I_3$  lies in  $SU(2, 1)$ , has order 2 and polar vector  $n_3$ .

**Lemma 3.1.** *Let  $I_1, I_2$  and  $I_3$  be given by (3-1) and (3-2). If  $I_1 I_3$  and  $I_2 I_3$  have order 3 then  $\theta = \pi/2$  and*

$$(3-3) \quad c(a^2 - b^2) = d(d - 1).$$

*Proof.* The condition that  $I_1 I_3$  and  $I_2 I_3$  have order 3 is equivalent to  $\text{tr}(I_1 I_3) = \text{tr}(I_2 I_3) = 0$ . That is,

$$\frac{-c(a^2 - b^2) + 2sab \cos \theta}{d - 1} + d = \frac{-c(a^2 - b^2) - 2sab \cos \theta}{d - 1} + d = 0.$$

The result follows directly. □

From now on, we write  $\theta = \pi/2$  in (3-2). Since we know  $a^2 + b^2 = d^2 - 1$  and  $a^2 - b^2 = d(d - 1)/c$ , we immediately have

$$(3-4) \quad a^2 = (d - 1)(1 + d + d/c)/2, \quad b^2 = (d - 1)(1 + d - d/c)/2.$$

**Corollary 3.2.** *Let*

$$\iota : \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix} \mapsto \begin{bmatrix} \bar{z}_1 \\ -\bar{z}_2 \\ \bar{z}_3 \end{bmatrix}.$$

*Then  $\iota$  has order 2 and*

$$\iota I_1 \iota = I_2, \quad \iota I_2 \iota = I_1, \quad \iota I_3 \iota = I_3.$$

*Proof.* It is easy to see that  $\iota^2$  is the identity. A simple calculation shows  $\iota(\mathbf{n}_1) = \mathbf{n}_2$  and  $\iota(\mathbf{n}_3) = \mathbf{n}_3$ , using  $e^{i\theta} = i$ . □

**Lemma 3.3.** *The group  $\langle I_1, I_2, I_3 \rangle$  is determined up to conjugacy by the variable  $d$ , which lies in the interval  $1 < d \leq c/(1 - c)$ . Moreover,  $\langle I_1, I_2, I_3 \rangle$  lies in  $SO(2, 1)$  when  $d = c/(1 - c)$ .*

*Proof.* We have conjugated so that  $I_1$  and  $I_2$  have the form (3-1), and  $I_3$  has the form (3-2) with  $\theta = \pi/2$ . After this conjugation, the only remaining parameters are the nonnegative real numbers  $a, b$  and  $d$ . Using (3-4) these are completely determined by  $d$ . Moreover, again using (3-4) we see that  $a^2$  and  $b^2$  are nonnegative if and only if  $d \geq 1$  and  $d \leq c/(1 - c)$ . We cannot have  $d = 1$  or else  $\mathbf{n}_3$  is the zero vector. Thus  $1 < d \leq c/(1 - c)$ . Finally, when  $d = c/(1 - c)$ , we have  $b = 0$  and the entries of  $I_3$  are all real. □

**Lemma 3.4.** *Let  $I_1, I_2$  and  $I_3$  be given by (3-1) and (3-2). Suppose  $I_1 I_3$  and  $I_2 I_3$  have order 3. Then  $I_1 I_3 I_2 I_3$  is elliptic if and only if  $d < 3/(4s^2)$ .*

*Proof.* Calculating directly, we see that

$$\begin{aligned} \text{tr}(I_1 I_3 I_2 I_3) &= \frac{c^2(a^2 - b^2)^2}{(d - 1)^2} + \frac{2(c^2 - s^2)(d - 1 - a^2 - b^2)}{d - 1} - 2c(a^2 - b^2) + d^2 \\ &= 4s^2 d. \end{aligned}$$

(We could have derived this using the formulae in [Pratoussevitch 2005].) The condition that  $I_1 I_3 I_2 I_3$  is elliptic is equivalent to  $3 > \text{tr}(I_1 I_3 I_2 I_3) = 4s^2 d$ . □

Thus, our parameter space for  $\langle I_1, I_2, I_3 \rangle$  with  $I_1 I_3 I_2 I_3$  nonelliptic is given by

$$(3-5) \quad \frac{3}{4s^2} \leq d \leq \frac{c}{1 - c}.$$

Note that the condition  $n > 3$  implies both  $3/(4s^2) > 1$  and  $c/(1 - c) > 1$ . For example, when  $n = 4$  we have  $c = s = 1/\sqrt{2}$  and our range becomes

$$3/2 \leq d \leq \sqrt{2} + 1.$$

### 4. The bisectors

We define a polyhedron  $D$  bounded by sides contained in  $2n$  bisectors.

**Definition 4.1.** For  $k \in \mathbb{Z}$ , define the involution  $A_k \in \langle I_1, I_2, I_3 \rangle$  as follows:

- (1) If  $k = 2m$  is an even integer then  $A_k = (I_2 I_1)^{k/2} I_3 (I_1 I_2)^{k/2}$ .
- (2) If  $k = 2m + 1$  is an odd integer then  $A_k = (I_2 I_1)^{(k-1)/2} I_2 I_3 I_2 (I_1 I_2)^{(k-1)/2}$ .

Let  $o$  be the fixed point of  $I_1 I_2$  in  $\mathbf{H}_{\mathbb{C}}^2$ . For all integers  $k$ , the bisector  $\mathcal{B}_k$  is defined to be the bisector equidistant from  $o$  and  $A_k(o)$ . Note that in both cases  $A_{k+2n} = A_k$  and so  $\mathcal{B}_{k+2n} = \mathcal{B}_k$ . This gives  $2n$  bisectors  $\mathcal{B}_{-n+1}$  to  $\mathcal{B}_n$  and we may take the index  $k \bmod 2n$ .

The following lemma follows immediately from the definition.

**Lemma 4.2.** Let  $\mathcal{B}_{-n+1}$  to  $\mathcal{B}_n$  be as defined in Definition 4.1. Then for each  $k \bmod 2n$  and each  $m \bmod n$ :

- (1) The map  $(I_2 I_1)^m$  sends  $\mathcal{B}_k$  to  $\mathcal{B}_{2m+k}$ .
- (2) The map  $(I_2 I_1)^m I_2$  sends  $\mathcal{B}_k$  to  $\mathcal{B}_{2m+1-k}$ . In particular, the map  $(I_2 I_1)^k I_2$  sends  $\mathcal{B}_k$  to  $\mathcal{B}_{k+1}$ .
- (3) The antiholomorphic involution  $\iota$  defined in Corollary 3.2 sends  $\mathcal{B}_k$  to  $\mathcal{B}_{-k}$ . In particular, the map  $(I_2 I_1)^m I_2 \iota$  sends  $\mathcal{B}_k$  to  $\mathcal{B}_{2m+1+k}$ .

The main result of this section is that the combinatorial configuration of the bisectors does not change as  $d$  decreases from  $c/(1-c)$  to  $3/(4s^2)$ . More precisely:

**Theorem 4.3.** Let  $\mathcal{B}_{-n+1}$  to  $\mathcal{B}_n$  be as defined in Definition 4.1. Suppose that  $3/(4s^2) \leq d \leq c/(1-c)$ . Then, taking the indices  $\bmod 2n$ , for each  $k$ :

- (1) The bisector  $\mathcal{B}_k$  intersects  $\mathcal{B}_{k\pm 1}$  in a Giraud disc. This Giraud disc is preserved by  $A_k A_{k\pm 1}$ , which has order 3.
- (2) The intersection of  $\mathcal{B}_k$  with  $\mathcal{B}_{k\pm 2}$  is contained in the halfspace bounded by  $\mathcal{B}_{k\pm 1}$  not containing  $o$ .
- (3) The bisector  $\mathcal{B}_k$  does not intersect  $\mathcal{B}_{k\pm \ell}$  for  $3 \leq \ell \leq n$ . Moreover, the boundaries of these bisectors are disjoint except for when  $\ell = 3$  and  $d = 3/(4s^2)$ , in which case the boundaries intersect in a single point, which is a parabolic fixed point.

As a corollary to this theorem, we can use the Poincaré polyhedron theorem to prove the “if” part of Theorem 1.6.

**Corollary 4.4.** Let  $A_{-n+1}$  to  $A_n$  and  $\mathcal{B}_{-n+1}$  to  $\mathcal{B}_n$  be as in Theorem 4.3. Suppose that  $3/(4s^2) \leq d \leq c/(1-c)$ . Let  $D$  be the polyhedron in  $\mathbf{H}_{\mathbb{C}}^2$  containing  $o$  and bounded by  $\mathcal{B}_{-n+1}$  to  $\mathcal{B}_n$ . Then the maps  $A_{-n+1}$  to  $A_n$  form a side paring for  $D$  that satisfies the conditions of the Poincaré polyhedron theorem, Theorem 2.3. In particular,  $\langle I_1, I_2, I_3 \rangle$  is a discrete and faithful representation of  $\Delta_{3,3,n}$ .

*Proof.* Since  $A_k$  is an involution, it is clear that the  $\{A_k\}$  form a side pairing for  $D$ . Now consider the ridge  $r_k = \mathcal{B}_k \cap \mathcal{B}_{k+1}$ . Applying either of the side pairing maps  $A_k$  or  $A_{k+1}$  sends this ridge to itself. We then apply  $P_k = (I_2 I_1)^k I_2$  to obtain the cycle transformation  $P_k A_k$ . When  $k$  is even,

$$P_k A_k = (I_2 I_1)^k I_2 (I_2 I_1)^{k/2} I_3 (I_1 I_2)^{k/2} = (I_2 I_1)^{k/2} I_2 I_3 (I_1 I_2)^{k/2},$$

and when  $k$  is odd,

$$\begin{aligned} P_k A_k &= (I_2 I_1)^k I_2 (I_2 I_1)^{(k-1)/2} I_2 I_3 I_2 (I_1 I_2)^{(k-1)/2} \\ &= (I_2 I_1)^{(k+1)/2} I_3 I_1 (I_1 I_2)^{(k+1)/2}. \end{aligned}$$

In both cases,  $P_k A_k$  is equal to  $A_k A_{k+1}$ , which has order 3. There is a neighbourhood  $U_k$  of the ridge  $r_k$  for which the intersection of  $U_k$  with  $D$  is the same as its intersection with the Dirichlet domain for  $\langle P_k A_k \rangle$ . Therefore, we have local tessellation around all the ridges of  $D$  using the argument of [Proposition 2.1](#).

All the other sides of  $D$  are disjoint, apart from when  $d = 3/(4s^2)$ , in which case  $\mathcal{B}_k$  and  $\mathcal{B}_{k\pm 3}$  are asymptotic at a point of  $\partial \mathbf{H}_{\mathbb{C}}^2$ . This point is a parabolic fixed point, as required.

Finally, each side yields the reflection relation  $A_k^2$ , which is conjugate to  $I_3^2$ . The cycle relations give  $(P_k A_k)^3$ , which are conjugate to  $(I_2 I_3)^3$  when  $k$  is even and  $(I_3 I_1)^3$  when  $k$  is odd. In addition we have the relations from  $\Upsilon = \langle I_1, I_2 \rangle$ , which are  $I_1^2, I_2^2$  and  $(I_1 I_2)^n$ . From the Poincaré theorem, all other relations may be deduced from these. Thus  $\langle I_1, I_2, I_3 \rangle$  is a faithful representation of  $\Delta_{3,3,n}$ .  $\square$

Write  $c_k = \cos(k\pi/n)$  and  $s_k = \sin(k\pi/n)$ . Then

$$(I_2 I_1)^m = \begin{bmatrix} c_{2m} & -s_{2m} & 0 \\ s_{2m} & c_{2m} & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad (I_2 I_1)^m I_2 = \begin{bmatrix} -c_{2m+1} & -s_{2m+1} & 0 \\ -s_{2m+1} & c_{2m+1} & 0 \\ 0 & 0 & -1 \end{bmatrix}.$$

We have

$$(I_2 I_1)^m I_3(o) = \begin{bmatrix} -c_{2m}a + s_{2m}bi \\ -s_{2m}a - c_{2m}bi \\ -d \end{bmatrix}, \quad (I_1 I_2)^m I_3(o) = \begin{bmatrix} -c_{2m}a - s_{2m}bi \\ s_{2m}a - c_{2m}bi \\ -d \end{bmatrix}.$$

Also

$$(I_2 I_1)^m I_2 I_3(o) = \begin{bmatrix} c_{2m+1}a + s_{2m+1}bi \\ s_{2m+1}a - c_{2m+1}bi \\ d \end{bmatrix}, \quad (I_1 I_2)^m I_1 I_3(o) = \begin{bmatrix} c_{2m+1}a - s_{2m+1}bi \\ -s_{2m+1}a - c_{2m+1}bi \\ d \end{bmatrix}.$$

We begin by proving [Theorem 4.3\(1\)](#).

**Proposition 4.5.** *For each  $-n+1 \leq k \leq n$ , the bisectors  $\mathcal{B}_k$  and  $\mathcal{B}_{k\pm 1}$  (with indices taken mod  $2n$ ) intersect in  $\mathbf{H}_{\mathbb{C}}^2$  in a Giraud disc. This Giraud disc is preserved by  $(I_2 I_1)^{k/2} (I_2 I_3) (I_1 I_2)^{k/2}$  when  $k$  is even and  $(I_2 I_1)^{(k+1)/2} (I_3 I_1) (I_1 I_2)^{(k+1)/2}$  when  $k$  is odd.*

*Proof.* Using Lemma 4.2 we need only consider  $k = 0$  and  $k = 1$ . The bisectors  $\mathcal{B}_0$  and  $\mathcal{B}_1$  are equidistant from  $o$  and from  $I_3(o) = I_3I_2(o)$  and from  $I_2I_3(o)$  respectively. Observe that  $I_2I_3$  does not fix  $o$ . Since the map  $I_2I_3$  has order 3, the Dirichlet domain with centre  $o$  for the cyclic group  $\langle I_2I_3 \rangle$  only contains faces contained in these two bisectors. The intersection is a Giraud disc invariant under powers of  $I_2I_3$  by construction.  $\square$

Next we prove Theorem 4.3(3) in the case where  $\ell = 2m + 1$  is odd.

**Proposition 4.6.** *Suppose that  $3/(4s^2) \leq d \leq c/(1 - c)$ . For each  $-n + 1 \leq k \leq n$  and  $1 \leq m \leq (n - 1)/2$ , the bisectors  $\mathcal{B}_k$  and  $\mathcal{B}_{k \pm (2m+1)}$  (with indices taken mod  $2n$ ) do not intersect in  $\mathbf{H}_{\mathbb{C}}^2$ . Moreover, their closures intersect on  $\partial \mathbf{H}_{\mathbb{C}}^2$  if and only if  $d = 3/(4s^2)$  and  $m = 1$ . In the latter case, the closures intersect in a unique point, which is a parabolic fixed point.*

*Proof.* Using Lemma 4.2 we need only consider  $\mathcal{B}_0$  and  $\mathcal{B}_{2m+1}$ . These bisectors are equidistant from  $o$  and  $I_3(o) = I_3I_2(I_1I_2)^m(o)$  and from  $(I_2I_1)^mI_2I_3(o)$  respectively. Consider the Dirichlet domain with centre  $o$  for the cyclic group  $\langle (I_2I_1)^mI_2I_3 \rangle$ . We claim that this Dirichlet domain has exactly two sides and these sides are disjoint. To do so, we use Phillips’ theorem, Proposition 2.2.

A brief calculation shows that

$$\text{tr}((I_2I_1)^mI_2I_3) = -c_{2m+1} \frac{a^2 - b^2}{d - 1} + d = \frac{d(c - c_{2m+1})}{c} = \frac{2ds_{m+1}s_m}{c}.$$

When  $1 \leq m \leq (n - 1)/2$ , we have

$$s_ms_{m+1} \geq ss_2 = 2s^2c$$

with equality if and only if  $m = 1$ . Therefore,

$$\text{tr}((I_2I_1)^mI_2I_3) = 2ds_{m+1}s_m/c \geq 4ds^2$$

with equality if and only if  $m = 1$ . Hence, when  $4ds^2 \geq 3$ , we have  $(I_2I_1)^mI_2I_3$  is nonelliptic with real trace, and is loxodromic unless  $m = 1$  and  $d = 3/(4s^2)$ . By Phillips’ theorem we see that any Dirichlet domain for  $\langle (I_2I_1)^mI_2I_3 \rangle$  has two faces and these faces do not intersect in  $\mathbf{H}_{\mathbb{C}}^2$ .

In fact, when  $d = 3/(4s^2)$  and  $m = 1$ , the bisectors  $\mathcal{B}_0$  and  $\mathcal{B}_3$  are asymptotic on the boundary of  $\mathbf{H}_{\mathbb{C}}^2$  at the (parabolic) fixed point of  $I_2I_1I_2I_3$ .  $\square$

**Proposition 4.7.** (i) *Suppose  $p = [z, w, 1]^t$  lies on  $\mathcal{B}_{2\ell} \cap \mathcal{B}_{-2\ell}$ . Then for some angles  $\theta, \phi$ , we have*

$$z = \frac{s_{2\ell}a(\cos \theta e^{i\phi} + d) - c_{2\ell}b \sin \theta e^{i\phi}}{c_{2\ell}s_{2\ell}(a^2 - b^2)},$$

$$w = \frac{-s_{2\ell}bi(\cos \theta e^{i\phi} + d) + c_{2\ell}ai \sin \theta e^{i\phi}}{c_{2\ell}s_{2\ell}(a^2 - b^2)}.$$

(ii) Suppose  $p = [z, w, 1]^t$  lies on  $\mathcal{B}_{2\ell+1} \cap \mathcal{B}_{-2\ell-1}$ . Then for some angles  $\theta, \phi$ , we have

$$z = \frac{s_{2\ell+1}a(\cos \theta e^{i\phi} + d) - c_{2\ell+1}b \sin \theta e^{i\phi}}{c_{2\ell+1}s_{2\ell+1}(a^2 - b^2)},$$

$$w = \frac{s_{2\ell+1}bi(\cos \theta e^{i\phi} + d) - c_{2\ell+1}ai \sin \theta e^{i\phi}}{c_{2\ell+1}s_{2\ell+1}(a^2 - b^2)}.$$

*Proof.* First consider the bisector intersection from (i). Then  $z$  and  $w$  satisfy

$$1 = |z(-c_{2\ell}a + s_{2\ell}bi) + w(s_{2\ell}a + c_{2\ell}bi) + d|,$$

$$1 = |z(-c_{2\ell}a - s_{2\ell}bi) + w(-s_{2\ell}a + c_{2\ell}bi) + d|.$$

Expanding out, adding and subtracting yields

$$1 = |zc_{2\ell}a - wc_{2\ell}bi - d|^2 + |zs_{2\ell}bi + ws_{2\ell}a|^2,$$

$$0 = 2 \operatorname{Re}((zc_{2\ell}a - wc_{2\ell}bi - d)(-\bar{z}s_{2\ell}bi + \bar{w}s_{2\ell}a)).$$

Thus we can write

$$zc_{2\ell}a - wc_{2\ell}bi - d = \cos \theta e^{i\phi},$$

$$zs_{2\ell}bi + ws_{2\ell}a = i \sin \theta e^{i\phi}.$$

Inverting these equations yields

$$z = \frac{s_{2\ell}a(\cos \theta e^{i\phi} + d) - c_{2\ell}b \sin \theta e^{i\phi}}{c_{2\ell}s_{2\ell}(a^2 - b^2)},$$

$$w = \frac{-s_{2\ell}bi(\cos \theta e^{i\phi} + d) + c_{2\ell}ai \sin \theta e^{i\phi}}{c_{2\ell}s_{2\ell}(a^2 - b^2)}.$$

For the second bisector intersection, we have

$$1 = |z(c_{2\ell+1}a + s_{2\ell+1}bi) + w(-s_{2\ell+1}a + c_{2\ell+1}bi) - d|^2,$$

$$1 = |z(c_{2\ell+1}a - s_{2\ell+1}bi) + w(s_{2\ell+1}a + c_{2\ell+1}bi) - d|^2.$$

Expanding out, adding and subtracting yields

$$1 = |zc_{2\ell+1}a + wc_{2\ell+1}bi - d|^2 + |-zs_{2\ell+1}bi + ws_{2\ell+1}a|^2,$$

$$0 = 2 \operatorname{Re}((zc_{2\ell+1}a + wc_{2\ell+1}bi - d)(\bar{z}s_{2\ell+1}bi + \bar{w}s_{2\ell+1}a)).$$

So once again we have

$$zc_{2\ell+1}a + wc_{2\ell+1}bi - d = \cos \theta e^{i\phi},$$

$$-zs_{2\ell+1}bi + ws_{2\ell+1}a = -i \sin \theta e^{i\phi}.$$



Thus,

$$z = \frac{s_{2\ell+1}a(\cos \theta e^{i\phi} + d) - c_{2\ell+1}b \sin \theta e^{i\phi}}{c_{2\ell+1}s_{2\ell+1}(a^2 - b^2)},$$

$$w = \frac{s_{2\ell+1}bi(\cos \theta e^{i\phi} + d) - c_{2\ell+1}ai \sin \theta e^{i\phi}}{c_{2\ell+1}s_{2\ell+1}(a^2 - b^2)}. \quad \square$$

We can now prove [Theorem 4.3\(3\)](#) in the case where  $\ell = 2m$  is even.

**Proposition 4.8.** *Suppose that  $3/(4s^2) \leq d \leq c/(1 - c)$ . For each  $-n + 1 \leq k \leq n$  and  $2 \leq m \leq n/2$ , the bisectors  $\mathcal{B}_k$  and  $\mathcal{B}_{k\pm 2m}$  (with indices taken mod  $2n$ ) do not intersect in complex hyperbolic space.*

*Proof.* Using [Lemma 4.2](#), we need only consider  $\mathcal{B}_m$  and  $\mathcal{B}_{-m}$  where  $2 \leq m \leq n/2$ .

Using [Proposition 4.7](#) we see that an intersection point  $p = [z, w, 1]^t$  of  $\mathcal{B}_m$  and  $\mathcal{B}_{-m}$  must satisfy

$$z = \frac{s_m a(\cos \theta e^{i\phi} + d) - c_m b \sin \theta e^{i\phi}}{c_m s_m (a^2 - b^2)},$$

$$w = \pm \frac{-s_m bi(\cos \theta e^{i\phi} + d) + c_m ai \sin \theta e^{i\phi}}{c_m s_m (a^2 - b^2)}.$$

We claim that  $|z|^2 + |w|^2 \geq 1$  and so such a point does not lie in  $\mathbf{H}_{\mathbb{C}}^2$ . We have

$$\begin{aligned} & c_m^2 s_m^2 (a^2 - b^2)^2 (|z|^2 + |w|^2 - 1) \\ &= |s_m a(\cos \theta e^{i\phi} + d) - c_m b \sin \theta e^{i\phi}|^2 \\ &\quad + |-s_m bi(\cos \theta e^{i\phi} + d) + c_m ai \sin \theta e^{i\phi}|^2 - c_m^2 s_m^2 (a^2 - b^2)^2 \\ &= s_m^2 (a^2 + b^2) (\cos^2 \theta + 2d \cos \theta \cos \phi + d^2) \\ &\quad - 2c_m s_m ab (2 \cos \theta \sin \theta + 2d \sin \theta \cos \phi) \\ &\quad + c_m^2 (a^2 + b^2) \sin^2 \theta - c_m^2 s_m^2 (a^2 + b^2)^2 + 4c_m^2 s_m^2 a^2 b^2 \\ &= s_m^2 (d^2 - 1) (\cos^2 \theta + 2d \cos \theta \cos \phi + d^2) \\ &\quad - 4c_m s_m ab (\cos \theta \sin \theta + d \sin \theta \cos \phi) \\ &\quad + c_m^2 (d^2 - 1) \sin^2 \theta - c_m^2 s_m^2 (d^2 - 1)^2 + 4c_m^2 s_m^2 a^2 b^2 \\ &= (\cos \theta \sin \theta + d \sin \theta \cos \phi - 2c_m s_m ab)^2 + d^2 \sin^2 \theta \sin^2 \phi \\ &\quad + (s_m^2 (d^2 - 1) - \sin^2 \theta) (\cos^2 \theta + 2d \cos \theta \cos \phi + d^2 - c_m^2 (d^2 - 1)) \\ &\geq (s_m^2 (d^2 - 1) - \sin^2 \theta) (\cos^2 \theta + 2d \cos \theta \cos \phi + d^2 - c_m^2 (d^2 - 1)). \end{aligned}$$

Therefore, it is sufficient to prove

$$(4-1) \quad 0 < s_m^2 (d^2 - 1) - \sin^2 \theta,$$

$$(4-2) \quad 0 < \cos^2 \theta + 2d \cos \theta \cos \phi + d^2 - c_m^2 (d^2 - 1).$$

In order to prove these inequalities, we need to use the lower bound on  $d$ . Using  $m \geq 2$  and  $d \geq 3/(4s^2)$ , we have

$$(4-3) \quad (1 - c_m)d \geq (1 - c_2)d = 2s^2d \geq 3/2.$$

We also use  $s_m^2 = 1 - c_m^2 = (1 - c_m)(1 + c_m)$  and  $c_m \geq 0$  (the latter uses  $m \leq n/2$ ).

First, we consider (4-1):

$$\begin{aligned} s_m^2(d^2 - 1) - \sin^2 \theta &= \frac{1 + c_m}{1 - c_m}((1 - c_m)d)^2 - 2 + c_m^2 + \cos^2 \theta \\ &\geq ((1 - c_m)d)^2 - 2 \\ &\geq 1/4, \end{aligned}$$

where the last inequality follows from (4-3). This proves (4-1).

Now consider (4-2):

$$\begin{aligned} \cos^2 \theta + 2d \cos \theta \cos \phi + d^2 - c_m^2(d^2 - 1) &= \frac{(d(1 - c_m) + \cos \theta \cos \phi)^2 + \cos^2 \theta \sin^2 \phi}{1 - c_m} \\ &\quad + \frac{c_m}{1 - c_m}((d(1 - c_m))^2 - \cos^2 \theta) + c_m^2 \\ &\geq \frac{c_m}{1 - c_m}(9/4 - \cos^2 \theta) \\ &> 0. \end{aligned}$$

Again we used (4-3). This proves (4-2) and so establishes the result.  $\square$

Propositions 4.6 and 4.8 complete the proof of Theorem 4.3(3). It remains to prove Theorem 4.3(2). That is, we must consider the intersection of  $\mathcal{B}_k$  and  $\mathcal{B}_{k \pm 2}$ .

Consider  $\mathcal{B}_1 \cap \mathcal{B}_{-1}$ . We claim that the fixed point of  $I_3 I_1 I_2 I_3$  (that is  $I_3(o)$ ) lies on  $\mathcal{B}_1 \cap \mathcal{B}_{-1}$ . The bisector  $\mathcal{B}_1$  consists of all points equidistant from  $o$  and  $A_1(o) = I_2 I_3 I_2(o) = I_2 I_3(o)$ . We have

$$\rho(I_3(o), I_2 I_3(o)) = \rho(o, I_3 I_2 I_3(o)) = \rho(o, I_2 I_3(o)).$$

The first equality follows since  $I_3$  is an isometry and the second since  $I_3 I_2 I_3 = I_2 I_3 I_2$  and  $I_2(o) = o$ . Thus  $I_3(o)$  lies on  $\mathcal{B}_1$ . A similar argument shows

$$\rho(I_3(o), I_1 I_3(o)) = \rho(o, I_1 I_3(o)).$$

and so  $I_3(o)$  lies on  $\mathcal{B}_{-1}$  as well. Thus  $\mathcal{B}_1 \cap \mathcal{B}_{-1}$  is nonempty, which can be seen in Figure 1. By symmetry, this comment also applies to the intersection of  $\mathcal{B}_k$  and  $\mathcal{B}_{k \pm 2}$ . We must show that this intersection never contributes a ridge of  $D$ .

**Proposition 4.9.** *Suppose that  $3/(4s^2) \leq d \leq c/(1 - c)$ . For each  $-n + 1 \leq k \leq n$ , all points of  $\mathcal{B}_k \cap \mathcal{B}_{k \pm 2}$  lie in the halfspace bounded by  $\mathcal{B}_{k \pm 1}$  not containing  $o$ .*

*Proof.* Using [Lemma 4.2](#) as before, it suffices to consider  $\mathcal{B}_1$  and  $\mathcal{B}_{-1}$ . We need to show that all points of  $\mathcal{B}_1 \cap \mathcal{B}_{-1}$  lie in the halfspace closer to  $I_3(o)$  than to  $o$ .

Suppose that  $p = [z, w, 1]^t$  lies on  $\mathcal{B}_1 \cap \mathcal{B}_{-1}$ . Using [Proposition 4.7\(ii\)](#) with  $m = 0$ , and using [\(3-3\)](#) to write  $c(a^2 - b^2) = d(d - 1)$ , we find

$$(4-4) \quad z = \frac{sa(\cos \theta e^{i\phi} + d) - cb \sin \theta e^{i\phi}}{sd(d-1)},$$

$$(4-5) \quad w = \frac{sbi(\cos \theta e^{i\phi} + d) - cai \sin \theta e^{i\phi}}{sd(d-1)}.$$

Note that we used [\(3-3\)](#) to simplify the denominator.

The point  $p = [z, w, 1]^t$  lies in the halfspace closer to  $I_3(o)$  than to  $o$  if and only if  $1 > |za - wbi - d|$ . We want to give this inequality in terms of  $\theta$ ,  $\phi$  and  $d$ . Suppose  $z$  and  $w$  satisfy [\(4-4\)](#) and [\(4-5\)](#) and consider  $za - wbi - d$ :

$$\begin{aligned} za - wbi - d &= \frac{sa^2(\cos \theta e^{i\phi} + d) - cab \sin \theta e^{i\phi}}{sd(d-1)} \\ &\quad + \frac{sb^2(\cos \theta e^{i\phi} + d) - cab \sin \theta e^{i\phi}}{sd(d-1)} - d \\ &= \frac{s(a^2 + b^2) \cos \theta e^{i\phi}}{sd(d-1)} - \frac{2cab \sin \theta e^{i\phi}}{sd(d-1)} + \frac{s(a^2 + b^2)d}{sd(d-1)} - d \\ &= \frac{s(d^2 - 1) \cos \theta e^{i\phi}}{sd(d-1)} - \frac{2cab \sin \theta e^{i\phi}}{sd(d-1)} + \frac{s(d^2 - 1)d}{sd(d-1)} - d \\ &= \frac{(d+1) \cos \theta e^{i\phi}}{d} - \frac{\sqrt{c^2(d+1)^2 - d^2} \sin \theta e^{i\phi}}{sd} + 1. \end{aligned}$$

Therefore,

$$\begin{aligned} |za - wbi - d|^2 - 1 &= \frac{(d+1)^2 \cos^2 \theta}{d^2} + \frac{c^2(d+1)^2 \sin^2 \theta}{s^2 d^2} - \frac{\sin^2 \theta}{s^2} \\ &\quad - \frac{2(d+1)\sqrt{c^2(d+1)^2 - d^2} \cos \theta \sin \theta}{sd^2} \\ &\quad + \frac{2(d+1) \cos \theta \cos \phi}{d} - \frac{2\sqrt{c^2(d+1)^2 - d^2} \sin \theta \cos \phi}{sd}. \end{aligned}$$

Arguing as in the proof of [Proposition 4.8](#), we have

$$\begin{aligned} |z|^2 + |w|^2 - 1 &= \left| \frac{sa(\cos \theta e^{i\phi} + d) - cb \sin \theta e^{i\phi}}{sd(d-1)} \right|^2 + \left| \frac{sbi(\cos \theta e^{i\phi} + d) - cai \sin \theta e^{i\phi}}{sd(d-1)} \right|^2 - 1 \end{aligned}$$

$$\begin{aligned}
 &= \frac{s^2(a^2 + b^2)|\cos \theta e^{i\phi} + d|^2}{s^2d^2(d-1)^2} + \frac{c^2(a^2 + b^2)\sin^2 \theta}{s^2d^2(d-1)^2} - 1 \\
 &\quad + \frac{isc(2abi)(2\cos \theta \sin \theta + 2d\sin \theta \cos \phi)}{s^2d^2(d-1)^2} \\
 &= \frac{(d+1)\cos^2 \theta}{d^2(d-1)} + \frac{2(d+1)\cos \theta \cos \phi}{d(d-1)} + \frac{d+1}{d-1} + \frac{c^2(d+1)\sin^2 \theta}{s^2d^2(d-1)} - 1 \\
 &\quad - \frac{2\sqrt{c^2(d+1)^2 - d^2}\cos \theta \sin \theta}{sd^2(d-1)} - \frac{2\sqrt{c^2(d+1)^2 - d^2}\sin \theta \cos \phi}{sd(d-1)} \\
 &= \frac{2}{d-1} + \frac{(d+1)\cos^2 \theta}{d^2(d-1)} + \frac{c^2(d+1)\sin^2 \theta}{s^2d^2(d-1)} - \frac{2\sqrt{c^2(d+1)^2 - d^2}\cos \theta \sin \theta}{sd^2(d-1)} \\
 &\quad + \frac{2(d+1)\cos \theta \cos \phi}{d(d-1)} - \frac{2\sqrt{c^2(d+1)^2 - d^2}\sin \theta \cos \phi}{sd(d-1)}.
 \end{aligned}$$

Now we eliminate  $\cos \phi$  using the equation for  $|za - wbi - d|^2$  derived above:

$$\begin{aligned}
 |z|^2 + |w|^2 - 1 &= \frac{1}{d-1}(|za - wbi - d|^2 - 1) + \frac{2\cos^2 \theta}{d-1} + \frac{2\sin^2 \theta}{d-1} \\
 &\quad + \frac{(d+1)\cos^2 \theta}{d^2(d-1)} + \frac{c^2(d+1)\sin^2 \theta}{s^2d^2(d-1)} - \frac{2\sqrt{c^2(d+1)^2 - d^2}\cos \theta \sin \theta}{sd^2(d-1)} \\
 &\quad - \frac{(d+1)^2\cos^2 \theta}{d^2(d-1)} - \frac{c^2(d+1)^2\sin^2 \theta}{s^2d^2(d-1)} + \frac{\sin^2 \theta}{s^2(d-1)} \\
 &\quad + \frac{2(d+1)\sqrt{c^2(d+1)^2 - d^2}\cos \theta \sin \theta}{sd^2(d-1)} \\
 &= \frac{1}{d-1}(|za - wbi - d|^2 - 1) \\
 &\quad + \frac{1}{d} \left( \cos \theta + \frac{\sqrt{c^2(d+1)^2 - d^2}\sin \theta}{s(d-1)} \right)^2 + \frac{(4s^2d - 3)\sin^2 \theta}{s^2(d-1)^2}.
 \end{aligned}$$

Since the last two terms are nonnegative, all points  $p = [z, w, 1]^t$  with  $z$  and  $w$  given by (4-4) and (4-5) and that satisfy  $|z|^2 + |w|^2 < 1$  must also satisfy  $|za - wbi - d| < 1$ . Geometrically, this means that all points in  $\mathbf{H}_{\mathbb{C}}^3$  that are on  $\mathcal{B}_1 \cap \mathcal{B}_{-1}$  are in the halfspace closer to  $I_3(o)$  than to  $o$ . This proves the result.  $\square$

This completes the proof of [Theorem 4.3](#).

### Acknowledgements

This research was carried out while John Parker and Jieyan Wang were visiting the Department of Mathematics at Hunan University. We would like to thank Yueping Jiang for making this visit possible. We would like to thank the referee for his comments which improved a previous version of this paper.

## References

- [Brehm 1990] U. Brehm, “The shape invariant of triangles and trigonometry in two-point homogeneous spaces”, *Geom. Dedicata* **33**:1 (1990), 59–76. MR 91c:53048 Zbl 0695.53038
- [Deraux 2015] M. Deraux, “On spherical CR uniformization of 3-manifolds”, *Exp. Math.* **24**:3 (2015), 355–370. MR 3359222
- [Deraux and Falbel 2015] M. Deraux and E. Falbel, “Complex hyperbolic geometry of the figure-eight knot”, *Geom. Topol.* **19**:1 (2015), 237–293. MR 3318751 Zbl 06413574
- [Deraux et al. 2015] M. Deraux, J. R. Parker, and J. Paupert, “New non-arithmetic complex hyperbolic lattices”, *Invent. Math.* (online publication May 2015).
- [Falbel and Wang 2014] E. Falbel and J. Wang, “Branched spherical CR structures on the complement of the figure-eight knot”, *Michigan Math. J.* **63**:3 (2014), 635–667. MR 3255694 Zbl 1300.32034
- [Falbel et al. 2015] E. Falbel, P.-V. Koseleff, and F. Rouillier, “Representations of fundamental groups of 3-manifolds into  $\mathrm{PGL}(3, \mathbb{C})$ : Exact computations in low complexity”, *Geom. Dedicata* **177**:1 (2015), 229–255. MR 3370032 Zbl 06468790
- [Goldman and Parker 1992] W. M. Goldman and J. R. Parker, “Complex hyperbolic ideal triangle groups”, *J. Reine Angew. Math.* **425** (1992), 71–86. MR 93c:20076 Zbl 0739.53055
- [Grossi 2007] C. H. Grossi, “On the type of triangle groups”, *Geom. Dedicata* **130** (2007), 137–148. MR 2008k:20092 Zbl 1139.53005
- [Mostow 1980] G. D. Mostow, “On a remarkable class of polyhedra in complex hyperbolic space”, *Pacific J. Math.* **86**:1 (1980), 171–276. MR 82a:22011 Zbl 0456.22012
- [Parker and Will 2015a] J. R. Parker and P. Will, “Complex hyperbolic free groups with many parabolic elements”, pp. 327–348 in *Groups, geometry and dynamics*, edited by C. S. Aravinda et al., Contemp. Math. **639**, Amer. Math. Soc., Providence, RI, 2015.
- [Parker and Will 2015b] J. R. Parker and P. Will, “A complex hyperbolic Riley slice”, preprint, 2015. arXiv 1510.01505
- [Phillips 1992] M. B. Phillips, “Dirichlet polyhedra for cyclic groups in complex hyperbolic space”, *Proc. Amer. Math. Soc.* **115**:1 (1992), 221–228. MR 93a:32042 Zbl 0768.53033
- [Pratoussevitch 2005] A. Pratoussevitch, “Traces in complex hyperbolic triangle groups”, *Geom. Dedicata* **111**:1 (2005), 159–185. MR 2006d:32036 Zbl 1115.32015
- [Schwartz 2001a] R. E. Schwartz, “Degenerating the complex hyperbolic ideal triangle groups”, *Acta Math.* **186**:1 (2001), 105–154. MR 2002e:20079 Zbl 0998.53050
- [Schwartz 2001b] R. E. Schwartz, “Ideal triangle groups, dented tori, and numerical analysis”, *Ann. of Math.* (2) **153**:3 (2001), 533–598. MR 2002j:57031 Zbl 1055.20040
- [Schwartz 2002] R. E. Schwartz, “Complex hyperbolic triangle groups”, pp. 339–349 in *Proceedings of the international congress of mathematicians, II: Invited lectures* (Beijing, 2002), edited by T. T. Li et al., Higher Ed. Press, Beijing, 2002. MR 2004b:57002 Zbl 1022.53034
- [Schwartz 2005] R. E. Schwartz, “A better proof of the Goldman–Parker conjecture”, *Geom. Topol.* **9** (2005), 1539–1601. MR 2006j:20063 Zbl 1098.20034
- [Schwartz 2007] R. E. Schwartz, *Spherical CR geometry and Dehn surgery*, Annals of Mathematics Studies **165**, Princeton Univ. Press, 2007. MR 2009a:32047 Zbl 1116.57016
- [Thompson 2010] J. M. Thompson, *Complex hyperbolic triangle groups*, Ph.D. thesis, Durham University, 2010, available at <http://core.ac.uk/download/pdf/85700.pdf>.

JOHN R. PARKER  
DEPARTMENT OF MATHEMATICAL SCIENCES  
DURHAM UNIVERSITY  
DURHAM  
DH1 3LE  
UNITED KINGDOM  
[j.r.parker@durham.ac.uk](mailto:j.r.parker@durham.ac.uk)

JIEYAN WANG  
ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE  
CHINESE ACADEMY OF SCIENCES  
BEIJING, 100190  
CHINA  
[jywang@hnu.edu.cn](mailto:jywang@hnu.edu.cn)

BAOHUA XIE  
COLLEGE OF MATHEMATICS AND ECONOMETRICS  
HUNAN UNIVERSITY  
CHANGSHA, 410082  
CHINA  
[xiexbh@hnu.edu.cn](mailto:xiexbh@hnu.edu.cn)

# PACIFIC JOURNAL OF MATHEMATICS

[msp.org/pjm](http://msp.org/pjm)

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2016 is US \$440/year for the electronic version, and \$600/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).


---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 280    No. 2    February 2016

---

Topological Molino's theory	257
JESÚS A. ÁLVAREZ LÓPEZ and MANUEL F. MOREIRA GALICIA	
Equivariant principal bundles and logarithmic connections on toric varieties	315
INDRANIL BISWAS, ARIJIT DEY and MAINAK PODDAR	
On a spectral theorem in paraorthogonality theory	327
KENIER CASTILLO, RUYMÁN CRUZ-BARROSO and FRANCISCO PERDOMO-PÍO	
Sigma theory and twisted conjugacy, II: Houghton groups and pure symmetric automorphism groups	349
DACIBERG L. GONÇALVES and PARAMESWARAN SANKARAN	
The second CR Yamabe invariant	371
PAK TUNG HO	
No hyperbolic pants for the 4-body problem with strong potential	401
CONNOR JACKMAN and RICHARD MONTGOMERY	
Unions of Lebesgue spaces and $A_1$ majorants	411
GREG KNESE, JOHN E. M <sup>C</sup> CARTHY and KABE MOEN	
Complex hyperbolic $(3, 3, n)$ triangle groups	433
JOHN R. PARKER, JIEYAN WANG and BAOHUA XIE	
Topological aspects of holomorphic mappings of hyperquadrics from $\mathbb{C}^2$ to $\mathbb{C}^3$	455
MICHAEL REITER	
2-Blocks with minimal nonabelian defect groups III	475
BENJAMIN SAMBALE	
Number of singularities of stable maps on surfaces	489
TAKAHIRO YAMAMOTO	