

*Pacific
Journal of
Mathematics*

***E*-POLYNOMIAL OF THE $SL(3, \mathbb{C})$ -CHARACTER VARIETY
OF FREE GROUPS**

SEAN LAWTON AND VICENTE MUÑOZ

***E*-POLYNOMIAL OF THE $SL(3, \mathbb{C})$ -CHARACTER VARIETY OF FREE GROUPS**

SEAN LAWTON AND VICENTE MUÑOZ

We compute the *E*-polynomial of the character variety of representations of a rank r free group in $SL(3, \mathbb{C})$. Expanding upon techniques of Logares, Muñoz and Newstead (*Rev. Mat. Complut.* **26:2** (2013), 635–703), we stratify the space of representations and compute the *E*-polynomial of each geometrically described stratum using fibrations. Consequently, we also determine the *E*-polynomial of its smooth, singular, and abelian loci and the corresponding Euler characteristic in each case. Along the way, we give a new proof of results of Cavazos and Lawton (*Int. J. Math.* **25:6** (2014), 1450058).

1. Introduction

Let Γ be a finitely generated group, and let G be a complex reductive algebraic group. The space of G -representations is

$$\mathcal{R}(\Gamma, G) = \{\rho : \Gamma \rightarrow G \mid \rho \text{ is a group morphism}\}.$$

Writing a presentation $\Gamma = \langle x_1, \dots, x_n \mid R_1, \dots, R_s \rangle$, we have that $\rho \in \mathcal{R}(\Gamma, G)$ is determined by the images $A_i = \rho(x_i)$, $1 \leq i \leq n$. Hence we can write $\rho = (A_1, \dots, A_n)$. These matrices are subject to the relations $R_j(A_1, \dots, A_n) = \text{Id}$, $1 \leq j \leq s$. Hence

$$\mathcal{R}(\Gamma, G) \cong \{(A_1, \dots, A_n) \in G^n \mid R_1(A_1, \dots, A_n) = \dots = R_s(A_1, \dots, A_n) = \text{Id}\}$$

is an affine algebraic set, since G is algebraic.

There is an action of G by conjugation on $\mathcal{R}(\Gamma, G)$, which is equivalent to the action of $PG = G/Z(G)$, where $Z(G)$ is the center of G , since the center acts trivially. The *G*-character variety of Γ is the GIT quotient

$$\mathcal{M}(\Gamma, G) = \mathcal{R}(\Gamma, G) // G,$$

which is an affine algebraic set by construction. Note that if we write $X := \mathcal{R}(\Gamma, G) = \text{Spec}(S)$, then $X // G = \text{Spec}(S^G)$.

MSC2010: 14D20, 20C15, 14L30, 20E05.

Keywords: *E*-polynomial, free group, $SL(3, \mathbb{C})$, character variety.

Every element $g \in \Gamma$ determines a character $\chi_g : X \rightarrow \mathbb{C}$, $\chi_g(\rho) = \text{tr}(\rho(g))$, with respect to an embedding $G \hookrightarrow \text{GL}(n, \mathbb{C})$. These regular functions $\chi_g \in S$ are invariant by conjugation, and hence $\chi_g \in S^G$. Consider the algebra of characters

$$T = \mathbb{C}[\chi_g \mid g \in \Gamma] \subset S^G,$$

and let $\chi(\Gamma, G) = \text{Spec}(T)$. There is a well-defined surjective map $\mathcal{M}(\Gamma, G) \rightarrow \chi(\Gamma, G)$, which is an isomorphism when $G = \text{SL}(n, \mathbb{C})$ among other examples; see [Sikora 2013].

In this paper we are interested in the character variety for the free group on r elements $\Gamma = F_r$ and for the group $G = \text{SL}(3, \mathbb{C})$. We compute the E -polynomial (also known as Hodge–Deligne polynomial) of $\mathcal{M}(F_r, \text{SL}(3, \mathbb{C}))$. The E -polynomial of $\mathcal{M}(F_r, \text{SL}(2, \mathbb{C}))$ has been computed in [Cavazos and Lawton 2014] by arithmetic methods (using the Weil conjectures). Recently, in [Mozgovoy and Reineke 2015], the E -polynomials of $\mathcal{M}(F_r, \text{PGL}(n, \mathbb{C}))$ have also been computed by arithmetic methods, where the result is given in the form of a generating function.

Here we use a geometric technique, introduced in [Logares et al. 2013], to compute E -polynomials of character varieties. This consists of stratifying the space of representations geometrically, and computing the E -polynomials of each stratum using the behavior of E -polynomials with fibrations. This technique is used in [Logares et al. 2013] for the case of $\Gamma = \pi_1(X)$ for a surface X of genus $g = 1, 2$ and $G = \text{SL}(2, \mathbb{C})$ (and also with one puncture, fixing the holonomy around the puncture). The case of $g = 3$ is worked out in [Martínez and Muñoz 2015a], the case of $g \geq 4$ in [Martínez and Muñoz 2015b], and the case of $g = 1$ with two punctures appears in [Logares and Muñoz 2014]. To implement this geometric technique for character varieties for $\text{SL}(n, \mathbb{C})$, for $n \geq 3$, we need to introduce the *equivariant Hodge–Deligne polynomial* with respect to a finite group action on an affine variety. This will be useful for studying character varieties of surface groups in $\text{SL}(n, \mathbb{C})$, $n \geq 3$.

We start by recovering the E -polynomials $e(\mathcal{M}(F_r, \text{SL}(2, \mathbb{C})))$ of [Cavazos and Lawton 2014] and $e(\mathcal{M}(F_r, \text{PGL}(2, \mathbb{C})))$ of [Mozgovoy and Reineke 2015], verifying that they are equal. Then we move to rank 3 to compute $e(\mathcal{M}(F_r, \text{SL}(3, \mathbb{C})))$ and $e(\mathcal{M}(F_r, \text{PGL}(3, \mathbb{C})))$. They turn out to be equal again. The latter one coincides, as expected, with the polynomial obtained in [Mozgovoy and Reineke 2015].

Unlike the methods used to obtain $e(\mathcal{M}(F_r, \text{PGL}(3, \mathbb{C})))$ in [Mozgovoy and Reineke 2015], our method provides an explicit geometric description of, and the E -polynomial for, each stratum. By results in [Florentino and Lawton 2012] this additional information determines the E -polynomial of the smooth and singular loci of $\mathcal{M}(F_r, \text{SL}(3, \mathbb{C}))$, and by [Florentino and Lawton 2014] also determines the E -polynomial of the abelian character variety $\mathcal{M}(\mathbb{Z}^r, \text{SL}(3, \mathbb{C}))$.

Our main theorem is thus:

Theorem 1. *The E -polynomials $e(\mathcal{M}(F_r, \mathrm{SL}(3, \mathbb{C})))$ and $e(\mathcal{M}(F_r, \mathrm{PGL}(3, \mathbb{C})))$ are both equal to*

$$(q^8 - q^6 - q^5 + q^3)^{r-1} + (q - 1)^{2r-2}(q^{3r-3} - q^r) + \frac{1}{6}(q - 1)^{2r-2}q(q + 1) + \frac{1}{2}(q^2 - 1)^{r-1}q(q - 1) + \frac{1}{3}(q^2 + q + 1)^{r-1}q(q + 1) - (q - 1)^{r-1}q^{r-1}(q^2 - 1)^{r-1}(2q^{2r-2} - q).$$

From the definition of the E -polynomial of a variety X , the classical Euler characteristic is given by $\chi(X) = e(X; 1, 1)$. Consequently, we deduce:

Corollary 2. *Let $r \geq 2$. Then $\mathcal{M}(F_r, \mathrm{SL}(3, \mathbb{C}))$, $\mathcal{M}(F_r, \mathrm{PGL}(3, \mathbb{C}))$, and (by [Florentino and Lawton 2009]) $\mathcal{M}(F_r, \mathrm{SU}(3))$, have Euler characteristic given by $2 \cdot 3^{r-2}$. The Euler characteristic of $\mathcal{M}(\mathbb{Z}^r, \mathrm{SL}(3, \mathbb{C}))$, and (by [Florentino and Lawton 2014]) also $\mathcal{M}(\mathbb{Z}^r, \mathrm{SU}(3))$, is given 3^{r-2} .*

2. Hodge structures and E -polynomials

Our main goal is to compute the E -polynomial (Hodge–Deligne polynomial) of the $\mathrm{SL}(3, \mathbb{C})$ -character variety of a free group. We will follow the methods in [Logares et al. 2013], so we collect some basic results from [loc. cit.] in this section.

We start by reviewing the definition of the Hodge–Deligne polynomial. A pure Hodge structure of weight k consists of a finite dimensional complex vector space H with a real structure, and a decomposition $H = \bigoplus_{k=p+q} H^{p,q}$ such that $H^{q,p} = \overline{H^{p,q}}$, the bar meaning complex conjugation on H . A Hodge structure of weight k gives rise to the so-called Hodge filtration, which is a descending filtration $F^p = \bigoplus_{s \geq p} H^{s,k-s}$. We define $\mathrm{Gr}_F^p(H) := F^p / F^{p+1} = H^{p,k-p}$.

A mixed Hodge structure consists of a finite dimensional complex vector space H with a real structure, an ascending (weight) filtration $\dots \subset W_{k-1} \subset W_k \subset \dots \subset H$ (defined over \mathbb{R}) and a descending (Hodge) filtration F such that F induces a pure Hodge structure of weight k on each $\mathrm{Gr}_k^W(H) = W_k / W_{k-1}$. We define $H^{p,q} := \mathrm{Gr}_F^p \mathrm{Gr}_{p+q}^W(H)$ and write $h^{p,q}$ for the *Hodge number* $h^{p,q} := \dim H^{p,q}$.

Let Z be any quasiprojective algebraic variety (possibly nonsmooth or noncompact). The cohomology groups $H^k(Z)$ and the cohomology groups with compact support $H_c^k(Z)$ are endowed with mixed Hodge structures [Deligne 1971; 1974]. We define the *Hodge numbers* of Z by

$$h_c^{k,p,q}(Z) = h^{p,q}(H_c^k(Z)) = \dim \mathrm{Gr}_F^p \mathrm{Gr}_{p+q}^W H_c^k(Z).$$

The Hodge–Deligne polynomial, or E -polynomial, is defined as

$$e(Z) = e(Z)(u, v) := \sum_{p,q,k} (-1)^k h_c^{k,p,q}(Z) u^p v^q.$$

The key property of Hodge–Deligne polynomials that permits their calculation is that they are additive for stratifications of Z . If Z is a complex algebraic variety and $Z = \bigsqcup_{i=1}^n Z_i$, where all Z_i are locally closed in Z , then

$$e(Z) = \sum_{i=1}^n e(Z_i).$$

Also, by [Logares et al. 2013, Remark 2.5], if $G \rightarrow X \rightarrow B$ is a principal fiber bundle with G a connected algebraic group, then $e(X) = e(G)e(B)$. In general we shall use this as $e(X/G) = e(X)/e(G)$ when $B = X/G$. In particular, if Z is a G -space, and there is a subspace $B \subset Z$ such that $B \times G \rightarrow Z$ is surjective and it is an H -homogeneous space for a connected subgroup $H \subset G$, then

$$(1) \quad e(Z) = e(B)e(G)/e(H).$$

Definition 3. Let X be a complex quasiprojective variety on which a finite group F acts. Then F also acts on the cohomology $H_c^*(X)$ respecting the mixed Hodge structure. So $[H_c^*(X)] \in R(F)$, the representation ring of F . The *equivariant Hodge–Deligne polynomial* is defined as

$$e_F(X) = \sum_{p,q,k} (-1)^k [H_c^{k,p,q}(X)] u^p v^q \in R(F)[u, v].$$

Note that the map $\dim : R(F) \rightarrow \mathbb{Z}$ gives $\dim(e_F(X)) = e(X)$.

For instance, for an action of \mathbb{Z}_2 , there are two irreducible representations T, N , where T is the trivial representation, and N is the nontrivial representation. Then $e_{\mathbb{Z}_2}(X) = aT + bN$. Clearly

$$e(X) = a + b, \quad e(X/\mathbb{Z}_2) = a.$$

In the notation of [Logares et al. 2013, Section 2], $a = e(X)^+$, $b = e(X)^-$. Note that if X, X' are spaces with \mathbb{Z}_2 -actions, then writing

$$e_{\mathbb{Z}_2}(X) = aT + bN \quad \text{and} \quad e_{\mathbb{Z}_2}(X') = a'T + b'N,$$

we have $e_{\mathbb{Z}_2}(X \times X') = (aa' + bb')T + (ab' + ba')N$ and so

$$(2) \quad e((X \times X')/\mathbb{Z}_2) = aa' + bb' = e(X)^+ e(X')^+ + e(X)^- e(X')^-.$$

When $h_c^{k,p,q} = 0$ for $p \neq q$, the polynomial $e(Z)$ depends only on the product uv . This will happen in all the cases that we shall investigate here. In this situation, it is conventional to use the variable $q = uv$. If this happens, we say that the variety is of *balanced type*. For instance, $e(\mathbb{C}^n) = q^n$.

3. *E*-polynomial of the $\mathrm{SL}(2, \mathbb{C})$ -character variety of free groups

Let F_r denote the free group on r generators. Then the space of representations of F_r in the group $\mathrm{SL}(2, \mathbb{C})$ is

$$\mathcal{R}_{r,2} = \mathrm{Hom}(F_r, \mathrm{SL}(2, \mathbb{C})) = \{(A_1, \dots, A_r) \mid A_i \in \mathrm{SL}(2, \mathbb{C})\} = \mathrm{SL}(2, \mathbb{C})^r.$$

The group $\mathrm{PGL}(2, \mathbb{C})$ acts on $\mathcal{R}_{r,2}$ by simultaneous conjugation of all matrices, and the character variety is defined as the GIT quotient

$$\mathcal{M}_{r,2} = \mathcal{R}_{r,2} // \mathrm{PGL}(2, \mathbb{C}).$$

We aim to compute the *E*-polynomial of $\mathcal{M}_{r,2}$ using the methods developed in [Logares et al. 2013] and to recover the results of [Cavazos and Lawton 2014]. We have the following sets:

- Reducible representations $\mathcal{R}_{r,2}^{\mathrm{red}} \subset \mathcal{R}_{r,2}$ and the corresponding set $\mathcal{M}_{r,2}^{\mathrm{red}} \subset \mathcal{M}_{r,2}$ of characters of reducible representations. A representation $\rho = (A_1, \dots, A_r)$ is reducible if and only if all A_i share at least one eigenvector.
- Irreducible representations $\mathcal{R}_{r,2}^{\mathrm{irr}} \subset \mathcal{R}_{r,2}$ and the corresponding set $\mathcal{M}_{r,2}^{\mathrm{irr}} \subset \mathcal{M}_{r,2}$ of characters of irreducible representations. This is the complement of $\mathcal{R}_{r,2}^{\mathrm{red}}$. It consists of the representations ρ such that $\mathrm{PGL}(2, \mathbb{C})$ acts freely on ρ , and the orbit $\mathrm{PGL}(2, \mathbb{C}) \cdot \rho$ is closed. Therefore $\mathcal{M}_{r,2}^{\mathrm{irr}} = \mathcal{R}_{r,2}^{\mathrm{irr}} / \mathrm{PGL}(2, \mathbb{C})$.

3.1. The reducible locus. Let us start by computing $e(\mathcal{M}_{r,2}^{\mathrm{red}})$. For a reducible representation, we have a basis of \mathbb{C}^2 in which

$$\rho = \left(\begin{pmatrix} \lambda_1 & * \\ 0 & \lambda_1^{-1} \end{pmatrix}, \begin{pmatrix} \lambda_2 & * \\ 0 & \lambda_2^{-1} \end{pmatrix}, \dots, \begin{pmatrix} \lambda_r & * \\ 0 & \lambda_r^{-1} \end{pmatrix} \right).$$

The associated point is determined by $(\lambda_1, \dots, \lambda_r) \in (\mathbb{C}^*)^r$, modulo $(\lambda_1, \dots, \lambda_r) \sim (\lambda_1^{-1}, \dots, \lambda_r^{-1})$. Note that the action of $\lambda \mapsto \lambda^{-1}$ on $X = \mathbb{C}^*$ has $e(X)^+ = q$ and $e(X)^- = -1$. Writing $X_i = \mathbb{C}^*$, $i = 1, \dots, r$, we have that

$$\begin{aligned} e(X_1 \times \dots \times X_r)^+ &= \sum_{\epsilon \in A} \prod_{i=1}^r e(X_i)^{\epsilon_i} \\ &= q^r + \binom{r}{2} q^{r-2} + \binom{r}{4} q^{r-4} + \dots + \binom{r}{2[r/2]} q^{r-2[r/2]} \\ &= \frac{1}{2}((q+1)^r + (q-1)^r), \end{aligned}$$

where $A = \{(\epsilon_1, \dots, \epsilon_r) \in (\pm 1)^r \mid \prod \epsilon_i = +1\}$. Also

$$\begin{aligned} e(X_1 \times \dots \times X_r)^- &= e(X_1 \times \dots \times X_r) - e(X_1 \times \dots \times X_r)^+ \\ &= (q-1)^r - \frac{1}{2}((q+1)^r + (q-1)^r) \\ &= \frac{1}{2}((q-1)^r - (q+1)^r). \end{aligned}$$

Also note that $e(\mathcal{M}_{r,2}^{\text{red}}) = e((X_1 \times \cdots \times X_r)/\mathbb{Z}_2) = e(X_1 \times \cdots \times X_r)^+$.

3.2. The reducible representations. Now we move to the computation of $e(\mathcal{R}_{r,2}^{\text{red}})$.

We stratify the space as $\mathcal{R}_{r,2}^{\text{red}} = R_0 \cup R_1 \cup R_2 \cup R_3$, where:

- R_0 consists of $(A_1, \dots, A_r) = (\pm \text{Id}, \dots, \pm \text{Id})$. So $e(R_0) = 2^r$.
- R_1 consists of

$$\rho \sim \left(\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_1^{-1} \end{pmatrix}, \begin{pmatrix} \lambda_2 & 0 \\ 0 & \lambda_2^{-1} \end{pmatrix}, \dots, \begin{pmatrix} \lambda_r & 0 \\ 0 & \lambda_r^{-1} \end{pmatrix} \right),$$

that is, abelian representations (all matrices are diagonalizable with respect to the same basis). Here $(\lambda_1, \dots, \lambda_r) \neq (\pm 1, \dots, \pm 1)$. Therefore this space is parametrized by

$$(\text{PGL}(2, \mathbb{C})/D \times ((\mathbb{C}^*)^r - \{(\pm 1, \dots, \pm 1)\})) / \mathbb{Z}_2,$$

where D is the space of diagonal matrices. We know that $e(\text{PGL}(2, \mathbb{C})/D)^+ = q^2$, $e(\text{PGL}(2, \mathbb{C})/D)^- = q$ by [Logares et al. 2013, Proposition 3.2]. For $B = (\mathbb{C}^*)^r - \{(\pm 1, \dots, \pm 1)\}$, we have $e(B)^+ = \frac{1}{2}((q+1)^r + (q-1)^r) - 2^r$ and $e(B)^- = \frac{1}{2}((q-1)^r - (q+1)^r)$, by our computation above. Therefore

$$\begin{aligned} e(R_1) &= e(\text{PGL}(2, \mathbb{C})/D)^+ e(B)^+ + e(\text{PGL}(2, \mathbb{C})/D)^- e(B)^- \\ &= q^2 \frac{1}{2}((q+1)^r + (q-1)^r - 2^r) + q \frac{1}{2}((q-1)^r - (q+1)^r) \\ &= \frac{1}{2}(q^2 - q)(q+1)^r + \frac{1}{2}(q^2 + q)(q-1)^r - q^2 2^r. \end{aligned}$$

- R_2 consists of

$$\rho \sim \left(\begin{pmatrix} \pm 1 & a_1 \\ 0 & \pm 1 \end{pmatrix}, \begin{pmatrix} \pm 1 & a_2 \\ 0 & \pm 1 \end{pmatrix}, \dots, \begin{pmatrix} \pm 1 & a_r \\ 0 & \pm 1 \end{pmatrix} \right),$$

where $(a_1, \dots, a_r) \in \mathbb{C}^r - \{0\}$. Let B_2 be the space of representations as above with respect to the canonical basis. Therefore, there is a canonical surjective map $B_2 \times \text{PGL}(2, \mathbb{C}) \rightarrow R_2$. The fibers of this map are given by $H_2 = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \right\} \cong \mathbb{C}^* \times \mathbb{C}$. That is, $H_2 \rightarrow B_2 \times \text{PGL}(2, \mathbb{C}) \rightarrow R_2$ is a fibration to which we apply Formula (1) to obtain

$$e(R_2) = \frac{e(B_2)e(\text{PGL}(2, \mathbb{C}))}{e(H_2)} = \frac{2^r(q^r - 1)(q^3 - q)}{q(q-1)} = 2^r(q^r - 1)(q+1).$$

- R_3 consists of

$$\rho \sim \left(\begin{pmatrix} \lambda_1 & b_1 \\ 0 & \lambda_1^{-1} \end{pmatrix}, \begin{pmatrix} \lambda_2 & b_2 \\ 0 & \lambda_2^{-1} \end{pmatrix}, \dots, \begin{pmatrix} \lambda_r & b_r \\ 0 & \lambda_r^{-1} \end{pmatrix} \right),$$

where $\lambda_i \in \mathbb{C}^*$, $(\lambda_1, \dots, \lambda_r) \neq (\pm 1, \dots, \pm 1)$. Here, $(b_1, \dots, b_r) \in \mathbb{C}^r$ and the upper diagonal matrices $\begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix}$ transform

$$(b_1, \dots, b_r) \mapsto (b_1 + y(\lambda_1 - \lambda_1^{-1}), \dots, b_r + y(\lambda_r - \lambda_r^{-1})).$$

As $(\lambda_1, \dots, \lambda_r) \neq (\pm 1, \dots, \pm 1)$, this action is nontrivial. Note that (b_1, \dots, b_r) does not live in the line spanned by $(\lambda_1 - \lambda_1^{-1}, \dots, \lambda_r - \lambda_r^{-1})$. There is a fibration $H_3 \rightarrow B_3 \times \mathrm{PGL}(2, \mathbb{C}) \rightarrow R_3$ where $H_3 = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \right\} \cong \mathbb{C}^* \times \mathbb{C}$. Thus

$$\begin{aligned} e(R_3) &= (q^r - q)((q - 1)^r - 2^r)e(\mathrm{PGL}(2, \mathbb{C}))/q(q - 1) \\ &= \frac{q^{r-1} - 1}{q - 1}((q - 1)^r - 2^r)(q^3 - q) \\ &= (q^{r-1} - 1)(q - 1)^{r-1}(q^3 - q) - 2^r \frac{q^{r-1} - 1}{q - 1}(q^3 - q). \end{aligned}$$

Now we add all the subsets together:

$$\begin{aligned} e(\mathcal{R}_{r,2}^{\mathrm{red}}) &= e(R_0) + e(R_1) + e(R_2) + e(R_3) \\ &= \frac{1}{2}(q^2 - q)(q + 1)^r + \frac{1}{2}(q^2 + q)(q - 1)^r \\ &\quad + (q^{r-1} - 1)(q - 1)^{r-1}(q^3 - q). \end{aligned}$$

3.3. The irreducible locus. Recall that $\mathcal{R}_{r,2}^{\mathrm{irr}} = \mathrm{SL}(2, \mathbb{C})^r - \mathcal{R}_{r,2}^{\mathrm{red}}$, so

$$\begin{aligned} e(\mathcal{R}_{r,2}^{\mathrm{irr}}) &= (q^3 - q)^r - \frac{1}{2}(q^2 - q)(q + 1)^r - \frac{1}{2}(q^2 + q)(q - 1)^r \\ &\quad - (q^{r-1} - 1)(q - 1)^{r-1}(q^3 - q), \end{aligned}$$

and

$$\begin{aligned} e(\mathcal{M}_{r,2}^{\mathrm{irr}}) &= \frac{e(\mathcal{R}_{r,2}^{\mathrm{irr}})}{q^3 - q} \\ &= (q^3 - q)^{r-1} - \frac{1}{2}(q + 1)^{r-1} - \frac{1}{2}(q - 1)^{r-1} - (q^{r-1} - 1)(q - 1)^{r-1}. \end{aligned}$$

Finally,

$$\begin{aligned} e(\mathcal{M}_{r,2}) &= e(\mathcal{M}_{r,2}^{\mathrm{irr}}) + e(\mathcal{M}_{r,2}^{\mathrm{red}}) = e(\mathcal{M}_{r,2}^{\mathrm{irr}}) + \frac{1}{2}((q + 1)^r + (q - 1)^r) \\ &= (q^3 - q)^{r-1} + \frac{1}{2}q(q + 1)^{r-1} + \frac{1}{2}q(q - 1)^{r-1} - q^{r-1}(q - 1)^{r-1}. \end{aligned}$$

This agrees with [Cavazos and Lawton 2014].

4. E-polynomial of the $\mathrm{PGL}(2, \mathbb{C})$ -character variety of free groups

Let us compute the E -polynomial of $\mathcal{M}(F_r, \mathrm{PGL}(2, \mathbb{C}))$. The space of representations will be denoted

$$\bar{\mathcal{R}}_{r,2} = \mathrm{Hom}(F_r, \mathrm{PGL}(2, \mathbb{C})) = \{(A_1, \dots, A_r) \mid A_i \in \mathrm{PGL}(2, \mathbb{C})\} = \mathrm{PGL}(2, \mathbb{C})^r.$$

Note that $\mathrm{PGL}(2, \mathbb{C}) = \mathrm{SL}(2, \mathbb{C})/\{\pm \mathrm{Id}\}$, so $\bar{\mathcal{R}}_{r,2} = \mathcal{R}_{r,2}/\{(\pm \mathrm{Id}, \dots, \pm \mathrm{Id})\}$. The character variety is

$$\bar{\mathcal{M}}_{r,2} = \bar{\mathcal{R}}_{r,2} // \mathrm{PGL}(2, \mathbb{C}).$$

We denote by $\bar{\mathcal{R}}_{r,2}^{\text{red}}$ and $\bar{\mathcal{R}}_{r,2}^{\text{irr}}$ the subsets of reducible and irreducible representations, respectively, of $\bar{\mathcal{R}}_{r,2}$. We denote by $\bar{\mathcal{M}}_{r,2}^{\text{red}}$ and $\bar{\mathcal{M}}_{r,2}^{\text{irr}}$ the corresponding spaces in $\bar{\mathcal{M}}_{r,2}$.

The reducible locus. We first compute $e(\bar{\mathcal{M}}_{r,2}^{\text{red}})$. A reducible representation in $\bar{\mathcal{M}}_{r,2}^{\text{red}}$ is determined by the eigenvalues $(\lambda_1, \dots, \lambda_r) \in (\mathbb{C}^*)^r$, modulo $\lambda_i \sim -\lambda_i$, $1 \leq i \leq r$, and $(\lambda_1, \dots, \lambda_r) \sim (\lambda_1^{-1}, \dots, \lambda_r^{-1})$. So it is determined by $(\lambda_1^2, \dots, \lambda_r^2) \in (\mathbb{C}^*)^r$, modulo $(\lambda_1^2, \dots, \lambda_r^2) \sim (\lambda_1^{-2}, \dots, \lambda_r^{-2})$. This space is isomorphic to the one in [Section 3.1](#), so $e(\bar{\mathcal{M}}_{r,2}^{\text{red}}) = \frac{1}{2}((q+1)^r + (q-1)^r)$.

The reducible representations. Now we compute $e(\bar{\mathcal{R}}_{r,2}^{\text{red}})$. We stratify it as

$$\bar{\mathcal{R}}_{r,2}^{\text{red}} = \bar{R}_0 \cup \bar{R}_1 \cup \bar{R}_2 \cup \bar{R}_3,$$

where:

- \bar{R}_0 consists of one point $(A_1, \dots, A_r) = (\text{Id}, \dots, \text{Id})$. So $e(R_0) = 1$.
- \bar{R}_1 consists of

$$\rho \sim \left(\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_1^{-1} \end{pmatrix}, \begin{pmatrix} \lambda_2 & 0 \\ 0 & \lambda_2^{-1} \end{pmatrix}, \dots, \begin{pmatrix} \lambda_r & 0 \\ 0 & \lambda_r^{-1} \end{pmatrix} \right),$$

where the eigenvalues are determined by $(\lambda_1^2, \dots, \lambda_r^2) \neq (1, \dots, 1)$. This space is parametrized by $(\text{PGL}(2, \mathbb{C})/D \times ((\mathbb{C}^*)^r - \{(1, \dots, 1)\})) / \mathbb{Z}_2$, where D is the space of diagonal matrices. Using that $e(\text{PGL}(2, \mathbb{C})/D)^+ = q^2$, $e(\text{PGL}(2, \mathbb{C})/D)^- = q$, and $e(B)^+ = \frac{1}{2}((q+1)^r + (q-1)^r) - 1$, $e(B)^- = \frac{1}{2}((q-1)^r - (q+1)^r)$, for $B = ((\mathbb{C}^*)^r - \{(1, \dots, 1)\})$, we have

$$\begin{aligned} e(\bar{R}_1) &= e(\text{PGL}(2, \mathbb{C})/D)^+ e(B)^+ + e(\text{PGL}(2, \mathbb{C})/D)^- e(B)^- \\ &= q^2 \frac{1}{2}((q+1)^r + (q-1)^r - 1) + q \frac{1}{2}((q-1)^r - (q+1)^r) \\ &= \frac{1}{2}(q^2 - q)(q+1)^r + \frac{1}{2}(q^2 + q)(q-1)^r - q^2. \end{aligned}$$

- \bar{R}_2 consists of

$$\rho \sim \left(\begin{pmatrix} 1 & a_1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & a_2 \\ 0 & 1 \end{pmatrix}, \dots, \begin{pmatrix} 1 & a_r \\ 0 & 1 \end{pmatrix} \right),$$

where $(a_1, \dots, a_r) \in \mathbb{C}^r - \{0\}$. Then

$$e(\bar{R}_2) = e(R_2)/2^r = (q^r - 1)(q+1).$$

- \bar{R}_3 consists of

$$\rho \sim \left(\begin{pmatrix} \lambda_1 & b_1 \\ 0 & \lambda_1^{-1} \end{pmatrix}, \begin{pmatrix} \lambda_2 & b_2 \\ 0 & \lambda_2^{-1} \end{pmatrix}, \dots, \begin{pmatrix} \lambda_r & b_r \\ 0 & \lambda_r^{-1} \end{pmatrix} \right),$$

where $\lambda_i \in \mathbb{C}^*$, $(\lambda_1^2, \dots, \lambda_r^2) \neq (1, \dots, 1)$. Here

$$(b_1, \dots, b_r) \in \mathbb{C}^r - \langle (\lambda_1 - \lambda_1^{-1}, \dots, \lambda_r - \lambda_r^{-1}) \rangle.$$

There is a fibration $H_3 \rightarrow B_3 \times \text{PGL}(2, \mathbb{C}) \rightarrow \bar{R}_3$ where B_3 parametrizes $(\lambda_1^2, \dots, \lambda_r^2)$ and (b_1, \dots, b_r) , and $H_3 = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \right\} \cong \mathbb{C}^* \times \mathbb{C}$. Then

$$\begin{aligned} e(\bar{R}_3) &= (q^r - q)((q - 1)^r - 1)e(\text{PGL}(2, \mathbb{C}))/q(q - 1) \\ &= \frac{q^{r-1} - 1}{q - 1}((q - 1)^r - 1)(q^3 - q). \end{aligned}$$

Now we add all subsets together to obtain:

$$\begin{aligned} e(\bar{\mathcal{R}}_{r,2}^{\text{red}}) &= e(\bar{R}_0) + e(\bar{R}_1) + e(\bar{R}_2) + e(\bar{R}_3) \\ &= \frac{1}{2}(q^2 - q)(q + 1)^r + \frac{1}{2}(q^2 + q)(q - 1)^r + (q^{r-1} - 1)(q - 1)^{r-1}(q^3 - q) \\ &= e(\mathcal{R}_{r,2}^{\text{red}}). \end{aligned}$$

The irreducible locus. Clearly, $e(\text{SL}(2, \mathbb{C})) = q^3 - q = e(\text{PGL}(2, \mathbb{C}))$ and $e(\bar{\mathcal{R}}_{r,2}^{\text{red}}) = e(\mathcal{R}_{r,2}^{\text{red}})$, we have that $e(\bar{\mathcal{R}}_{r,2}^{\text{irr}}) = e(\mathcal{R}_{r,2}^{\text{irr}})$. Therefore $e(\bar{\mathcal{M}}_{r,2}^{\text{irr}}) = e(\mathcal{M}_{r,2}^{\text{irr}})$. Finally, since $e(\bar{\mathcal{M}}_{r,2}^{\text{red}}) = e(\mathcal{M}_{r,2}^{\text{red}})$, we have that

$$\begin{aligned} e(\bar{\mathcal{M}}_{r,2}) &= e(\mathcal{M}_{r,2}) \\ &= (q^3 - q)^{r-1} + \frac{1}{2}q(q + 1)^{r-1} + \frac{1}{2}q(q - 1)^{r-1} - q^{r-1}(q - 1)^{r-1}. \end{aligned}$$

5. E-polynomial of the $\text{SL}(3, \mathbb{C})$ -character variety for F_1

Having given a new geometric derivation of the E -polynomial for $\mathcal{M}_{r,2}$ and $\bar{\mathcal{M}}_{r,2}$, in the next sections we work out the E -polynomial of $\mathcal{M}_{r,3}$ and $\bar{\mathcal{M}}_{r,3}$ in a similar fashion.

However, in this section we first address the $r = 1$ case. Although it is easy to see that $\mathcal{M}_{r,n} \cong \mathbb{C}^{n-1}$ via the coefficients of the characteristic polynomial, and hence $e(\mathcal{M}_{r,n}) = q^{n-1}$, this case will motivate the more complicated stratification, and the use of the *equivariant* E -polynomial, needed to compute the general E -polynomials for $\mathcal{M}_{r,3}$ and $\bar{\mathcal{M}}_{r,3}$ when $r \geq 2$.

We begin with the E -polynomials for $\text{GL}(3, \mathbb{C})$, $\text{SL}(3, \mathbb{C})$, and $\text{PGL}(3, \mathbb{C})$. Like in the previous sections, we then stratify $\text{Hom}(F_1, \text{SL}(3, \mathbb{C}))$ by orbit type and compute the E -polynomial for each strata.

Lemma 4. $e(\text{SL}(3, \mathbb{C})) = e(\text{PGL}(3, \mathbb{C})) = (q^3 - 1)(q^3 - q)q^2 = q^8 - q^6 - q^5 + q^3$.

Proof. Consider \mathbb{C}^n , and let V_k be the Stiefel manifold of k linearly independent vectors in \mathbb{C}^n . Then, there is a (Zariski locally trivial) fibration $\mathbb{C}^n - \mathbb{C}^{k-1} \rightarrow V_k \rightarrow V_{k-1}$. Therefore $e(V_k) = \prod_{i=0}^{k-1} (q^n - q^i)$. So $e(\text{GL}(n, \mathbb{C})) = e(V_n) = \prod_{i=0}^{n-1} (q^n - q^i)$.

Now there is a (Zariski locally trivial) fibration $\mathbb{C}^* \rightarrow \mathrm{GL}(n, \mathbb{C}) \rightarrow \mathrm{PGL}(n, \mathbb{C})$, hence $e(\mathrm{PGL}(n, \mathbb{C})) = e(\mathrm{GL}(n, \mathbb{C})) / (q - 1) = q^{n-1} \prod_{i=0}^{n-2} (q^n - q^i)$.

For $\mathrm{SL}(n, \mathbb{C})$, the choice of $(v_1, \dots, v_{n-1}) \in V_{n-1}$ determines an affine hyperplane

$$\{v \in \mathbb{C}^n \mid \det(v_1, \dots, v_{n-1}, v) = 1\}.$$

This gives a (Zariski locally trivial) affine bundle $\mathbb{C}^{n-1} \rightarrow \mathrm{SL}(n, \mathbb{C}) \rightarrow V_{n-1}$, and hence $e(\mathrm{SL}(n, \mathbb{C})) = q^{n-1} \prod_{i=0}^{n-2} (q^n - q^i)$. \square

Now let us consider the representations of F_1 to $\mathrm{SL}(3, \mathbb{C})$. This is equivalent to studying the conjugation action of $\mathrm{PGL}(3, \mathbb{C})$ on $X := \mathrm{SL}(3, \mathbb{C})$. For this action, there are 6 strata types. In the following list, we write down all 6 strata, but include the computation of their E -polynomials for only the first 5. This is because the computation is apparent from the geometric description of each stratum alone in those cases.

- X_0 is formed by matrices of type

$$\begin{pmatrix} \xi & 0 & 0 \\ 0 & \xi & 0 \\ 0 & 0 & \xi \end{pmatrix}.$$

Here $\xi^3 = 1$, so X_0 consists of 3 points and $e(X_0) = 3$.

- X_1 is formed by matrices of type

$$\begin{pmatrix} \xi & 0 & 0 \\ 0 & \xi & 1 \\ 0 & 0 & \xi \end{pmatrix}.$$

Here $\xi^3 = 1$, so ξ admits 3 values. The stabilizer of this matrix is

$$U_1 = \left\{ \begin{pmatrix} \mu^{-2} & 0 & b \\ a & \mu & c \\ 0 & 0 & \mu \end{pmatrix} \right\} \cong \mathbb{C}^* \times \mathbb{C}^3.$$

So

$$\begin{aligned} e(X_1) &= 3e(\mathrm{PGL}(3, \mathbb{C})/U_1) \\ &= 3(q^3 - 1)(q^3 - q)q^2/q^3(q - 1) = 3q^4 + 3q^3 - 3q - 3. \end{aligned}$$

- X_2 is formed by matrices of type

$$\begin{pmatrix} \xi & 1 & 0 \\ 0 & \xi & 1 \\ 0 & 0 & \xi \end{pmatrix}.$$

Here $\xi^3 = 1$, so ξ admits 3 values. The stabilizer of this matrix is

$$U_2 = \left\{ \begin{pmatrix} 1 & b & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \right\} \cong \mathbb{C}^2.$$

So

$$\begin{aligned} e(X_1) &= 3e(\mathrm{PGL}(3, \mathbb{C})/U_2) \\ &= 3(q^3 - 1)(q^3 - q)q^2/q^2 = 3q^6 - 3q^4 - 3q^3 + 3q. \end{aligned}$$

- X_3 is formed by matrices of type

$$\begin{pmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda^{-2} \end{pmatrix},$$

where $\lambda \in \mathbb{C}^* - \{\xi \mid \xi^3 = 1\}$. The stabilizer of this matrix is

$$U_3 = \left\{ \begin{pmatrix} A & 0 \\ 0 & (\det A)^{-1} \end{pmatrix} \mid A \in \mathrm{GL}(2, \mathbb{C}) \right\} \cong \mathrm{GL}(2, \mathbb{C}).$$

So

$$\begin{aligned} e(X_3) &= (q - 4)e(\mathrm{PGL}(3, \mathbb{C})/U_3) \\ &= (q - 4)(q^3 - 1)(q^3 - q)q^2/(q^2 - 1)(q^2 - q) = q^5 - 3q^4 - 3q^3 - 4q^2. \end{aligned}$$

- X_4 is formed by matrices of type

$$\begin{pmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda^{-2} \end{pmatrix},$$

where $\lambda \in \mathbb{C}^* - \{\xi \mid \xi^3 = 1\}$. The stabilizer of this matrix is

$$U_4 = \left\{ \begin{pmatrix} \mu & b & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \mu^{-2} \end{pmatrix} \right\} \cong \mathbb{C}^* \times \mathbb{C}.$$

So

$$\begin{aligned} e(X_4) &= (q - 4)e(\mathrm{PGL}(3, \mathbb{C})/U_4) \\ &= (q - 4)(q^3 - 1)(q^3 - q)q^2/q(q - 1) \\ &= q^7 - 3q^6 - 4q^5 - q^4 + 3q^3 + 4q^2. \end{aligned}$$

- X_5 is formed by matrices of type

$$\begin{pmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \gamma \end{pmatrix},$$

where $\lambda, \mu, \gamma \in \mathbb{C}^*$ are different and $\lambda\mu\gamma = 1$. The stabilizer is isomorphic to the diagonal matrices $D \cong \mathbb{C}^* \times \mathbb{C}^*$. The parameter space is

$$B = \{(\lambda, \mu) \in (\mathbb{C}^*)^2 \mid \lambda \neq \mu^{-2}, \mu \neq \lambda^{-2}, \mu \neq \lambda\}.$$

The map $\mathrm{PGL}(3, \mathbb{C})/D \times B \rightarrow X_5$ is a 6:1 cover. Moreover,

$$X_5 \cong (\mathrm{PGL}(3, \mathbb{C})/D \times B)/\Sigma_3,$$

where the symmetric group Σ_3 acts on $\mathrm{PGL}(3, \mathbb{C})$ by permuting the columns and acts on the triple $(\lambda, \mu, \gamma = \lambda^{-1}\mu^{-1})$ by permuting the entries.

We now compute $e(X_5)$ using the *equivariant E-polynomial*. Consider the finite group $F = \Sigma_3$. The representation ring $R(F)$ is generated by three irreducible representations:

- T is the (one-dimensional) trivial representation.
- S is the sign representation. This is one-dimensional and given by the sign map $\Sigma_3 \rightarrow \{\pm 1\} \subset \mathrm{GL}(1, \mathbb{C})$.
- V is the two-dimensional representation given as follows. Take $St = \mathbb{C}^3$ the standard 3-dimensional representation. This is generated by e_1, e_2, e_3 and Σ_3 acts by permuting the elements of the basis. Then $T = \langle e_1 + e_2 + e_3 \rangle$ and we can decompose $St = T \oplus V$.

The representation ring $R(\Sigma_3)$ has a multiplicative structure given by: $T \otimes T = T$, $T \otimes S = S$, $T \otimes V = V$, $S \otimes S = T$, $S \otimes V = V$, $V \otimes V = T \oplus S \oplus V$.

Lemma 5.

$$\begin{aligned} e_{\Sigma_3}(B) &= (q^2 - q + 1)T + S - 2(q - 2)V. \\ e_{\Sigma_3}(\mathrm{PGL}(3, \mathbb{C})/D) &= q^6T + q^3S + (q^5 + q^4)V. \end{aligned}$$

Proof. Write $e_{\Sigma_3}(X) = aT + bS + cV$, for a quasiprojective variety X with a Σ_3 -action. Then $a = e(X/\Sigma_3)$. If we consider the cycle $(1, 2)$ and the subgroup $H = \langle (1, 2) \rangle$, there is a map $R(F) \rightarrow R(H)$ which sends $T \mapsto T$, $S \mapsto N$ and $V \mapsto T + N$. Then $e_H(X) = aT + bN + c(T + N) = (a + c)T + (b + c)N$. Therefore, $a + c = e(X/H)$. As $e(X) = a + b + 2c$, we can compute a, b, c by knowing these E -polynomials.

For $B = \{(\lambda, \mu) \in (\mathbb{C}^*)^2 \mid \lambda \neq \mu^{-2}, \mu \neq \lambda^{-2}, \mu \neq \lambda\}$, the three curves $\lambda = \mu^{-2}$, $\mu = \lambda^{-2}$, $\mu = \lambda$ intersect at the three points $\{(\xi, \xi) \mid \xi^3 = 1\}$. Hence $e(B) = (q - 1)^2 - 3(q - 4) - 3 = q^2 - 5q + 10$.

Now Σ_3 acts on (λ, μ, γ) and the quotient space is parametrized by $s = \lambda + \mu + \gamma$, $t = \lambda\mu + \lambda\gamma + \mu\gamma$ and $p = \lambda\mu\gamma = 1$, that is, by $(s, t) \in \mathbb{C}^2$. We have to remove the cases $s = \lambda + \lambda^{-2} + \lambda$, $t = \lambda^{-1} + \lambda^2 + \lambda^{-1}$. This defines a rational curve in \mathbb{C}^2 . It has two points at infinity. The map $\lambda \mapsto (2\lambda + \lambda^{-2}, 2\lambda^{-1} + \lambda^2)$ is an embedding. Therefore $e(B/\Sigma_3) = q^2 - (q - 1) = q^2 - q + 1$.

The action by H permutes (λ, μ) , hence the quotient is parametrized by $s' = \lambda + \mu$, $p' = \lambda\mu \neq 0$. We have to remove the cases $s' = \lambda + \lambda^{-2}$, $p' = \lambda^{-1}$, that is, $s' = (p')^{-1} + (p')^2$; and $s' = 2\lambda$, $p' = \lambda^2$, i.e., $4p' = (s')^2$. They intersect at three points. Then $e(B/H) = q(q-1) - 2(q-1) + 3 = q^2 - 3q + 5$.

Thus

$$e_{\Sigma_3}(B) = (q^2 - q + 1)T + S - 2(q - 2)V.$$

For $C = \text{PGL}(3, \mathbb{C})/D$, the space C consists of points in $(\mathbb{P}^2)^3 - \Delta$, where Δ is the diagonal (triples of coplanar points). Certainly, a matrix in $\text{GL}(3, \mathbb{C})$ can be written as (v_1, v_2, v_3) , where v_1, v_2, v_3 are linearly independent vectors. Taking a quotient by the diagonal matrices corresponds to the vectors up to a scalar: $[v_1], [v_2], [v_3]$. Therefore, $e(C) = (q^3 - 1)(q^3 - q)q^2 / (q - 1)^2 = q^6 + 2q^5 + 2q^4 + q^3$.

The group Σ_3 acts by permuting the vectors, so $C/\Sigma_3 = \text{Sym}^3 \mathbb{P}^2 - \bar{\Delta}$, where $\bar{\Delta}$ consists of linearly dependent triples $([v_1], [v_2], [v_3])$. If they are equal, the set has $e(\mathbb{P}^2) = q^2 + q + 1$. If they are collinear, there is a fibration with fiber $\text{Sym}^3(\mathbb{P}^1) - \Delta$ and base $(\mathbb{P}^2)^\vee$. This has E -polynomial $(1 + q + q^2 + q^3 - 1 - q)(1 + q + q^2) = q^5 + 2q^4 + 2q^3 + q^2$. Also $e(\text{Sym}^3 \mathbb{P}^2) = q^6 + q^5 + 2q^4 + 2q^3 + 2q^2 + q + 1$. Therefore

$$e(C/\Sigma_3) = q^6 + q^5 + 2q^4 + 2q^3 + 2q^2 + q + 1 - (q^5 + 2q^4 + 2q^3 + q^2 + q^2 + q + 1) = q^6.$$

The group H acts by permuting the first two vectors, so $C/H = \text{Sym}^2 \mathbb{P}^2 \times \mathbb{P}^2 - \bar{\Delta}'$, where $\bar{\Delta}'$ consists of linearly dependent triples $([v_1], [v_2], [v_3])$. If $[v_1] = [v_2]$, we have the E -polynomial $(q^2 + q + 1)(q^2 + q + 1) = q^4 + 2q^3 + 3q^2 + 2q + 1$. If $[v_1] \neq [v_2]$, they lie in $\text{Sym}^2 \mathbb{P}^2 - \Delta$ and we have the E -polynomial

$$(q^4 + q^3 + 2q^2 + q + 1 - (q^2 + q + 1))(q + 1) = q^5 + 2q^4 + 2q^3 + q^2.$$

Also $e(\text{Sym}^2 \mathbb{P}^2 \times \mathbb{P}^2) = (q^4 + q^3 + 2q^2 + q + 1)(q^2 + q + 1)$, so

$$\begin{aligned} e(C/H) &= q^6 + 2q^5 + 4q^4 + 4q^3 + 4q^2 + 2q + 1 \\ &\quad - (q^5 + 2q^4 + 2q^3 + q^2 + q^4 + 2q^3 + 3q^2 + 2q + 1) \\ &= q^6 + q^5 + q^4. \end{aligned}$$

This produces the polynomial

$$e_{\Sigma_3}(C) = q^6 T + q^3 S + (q^5 + q^4) V. \quad \square$$

Remark 6. If we consider $B' = \{(\lambda, \mu) \in (\mathbb{C}^*)^2\}$, then the proof of [Lemma 5](#) says that $e_{\Sigma_3}(B') = q^2 T + S - q V$.

Now suppose $e_{\Sigma_3}(X) = aT + bS + cV$ and $e_{\Sigma_3}(X') = a'T + b'S + c'V$. Then $e_{\Sigma_3}(X \times X') = (aa' + bb' + cc')T + (ab' + ba' + cc')S + (ac' + ca' + bc' + cb' + cc')V$,

and hence

$$(3) \quad e((X \times X')/\Sigma_3) = aa' + bb' + cc'.$$

We finally obtain the E -polynomial for the sixth strata X_5 :

$$\begin{aligned} e(X_5) &= e((B \times C)/\Sigma_3) \\ &= (q^2 - q + 1)q^6 + q^3 - 2(q - 2)(q^5 + q^4) \\ &= q^8 - q^7 - q^6 + 2q^5 + 4q^4 + q^3. \end{aligned}$$

Now we add the strata together:

$$e(X_0) + e(X_1) + e(X_2) + e(X_3) + e(X_4) + e(X_5) = q^8 - q^6 - q^5 + q^3 = e(\mathrm{SL}(3, \mathbb{C})),$$

as expected.

Remark 7. All elements of $X = \mathrm{SL}(3, \mathbb{C})$ are reducible. The semisimple ones are given by diagonal matrices with entries λ, μ, γ with $\lambda\mu\gamma = 1$. So they are parametrized by $s = \lambda + \mu + \gamma$, $t = \lambda\mu + \lambda\gamma + \mu\gamma = \lambda^{-1} + \gamma^{-1} + \mu^{-1}$, for $(s, t) \in \mathbb{C}^2$. Hence $e(\mathcal{M}_{1,3}) = q^2$, as noted at the beginning of this section.

6. E -polynomials of character varieties for F_r , $r > 1$, and $\mathrm{SL}(3, \mathbb{C})$

In this section we prove (most of) our main theorem ([Theorem 1](#)) by computing the E -polynomial for $\mathcal{M}_{r,3}$; the rest of [Theorem 1](#) is proved in [Section 7](#). The computation is similar to the computation in [Section 3](#) except the stratification is more complicated and the *equivariant* E -polynomial is needed, as was demonstrated in [Section 5](#).

Indeed, we want to study the space of representations

$$\begin{aligned} \mathcal{R}_{r,3} &= \mathrm{Hom}(F_r, \mathrm{SL}(3, \mathbb{C})) = \{\rho : F_r \rightarrow \mathrm{SL}(3, \mathbb{C})\} \\ &= \{(A_1, \dots, A_r) \mid A_i \in \mathrm{SL}(3, \mathbb{C})\} = \mathrm{SL}(3, \mathbb{C})^r \end{aligned}$$

and the corresponding character variety

$$\mathcal{M}_{r,3} = \mathrm{Hom}(F_r, \mathrm{SL}(3, \mathbb{C})) // \mathrm{PGL}(3, \mathbb{C}).$$

Much of the algebraic structure of $\mathcal{M}_{r,3}$ has been worked out in [[Lawton 2007](#); [2008](#); [2010](#)].

Let us start by computing the E -polynomial of the space of reducible representations $\mathcal{R}_{r,3}^{\mathrm{red}} \subset \mathrm{Hom}(F_r, \mathrm{SL}(3, \mathbb{C}))$.

We now list the stratification and the computation of the E -polynomial for each stratum for $\mathcal{R}_{r,3}^{\mathrm{red}}$.

(i) $R_0 = R_{01} \cup R_{02}$. R_{01} is formed by representations $\rho = (A_1, \dots, A_r)$ which have a common eigenvector and such that the quotient representation is irreducible, that is,

$$A_i = \begin{pmatrix} \lambda_i^{-2} & b_i & c_i \\ 0 & & \lambda_i B_i \\ 0 & & \end{pmatrix},$$

where $(B_1, \dots, B_r) \in \mathcal{R}_{2,r}^{\text{irr}}$. Let B_{01} be the space of representations of such form with respect to the standard basis. The stabilizer of B_{01} (i.e., the set $H_{01} \subset \text{PGL}(3, \mathbb{C})$ sending B_{01} to itself) is

$$H_{01} = \left\{ \begin{pmatrix} (\det B)^{-1} & a & b \\ 0 & & B \\ 0 & & \end{pmatrix} \right\} \cong \text{GL}(2, \mathbb{C}) \times \mathbb{C}^2.$$

This means that there is a fibration $H_{01} \rightarrow B_{01} \times \text{PGL}(3, \mathbb{C}) \rightarrow R_{01}$. Hence

$$e(R_{01}) = (q-1)^r q^{2r} e(\mathcal{R}_{2,r}^{\text{irr}}) \frac{e(\text{PGL}(3, \mathbb{C}))}{q^2 e(\text{GL}(2, \mathbb{C}))}.$$

R_{02} is formed by representations $\rho = (A_1, \dots, A_r)$ which have a common two-dimensional space and upon which it acts irreducibly, that is,

$$A_i = \begin{pmatrix} \lambda_i B_i & 0 \\ b_i & c_i & \lambda_i^{-2} \end{pmatrix},$$

where $(B_1, \dots, B_r) \in \mathcal{R}_{2,r}^{\text{irr}}$. The stabilizer is now

$$H_{02} = \left\{ \begin{pmatrix} B & 0 \\ a & b & (\det B)^{-1} \end{pmatrix} \right\} \cong \text{GL}(2, \mathbb{C}) \times \mathbb{C}^2.$$

Hence

$$e(R_{02}) = (q-1)^r q^{2r} e(\mathcal{R}_{2,r}^{\text{irr}}) \frac{e(\text{PGL}(3, \mathbb{C}))}{q^2 e(\text{GL}(2, \mathbb{C}))}.$$

The intersection $R_{01} \cap R_{02}$ consists of those representations with $b_i = c_i = 0$, which have stabilizer $\text{GL}(2, \mathbb{C})$, hence

$$e(R_{01} \cap R_{02}) = (q-1)^r e(\mathcal{R}_{2,r}^{\text{irr}}) \frac{e(\text{PGL}(3, \mathbb{C}))}{e(\text{GL}(2, \mathbb{C}))}.$$

Finally $e(R_0) = e(R_{01}) + e(R_{02}) - e(R_{01} \cap R_{02}) = 2e(R_{01}) - e(R_{01} \cap R_{02})$. Note that the remaining representations have a full invariant flag.

(ii) R_1 is formed by representations $\rho = (A_1, \dots, A_r)$ such that the eigenvalues of all A_i are equal (and hence cubic roots of unity). This consists of the following substrata:

- R_{11} consisting of matrices

$$A_i = \begin{pmatrix} \xi_i & 0 & 0 \\ 0 & \xi_i & 0 \\ 0 & 0 & \xi_i \end{pmatrix},$$

where $\xi_i^3 = 1$. So $e(R_{11}) = 3^r$.

- R_{12} formed by matrices of type

$$A_i = \begin{pmatrix} \xi_i & 0 & 0 \\ 0 & \xi_i & a_i \\ 0 & 0 & \xi_i \end{pmatrix},$$

with $\xi_i^3 = 1$ and $(a_1, \dots, a_r) \neq 0$. Then the stabilizer is

$$H_{12} = \left\{ \begin{pmatrix} \mu^{-1}\gamma^{-1} & 0 & b \\ a & \mu & c \\ 0 & 0 & \gamma \end{pmatrix} \right\} \cong (\mathbb{C}^*)^2 \times \mathbb{C}^3.$$

So

$$e(R_{12}) = 3^r (q^r - 1) \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q-1)^2 q^3}.$$

- R_{13} formed by matrices of type

$$A_i = \begin{pmatrix} \xi_i & 0 & a_i \\ 0 & \xi_i & b_i \\ 0 & 0 & \xi_i \end{pmatrix},$$

with $\xi_i^3 = 1$ and $(a_1, \dots, a_r), (b_1, \dots, b_r)$ linearly independent. Note that when they are linearly dependent, one may arrange a basis so that it belongs to the stratum R_{12} . Then the stabilizer is

$$H_{13} = \left\{ \begin{pmatrix} A & b \\ 0 & 0 & (\det A)^{-1} \end{pmatrix} \right\} \cong \mathrm{GL}(2, \mathbb{C}) \times \mathbb{C}^2.$$

Hence

$$e(R_{13}) = 3^r (q^r - 1)(q^r - q) \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q^2 - 1)(q^2 - q)q^2}.$$

- R_{14} formed by matrices of type

$$A_i = \begin{pmatrix} \xi_i & a_i & b_i \\ 0 & \xi_i & 0 \\ 0 & 0 & \xi_i \end{pmatrix},$$

with $\xi_i^3 = 1$ and $(a_1, \dots, a_r), (b_1, \dots, b_r)$ linearly independent. Note again that when they are linearly dependent, one may arrange a basis so that it belongs to the stratum R_{12} . Then the stabilizer is

$$H_{14} = \left\{ \begin{pmatrix} (\det A)^{-1} & b & c \\ 0 & & A \end{pmatrix} \right\} \cong \text{GL}(2, \mathbb{C}) \times \mathbb{C}^2.$$

Hence

$$e(R_{14}) = 3^r (q^r - 1)(q^r - q) \frac{e(\text{PGL}(3, \mathbb{C}))}{(q^2 - 1)(q^2 - q)q^2}.$$

- R_{15} formed by matrices of type

$$A_i = \begin{pmatrix} \xi_i & a_i & b_i \\ 0 & \xi_i & c_i \\ 0 & 0 & \xi_i \end{pmatrix},$$

with $\xi_i^3 = 1$ and $(a_1, \dots, a_r), (c_1, \dots, c_r)$ are both nonzero (if one of them is zero, then we are back in the case R_{13}). Then the stabilizer is

$$H_{15} = \left\{ \begin{pmatrix} a & b & c \\ 0 & d & e \\ 0 & 0 & f \end{pmatrix} \right\}.$$

Hence

$$e(R_{14}) = 3^r (q^r - 1)^2 q^r \frac{e(\text{PGL}(3, \mathbb{C}))}{(q - 1)^2 q^3}.$$

All together, we have

$$e(R_1) = 3^r (1 + (1 + q + q^2)(q^{3r+1} + q^{3r} - 2q^{2r+1} + q - 1)).$$

(iii) R_2 is formed by matrices with eigenvalues $(\lambda_i, \lambda_i, \mu_i)$. Let $\lambda = (\lambda_1, \dots, \lambda_r)$, $\mu = (\mu_1, \dots, \mu_r)$, with $\lambda - \mu \neq \mathbf{0}$. Note that $\mu_i = \lambda_i^{-2}$, so the parameter space has E -polynomial $(q - 1)^r - 3^r$. We have the following substrata:

- R_{21} consists of representations of type

$$A_i = \begin{pmatrix} \lambda_i & 0 & 0 \\ 0 & \lambda_i & 0 \\ 0 & 0 & \mu_i \end{pmatrix}.$$

The stabilizer is $P(\mathrm{GL}(2, \mathbb{C}) \times \mathbb{C}^*) \cong \mathrm{GL}(2, \mathbb{C})$, so

$$e(R_{21}) = ((q-1)^r - 3^r) \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q^2-1)(q^2-q)}.$$

- R_{22} consists of representations of type

$$A_i = \begin{pmatrix} \lambda_i & a_i & b_i \\ 0 & \lambda_i & c_i \\ 0 & 0 & \mu_i \end{pmatrix},$$

with $\mathbf{a} \neq \mathbf{0}$, $\mathbf{b} = (b_1, \dots, b_r)$, $\mathbf{c} = (c_1, \dots, c_r) \in \mathbb{C}^r$. The stabilizer is

$$H_{22} = \left\{ \begin{pmatrix} a & b & c \\ 0 & d & e \\ 0 & 0 & f \end{pmatrix} \right\}.$$

Hence

$$e(R_{22}) = ((q-1)^r - 3^r)(q^r - 1)q^{2r} \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q-1)^2q^3}.$$

- R_{23} consists of representations of type

$$A_i = \begin{pmatrix} \lambda_i & 0 & 0 \\ 0 & \lambda_i & a_i \\ 0 & 0 & \mu_i \end{pmatrix},$$

with $\mathbf{a} \notin \langle \boldsymbol{\lambda} - \boldsymbol{\mu} \rangle$. If $\mathbf{a} = x(\boldsymbol{\lambda} - \boldsymbol{\mu})$, $x \in \mathbb{C}$, then we can arrange a basis so that this belongs to the stratum R_{21} . The stabilizer is

$$H_{23} = \left\{ \begin{pmatrix} a & 0 & 0 \\ b & c & d \\ 0 & 0 & e \end{pmatrix} \right\}.$$

So

$$e(R_{23}) = ((q-1)^r - 3^r)(q^r - q) \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q-1)^2q^2}.$$

- R_{24} consists of representations of type

$$A_i = \begin{pmatrix} \lambda_i & 0 & a_i \\ 0 & \lambda_i & b_i \\ 0 & 0 & \mu_i \end{pmatrix},$$

with \mathbf{a} , \mathbf{b} and $\boldsymbol{\lambda} - \boldsymbol{\mu}$ linearly independent (if they were linearly dependent, one can arrange a basis so that we go back to case R_{23}). The stabilizer is

$$H_{24} = \left\{ \begin{pmatrix} A & \mathbf{a} \\ 0 & 0 & (\det A)^{-1} \end{pmatrix} \right\}.$$

Hence

$$e(R_{24}) = ((q-1)^r - 3^r)(q^r - q)(q^r - q^2) \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q^2 - 1)(q^2 - q)q^2}.$$

- R_{25} consists of representations of type

$$A_i = \begin{pmatrix} \mu_i & a_i & b_i \\ 0 & \lambda_i & c_i \\ 0 & 0 & \lambda_i \end{pmatrix},$$

with $\mathbf{a} \notin \langle \boldsymbol{\lambda} - \boldsymbol{\mu} \rangle$, $\mathbf{c} \neq \mathbf{0}$. The stabilizer is

$$H_{25} = \left\{ \begin{pmatrix} a & b & c \\ 0 & d & e \\ 0 & 0 & f \end{pmatrix} \right\}.$$

Hence

$$e(R_{25}) = ((q-1)^r - 3^r)(q^r - 1)(q^r - q)q^r \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q-1)^2 q^3}.$$

- R_{26} consists of representations of type

$$A_i = \begin{pmatrix} \mu_i & 0 & b_i \\ 0 & \lambda_i & c_i \\ 0 & 0 & \lambda_i \end{pmatrix},$$

with $\mathbf{b} \notin \langle \boldsymbol{\lambda} - \boldsymbol{\mu} \rangle$, $\mathbf{c} \neq \mathbf{0}$. (If \mathbf{b} is a multiple of $\boldsymbol{\lambda} - \boldsymbol{\mu}$, then we can arrange with a suitable basis that $\mathbf{b} = \mathbf{0}$, and this belongs to the substrata R_{22}). The stabilizer is

$$H_{26} = \left\{ \begin{pmatrix} a & 0 & b \\ 0 & c & d \\ 0 & 0 & e \end{pmatrix} \right\}.$$

Hence

$$e(R_{26}) = ((q-1)^r - 3^r)(q^r - 1)(q^r - q) \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q-1)^2 q^2}.$$

- R_{27} consists of representations of type

$$A_i = \begin{pmatrix} \mu_i & a_i & 0 \\ 0 & \lambda_i & 0 \\ 0 & 0 & \lambda_i \end{pmatrix},$$

with $a \notin \langle \lambda - \mu \rangle$. The stabilizer is

$$H_{27} = \left\{ \begin{pmatrix} a & b & 0 \\ 0 & c & 0 \\ 0 & d & e \end{pmatrix} \right\}.$$

Hence

$$e(R_{27}) = ((q-1)^r - 3^r)(q^r - q) \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q-1)^2 q^2}.$$

- R_{28} consists of representations of type

$$A_i = \begin{pmatrix} \mu_i & a_i & b_i \\ 0 & \lambda_i & 0 \\ 0 & 0 & \lambda_i \end{pmatrix},$$

with $a, b, \lambda - \mu$ linearly independent (otherwise we can reduce to the case R_{27}). The stabilizer is

$$H_{28} = \left\{ \begin{pmatrix} (\det A)^{-1} & b & c \\ 0 & & A \\ 0 & & \end{pmatrix} \right\}.$$

Hence

$$e(R_{28}) = ((q-1)^r - 3^r)(q^r - q)(q^r - q^2) \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q^2 - q)(q^2 - 1)q^2}.$$

- R_{29} consists of representations of type

$$A_i = \begin{pmatrix} \lambda_i & a_i & b_i \\ 0 & \mu_i & c_i \\ 0 & 0 & \lambda_i \end{pmatrix},$$

with $a, c \notin \langle \lambda - \mu \rangle$. The stabilizer is

$$H_{29} = \left\{ \begin{pmatrix} a & b & c \\ 0 & d & e \\ 0 & 0 & f \end{pmatrix} \right\}.$$

Hence

$$e(R_{29}) = ((q-1)^r - 3^r)(q^r - q)^2 q^r \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q-1)^2 q^3}.$$

All together, we have

$$e(R_2) = ((q-1)^r - 3^r)(q^2 + q + 1)(3q^{3r+1} + 3q^{3r} - 2q^{2r+2} - 4q^{2r+1} + q^3).$$

(iv) R_3 is formed by matrices with eigenvalues $(\lambda_i, \mu_i, \gamma_i)$ where $\lambda = (\lambda_1, \dots, \lambda_r)$, $\mu = (\mu_1, \dots, \mu_r)$, and $\gamma = (\gamma_1, \dots, \gamma_r)$ are distinct. Note that $\lambda_i \mu_i \gamma_i = 1$ for all $1 \leq i \leq r$. The base B_r parametrizing (λ, μ, γ) has E -polynomial $e(B_r) = (q-1)^{2r} - 3(q-1)^r + 2 \cdot 3^r$.

- R_{31} consists of representations of type

$$A_i = \begin{pmatrix} \lambda_i & 0 & 0 \\ 0 & \mu_i & 0 \\ 0 & 0 & \gamma_i \end{pmatrix}.$$

Then the stabilizer is $D \times \Sigma_3$, where D is the diagonal matrices. So we have to compute the E -polynomial of the quotient $R_{31} = (\mathrm{PGL}(3, \mathbb{C})/D \times B_r)/\Sigma_3$. We start by computing $e_{\Sigma_3}(B_r)$. Let $B'_r = \{(\lambda, \mu, \gamma) \in (\mathbb{C}^*)^{3r} \mid \lambda \mu \gamma = (1, \dots, 1)\}$. This is $B'_r = (B')^r$, in the notation of [Remark 6](#). Then

$$e_{\Sigma_3}(B'_r) = e_{\Sigma_3}(B')^r = (q^2 T + S - qV)^r.$$

Using the properties $T \otimes T = T$, $T \otimes S = S$, $T \otimes V = V$, $S \otimes S = T$, $S \otimes V = V$, $V \otimes V = T \oplus S \oplus V$, it is easy to see that $V^b = a_b V + a_{b-1}(T + S)$, where $a_b = a_{b-1} + 2a_{b-2}$, with $a_0 = 0$, $a_1 = 1$. This recurrence solves as $a_b = (2^b - (-1)^b)/3$. Therefore

$$\begin{aligned} (4) \quad e_{\Sigma_3}(B'_r) &= (q^2 T + S - qV)^r \\ &= \sum \frac{r!}{(r-a-b)!a!b!} q^{2(r-a-b)} S^a (-q)^b V^b \\ &= \sum \frac{r!}{(r-a)!a!} q^{2(r-a)} S^a \\ &\quad + \sum_{b>0} \frac{r!}{(r-a-b)!a!b!} q^{2(r-a-b)} S^a (-q)^b V^b \\ &= \frac{1}{2}((q^2 + 1)^r + (q^2 - 1)^r)T + \frac{1}{2}((q^2 + 1)^r - (q^2 - 1)^r)S \\ &\quad + \sum_{b>0} \frac{r!}{(r-a-b)!a!b!} q^{2(r-a-b)} S^a (-q)^b \\ &\quad \quad \times \left(\frac{1}{3}(2^b - (-1)^b)V + \frac{1}{3}(2^{b-1} - (-1)^{b-1})(T + S) \right) \\ &= \frac{1}{2}((q^2 + 1)^r + (q^2 - 1)^r)T + \frac{1}{2}((q^2 + 1)^r - (q^2 - 1)^r)S \\ &\quad + \frac{1}{3}((q^2 - 2q + 1)^r - (q^2 + q + 1)^r)V \\ &\quad + \frac{1}{3}\left(\frac{1}{2}(q^2 - 2q + 1)^r + (q^2 + q + 1)^r - \frac{3}{2}(q^2 + 1)^r\right)(T + S) \end{aligned}$$

$$\begin{aligned}
&= \left(\frac{1}{2}(q^2 - 1)^r + \frac{1}{6}(q - 1)^{2r} + \frac{1}{3}(q^2 + q + 1)^r\right)T \\
&\quad + \left(-\frac{1}{2}(q^2 - 1)^r + \frac{1}{6}(q - 1)^{2r} + \frac{1}{3}(q^2 + q + 1)^r\right)S \\
&\quad + \frac{1}{3}\left((q - 1)^{2r} - (q^2 + q + 1)^r\right)V.
\end{aligned}$$

Now we have to look at the part that we removed:

$$C_r = \{(\lambda, \lambda, \lambda^{-2}) \mid \lambda \in (\mathbb{C}^*)^r\} \cup \{(\lambda, \lambda^{-2}, \lambda) \mid \lambda \in (\mathbb{C}^*)^r\} \cup \{(\lambda^{-2}, \lambda, \lambda) \mid \lambda \in (\mathbb{C}^*)^r\}.$$

Then $e(C_r) = 3(q - 1)^r - 2 \cdot 3^r$. The quotient $C_r/\Sigma_3 \cong (\mathbb{C}^*)^r$, so $e(C_r/\Sigma_3) = (q - 1)^r$. And for $H = \langle(1, 2)\rangle$ we have $C_r/H \cong \{(\lambda, \lambda, \lambda^{-2}) \mid \lambda \in (\mathbb{C}^*)^r\} \cup \{(\lambda, \lambda^{-2}, \lambda) \mid \lambda \in (\mathbb{C}^*)^r\}$, so $e(C_r/H) = 2(q - 1)^r - 3^r$. Hence,

$$e_{\Sigma_3}(C_r) = (q - 1)^r T + ((q - 1)^r - 3^r)V.$$

For $B_r = B'_r - C_r$, we have

$$\begin{aligned}
(5) \quad e_{\Sigma_3}(B_r) &= \left(\frac{1}{2}(q^2 - 1)^r + \frac{1}{6}(q - 1)^{2r} + \frac{1}{3}(q^2 + q + 1)^r - (q - 1)^r\right)T \\
&\quad + \left(-\frac{1}{2}(q^2 - 1)^r + \frac{1}{6}(q - 1)^{2r} + \frac{1}{3}(q^2 + q + 1)^r\right)S \\
&\quad + \left(\frac{1}{3}(q - 1)^{2r} - \frac{1}{3}(q^2 + q + 1)^r - (q - 1)^r + 3^r\right)V.
\end{aligned}$$

Hence Formula (3) and Lemma 5 imply

$$\begin{aligned}
e(R_{31}) &= aa' + bb' + cc' \\
&= \left(\frac{1}{2}(q^2 - 1)^r + \frac{1}{6}(q - 1)^{2r} + \frac{1}{3}(q^2 + q + 1)^r - (q - 1)^r\right)q^6 \\
&\quad + \left(-\frac{1}{2}(q^2 - 1)^r + \frac{1}{6}(q - 1)^{2r} + \frac{1}{3}(q^2 + q + 1)^r\right)q^3 \\
&\quad + \left(\frac{1}{3}(q - 1)^{2r} - \frac{1}{3}(q^2 + q + 1)^r - (q - 1)^r + 3^r\right)(q^5 + q^4).
\end{aligned}$$

- R_{32} consists of representations of type

$$A_i = \begin{pmatrix} \lambda_i & 0 & 0 \\ 0 & \mu_i & a_i \\ 0 & 0 & \gamma_i \end{pmatrix},$$

with $a \notin \langle \mu - \gamma \rangle$. The stabilizer is

$$H_{32} = \left\{ \begin{pmatrix} a & 0 & 0 \\ 0 & b & c \\ 0 & 0 & d \end{pmatrix} \right\}.$$

Hence,

$$e(R_{32}) = ((q - 1)^{2r} - 3(q - 1)^r + 2 \cdot 3^r)(q^r - q) \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q - 1)^2 q}.$$

- R_{33} consists of representations of type

$$A_i = \begin{pmatrix} \lambda_i & 0 & a_i \\ 0 & \mu_i & b_i \\ 0 & 0 & \gamma_i \end{pmatrix},$$

with $\mathbf{a} \notin \langle \boldsymbol{\lambda} - \boldsymbol{\gamma} \rangle$, $\mathbf{b} \notin \langle \boldsymbol{\mu} - \boldsymbol{\gamma} \rangle$. The stabilizer is

$$H_{33} = \left\{ \begin{pmatrix} a & 0 & b \\ 0 & c & d \\ 0 & 0 & e \end{pmatrix} \right\} \times \mathbb{Z}_2,$$

where \mathbb{Z}_2 permutes the eigenvalues λ_i, μ_i . Therefore,

$$R_{33} = (B_r \times (\mathbb{C}^r - \mathbb{C})^2 \times (\mathrm{PGL}(3, \mathbb{C})/H_{33}))/\mathbb{Z}_2.$$

By (5), we have that

$$\begin{aligned} e_H(B_r) &= \left(\frac{1}{2}(q^2 - 1)^r + \frac{1}{2}(q - 1)^{2r} - 2(q - 1)^r + 3^r \right) T \\ &\quad + \left(-\frac{1}{2}(q^2 - 1)^r + \frac{1}{2}(q - 1)^{2r} - (q - 1)^r + 3^r \right) N, \end{aligned}$$

since under $H \subset \Sigma_3$, we have $T \mapsto T$, $S \mapsto N$, $V \mapsto T + N$. For the second factor, $e((\mathbb{C}^r - \mathbb{C})^2) = (q^r - q)^2$ and $e(\mathrm{Sym}^2(\mathbb{C}^r - \mathbb{C})) = q^{2r} - q^{r+1}$, so

$$e_H((\mathbb{C}^r - \mathbb{C})^2) = (q^{2r} - q^{r+1})T + (q^2 - q^{r+1})N.$$

Finally, $\mathrm{PGL}(3, \mathbb{C})/H_{33} \cong \mathbb{P}^2 \times \mathbb{P}^2 - \Delta$, by considering the first two columns of the matrix, where Δ is the diagonal. As $e(\mathbb{P}^2 \times \mathbb{P}^2 - \Delta) = (1 + q + q^2)(q + q^2)$ and $e(\mathrm{Sym}^2 \mathbb{P}^2 - \bar{\Delta}) = q^4 + q^3 + q^2$, we have

$$e_H(\mathrm{PGL}(3, \mathbb{C})/H_{33}) = (q^4 + q^3 + q^2)T + (q^3 + q^2 + q)N.$$

Hence,

$$\begin{aligned} e(R_{33}) &= \left(\frac{1}{2}(q^2 - 1)^r + \frac{1}{2}(q - 1)^{2r} - 2(q - 1)^r + 3^r \right) \\ &\quad \times \left((q^{2r} - q^{r+1})(q^4 + q^3 + q^2) + (q^2 - q^{r+1})(q^3 + q^2 + q) \right) \\ &\quad + \left(-\frac{1}{2}(q^2 - 1)^r + \frac{1}{2}(q - 1)^{2r} - (q - 1)^r + 3^r \right) \\ &\quad \times \left((q^{2r} - q^{r+1})(q^3 + q^2 + q) + (q^2 - q^{r+1})(q^4 + q^3 + q^2) \right). \end{aligned}$$

- R_{34} consists of representations of type

$$A_i = \begin{pmatrix} \lambda_i & a_i & b_i \\ 0 & \mu_i & 0 \\ 0 & 0 & \gamma_i \end{pmatrix},$$

with $a \notin \langle \lambda - \mu \rangle$, $b \notin \langle \lambda - \gamma \rangle$. The stabilizer is

$$H_{34} = \left\{ \begin{pmatrix} a & b & c \\ 0 & d & 0 \\ 0 & 0 & e \end{pmatrix} \right\}.$$

The computations are analogous to the case of R_{33} , so $e(R_{33}) = e(R_{34})$.

- R_{35} consists of representations of type

$$A_i = \begin{pmatrix} \lambda_i & a_i & b_i \\ 0 & \mu_i & c_i \\ 0 & 0 & \gamma_i \end{pmatrix},$$

with $a \notin \langle \lambda - \mu \rangle$, $c \notin \langle \mu - \gamma \rangle$. The stabilizer is

$$H_{35} = \left\{ \begin{pmatrix} a & b & c \\ 0 & d & e \\ 0 & 0 & f \end{pmatrix} \right\}.$$

Hence

$$e(R_{35}) = ((q-1)^{2r} - 3(q-1)^r + 2 \cdot 3^r)(q^r - q)^2 q^r \frac{e(\mathrm{PGL}(3, \mathbb{C}))}{(q-1)^2 q^3}.$$

All together, we have

$$\begin{aligned} e(R_3) &= (2 \cdot 3^r - 3(q-1)^r + (q-1)^{2r}) \\ &\quad \times (q+1)(q^2 + q + 1)(q^r - q)(q^2 + q^{2r} - q^{1+r}) \\ &\quad + (2 \cdot 3^r - 2(q-1)^r + (q-1)^{2r} - (q^2 - 1)^r) \\ &\quad \times q(q^2 + q + 1)(q^r - q)(q^r - q^2) \\ &\quad + (2 \cdot 3^r - 4(q-1)^r + (q-1)^{2r} + (q^2 - 1)^r) \\ &\quad \times q^2(q^2 + q + 1)(q^r - 1)(q^r - q) \\ &\quad + \frac{1}{6}q^3((q-1)^{2r} - 3(q^2 - 1)^r + 2(q^2 + q + 1)^r \\ &\quad \quad + 2q(q+1)(3^{r+1} - 3(q-1)^r + (q-1)^{2r} - (q^2 + q + 1)^r)) \\ &\quad + \frac{1}{6}q^6(-6(q-1)^r + (q-1)^{2r} + 3(q^2 - 1)^r + 2(q^2 + q + 1)^r). \end{aligned}$$

Therefore,

$$\begin{aligned} e(\mathcal{R}_{r,3}^{\mathrm{red}}) &= \frac{1}{3}(q^2 + q + 1)^r (q-1)^2 q^3 (q+1) \\ &\quad + (q^2 + q + 1)(2q^{2r} - q^2)(q-1)^{2r} q^r (q+1)^r \\ &\quad - \frac{1}{3}(q-1)^{2r} (q+1)(q^2 + q + 1)(3q^{3r} - 3q^{r+2} + q^3), \end{aligned}$$

and so,

$$e(\mathcal{R}_{r,3}^{\text{irr}}) = e(\mathcal{R}_{r,3}) - e(\mathcal{R}_{r,3}^{\text{red}}) = e(\text{SL}(3, \mathbb{C}))^r - e(\mathcal{R}_{r,3}^{\text{red}}),$$

and consequently,

$$e(\mathcal{M}_{r,3}^{\text{irr}}) = e(\mathcal{R}_{r,3}^{\text{irr}})/e(\text{PGL}(3, \mathbb{C})) = e(\text{SL}(3, \mathbb{C}))^{r-1} - e(\mathcal{R}_{r,3}^{\text{red}})/e(\text{SL}(3, \mathbb{C})).$$

E-polynomial of the moduli of reducible representations. To compute $e(\mathcal{M}_{r,3})$, it remains to compute the moduli space of reducible representations $\mathcal{M}_{r,3}^{\text{red}}$. This is formed by two strata:

- (i) M_0 formed by semisimple representations which split into irreducible representations of ranks 1 and 2, that is, of the form:

$$A_i = \begin{pmatrix} \lambda_i^{-2} & 0 & 0 \\ 0 & \lambda_i B_i \\ 0 & 0 & 0 \end{pmatrix},$$

where $(B_1, \dots, B_r) \in \mathcal{M}_{r,2}^{\text{irr}}$. So $e(M_0) = (q - 1)^r e(\mathcal{M}_{r,2}^{\text{irr}})$.

- (ii) M_1 formed by semisimple representations which split into three irreducible representations of rank 1. These are given by eigenvalues $\lambda = (\lambda_1, \dots, \lambda_r)$, $\mu = (\mu_1, \dots, \mu_r)$, $\gamma = (\gamma_1, \dots, \gamma_r)$ in $(\mathbb{C}^*)^r$, where $\lambda_i \mu_i \gamma_i = 1$ for all $1 \leq i \leq r$. This is the space B'_r whose E -polynomial has been computed in (4). Thus

$$e(M_1) = e(B'_r/\Sigma_3) = \frac{1}{2}(q^2 - 1)^r + \frac{1}{6}(q - 1)^{2r} + \frac{1}{3}(q^2 + q + 1)^r.$$

Finally, $e(\mathcal{M}_{r,3}^{\text{red}}) = e(M_0) + e(M_1)$, and adding up everything we get

$$\begin{aligned} e(\mathcal{M}_{r,3}) &= e(\mathcal{M}_{r,3}^{\text{irr}}) + e(\mathcal{M}_{r,3}^{\text{red}}) \\ &= (q^8 - q^6 - q^5 + q^3)^{r-1} + (q - 1)^{2r-2}(q^{3r-3} - q^r) \\ &\quad + \frac{1}{6}(q - 1)^{2r-2}q(q + 1) + \frac{1}{2}(q^2 - 1)^{r-1}q(q - 1) \\ &\quad + \frac{1}{3}(q^2 + q + 1)^{r-1}q(q + 1) \\ &\quad - (q - 1)^{r-1}q^{r-1}(q^2 - 1)^{r-1}(2q^{2r-2} - q). \end{aligned}$$

This completes the main part of the proof of [Theorem 1](#). It remains to show that $e(\mathcal{M}_{r,3}) = e(\overline{\mathcal{M}}_{r,3})$, which we do in the following section.

Remark 8. By [\[Florentino and Lawton 2012\]](#), the singular locus of $\mathcal{M}_{r,3}$ is exactly the reducible locus (and so the smooth locus is its complement). Therefore, the above computation of M_0 and M_1 gives the E -polynomial of the singular locus of $\mathcal{M}_{r,3}$. Likewise, $e(\mathcal{M}_{r,3}^{\text{irr}})$ is the E -polynomial of the smooth locus of $\mathcal{M}_{r,3}$. Moreover, by [\[Florentino and Lawton 2014\]](#), the abelian character variety $\mathcal{M}(\mathbb{Z}^r, \text{SL}(3, \mathbb{C}))$ is exactly the diagonalizable representations in $\mathcal{M}_{r,3}$. The above

computation of M_1 gives the E -polynomial of $\mathcal{M}(\mathbb{Z}^r, \mathrm{SL}(3, \mathbb{C}))$. In each case, setting $q = 1$ gives the Euler characteristic of the corresponding space.

7. E -polynomials of character varieties for $F_r, r > 1$, and $\mathrm{PGL}(3, \mathbb{C})$

In this final section, we focus on the space of representations

$$\begin{aligned} \bar{\mathcal{R}}_{r,3} &= \mathrm{Hom}(F_r, \mathrm{PGL}(3, \mathbb{C})) = \{\rho : F_r \rightarrow \mathrm{PGL}(3, \mathbb{C})\} \\ &= \{(A_1, \dots, A_r) \mid A_i \in \mathrm{PGL}(3, \mathbb{C})\} = \mathrm{PGL}(3, \mathbb{C})^r \end{aligned}$$

and the character variety

$$\bar{\mathcal{M}}_{r,3} = \mathrm{Hom}(F_r, \mathrm{PGL}(3, \mathbb{C})) // \mathrm{PGL}(3, \mathbb{C}).$$

Let $\zeta = e^{2\pi\sqrt{-1}/3}$, and let $\mathbb{Z}_3 = \{1, \zeta, \zeta^2\}$ be the space of cubic roots of unity. Then $\mathrm{PGL}(3, \mathbb{C}) = \mathrm{SL}(3, \mathbb{C})/\mathbb{Z}_3$,

$$\bar{\mathcal{R}}_{r,3} = \mathcal{R}_{r,3}/(\mathbb{Z}_3)^r, \quad \text{and} \quad \bar{\mathcal{M}}_{r,3} = \mathcal{M}_{r,3}/(\mathbb{Z}_3)^r,$$

where $(\zeta^{a_1}, \dots, \zeta^{a_r})$ acts as $(A_1, \dots, A_r) \mapsto (\zeta^{a_1} A_1, \dots, \zeta^{a_r} A_r)$. Clearly $\bar{\mathcal{R}}_{r,3}^{\mathrm{red}} = \mathcal{R}_{r,3}^{\mathrm{red}}/(\mathbb{Z}_3)^r$ and $\bar{\mathcal{R}}_{r,3}^{\mathrm{irr}} = \mathcal{R}_{r,3}^{\mathrm{irr}}/(\mathbb{Z}_3)^r$.

We know from [Lemma 4](#) that $e(\mathrm{PGL}(3, \mathbb{C})) = e(\mathrm{SL}(3, \mathbb{C}))$. Let us see now that $e(\bar{\mathcal{R}}_{r,3}^{\mathrm{red}}) = e(\mathcal{R}_{r,3}^{\mathrm{red}})$. We stratify $\bar{\mathcal{R}}_{r,3}^{\mathrm{red}} = \bar{R}_0 \sqcup \bar{R}_1 \sqcup \bar{R}_2 \sqcup \bar{R}_3$, where $\bar{R}_i = R_i/(\mathbb{Z}_3)^r$ and the $R_i, i = 0, 1, 2, 3$, have been defined in [Section 6](#).

We now list the strata with the computation of their E -polynomials:

(i) $\bar{R}_0 = \bar{R}_{01} \cup \bar{R}_{02}$, where $\bar{R}_{0j} = R_{0j}/(\mathbb{Z}_3)^r, j = 1, 2$. To compute $e(\bar{R}_{01})$, recall that R_{01} is formed by representations $\rho = (A_1, \dots, A_r)$ with

$$A_i = \begin{pmatrix} \lambda_i^{-2} & b_i & c_i \\ 0 & \lambda_i B_i & \\ 0 & & \end{pmatrix},$$

where $(B_1, \dots, B_r) \in \mathcal{R}_{2,r}^{\mathrm{irr}}$. The action of ζ^{a_i} on A_i is given by $(\lambda_i, b_i, c_i, B_i) \mapsto (\zeta^{a_i} \lambda_i, \zeta^{a_i} b_i, \zeta^{a_i} c_i, \zeta^{a_i} B_i)$. Note that $\mathbb{C}/\mathbb{Z}_3 \cong \mathbb{C}$ and $\mathbb{C}^*/\mathbb{Z}_3 \cong \mathbb{C}^*$, so the relevant cohomology is invariant. Therefore

$$e((\mathbb{C}^*)^r \times \mathbb{C}^r \times \mathbb{C}^r \times \mathcal{R}_{2,r}^{\mathrm{irr}}/(\mathbb{Z}_3)^r) = e(\mathbb{C}^*)^r e(\mathbb{C})^r e(\mathbb{C})^r e(\mathcal{R}_{2,r}^{\mathrm{irr}}/(\mathbb{Z}_3)^r).$$

This means that $e(\bar{R}_{01}) = e(R_{01})$. Analogously $e(\bar{R}_{02}) = e(R_{02})$ and $e(\bar{R}_{01} \cap \bar{R}_{02}) = e(R_{01} \cap R_{02})$, so $e(\bar{R}_0) = e(R_0)$.

(ii) $\bar{R}_1 = R_1/(\mathbb{Z}_3)^r$. Note that R_1 is formed by 3^r copies of the same subvariety. Hence

$$e(\bar{R}_1) = \frac{e(R_1)}{3^r} = 1 + (1 + q + q^2)(q^{3r+1} + q^{3r} - 2q^{2r+1} + q - 1).$$

(iii) $\bar{R}_2 = R_2/(\mathbb{Z}_3)^r$. Recall that R_2 is formed by matrices with eigenvalues $(\lambda_i, \lambda_i, \mu_i)$ where $\lambda = (\lambda_1, \dots, \lambda_r) \in P = (\mathbb{C}^*)^r - \{1, \zeta, \zeta^2\}^r$. Now

$$\bar{P} = P/(\mathbb{Z}_3)^r \cong (\mathbb{C}^*)^r - \{(1, 1, \dots, 1)\},$$

so $e(\bar{P}) = (q-1)^r - 1$. It is more or less straightforward to see that \bar{R}_2 can be stratified by $\bar{R}_{2j} = R_{2j}/(\mathbb{Z}_3)^r$, $j = 1, 2, \dots, 9$. For each \bar{R}_{2j} the computation of $e(\bar{R}_{2j})$ is the same as that of $e(R_{2j})$, but replacing $e(P) = (q-1)^r - 3^r$ by $e(\bar{P}) = (q-1)^r - 1$. Hence

$$e(\bar{R}_2) = ((q-1)^r - 1)(q^2 + q + 1)(3q^{3r+1} + 3q^{3r} - 2q^{2r+2} - 4q^{2r+1} + q^3).$$

(iv) $\bar{R}_3 = R_3/(\mathbb{Z}_3)^r$. We follow the lines of the computation of $e(R_3)$. The base for the space of eigenvalues is $\bar{B}_r = B_r/(\mathbb{Z}_3)^r$ with $e(\bar{B}_r) = (q-1)^{2r} - 3(q-1)^r + 2$.

- Let $\bar{R}_{31} = R_{31}/(\mathbb{Z}_3)^r \cong (\text{PGL}(3, \mathbb{C})/D \times \bar{B}_r)/\Sigma_3$. If $\bar{B}'_r = B'_r/(\mathbb{Z}_3)^r$, then easily $e_{\Sigma_3}(\bar{B}'_r) = e_{\Sigma_3}(\bar{B}'_r)^r = (q^2T + S - qV)^r = e_{\Sigma_3}(B'_r)$. For $\bar{C}_r = C_r/(\mathbb{Z}_3)^r$, we have instead that $e_{\Sigma_3}(\bar{C}_r) = (q-1)^rT + ((q-1)^r - 1)V$, so $\bar{B}_r = \bar{B}'_r - \bar{C}_r$ has

$$\begin{aligned} e_{\Sigma_3}(B_r) &= \left(\frac{1}{2}(q^2 - 1)^r + \frac{1}{6}(q-1)^{2r} + \frac{1}{3}(q^2 + q + 1)^r - (q-1)^r\right)T \\ &\quad + \left(-\frac{1}{2}(q^2 - 1)^r + \frac{1}{6}(q-1)^{2r} + \frac{1}{3}(q^2 + q + 1)^r\right)S \\ &\quad + \left(\frac{1}{3}(q-1)^{2r} - \frac{1}{3}(q^2 + q + 1)^r - (q-1)^r + 1\right)V, \end{aligned}$$

and

$$\begin{aligned} e(\bar{R}_{31}) &= \left(\frac{1}{2}(q^2 - 1)^r + \frac{1}{6}(q-1)^{2r} + \frac{1}{3}(q^2 + q + 1)^r - (q-1)^r\right)q^6 \\ &\quad + \left(-\frac{1}{2}(q^2 - 1)^r + \frac{1}{6}(q-1)^{2r} + \frac{1}{3}(q^2 + q + 1)^r\right)q^3 \\ &\quad + \left(\frac{1}{3}(q-1)^{2r} - \frac{1}{3}(q^2 + q + 1)^r - (q-1)^r + 1\right)(q^5 + q^4). \end{aligned}$$

- $\bar{R}_{32} = R_{32}/(\mathbb{Z}_3)^r$ has

$$e(\bar{R}_{32}) = ((q-1)^{2r} - 3(q-1)^r + 2)(q^r - q) \frac{e(\text{PGL}(3, \mathbb{C}))}{(q-1)^2q}.$$

- $\bar{R}_{33} = R_{33}/(\mathbb{Z}_3)^r \cong (\bar{B}_r \times (\mathbb{C} - \mathbb{C})^2 \times (\text{PGL}(3, \mathbb{C})/H_{33}))/\mathbb{Z}_2$, where $H = \mathbb{Z}_2$ acts by swapping the first two eigenvalues. Now

$$\begin{aligned} e_H(\bar{B}_r) &= \left(\frac{1}{2}(q^2 - 1)^r + \frac{1}{2}(q-1)^{2r} - 2(q-1)^r + 1\right)T \\ &\quad + \left(-\frac{1}{2}(q^2 - 1)^r + \frac{1}{2}(q-1)^{2r} - (q-1)^r + 1\right)N, \end{aligned}$$

so

$$\begin{aligned} e(\bar{R}_{33}) &= \left(\frac{1}{2}(q^2 - 1)^r + \frac{1}{2}(q - 1)^{2r} - 2(q - 1)^r + 1\right) \\ &\quad \times \left((q^{2r} - q^{r+1})(q^4 + q^3 + q^2) + (q^2 - q^{r+1})(q^3 + q^2 + q)\right) \\ &\quad + \left(-\frac{1}{2}(q^2 - 1)^r + \frac{1}{2}(q - 1)^{2r} - (q - 1)^r + 1\right) \\ &\quad \times \left((q^{2r} - q^{r+1})(q^3 + q^2 + q) + (q^2 - q^{r+1})(q^4 + q^3 + q^2)\right). \end{aligned}$$

• $\bar{R}_{34} = R_{34}/(\mathbb{Z}_3)^r$ has $e(\bar{R}_{34}) = e(\bar{R}_{33})$.

• $\bar{R}_{35} = R_{35}/(\mathbb{Z}_3)^r$ has

$$e(\bar{R}_{35}) = ((q - 1)^{2r} - 3(q - 1)^r + 2)(q^r - q)^2 q^r \frac{e(\text{PGL}(3, \mathbb{C}))}{(q - 1)^2 q^3}.$$

All together, we have:

$$\begin{aligned} e(\bar{R}_3) &= (2 - 3(q - 1)^r + (q - 1)^{2r})(q + 1)(q^2 + q + 1) \\ &\quad \times (q^r - q)(q^2 + q^{2r} - q^{r+1}) \\ &\quad + (2 - 2(q - 1)^r + (q - 1)^{2r} - (q^2 - 1)^r) \\ &\quad \times q(q^2 + q + 1)(q^r - q)(q^r - q^2) \\ &\quad + (2 - 4(q - 1)^r + (q - 1)^{2r} + (q^2 - 1)^r) \\ &\quad \times q^2(q^2 + q + 1)(q^r - 1)(q^r - q) \\ &\quad + \frac{1}{6}q^3((q - 1)^{2r} - 3(q^2 - 1)^r + 2(q^2 + q + 1)^r \\ &\quad \quad + 2q(q + 1)(3 - 3(q - 1)^r + (q - 1)^{2r} - (q^2 + q + 1)^r)) \\ &\quad + \frac{1}{6}q^6(-6(q - 1)^r + (q - 1)^{2r} + 3(q^2 - 1)^r + 2(q^2 + q + 1)^r). \end{aligned}$$

Adding up all the contributions we get:

$$\begin{aligned} e(\bar{\mathcal{R}}_{r,3}^{\text{red}}) &= \frac{1}{3}(q^2 + q + 1)^r (q - 1)^2 q^3 (q + 1) \\ &\quad + (q^2 + q + 1)(2q^{2r} - q^2)(q - 1)^{2r} q^r (q + 1)^r \\ &\quad - \frac{1}{3}(q - 1)^{2r} (q + 1)(q^2 + q + 1)(3q^{3r} - 3q^{r+2} + q^3) \\ &= e(\mathcal{R}^{\text{red}}). \end{aligned}$$

From this $e(\bar{\mathcal{R}}_{r,3}^{\text{irr}}) = e(\mathcal{R}_{r,3}^{\text{irr}})$ and $e(\bar{\mathcal{M}}_{r,3}^{\text{irr}}) = e(\mathcal{M}_{r,3}^{\text{irr}})$.

The remaining thing to compute is $e(\bar{\mathcal{M}}_{r,3}^{\text{red}})$. This is formed by two strata:

(i) $\bar{M}_0 = M_0/(\mathbb{Z}_3)^r \cong ((\mathbb{C}^*)^r \times \mathcal{M}_{r,2}^{\text{irr}})/(\mathbb{Z}_3)^r$. Hence

$$e(\bar{M}_0) = (q - 1)^r e(\bar{\mathcal{M}}_{r,2}^{\text{irr}}) = e(M_0).$$

(ii) $\bar{M}_1 = M_1/(\mathbb{Z}_3)^r \cong ((\mathbb{C}^*)^r/(\mathbb{Z}_3)^r)/\Sigma_3 \cong (\mathbb{C}^*)^r/\Sigma_3$. So $e(\bar{M}_1) = e(M_1)$.

We get finally $e(\overline{\mathcal{M}}_{r,3}^{\text{red}}) = e(\mathcal{M}_{r,3}^{\text{red}})$. This concludes the proof of the equality $e(\overline{\mathcal{M}}_{r,3}) = e(\mathcal{M}_{r,3})$.

Remark 9. There is an arithmetic argument communicated to us by S. Mozgovoy to prove that $e(\overline{\mathcal{M}}_{r,n}) = e(\mathcal{M}_{r,n})$ for n odd. It goes as follows: find infinitely many primes p such that $p - 1$ and n are coprime (by Dirichlet's theorem on arithmetic progressions); then $\text{SL}(n, \mathbb{F}_p) \rightarrow \text{PGL}(n, \mathbb{F}_p)$ is bijective and one gets a bijection between corresponding character varieties over \mathbb{F}_p . So the count number of points of $\mathcal{M}_{r,n}$ and $\overline{\mathcal{M}}_{r,n}$ over \mathbb{F}_p coincide, and hence the E -polynomials coincide.

However this argument cannot be used for even n . Despite this, the E -polynomials for the $\text{SL}(2, \mathbb{C})$ -character varieties of free groups do equal those of $\text{PGL}(2, \mathbb{C})$. We expect to address the case of $\text{SL}(4, \mathbb{C})$ in future work.

Acknowledgments

We are grateful to S. Mozgovoy and M. Reineke for providing us with a copy of [Mozgovoy and Reineke 2015], and also for giving us an explicit formula for $e(\mathcal{M}(F_r, \text{PGL}(3, \mathbb{C})))$ for checking against our polynomials. We also thank the referee for helping improve the exposition of this article. Lawton was supported by the Simons Foundation Collaboration grant 245642, and the U.S. NSF grant DMS 1309376. Muñoz was partially supported by Project MICINN (Spain) MTM2010-17389. We also acknowledge support from U.S. NSF grants DMS 1107452, 1107263, 1107367 “RNMS: GEometric structures And Representation varieties” (the GEAR Network).

References

- [Cavazos and Lawton 2014] S. Cavazos and S. Lawton, “ E -polynomial of $\text{SL}_2(\mathbb{C})$ -character varieties of free groups”, *Int. J. Math.* **25**:6 (2014), 1450058. MR 3225582 Zbl 1325.14065
- [Deligne 1971] P. Deligne, “Théorie de Hodge, II”, *Inst. Hautes Études Sci. Publ. Math.* 40 (1971), 5–57. MR 58 #16653a Zbl 0219.14007
- [Deligne 1974] P. Deligne, “Théorie de Hodge, III”, *Inst. Hautes Études Sci. Publ. Math.* 44 (1974), 5–77. MR 58 #16653b Zbl 0237.14003
- [Florentino and Lawton 2009] C. Florentino and S. Lawton, “The topology of moduli spaces of free group representations”, *Math. Ann.* **345**:2 (2009), 453–489. MR 2010h:14075 Zbl 1200.14093
- [Florentino and Lawton 2012] C. Florentino and S. Lawton, “Singularities of free group character varieties”, *Pacific J. Math.* **260**:1 (2012), 149–179. MR 3001789 Zbl 1264.14064
- [Florentino and Lawton 2014] C. Florentino and S. Lawton, “Topology of character varieties of Abelian groups”, *Topology Appl.* **173** (2014), 32–58. MR 3227204 Zbl 1300.14045
- [Lawton 2007] S. Lawton, “Generators, relations and symmetries in pairs of 3×3 unimodular matrices”, *J. Algebra* **313**:2 (2007), 782–801. MR 2008k:16039 Zbl 1119.13004
- [Lawton 2008] S. Lawton, “Minimal affine coordinates for $\text{SL}(3, \mathbb{C})$ character varieties of free groups”, *J. Algebra* **320**:10 (2008), 3773–3810. MR 2009j:20060 Zbl 1157.14030

- [Lawton 2010] S. Lawton, “Algebraic independence in $SL(3, \mathbb{C})$ character varieties of free groups”, *J. Algebra* **324**:6 (2010), 1383–1391. MR 2011g:20071 Zbl 1209.14036
- [Logares and Muñoz 2014] M. Logares and V. Muñoz, “Hodge polynomials of the $SL(2, \mathbb{C})$ -character variety of an elliptic curve with two marked points”, *Int. J. Math.* **25**:14 (2014), 1450125. MR 3306833 Zbl 1316.14025
- [Logares et al. 2013] M. Logares, V. Muñoz, and P. E. Newstead, “Hodge polynomials of $SL(2, \mathbb{C})$ -character varieties for curves of small genus”, *Rev. Mat. Complut.* **26**:2 (2013), 635–703. MR 3068615
- [Martínez and Muñoz 2015a] J. Martínez and V. Muñoz, “E-polynomial of $SL(2, \mathbb{C})$ -character varieties of complex curves of genus 3”, preprint, 2015, Available at <http://www.math.sci.osaka-u.ac.jp/ojm/pdf/3905.pdf>. To appear in *Osaka J. Math.*
- [Martínez and Muñoz 2015b] J. Martínez and V. Muñoz, “E-polynomials of the $SL(2, \mathbb{C})$ -character varieties of surface groups”, *Int. Math. Res. Not.* **2015** (online publication June 2015).
- [Mozgovoy and Reineke 2015] S. Mozgovoy and M. Reineke, “Arithmetic of character varieties of free groups”, *Int. J. Math.* **26**:12 (2015), 1550100.
- [Sikora 2013] A. S. Sikora, “Generating sets for coordinate rings of character varieties”, *J. Pure Appl. Algebra* **217**:11 (2013), 2076–2087. MR 3057078 Zbl 1200.14093

Received March 24, 2015. Revised July 29, 2015.

SEAN LAWTON
DEPARTMENT OF MATHEMATICAL SCIENCES
GEORGE MASON UNIVERSITY
4400 UNIVERSITY DRIVE
FAIRFAX, VA 22030
UNITED STATES
slawton3@gmu.edu

VICENTE MUÑOZ
FACULTAD DE MATEMÁTICAS
UNIVERSIDAD COMPLUTENSE DE MADRID
PLAZA DE CIENCIAS 3
28040 MADRID
SPAIN
vicente.munoz@mat.ucm.es

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2016 is US \$440/year for the electronic version, and \$600/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY



mathematical sciences publishers

nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 282 No. 1 May 2016

On the half-space theorem for minimal surfaces in Heisenberg space	1
TRISTAN ALEX	
Extending smooth cyclic group actions on the Poincaré homology sphere	9
NIMA ANVARI	
A short proof of the existence of supercuspidal representations for all reductive p -adic groups	27
RAPHAËL BEUZART-PLESSIS	
Quantum groups and generalized circular elements	35
MICHAEL BRANNAN and KAY KIRKPATRICK	
Volumes of Montesinos links	63
KATHLEEN FINLINSON and JESSICA S. PURCELL	
Minimal surfaces with two ends which have the least total absolute curvature	107
SHOICHI FUJIMORI and TOSHIHIRO SHODA	
Multiplicité du spectre de Steklov sur les surfaces et nombre chromatique	145
PIERRE JAMMES	
E -polynomial of the $SL(3, \mathbb{C})$ -character variety of free groups	173
SEAN LAWTON and VICENTE MUÑOZ	
The Blum–Hanson property for $\mathcal{C}(K)$ spaces	203
PASCAL LEFÈVRE and ÉTIENNE MATHERON	
Crossed product algebras and direct integral decomposition for Lie supergroups	213
KARL-HERMANN NEEB and HADI SALMASIAN	
Associated primes of local cohomology modules over regular rings	233
TONY J. PUTHENPURAKAL	