

*Pacific  
Journal of  
Mathematics*

**A FUNCTIONAL CALCULUS AND  
RESTRICTION THEOREM ON H-TYPE GROUPS**

HEPING LIU AND MANLI SONG

Volume 286    No. 2

February 2017



## A FUNCTIONAL CALCULUS AND RESTRICTION THEOREM ON H-TYPE GROUPS

HEPING LIU AND MANLI SONG

**Let  $L$  be the sublaplacian and  $T$  the partial laplacian with respect to central variables on H-type groups. We investigate a class of invariant differential operators by the joint functional calculus of  $L$  and  $T$ . We establish Stein–Tomas type restriction theorems for these operators. In particular, the asymptotic behaviors of restriction estimates are given.**

### 1. Introduction

The restriction theorem for the Fourier transform plays an important role in harmonic analysis as well as in the theory of partial differential equations. The original version is credited to E. M. Stein and P. A. Tomas, and states that the transform of an  $L^p$ -function on  $\mathbb{R}^n$  has a well-defined restriction to the unit sphere  $S^{n-1}$  which is square integrable on  $S^{n-1}$ . The result is listed as follows:

**Theorem 1.1** [Stein 1993; Tomas 1975]. *Let  $1 \leq p \leq \frac{2n+2}{n+3}$ . Then the estimate*

$$(1-1) \quad \|\hat{f}\|_{L^2(S^{n-1})} \leq C \|f\|_{L^p(\mathbb{R}^n)}$$

*holds for all functions  $f \in L^p(\mathbb{R}^n)$ .*

A simple duality argument shows that the estimate (1-1) is equivalent to the following estimate:

$$(1-2) \quad \|f * \widehat{d\sigma_r}\|_{p'} \leq C_r \|f\|_p$$

for all Schwartz functions  $f$  on  $\mathbb{R}^n$ , where  $1/p + 1/p' = 1$  and  $d\sigma_r$  is the surface measure on the sphere with radius  $r$ .

---

The first author is supported by the National Natural Science Foundation of China under Grant #11371036 and the Specialized Research Fund for the Doctoral Program of Higher Education of China under Grant #2012000110059. The second author is supported by the China Scholarship Council under Grant #201206010098 and the Fundamental Research Funds for the Central Universities under Grant #3102015ZY068. Manli Song is the corresponding author.

*MSC2010:* primary 43A65; secondary 42B10, 47A60.

*Keywords:* H-type group, scaled special Hermite expansion, joint functional calculus, restriction operator.

Moreover, according to the Knapp example [Stein 1993], the estimates (1-1) and (1-2) fail if  $(2n + 2)/(n + 3) < p \leq 2$ .

Many authors have worked on the topic and various new restriction theorems have been proved. The study of restriction theorems has recently obtained more and more attention. A survey of recent progress on restriction theorems can be found in [Tao 2004]. To generalize the restriction theorem on the Heisenberg group, D. Müller [1990] established the boundedness of the restriction operator with respect to the mixed  $L^p$ -norm and also gave a counterexample to show that the estimate between Lebesgue spaces for the restriction operator was necessarily trivial, due to the fact that the center of the Heisenberg group was of dimension one. Some extensions have been treated by S. Thangavelu [1991a; 1991b]. Restriction theorems have been also studied in the case of the Heisenberg motion group by P. K. Ratnakumar, R. Rawat and S. Thangavelu [Ratnakumar et al. 1997], where groups with center with dimension higher than one were first considered.

On an H-type group, let  $T$  be the laplacian on the center and  $L$  the sublaplacian. It is well known that  $L$  is positive and essentially self-adjoint. Let  $L = \int_0^\infty \lambda dE(\lambda)$  be the spectral decomposition of  $L$ . Then the restriction operator can be formally written  $\mathcal{P}_\lambda f = \delta_\lambda(L)f = \lim_{\epsilon \rightarrow \infty} \chi_{(\lambda-\epsilon, \lambda+\epsilon)}(L)f$  which is well defined for a Schwartz function  $f$ , where  $\chi_{(\lambda-\epsilon, \lambda+\epsilon)}$  is the characteristic function of the interval  $(\lambda - \epsilon, \lambda + \epsilon)$ . Liu and Wang [2011] investigated the restriction theorem for the sublaplacian  $L$  on H-type groups with center whose dimension was greater than one. They gave the following result:

**Theorem 1.2.** *Let  $G$  be an H-type group with the underlying manifold  $\mathbb{R}^{2n+m}$ , where  $m > 1$  is the dimension of the center. Suppose  $1 \leq p \leq (2m + 2)/(m + 3)$ . Then the following estimate*

$$\|\mathcal{P}_\lambda f\|_{p'} \leq C \lambda^{2(n+m)\left(\frac{1}{p}-\frac{1}{2}\right)-1} \|f\|_p, \lambda > 0$$

*holds for all Schwartz functions  $f$  on  $G$ .*

V. Casarino and P. Ciatti [2013a; 2013b] extended the results of Müller, Liu and Wang to Métivier groups. They proved the restriction theorem for the sublaplacian and the full laplacian on Métivier groups. In fact, they also investigated the joint functional calculus of  $L$  and  $T$ . The invariant differential operators related to the joint functional calculus of  $L$  and  $T$  on H-type groups do not have the homogeneous properties in general. Thus the asymptotic behaviors of restriction estimates for these operators are also interesting. Casarino and Ciatti [2013a; 2013b] did not discuss the asymptotic behavior of the full laplacian. In this article we will show restriction estimates for these operators on H-type groups. In particular, the asymptotic behaviors of restriction estimates are given.

The outline of the paper is as follows. In the second section, we provide the necessary background for the H-type group. In the third section, by introducing the joint functional calculus of  $L$  and  $T$ , the restriction operator can be computed explicitly. In the fourth section, we prove the restriction theorem on H-type groups. In the fifth section, we describe the restriction theorems for other operators with the form of the joint functional calculus of  $L$  and  $T$ . Finally, in the last section, we show that the range of  $p$  in the restriction theorem is sharp.

### 2. Preliminaries

**Definition 2.1** (H-type group). Let  $\mathfrak{g}$  be a two step nilpotent Lie algebra endowed with an inner product  $\langle \cdot, \cdot \rangle$ . Its center is denoted by  $\mathfrak{z}$ . The algebra  $\mathfrak{g}$  is said to be of H-type if  $[\mathfrak{z}^\perp, \mathfrak{z}^\perp] = \mathfrak{z}$  and for every  $t \in \mathfrak{z}$ , the map  $J_t : \mathfrak{z}^\perp \rightarrow \mathfrak{z}^\perp$  defined by

$$\langle J_t u, w \rangle := \langle t, [u, w] \rangle \text{ for all } u, w \in \mathfrak{z}^\perp$$

is an orthogonal map whenever  $|t| = 1$ .

An H-type group is a connected and simply connected Lie group  $G$  whose Lie algebra is of H-type.

For a given  $0 \neq a \in \mathfrak{z}^*$ , the dual of  $\mathfrak{z}$ , we can define a skew-symmetric mapping  $B(a)$  on  $\mathfrak{z}^\perp$  by

$$\langle B(a)u, w \rangle = a([u, w]) \text{ for all } u, w \in \mathfrak{z}^\perp.$$

We denote by  $z_a$  the element of  $\mathfrak{z}$  determined by

$$\langle B(a)u, w \rangle = a([u, w]) = \langle J_{z_a} u, w \rangle.$$

Since  $B(a)$  is skew-symmetric and nondegenerate, the dimension of  $\mathfrak{z}^\perp$  is even, i.e.,  $\dim \mathfrak{z}^\perp = 2n$ .

For a given  $0 \neq a \in \mathfrak{z}^*$ , we can choose an orthonormal basis

$$\{E_1(a), E_2(a), \dots, E_n(a), \bar{E}_1(a), \bar{E}_2(a), \dots, \bar{E}_n(a)\}$$

of  $\mathfrak{z}^\perp$  such that

$$B(a)E_i(a) = |z_a| \frac{J_{z_a}}{|z_a|} E_i(a) = |a| \bar{E}_i(a)$$

and

$$B(a)\bar{E}_i(a) = -|a|E_i(a).$$

We set  $m = \dim \mathfrak{z}$ . Throughout this paper we assume that  $m > 1$ . We can choose an orthonormal basis  $\{\epsilon_1, \epsilon_2, \dots, \epsilon_m\}$  of  $\mathfrak{z}$  such that  $a(\epsilon_1) = |a|$  and  $a(\epsilon_j) = 0$ ,  $j = 2, 3, \dots, m$ . Then we can denote the elements of  $\mathfrak{g}$  by

$$(z, t) = (x, y, t) = \sum_{i=1}^n (x_i E_i + y_i \bar{E}_i) + \sum_{j=1}^m t_j \epsilon_j.$$

We identify  $G$  with its Lie algebra  $\mathfrak{g}$  via the exponential map. The group law on H-type group  $G$  has the form

$$(2-1) \quad (z, t)(z', t') = (z + z', t + t' + \frac{1}{2}[z, z']),$$

where  $[z, z']_j = \langle z, U^j z' \rangle$  for a suitable skew-symmetric matrix  $U^j, j = 1, 2, \dots, m$ .

**Theorem 2.2.**  *$G$  is an H-type group with underlying manifold  $\mathbb{R}^{2n+m}$ , with the group law (2-1) and the matrix  $U^j, j = 1, 2, \dots, m$  satisfies the following conditions:*

- (i)  $U^j$  is a  $2n \times 2n$  skew-symmetric and orthogonal matrix,  $j = 1, 2, \dots, m$ .
- (ii)  $U^i U^j + U^j U^i = 0$ , where  $i, j = 1, 2, \dots, m$  with  $i \neq j$ .

*Proof.* See [Bonfiglioli and Uguzzoni 2004]. □

**Remark 2.3.** In particular,  $\langle z, U^1 z' \rangle = \sum_{j=1}^n (x'_j y_j - y'_j x_j)$ .

**Remark 2.4.** All the above expressions depend on a given  $0 \neq a \in \mathfrak{z}^*$ , but we will suppress  $a$  from them for simplification.

**Remark 2.5.** It is well know that H-type algebras are closely related to Clifford modules [Reimann 2001]. H-type algebras can be classified by the standard theory of Clifford algebras. Especially, on the H-type group  $G$ , there is a relation between the dimension of the center and its orthogonal complement space. That is  $m + 1 \leq 2n$  (see [Kaplan and Ricci 1983]).

The left invariant vector fields which agree respectively with  $\partial/\partial x_j, \partial/\partial y_j$  at the origin are given by

$$X_j = \frac{\partial}{\partial x_j} + \frac{1}{2} \sum_{k=1}^m \left( \sum_{l=1}^{2n} z_l U_{l,j}^k \right) \frac{\partial}{\partial t_k},$$

$$Y_j = \frac{\partial}{\partial y_j} + \frac{1}{2} \sum_{k=1}^m \left( \sum_{l=1}^{2n} z_l U_{l,j+n}^k \right) \frac{\partial}{\partial t_k},$$

where  $z_l = x_l, z_{l+n} = y_l, l = 1, 2, \dots, n$ .

The vector fields  $T_k = \partial/\partial t_k, k = 1, 2, \dots, m$  correspond to the center of  $G$ . In terms of these vector fields, we introduce the sublaplacian  $L$  and full laplacian  $\Delta$  respectively

$$(2-2) \quad L = - \sum_{j=1}^n (X_j^2 + Y_j^2) = -\Delta_z + \frac{1}{4}|z|^2 T - \sum_{k=1}^m \langle z, U^k \nabla_z \rangle T_k,$$

$$(2-3) \quad \Delta = L + T,$$

where

$$\Delta_z = \sum_{j=1}^{2n} \frac{\partial^2}{\partial z_j^2}, \quad T = - \sum_{k=1}^m \frac{\partial^2}{\partial t_k^2}, \quad \nabla_z = \left( \frac{\partial}{\partial z_1}, \frac{\partial}{\partial z_2}, \dots, \frac{\partial}{\partial z_{2n}} \right)^t.$$

### 3. The restriction operator

First we recall some results about the scaled special Hermite expansion. We refer the reader to [Thangavelu 1993, 2004] for details. Letting  $\lambda > 0$ , the twisted laplacian (or the scaled special Hermite expansion)  $L_\lambda$  is defined by

$$L_\lambda = -\Delta_z + \frac{\lambda^2 |z|^2}{4} - i\lambda \sum_{j=1}^n \left( x_j \frac{\partial}{\partial y_j} - y_j \frac{\partial}{\partial x_j} \right),$$

where we identify  $z = x + iy \in \mathbb{C}^n$  with  $z = (x, y) \in \mathbb{R}^{2n}$ .

For  $f, g \in L^1(\mathbb{C}^n)$ , we define the  $\lambda$ -twisted convolution by

$$f \times_\lambda g = \int_{\mathbb{C}^n} f(z-w)g(w)e^{\frac{1}{2}i\lambda \operatorname{Im} z \cdot \bar{w}} dw.$$

Set Laguerre function  $\varphi_k^\lambda(z) = L_k^{n-1}(\frac{1}{2}\lambda|z|^2)e^{-\frac{1}{4}\lambda|z|^2}$ ,  $k = 0, 1, 2, \dots$ , where  $L_k^{n-1}$  is the Laguerre polynomial of type  $(n-1)$  and degree  $k$ .

For any Schwartz function  $f$  on  $\mathbb{C}^n$ , we have the scaled special Hermite expansion

$$(3-1) \quad f(z) = \left( \frac{\lambda}{2\pi} \right)^n \sum_{k=0}^{\infty} f \times_\lambda \varphi_k^\lambda(z),$$

which is an orthogonal form. We also have

$$(3-2) \quad \|f\|^2 = \left( \frac{\lambda}{2\pi} \right)^n \sum_{k=0}^{\infty} \|f \times_\lambda \varphi_k^\lambda\|^2.$$

Moreover,  $f \times_\lambda \varphi_k^\lambda$  is an eigenfunction of  $L_\lambda$  with the eigenvalue  $(2k+n)\lambda$  and

$$(3-3) \quad \|f \times_\lambda \varphi_k^\lambda\|_2 \leq (2k+n)^{n(\frac{1}{p}-\frac{1}{2})-\frac{1}{2}} \lambda^{n(\frac{1}{p}-\frac{3}{2})} \|f\|_p \quad \text{for } 1 \leq p < \frac{6n+2}{3n+4}$$

(see [Thangavelu 1991b]).

Now we turn to the expression for the restriction operator. We may identify  $\mathfrak{z}^*$  with  $\mathfrak{z}$ . Therefore, we will write  $\langle a, t \rangle$  instead of  $a(t)$  for  $a \in \mathfrak{z}^*$  and  $t \in \mathfrak{z}$ .

**Lemma 3.1.** *Let  $0 \neq a \in \mathfrak{z}^*$ . If  $f(z, t) = e^{-i\langle a, t \rangle} \varphi(z)$ , then*

$$Lf(z, t) = e^{-i\langle a, t \rangle} L_{|a|} \varphi(z).$$

*Proof.* Because  $\langle a, t \rangle = |a|t_1$  and  $\langle z, U^1 \nabla_z \rangle = \sum_{j=1}^n (y_j \frac{\partial}{\partial x_j} - x_j \frac{\partial}{\partial y_j})$ , Lemma 3.1 is easily deduced from the expression (2-2). □

Set  $e_k^a(z, t) = e^{-i\langle a, t \rangle} \varphi_k^{|a|}(z)$ . For  $f \in \mathcal{S}(G)$ , let

$$f^a(z) = \int_{\mathbb{R}^m} f(z, t) e^{i\langle a, t \rangle} dt$$

be the Fourier transform of  $f$  with respect to the central variable  $t$ . It is easy to obtain

$$(3-4) \quad f * e_k^a(z, t) = e^{-i\langle a, t \rangle} f^a \times_{|a|} \varphi_k^{|a|}(z).$$

Note that  $f * e_k^a$  is an eigenfunction of  $T$  with the eigenvalue  $|a|^2$ . Furthermore, it follows from Lemma 3.1 that  $f * e_k^a$  is an eigenfunction of  $L$  with the eigenvalue  $(2k + n)|a|$ . Thus  $f * e_k^a$  is a joint eigenfunction of the operators  $L$  and  $T$ .

For a Schwartz function  $f$  on an H-type group, using the inversion formula for the Fourier transform together with (3-1) and (3-4), we have

$$\begin{aligned} f(z, t) &= \frac{1}{(2\pi)^m} \int_{\mathbb{R}^m} f^a(z) e^{-i\langle a, t \rangle} da \\ &= \frac{1}{(2\pi)^m} \int_{\mathbb{R}^m} \left( \frac{|a|^n}{(2\pi)^n} \sum_{k=0}^{\infty} f^a \times_{|a|} \varphi_k^{|a|}(z) \right) e^{-i\langle a, t \rangle} da \\ &= \frac{1}{(2\pi)^{n+m}} \int_{\mathbb{R}^m} \sum_{k=0}^{\infty} f * e_k^a(z, t) |a|^n da \\ &= \int_0^{\infty} \left( \frac{1}{(2\pi)^{n+m}} \sum_{k=0}^{\infty} \lambda^{n+m-1} \int_{S^{m-1}} f * e_k^{\lambda \tilde{a}}(z, t) d\sigma(\tilde{a}) \right) d\lambda. \end{aligned}$$

The operators  $L$  and  $T$  extend to a pair of strongly commuting self-adjoint operators. Therefore, they admit a joint spectral decomposition. By the spectral theorem, we can define the joint functional calculus of  $L$  and  $T$ . The joint functional calculus of  $L$  and  $T$  was investigated in [Casarino and Ciatti 2013a]. As in that paper, we define the operator  $\delta_\mu(h(L, T))$  for a suitable function  $h : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$  as

$$h(L, T)f(z, t) = \int_0^{\infty} \left( \frac{1}{(2\pi)^{n+m}} \sum_{k=0}^{\infty} h((2k + n)\lambda, \lambda^2) \lambda^{n+m-1} \int_{S^{m-1}} f * e_k^{\lambda \tilde{a}}(z, t) d\sigma(\tilde{a}) \right) d\lambda,$$

where we make the assumption on  $h$  that the expression on the right-hand side is a well-defined distribution for all Schwartz functions  $f$ . We also suppose  $h((2k + n)\lambda, \lambda^2)$  is a strictly monotonic differentiable positive function of  $\lambda$  on  $\mathbb{R}_+$ , with the domain  $(A, B)$  where  $0 \leq A < B \leq \infty$ . Then for each  $\mu \in (A, B)$ , the equation

$$h((2k + n)\lambda, \lambda^2) = \mu$$



may be solved for each  $k$ . We denote the solution by  $\lambda = \lambda_k(\mu)$  and  $\lambda'_k$  denotes the derivative of  $\lambda_k$ . Replacing  $\lambda$  with  $\mu$  in the integral, we obtain

$$h(L, T)f(z, t) = \int_A^B \mu \left( \frac{1}{(2\pi)^{n+m}} \sum_{k=0}^{\infty} \lambda_k^{n+m-1}(\mu) |\lambda'_k(\mu)| \int_{S^{m-1}} f * e_k^{\lambda_k(\mu)\tilde{a}}(z, t) d\sigma(\tilde{a}) \right) d\mu,$$

which is the spectral decomposition of  $h(L, T)$ .

Thus, given a Schwartz function  $f$ , the spectral decomposition with respect to  $h(L, T)$  is

$$f(z, t) = \int_A^B \left( \frac{1}{(2\pi)^{n+m}} \sum_{k=0}^{\infty} \lambda_k^{n+m-1}(\mu) |\lambda'_k(\mu)| \int_{S^{m-1}} f * e_k^{\lambda_k(\mu)\tilde{a}}(z, t) d\sigma(\tilde{a}) \right) d\mu.$$

We can also use this equation to introduce the spectral resolution of  $h(L, T)$ , which is defined by

$$(3-5) \quad \mathcal{P}_\mu^h f(z, t) = \delta_\mu(h(L, T))f(z, t) = \lim_{\epsilon \rightarrow 0^+} \frac{1}{2\epsilon} \chi_{(\mu-\epsilon, \mu+\epsilon)}(h(L, T))f,$$

where  $f$  is a Schwartz function and  $\chi_{(\mu-\epsilon, \mu+\epsilon)}$  is the characteristic function of the interval  $(\mu - \epsilon, \mu + \epsilon)$ . We easily find

$$\mathcal{P}_\mu^h f(z, t) = \frac{1}{(2\pi)^{n+m}} \sum_{k=0}^{\infty} \lambda_k^{n+m-1}(\mu) |\lambda'_k(\mu)| \int_{S^{m-1}} f * e_k^{\lambda_k(\mu)\tilde{a}}(z, t) d\sigma(\tilde{a}).$$

Specifically, for the full laplacian  $\Delta$ ,  $h(\xi, \eta) = \xi + \eta$ , so we have  $\mu = (2k + n)\lambda + \lambda^2$ , which yields

$$(3-6) \quad \lambda_k(\mu) = \frac{1}{2} \sqrt{4\mu + (2k + n)^2} - \frac{2k + n}{2} \quad \text{and} \quad \lambda'_k(\mu) = \frac{1}{\sqrt{4\mu + (2k + n)^2}}.$$

Therefore,

$$\mathcal{P}_\mu^\Delta f(z, t) = \frac{1}{(2\pi)^{n+m}} \sum_{k=0}^{\infty} \lambda_k^{n+m-1}(\mu) \lambda'_k(\mu) \int_{S^{m-1}} f * e_k^{\lambda_k(\mu)\tilde{a}}(z, t) d\sigma(\tilde{a}).$$

#### 4. The restriction theorem

Our main result is the following theorem.

**Theorem 4.1.** *Let  $G$  be an H-type group with the underlying manifold  $\mathbb{R}^{2n+m}$ , where  $m > 1$  is the dimension of the center. Let  $h(\xi, \eta) = \xi^\alpha + \eta^\beta$ ,  $\alpha, \beta > 0$ . Then*

for  $1 \leq p \leq (2m + 2)/(m + 3)$ , we have for all Schwartz functions  $f$ :

$$\begin{aligned}
 \text{if } \alpha < 2\beta & \begin{cases} \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{\frac{2}{\alpha}(n+\frac{\alpha}{2\beta}m)}(\frac{1}{p}-\frac{1}{2})^{-1} \|f\|_p, & \mu > 1, \\ \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{\frac{2}{\alpha}(n+m)}(\frac{1}{p}-\frac{1}{2})^{-1} \|f\|_p, & 0 < \mu \leq 1, \end{cases} \\
 \text{if } \alpha > 2\beta & \begin{cases} \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{\frac{2}{\alpha}(n+m)}(\frac{1}{p}-\frac{1}{2})^{-1} \|f\|_p, & \mu > 1, \\ \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{\frac{2}{\alpha}(n+\frac{\alpha}{2\beta}m)}(\frac{1}{p}-\frac{1}{2})^{-1} \|f\|_p, & 0 < \mu \leq 1, \end{cases} \\
 \text{if } \alpha = 2\beta & \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{\frac{2}{\alpha}(n+m)}(\frac{1}{p}-\frac{1}{2})^{-1} \|f\|_p, \quad 0 < \mu < \infty.
 \end{aligned}$$

First, we have the following abstract statement.

**Proposition 4.2.** *The function  $h((2k + n)\lambda, \lambda^2)$  is a strictly monotonic differentiable positive function of  $\lambda$  on  $\mathbb{R}_+$ , with the domain  $(A, B)$  where  $0 \leq A < B \leq \infty$ . Then for  $1 \leq p \leq (2m + 2)/(m + 3)$ , the estimate*

$$\|\mathcal{P}_\mu^h f\|_{p'} \leq C_\mu \|f\|_p$$

holds, where

$$(4-1) \quad C_\mu \leq C \sum_{k=0}^\infty (2k + n)^{2n} (\frac{1}{p}-\frac{1}{2})^{-1} \lambda_k^{2(n+m)} (\frac{1}{p}-\frac{1}{2})^{-1} (\mu) |\lambda'_k(\mu)|$$

for all Schwartz functions  $f$  and all positive  $\mu \in (A, B)$ .

The proof of Proposition 4.2 coincides essentially with Theorem 4.1 in [Casarino and Ciatti 2013a] and we omit it. To obtain our Theorem 4.1, it suffices to show the convergence of the series in (4-1). Next we will exploit the following estimates, which can be easily proved by comparing the sums with the corresponding integrals:

**Lemma 4.3.** *Fix  $v \in \mathbb{R}$ . There exists  $C_v > 0$  such that for  $A > 0$  and  $n \in \mathbb{Z}_+$ , we have*

$$(4-2) \quad \sum_{\substack{m \in \mathbb{N} \\ 2m+n \geq A}} (2m + n)^v \leq C_v A^{v+1}, \quad v < -1;$$

$$(4-3) \quad \sum_{\substack{m \in \mathbb{N} \\ 2m+n \leq A}} (2m + n)^v \leq C_v A^{v+1}, \quad v > -1.$$

Now Theorem 4.1 follows from the result in the following lemma.

**Lemma 4.4.** *Let  $h(\xi, \eta) = \xi^\alpha + \eta^\beta$ ,  $\alpha, \beta > 0$ . The series in (4-1) has the estimate*

$$\begin{aligned}
 \text{if } \alpha < 2\beta & \begin{cases} C_\mu \leq C\mu^{\frac{2}{\alpha}(n+\frac{\alpha}{2\beta}m)(\frac{1}{p}-\frac{1}{2})-1}, & \mu > 1, \\ C_\mu \leq C\mu^{\frac{2}{\alpha}(n+m)(\frac{1}{p}-\frac{1}{2})-1}, & 0 < \mu \leq 1, \end{cases} \\
 \text{if } \alpha > 2\beta & \begin{cases} C_\mu \leq C\mu^{\frac{2}{\alpha}(n+m)(\frac{1}{p}-\frac{1}{2})-1}, & \mu > 1, \\ C_\mu \leq C\mu^{\frac{2}{\alpha}(n+\frac{\alpha}{2\beta}m)(\frac{1}{p}-\frac{1}{2})-1}, & 0 < \mu \leq 1, \end{cases} \\
 \text{if } \alpha = 2\beta & \begin{cases} C_\mu \leq C\mu^{\frac{2}{\alpha}(n+m)(\frac{1}{p}-\frac{1}{2})-1}, & 0 < \mu < \infty. \end{cases}
 \end{aligned}$$

*Proof.* The function  $h(\xi, \eta) = \xi^\alpha + \eta^\beta$ ,  $\alpha, \beta > 0$ , so  $\mu = (2k+n)^\alpha \lambda_k^\alpha(\mu) + \lambda_k^{2\beta}(\mu)$ , which yields

$$\lambda'_k(\mu) = \frac{1}{\alpha(2k+n)^\alpha \lambda_k^{\alpha-1}(\mu) + 2\beta \lambda_k^{2\beta-1}(\mu)}.$$

To study the convergence of this series, we need to distinguish three cases according to the relation of  $\alpha$  and  $2\beta$ :  $\alpha < 2\beta$ ,  $\alpha > 2\beta$  and  $\alpha = 2\beta$ . In order not to burden the exposition, we only prove the case  $\alpha < 2\beta$ , and the other cases are analogous.

If  $\alpha < 2\beta$ , then when  $\mu \leq 1$ , it is easy to see that  $\lambda_k(\mu) \sim \mu^{\frac{1}{\alpha}}/(2k+n)$  and  $\lambda'_k(\mu) \sim \mu^{\frac{1}{\alpha}-1}/(2k+n)$ , so that the series

$$\begin{aligned}
 (4-4) \quad C_\mu & \leq C \sum_{k=0}^{\infty} (2k+n)^{2n(\frac{1}{p}-\frac{1}{2})-1} \lambda_k^{2(n+m)(\frac{1}{p}-\frac{1}{2})-1}(\mu) |\lambda'_k(\mu)| \\
 & \leq C \sum_{k=0}^{\infty} (2k+n)^{2n(\frac{1}{p}-\frac{1}{2})-1} \left( \frac{\mu^{\frac{1}{\alpha}}}{2k+n} \right)^{2(n+m)(\frac{1}{p}-\frac{1}{2})-1} \frac{\mu^{\frac{1}{\alpha}-1}}{2k+n} \\
 & \leq C\mu^{\frac{2}{\alpha}(n+m)(\frac{1}{p}-\frac{1}{2})-1} \sum_{k=0}^{\infty} \frac{1}{(2k+n)^{2m(\frac{1}{p}-\frac{1}{2})+1}} \\
 & \leq C\mu^{\frac{2}{\alpha}(n+m)(\frac{1}{p}-\frac{1}{2})-1}
 \end{aligned}$$

converges.

When  $\mu > 1$ , we split the sum into two parts, the sum over those  $k$  such that  $(2k+n)^\alpha \lambda_k^\alpha(\mu) \geq \lambda^{2\beta}(\mu)$  and those such that  $(2k+n)^\alpha \lambda_k^\alpha(\mu) < \lambda^{2\beta}(\mu)$ . They are denoted by I and II respectively.

For the first part,  $(2k+n)^\alpha \lambda_k^\alpha(\mu) \geq \lambda^{2\beta}(\mu)$  implies

$$\lambda_k(\mu) \sim \frac{\mu^{\frac{1}{\alpha}}}{2k+n}, \quad \lambda'_k(\mu) \sim \frac{\mu^{\frac{1}{\alpha}-1}}{2k+n}, \quad \text{and} \quad 2k+n \geq \mu^{\frac{2\beta-\alpha}{2\alpha\beta}}.$$

Then we control the first part I by

$$\begin{aligned}
I &\leq C \sum_{2k+n \geq \mu^{\frac{2\beta-\alpha}{2\alpha\beta}}} (2k+n)^{2n\left(\frac{1}{p}-\frac{1}{2}\right)-1} \lambda_k^{2(n+m)\left(\frac{1}{p}-\frac{1}{2}\right)-1}(\mu) |\lambda'_k(\mu)| \\
&\leq C \sum_{2k+n \geq \mu^{\frac{2\beta-\alpha}{2\alpha\beta}}} (2k+n)^{2n\left(\frac{1}{p}-\frac{1}{2}\right)-1} \left(\frac{\mu^{\frac{1}{\alpha}}}{2k+n}\right)^{2(n+m)\left(\frac{1}{p}-\frac{1}{2}\right)-1} \frac{\mu^{\frac{1}{\alpha}-1}}{2k+n} \\
&\leq C \mu^{\frac{2}{\alpha}(n+m)\left(\frac{1}{p}-\frac{1}{2}\right)-1} \sum_{2k+n \geq \mu^{\frac{2\beta-\alpha}{2\alpha\beta}}} \frac{1}{(2k+n)^{2m\left(\frac{1}{p}-\frac{1}{2}\right)+1}}.
\end{aligned}$$

By (4-2), we have

$$(4-5) \quad I \leq C \mu^{\frac{2}{\alpha}(n+m)\left(\frac{1}{p}-\frac{1}{2}\right)-1} \frac{1}{\mu^{\frac{2\beta-\alpha}{2\alpha\beta}\left(2m\left(\frac{1}{p}-\frac{1}{2}\right)\right)}} \leq C \mu^{\frac{2}{\alpha}\left(n+\frac{\alpha}{2\beta}m\right)\left(\frac{1}{p}-\frac{1}{2}\right)-1}.$$

For the second part,  $(2k+n)^\alpha \lambda_k^\alpha(\mu) < \lambda^{2\beta}(\mu)$  implies

$$\lambda_k(\mu) \sim \mu^{\frac{1}{2\beta}}, \quad \lambda'_k(\mu) \sim \mu^{\frac{1}{2\beta}-1}, \quad \text{and} \quad 2k+n < \mu^{\frac{2\beta-\alpha}{2\alpha\beta}}.$$

Then we control the second part II by

$$\begin{aligned}
II &\leq C \sum_{2k+n < \mu^{\frac{2\beta-\alpha}{2\alpha\beta}}} (2k+n)^{2n\left(\frac{1}{p}-\frac{1}{2}\right)-1} \lambda_k^{2(n+m)\left(\frac{1}{p}-\frac{1}{2}\right)-1}(\mu) |\lambda'_k(\mu)| \\
&\leq C \sum_{2k+n < \mu^{\frac{2\beta-\alpha}{2\alpha\beta}}} (2k+n)^{2n\left(\frac{1}{p}-\frac{1}{2}\right)-1} \left(\mu^{\frac{1}{2\beta}}\right)^{2(n+m)\left(\frac{1}{p}-\frac{1}{2}\right)-1} \mu^{\frac{1}{2\beta}-1} \\
&\leq C \mu^{\frac{1}{\beta}(n+m)\left(\frac{1}{p}-\frac{1}{2}\right)-1} \sum_{2k+n < \mu^{\frac{2\beta-\alpha}{2\alpha\beta}}} (2k+n)^{2n\left(\frac{1}{p}-\frac{1}{2}\right)-1}.
\end{aligned}$$

Because  $1 \leq p \leq (2m+2)/(m+3)$ , we obtain  $2n\left(\frac{1}{p}-\frac{1}{2}\right)-1 \geq -1$ . Hence, by (4-3) we get

$$\sum_{2k+n < \mu^{\frac{2\beta-\alpha}{2\alpha\beta}}} (2k+n)^{2n\left(\frac{1}{p}-\frac{1}{2}\right)-1} \lesssim \mu^{\frac{2\beta-\alpha}{2\alpha\beta}\left(2n\left(\frac{1}{p}-\frac{1}{2}\right)\right)} = \mu^{\left(\frac{2}{\alpha}-\frac{1}{\beta}\right)n\left(\frac{1}{p}-\frac{1}{2}\right)}.$$

Thus, for the second part we also have

$$(4-6) \quad II \leq C \mu^{\frac{2}{\alpha}\left(n+\frac{\alpha}{2\beta}m\right)\left(\frac{1}{p}-\frac{1}{2}\right)-1}.$$

Finally, the estimate for the case  $\alpha < 2\beta$  follows from (4-4), (4-5) and (4-6). This completes the proof of the first case.

Combining Proposition 4.2 and Lemma 4.4, Theorem 4.1 comes out easily.  $\square$

Especially, in the case  $\Delta = L + T$ ,  $h(\xi, \eta) = \xi + \eta$ , we obtain the restriction theorem associated with the full laplacian on H-type groups.

**Corollary 4.5.** *For  $1 \leq p \leq (2m + 2)/(m + 3)$ , the estimates*

$$\|\mathcal{P}_\mu^\Delta f\|_{p'} \leq C\mu^{(2n+m)(\frac{1}{p}-\frac{1}{2})-1} \|f\|_p, \mu > 1$$

and

$$\|\mathcal{P}_\mu^\Delta f\|_{p'} \leq C\mu^{2(n+m)(\frac{1}{p}-\frac{1}{2})-1} \|f\|_p, 0 < \mu \leq 1$$

hold for all Schwartz functions  $f$ .

### 5. Examples

Similarly to what we have done so far in Theorem 4.1, we now discuss other operators with the form of the joint functional calculus of  $L$  and  $T$ . We obtain the following results. We omit the arguments which are really similar to that of Theorem 4.1.

**Example 5.1.** Let  $h(\xi, \eta) = (\xi^\alpha + \eta^\beta)^{-1}$ ,  $\alpha, \beta > 0$ . For  $1 \leq p \leq (2m + 2)/(m + 3)$ , we have for all Schwartz functions  $f$ :

$$\begin{aligned} \text{if } \alpha < 2\beta & \begin{cases} \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha}(n+m)(\frac{1}{p}-\frac{1}{2})-1} \|f\|_p, & \mu > 1, \\ \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha}(n+\frac{\alpha}{2\beta}m)(\frac{1}{p}-\frac{1}{2})-1} \|f\|_p, & 0 < \mu \leq 1, \end{cases} \\ \text{if } \alpha > 2\beta & \begin{cases} \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha}(n+\frac{\alpha}{2\beta}m)(\frac{1}{p}-\frac{1}{2})-1} \|f\|_p, & \mu > 1, \\ \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha}(n+m)(\frac{1}{p}-\frac{1}{2})-1} \|f\|_p, & 0 < \mu \leq 1, \end{cases} \\ \text{if } \alpha = 2\beta & \begin{cases} \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha}(n+m)(\frac{1}{p}-\frac{1}{2})-1} \|f\|_p, & 0 < \mu < \infty. \end{cases} \end{aligned}$$

**Example 5.2.** Let  $h(\xi, \eta) = (1 + \xi)^{-1}$ . For  $1 \leq p \leq (2m + 2)/(m + 3)$ , the estimates

$$\|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-2(n+m)(\frac{1}{p}-\frac{1}{2})-1} \|f\|_p, \text{ when } \mu \rightarrow 0^+,$$

and

$$\|\mathcal{P}_\mu^h f\|_{p'} \leq C(1 - \mu)^{2(n+m)(\frac{1}{p}-\frac{1}{2})-1} \|f\|_p, \text{ when } \mu \rightarrow 1^-,$$

hold for all Schwartz functions  $f$ .

More generally, we have:

**Example 5.3.** Let  $h(\xi, \eta) = (\xi^\alpha + \eta^\beta)^\gamma$ ,  $\alpha, \beta, \gamma > 0$ . For  $1 \leq p \leq (2m+2)/(m+3)$ , we have for all Schwartz functions  $f$ :

$$\begin{aligned} \text{if } \alpha < 2\beta & \begin{cases} \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{\frac{2}{\alpha\gamma}(n+\frac{\alpha}{2\beta}m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p, & \mu > 1, \\ \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{\frac{2}{\alpha\gamma}(n+m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p, & 0 < \mu \leq 1, \end{cases} \\ \text{if } \alpha > 2\beta, & \begin{cases} \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{\frac{2}{\alpha\gamma}(n+m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p, & \mu > 1, \\ \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{\frac{2}{\alpha\gamma}(n+\frac{\alpha}{2\beta}m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p, & 0 < \mu \leq 1, \end{cases} \\ \text{if } \alpha = 2\beta & \left\{ \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{\frac{2}{\alpha\gamma}(n+m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p, \quad 0 < \mu < \infty. \right. \end{aligned}$$

**Example 5.4.** Letting  $h(\xi, \eta) = (\xi^\alpha + \eta^\beta)^{-\gamma}$ ,  $\alpha, \beta, \gamma > 0$ , then for  $1 \leq p \leq (2m+2)/(m+3)$ , we have for all Schwartz functions  $f$ :

$$\begin{aligned} \text{if } \alpha < 2\beta & \begin{cases} \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha\gamma}(n+m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p, & \mu > 1, \\ \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha\gamma}(n+\frac{\alpha}{2\beta}m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p, & 0 < \mu \leq 1, \end{cases} \\ \text{if } \alpha > 2\beta & \begin{cases} \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha\gamma}(n+\frac{\alpha}{2\beta}m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p, & \mu > 1, \\ \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha\gamma}(n+m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p, & 0 < \mu \leq 1, \end{cases} \\ \text{if } \alpha = 2\beta & \left\{ \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha\gamma}(n+m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p, \quad 0 < \mu < \infty. \right. \end{aligned}$$

**Example 5.5.** Let  $h(\xi, \eta) = (1 + \xi^\alpha + \eta^\beta)^{-\gamma}$ ,  $\alpha, \beta, \gamma > 0$ . Then for  $1 \leq p \leq (2m+2)/(m+3)$ , we have for all Schwartz functions  $f$ :

$$\begin{aligned} \text{if } \alpha \leq 2\beta & \begin{cases} \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha\gamma}(n+\frac{\alpha}{2\beta}m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p & \text{when } \mu \rightarrow 0^+, \\ \|\mathcal{P}_\mu^h f\|_{p'} \leq C(1-\mu^{\frac{1}{\gamma}})^{\frac{2}{\alpha}(n+m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p & \text{when } \mu \rightarrow 1^-, \end{cases} \\ \text{if } \alpha > 2\beta & \begin{cases} \|\mathcal{P}_\mu^h f\|_{p'} \leq C\mu^{-\frac{2}{\alpha\gamma}(n+m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p & \text{when } \mu \rightarrow 0^+, \\ \|\mathcal{P}_\mu^h f\|_{p'} \leq C(1-\mu^{\frac{1}{\gamma}})^{\frac{2}{\alpha}(n+\frac{\alpha}{2\beta}m)}(\frac{1}{p}-\frac{1}{2})^{-1}\|f\|_p & \text{when } \mu \rightarrow 1^-. \end{cases} \end{aligned}$$

### 6. Sharpness of the range $p$

In this section we only give an example to show that the range of  $p$  in the restriction theorem associated with the full laplacian  $\Delta$  is sharp. The example is constructed similarly to the counterexample of Müller [1990], which shows that the estimates between Lebesgue spaces for the operators  $\mathcal{P}_\mu^\Delta$  are necessarily trivial.

Let  $\varphi \in C_c^\infty(\mathbb{R}^m)$  be a radial function such that  $\varphi(a) = \psi(|a|)$ , where  $\psi \in C_c^\infty(\mathbb{R})$ , with  $\psi = 1$  on a neighborhood of the point  $n$  and  $\psi = 0$  near 0. Let  $h$  be a Schwartz

function on  $\mathbb{R}^m$  and define

$$f(z, t) = \int_{\mathbb{R}^m} \varphi(a) \hat{h}(a) e^{-\frac{|a|}{4}|z|^2} e^{-i\langle a, t \rangle} |a|^n da.$$

Denote

$$\begin{aligned} g(z, t) &= \int_{\mathbb{R}^m} \varphi(a) e^{-\frac{|a|}{4}|z|^2} e^{-i\langle a, t \rangle} |a|^n da \\ &= \int_{\mathbb{R}^{m+2n}} \varphi(a) e^{-\frac{|\xi|^2}{|a|}} e^{-i\langle (a, t) + (\xi, z) \rangle} d\xi da. \end{aligned}$$

Hence  $\widehat{g(\xi, a)} = \varphi(a) e^{-\frac{|\xi|^2}{|a|}}$ , which shows that  $\hat{g}$  and consequently  $g$  are Schwartz functions. On the other hand, we have  $f = h *_t g$ , where  $*_t$  denotes the involution about the central variable. By Lemma 3.1, we have  $\Delta(e^{-i\langle a, t \rangle} e^{-\frac{1}{4}|a||z|^2}) = (n\lambda + \lambda^2) e^{-i\langle a, t \rangle} e^{-\frac{1}{4}|a||z|^2}$ . Therefore, we write  $f$  by the integration with polar coordinates as

$$\begin{aligned} f(z, t) &= \int_0^\infty \left( \lambda^{n+m-1} \psi(\lambda) e^{-\frac{\lambda}{4}|z|^2} \int_{S^{m-1}} \hat{h}(\lambda w) e^{-i\lambda\langle w, t \rangle} d\sigma(w) \right) d\lambda \\ &= \int_0^\infty \left( \lambda_\Delta(\mu)^{n+m-1} \lambda'_\Delta(\mu) \psi(\lambda_\Delta(\mu)) e^{-\frac{\lambda_\Delta(\mu)}{4}|z|^2} \right. \\ &\quad \left. \int_{S^{m-1}} \hat{h}(\lambda_\Delta(\mu) w) e^{-i\lambda_\Delta(\mu)\langle w, t \rangle} d\sigma(w) \right) d\mu \\ &= \int_0^\infty \mathcal{P}_\mu^\Delta f(z, t) d\mu, \end{aligned}$$

where

$$\begin{aligned} \mathcal{P}_\mu^\Delta f(z, t) &= \lambda_\Delta(\mu)^{n+m-1} \lambda'_\Delta(\mu) \psi(\lambda_\Delta(\mu)) e^{-\frac{\lambda_\Delta(\mu)}{4}|z|^2} \\ &\quad \times \int_{S^{m-1}} \hat{h}(\lambda_\Delta(\mu) w) e^{-i\lambda_\Delta(\mu)\langle w, t \rangle} d\sigma(w), \\ \lambda_\Delta(\mu) &= \frac{\sqrt{n^2 + 4\mu} - n}{2}. \end{aligned}$$

Therefore, letting  $\mu = 2n^2$ , we have  $\lambda_\Delta(2n^2) = n$ ,  $\lambda'_\Delta(2n^2) = 1/(3n)$  and

$$\begin{aligned} \mathcal{P}_{2n^2}^\Delta f(z, t) &= \frac{1}{3} n^{n+m-2} e^{-\frac{n|z|^2}{4}} \int_{S^{m-1}} \hat{h}(nw) e^{-in\langle w, t \rangle} d\sigma(w) \\ &= \frac{1}{3} n^{n-1} e^{-\frac{n|z|^2}{4}} h * \widehat{d\sigma_n}(t). \end{aligned}$$

From the restriction theorem associated the full laplacian on H-type groups, we have the estimate  $\|\mathcal{P}_{2n^2}^\Delta f\|_{L^{p'}(G)} \leq C \|f\|_{L^p(G)}$ .

Because of

$$(6-1) \quad \|\mathcal{P}_{2n^2}^\Delta f\|_{L^{p'}(G)} = C \|h * \widehat{d\sigma_n}\|_{L^{p'}(\mathbb{R}^m)}$$

and

$$(6-2) \quad \|f\|_{L^p(G)} \leq \|h\|_{L^p(\mathbb{R}^m)} \|g\|_{L_t^1 L_z^p} \lesssim \|h\|_{L^p(\mathbb{R}^m)},$$

where the mixed Lebesgue norm is defined by

$$\|g\|_{L_t^1 L_z^p} = \left( \int_{\mathbb{R}^{2n}} \left( \int_{\mathbb{R}^m} |f(z, t)| dt \right)^p dz \right)^{\frac{1}{p}},$$

we have  $\|h * \widehat{d\sigma_n}\|_{L^{p'}(\mathbb{R}^m)} \leq C \|h\|_{L^p(\mathbb{R}^m)}$ .

From the sharpness of the Stein–Tomas theorem which is guaranteed by the Knapp counterexample, this would imply  $p \leq (2m + 2)/(m + 3)$ . Hence the range of  $p$  can not be extended. With the same tricks we can prove the range of  $p$  for the restriction theorem associated with the functional calculus is also sharp.

### Acknowledgements

This work was performed while the second author studied as a joint Ph.D. student in the mathematics department of Christian-Albrechts-Universität zu Kiel. She is deeply grateful to Professor Detlef Müller for generous discussions and his continuous encouragement.

### References

- [Bonfiglioli and Uguzzoni 2004] A. Bonfiglioli and F. Uguzzoni, “Nonlinear Liouville theorems for some critical problems on H-type groups”, *J. Funct. Anal.* **207**:1 (2004), 161–215. MR Zbl
- [Casarino and Ciatti 2013a] V. Casarino and P. Ciatti, “Restriction estimates for the full Laplacian on Métivier groups”, *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **24**:2 (2013), 165–179. MR Zbl
- [Casarino and Ciatti 2013b] V. Casarino and P. Ciatti, “A restriction theorem for Métivier groups”, *Adv. Math.* **245** (2013), 52–77. MR Zbl
- [Kaplan and Ricci 1983] A. Kaplan and F. Ricci, “Harmonic analysis on groups of Heisenberg type”, pp. 416–435 in *Harmonic analysis* (Cortona, 1982), edited by G. Manceri et al., Lecture Notes in Math. **992**, Springer, 1983. MR Zbl
- [Liu and Wang 2011] H. Liu and Y. Wang, “A restriction theorem for the H-type groups”, *Proc. Amer. Math. Soc.* **139**:8 (2011), 2713–2720. MR Zbl
- [Müller 1990] D. Müller, “A restriction theorem for the Heisenberg group”, *Ann. of Math. (2)* **131**:3 (1990), 567–587. MR Zbl
- [Ratnakumar et al. 1997] P. K. Ratnakumar, R. Rawat, and S. Thangavelu, “A restriction theorem for the Heisenberg motion group”, *Studia Math.* **126**:1 (1997), 1–12. MR Zbl
- [Reimann 2001] H. M. Reimann, “H-type groups and Clifford modules”, *Adv. Appl. Clifford Algebras* **11**:S2 (2001), 277–287. MR Zbl



- [Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series **43**, Princeton University Press, 1993. MR Zbl
- [Tao 2004] T. Tao, “Some recent progress on the restriction conjecture”, pp. 217–243 in *Fourier analysis and convexity*, edited by L. Brandolini et al., Birkhäuser, Boston, 2004. MR Zbl
- [Thangavelu 1991a] S. Thangavelu, “Restriction theorems for the Heisenberg group”, *J. Reine Angew. Math.* **414** (1991), 51–65. MR Zbl
- [Thangavelu 1991b] S. Thangavelu, “Some restriction theorems for the Heisenberg group”, *Studia Math.* **99**:1 (1991), 11–21. MR Zbl
- [Thangavelu 1993] S. Thangavelu, *Lectures on Hermite and Laguerre expansions*, Mathematical Notes **42**, Princeton University Press, 1993. MR Zbl
- [Thangavelu 2004] S. Thangavelu, *An introduction to the uncertainty principle: Hardy’s theorem on Lie groups*, Progress in Mathematics **217**, Birkhäuser, 2004. MR Zbl
- [Tomas 1975] P. A. Tomas, “A restriction theorem for the Fourier transform”, *Bull. Amer. Math. Soc.* **81** (1975), 477–478. MR

Received October 30, 2013. Revised February 6, 2016.

HEPING LIU  
LMAM, SCHOOL OF MATHEMATICAL SCIENCES  
PEKING UNIVERSITY  
BEIJING, 100871  
CHINA  
hpliu@pku.edu.cn

MANLI SONG  
SCHOOL OF NATURAL AND APPLIED SCIENCES  
NORTHWESTERN POLYTECHNICAL UNIVERSITY  
XI’AN, 710129  
CHINA  
mlsong@nwpu.edu.cn



# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Igor Pak  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pak.pjm@gmail.com](mailto:pak.pjm@gmail.com)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 286 No. 2 February 2017

---

Almost everywhere convergence for modified Bochner–Riesz means at the critical index for $p \geq 2$	257
MARCO ANNONI	
Uniqueness of conformal Ricci flow using energy methods	277
THOMAS BELL	
A functional calculus and restriction theorem on H-type groups	291
HEPING LIU and MANLI SONG	
Identities involving cyclic and symmetric sums of regularized multiple zeta values	307
TOMOYA MACHIDE	
Conformally Kähler Ricci solitons and base metrics for warped product Ricci solitons	361
GIDEON MASCHLER	
Calculating Greene’s function via root polytopes and subdivision algebras	385
KAROLA MÉSZÁROS	
Classifying resolving subcategories	401
WILLIAM SANDERS	
The symplectic plactic monoid, crystals, and MV cycles	439
JACINTA TORRES	
A note on torus actions and the Witten genus	499
MICHAEL WIEMELER	



0030-8730(201702)286:2;1-X