

*Pacific
Journal of
Mathematics*

ON LOCALLY COHERENT HEARTS

MANUEL SAORÍN

Volume 287 No. 1

March 2017

ON LOCALLY COHERENT HEARTS

MANUEL SAORÍN

Let \mathcal{G} be a locally coherent Grothendieck category. We show that, under particular conditions, if a t-structure τ in the unbounded derived category $\mathcal{D}(\mathcal{G})$ restricts to the bounded derived category $\mathcal{D}^b(\text{fp}(\mathcal{G}))$ of its category of finitely presented (i.e. coherent) objects, then its heart \mathcal{H}_τ is a locally coherent Grothendieck category on which $\mathcal{H}_\tau \cap \mathcal{D}^b(\text{fp}(\mathcal{G}))$ is the class of finitely presented objects. Those particular conditions are always satisfied when \mathcal{G} is arbitrary and τ is the Happel–Reiten–Smalø t-structure in $\mathcal{D}(\mathcal{G})$ associated to a torsion pair in $\text{fp}(\mathcal{G})$ or when $\mathcal{G} = \text{Qcoh}(\mathbb{X})$ is the category of quasicoherent sheaves on a noetherian affine scheme \mathbb{X} and τ is any compactly generated t-structure in $\mathcal{D}(\mathbb{X}) := \mathcal{D}(\text{Qcoh}(\mathbb{X}))$ which restricts to $\mathcal{D}^b(\mathbb{X}) := \mathcal{D}^b(\text{coh}(\mathbb{X}))$. In particular, the heart of any t-structure in $\mathcal{D}^b(\mathbb{X})$ is the category of finitely presented objects of a locally coherent Grothendieck category.

1. Introduction

Beilinson, Bernstein and Deligne [1982] introduced the notion of a t-structure in a triangulated category in their study of perverse sheaves on an algebraic or analytic variety. If \mathcal{D} is such a triangulated category, a t-structure is a pair of full subcategories satisfying some axioms which guarantee that their intersection is an abelian category \mathcal{H} , called the heart of the t-structure. This category comes with a cohomological functor $\mathcal{D} \rightarrow \mathcal{H}$. Roughly speaking, a t-structure allows to develop an intrinsic (co)homology theory, where the homology “spaces” are again objects of \mathcal{D} itself.

Nowadays, t-structures are used in several branches of mathematics, with special impact in algebraic geometry, homotopical algebra, and representation theory of groups and algebras. When dealing with t-structures, a natural question arises. It

This work is backed by research projects from the Ministerio de Economía y Competitividad of Spain (MTM201346837-P) and the Fundación “Séneca” of Murcia (19880/GERM/15), both partly supported by FEDER funds. The author thanks these institutions for their support. He also thanks Carlos Parra and the referee for their comments and careful reading of the paper.

MSC2010: 18E15, 18E30, 13DXX, 14AXX, 16EXX.

Keywords: locally coherent Grothendieck category, triangulated category, derived category, t-structure, heart of a t-structure.

asks under which conditions the heart of a given t-structure is a “nice” abelian category. Using a classical hierarchy for abelian categories introduced by Grothendieck, one may think of Grothendieck and module categories as the nicest possible abelian categories. It is therefore not surprising that the question of when the heart of a t-structure is a Grothendieck or module category received much attention in recent times (see, e.g., [Hoshino et al. 2002; Colpi et al. 2007; 2011; Colpi and Gregorio 2010; Mantese and Tonolo 2012; Parra and Saorín 2016b; 2015; Psaroudakis and Vitória 2015; Nicolás et al. 2015]).

Among Grothendieck categories, the most studied ones are those that have finiteness conditions (e.g., those which are locally coherent, locally noetherian or even locally finite). Module categories over noetherian or coherent rings or over Artin algebras, or the categories of quasicohherent sheaves over coherent or noetherian schemes provide examples of such categories. A natural subsequent question would ask when a given t-structure has a heart which is a Grothendieck category with good finiteness conditions. In this paper, we tackle the question for the locally coherent condition, assuming that the t-structure lives in the (unbounded) derived category $\mathcal{D}(\mathcal{G})$ of a Grothendieck category \mathcal{G} which is itself locally coherent. Although to find a general answer seems to be hopeless, it is not so when the t-structure restricts to $\mathcal{D}^b(\text{fp}(\mathcal{G}))$, the bounded derived category of the category of finitely presented (i.e., coherent) objects. Our basic technical result in the paper, [Proposition 4.5](#), gives a precise list of sufficient conditions on a t-structure in $\mathcal{D}(\mathcal{G})$ so that its heart \mathcal{H} is a locally coherent Grothendieck category on which $\mathcal{H} \cap \mathcal{D}^b(\text{fp}(\mathcal{G}))$ is the class of its finitely presented objects. As an application, we give the main results of the paper, referring the reader to the next section for the notation and terminology used:

- (1) ([Theorem 5.2](#)) Let \mathcal{G} be a locally coherent Grothendieck category and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a torsion pair in \mathcal{G} . The associated Happel–Reiten–Smalø t-structure in $\mathcal{D}(\mathcal{G})$ restricts to $\mathcal{D}^b(\text{fp}(\mathcal{G}))$ and has a heart which is a locally coherent Grothendieck category if, and only if, \mathcal{F} is closed under taking direct limits in \mathcal{G} and \mathbf{t} restricts to $\text{fp}(\mathcal{G})$.
- (2) ([Theorem 6.3](#)) If R is a commutative noetherian ring, then any compactly generated t-structure in $\mathcal{D}(R)$ which restricts to $\mathcal{D}_{\text{fg}}^b(R) \cong \mathcal{D}^b(R\text{-mod})$ has a heart \mathcal{H} which is a locally coherent Grothendieck category on which $\mathcal{H} \cap \mathcal{D}_{\text{fg}}^b(R)$ is the class of its finitely presented objects.
- (3) ([Corollary 6.4](#)) If R is a commutative noetherian ring, then the heart of each t-structure in $\mathcal{D}_{\text{fg}}^b(R)$ is equivalent to the category of finitely presented objects of some locally coherent Grothendieck category.

Of course, when taking the affine scheme $\mathbb{X} = \text{Spec } R$ in (2) and (3), one obtains the geometric versions mentioned in the abstract (see also [Corollary 6.5](#)).

The organization of the paper is as follows. [Section 2](#) introduces all the concepts and terminology used in the paper. In [Section 3](#) we give some general results about locally coherent Grothendieck categories which are used later. [Section 4](#) contains the technical [Proposition 4.5](#), which is central to the paper, and a few auxiliary results needed for its proof. [Section 5](#) is dedicated to the Happel–Reiten–Smalø t-structure and the proof of [Theorem 5.2](#). The final [Section 6](#) gives [Theorem 6.3](#), of which [Corollary 6.4](#) is a direct consequence, and two lemmas needed for its proof.

2. Preliminaries and terminology

All categories in this paper will be additive and all rings will be supposed to be associative with unit, unless otherwise specified. Whenever the term “module” is used over a noncommutative ring, it will mean “left module” and, for a given ring R , we will denote by $R\text{-Mod}$ the category of all R -modules. Let \mathcal{A} be an additive category in the rest of the paragraph. If \mathcal{C} is any class of objects in \mathcal{A} , the symbol \mathcal{C}^\perp (resp. ${}^\perp\mathcal{C}$) will denote the full subcategory of \mathcal{A} whose objects are those $X \in \text{Ob}(\mathcal{A})$ such that $\text{Hom}_{\mathcal{A}}(C, X) = 0$ (resp. $\text{Hom}_{\mathcal{A}}(X, C) = 0$), for all $C \in \mathcal{C}$. The expression “ \mathcal{A} has products (resp. coproducts)” will mean that \mathcal{A} has arbitrary set-indexed products (coproducts). If \mathcal{S} is a set of objects in \mathcal{A} , we denote by $\text{sum}(\mathcal{S})$ the class of objects which are isomorphic to a finite coproduct of objects of \mathcal{S} , and by $\text{add}(\mathcal{S})$ the class of objects isomorphic to a direct summand of a finite coproduct of objects of \mathcal{S} . When \mathcal{A} has coproducts, we shall say that an object X is a *compact* (or *small*) *object* when the functor $\text{Hom}_{\mathcal{A}}(X, ?) : \mathcal{A} \rightarrow \text{Ab}$ preserves coproducts.

Two types of additive categories will get most of our interest in this paper: *abelian categories* (see [[Popescu 1973](#)]) and *triangulated categories* (see [[Neeman 2001](#)]). Diverting from the terminology in this latter reference, for a triangulated category \mathcal{D} , the shift or suspension functor will be denoted by $?[1]$, putting $?[k]$ for its k -th power, for each $k \in \mathbb{Z}$. We shall use the term *class* (resp. *set*) of *generators* with two different meanings, depending on whether we are in the abelian or the triangulated context. When \mathcal{A} is an abelian category with coproducts, a class (resp. set) of generators \mathcal{S} is a class (set) of objects such that each object in \mathcal{A} is an epimorphic image of a coproduct of objects in \mathcal{S} . When \mathcal{S} is a class (set) of objects in the triangulated category \mathcal{D} , we shall say that it is a class (set) of generators if an object X of \mathcal{D} is zero exactly when $\text{Hom}_{\mathcal{D}}(\mathcal{S}[k], X) = 0$, for all $S \in \mathcal{S}$ and all $k \in \mathbb{Z}$.

Given a triangulated category \mathcal{D} , a subcategory \mathcal{E} will be called a *triangulated subcategory* when it is closed under taking extensions and $\mathcal{E}[1] = \mathcal{E}$. If, in addition, it is closed under taking direct summands, we will say that \mathcal{E} is a *thick subcategory* of \mathcal{D} . When \mathcal{S} is a set of objects of \mathcal{D} , we shall denote by $\text{tria}_{\mathcal{D}}(\mathcal{S})$ the smallest triangulated subcategory of \mathcal{D} that contains \mathcal{S} , and by $\text{thick}_{\mathcal{D}}(\mathcal{S})$ the smallest thick

subcategory of \mathcal{D} that contains \mathcal{S} .

For an additive category \mathcal{A} , we will denote by $\mathcal{C}(\mathcal{A})$ and $\mathcal{K}(\mathcal{A})$ the category of chain complexes of objects of \mathcal{A} and the homotopy category of \mathcal{A} . Diverting from the classical notation, we will write superindices for chains, cycles and boundaries in ascending order. We will denote by $\mathcal{C}^-(\mathcal{A})$, $\mathcal{C}^+(\mathcal{A})$, and $\mathcal{C}^b(\mathcal{A})$ the full subcategories of $\mathcal{C}(\mathcal{A})$ consisting of those objects isomorphic to upper bounded, lower bounded, and upper and lower bounded complexes, respectively, and similarly for $\mathcal{K}(\mathcal{A})$, $\mathcal{K}^-(\mathcal{A})$, $\mathcal{K}^+(\mathcal{A})$, and $\mathcal{K}^b(\mathcal{A})$. Note that $\mathcal{K}(\mathcal{A})$ is always a triangulated category of which $\mathcal{K}^-(\mathcal{A})$, $\mathcal{K}^+(\mathcal{A})$ and $\mathcal{K}^b(\mathcal{A})$ are triangulated subcategories. When \mathcal{A} is an abelian category, we will denote by $\mathcal{D}(\mathcal{A})$ its *derived category*, which is the one obtained from $\mathcal{C}(\mathcal{A})$ by keeping the same objects and formally inverting the quasi-isomorphisms (see [Verdier 1996] for the details). We shall denote by $\mathcal{D}^-(\mathcal{A})$, $\mathcal{D}^+(\mathcal{A})$, and $\mathcal{D}^b(\mathcal{A})$ the full subcategory of $\mathcal{D}(\mathcal{A})$ consisting of those complexes X such that $H^k(X) = 0$, for all $k \gg 0$, $k \ll 0$, and $|k| \gg 0$, respectively, where $H^k : \mathcal{D}(\mathcal{A}) \rightarrow \mathcal{A}$ denotes the k -th homology functor, for each $k \in \mathbb{Z}$. The objects of $\mathcal{D}^-(\mathcal{A})$ (resp. $\mathcal{D}^+(\mathcal{A})$, $\mathcal{D}^b(\mathcal{A})$) will be called *homologically upper bounded* (resp. *homologically lower bounded*, *homologically bounded*) complexes. For integers $m \leq n$, we will denote by $\mathcal{D}^{[m,n]}(\mathcal{A})$ the full subcategory of $\mathcal{D}(\mathcal{A})$ consisting of the complexes X such that $H^k(X) = 0$ for integers k not in the closed interval $[m, n]$. We will also use $\mathcal{D}^{\leq n}(\mathcal{A})$, $\mathcal{D}^{< n}(\mathcal{A})$, $\mathcal{D}^{\geq n}(\mathcal{A})$, and $\mathcal{D}^{> n}(\mathcal{A})$ to denote the full subcategories consisting of the complexes X such that $H^i(X) = 0$, for all $i > n$, $i \geq n$, $i < n$, and $i \leq n$, respectively.

Strictly speaking, for a general abelian category \mathcal{A} , the category $\mathcal{D}(\mathcal{A})$ need not exist since the morphisms between two given objects could form a proper class and not just a set. However, this problem disappears when $\mathcal{A} = \mathcal{G}$ is a *Grothendieck category*. This is a cocomplete abelian category with a set of generators on which direct limits are exact. In a Grothendieck category \mathcal{G} an object S is called *finitely presented* when $\text{Hom}_{\mathcal{G}}(S, ?) : \mathcal{G} \rightarrow \text{Ab}$ preserves direct limits. We say that \mathcal{G} is *locally finitely presented* when it has a set of finitely presented generators. The reader is referred to [Crawley-Boevey 1994] for the corresponding more general concept of locally finitely presented additive categories with direct limits and is invited to check that, in the case of Grothendieck categories, it coincides with the one given here. Recall that an object in a Grothendieck category is called *noetherian* when it satisfies the ascending chain condition on subobjects. A *locally noetherian* Grothendieck category is a Grothendieck category which has a set of noetherian generators. When \mathcal{G} is locally finitely presented and locally noetherian, an object N of \mathcal{G} is noetherian if and only if it is finitely presented. (See [Krause 1997, Proposition A.11] for one direction, the reverse one being obvious since each noetherian object in such a category is an epimorphic image of a finitely presented one and the kernel of this epimorphism is again noetherian.)

Recall that if \mathcal{D} and \mathcal{A} are a triangulated and an abelian category, respectively, then an additive functor $H : \mathcal{D} \rightarrow \mathcal{A}$ is a *cohomological functor* when, given any triangle $X \rightarrow Y \rightarrow Z \xrightarrow{+}$, one gets an induced long exact sequence in \mathcal{A} :

$$\dots \rightarrow H^{n-1}(Z) \rightarrow H^n(X) \rightarrow H^n(Y) \rightarrow H^n(Z) \rightarrow H^{n+1}(X) \rightarrow \dots,$$

where $H^n := H \circ (?[n])$, for each $n \in \mathbb{Z}$.

A *torsion pair* in the abelian category \mathcal{A} is a pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ of full subcategories such that $\text{Hom}_{\mathcal{A}}(T, F) = 0$, for all $T \in \mathcal{T}$ and $F \in \mathcal{F}$, and each object X of \mathcal{A} fits into an exact sequence $0 \rightarrow T_X \rightarrow X \rightarrow F_X \rightarrow 0$, where $T_X \in \mathcal{T}$ and $F_X \in \mathcal{F}$. In this latter case the assignments $X \rightsquigarrow T_X$ and $X \rightsquigarrow F_X$ extend to endofunctors $t, (1 : t) : \mathcal{A} \rightarrow \mathcal{A}$. The functor t is usually called the *torsion radical* associated to \mathbf{t} . The torsion pair \mathbf{t} will be called *hereditary* when \mathcal{T} is closed under taking subobjects in \mathcal{A} .

Now let \mathcal{D} be a triangulated category. A *t-structure* in \mathcal{D} (see [Beilinson et al. 1982, Section 1]) is a pair $\tau = (\mathcal{U}, \mathcal{W})$ of full subcategories, closed under taking direct summands in \mathcal{D} , which satisfy the following properties:

- (i) $\text{Hom}_{\mathcal{D}}(U, W[-1]) = 0$, for all $U \in \mathcal{U}$ and $W \in \mathcal{W}$.
- (ii) $\mathcal{U}[1] \subseteq \mathcal{U}$.
- (iii) For each $X \in \text{Ob}(\mathcal{D})$, there is a triangle $U \rightarrow X \rightarrow V \xrightarrow{+}$ in \mathcal{D} , where $U \in \mathcal{U}$ and $V \in \mathcal{W}[-1]$.

In this case $\mathcal{W} = \mathcal{U}^\perp[1]$ and $\mathcal{U} = {}^\perp(\mathcal{W}[-1]) = {}^\perp(\mathcal{U}^\perp)$ and, for this reason, we will write a t-structure as $\tau = (\mathcal{U}, \mathcal{U}^\perp[1])$. We will call \mathcal{U} and \mathcal{U}^\perp the *aisle* and the *co-aisle* of the t-structure. The objects U and V in the above triangle are uniquely determined by X , up to isomorphism, and define functors $\tau_{\mathcal{U}} : \mathcal{D} \rightarrow \mathcal{U}$ and $\tau^{\mathcal{U}^\perp} : \mathcal{D} \rightarrow \mathcal{U}^\perp$ which are right and left adjoints to the respective inclusion functors. We call them the *left and right truncation functors* with respect to the given t-structure. The full subcategory $\mathcal{H} = \mathcal{U} \cap \mathcal{W} = \mathcal{U} \cap \mathcal{U}^\perp[1]$ is called the *heart* of the t-structure and it is an abelian category, where the short exact sequences “are” the triangles in \mathcal{D} with its three terms in \mathcal{H} . Moreover, with the obvious abuse of notation, the assignments $X \rightsquigarrow (\tau_{\mathcal{U}} \circ \tau^{\mathcal{U}^\perp[1]})(X)$ and $X \rightarrow (\tau^{\mathcal{U}^\perp[1]} \circ \tau_{\mathcal{U}})(X)$ define naturally isomorphic functors $\mathcal{D} \rightarrow \mathcal{H}$ which are cohomological (see [Beilinson et al. 1982]). We will identify them and denote the corresponding functor by \tilde{H} . When \mathcal{D} has coproducts, the t-structure τ will be called *compactly generated* when there is a set $\mathcal{S} \subseteq \mathcal{U}$, formed by compact objects in \mathcal{D} , such that $\mathcal{W}[-1] = \mathcal{U}^\perp$ consists of the objects Y such that $\text{Hom}_{\mathcal{D}}(\mathcal{S}[k], Y) = 0$, for all $S \in \mathcal{S}$ and integers $k \geq 0$.

When \mathcal{D} is a triangulated category with coproducts, we will use the term *Milnor colimit* of a sequence of morphisms $X_0 \xrightarrow{x_1} X_1 \xrightarrow{x_2} \dots \xrightarrow{x_n} X_n \xrightarrow{x_{n+1}} \dots$, which in [Neeman 2001] is called the homotopy colimit. It will be denoted $\text{Mcolim}(X_n)$, without reference to the x_n .

3. Generalities about locally coherent Grothendieck categories

In this section we are interested in a particular case of locally finitely presented Grothendieck categories. Let us start with the following result which is folklore.

Lemma 3.1. *Let \mathcal{A} be an abelian category and \mathcal{B} be a full additive subcategory. The following assertions are equivalent:*

- (1) \mathcal{B} is an abelian category such that the inclusion functor $\mathcal{B} \hookrightarrow \mathcal{A}$ is exact.
- (2) \mathcal{B} is closed under taking finite (co)products, kernels and cokernels in \mathcal{A} .

In this case we will say that \mathcal{B} is an abelian exact subcategory of \mathcal{A} .

Note that if \mathcal{G} is a locally finitely presented Grothendieck category, then the class $\text{fp}(\mathcal{G})$ of finitely presented objects is skeletally small and is closed under taking cokernels and finite coproducts.

Definition. A Grothendieck category \mathcal{G} is called *locally coherent* when it is locally finitely presented and the subcategory $\text{fp}(\mathcal{G})$ is an abelian exact subcategory of \mathcal{G} (equivalently, when $\text{fp}(\mathcal{G})$ is closed under taking kernels).

Recall that a *pseudokernel* of a morphism $f : X \rightarrow Y$ in the additive category \mathcal{A} is a morphism $u : Z \rightarrow X$ such that the sequence of contravariant functors $\text{Hom}_{\mathcal{A}}(?, Z) \xrightarrow{u_*} \text{Hom}_{\mathcal{A}}(?, X) \xrightarrow{f_*} \text{Hom}_{\mathcal{A}}(?, Y)$ is exact, and similarly, a *pseudocokernel* of a morphism $f : X \rightarrow Y$ in the additive category \mathcal{A} is a morphism $v : Y \rightarrow Z$ such that the sequence of covariant functors

$$\text{Hom}_{\mathcal{A}}(Z, ?) \xrightarrow{v^*} \text{Hom}_{\mathcal{A}}(Y, ?) \xrightarrow{f^*} \text{Hom}_{\mathcal{A}}(X, ?)$$

is exact. We say that \mathcal{A} has *pseudokernels* (resp. *pseudocokernels*) when each morphism in \mathcal{A} has a pseudokernel (pseudocokernel).

Examples 3.2. Here are some locally coherent Grothendieck categories to which the results in this and next section apply. The first is well-known; for the others we provide a brief justification.

1. $R\text{-Mod}$, when R is a left coherent ring R (i.e., when each finitely generated left ideal of R is finitely presented).
2. The category $[\mathcal{C}, \text{Ab}]$ (resp. $[\mathcal{C}^{\text{op}}, \text{Ab}]$) of covariant (resp. contravariant) additive functors $\mathcal{C} \rightarrow \text{Ab}$, where \mathcal{C} is a skeletally small additive category with pseudocokernels (pseudokernels). In particular, when \mathcal{C} is a skeletally small abelian or triangulated category, both $[\mathcal{C}, \text{Ab}]$ and $[\mathcal{C}^{\text{op}}, \text{Ab}]$ are locally coherent Grothendieck categories.

The covariant version follows from Propositions 1.3 and 2.1 of [Herzog 1997], taking into account that, in the second of these, the proof that each representable functor $(X, ?)$ is a coherent object only requires that each morphism $X \rightarrow Y$ has a pseudocokernel. The contravariant version follows by duality.

3. The category $\text{Qcoh}(\mathbb{X})$ of quasicoherent sheaves, where \mathbb{X} is a *coherent scheme*, i.e., a quasicompact and quasiseparated scheme admitting a covering $\mathbb{X} = \bigcup_{i \in I} U_i$ by affine open subschemes U_i such that $U_i = \text{Spec } A_i$, for a commutative coherent ring A_i , for each $i \in I$.

For the proof, see [Garkusha 2009, Proposition 40], and also [Sitte 2014, Example 1.1.6.iv].

4. Any locally noetherian and locally finitely presented Grothendieck category.

This is clear, since $\text{fp}(\mathcal{G})$ coincides with the class of noetherian objects in that case, and this latter class is always closed under taking kernels (even subobjects).

Lemma 3.3. *Let \mathcal{G} be a locally coherent Grothendieck category, let \mathcal{S} be a set of finitely presented generators of \mathcal{G} and let M be any object in $\mathcal{D}(\mathcal{G})$. The following assertions hold:*

- (1) *M is a homologically upper bounded complex whose homology objects are finitely presented if, and only if, M is isomorphic in $\mathcal{D}(\mathcal{G})$ to an upper bounded complex N of objects in $\text{sum}(\mathcal{S})$. Moreover, N can be chosen such that $\max\{i \in \mathbb{Z} : N^i \neq 0\} = \max\{i \in \mathbb{Z} : H^i(M) \neq 0\}$.*
- (2) *M is homologically bounded and its homology objects are finitely presented if, and only if, M is isomorphic in $\mathcal{D}(\mathcal{G})$ to a bounded complex*

$$\dots 0 \rightarrow N^m \rightarrow N^{m+1} \rightarrow N^{n-1} \rightarrow N^n \rightarrow 0 \dots,$$

where the N^i are finitely presented objects (and $N^i \in \text{sum}(\mathcal{S})$, for $m < i \leq n$).

If, moreover, the objects of \mathcal{S} form a set of compact generators of $\mathcal{D}(\mathcal{G})$, then the following assertion also holds:

- (3) *The compact objects of $\mathcal{D}(\mathcal{G})$ are those isomorphic to direct summands of bounded complexes of objects in $\text{add}(\mathcal{S})$.*

Proof. We will frequently use the fact that if M is a complex whose homology objects are all finitely presented, then a given k -cycle object $Z^k = Z^k(M)$ is finitely presented if and only if so is the k -boundary object $B^k = B^k(M)$.

(1) The proof of this assertion is reminiscent of the dual of the proof of Lemma 4.6(3) in [Hartshorne 1966, Chapter I], with $\mathcal{A}' = \text{fp}(\mathcal{G})$ and $\mathcal{A} = \mathcal{G}$, although the assumptions of that lemma do not hold in our situation. By truncating at the greatest integer i such that $H^i(M) \neq 0$ and shifting if necessary, we can assume without loss of generality that M is concentrated in degrees ≤ 0 and that $H^0(M) \neq 0$. We then inductively construct a sequence in $\mathcal{C}(\mathcal{G})$

$$\dots M_n \xrightarrow{f_n} M_{n-1} \rightarrow \dots \rightarrow M_1 \xrightarrow{f_1} M_0 \xrightarrow{f_0} M$$

satisfying the following properties:

- (a) Each M_n is concentrated in degrees ≤ 0 .
- (b) The connecting chain maps $f_n : M_n \rightarrow M_{n-1}$ are quasi-isomorphisms, for all $n \in \mathbb{N}$ (with the convention that $M_{-1} = M$).
- (c) Given $n \in \mathbb{N}$, one has $M_n^{-k} \in \text{sum}(\mathcal{S})$ for $0 \leq k \leq n$.
- (d) Given any $k \in \mathbb{N}$, the morphism $f_n^{-k} : M_n^{-k} \rightarrow M_{n-1}^{-k}$ is the identity map, for all $n > k$.

Once the sequence has been constructed, we clearly see that the inverse limit of the sequence, $X := \varprojlim_{C(\mathcal{G})} (M_n)$, is a complex of objects in $\text{sum}(\mathcal{S})$ concentrated in degrees ≤ 0 such that the induced chain map $X \rightarrow M$ is a quasi-isomorphism.

We now move on to construct the mentioned sequence. At the initial step, one easily gets a morphism $f : X^0 \rightarrow M^0$ such that $X^0 \in \text{add}(\mathcal{S})$ and the composition $X^0 \xrightarrow{f} M^0 \xrightarrow{p} H^0(M)$ is an epimorphism, where p is the projection. Now, taking the pullback of f and the differential $M^{-1} \rightarrow M^0$, we easily get a quasi-isomorphism $f_0 : M_0 \rightarrow M$, where $f_0^{-k} : M_0^{-k} = M^{-k} \rightarrow M^{-k}$ is the identity map for all $k \geq 2$, and $f_0^0 : M_0^0 = X^0 \rightarrow M^0$ is f .

Assume now that $n > 0$ and that the quasi-isomorphisms

$$M_{n-1} \xrightarrow{f_{n-1}} M_{n-2} \rightarrow \cdots \rightarrow M_1 \xrightarrow{f_1} M_0 \xrightarrow{f_0} M$$

have already been constructed, satisfying the requirements. Note that $Z^{-k} := Z^{-k}(M_{n-1})$, and hence also $B^{-k} := B^{-k}(M_{n-1})$, are finitely presented objects for $k = 0, 1, \dots, n-1$. Let us fix a direct system $(Y_i)_{i \in I}$ in $\text{fp}(\mathcal{G})$ such that $\varinjlim Y_i \cong M_{n-1}^{-n}$. Replacing the directed set I by a cofinal subset if necessary, there is no loss of generality in assuming that the composition

$$Y_j \xrightarrow{u_j} \varinjlim Y_i \cong M_{n-1}^{-n} \xrightarrow{d^{-n}} B^{-n+1}$$

is an epimorphism, for all $j \in I$, where u_j is the canonical morphism to the direct limit. It is seen in a straightforward way that we have a direct system of exact sequences

$$0 \rightarrow u_i^{-1}(Z^{-n}) \rightarrow Y_i \xrightarrow{d^{-n} \circ u_i} B^{-n+1} \rightarrow 0 \quad (i \in I)$$

whose direct limit is precisely the canonical exact sequence

$$0 \rightarrow Z^{-n} \rightarrow X^{-n} \xrightarrow{d^{-n}} B^{-n+1} \rightarrow 0.$$

Due to the fact that $H^{-n} := H^{-n}(M_{n-1})$ is finitely presented, there is some index $j \in I$ such that the composition $u_j^{-1}(Z^{-n}) \xrightarrow{u_j} Z^{-n} \xrightarrow{p} H^{-n}$ is an epimorphism. We fix such an index j and choose any epimorphism $\epsilon : X^{-n} \twoheadrightarrow Y_j$, with $X^{-n} \in \text{sum}(\mathcal{S})$. Putting $M_n^{-n} := X^{-n}$, the composition $g : M_n^{-n} \xrightarrow{\epsilon} Y_j \xrightarrow{u_j} \varinjlim (Y_i) \cong M_{n-1}^{-n}$ is then a morphism which leads to the following commutative diagram, where all squares are bicartesian:

$$\begin{array}{ccccccc}
M_n^{-n-1} & \longrightarrow & \widetilde{B}^{-n} & \hookrightarrow & g^{-1}(Z^{-n}) & \hookrightarrow & M_n^{-n} \\
\downarrow g' & & \downarrow & & \downarrow & & \downarrow g \\
M_{n-1}^{-n-1} & \xrightarrow{d^{-n-1}} & B^{-n} & \hookrightarrow & Z^{-n} & \hookrightarrow & M_{n-1}^{-n}
\end{array}$$

We derive a quasi-isomorphism $h : M_n \rightarrow M_{n-1}$, where $h^{-k} : M_n^{-k} = M_{n-1}^{-k} \rightarrow M_{n-1}^{-k}$ is the identity map, for $k \geq 0$ and $k \neq n, n+1$, and where $h^{-n-1} = g'$ and $h^{-n} = g$ are the morphisms from the last diagram.

(2) By assertion (1), we can assume that M is of the form

$$\dots \rightarrow N^k \rightarrow N^{k+1} \rightarrow \dots \rightarrow N^{n-1} \rightarrow N^n \rightarrow 0 \dots,$$

where the N^i are in $\text{sum}(\mathcal{S})$. Let us assume that $m = \min\{j \in \mathbb{Z} : H^j(M) \neq 0\}$. Then the intelligent truncation at m gives the complex

$$\tau^{\geq m} M : \dots 0 \rightarrow B^m \hookrightarrow N^m \rightarrow N^{m+1} \rightarrow \dots \rightarrow N^{n-1} \rightarrow N^n \rightarrow 0 \dots,$$

where B^m is an m -boundary object of M . But B^m is finitely presented because $Z^m = \text{Ker}(N^m \rightarrow N^{m+1})$ is. We then take $N^m = B^m$ and the proof of the implication is complete because the canonical map $\tau^{\geq m} M \rightarrow M$ is an isomorphism in $\mathcal{D}(\mathcal{G})$.

In the rest of the proof, we assume that \mathcal{S} is a set of compact generators of $\mathcal{D}(\mathcal{G})$.

(3) Note that each bounded complex of objects in $\text{add}(\mathcal{S})$ is compact in $\mathcal{D}(\mathcal{G})$ since it is a finite iterated extension of stalks $X[k]$, with $X \in \text{add}(\mathcal{S})$. Conversely, suppose that M is a compact object in $\mathcal{D}(\mathcal{G})$. It follows from [Keller 1994, Theorem 5.3] that it is a direct summand of a finite iterated extension of complexes of the form $S[k]$, with $S \in \mathcal{S}$ and $k \in \mathbb{Z}$. In particular M has bounded and finitely presented homology. If we fix now a quasi-isomorphism $f : P \rightarrow M$ such that P is a bounded above complex of objects in $\text{add}(\mathcal{S})$, then we can assume without loss of generality that $P^0 \neq 0 = P^k$, for all $k > 0$. Note that then P is the Milnor colimit of the stupid truncations $\sigma^{\geq -n} P$. Since P is compact in $\mathcal{D}(\mathcal{G})$, an argument as in the proof of [Keller 1994, Theorem 5.3] shows that the identity map 1_P factors in the form $P \rightarrow \sigma^{\geq -n} P \rightarrow P$, for some $n \in \mathbb{N}$. It follows that $M \cong P$ is isomorphic in $\mathcal{D}(\mathcal{G})$ to a direct summand of a bounded complex of objects in $\text{add}(\mathcal{S})$. \square

When \mathcal{G} is a locally coherent Grothendieck category, one easily gets from assertions (1) and (2) of the last lemma that $\mathcal{D}^b(\text{fp}(\mathcal{G}))$ is equivalent, as a triangulated category, to the full subcategory $\mathcal{D}_{\text{fp}}^b(\mathcal{G})$ of $\mathcal{D}(\mathcal{G})$ consisting of those complexes $M \in \mathcal{D}^b(\mathcal{G})$ such that $H^i(M) \in \text{fp}(\mathcal{G})$, for all $i \in \mathbb{Z}$. In the sequel we will identify these equivalent triangulated categories, viewing $\mathcal{D}^b(\text{fp}(\mathcal{G}))$ as a full triangulated subcategory of $\mathcal{D}(\mathcal{G})$.

Definition. Let \mathcal{G} be a locally finitely presented Grothendieck category. An object Y of \mathcal{G} will be called *fp-injective* when $\text{Ext}_{\mathcal{G}}^1(?, Y)$ vanishes on finitely presented objects.

The following is an easy consequence of the proof of implication $1) \Rightarrow 2)$ in [Šťovíček 2014, Proposition B.3], after a clear induction argument:

Lemma 3.4. *Let \mathcal{G} be a locally coherent Grothendieck category. If Y is an fp-injective object of \mathcal{G} , then $\text{Ext}_{\mathcal{G}}^k(?, Y)$ vanishes on finitely presented objects, for all $k > 0$.*

Recall that if $F : \mathcal{G} \rightarrow \text{Ab}$ is any left exact functor, then an object Y of \mathcal{G} is *F-acyclic* when the right derived functors $\mathbf{R}^k F : \mathcal{G} \rightarrow \text{Ab}$ vanish on Y , for all $k > 0$. Recall also that, for each $X \in \text{Ob}(\mathcal{G})$, one can calculate $\mathbf{R}^k F(X)$ by considering *F-acyclic* resolutions. That is, if one picks an exact sequence $0 \rightarrow X \rightarrow Y^0 \xrightarrow{d^0} Y^1 \xrightarrow{d^1} \dots Y^n \xrightarrow{d^n} \dots$, where all the Y^k are *F-acyclic*, then $\mathbf{R}^k F(X)$ is the k -th homology group of the complex

$$\dots 0 \rightarrow F(Y^0) \xrightarrow{F(d^0)} F(Y^1) \xrightarrow{F(d^1)} \dots F(Y^n) \xrightarrow{F(d^n)} \dots,$$

for each integer $k \geq 0$. The following result seems to be well-known (see [Gillespie 2016, Introduction] or [Prest 2009, Chapter 11]), but we include a proof after not finding an explicit one in the literature.

Proposition 3.5. *Let \mathcal{G} be a locally finitely presented Grothendieck category, let X be a finitely presented object, let $(M_i)_{i \in I}$ be a direct system in \mathcal{G} and consider the canonical map $\mu_k : \varinjlim \text{Ext}_{\mathcal{G}}^k(X, M_i) \rightarrow \text{Ext}_{\mathcal{G}}^k(X, \varinjlim M_i)$, for each integer $k \geq 0$.*

- (1) μ_0 is an isomorphism and μ_1 is a monomorphism.
- (2) When \mathcal{G} is locally coherent, μ_k is an isomorphism, for all $k \geq 0$.

Proof. (1) The case $k = 0$ follows from the definition of a finitely presented object. An element of $\varinjlim \text{Ext}_{\mathcal{G}}^1(X, M_i)$ is represented by a direct system $(\epsilon_i)_{i \in I}$ of exact sequences

$$\epsilon_i : 0 \rightarrow M_i \rightarrow N_i \rightarrow X \rightarrow 0$$

whose “projection” on the first component is precisely the direct system $(M_i)_{i \in I}$ and where X is viewed as a constant direct system. The image of (ϵ_i) by the canonical map $\varinjlim \text{Ext}_{\mathcal{G}}^1(X, M_i) \rightarrow \text{Ext}_{\mathcal{G}}^1(X, \varinjlim M_i)$ is the induced exact sequence

$$0 \rightarrow \varinjlim M_i \rightarrow \varinjlim N_i \xrightarrow{\pi} X \rightarrow 0.$$

If this latter sequence splits and we fix a section $\mu : X \rightarrow \varinjlim N_i$ for π , then, since X is a finitely presented object, μ factors in the form $X \xrightarrow{\mu_j} N_j \xrightarrow{u_j} \varinjlim N_i$, for some $j \in I$, where u_j is the canonical morphism to the direct limit. This immediately

implies that the j -th sequence $\epsilon_j : 0 \rightarrow M_j \rightarrow N_j \rightarrow X \rightarrow 0$ splits and, hence, that $(\epsilon_i)_{i \in I}$ is the zero element of $\varinjlim \text{Ext}_{\mathcal{G}}^1(X, M_i)$.

(2) By [Adámek and Rosický 1994, Corollary 1.7 and subsequent remark], we can assume without loss of generality that $I = \lambda = \{\alpha \text{ ordinal} : \alpha < \lambda\}$ is an infinite limit ordinal and that, for each limit ordinal $\alpha < \lambda$, one has $M_\alpha = \varinjlim_{\beta < \alpha} M_\beta$. We now construct a direct system $(E_\alpha)_{\alpha < \lambda}$ in the category $\mathcal{C}(\mathcal{G})$ of complexes, satisfying the following properties:

- (a) $E_\alpha : \cdots \rightarrow E_\alpha^0 \rightarrow E_\alpha^1 \rightarrow \cdots \rightarrow E_\alpha^n \rightarrow \cdots$ is a complex concentrated in degrees ≥ 0 and $H^k(E_\alpha) = 0$, for all $\alpha < \lambda$ and all $k \neq 0$.
- (b) E_α^n is an fp-injective object, for all $\alpha < \lambda$ and all integers $n \geq 0$.
- (c) The direct system $(H^0(E_\alpha))_{\alpha < \lambda}$ in \mathcal{G} is isomorphic to $(M_\alpha)_{\alpha < \lambda}$.

Once the direct system $(E_\alpha)_{\alpha < \lambda}$ is constructed, the exactness of the direct limit functor in \mathcal{G} and the fact that the class of fp-injective objects is closed under taking direct limits (see [Šťovíček 2014, Proposition B.3]) will give that $E_\lambda := \varinjlim_{\mathcal{C}(\mathcal{G})} E_\alpha$ is a complex of fp-injective objects concentrated in degrees ≥ 0 whose only nonzero homology object is $H^0(E_\lambda) \cong \varinjlim_{\alpha < \lambda} M_\alpha$. That is, E_λ is a (deleted) fp-injective resolution of $M := \varinjlim_{\alpha < \lambda} M_\alpha$. By the previous lemma, we know that each fp-injective object is $\text{Hom}_{\mathcal{G}}(X, ?)$ -acyclic, whenever $X \in \text{fp}(\mathcal{G})$. It follows that, for such an X , we have that $\text{Ext}_{\mathcal{G}}^k(X, M)$ is the k -th homology abelian group of the complex $\text{Hom}_{\mathcal{G}}(X, E_\lambda)$. But, by definition of E_λ and the fact that $\text{Hom}_{\mathcal{G}}(X, ?)$ preserves direct limits, we have an isomorphism of complexes of abelian groups $\varinjlim_{\mathcal{C}(\text{Ab})} (\text{Hom}_{\mathcal{G}}(X, E_\alpha)) \cong \text{Hom}_{\mathcal{G}}(X, E_\lambda)$. Then the k -th homology map will give the desired isomorphism

$$\varinjlim \text{Ext}_{\mathcal{G}}^k(X, M_\alpha) \xrightarrow{\cong} \text{Ext}_{\mathcal{G}}^k(X, M) = \text{Ext}_{\mathcal{G}}^k(X, \varinjlim_{\alpha < \lambda} M_\alpha).$$

It remains to construct the direct system $(E_\alpha)_{\alpha < \lambda}$ in $\mathcal{C}(\mathcal{G})$. Let $u_\alpha : M_\alpha \rightarrow M_{\alpha+1}$ denote the morphism from the direct system $(M_\alpha)_{\alpha < \lambda}$. For a nonlimit ordinal α , E_α will be the (deleted) minimal injective resolution of M_α . If α is a limit ordinal and we already have defined the direct system $(E_\beta)_{\beta < \alpha}$, then $E_\alpha = \varinjlim_{\beta < \alpha} E_\beta$, where the direct limit is taken in $\mathcal{C}(\mathcal{G})$. Note that $H^0(E_\alpha) \cong \varinjlim_{\beta < \alpha} M_\beta = M_\alpha$. For the construction of $(E_\alpha)_{\alpha < \lambda}$ one just needs to define the connecting chain map $E_\alpha \rightarrow E_{\alpha+1}$, when $\alpha < \lambda$ is any ordinal for which E_α is already defined. This connecting chain map is defined by choosing a family $(f_\alpha^n : E_\alpha^n \rightarrow E_{\alpha+1}^n)_{n \geq 0}$ of morphisms in \mathcal{G} such that the following diagram is commutative and the induced

map $\text{Ker}(E_\alpha^0 \rightarrow E_\alpha^1) \rightarrow \text{Ker}(E_{\alpha+1}^0 \rightarrow E_{\alpha+1}^1)$ is the morphism $u_\alpha : M_\alpha \rightarrow M_{\alpha+1}$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & E_\alpha^0 & \longrightarrow & E_\alpha^1 & \longrightarrow & \dots \\ & & \downarrow f_\alpha^0 & & \downarrow f_\alpha^1 & & \\ 0 & \longrightarrow & E_{\alpha+1}^0 & \longrightarrow & E_{\alpha+1}^1 & \longrightarrow & \dots \end{array}$$

The reader is invited to check that the direct system $(E_\alpha)_{\alpha < \lambda}$ satisfies all the requirements. □

4. Some sufficient conditions for the heart to be a locally coherent Grothendieck category

Definition. Let \mathcal{D}' be a full triangulated subcategory of the triangulated category \mathcal{D} and let $(\mathcal{U}, \mathcal{U}^\perp[1])$ be a t-structure in \mathcal{D} . We say that this t-structure *restricts to* \mathcal{D}' when $(\mathcal{U} \cap \mathcal{D}', (\mathcal{U}^\perp \cap \mathcal{D}') [1])$ is a t-structure of \mathcal{D}' . This is equivalent to saying that, for each object X of \mathcal{D}' , the truncation triangle $\tau_{\mathcal{U}}(X) \rightarrow X \rightarrow \tau^{\mathcal{U}^\perp}(X) \xrightarrow{+}$ has its three vertices in \mathcal{D}' .

Lemma 4.1. *Let \mathcal{D}' be a full triangulated subcategory of \mathcal{D} and let $(\mathcal{U}, \mathcal{U}^\perp[1])$ be a t-structure in \mathcal{D} whose heart is \mathcal{H} . If the t-structure restricts to \mathcal{D}' , then $\mathcal{H} \cap \mathcal{D}'$ is an abelian exact subcategory of \mathcal{H} .*

Proof. Let $f : X \rightarrow Y$ be a morphism in $\mathcal{H} \cap \mathcal{D}'$ and complete it to a triangle, which is in \mathcal{D}' :

$$X \xrightarrow{f} Y \rightarrow Z \xrightarrow{+} .$$

Note that then $Z \in \mathcal{U} \cap \mathcal{U}^\perp[2]$ and hence $Z[-1] \in \mathcal{U}^\perp[1]$. According to [Parra and Saorín 2015, Lemma 3.1], we have $\tilde{H}(Z) = \tau^{\mathcal{U}^\perp}(Z[-1])[1]$ and $\tilde{H}(Z[-1]) = \tau_{\mathcal{U}}(Z[-1])$. Moreover, since the t-structure restricts to \mathcal{D}' we get that both $\tilde{H}(Z)$ and $\tilde{H}(Z[-1])$ are in $\mathcal{H} \cap \mathcal{D}'$. But we then have a triangle

$$\tilde{H}(Z[-1])[1] \rightarrow Z \rightarrow \tilde{H}(Z) \xrightarrow{+} .$$

By [Beilinson et al. 1982], we have isomorphisms $\text{Ker}_{\mathcal{H}}(f) \cong \tilde{H}(Z[-1])$ and $\text{Coker}_{\mathcal{H}}(f) \cong \tilde{H}(Z)$ and, hence, $\mathcal{H} \cap \mathcal{D}'$ is closed under taking kernels and cokernels in \mathcal{H} . That it is also closed under taking finite coproducts is clear. □

Setting 4.2. In the rest of the section we assume that \mathcal{G} is a locally coherent Grothendieck category and we fix a set \mathcal{S} of finitely presented generators of \mathcal{G} . Recall that then \mathcal{S} is also a set of generators of $\mathcal{D}(\mathcal{G})$ as a triangulated category (see [Nicolás et al. 2015, Lemma 9] or [Psaroudakis and Vitória 2015, Lemma 4.10]).

Lemma 4.3. *Let $X \in \mathcal{D}^{\leq 0}(\mathcal{G})$ have bounded finitely presented homology (i.e., X is homologically bounded and $H^k(X) \in \text{fp}(\mathcal{G})$, for all $k \in \mathbb{Z}$) and let n be a natural*

number. There is a complex $P \in \mathcal{C}^b(\text{sum}(\mathcal{S}))$ together with a morphism $g : P \rightarrow X$ in $\mathcal{D}(\mathcal{G})$ such that the restriction of the natural transformation $g^* : \text{Hom}_{\mathcal{D}(\mathcal{G})}(X, ?) \rightarrow \text{Hom}_{\mathcal{D}(\mathcal{G})}(P, ?)$ to $\mathcal{D}^{[-n, 0]}(\mathcal{G})$ is a natural isomorphism.

Proof. By [Lemma 3.3](#), there is an isomorphism $p : Q \rightarrow X$ in $\mathcal{D}(\mathcal{G})$ such that Q is a complex of objects in $\text{sum}(\mathcal{S})$ concentrated in degrees ≤ 0 . We have that $p^* : \text{Hom}_{\mathcal{D}(\mathcal{G})}(X, ?) \xrightarrow{\cong} \text{Hom}_{\mathcal{D}(\mathcal{G})}(Q, ?)$ is a natural isomorphism of functors $\mathcal{D}(\mathcal{G}) \rightarrow \text{Ab}$. Stupid truncation at $-n - 2$ gives a triangle in $\mathcal{K}(\mathcal{G})$

$$\sigma^{>-n-2}Q \xrightarrow{h} Q \rightarrow \sigma^{\leq -n-2}Q \xrightarrow{+},$$

where the left vertex is in $\mathcal{C}^b(\text{sum}(\mathcal{S}))$. Since $\text{Hom}_{\mathcal{D}(\mathcal{G})}(\sigma^{\leq -n-2}Q[k], ?)$ vanishes on $\mathcal{D}^{[-n, 0]}(\mathcal{G})$, for $k = -1, 0$, we get that the restriction of the natural transformation

$$h^* : \text{Hom}_{\mathcal{D}(\mathcal{G})}(Q, ?) \rightarrow \text{Hom}_{\mathcal{D}(\mathcal{G})}(\sigma^{>-n-2}Q, ?)$$

to $\mathcal{D}^{[-n, 0]}(\mathcal{G})$ is an isomorphism. Putting $P := \sigma^{>-n-2}Q$, the desired morphism g is the composition $P \xrightarrow{h} Q \xrightarrow{p} X$. \square

Remark 4.4. Let $(\mathcal{U}, \mathcal{U}^\perp[1])$ be a t-structure in any triangulated category \mathcal{D} and suppose that it restricts to a full triangulated subcategory \mathcal{D}' . If $\tilde{H} : \mathcal{D} \rightarrow \mathcal{H}$ is the associated cohomological functor, then $\tilde{H}(M)$ is in $\mathcal{H} \cap \mathcal{D}'$, for all $M \in \mathcal{D}'$. This is because $\tau_{\mathcal{U}}(\mathcal{D}') \subseteq \mathcal{D}'$ and $\tau^{\mathcal{U}^\perp[1]}(\mathcal{D}') \subseteq \mathcal{D}'$.

The following technical result is crucial for the main results of the paper.

Proposition 4.5. *Let \mathcal{G} and \mathcal{S} be as in [Setting 4.2](#), let $(\mathcal{U}, \mathcal{U}^\perp[1])$ be a t-structure in $\mathcal{D}(\mathcal{G})$, with heart \mathcal{H} , and let $\tilde{H} : \mathcal{D}(\mathcal{G}) \rightarrow \mathcal{H}$ be the associated cohomological functor. Suppose that the following conditions hold:*

- (1) $(\mathcal{U}, \mathcal{U}^\perp[1])$ restricts to $\mathcal{D}^b(\text{fp}(\mathcal{G}))$.
- (2) There exist integers $m \leq n$ such that $\mathcal{D}^{\leq m}(\mathcal{G}) \subseteq \mathcal{U} \subseteq \mathcal{D}^{\leq n}(\mathcal{G})$.
- (3) $\mathcal{H} \cap \mathcal{D}^b(\text{fp}(\mathcal{G}))$ is a (skeletally small) class of generators of \mathcal{H} .
- (4) For each direct system $(M_i)_{i \in I}$ in \mathcal{H} , for each $S \in \mathcal{S}$ and for each $k \in \mathbb{Z}$, the canonical map $\eta_{S[k]} : \varinjlim \text{Hom}_{\mathcal{D}(\mathcal{G})}(S[k], M_i) \rightarrow \text{Hom}_{\mathcal{D}(\mathcal{G})}(S[k], \varinjlim_{\mathcal{H}} M_i)$ is an isomorphism.

Then \mathcal{H} is a locally coherent Grothendieck category on which $\mathcal{H} \cap \mathcal{D}^b(\mathcal{G})$ is the class of its finitely presented objects.

Proof. Take the cohomological functor $H' := \coprod_{S \in \mathcal{S}} \text{Hom}_{\mathcal{D}(\mathcal{G})}(S, ?) : \mathcal{D}(\mathcal{G}) \rightarrow \text{Ab}$. Using condition (4) and the fact that \mathcal{S} is a set of generators of $\mathcal{D}(\mathcal{G})$, we see that, with the terminology of [\[Parra and Saorín 2015, Section 3\]](#), the pair $(H', +\infty)$ is a cohomological datum in $\mathcal{D}(\mathcal{G})$ for \mathcal{H} . Then [\[Parra and Saorín 2015, Proposition 3.4\]](#) says that \mathcal{H} is an AB5 abelian category. But condition (3) says that it has a set of generators, so that \mathcal{H} is a Grothendieck category.

Fix a direct system $(M_i)_{i \in I}$ in \mathcal{H} in the sequel and consider the full subcategory \mathcal{C} of $\mathcal{D}(\mathcal{G})$ consisting of those complexes X such that

$$\eta_{X[k]} : \varinjlim \text{Hom}_{\mathcal{D}(\mathcal{G})}(X[k], M_i) \rightarrow \text{Hom}_{\mathcal{D}(\mathcal{G})}(X[k], \varinjlim_{\mathcal{H}} M_i)$$

is an isomorphism, for all $k \in \mathbb{Z}$. Using the Five Lemma, one readily sees that \mathcal{C} is a thick subcategory of $\mathcal{D}(\mathcal{G})$ which, by condition (4), contains \mathcal{S} . We then have $\text{thick}_{\mathcal{D}(\mathcal{G})}(\mathcal{S}) \subseteq \mathcal{C}$. In particular, if a complex $X \in \mathcal{C}^b(\text{sum}(\mathcal{S}))$ is viewed as an object of $\mathcal{D}(\mathcal{G})$, then $X \in \mathcal{C}$.

We now claim that η_X is also an isomorphism, for each $X \in \mathcal{D}^b(\text{fp}(\mathcal{G}))$. Indeed, condition (2) implies that $\mathcal{H} \subseteq \mathcal{D}^{[m,n]}(\mathcal{G})$. Let $X \in \mathcal{D}^b(\text{fp}(\mathcal{G}))$ be arbitrary. Replacing n by a larger integer if necessary, we can assume that $X \in \mathcal{D}^{\leq n}(\mathcal{G})$. Then the obvious generalization of Lemma 4.3 says that there exist a $P \in \mathcal{C}^b(\text{sum}(\mathcal{S}))$ and a morphism $g : P \rightarrow X$ in $\mathcal{D}(\mathcal{G})$ such that the natural transformation

$$g^* : \text{Hom}_{\mathcal{D}(\mathcal{G})}(X, ?) \rightarrow \text{Hom}_{\mathcal{D}(\mathcal{G})}(P, ?)$$

is an isomorphism when evaluated on objects of $\mathcal{D}^{[m,n]}(\mathcal{G})$. We then have the following commutative diagram

$$\begin{CD} \varinjlim \text{Hom}_{\mathcal{D}(\mathcal{G})}(X, M_i) @>\eta_X>> \text{Hom}_{\mathcal{D}(\mathcal{G})}(X, \varinjlim_{\mathcal{H}} M_i) \\ @Vg^*VV @VVg^*V \\ \varinjlim \text{Hom}_{\mathcal{D}(\mathcal{G})}(P, M_i) @>\eta_P>> \text{Hom}_{\mathcal{D}(\mathcal{G})}(P, \varinjlim_{\mathcal{H}} M_i) \end{CD}$$

where the vertical arrows are isomorphisms and, due to the previous paragraph, the lower horizontal arrow is an isomorphism also. This settles our claim. In particular, it implies that $\mathcal{H} \cap \mathcal{D}^b(\mathcal{G})$ is a class of finitely presented objects in \mathcal{H} and, by conditions (1) and (3), it is a class of generators of \mathcal{H} (see Remark 4.4). In particular \mathcal{H} is locally finitely presented. Note also that, by condition (1) and Lemma 4.1, we know that $\mathcal{H} \cap \mathcal{D}^b(\text{fp}(\mathcal{G}))$ is closed under taking cokernels (and kernels) in \mathcal{H} . It immediately follows that each finitely presented object of \mathcal{H} is in $\mathcal{H} \cap \mathcal{D}^b(\text{fp}(\mathcal{G}))$ since it is the cokernel of a morphism in this latter category. Then we have that $\mathcal{H} \cap \mathcal{D}^b(\text{fp}(\mathcal{G})) = \text{fp}(\mathcal{H})$, and this is an abelian exact subcategory of \mathcal{H} . Therefore \mathcal{H} is locally coherent. \square

Remark 4.6. Condition (1) of the last proposition is not necessary for the heart to be a locally coherent Grothendieck category. Indeed, by [Parra and Saorín 2014, Corollary 5.12] and using the terminology of that reference, if R is a commutative noetherian ring and $Z \subsetneq \text{Spec } R$ is a perfect sp-subset, then $(\mathcal{U}, \mathcal{U}^\perp[1])$ is a t -structure whose heart is equivalent to $R_Z\text{-Mod}$, where \mathcal{U} consists of the complexes U such that $\text{Supp}(H^j(U)) \subseteq Z$, for all $j > -1$. Then the heart is locally coherent since R_Z is a noetherian commutative ring. But the associated sp-filtration $\phi = \phi_{\mathcal{U}}$

of $\text{Spec } R$ (see [Alonso et al. 2010, Section 2.8 and Theorem 3.11]) is given by $\phi(i) = \text{Spec } R$, for $i \leq -1$, and $\phi(i) = Z$, for all $i > -1$. This sp-filtration does not satisfy in general the weak Cousin condition, in whose case $(\mathcal{U}, \mathcal{U}^\perp[1])$ does not restrict to $\mathcal{D}^b(\text{fp}(R\text{-Mod})) \cong \mathcal{D}_{\text{fg}}^b(R)$ (see [Alonso et al. 2010, Corollary 4.5]). As an example of the last situation, consider $R = \mathbb{Z}$ and $Z = \text{Spec } \mathbb{Z} \setminus \{0\}$, so that $R_Z = \mathbb{Q}$. We have a canonical triangle $\mathbb{Q}/\mathbb{Z}[-1] \rightarrow \mathbb{Z} \rightarrow \mathbb{Q} \xrightarrow{+}$, where $\mathbb{Q}/\mathbb{Z}[-1] \in \mathcal{U}$ and $\mathbb{Q} \in \mathcal{U}^\perp$.

5. The case of the Happel–Reiten–Smalø t-structure

Recall (see [Happel et al. 1996]) that if \mathcal{A} is any abelian category and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ is a torsion pair in \mathcal{A} , then $(\mathcal{U}_\mathbf{t}, \mathcal{U}_\mathbf{t}^\perp[1]) = (\mathcal{U}_\mathbf{t}, \mathcal{V}_\mathbf{t})$ is a t-structure in $\mathcal{D}(\mathcal{A})$, where

$$\mathcal{U}_\mathbf{t} = \{U \in \mathcal{D}^{\leq 0}(\mathcal{A}) : H^0(U) \in \mathcal{T}\} \quad \text{and} \quad \mathcal{V}_\mathbf{t} = \{V \in \mathcal{D}^{\geq -1}(\mathcal{A}) : H^{-1}(V) \in \mathcal{F}\}.$$

This t-structure will be called the *Happel–Reiten–Smalø* (or just *HRS*) *t-structure associated to \mathbf{t}* . In this paper we are only interested in the case when $\mathcal{A} = \mathcal{G}$ is a locally coherent Grothendieck category.

Therefore, all throughout this section, \mathcal{G} will be a locally coherent Grothendieck category and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ will be a torsion pair in \mathcal{G} . Recall that \mathbf{t} is said to be of *finite type* when the torsion radical $t : \mathcal{G} \rightarrow \mathcal{T}$ preserves direct limits or, equivalently, when \mathcal{F} is closed under taking direct limits in \mathcal{G} (see [Krause 1997, Section 2]). We shall say that \mathbf{t} *restricts to $\text{fp}(\mathcal{G})$* when $t(X)$ is in $\text{fp}(\mathcal{G})$, for each $X \in \text{fp}(\mathcal{G})$. Note that this is equivalent to saying that $\mathbf{t}' = (\mathcal{T} \cap \text{fp}(\mathcal{G}), \mathcal{F} \cap \text{fp}(\mathcal{G}))$ is a torsion pair in $\text{fp}(\mathcal{G})$.

Proposition 5.1. *Let $(\mathcal{U}_\mathbf{t}, \mathcal{U}_\mathbf{t}^\perp[1])$ be the HRS t-structure in $\mathcal{D}(\mathcal{G})$ associated to \mathbf{t} . The following assertions are equivalent:*

- (1) *The t-structure $(\mathcal{U}_\mathbf{t}, \mathcal{U}_\mathbf{t}^\perp[1])$ restricts to $\mathcal{D}^b(\text{fp}(\mathcal{G}))$.*
- (2) *The torsion pair \mathbf{t} restricts to $\text{fp}(\mathcal{G})$.*

In particular, if \mathcal{G} is locally noetherian then $(\mathcal{U}_\mathbf{t}, \mathcal{U}_\mathbf{t}^\perp[1])$ restricts to $\mathcal{D}^b(\text{fp}(\mathcal{G}))$.

Proof. Given $M \in \mathcal{D}^b(\text{fp}(\mathcal{G}))$, we have canonical triangles in $\mathcal{D}(\mathcal{G})$

$$\begin{aligned} \tau^{\leq -1} M \rightarrow M \rightarrow \tau^{\geq 0} M \xrightarrow{+}, \\ t(H^0(M))[0] \rightarrow \tau^{\geq 0} M \rightarrow W \xrightarrow{+}, \end{aligned}$$

where $W \in \mathcal{D}^{\geq 0}(\mathcal{G})$, $H^0(W) \cong (H^0(M))/t(H^0(M))$ and $H^k(W) = H^k(M)$, for all $k > 0$. Then $W \in \mathcal{U}_\mathbf{t}^\perp = \mathcal{V}_\mathbf{t}[-1]$. Applying the octahedron axiom to the last two triangles, we obtain two new triangles

$$\begin{aligned} \tau^{\leq -1} M \rightarrow U \rightarrow t(H^0(M))[0] \xrightarrow{+}, \\ U \rightarrow M \rightarrow W \xrightarrow{+}. \end{aligned}$$

It follows from the first triangle that $U \in \mathcal{U}_t$ since the outer vertices of the triangle are in \mathcal{U}_t . We then conclude that the second triangle is precisely the truncation triangle of M with respect to $(\mathcal{U}_t, \mathcal{U}_t^\perp[1])$.

The last truncation triangle is in $\mathcal{D}^b(\mathrm{fp}(\mathcal{G}))$ if, and only if, $U \in \mathcal{D}^b(\mathrm{fp}(\mathcal{G}))$. But this happens exactly when $t(H^0(M))[0] \in \mathcal{D}^b(\mathrm{fp}(\mathcal{G}))$. That is, exactly when $t(H^0(M))$ is a finitely presented object. The equivalence of assertions (1) and (2) is now clear.

Noting that \mathcal{G} is locally coherent all throughout this section, when \mathcal{G} is also locally noetherian we have that $\mathrm{fp}(\mathcal{G})$ coincides with the class $\mathrm{noeth}(\mathcal{G})$ of noetherian objects, which is obviously closed under taking subobjects. Therefore t always restricts to $\mathrm{fp}(\mathcal{G})$. \square

We are now ready to prove the first main result of the paper.

Theorem 5.2. *Let \mathcal{G} be a locally coherent Grothendieck category, let $t = (\mathcal{T}, \mathcal{F})$ be a torsion pair in \mathcal{G} , let $(\mathcal{U}_t, \mathcal{U}_t^\perp[1])$ be the associated t -structure in $\mathcal{D}(\mathcal{G})$ and let \mathcal{H}_t be its heart. The following assertions are equivalent:*

- 1) $(\mathcal{U}_t, \mathcal{U}_t^\perp[1])$ restricts to $\mathcal{D}^b(\mathrm{fp}(\mathcal{G}))$ and \mathcal{H}_t is a locally coherent Grothendieck category (with $\mathcal{H}_t \cap \mathcal{D}^b(\mathrm{fp}(\mathcal{G}))$ as the class of finitely presented objects).
- 2) t is of finite type and restricts to $\mathrm{fp}(\mathcal{G})$.
- 3) There exists a torsion pair $t' = (\mathcal{T}', \mathcal{F}')$ in $\mathrm{fp}(\mathcal{G})$ such that $t = (\varinjlim \mathcal{T}', \varinjlim \mathcal{F}')$.

When in addition \mathcal{G} is locally noetherian, these assertions are also equivalent to:

- 4) t is of finite type.

Proof. All throughout the proof, we fix a set \mathcal{S} of finitely presented generators of \mathcal{G} .

1) \Rightarrow 2) By [Proposition 5.1](#), we know that t restricts to $\mathrm{fp}(\mathcal{G})$ and, by [\[Parra and Saorín 2015, Theorem 4.8\]](#), we know that t is of finite type.

2) \Rightarrow 3) If we put $\mathcal{T}' = \mathcal{T} \cap \mathrm{fp}(\mathcal{G})$ and $\mathcal{F}' = \mathcal{F} \cap \mathrm{fp}(\mathcal{G})$, then $t' = (\mathcal{T}', \mathcal{F}')$ is a torsion pair in $\mathrm{fp}(\mathcal{G})$ since t restricts to $\mathrm{fp}(\mathcal{G})$. By [\[Crawley-Boevey 1994, Lemma 4.4\]](#), we know that $(\varinjlim \mathcal{T}', \varinjlim \mathcal{F}')$ is a torsion pair in \mathcal{G} . But \mathcal{T} and \mathcal{F} are closed under taking direct limits in \mathcal{G} , which implies that $\varinjlim \mathcal{T}' \subseteq \mathcal{T}$ and $\varinjlim \mathcal{F}' \subseteq \mathcal{F}$. Since we always have $\mathcal{F} = \mathcal{T}^\perp \subseteq (\varinjlim \mathcal{T}')^\perp = \varinjlim \mathcal{F}'$ we conclude that $(\mathcal{T}, \mathcal{F}) = (\varinjlim \mathcal{T}', \varinjlim \mathcal{F}')$.

3) \Rightarrow 2) is clear.

2) \Rightarrow 1) The finite type condition of t implies that \mathcal{H}_t is a Grothendieck category (see [\[Parra and Saorín 2016a, Theorem 1.2\]](#)). Now, let $(M_i)_{i \in I}$ be a direct system in \mathcal{H}_t . Bearing in mind that \mathcal{F} is closed under taking direct limits in \mathcal{G} and using [\[Parra and Saorín 2015, Proposition 4.2\]](#), we get an exact sequence in \mathcal{H}_t :

$$0 \rightarrow (\varinjlim H^{-1}(M_i))[1] \rightarrow \varinjlim_{\mathcal{H}_t} M_i \rightarrow (\varinjlim H^0(M_i))[0] \rightarrow 0.$$

To abbreviate, let us put $(X, Y) = \text{Hom}_{\mathcal{D}(\mathcal{G})}(X, Y)$, for all $X, Y \in \mathcal{D}(\mathcal{G})$. Then, for each $S \in \mathcal{S}$ and each $k \in \mathbb{Z}$, we have a commutative diagram of abelian groups with exact columns, where the horizontal arrows are the canonical morphisms:

$$\begin{array}{ccc}
 (S[k], (\varinjlim H^0(M_i))[-1]) & \longleftarrow & \varinjlim(S[k], H^0(M_i)[-1]) \\
 \downarrow & & \downarrow \\
 (S[k], (\varinjlim H^{-1}(M_i))[1]) & \longleftarrow & \varinjlim(S[k], H^{-1}(M_i)[1]) \\
 \downarrow & & \downarrow \\
 (S[k], \varinjlim_{\mathcal{H}_t} M_i) & \longleftarrow & \varinjlim(S[k], M_i) \\
 \downarrow & & \downarrow \\
 (S[k], (\varinjlim H^0(M_i))[0]) & \longleftarrow & \varinjlim(S[k], H^0(M_i)[0]) \\
 \downarrow & & \downarrow \\
 (S[k], (\varinjlim H^{-1}(M_i))[2]) & \longleftarrow & \varinjlim(S[k], H^{-1}(M_i)[2])
 \end{array}$$

By [Proposition 3.5](#), we have that the two uppermost and the two lowermost horizontal arrows are isomorphisms, which implies the canonical map $\varinjlim(S[k], M_i) \rightarrow (S[k], \varinjlim_{\mathcal{H}_t} M_i)$ is also an isomorphism.

We will check now that all conditions (1)–(4) of [Proposition 4.5](#) are satisfied by $(\mathcal{U}_t, \mathcal{U}_t[1])$. By [Proposition 5.1](#), we know that $(\mathcal{U}_t, \mathcal{U}_t^\perp[1])$ restricts to $\mathcal{D}^b(\text{fp}(\mathcal{G}))$ and, by definition of the HRS t-structure, we know that $\mathcal{D}^{\leq -1}(\mathcal{G}) \subseteq \mathcal{U}_t \subseteq \mathcal{D}^{\leq 0}(\mathcal{G})$, so that conditions (1) and (2) of [Proposition 4.5](#) hold. Moreover, the previous paragraph says that condition (4) also holds.

We will finally check that each object of \mathcal{H}_t is an epimorphic image of a coproduct of objects of $\mathcal{H}_t \cap \mathcal{D}^b(\text{fp}(\mathcal{G}))$, which will give condition (3) of [Proposition 5.1](#) and will end the proof. Let M be any object of \mathcal{H}_t and let us write $H^0(M) = \varinjlim T_i$, for some direct system $(T_i)_{i \in I}$ in $\mathcal{T} \cap \text{fp}(\mathcal{G})$. Note that this is possible since $\mathcal{T} = \varinjlim(\mathcal{T} \cap \text{fp}(\mathcal{G}))$. Considering the canonical exact sequence

$$0 \rightarrow H^{-1}(M)[1] \rightarrow M \rightarrow H^0(M)[0] \rightarrow 0$$

and pulling it back, for each $i \in I$, along the obvious map $T_i[0] \rightarrow H^0(M)[0]$, we get a direct system of exact sequences in \mathcal{H}_t :

$$0 \rightarrow H^{-1}(M)[1] \rightarrow M_i \rightarrow T_i[0] \rightarrow 0.$$

Since \mathcal{H}_t is a Grothendieck category it immediately follows that $M = \varinjlim_{\mathcal{H}_t} M_i$, so that M is an epimorphic image of $\coprod_{i \in I} M_i$. Replacing M by any of the M_i , we can and shall assume in the rest of the proof that $H^0(M) \in \mathcal{T} \cap \text{fp}(\mathcal{G})$. We then write

M as a complex $\cdots 0 \rightarrow M^{-1} \rightarrow M^0 \rightarrow 0 \cdots$ concentrated in degrees -1 and 0 . Note that if we put $M^0 = \varinjlim M_i^0$, where $(M_i^0)_{i \in I}$ is a direct system in $\text{fp}(\mathcal{G})$, then some composition $M_j^0 \xrightarrow{t_j} \varinjlim M_i^0 = M^0 \xrightarrow{p} H^0(M)$ should be an epimorphism, because $H^0(M)$ is finitely presented. Replacing M^0 by M_j^0 if necessary, we can assume in the sequel that M^0 is also finitely presented.

Once we assume that $H^0(M)$ and M^0 are both finitely presented, we follow the lines of the proof of [Parra and Saorín 2015, Proposition 4.7] with an easy adaptation. The details are left to the reader. Since M^{-1} is a direct limit of finitely presented objects, we can fix an epimorphism $\coprod_{j \in J} X_j \twoheadrightarrow M^{-1}$ in \mathcal{G} , where $X_j \in \text{fp}(\mathcal{G})$ for all $j \in J$. Now we construct a four-row commutative diagram as in the mentioned proof, where $G^{(J)}$ and $G^{(F)}$ are replaced in our case by $\coprod_{j \in J} X_j$ and $\coprod_{j \in F} X_j$, respectively. The key point now is that the appearing U_F and X_F are finitely presented objects. Since t restricts to $\text{fp}(\mathcal{G})$, we also know that $t(X_F)$ and M_F^0 are finitely presented, for each finite subset $F \subseteq J$. If now $L = \tilde{H}_{\mathcal{U}_t} : \mathcal{U}_t \rightarrow \mathcal{H}_t$ is the left adjoint to the inclusion functor (see [Parra and Saorín 2015, Lemma 3.1]), the mentioned proof shows that we have epimorphisms $\coprod_{F \subseteq J, F \text{ finite}} L(K_F) \twoheadrightarrow L(K_J)$ and $L(K_J) \twoheadrightarrow M$ in \mathcal{H}_t , where $L(K_F)$ is the object of \mathcal{H}_t represented by the complex

$$\cdots 0 \rightarrow \frac{\coprod_{j \in F} X_j}{t(U_F)} \rightarrow M_F^0 \rightarrow 0 \cdots,$$

concentrated in degrees -1 and 0 . But $t(U_F)$ is finitely presented, because so is U_F . It follows that the latter complex is a complex of finitely presented objects, and hence $L(K_F) \in \mathcal{H}_t \cap \mathcal{D}^b(\text{fp}(\mathcal{G}))$.

4) \Rightarrow 2) = 3) If \mathcal{G} is locally noetherian, each torsion pair restricts to its subcategory of noetherian objects, that is, to $\text{fp}(\mathcal{G})$. □

6. The heart of a restricted t-structure in the derived category of a commutative noetherian ring

All throughout this section R is a commutative noetherian ring. To apply the results of earlier sections, we will consider $\mathcal{G} = R\text{-Mod}$ the category of all R -modules, which is a locally noetherian Grothendieck category. Then we have that $\text{fp}(\mathcal{G}) = R\text{-mod}$ is the subcategory of finitely generated R -modules and, as usual (see comments on page 207), we identify $\mathcal{D}_{\text{fg}}^b(R) := \mathcal{D}_{\text{fp}}^b(R\text{-Mod})$ with $\mathcal{D}^b(R\text{-mod})$.

Recall that a *filtration by supports* or *sp-filtration* of $\text{Spec } R$ is a decreasing map $\phi : \mathbb{Z} \rightarrow \mathcal{P}(\text{Spec } R)$ such that $\phi(i) \subseteq \text{Spec } R$ is a stable under specialization subset, for each $i \in \mathbb{Z}$. Filtrations by supports turn out to be in bijection with the compactly generated t-structures in $\mathcal{D}(R)$ (see [Alonso et al. 2010, Theorem 3.11]). Concretely, given an sp-filtration ϕ and putting

$$\mathcal{U}_\phi = \{U \in \mathcal{D}(R) : \text{Supp}(H^i(U)) \subseteq \phi(i), \text{ for all } i \in \mathbb{Z}\},$$

we get a compactly generated t-structure $\tau_\phi = (\mathcal{U}_\phi, \mathcal{U}_\phi^\perp[1])$ and the assignment $\phi \rightsquigarrow \tau_\phi$ gives the mentioned bijection. All through this section, the reader is referred to [Alonso et al. 2010] for all nondefined terms that we might use.

Lemma 6.1. *Let $X \in \mathcal{D}_{\text{fg}}^b(R)$ and $Y \in \mathcal{D}^+(R)$. For each $\mathfrak{p} \in \text{Spec } R$, the canonical map*

$$\text{Hom}_{\mathcal{D}(R)}(X, Y)_{\mathfrak{p}} \rightarrow \text{Hom}_{\mathcal{D}(R_{\mathfrak{p}})}(X_{\mathfrak{p}}, Y_{\mathfrak{p}})$$

is an isomorphism.

Proof. Let us fix $Y \in \mathcal{D}^+(R)$, which we consider to be a bounded below complex of injective R -modules. For each Z in $\mathcal{D}_{\text{fg}}^b(R)$, we denote by η_Z the canonical map $\text{Hom}_{\mathcal{D}(R)}(Z, Y)_{\mathfrak{p}} \rightarrow \text{Hom}_{\mathcal{D}(R_{\mathfrak{p}})}(Z_{\mathfrak{p}}, Y_{\mathfrak{p}})$. We then consider the full subcategory \mathcal{C} of $\mathcal{D}_{\text{fg}}^b(R)$ consisting of those Z such that $\eta_{Z[k]}$ is an isomorphism, for all $k \in \mathbb{Z}$. It is clear that \mathcal{C} is a thick subcategory of $\mathcal{D}_{\text{fg}}^b(R)$.

We claim that $M[0] \in \mathcal{C}$, for each finitely generated R -module M . Once this is proved, the proof will be finished. Indeed, we will conclude that $\mathcal{C} = \mathcal{D}_{\text{fg}}^b(R)$ since each $Z \in \mathcal{D}_{\text{fg}}^b(R)$ is a finite iterated extension of the stalk complexes $H^{-k}(Z)[k]$, and each $H^{-k}(Z)$ is finitely generated. Recall that $\text{Hom}_{\mathcal{D}(R)}(M[-k], Y)$ is the k -th homology module of the complex of R -modules $\text{Hom}_R(M, Y)$. Similarly, $\text{Hom}_{\mathcal{D}(R_{\mathfrak{p}})}(M_{\mathfrak{p}}[-k], Y_{\mathfrak{p}})$ is the k -th homology module of the complex of $R_{\mathfrak{p}}$ -modules $\text{Hom}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}, Y_{\mathfrak{p}})$ since $Y_{\mathfrak{p}}$ is a bounded below complex of injective $R_{\mathfrak{p}}$ -modules. The claim follows from the exactness of the localization at \mathfrak{p} and from the truth of the result when Y is a module (see, e.g., [Kunz 1985, Proposition IV.1.10]). \square

Lemma 6.2. *Let R be connected, let $(\mathcal{U}, \mathcal{U}^\perp[1])$ be a compactly generated t-structure in $\mathcal{D}(R)$ which restricts to $\mathcal{D}_{\text{fg}}^b(R)$, let \mathcal{H} be its heart and let $U \in \mathcal{D}^-(R) \cap \mathcal{U}$ be a complex with finitely generated homology modules. Then $\tilde{H}(U)$ is in $\mathcal{H} \cap \mathcal{D}_{\text{fg}}^b(R)$.*

Proof. Let ϕ be the sp-filtration of $\text{Spec } R$ associated to $(\mathcal{U}, \mathcal{U}^\perp[1])$. By Corollaries 4.5 and 4.8 of [Alonso et al. 2010], we know that there exists some $j_0 \in \mathbb{Z}$ such that $\phi(j_0) = \text{Spec } R$. Without loss of generality, we assume that $j_0 = 0$. We then have

$$\mathcal{D}^{\leq 0}(R) \subseteq \mathcal{U} \quad \text{and} \quad \mathcal{H} \subseteq \mathcal{D}^{\geq 0}(R).$$

By considering now for the object U of the statement the canonical truncation triangle

$$\tau^{\leq 0}(U[-1]) \rightarrow U[-1] \xrightarrow{g} \tau^{> 0}(U[-1]) \rightarrow$$

and applying the octahedron axiom, we see that

$$\tau^{\mathcal{U}^\perp}(g) : \tau^{\mathcal{U}^\perp}(U[-1]) \rightarrow \tau^{\mathcal{U}^\perp}(\tau^{> 0}(U[-1]))$$

is an isomorphism. However, the codomain of this morphism is in $\mathcal{D}_{\text{fg}}^b(R)$ since

$\tau^{>0}(U[-1]) \in \mathcal{D}_{\text{fg}}^b(R)$ and the t -structure $(\mathcal{U}, \mathcal{U}^\perp[1])$ restricts to $\mathcal{D}_{\text{fg}}^b(R)$. Then $\widetilde{H}(U) = \tau^{\mathcal{U}^\perp}(U[-1])[1]$ is in $\mathcal{D}_{\text{fg}}^b(R)$ (see [Parra and Saorín 2015, Lemma 3.1]). □

We are now ready to prove the main result of the paper.

Theorem 6.3. *Let R be a commutative noetherian ring and let $(\mathcal{U}, \mathcal{U}^\perp[1])$ be a compactly generated t -structure in $\mathcal{D}(R)$ which restricts to $\mathcal{D}_{\text{fg}}^b(R)$. The heart \mathcal{H} of this t -structure is a locally coherent Grothendieck category where $\mathcal{H} \cap \mathcal{D}_{\text{fg}}^b(R)$ is the subcategory of its finitely presented objects.*

Proof. All throughout the proof, without loss of generality, we assume that R is connected. Remember that then the associated sp -filtration ϕ satisfies the weak Cousin condition and, hence, has the property that $\phi(i) = \text{Spec } R$, for $i \ll 0$ (see [Alonso et al. 2010, Theorem 4.4 and Corollary 4.8]). This in turn implies that $\mathcal{H} = \mathcal{H}_\phi \subseteq \mathcal{D}^{\geq m}(R)$, for some $m \in \mathbb{Z}$. Moreover, by [Parra and Saorín 2014, Theorem 4.10], we know that $\mathcal{H} = \mathcal{H}_\phi$ is a Grothendieck category.

Step 1: $\mathcal{H} \cap \mathcal{D}_{\text{fg}}^b(R)$ is a (skeletally small) class of generators of \mathcal{H} : Let \mathcal{U}' denote the full subcategory of \mathcal{U} consisting of complexes in $\mathcal{U} \cap \mathcal{D}^-(R)$ which have finitely generated homology modules. Each object of \mathcal{U}' is isomorphic in $\mathcal{D}(R)$ to a bounded above complex of finitely generated R -modules. Let $L = \widetilde{H}_{|\mathcal{U}'} : \mathcal{U} \rightarrow \mathcal{H}$ be the left adjoint to the inclusion functor $\mathcal{H} \hookrightarrow \mathcal{U}$. A slight modification of the proof of [Parra and Saorín 2014, Proposition 3.10] shows that $\mathcal{X} := L(\mathcal{U}')$ is a skeletally small class of generators of \mathcal{H} . By Lemma 6.2, we get that $\mathcal{X} \subseteq \mathcal{H} \cap \mathcal{D}_{\text{fg}}^b(R)$, which ends this first step.

Step 2: *The result is true when ϕ is eventually trivial* (i.e., when $\phi(i) = \emptyset$, for some $i \in \mathbb{Z}$): We shall check all conditions (1)–(4) of Proposition 4.5. Without loss of generality, we assume that the filtration is

$$\text{Spec } R = \cdots \phi(-n-1) = \phi(-n) \supsetneq \phi(-n+1) \supseteq \cdots \supseteq \phi(0) \supsetneq \phi(1) = \phi(2) = \cdots = \emptyset,$$

in which case we have that $\mathcal{D}^{\leq -n}(R) \subseteq \mathcal{U} \subseteq \mathcal{D}^{\leq 0}(R)$ and $\mathcal{H} = \mathcal{H}_\phi \subseteq \mathcal{D}^{[-n, 0]}(R)$ (see [Parra and Saorín 2014, Lemma 4.1]). Then condition (2) of Proposition 4.5 holds and condition (1) holds by hypothesis. Moreover, Step 1 of this proof gives condition (3) of that proposition. Finally, bearing in mind that we have a natural isomorphism $H^k \cong \text{Hom}_{\mathcal{D}(R)}(R[-k], ?)$ of functors $\mathcal{D}(R) \rightarrow R\text{-Mod}$, by taking $\mathcal{S} = \{R\}$ and using [Parra and Saorín 2014, Theorem 4.9] we also get that condition (4) holds.

Step 3: *The general case.* The proof reduces to checking that $\mathcal{H} \cap \mathcal{D}_{\text{fg}}^b(R) \subseteq \text{fp}(\mathcal{H})$. Indeed, if this is proved, then Step 1 implies that \mathcal{H} is locally finitely presented and that each object in $\text{fp}(\mathcal{H})$ is the cokernel of a morphism in $\mathcal{H} \cap \mathcal{D}_{\text{fg}}^b(R)$. It will follow from Lemma 4.1 that $\text{fp}(\mathcal{H}) = \mathcal{H} \cap \mathcal{D}_{\text{fg}}^b(R)$ and that this is an abelian exact

subcategory of \mathcal{H} . That is, \mathcal{H} will be a locally coherent Grothendieck category with $\mathcal{H} \cap \mathcal{D}_{\text{fg}}^b(R)$ as its class of finitely presented objects.

We then prove the inclusion $\mathcal{H} \cap \mathcal{D}_{\text{fg}}^b(R) \subseteq \text{fp}(\mathcal{H})$. Let $(M_i)_{i \in I}$ be a direct system in \mathcal{H} and let $X \in \mathcal{H} \cap \mathcal{D}_{\text{fg}}^b(R)$ be any object. We consider the canonical morphism

$$\eta_X : \varinjlim \text{Hom}_{\mathcal{D}(R)}(X, M_i) \rightarrow \text{Hom}_{\mathcal{D}(R)}(X, \varinjlim_{\mathcal{H}} M_i),$$

which is a morphism in $R\text{-Mod}$. Localization at any prime ideal \mathfrak{p} preserves direct limits and, by [Parra and Saorín 2014, Proposition 3.11], we also have that $(\varinjlim_{\mathcal{H}} M_i)_{\mathfrak{p}} \cong \varinjlim_{\mathcal{H}_{\mathfrak{p}}} (M_i)_{\mathfrak{p}}$. Here if $\mathcal{H} = \mathcal{H}_{\phi}$, then we put $\mathcal{H}_{\mathfrak{p}} = \mathcal{H}_{\phi_{\mathfrak{p}}}$, using the terminology of [Parra and Saorín 2014]. Therefore, using Lemma 6.1, we can identify $(\eta_X)_{\mathfrak{p}} : (\varinjlim \text{Hom}_{\mathcal{D}(R)}(X, M_i))_{\mathfrak{p}} \rightarrow (\text{Hom}_{\mathcal{D}(R)}(X, \varinjlim_{\mathcal{H}} M_i))_{\mathfrak{p}}$ with the canonical morphism

$$\eta_{X_{\mathfrak{p}}} : \varinjlim \text{Hom}_{\mathcal{D}(R_{\mathfrak{p}})}(X_{\mathfrak{p}}, (M_i)_{\mathfrak{p}}) \rightarrow \text{Hom}_{\mathcal{D}(R_{\mathfrak{p}})}(X_{\mathfrak{p}}, \varinjlim_{\mathcal{H}_{\mathfrak{p}}} (M_i)_{\mathfrak{p}}).$$

But the sp-filtration $\phi_{\mathfrak{p}}$ of $\text{Spec } R_{\mathfrak{p}}$ also satisfies the weak Cousin condition and, since $R_{\mathfrak{p}}$ has finite Krull dimension, we get that $\phi_{\mathfrak{p}}$ is eventually trivial (see [Alonso et al. 2010, Corollary 4.8]). The truth of the theorem when the associated filtration is eventually trivial implies that $\eta_{X_{\mathfrak{p}}}$ is an isomorphism, for all $\mathfrak{p} \in \text{Spec } R$, because $X_{\mathfrak{p}} \in \text{fp}(\mathcal{H}_{\mathfrak{p}})$. Therefore the kernel and cokernel of η_X are R -modules with empty support. Then they are both zero, so that η_X is an isomorphism, and hence X is in $\text{fp}(\mathcal{H})$ as desired. \square

Corollary 6.4. *Let R be a commutative noetherian ring. The heart of any t-structure in $\mathcal{D}_{\text{fg}}^b(R)$ is equivalent to the category of finitely presented objects of a locally coherent Grothendieck category.*

Proof. Each t-structure in $\mathcal{D}_{\text{fg}}^b(R)$ is the restriction of the t-structure τ_{ϕ} in $\mathcal{D}(R)$ associated to an sp-filtration (see [Alonso et al. 2010, Corollary 3.12]). The result is then an immediate consequence of the last theorem, using [Alonso et al. 2010, Theorem 3.10]. \square

As a final comment, we give the geometric translation of the last theorem and corollary:

Corollary 6.5. *Let \mathbb{X} be an affine noetherian scheme and let $(\mathcal{U}, \mathcal{U}^{\perp}[1])$ be a t-structure in $\mathcal{D}(\mathbb{X}) := \mathcal{D}(\text{Qcoh}(\mathbb{X}))$ which restricts to $\mathcal{D}_{\text{coh}}^b(\mathbb{X}) \cong \mathcal{D}^b(\text{coh}(\mathbb{X}))$. The heart \mathcal{H} of the t-structure is a locally coherent Grothendieck category on which $\mathcal{H} \cap \mathcal{D}_{\text{coh}}^b(\mathbb{X})$ is the class of finitely presented objects. In particular, the heart of each t-structure in $\mathcal{D}^b(\text{coh}(\mathbb{X}))$ is equivalent to the category of finitely presented objects of a locally coherent Grothendieck category.*

References

- [Adámek and Rosický 1994] J. Adámek and J. Rosický, *Locally presentable and accessible categories*, London Mathematical Society Lecture Note Series **189**, Cambridge University Press, 1994. [MR](#) [Zbl](#)
- [Alonso et al. 2010] L. Alonso Tarrío, A. Jeremías López, and M. Saorín, “Compactly generated t -structures on the derived category of a Noetherian ring”, *J. Algebra* **324**:3 (2010), 313–346. [MR](#) [Zbl](#)
- [Beilinson et al. 1982] A. A. Beilinson, J. Bernstein, and P. Deligne, “Faisceaux pervers”, pp. 5–171 in *Analysis and topology on singular spaces, I* (Luminy, 1981), Astérisque **100**, Société Mathématique de France, Paris, 1982. [MR](#) [Zbl](#)
- [Colpi and Gregorio 2010] R. Colpi and E. Gregorio, “The heart of cotilting theory pair is a Grothendieck category”, preprint, 2010.
- [Colpi et al. 2007] R. Colpi, E. Gregorio, and F. Mantese, “On the heart of a faithful torsion theory”, *J. Algebra* **307**:2 (2007), 841–863. [MR](#) [Zbl](#)
- [Colpi et al. 2011] R. Colpi, F. Mantese, and A. Tonolo, “When the heart of a faithful torsion pair is a module category”, *J. Pure Appl. Algebra* **215**:12 (2011), 2923–2936. [MR](#) [Zbl](#)
- [Crawley-Boevey 1994] W. Crawley-Boevey, “Locally finitely presented additive categories”, *Comm. Algebra* **22**:5 (1994), 1641–1674. [MR](#) [Zbl](#)
- [Garkusha 2009] G. A. Garkusha, “Classifying finite localizations of quasi-coherent sheaves”, *Algebra i Analiz* **21**:3 (2009), 93–129. In Russian; translated in *St. Petersburg Math. J.* **21**:3 (2010), 433–458. [MR](#) [Zbl](#)
- [Gillespie 2016] J. Gillespie, “Models for homotopy categories of injectives and Gorenstein injectives”, *Comm. Algebra* (online publication October 2016).
- [Happel et al. 1996] D. Happel, I. Reiten, and S. O. Smalø, *Tilting in abelian categories and quasitilted algebras*, Mem. Amer. Math. Soc. **575**, Amer. Math. Soc., Providence, RI, 1996. [MR](#) [Zbl](#)
- [Hartshorne 1966] R. Hartshorne, *Residues and duality*, Lecture Notes in Mathematics **20**, Springer, 1966. [MR](#) [Zbl](#)
- [Herzog 1997] I. Herzog, “The Ziegler spectrum of a locally coherent Grothendieck category”, *Proc. London Math. Soc.* (3) **74**:3 (1997), 503–558. [MR](#) [Zbl](#)
- [Hoshino et al. 2002] M. Hoshino, Y. Kato, and J.-I. Miyachi, “On t -structures and torsion theories induced by compact objects”, *J. Pure Appl. Algebra* **167**:1 (2002), 15–35. [MR](#) [Zbl](#)
- [Keller 1994] B. Keller, “Deriving DG categories”, *Ann. Sci. École Norm. Sup.* (4) **27**:1 (1994), 63–102. [MR](#) [Zbl](#)
- [Krause 1997] H. Krause, “The spectrum of a locally coherent category”, *J. Pure Appl. Algebra* **114**:3 (1997), 259–271. [MR](#) [Zbl](#)
- [Kunz 1985] E. Kunz, *Introduction to commutative algebra and algebraic geometry*, Birkhäuser, Boston, 1985. [MR](#) [Zbl](#)
- [Mantese and Tonolo 2012] F. Mantese and A. Tonolo, “On the heart associated with a torsion pair”, *Topology Appl.* **159**:9 (2012), 2483–2489. [MR](#) [Zbl](#)
- [Neeman 2001] A. Neeman, *Triangulated categories*, Annals of Mathematics Studies **148**, Princeton University Press, 2001. [MR](#) [Zbl](#)
- [Nicolás et al. 2015] P. Nicolás, M. Saorín, and A. Zvonareva, “Silting theory in triangulated categories with coproducts”, preprint, 2015. [arXiv](#)
- [Parra and Saorín 2014] C. E. Parra and M. Saorín, “Hearts of t -structures in the derived category of a commutative Noetherian ring”, preprint, 2014. To appear in *Trans. Amer. Math. Soc.* [arXiv](#)

- [Parra and Saorín 2015] C. E. Parra and M. Saorín, “Direct limits in the heart of a t-structure: the case of a torsion pair”, *J. Pure Appl. Algebra* **219**:9 (2015), 4117–4143. Addendum in **220**:6 (2016), 2467–2469. [MR](#) [Zbl](#)
- [Parra and Saorín 2016a] C. E. Parra and M. Saorín, “Addendum to ‘Direct limits in the heart of a t-structure: the case of a torsion pair’”, *J. Pure Appl. Algebra* **220**:6 (2016), 2467–2469. [MR](#) [Zbl](#)
- [Parra and Saorín 2016b] C. E. Parra and M. Saorín, “On hearts which are module categories”, *J. Math. Soc. Japan* **68**:4 (2016), 1421–1460.
- [Popescu 1973] N. Popescu, *Abelian categories with applications to rings and modules*, London Mathematical Society Monographs **3**, Academic Press, London, 1973. [MR](#) [Zbl](#)
- [Prest 2009] M. Prest, *Purity, spectra and localisation*, Encyclopedia of Mathematics and its Applications **121**, Cambridge University Press, 2009. [MR](#) [Zbl](#)
- [Psaroudakis and Vitória 2015] C. Psaroudakis and J. Vitória, “Realisation functors in tilting theory”, preprint, 2015. [arXiv](#)
- [Sitte 2014] T. Sitte, *Local cohomology sheaves on algebraic stacks*, Ph.D. thesis, Universität Regensburg, 2014, available at <http://epub.uni-regensburg.de/30539/1/thesis.pdf>.
- [Šťovíček 2014] J. Šťovíček, “On purity and applications to coderived and singularity categories”, preprint, 2014. [arXiv](#)
- [Verdier 1996] Jean-Louis Verdier, *Des catégories dérivées des catégories abéliennes*, edited by G. Maltsiniotis, Astérisque **239**, Société Mathématique de France, Paris, 1996. [MR](#) [Zbl](#)

Received May 19, 2016. Revised September 20, 2016.

MANUEL SAORÍN
DEPARTAMENTO DE MATEMÁTICAS
UNIVERSIDAD DE MURCIA
APTDO. 4021
30100 MURCIA
SPAIN
msaorinc@um.es

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 287 No. 1 March 2017

Operator ideals related to absolutely summing and Cohen strongly summing operators	1
GERALDO BOTELHO, JAMILSON R. CAMPOS and JOEDSON SANTOS	
Homology for quandles with partial group operations	19
SCOTT CARTER, ATSUSHI ISHII, MASAHICO SAITO and KOKORO TANAKA	
Three-dimensional discrete curvature flows and discrete Einstein metrics	49
HUABIN GE, XU XU and SHIJIN ZHANG	
Inclusion of configuration spaces in Cartesian products, and the virtual cohomological dimension of the braid groups of S^2 and $\mathbb{R}P^2$	71
DACIBERG LIMA GONÇALVES and JOHN GUASCHI	
Groups of PL-homeomorphisms admitting nontrivial invariant characters	101
DACIBERG L. GONÇALVES, PARAMESWARAN SANKARAN and RALPH STREBEL	
Bernstein-type theorems for spacelike stationary graphs in Minkowski spaces	159
XIANG MA, PENG WANG and LING YANG	
Comparison results for derived Deligne–Mumford stacks	177
MAURO PORTA	
On locally coherent hearts	199
MANUEL SAORÍN	
Approximability of convex bodies and volume entropy in Hilbert geometry	223
CONSTANTIN VERNICOS	