

*Pacific  
Journal of  
Mathematics*

**$C^1$ -UMBILICS WITH ARBITRARILY HIGH INDICES**

NAOYA ANDO, TOSHIFUMI FUJIYAMA AND MASAOKI UMEHARA

Volume 288 No. 1

May 2017



**$C^1$ -UMBILICS WITH ARBITRARILY HIGH INDICES**

NAOYA ANDO, TOSHIFUMI FUJIYAMA AND MASAACKI UMEHARA

**We show that  $C^1$ -umbilics with arbitrarily high indices exist. This implies that more than  $C^1$ -regularity is required to prove Loewner's conjecture.**

**1. Introduction**

The *index* of an isolated umbilic on a given regular surface is the index of the curvature line flow of the surface at that point, which takes values in the set of half-integers. *Loewner's conjecture* asserts that any isolated umbilic on an immersed surface must have index at most 1. *Carathéodory's conjecture* asserts the existence of at least two umbilics on an immersed sphere in  $\mathbb{R}^3$ , which follows immediately from Loewner's conjecture. Although this problem was investigated mainly on real-analytic surfaces after Hamburger's work [1940; 1941a; 1941b], several geometers recently became interested in nonanalytic cases; see [Ando 2003; Bates 2001; Ghomi and Howard 2012; Gutierrez et al. 1996; Smyth and Xavier 1992]. In particular, Smyth and Xavier [1992] observed that Enneper's minimal surface is inverted to a branched sphere such that the index of the curvature line flow at the branch point is equal to two. Bates [2001] found that the graph of the function

$$(1-1) \quad B(x, y) := 2 + \frac{xy}{\sqrt{1+x^2}\sqrt{1+y^2}}$$

has no umbilics on  $\mathbb{R}^2$  and inversion of it gives a genus zero surface without self-intersections, which is differentiable at the image of infinity under that inversion. Ghomi and Howard [2012] gave similar examples of genus zero surfaces using inversion. Moreover, they showed that Carathéodory's conjecture for closed convex surfaces can be reduced to the problem of existence of umbilics of certain entire graphs over  $\mathbb{R}^2$ . A brief history of Carathéodory's conjecture and recent developments are written also in [Ghomi and Howard 2012]. Recently, Guilfoyle and

---

Ando was partly supported by the Grant-in-Aid for Young Scientists (B) 24740048, Japan Society for the Promotion of Science. Umehara was partly supported by the Grant-in-Aid for Scientific Research (A) 262457005, Japan Society for the Promotion of Science.

*MSC2010:* primary 53A05, 53C45; secondary 57R42, 37C10, 53A30.

*Keywords:* umbilic, curvature line, Carathéodory's conjecture, Loewner's conjecture.

Klingenberg [2008; 2012] gave an approach to proving the Carathéodory and the Loewner conjectures in the smooth case.

Let  $P : U \rightarrow \mathbb{R}^3$  be a  $C^1$ -immersion defined on an open subset  $U$  of  $\mathbb{R}^2$  such that  $P$  is  $C^\infty$ -differentiable on  $U \setminus \{q\}$  and not  $C^2$ -differentiable at  $q$ . Then the point  $q \in U$  is called a  $C^1$ -umbilic if the umbilics of  $P$  on  $U \setminus \{q\}$  do not accumulate to  $q$ . At that point  $q$ , we can compute the index of the curvature line flow of  $P$ . In this paper, we prove the following assertion:

**Theorem 1.1.** *Let  $U_1 \subset \mathbb{R}^2$  be the unit disk centered at the origin. For each positive integer  $m$ , there exists a  $C^1$ -function  $f : U_1 \rightarrow \mathbb{R}$  satisfying the following properties:*

- (1)  $f$  is real-analytic on  $U_1^* := U_1 \setminus \{(0, 0)\}$ ,
- (2)  $(0, 0, f(0, 0))$  is a  $C^1$ -umbilic of the graph of  $f$  with index  $1 + (m/2)$ .

It should be remarked that the inversion of the graph of Bates' function  $B(x, y)$  has a differentiable umbilic of index 2 although not of class  $C^1$  (see Example 2.3). It was classically known that curvature line flows are closely related to the eigenflows of the Hessian matrices of functions (see Appendix A). As an application of the above result, we can show the following:

**Corollary 1.2.** *For each  $m \geq 1$ , there exists a  $C^1$ -function  $\lambda : U_1 \rightarrow \mathbb{R}$  satisfying*

- (1)  $\lambda$  is real-analytic on  $U_1^*$ , and
- (2) the eigenflow of the Hessian matrix of  $\lambda$  has an isolated singular point  $(0, 0)$  with index  $1 + (m/2)$ .

When we consider the eigenflow of the Hessian matrix of  $f$ , it is well known that the index of the flow at an isolated singular point is equal to half of the index of the vector field

$$(1-2) \quad d_f := 2f_{xy} \frac{\partial}{\partial x} + (f_{yy} - f_{xx}) \frac{\partial}{\partial y}.$$

In addition, if  $o := (0, 0)$  is an isolated singular point of the eigenflow of the Hessian matrix of  $f$ , then its index is equal to  $1 + \text{ind}_o(\delta_f)/2$  (see Appendix B), where  $\text{ind}_o(\delta_f)$  is the index of the vector field

$$(1-3) \quad \delta_f := 2(rf_{r\theta} - f_\theta) \frac{\partial}{\partial x} + (-r^2 f_{rr} + rf_r + f_{\theta\theta}) \frac{\partial}{\partial y}$$

at  $o$ , and  $x = r \cos \theta$ ,  $y = r \sin \theta$ . In order to prove the above theorem, we introduce vector fields  $D_f$  and  $\Delta_f$  analogous to  $d_f$  and  $\delta_f$ , respectively (see Propositions 3.3 and 4.2), and prove the theorem by computing the index of  $\Delta_f$  at infinity for each of the functions (see Section 5)

$$(1-4) \quad f = f_m(r, \theta) := 1 + \tanh(r^a \cos m\theta), \quad 0 < a < 1/4, \quad m = 1, 2, \dots$$

We also give an alternative proof of Theorem 1.1 without use of inversion, by an explicit example of  $\lambda$ , see (6-1), satisfying (1) and (2) of Corollary 1.2 (see Section 6).

## 2. The regularity of the inversion

Let  $R$  be a positive number. Consider a function  $f : \mathbb{R}^2 \setminus \Omega_R \rightarrow \mathbb{R}$ , where

$$(2-1) \quad \Omega_R := \{(x, y) \in \mathbb{R}^2; \sqrt{x^2 + y^2} \leq R\}.$$

Then  $F = (x, y, f(x, y))$  gives a parametrization of the graph of  $f$ . The inversion of  $F$  is given by  $F/(F \cdot F)$ , where the dot denotes the inner product on  $\mathbb{R}^3$ . We consider the following coordinate change:

$$(2-2) \quad x = \frac{u}{u^2 + v^2}, \quad y = \frac{v}{u^2 + v^2}.$$

Then

$$(2-3) \quad \Psi_f := \frac{1}{\rho^2 \hat{f}^2 + 1} (u, v, \rho^2 \hat{f}), \quad \hat{f}(u, v) := f\left(\frac{u}{\rho^2}, \frac{v}{\rho^2}\right)$$

gives a parametrization of the inversion, where  $\rho := \sqrt{u^2 + v^2}$ . The map  $\Psi_f$  is defined on the domain

$$(2-4) \quad U_{1/R}^* := U_{1/R} \setminus \{o\}, \quad \left( U_{1/R} := \left\{ (u, v) \in \mathbb{R}^2; \sqrt{u^2 + v^2} < \frac{1}{R} \right\} \right),$$

where  $o := (0, 0)$ . If we set

$$(2-5) \quad x = r \cos \theta, \quad y = r \sin \theta,$$

where  $r > 0$ , then (2-2) yields

$$(2-6) \quad \rho = \frac{1}{r}, \quad u = \rho \cos \theta, \quad v = \rho \sin \theta.$$

In particular, the angular parameter is common in the  $xy$ -plane and the  $uv$ -plane.

**Proposition 2.1.** *Let  $f : \mathbb{R}^2 \setminus \Omega_R \rightarrow \mathbb{R}$  be a  $C^\infty$ -function such that  $f/r$  is bounded. Then the inversion  $\Psi_f : U_{1/R}^* \rightarrow \mathbb{R}^3$  can be continuously extended to  $(0, 0)$ , and moreover, if*

$$(2-7) \quad \left| \frac{f^2 - 2rff_r}{r^2} \right| < 1, \quad r > R,$$

then the image of  $\Psi_f = (X, Y, Z)$  can be locally expressed as the graph of a function  $Z = Z_f(X, Y)$  on a neighborhood of  $(0, 0)$  in the  $XY$ -plane. Under the assumption (2-7), the function  $Z_f(X, Y)$  is differentiable if and only if

$$\lim_{r \rightarrow \infty} \frac{f}{r} = 0.$$

*Proof.* We can write

$$(2-8) \quad \Psi_f(u, v) = \frac{1}{1 + \varphi(u, v)^2} \left( u, v, \varphi(u, v) \sqrt{u^2 + v^2} \right),$$

where

$$(2-9) \quad \varphi(u, v) = \sqrt{u^2 + v^2} \hat{f}(u, v) = \frac{f(x, y)}{r}.$$

Since  $f/r$  is bounded, the function  $\varphi$  is bounded on  $U_{1/R}^*$ . Thus, using (2-8), we can prove  $\lim_{\rho \rightarrow 0} \Psi_f = (0, 0, 0)$ , i.e.,  $\Psi_f(u, v)$  can be continuously extended to  $(0, 0)$ . We denote by  $\Pi : \mathbb{R}^3 \ni (x, y, z) \mapsto (x, y) \in \mathbb{R}^2$  the orthogonal projection. Setting

$$\psi(\rho, \theta) := \frac{\rho}{1 + \varphi(\rho \cos \theta, \rho \sin \theta)^2},$$

it holds that

$$(2-10) \quad \Pi \circ \Psi_f(u, v) = (\psi(\rho, \theta) \cos \theta, \psi(\rho, \theta) \sin \theta).$$

Since  $\hat{f}(\rho \cos \theta, \rho \sin \theta) = f(\cos \theta / \rho, \sin \theta / \rho)$ , we have

$$\varphi_\rho = f - r f_r.$$

In particular, it holds that

$$\psi_\rho = \frac{1 - (f^2 - 2r f f_r) / r^2}{(1 + f^2 / r^2)^2}.$$

By (2-7), there exists  $\varepsilon > 0$  such that  $\rho \mapsto \psi(\rho, \theta)$ ,  $|\rho| \leq \varepsilon$ , is a monotone increasing function for each  $\theta$ . Thus, by (2-10), we can conclude that  $\Pi \circ \Psi_f : \bar{U}_\varepsilon \rightarrow \mathbb{R}^2$  is an injection. Since a continuous bijection from a compact space to a Hausdorff space is a homeomorphism, the inverse map  $G : \Omega \rightarrow U_\varepsilon$  of  $\Pi \circ \Psi_f|_{U_\varepsilon}$  is continuous, where  $\Omega$  is a neighborhood of the origin of the  $XY$ -plane in  $\mathbb{R}^3$ . Then the graph of

$$(2-11) \quad Z_f \left( = \frac{\rho \varphi}{1 + \varphi^2} \right) = \frac{\varphi(G(X, Y)) \rho(G(X, Y))}{1 + \varphi(G(X, Y))^2}$$

coincides with the image of  $\Psi_f = (X, Y, Z)$  around  $(0, 0, 0)$ . Then

$$X = \frac{u}{1 + \varphi^2}, \quad Y = \frac{v}{1 + \varphi^2}, \quad Z = \frac{\rho \varphi}{1 + \varphi^2}.$$

Since  $\rho \rightarrow 0$  as  $(X, Y) \rightarrow (0, 0)$ , we obtain

$$(2-12) \quad \lim_{(X, Y) \rightarrow (0, 0)} \frac{Z_f(X, Y)}{\sqrt{X^2 + Y^2}} = \lim_{(X, Y) \rightarrow (0, 0)} \frac{\varphi \rho}{\sqrt{u^2 + v^2}} = \lim_{\rho \rightarrow 0} \varphi = \lim_{r \rightarrow \infty} \frac{f}{r}. \quad \square$$

**Corollary 2.2.** *Suppose that  $f : \mathbb{R}^2 \setminus \Omega_R \rightarrow \mathbb{R}$  is a bounded  $C^\infty$ -function satisfying*

$$(2-13) \quad \lim_{r \rightarrow \infty} \frac{f_r}{r} = 0.$$

*Then the inversion  $\Psi_f : U_{1/R}^* \rightarrow \mathbb{R}^3$  can be continuously extended to  $(0, 0)$ , and moreover, the image of  $\Psi_f$  is locally a graph which is differentiable at  $(0, 0)$ .*

**Example 2.3.** Bates' example mentioned in the introduction is differentiable. In fact,  $B(x, y)$  in (1-1) is bounded and  $B_r/r$  converges to zero as  $r \rightarrow \infty$ . However, the inversion of  $(x, y, B(x, y))$  is not  $C^1$ . In fact, the unit normal vector field of the graph of  $B$  is not continuously extended to the point at infinity. Since the inversion preserves the angle, the unit normal vector field of its inversion cannot be continuously extended to  $(0, 0, 0)$ .

**Example 2.4.** Ghomi and Howard [2012] gave an example:

$$(2-14) \quad f_{\text{GH}} = 1 + \lambda \frac{1 + x + y^2}{\sqrt{1 + (x + y^2)^2}}, \quad (\lambda > 0).$$

The graph of  $f_{\text{GH}}$  is umbilic-free (see Example 3.5 in Section 3). The function  $f_{\text{GH}}$  is bounded. In addition, since  $(f_{\text{GH}})_r$  is bounded, (2-13) is obvious. Therefore, as pointed out in [Ghomi and Howard 2012], the inversion of  $(x, y, f_{\text{GH}}(x, y))$  is differentiable. However, it is not a  $C^1$ -map. In fact, the limit of the unit normal vector field along  $y = 0$  of the graph of  $f_{\text{GH}}$  is not equal to that along  $x + y^2 = 0$  at the point at infinity.

Next, we give a condition for  $\Psi_f$  to be extendable as a  $C^1$ -map to  $(0, 0)$ .

**Proposition 2.5.** *Suppose that  $f : \mathbb{R}^2 \setminus \Omega_R \rightarrow \mathbb{R}$  is a bounded  $C^\infty$ -function satisfying*

- (a)  $\lim_{r \rightarrow \infty} f_r = 0$ ,
- (b)  $\lim_{r \rightarrow \infty} f_\theta / r = 0$ .

*Then  $\Psi_f = (X, Y, Z)$  can be extended to  $(0, 0)$  as a  $C^1$ -map. Moreover, the map  $(u, v) \mapsto (X(u, v), Y(u, v))$  is a  $C^1$ -diffeomorphism from a neighborhood of the origin in the  $uv$ -plane onto a neighborhood of the origin in the  $XY$ -plane.*

To prove this, we prepare the following lemma.

**Lemma 2.6.** *The conditions (a) and (b) in Proposition 2.5 are equivalent to the following two conditions, respectively:*

- (1)  $\lim_{\rho \rightarrow 0} \rho^2 \hat{f}_\rho = 0$ ,
- (2)  $\lim_{\rho \rightarrow 0} \rho \hat{f}_\theta = 0$ .

*Proof.* The equivalency of (2) and (b) is obvious. The equivalency of (1) and (a) follows from the identity  $\hat{f}_\rho = -f_r / \rho^2$ . □

*Proof of Proposition 2.5.* We see by Corollary 2.2 that  $\Psi_f$  can be extended to  $(0, 0)$  as a differentiable map and that the map  $(u, v) \mapsto (X(u, v), Y(u, v))$  is a homeomorphism from a neighborhood of  $(0, 0)$  onto a neighborhood of  $(0, 0)$ . We set

$$(2-15) \quad h := \rho^2 \hat{f} (= \rho\varphi), \quad k := (\rho \hat{f})^2 (= \varphi^2).$$

By (2-3), we can write

$$(2-16) \quad \Psi_f = (X, Y, Z) = \frac{1}{k+1}(u, v, h).$$

To show that  $\Psi_f$  is a  $C^1$ -map at  $(0, 0)$ , it is sufficient to show that  $h$  and  $k$  are  $C^1$ -functions. Since  $h$  and  $k$  are  $C^\infty$ -functions on  $U_{1/R}^*$ , they satisfy

$$(2-17) \quad \begin{aligned} h_u &= \rho((2\hat{f} + \rho\hat{f}_\rho)\cos\theta - \hat{f}_\theta\sin\theta), \\ h_v &= \rho((2\hat{f} + \rho\hat{f}_\rho)\sin\theta + \hat{f}_\theta\cos\theta), \end{aligned}$$

$$(2-18) \quad \begin{aligned} k_u &= 2\hat{f}\rho(\cos\theta(\hat{f} + \rho\hat{f}_\rho) - \hat{f}_\theta\sin\theta), \\ k_v &= 2\hat{f}\rho(\sin\theta(\hat{f} + \rho\hat{f}_\rho) + \hat{f}_\theta\cos\theta), \end{aligned}$$

on  $U_{1/R}^*$ . Using (1), (2) in Lemma 2.6, (2-17) and (2-18), one can easily see that

$$(2-19) \quad \lim_{\rho \rightarrow 0} h_u = \lim_{\rho \rightarrow 0} h_v = \lim_{\rho \rightarrow 0} k_u = \lim_{\rho \rightarrow 0} k_v = 0,$$

which shows that  $\Psi_f$  extends to  $(0, 0)$  as a  $C^1$ -map. By (2-16) and (2-19), we have

$$X_u(0, 0) = 1, \quad X_v(0, 0) = 0, \quad Y_u(0, 0) = 0, \quad Y_v(0, 0) = 1.$$

Thus the second assertion follows from the inverse mapping theorem, because the Jacobi matrix of the map  $(u, v) \mapsto (X(u, v), Y(u, v))$  is regular at  $(0, 0)$ .  $\square$

In Section 5, we need the following:

**Proposition 2.7.** *Let  $f : \mathbb{R}^2 \setminus \Omega_R \rightarrow \mathbb{R}$  be a bounded  $C^\infty$ -function satisfying conditions (a) and (b) of Proposition 2.5. If there exists a constant  $0 \leq c < \frac{1}{2}$  such that*

$$r^{1-c/2}f_r, \quad r^{-c/2}f_\theta, \quad r^{2-c}f_{rr}, \quad r^{1-c}f_{r\theta}, \quad r^{-c}f_{\theta\theta}$$

*are bounded on  $\mathbb{R}^2 \setminus \Omega_R$ , then the map  $(u, v) \mapsto (X(u, v), Y(u, v))$  is a  $C^2$ -map at  $(0, 0)$ , where  $\Psi_f = (X, Y, Z)$ .*

We prepare the following lemmas:

**Lemma 2.8.** *The boundedness of the five functions in Proposition 2.7 is equivalent to the boundedness of the functions*

$$(2-20) \quad \rho^{1+c/2}\hat{f}_\rho, \quad \rho^{c/2}\hat{f}_\theta, \quad \rho^{2+c}\hat{f}_{\rho\rho}, \quad \rho^{1+c}\hat{f}_{\rho\theta}, \quad \rho^c\hat{f}_{\theta\theta}$$

*on  $U \setminus \{(0, 0)\}$ , where  $U$  is a sufficiently small neighborhood of  $(0, 0)$ .*

*Proof.* Differentiating  $\hat{f} = \hat{f}(\rho \cos\theta, \rho \sin\theta)$  by  $\rho$ , we get  $\rho\hat{f}_\rho = -rf_r$  and  $\rho^2\hat{f}_{\rho\rho} = 2rf_r + r^2f_{rr}$ , which can be used to check the assertion.  $\square$



**Lemma 2.9.** *Suppose that the five functions in (2-20) are bounded on  $U \setminus \{(0, 0)\}$ . Then  $\rho^{2c}k_{uu}$ ,  $\rho^{2c}k_{uv}$  and  $\rho^{2c}k_{vv}$  are also bounded on  $U \setminus \{(0, 0)\}$ , where  $k$  is the function given in (2-15).*

*Proof.* In fact, each of  $k_{uu}$ ,  $k_{uv}$ ,  $k_{vv}$  is written as a linear combination of

$$1, \quad \rho \hat{f}_\rho, \quad \hat{f}_\theta, \quad (\rho \hat{f}_\rho)^2, \quad \rho \hat{f}_\rho \hat{f}_\theta, \quad \hat{f}_\theta^2, \quad \rho^2 \hat{f}_{\rho\rho}, \quad \rho \hat{f}_{\rho\theta}, \quad \hat{f}_{\theta\theta},$$

with coefficients that are bounded functions. For example,

$$k_{uv} = \sin 2\theta (\rho^2 \hat{f}_\rho^2 + \hat{f}_\theta (\rho^2 \hat{f}_{\rho\rho} + 3\rho \hat{f}_\rho - \hat{f}_{\theta\theta}) - \hat{f}_\theta^2) + 2\cos 2\theta (\hat{f}_\theta (\rho \hat{f}_\rho + \hat{f}_\theta) + \rho \hat{f}_\theta \hat{f}_{\rho\theta}).$$

Thus, we get the assertion.  $\square$

*Proof of Proposition 2.7.* By Lemmas 2.8 and 2.9, the fact that  $2c < 1$  yields that

$$(2-21) \quad \lim_{\rho \rightarrow 0} \rho k_{uu} = \lim_{\rho \rightarrow 0} \rho k_{uv} = \lim_{\rho \rightarrow 0} \rho k_{vv} = 0.$$

Since

$$\begin{aligned} X_{uu} &= \frac{2uk_u^2 - 2(k+1)k_u - u(k+1)k_{uu}}{(k+1)^3}, \\ X_{uv} &= -\frac{k_v(-2uk_u + k+1) + u(k+1)k_{uv}}{(k+1)^3}, \\ X_{vv} &= -\frac{u((k+1)k_{vv} - 2k_v^2)}{(k+1)^3}, \end{aligned}$$

we have that  $X_{uu}$ ,  $X_{uv}$ ,  $X_{vv}$  tend to 0 as  $\rho \rightarrow 0$ . This implies that  $X_u$ ,  $X_v$  are  $C^1$ -functions. Similarly,  $Y_u$ ,  $Y_v$  are also  $C^1$ -functions.  $\square$

### 3. The pair of identifiers for umbilics

Let  $U$  be a domain on  $\mathbb{R}^2$ . Consider a flow (i.e., a 1-dimensional foliation)  $\mathcal{F}$  defined on  $U \setminus \{p_1, \dots, p_n\}$ , where  $p_1, \dots, p_n$  are distinct points in  $U$ . We are interested in the case where  $\mathcal{F}$  is

- the curvature line flow of an immersion  $P : U \rightarrow \mathbb{R}^3$ ,
- the eigenflow of a matrix-valued function on  $U$ , or
- the flow induced by a vector field on  $U$ .

We fix a simple closed smooth curve  $\gamma : T^1 \rightarrow U \setminus \{p_1, \dots, p_n\}$ , where  $T^1 := \mathbb{R}/2\pi\mathbb{Z}$ . We set

$$\partial_x := \frac{\partial}{\partial x}, \quad \partial_y := \frac{\partial}{\partial y}.$$

Then one can take a smooth vector field

$$V(t) := a(t)\partial_x + b(t)\partial_y$$

along the curve  $\gamma(t)$  such that  $V(t)$  is a nonzero tangent vector of  $\mathbb{R}^2$  at  $\gamma(t)$  which points in the direction of the flow  $\mathcal{F}$ . Then the map

$$(3-1) \quad \check{V} : T^1 \ni t \mapsto \frac{(a(t), b(t))}{\sqrt{a(t)^2 + b(t)^2}} \in S^1 := \{\mathbf{x} \in \mathbb{R}^2; |\mathbf{x}| = 1\}$$

is called the *Gauss map* of  $\mathcal{F}$  with respect to the curve  $\gamma$ . The mapping degree of the map  $\check{V}$  is called the *rotation index* of  $\mathcal{F}$  with respect to  $\gamma$  and denoted by  $\text{ind}(\mathcal{F}, \gamma)$ , which is a half-integer, in general. If  $\gamma$  surrounds only  $p_j$ , then  $\text{ind}(\mathcal{F}, \gamma)$  is independent of the choice of such a curve  $\gamma$ . So we call it the (*rotation*) *index* of the flow  $\mathcal{F}$  at  $p_j$ , and it is denoted by  $\text{ind}_{p_j}(\mathcal{F})$ . If the flow  $\mathcal{F}$  is generated by a vector field  $V$  defined on  $U \setminus \{p_1, \dots, p_n\}$ , then  $\text{ind}_{p_j}(\mathcal{F})$  is an integer, and we denote it by  $\text{ind}_{p_j}(V)$ .

We denote by  $S_2(\mathbb{R})$  the set of real symmetric 2-matrices. Let  $U$  be a domain in  $\mathbb{R}^2$ , and

$$A = \begin{pmatrix} a_{11}(x, y) & a_{12}(x, y) \\ a_{12}(x, y) & a_{22}(x, y) \end{pmatrix} : U \rightarrow S_2(\mathbb{R}),$$

a  $C^\infty$ -map. A point  $p \in U$  is called an *equidiagonal point* of  $A$  if  $a_{11} = a_{22}$  and  $a_{12} = 0$  at  $p$ . We now suppose that  $p$  is an isolated equidiagonal point. Without loss of generality, we may assume that  $A$  has no equidiagonal points on  $U \setminus \{p\}$ . Since two eigenflows of  $A$  are mutually orthogonal, the indices of the two eigenflows of the  $S_2(\mathbb{R})$ -valued function  $A$  are the same half-integer at  $p$ . We denote it by  $\text{ind}_p(A)$ .

It is well known that for an  $S_2(\mathbb{R})$ -valued function  $A$ , the formula

$$(3-2) \quad \text{ind}_p(A) = \frac{1}{2} \text{ind}_p(\mathbf{v}_A)$$

holds, where  $\mathbf{v}_A$  is the vector field on  $U$  given by

$$(3-3) \quad \mathbf{v}_A := (a_{11} - a_{22})\partial_x + a_{12}\partial_y.$$

We shall apply these facts to the computation of the indices of isolated umbilics on regular surfaces in  $\mathbb{R}^3$  as follows. Let  $f : U \rightarrow \mathbb{R}$  be a  $C^\infty$ -function. The symmetric matrices associated with the first and the second fundamental forms of the graph of  $f$  are given by

$$(3-4) \quad I := \begin{pmatrix} 1 + f_x^2 & f_x f_y \\ f_x f_y & 1 + f_y^2 \end{pmatrix}, \quad II := \begin{pmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{pmatrix}.$$

We consider a  $\text{GL}(2, \mathbb{R})$ -valued function

$$(3-5) \quad P := \begin{pmatrix} 0 & \sqrt{1 + f_x^2} \\ -\sqrt{(1 + f_x^2 + f_y^2)/(1 + f_x^2)} & f_x f_y / \sqrt{1 + f_x^2} \end{pmatrix},$$

which satisfies the identity  $PP^T = I$ , where  $P^T$  is the transpose of  $P$ . Then

$$A_f := P^{-1}II(P^T)^{-1} = P^T(I^{-1}II)(P^T)^{-1}$$

is an  $S_2(\mathbb{R})$ -valued function. The umbilics of the graph of  $f$  correspond to the equidiagonal points of  $A_f$ . We show the following:

**Proposition 3.1.** *The symmetric matrix  $A_f(p)$  is proportional to the identity matrix at  $p \in U$  if and only if  $p$  gives an umbilic of the graph of  $f$ . Moreover, if  $p$  is an isolated umbilic, then  $\text{ind}_p(A_f)$  coincides with the index of the umbilic  $p$ .*

*Proof.* The first assertion follows from the definition of  $A_f$ . Without loss of generality, we may assume that  $p$  coincides with the origin  $o := (0, 0)$ , and the graph of  $f$  has no umbilics other than  $o$  on  $U$ . Take a sufficiently small positive number  $\varepsilon > 0$  so that the circle

$$\gamma(t) = \varepsilon(\cos t, \sin t), \quad 0 \leq t \leq 2\pi,$$

is null-homotopic in  $U$ .

We denote by  $(a_1(t), b_1(t))^T$  and  $(a_2(t), b_2(t))^T$ , eigenvectors of  $I^{-1}II$  and  $A_f$  at  $\gamma(t)$ , respectively. We may suppose

$$(a_1(t), b_1(t))P(\gamma(t)) = (a_2(t), b_2(t)), \quad 0 \leq t \leq 2\pi.$$

We set

$$\mathbf{w}_i(t) := a_i(t)\partial_x + b_i(t)\partial_y, \quad i = 1, 2.$$

Then  $\mathbf{w}_1$  points in one of the principal directions of the graph of  $f$ . The matrix  $P(\gamma(t))$  takes values in the set

$$(3-6) \quad \mathcal{T} := \left\{ \begin{pmatrix} 0 & x \\ -y & z \end{pmatrix}; x, y > 0, z \in \mathbb{R} \right\}.$$

Since the set  $\mathcal{T}$  is null-homotopic, the mapping degree of  $\check{\mathbf{w}}_1(t)$  with respect to the origin is equal to that of  $\check{\mathbf{w}}_2(t)$ . Since the degree of  $\check{\mathbf{w}}_2(t)$  with respect to  $o$  coincides with  $\text{ind}_o(A_f)$ , we get the second assertion.  $\square$

By a straightforward calculation, one can get the following identity:

$$\tilde{A}_f := hk^3 A_f = \begin{pmatrix} f_x f_y (f_x f_y f_{xx} - 2hf_{xy}) + h^2 f_{yy} & lk \\ lk & k^2 f_{xx} \end{pmatrix},$$

where

$$h := 1 + f_x^2, \quad k := \sqrt{1 + f_x^2 + f_y^2}, \quad l := -hf_{xy} + f_x f_y f_{xx}.$$

Then the coefficients of the vector field

$$\mathbf{v}_{\tilde{A}_f} = v_1 \partial_x + v_2 \partial_y$$

defined as in (3-3) for  $A = \tilde{A}_f$  are given by

$$\begin{aligned} v_1 &= \tilde{a}_{11} - \tilde{a}_{22} = (-1 + f_x^2) f_y^2 f_{xx} - h f_{xx} - 2h f_x f_{xy} f_y + h^2 f_{yy}, \\ v_2 &= \tilde{a}_{12} = -k(h f_{xy} - f_x f_y f_{xx}), \end{aligned}$$

where  $\tilde{A}_f = (\tilde{a}_{ij})_{i,j=1,2}$ . Hence, we get the following identity:

$$v_1 = \frac{2f_x f_y}{k} v_2 + h(-f_{xx}(1 + f_y^2) + (1 + f_x^2) f_{yy}).$$

Consequently, we get the following fact (see [Ghomi and Howard 2012, (10)]):

**Fact 3.2.** *The graph of the function  $z = f(x, y)$  defined on  $U$  has an umbilic at  $p \in U$  if and only if the functions*

$$d_1(x, y) := (1 + f_x^2) f_{xy} - f_x f_y f_{xx}, \quad \text{and} \quad d_2(x, y) := (1 + f_x^2) f_{yy} - f_{xx}(1 + f_y^2)$$

*both vanish at  $p$ .*

We consider the vector field

$$D_f := d_1 \partial_x + d_2 \partial_y$$

defined on the domain  $U$  in the  $xy$ -plane. Suppose that  $p$  is a zero of  $D_f$ . The following assertion holds:

**Proposition 3.3.** *If  $p$  gives an isolated umbilic of the graph of  $f$ , then half of the index of the vector field  $D_f$  at  $p$  coincides with the index of the umbilic  $p$ .*

*Proof.* The half of the index of the vector field

$$X := -\mathbf{v}_{\tilde{A}_f} = (2f_x f_y d_1 - h d_2) \partial_x + k d_1 \partial_y$$

at  $p$  is equal to  $\text{ind}_p(\tilde{A}_f)$ . We now set

$$X_s := (\partial_x, \partial_y) \begin{pmatrix} \frac{2s f_x f_y}{\sqrt{1 + s(f_x^2 + f_y^2)}} & -1 - s f_x^2 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} d_1 \\ d_2 \end{pmatrix}, \quad 0 \leq s \leq 1.$$

Then  $X = X_1$  and  $X_0 = -d_2 \partial_x + d_1 \partial_y$ , and the rotation index of  $X_s$  at  $p$  does not depend on  $s \in [0, 1]$ . Since the rotation index of  $D_f = (d_1, d_2)$  at  $p$  coincides with that of  $X_0$ , we can conclude that  $X$  has the same rotation index as  $D_f$  at  $p$ .  $\square$

We call  $d_1, d_2$  the *Cartesian umbilic identifiers* of the function  $f$ .

**Example 3.4.** For a function  $f(x, y) := \text{Re}(z^3) = x^3 - 3xy^2$  ( $z = x + iy$ ), the Cartesian umbilic identifiers are given by  $d_1 = -6y\varphi_1$ ,  $d_2 = -6x\varphi_2$ , where

$$\varphi_1 := -9x^4 + 9y^4 + 1, \quad \varphi_2 := 9x^4 + 18x^2y^2 + 9y^4 + 2.$$

Since  $\varphi_i$ ,  $i = 1, 2$ , are positive at the origin  $(0, 0)$ , the vector field  $D_f$  can be continuously deformed into the vector field  $-y\partial_x - x\partial_y$  preserving the property that the origin is an isolated zero. Thus  $D_f$  is of index  $-1$ , and the graph of the function  $f$  has an isolated umbilic of index  $-\frac{1}{2}$  at the origin.

**Example 3.5.** Bates' function  $B(x, y)$  has no umbilics since  $d_1 > 0$  on  $\mathbb{R}^2$ . On the other hand, the identifier  $d_1$  with respect to Ghomi and Howard's function  $f_{GH}(x, y)$  in (2-14) vanishes if and only if  $y = 0$  or  $x = -y^2$ . Since  $d_2$  never vanishes on these two sets, the graph of  $f_{GH}$  also has no umbilics on  $\mathbb{R}^2$ .

#### 4. The pair of polar identifiers for umbilics

Let  $U$  be a domain in the  $xy$ -plane, and  $f : U \rightarrow \mathbb{R}$  a  $C^\infty$ -function. Let  $(r, \theta)$  be the polar coordinate system associated to  $(x, y)$  as in (2-5). Then

$$F(r, \theta) := (r \cos \theta, r \sin \theta, f(r \cos \theta, r \sin \theta))$$

gives a parametrization of the graph of  $f$  with the unit normal vector

$$v := \frac{1}{\sqrt{f_\theta^2 + r^2(1 + f_r^2)}}(f_\theta \sin \theta - r f_r \cos \theta, -r f_r \sin \theta - f_\theta \cos \theta, r).$$

Then

$$\hat{I} := \begin{pmatrix} 1 + f_r^2 & f_r f_\theta \\ f_r f_\theta & r^2 + f_\theta^2 \end{pmatrix}$$

is the symmetric matrix consisting of the coefficients of the first fundamental form of  $F$ . If we set

$$Q = \begin{pmatrix} 0 & \sqrt{1 + f_r^2} \\ -\sqrt{f_\theta^2 + r^2(1 + f_r^2)}/\sqrt{1 + f_r^2} & f_r f_\theta / \sqrt{1 + f_r^2} \end{pmatrix},$$

then  $QQ^T = \hat{I}$ . The symmetric matrix consisting of the coefficients of the second fundamental form is given by

$$\hat{\Pi} := \frac{1}{\sqrt{f_\theta^2 + r^2(1 + f_r^2)}} \begin{pmatrix} r f_{rr} & r f_{r\theta} - f_\theta \\ r f_{r\theta} - f_\theta & r(f_{\theta\theta} + r f_r) \end{pmatrix}.$$

Then the symmetric matrix

$$B_f = Q^{-1} \hat{\Pi} (Q^{-1})^T = Q^T (\hat{I}^{-1} \hat{\Pi}) (Q^T)^{-1}$$

satisfies

$$\tilde{B}_f = \hat{h} \hat{k}^3 B_f = \begin{pmatrix} r f_r^2 f_\theta^2 f_{rr} + \hat{h} f_r (-2r f_\theta f_{r\theta} + 2f_\theta^2 + r^2 \hat{h}) + r \hat{h}^2 f_{\theta\theta} & \hat{I} \hat{k} \\ \hat{I} \hat{k} & r \hat{k}^2 f_{rr} \end{pmatrix},$$

where

$$\hat{h} := 1 + f_r^2, \quad \hat{k} := \sqrt{f_\theta^2 + r^2(1 + f_r^2)}, \quad \hat{l} := f_\theta(\hat{h} + rf_r f_{rr}) - r\hat{h}f_{r\theta}.$$

The following holds:

**Proposition 4.1.** *The symmetric matrix  $\tilde{B}_f(p)$  is proportional to the identity matrix at  $p \in U \setminus \{o\}$  if and only if  $p$  gives an umbilic of the graph of  $f$ . Moreover, if  $o$  is an isolated umbilic of the graph of  $f$ , then the index of the umbilic at  $o$  is equal to  $1 + \text{ind}_o(\tilde{B}_f)$ .*

*Proof.* The first assertion follows from the above discussions. So we now prove the second assertion. Suppose  $o$  is an isolated umbilic. We take a simple closed smooth curve  $\gamma(t)$  in the  $xy$ -plane, where  $0 \leq t \leq 2\pi$ , which surrounds the origin  $o$  anticlockwisely, and does not surround any other umbilics. Let  $\mathbf{w}_1 : [0, 2\pi] \rightarrow \mathbb{R}^2$  be a vector field along  $\gamma$  such that  $\mathbf{w}_1(t)$  is an eigenvector of the matrix  $I^{-1}II$  at  $\gamma(t)$  for each  $t \in [0, 2\pi]$ . Since

$$\begin{aligned} \partial_r &= \cos \theta \partial_x + \sin \theta \partial_y, \\ \partial_\theta &= -r \sin \theta \partial_x + r \cos \theta \partial_y, \end{aligned}$$

we have that

$$(\partial_r, \partial_\theta) = (\partial_x, \partial_y)T_0, \quad T_0 := \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix}.$$

Then, it holds that

$$\hat{I}^{-1}\hat{II} = (T_0)^{-1}(I^{-1}II)T_0.$$

In particular,

$$\mathbf{w}_2(t) := T_0(\gamma(t))^{-1}\mathbf{w}_1(t), \quad 0 \leq t \leq 2\pi,$$

gives an eigenvector of the matrix  $\hat{I}^{-1}\hat{II}$  at  $\gamma(t)$ . Let  $T_s : U \rightarrow \text{GL}(2, \mathbb{R})$ ,  $0 \leq s \leq 1$ , be a map defined by

$$T_s := \begin{pmatrix} \cos \theta & -(r(1-s) + s) \sin \theta \\ \sin \theta & (r(1-s) + s) \cos \theta \end{pmatrix}, \quad 0 \leq s \leq 1.$$

Then it gives a continuous deformation of  $T_0$  to the rotation matrix  $T_1$ . Since the winding number of the curve  $\gamma(t)$  with respect to the origin  $o$  is equal to 1, the difference between the rotation indices of  $\mathbf{w}_1$  and  $\mathbf{w}_2$  is equal to 1. Since the eigenflow of the symmetric matrix  $\tilde{B}_f$  is associated with that of the matrix  $\hat{I}^{-1}\hat{II}$  by  $Q$ , the fact that  $Q$  takes values in the set  $\mathcal{T}$  in Section 3 yields that the index of the umbilic  $o$  is equal to  $1 + \text{ind}_o(\tilde{B}_f)$ .  $\square$

We now set

$$\delta_1 := -\tilde{b}_{12}/\hat{k} = -f_\theta(1 + f_r^2 + rf_r f_{rr}) + r(1 + f_r^2)f_{r\theta},$$

where  $\tilde{B}_f = (\tilde{b}_{ij})_{i,j=1,2}$ . Then we have

$$\tilde{b}_{11} - \tilde{b}_{22} = -2f_r f_\theta \delta_1 + r(1 + f_r^2)\delta_2,$$

where

$$\delta_2 := (1 + f_r^2)(rf_r + f_{\theta\theta}) - f_{rr}(r^2 + f_\theta^2).$$

Thus, as in the proof of Proposition 3.3, we get the following assertion:

**Proposition 4.2.** *Let  $U$  be a neighborhood of the origin  $o := (0, 0)$ . Let  $f : U \rightarrow \mathbb{R}$  be a  $C^\infty$ -function. Then the graph of  $f$  has an umbilic at  $p \in U \setminus \{o\}$  if and only if the two functions  $\delta_1(r, \theta)$ ,  $\delta_2(r, \theta)$  both vanish at  $p$ , where  $x = r \cos \theta$  and  $y = r \sin \theta$ . Further, if  $o$  is an isolated umbilic, then half of the index of the vector field*

$$\Delta_f := \delta_1 \partial_x + \delta_2 \partial_y$$

at  $o$  equals  $-1 + I_f(o)$ , where  $I_f(o)$  is the index of the umbilic  $o$ .

We call  $\delta_1, \delta_2$  the *polar umbilic identifiers* of the function  $f$ .

**Example 4.3.** Consider the function (where  $z = x + iy$ )

$$f(x, y) := \operatorname{Re}(z^2 \bar{z}) = x^3 + xy^2 = r^3 \cos \theta.$$

By straightforward calculations, we have


$$\delta_1 = -2r^3 \sin \theta, \quad \delta_2 = -2r^3(2 - 3r^4 - 6r^4 \cos 2\theta) \cos \theta.$$

Since  $2 - 3r^4 - 6r^4 \cos 2\theta$  is positive for sufficiently small  $r > 0$ , the vector field  $\Delta_f$  can be continuously deformed into the vector field  $-\sin \theta \partial_r - \cos \theta \partial_\theta$  preserving the property that the origin is an isolated zero. Thus the rotation index of  $\Delta_f$  at  $o$  is equal to  $-1$ , and  $I_f(o) = 1 - \frac{1}{2} = \frac{1}{2}$ .

We give a generalization of Proposition 4.2 for the computation of the index of the curvature line flow of a surface along an arbitrarily given simple closed curve surrounding several umbilics as follows. Let  $z = f(x, y)$  be a  $C^\infty$ -function defined on  $\mathbb{R}^2$  admitting only isolated umbilics. Suppose that  $\gamma : \mathbb{R} \rightarrow \mathbb{R}^2$  is a  $C^\infty$ -map satisfying  $\gamma(t + 2\pi) = \gamma(t)$  which gives a simple closed curve in the  $xy$ -plane such that it surrounds a bounded domain containing the origin  $o$  anticlockwisely. Moreover, we assume that  $\gamma(t)$  does not pass through any points corresponding to umbilics of the graph of  $f$ . We denote by  $I_f(\gamma)$  (resp.  $\operatorname{ind}_\gamma(\Delta_f)$ ) the rotation index of the curvature line flow (resp. of the vector field  $\Delta_f$ ) along the simple closed curve  $\gamma$ . Then the formula

$$(4-1) \quad I_f(\gamma) = 1 + \frac{\operatorname{ind}_\gamma(\Delta_f)}{2}$$

can be proved by modifying the proof of Proposition 4.2. Suppose that there exist at most finitely many points  $t = t_1, \dots, t_k \in [0, 2\pi]$  such that  $\delta_1(\gamma(t))$  vanishes



**Figure 1.** The inversion of the graph  $f_5$  for  $a = \frac{1}{5}$  (left) and its enlarged view (right). In these two figures, the  $z$ -axis points toward the downward direction.

at  $t = t_j$ . We now assume that  $\delta'_1(\gamma(t)) := d\delta_1(\gamma(t))/dt$  does not vanish at  $t = t_j$ , for  $j = 1, \dots, k$ . We set

$$\varepsilon(t_j) = \begin{cases} 0 & \text{for } \delta_2(\gamma(t_j)) < 0, \\ 1 & \text{for } \delta'_1(\gamma(t_j)) > 0 \text{ and } \delta_2(\gamma(t_j)) > 0, \\ -1 & \text{for } \delta'_1(\gamma(t_j)) < 0 \text{ and } \delta_2(\gamma(t_j)) > 0. \end{cases}$$

Then, it holds that

$$(4-2) \quad \text{ind}_\gamma(\Delta_f) = - \sum_{j=1}^k \varepsilon(t_j).$$

## 5. Proof of the main theorem

In this section, using the function  $f = f_m$  ( $m = 1, 2, 3, \dots$ ) given in (1-4), we prove Theorem 1.1 and Corollary 1.2 in the introduction. More generally, we consider the function

$$(5-1) \quad g = g_m(r, \theta) := 1 + F(r^a \cos m\theta), \quad 0 < a < 1/4, \quad m = 1, 2, 3, \dots,$$

which is defined on  $\{(r, \theta); r > R\}$ , where  $R$  is an arbitrarily fixed positive number, and  $F : \mathbb{R} \rightarrow \mathbb{R}$  is a bounded  $C^\infty$ -function satisfying the following conditions:

- (i)  $F(x)$  is an odd function, that is, it satisfies  $F(-x) = -F(x)$ ,
- (ii) the derivative  $F'(x)$  of  $F$  is a positive-valued bounded function on  $\mathbb{R}$ ,
- (iii) the second derivative  $F''(x)$  is a bounded function on  $\mathbb{R}$  such that  $F''(x) < 0$  for  $x > 0$ ,



(iv) there exist three constants  $\alpha$ ,  $\beta$  and  $\gamma$  ( $\beta \neq 0$ ,  $\gamma > 0$ ) such that

$$\lim_{x \rightarrow \infty} e^{\gamma x} F'(x) = \alpha, \quad \lim_{x \rightarrow \infty} e^{\gamma x} F''(x) = \beta.$$

One can easily construct a bounded  $C^\infty$ -function  $F(x)$  satisfying properties (i–iv). For example, one can construct an odd  $C^\infty$ -function satisfying (ii) and (iii) so that

$$F(x) = 1 - e^{-x}, \quad x \in [M, \infty),$$

for a positive number  $M$ . Then it satisfies (iv) also. However, to prove Theorem 1.1, we must choose the function  $F(x)$  to be real-analytic, and

$$F(x) := \tanh x$$

satisfies all of the properties required. In this case,  $g_m = f_m$  holds. From now on, we shall prove Theorem 1.1 and Corollary 1.2 using only the above four properties of  $F(x)$ .

The function  $g$  can be considered as a  $C^\infty$ -function on  $\mathbb{R}^2 \setminus \Omega_R$  in the  $xy$ -plane for any  $R > 0$ . The graph of  $g$  lies between two parallel planes orthogonal to the  $z$ -axis, and is symmetric under rotation by the angle  $2\pi/m$  with respect to the  $z$ -axis (the entire figure of the inversion of the graph of  $f_5$  is given in the left-hand side of Figure 1). The partial derivatives of the function  $g$  are given by

$$\begin{aligned} g_r &= ar^{a-1}c_m F'(r^a c_m), \\ g_\theta &= -mr^a s_m F'(r^a c_m), \\ (5-2) \quad g_{rr} &= ar^{a-2}c_m(ar^a c_m F''(r^a c_m) + (a-1)F'(r^a c_m)), \\ g_{r\theta} &= -amr^{a-1}s_m(r^a c_m F''(r^a c_m) + F'(r^a c_m)), \\ g_{\theta\theta} &= m^2 r^a (r^a s_m^2 F''(r^a c_m) - c_m F'(r^a c_m)), \end{aligned}$$

where

$$(5-3) \quad c_m := \cos m\theta, \quad s_m := \sin m\theta.$$

Since  $F(x)$  is a bounded function,  $g$  is bounded and satisfies (2-13), since  $a < 2$ . Therefore, the inversion  $\Psi_g$  can be expressed as a graph near  $(0, 0, 0)$ . Since  $0 < a < 1$ , the function  $g$  satisfies (a) and (b) of Proposition 2.5. Then  $Z = Z_f(X, Y)$  as in (2-11), where  $f := g$  is a  $C^1$ -function at  $(0, 0)$ . The graph of  $Z_g$  for  $g = f_5$  near  $(0, 0, 0)$  is indicated in the right-hand side of Figure 1. To prove Theorem 1.1, it is sufficient to show that  $(0, 0, 0)$  is a  $C^1$ -umbilic of the graph of  $Z_g(X, Y)$  with index  $1 + (m/2)$ . In the following discussions, we would like to show that there exists a positive number  $R$  such that the graph of  $g$  has no umbilics if  $r > R$ . We then compute the index  $I_g(\Gamma)$  with respect to the circle

$$(5-4) \quad \Gamma(\theta) := (r \cos \theta, r \sin \theta), \quad 0 \leq \theta \leq 2\pi, \quad r > R,$$

using (4-1) and (4-2), which does not depend on the choice of  $r > R$ , as follows. We set

$$(5-5) \quad \check{\delta}_j(\theta) := \delta_j(\Gamma(\theta)), \quad j = 1, 2.$$

The first polar identifier is given by

$$(5-6) \quad \delta_1 = -mr^a s_m (ar^a c_m F''(r^a c_m) + (a-1)F'(r^a c_m)).$$

Since  $0 < a < 1$ , condition (ii) yields that

$$(5-7) \quad (a-1)F'(r^a c_m) < 0.$$

On the other hand, by (i) and (iii), it holds that

$$(5-8) \quad xF''(x) \leq 0, \quad x := r^a c_m.$$

By (5-7) and (5-8), we can conclude that  $\check{\delta}_1(\theta)$  changes sign only at the zeros of the function  $\sin m\theta$ . Since the function  $g$  is symmetric with respect to rotation by angle  $2\pi/m$ , to compute the rotation index of  $\Delta_g$  along  $\Gamma$ , it is sufficient to check the sign changes of  $\check{\delta}_i(\theta)$ ,  $i = 1, 2$ , for  $\theta = 0$  and  $\theta = \pi/m$ . By (5-6), (5-7) and (5-8), we get the following:

$$(5-9) \quad \left. \frac{d\check{\delta}_1}{d\theta} \right|_{\theta=0} > 0, \quad \left. \frac{d\check{\delta}_1}{d\theta} \right|_{\theta=\pi/m} < 0.$$

The second polar identifier  $\delta_2$  is given by

$$r^{2-3a}\delta_2 = -r^{2-a}(a^2 c_m^2 - m^2 s_m^2)F''(c_m r^a) + ac_m(a^2 c_m^2 - am^2 + m^2 s_m^2)F'(c_m r^a)^3 - c_m r^{2-2a}(a^2 - 2a + m^2)F'(c_m r^a).$$

We need the sign of  $\check{\delta}_2(\theta)$  at  $\theta \in (\pi/m)\mathbb{Z}$ . In this case,  $s_m = 0$  and  $c_m = \pm 1$ . Substituting these relations and using the fact that  $F'$  (resp.  $F''$ ) is an even function (resp. an odd function), we have

$$r^{2-3a}\delta_2 = \mp r^{2-a} a^2 F''(r^a) \pm a^2 (a - m^2) F'(r^a)^3 \mp r^{2-2a} (a^2 - 2a + m^2) F'(r^a).$$

Since  $F'$  is bounded, the middle term is bounded. Hence, by (iv) and by the fact that  $0 < a < 1$ , there exists a positive number  $R$  such that the sign of  $\delta_2$  is determined by the sign of the first term  $\mp r^{2-a} a^2 F''(r^a)$  whenever  $r > R$ . Then, we have

$$(5-10) \quad -\check{\delta}_2(\pi/m) = \check{\delta}_2(0) > 0.$$

In particular, the image of the graph of  $g$  has no umbilics when  $r > R$ . By the  $2\pi/m$ -symmetry of  $g$ , (4-2), (5-9), and (5-10), the index  $\text{ind}_\Gamma(\Delta_g)$  is equal to  $-m$ . Then the index of the curvature line flow along  $\Gamma$  is equal to  $I_g(\Gamma) = 1 - m/2$

by (4-1). Then after inversion, the Poincaré–Hopf index formula yields that the index  $I_0$  of the umbilic of  $\Psi_g$  at the origin is

$$I_0 = 2 - I_g(\Gamma) = 1 + m/2.$$

If we choose  $F(x) := \tanh x$ , then the function  $Z_g(X, Y)$  satisfies the properties of Theorem 1.1.

We next prove the corollary. We set

$$(5-11) \quad \lambda := \frac{Z\sqrt{1 + Z_X^2 + Z_Y^2}}{1 + \sqrt{1 + Z_X^2 + Z_Y^2}},$$

where  $Z := Z_g$  is the function given in (2-11). Suppose that  $\lambda$  and  $\lambda\nu$  are a  $C^1$ -function and a  $C^1$ -vector field defined on a sufficiently small neighborhood of  $(0, 0)$  in the  $XY$ -plane, respectively, where  $\nu$  is a unit normal vector field of the graph of  $Z_g$ . Then the map

$$\Phi : (X, Y) \mapsto (\xi(X, Y), \eta(X, Y))$$

given by (A-4) for  $f = Z_{f_m}$  is a local  $C^1$ -diffeomorphism, and is real-analytic on  $U \setminus \{(0, 0)\}$ . Then the proof of Fact A.1 in Appendix A is valid in our situation, and we can conclude that the eigenflow of the Hessian matrix of  $\lambda(\xi, \eta)$  is equal to the curvature line flow of the map  $P(\xi, \eta)$  given by (A-8). Since the image of  $P(\xi, \eta)$  coincides with that of  $\Psi_{f_m}(u, v)$ , we get the proof of the corollary in the introduction.

Thus, it is sufficient to show that  $\lambda$  and  $\lambda\nu$  are  $C^1$  at  $(X, Y) = (0, 0)$ . By (5-11), we have the following expression

$$(5-12) \quad \lambda\nu = \frac{(ZZ_X, ZZ_Y, -Z)}{1 + \sqrt{1 + Z_X^2 + Z_Y^2}}.$$

By (5-11) and (5-12), we can say that  $\lambda(X, Y)$  and  $\lambda(X, Y)\nu(X, Y)$  are  $C^1$  at  $(0, 0)$  if

$$(5-13) \quad \lim_{(X,Y) \rightarrow (0,0)} ZZ_{XX} = \lim_{(X,Y) \rightarrow (0,0)} ZZ_{XY} = \lim_{(X,Y) \rightarrow (0,0)} ZZ_{YY} = 0$$

hold. So to prove the corollary, it is sufficient to show (5-13). It can be easily seen that all of  $r^{1-a}g_r, r^{-a}g_\theta, r^{2-2a}g_{rr}, r^{1-2a}g_{r\theta}$  and  $r^{-2a}g_{\theta\theta}$  are bounded functions on  $\mathbb{R}^2 \setminus \Omega_R$ . Since  $0 < a < \frac{1}{4}$ , Proposition 2.7 yields that the map  $(u, v) \mapsto (X, Y) = \Pi \circ \Psi_g(u, v)$  is a  $C^2$ -map. Then (5-13) is equivalent to

$$(5-14) \quad \lim_{(u,v) \rightarrow (0,0)} ZZ_{uu} = \lim_{(u,v) \rightarrow (0,0)} ZZ_{uv} = \lim_{(u,v) \rightarrow (0,0)} ZZ_{vv} = 0.$$

Since  $Z = h/(k+1)$ , (5-14) follows from (2-19), (2-21) and the fact that

$$\lim_{\rho \rightarrow 0} \rho h_{uu} = \lim_{\rho \rightarrow 0} \rho h_{uv} = \lim_{\rho \rightarrow 0} \rho h_{vv} = 0.$$

## 6. An alternative proof of the main theorem

In the previous section, we have proved Corollary 1.2. However, it is natural to expect that one can give an explicit description of the function with the desired properties. The function  $\lambda$  given in (5-11) does not have a simple expression. On the other hand, we will see that functions

$$(6-1) \quad \Lambda = \Lambda_m := r^2 \tanh(r^{-a} \cos m\theta), \quad m = 1, 2, 3, \dots,$$

satisfy (1) and (2) of Corollary 1.2 if  $0 < a < 1$ . We set

$$(\lambda :=) \lambda_m := r^2 F(r^{-a} \cos m\theta),$$

where  $\xi = r \cos \theta$ ,  $\eta = r \sin \theta$ , and  $F: \mathbb{R} \rightarrow \mathbb{R}$  is a function satisfying the properties (i–iv) given in the beginning of Section 5. Then  $\Lambda_m$  is a special case of  $\lambda_m$  for  $F(x) := \tanh x$ . It holds that

$$\begin{aligned} \lambda_r &= r(2F(r^{-a} c_m) - a c_m r^{-a} F'(r^{-a} c_m)), \\ \lambda_\theta &= -m r^{2-a} s_m F'(r^{-a} c_m), \\ \lambda_{rr} &= 2F(r^{-a} c_m) + a r^{-2a} c_m ((a-3)r^a F'(r^{-a} c_m) + a c_m F''(r^{-a} c_m)), \\ \lambda_{r\theta} &= m s_m r^{1-2a} ((a-2)r^a F'(r^{-a} c_m) + a c_m F''(r^{-a} c_m)), \\ \lambda_{\theta\theta} &= -m^2 r^{2-2a} (r^a c_m F'(r^{-a} c_m) - s_m^2 F''(r^{-a} c_m)), \end{aligned}$$

where  $c_m$  and  $s_m$  are defined in (5-3). We set


$$\zeta_1 := 2(r\lambda_{r\theta} - \lambda_\theta), \quad \zeta_2 := -r^2 \lambda_{rr} + r\lambda_r + \lambda_{\theta\theta}.$$

Then each component of the vector field  $\delta_\lambda := \zeta_1 \partial_x + \zeta_2 \partial_y$  is an identifier for the eigenflow of the Hessian matrix of  $\lambda$  at the origin given in the introduction; see (1-3). By a direct calculation, we have

$$\begin{aligned} \zeta_1 &= 2m r^{2-2a} s_m (a c_m F''(r^{-a} c_m) + (a-1)r^a F'(r^{-a} c_m)), \\ \zeta_2 &= -r^{2-2a} (a^2 c_m^2 - m^2 s_m^2) F''(r^{-a} c_m) - (a^2 - 2a + m^2) r^{2-a} c_m F'(r^{-a} c_m). \end{aligned}$$

By the property (ii) of  $F$ ,  $(a-1)r^a F'(r^{-a} c_m)$  is negative, and by (ii) and (iii),  $c_m F''(r^{-a} c_m)$  is also negative. So  $\zeta_1$  is positively proportional to  $-s_m (= -\sin m\theta)$ . In particular,  $\zeta_1$  vanishes only when  $s_m = 0$ . Moreover, for fixed  $r$ , it holds that  $d\zeta_1/d\theta < 0$  (resp.  $d\zeta_1/d\theta > 0$ ) if  $c_m = 1$  (resp.  $c_m = -1$ ).

On the other hand, if  $s_m = 0$  and  $r$  tends to zero, then  $c_m = \pm 1$  and  $F'(\pm r^{-a})$  and  $F''(\pm r^{-a})$  tend to zero with exponential order (see condition (iv) for  $F(x)$ ). Therefore, the leading term of  $\zeta_2$  for small  $r$  is  $-r^{2-2a} (a^2 c_m^2 - m^2 s_m^2) F''(r^{-a} c_m)$ . Hence, for a fixed sufficiently small  $r$ , the function  $\zeta_2$  is positive (resp. negative) if  $c_m = 1$  (resp.  $c_m = -1$ ). Summarizing these facts, one can easily show that the



**Figure 2.** The image of  $P$  ( $r \leq \frac{1}{2}$ ) for  $m = 2$  and  $a = \frac{1}{2}$ .

index of the vector field  $\delta_\lambda$  at  $o := (0, 0)$  is equal to  $m$ . So the index of the eigenflow of the Hessian matrix of  $\lambda$  at  $o$  is equal to  $1 + m/2$  (see Appendix B). One can easily check that  $\lambda$  is a  $C^1$ -function at  $o$  and the function  $\lambda$  satisfies (1) and (2) of Corollary 1.2. Since  $\Lambda$  is a special case of  $\lambda$ , we proved that  $\Lambda$  satisfies the desired properties.

To give an alternative proof of Theorem 1.1, we consider the real analytic map  $P : \mathbb{R}^2 \setminus \{o\} \rightarrow \mathbb{R}^3$  defined (see (A-8)) by

$$P(\xi, \eta) := (\xi, \eta, \Lambda(\xi, \eta)) - \Lambda(\xi, \eta)v(\xi, \eta),$$

where

$$(6-2) \quad v := \frac{1}{\Lambda_\xi^2 + \Lambda_\eta^2 + 1} (2\Lambda_\xi, 2\Lambda_\eta, \Lambda_\xi^2 + \Lambda_\eta^2 - 1).$$

One can easily verify that

$$\begin{aligned} \Lambda_\xi &= r^{1-a} \left( (ms_1s_m - ac_1c_m) \operatorname{sech}^2(r^{-a}c_m) + 2r^a c_1 \tanh(r^{-a}c_m) \right), \\ \Lambda_\eta &= r^{1-a} \left( 2r^a s_1 \tanh(r^{-a}c_m) - (as_1c_m + mc_1s_m) \operatorname{sech}^2(r^{-a}c_m) \right), \end{aligned}$$

where  $c_1 = \cos \theta$  and  $s_1 = \sin \theta$ . Using them, one can get the expressions

$$(6-3) \quad \Lambda_{\xi\xi} = \frac{1}{r^{2a}} h_1(r, \theta), \quad \Lambda_{\xi\eta} = \frac{1}{r^{2a}} h_2(r, \theta), \quad \Lambda_{\eta\eta} = \frac{1}{r^{2a}} h_3(r, \theta),$$

where  $h_i(r, \theta)$ ,  $i = 1, 2, 3$ , are continuous functions defined on  $\mathbb{R}^2$ . Using (6-2), (6-3) and the fact  $\lim_{r \rightarrow 0} \Lambda/r^{2a} = 0$ , we have

$$(6-4) \quad \lim_{r \rightarrow 0} \Lambda v_\xi = \lim_{r \rightarrow 0} \frac{\Lambda}{r^{2a}} (r^{2a} v_\xi) = 0,$$

and also

$$(6-5) \quad \lim_{r \rightarrow 0} \Lambda v_\eta = 0.$$

Using (6-4), (6-5) and the fact

$$d(\Lambda v) = (d\Lambda)v + \Lambda dv,$$

we can conclude that  $\Lambda v$  can be extended as a  $C^1$ -function at  $o$ . Thus  $P(\xi, \eta)$  can also be extended as a  $C^1$ -differentiable map at  $o$ . One can also easily check that

$$P_\xi(0, 0) = (1, 0, 0), \quad P_\eta(0, 0) = (0, 1, 0).$$

Hence  $P$  is an immersion at  $o$ , and

$$\Phi : (\xi, \eta) \mapsto (X(\xi, \eta), Y(\xi, \eta))$$

is a local  $C^1$ -diffeomorphism, where  $P = (X, Y, Z)$ . In particular,

$$Z_\Lambda := Z(\Phi^{-1}(X, Y))$$

gives a function defined on a neighborhood of  $(X, Y) = (0, 0)$ . By Fact A.1 in Appendix A, the index of the curvature line flow at  $(0, 0)$  of the graph of  $Z_\Lambda$  is equal to the index of the eigenflow of the Hessian matrix of  $\Lambda$ , which implies Theorem 1.1. The image of  $P$  for  $m = 3$  and  $a = \frac{1}{2}$  is given in Figure 2.

## 7. The duality of indices

At the end of this paper, we consider the index at infinity for eigenflows of Hessian matrices. Let

$$f : \mathbb{R}^2 \setminus \Omega_R \rightarrow \mathbb{R}, \quad g : U_{1/R} \setminus \{o\} \rightarrow \mathbb{R}$$

be  $C^2$ -functions, where  $\Omega_R$  and  $U_{1/R}$  are disks defined in Section 2. Let  $\mathcal{H}_f$  (resp.  $\mathcal{H}_g$ ) be the eigenflow of the Hessian matrix of  $f$  (resp.  $g$ ). If the Hessian matrix of  $f$  has no equidiagonal points, then we can consider the index  $\text{ind}(\mathcal{H}_f, \Gamma)$  with respect to the circle  $\Gamma$  given in (5-4) and it is independent of the choice of  $r > R$ . So we denote it by  $\text{ind}_\infty(\mathcal{H}_f)$ . Similarly, if the Hessian matrix of  $g$  has no equidiagonal points, then we can consider the index  $\text{ind}(\mathcal{H}_g, \Gamma')$  with respect to the circle  $\Gamma'(\theta) := (\rho \cos \theta, \rho \sin \theta)$ ,  $0 \leq \theta \leq 2\pi$ ,  $\rho < 1/R$ . Since it is independent of the choice of  $\rho < 1/R$ , we denote it by  $\text{ind}_o(\mathcal{H}_g)$ . Consider the plane-inversion

$$\iota : \mathbb{R}^2 \ni (u, v) \mapsto \frac{1}{u^2 + v^2} (u, v) \in \mathbb{R}^2.$$

Then the following assertion holds:

**Proposition 7.1** (duality of indices). *Let  $f : \mathbb{R}^2 \setminus \Omega_R \rightarrow \mathbb{R}$  be a  $C^2$ -function whose Hessian matrix has no equidiagonal points. Then the function  $g : \Omega_R \rightarrow \mathbb{R}$  defined by*

$$g(x, y) := (u^2 + v^2) f \circ \iota(u, v)$$

(called the dual of  $f$ ) satisfies

$$\text{ind}_o(\mathcal{H}_g) + \text{ind}_\infty(\mathcal{H}_f) = 2.$$

*Proof.* Using the identification of  $(u, v)$  and  $z = u + iv$ , it holds that  $u = (z + \bar{z})/2$  and  $v = (z - \bar{z})/(2i)$ . In particular,  $f$  can be considered as a function of variables  $z$  and  $\bar{z}$ , and can be denoted by  $f = f(z, \bar{z})$ . Since  $\iota(z) = 1/\bar{z}$ , we can write

$$g(z, \bar{z}) := z\bar{z}f(1/\bar{z}, 1/z).$$

Then

$$g_{z\bar{z}}(z, \bar{z}) = \frac{\bar{z}f_{\bar{z}\bar{z}}(1/\bar{z}, 1/z)}{z^3}$$

holds, where

$$\frac{\partial}{\partial z} := \frac{1}{2} \left( \frac{\partial}{\partial u} - i \frac{\partial}{\partial v} \right), \quad \frac{\partial}{\partial \bar{z}} := \frac{1}{2} \left( \frac{\partial}{\partial u} + i \frac{\partial}{\partial v} \right).$$

Since  $\Gamma(\theta) = re^{i\theta}$ , we have that

$$g_{z\bar{z}}(\Gamma(\theta)) = \frac{f_{\bar{z}\bar{z}}(\iota \circ \Gamma(\theta))}{r^2 e^{4i\theta}}.$$

Thus, it holds that

$$\text{ind}_o(g_{z\bar{z}}, \Gamma) = -4 + \text{ind}_o(f_{\bar{z}\bar{z}}, \iota \circ \Gamma).$$

By (B-1), we have

$$\begin{aligned} \text{ind}_o(g_{z\bar{z}}, \Gamma) &= -2 \text{ind}_o(\mathcal{H}_g), \\ \text{ind}_o(f_{\bar{z}\bar{z}}, \iota \circ \Gamma) &= -\text{ind}_o(f_{z\bar{z}}, \iota \circ \Gamma) = 2 \text{ind}_\infty(\mathcal{H}_f). \end{aligned}$$

Thus we get the assertion. □

Applying Proposition 7.1 for the function  $g = \Lambda_m$ , see (6-1), we get the following:

**Corollary 7.2.** *For each  $m \geq 1$ , there exists a  $C^1$ -function  $f : \mathbb{R}^2 \setminus \Omega_R \rightarrow \mathbb{R}$  satisfying*

- (1)  $f$  is real-analytic on  $\mathbb{R}^2 \setminus \Omega_R$ ,
- (2) the eigenflow of the Hessian matrix of  $f$  has no singular points, and
- (3) the index at infinity of the eigenflow of  $H_f$  is equal to  $1 - m/2$ .

The function  $\Lambda_m$  used in the second proof of Theorem 1.1 coincides with the dual of the function  $f_m - 1$  given in (1-4).

### Appendix A: The classical reduction

In this appendix we show the existence of a special coordinate system  $(\xi, \eta)$  of the graph of a function  $f(x, y)$  which reduces the curvature line flow to the Hessian of a certain function, called Ribaucour's parametrization (Umehara learned this from Konrad Voss at the conference of Thessaloniki 1997). Although, the existence of such a coordinate system was classically known, and a proof is in the appendix of [Scherbel 1993], the authors will give the proof here for the sake of convenience. We set  $P = (x, y, f(x, y))$ , and suppose that  $f(0, 0) = f_x(0, 0) = f_y(0, 0) = 0$ . Consider a sphere which is tangent to the graph of  $f$  at  $P$  and also tangent to the  $xy$ -plane at a point  $Q$ . Then, it holds that

$$(A-1) \quad Q + \lambda \mathbf{e}_3 = P + \lambda \nu,$$

where  $\mathbf{e}_3 = (0, 0, 1)$  and  $\nu = (f_x, f_y, -1)/\sqrt{1 + f_x^2 + f_y^2}$ . Taking the third component of (A-1), we get

$$(A-2) \quad \lambda = \frac{f\sqrt{1 + f_x^2 + f_y^2}}{1 + \sqrt{1 + f_x^2 + f_y^2}}.$$

In particular,  $\lambda(0, 0) = 0$ . Since  $f_x(0, 0) = f_y(0, 0) = 0$ , we have that

$$(A-3) \quad d\lambda(0, 0) = df(0, 0) = 0.$$

Taking the exterior derivative of (A-1), and using (A-3) and  $\lambda(0, 0) = 0$ , we have  $dP(0, 0) = dQ(0, 0)$ . So, if we set  $Q = (\xi(x, y), \eta(x, y), 0)$ , then it holds that

$$\begin{aligned} (\xi_x(0, 0)dx + \xi_y(0, 0)dy, \eta_x(0, 0)dx + \eta_y(0, 0)dy, 0) &= dQ \\ &= dP = (dx, dy, f_x(0, 0)dx + f_y(0, 0)dy) = (dx, dy, 0), \end{aligned}$$

which implies that the Jacobi matrix of the map

$$(A-4) \quad \Phi : (x, y) \mapsto (\xi(x, y), \eta(x, y))$$

is the identity matrix at  $(0, 0)$ . So we can take  $(\xi, \eta)$  as a new local coordinate system. Differentiating (A-1) by  $\xi$  and  $\eta$ , we get the following two identities:

$$Q_\xi + \lambda_\xi \mathbf{e}_3 = P_\xi + \lambda_\xi \nu + \lambda \nu_\xi, \quad Q_\eta + \lambda_\eta \mathbf{e}_3 = P_\eta + \lambda_\eta \nu + \lambda \nu_\eta.$$

Taking the inner products of them and  $\nu$ , these two equations yield

$$(A-5) \quad Q_\xi \cdot \nu + \lambda_\xi \nu_3 = \lambda_\xi, \quad Q_\eta \cdot \nu + \lambda_\eta \nu_3 = \lambda_\eta,$$

where we set  $\nu = (\nu_1, \nu_2, \nu_3)$ . Since  $Q = (\xi, \eta, 0)$ , we have that  $Q_\xi = (1, 0, 0)$  and  $Q_\eta = (0, 1, 0)$ . So  $Q_\xi \cdot \nu = \nu_1$  and  $Q_\eta \cdot \nu = \nu_2$ . Substituting this into (A-5), we have

$$(A-6) \quad \lambda_\xi = \frac{\nu_1}{1 - \nu_3}, \quad \lambda_\eta = \frac{\nu_2}{1 - \nu_3}.$$



This implies that  $(\lambda_\xi, \lambda_\eta)$  is the image of  $\nu$  via the stereographic projection, and

$$(A-7) \quad \nu = \frac{1}{1 + \lambda_\xi^2 + \lambda_\eta^2} (2\lambda_\xi, 2\lambda_\eta, \lambda_\xi^2 + \lambda_\eta^2 - 1).$$

By (A-1), we have

$$(A-8) \quad P = (\xi, \eta, 0) - \lambda\nu + (0, 0, \lambda).$$

We prove the following:

**Fact A.1.** *The curvature line flow of the graph  $z = f(x, y)$  coincides with the eigenflow of the Hessian of the function  $\lambda(\xi, \eta)$  given by (A-2).*

*Proof.* Noticing (A-8), we set

$$\Delta_{(\xi, \eta)} := \det \begin{pmatrix} \nu \\ dP \\ d\nu \end{pmatrix} = \det \begin{pmatrix} \nu \\ d\xi, d\eta, d\lambda \\ d\nu \end{pmatrix}.$$

Then this gives a map  $\Delta_{(\xi, \eta)} : T_{(\xi, \eta)}\mathbb{R}^2 \rightarrow \mathbb{R}$  such that

$$\Delta_{(\xi, \eta)} \left( a \frac{\partial}{\partial \xi} + b \frac{\partial}{\partial \eta} \right) = \det(\nu, aP_\xi(\xi, \eta) + bP_\eta(\xi, \eta), av_\xi(\xi, \eta) + bv_\eta(\xi, \eta)) \in \mathbb{R}.$$

It is well known that  $\mathbf{w} \in T_{(\xi, \eta)}\mathbb{R}^2$  points in a principal direction of  $P$  at  $(\xi, \eta)$  if and only if  $\Delta_{(\xi, \eta)}(\mathbf{w}) = 0$ . Since  $(v_1)^2 + (v_2)^2 + (v_3)^2 = 1$ , (A-6) yields that

$$\lambda_\xi v_1 + \lambda_\eta v_2 = \frac{(v_1)^2 + (v_2)^2}{1 - v_3} = \frac{1 - (v_3)^2}{1 - v_3} = 1 + v_3,$$

which implies  $v_3 = \lambda_\xi v_1 + \lambda_\eta v_2 - 1$ . We now set  $\mu = 2/(1 + \lambda_\xi^2 + \lambda_\eta^2)$ . Differentiating (A-7), we have

$$d\nu = \frac{d\mu}{\mu} \nu + \mu(d\lambda_\xi, d\lambda_\eta, \lambda_\xi d\lambda_\xi + \lambda_\eta d\lambda_\eta).$$

The first term of the right-hand side of the above equation is proportional to  $\nu$  and does not affect the computation of  $\Delta_{(\xi, \eta)}$ . So we have that

$$\begin{aligned} \Delta_{(\xi, \eta)} &= \mu \begin{vmatrix} v_1 & v_2 & \lambda_\xi v_1 + \lambda_\eta v_2 - 1 \\ d\xi & d\eta & \lambda_\xi d\xi + \lambda_\eta d\eta \\ d\lambda_\xi & d\lambda_\eta & \lambda_\xi d\lambda_\xi + \lambda_\eta d\lambda_\eta \end{vmatrix} \\ &= \mu \begin{vmatrix} v_1 & v_2 & -1 \\ d\xi & d\eta & 0 \\ d\lambda_\xi & d\lambda_\eta & 0 \end{vmatrix} = -\mu \begin{vmatrix} d\xi & d\eta \\ d\lambda_\xi & d\lambda_\eta \end{vmatrix} \\ &= \mu((\lambda_{\xi\xi} - \lambda_{\eta\eta})d\xi d\eta - \lambda_{\xi\eta}(d\xi^2 - d\eta^2)). \end{aligned}$$

Fact A.1 follows from this representation of  $\Delta_{(\xi, \eta)}$ . □

### Appendix B: Indices of eigenflows of Hessian matrices

Let  $g : \Omega_R \setminus \{o\} \rightarrow \mathbb{R}$  be a  $C^2$ -function, where  $\Omega_R$  is the closed disk of radius  $R$  centered at the origin  $o := (0, 0)$ ; see (2-1). The Hessian matrix of  $g$  is given by

$$H_g := \begin{pmatrix} g_{xx} & g_{xy} \\ g_{yx} & g_{yy} \end{pmatrix}.$$

We denote by  $\mathcal{H}_g$  the eigenflow of  $H_g$ . A point  $p \in \Omega_R \setminus \{o\}$  is called an *equidiagonal point* of  $\mathcal{H}_g$  if  $H_g(p)$  is proportional to the identity matrix. Consider the circle

$$\Gamma(\theta) := r(\cos \theta, \sin \theta), \quad 0 \leq \theta < 2\pi, r < R.$$

If there are no equidiagonal points on  $\Omega_R \setminus \{o\}$ , then we can define the index  $\text{ind}(\mathcal{H}_g, \Gamma)$  of the eigenflow  $\mathcal{H}_g$  with respect to  $\Gamma$ , which does not depend on the choice of  $r$ . We call it the index of  $\mathcal{H}_g$  at the origin and denote it by  $\text{ind}_o(\mathcal{H}_g)$ . Consider the vector field

$$d_g := 2g_{xy} \frac{\partial}{\partial x} + (g_{yy} - g_{xx}) \frac{\partial}{\partial y}.$$

It is well known that the mapping degree of the Gauss map, see (3-1),

$$\check{d}_g : T^1 := \mathbb{R}/2\pi\mathbb{Z} \ni \theta \mapsto \frac{d_g(\Gamma(\theta))}{|d_g(\Gamma(\theta))|} \in S^1 := \{(x, y) \in \mathbb{R}^2; x^2 + y^2 = 1\}$$

is equal to  $2 \text{ind}_o(\mathcal{H}_g)$ . Using the correspondence  $(x, y) \mapsto x + iy$ , we identify  $\mathbb{R}^2$  with  $\mathbb{C}$ , where  $i = \sqrt{-1}$ . Then

$$\begin{aligned} g_z &= \frac{1}{2}(g_x - ig_y), \\ g_{zz} &= \frac{1}{4}((g_{xx} - g_{yy}) - 2ig_{xy}), \end{aligned}$$

where  $g_z := \partial g / \partial z$ ,  $g_{zz} := \partial^2 g / \partial z^2$  and

$$\frac{\partial}{\partial z} := \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right).$$

Thus,  $d_g$  can be identified with the right-angle rotation of  $\overline{g_{zz}}$ . In particular,

$$(B-1) \quad \text{ind}_o(\mathcal{H}_g) = -\frac{1}{2} \text{ind}_o(g_{zz}).$$

Here  $g_{zz}$  is considered as a vector field and  $\text{ind}_o(g_{zz})$  is its index at the origin. Let  $(r, \theta)$  be as in (2-5). Then  $z = re^{i\theta}$  and

$$\begin{aligned} g_z &= \frac{e^{-i\theta}}{2r} (rg_r - ig_\theta), \\ g_{zz} &= \frac{e^{-2i\theta}}{4r^2} ((r^2 g_{rr} - rg_r - g_{\theta\theta}) + 2i(g_\theta - rg_{r\theta})). \end{aligned}$$

We consider the vector field defined by

$$(B-2) \quad \delta_g := 2(r g_{r\theta} - g_\theta) \frac{\partial}{\partial x} + (-r^2 g_{rr} + r g_r + g_{\theta\theta}) \frac{\partial}{\partial y}.$$

Since, from [Klotz 1959, (18)],

$$\text{ind}_o(\overline{g_{zz}}) = 2 + \text{ind}_o(\delta_g),$$

we obtain the following:

**Lemma B.1.** *The identity  $\text{ind}_o(\mathcal{H}_g) = 1 + \text{ind}_o(\delta_g)/2$  holds.*

### Acknowledgements

The authors thank Wayne Rossman for valuable comments. Umehara also thanks Udo Hertrich-Jeromin for fruitful conversations.

### References

- [Ando 2003] N. Ando, “An umbilical point on a non-real-analytic surface”, *Hiroshima Math. J.* **33**:1 (2003), 1–14. MR Zbl
- [Bates 2001] L. Bates, “A weak counterexample to the Carathéodory conjecture”, *Differential Geom. Appl.* **15**:1 (2001), 79–80. MR Zbl
- [Ghomi and Howard 2012] M. Ghomi and R. Howard, “Normal curvatures of asymptotically constant graphs and Carathéodory’s conjecture”, *Proc. Amer. Math. Soc.* **140**:12 (2012), 4323–4335. MR Zbl
- [Guilfoyle and Klingenberg 2008] B. Guilfoyle and W. Klingenberg, “Proof of the Carathéodory conjecture”, preprint, 2008. arXiv
- [Guilfoyle and Klingenberg 2012] B. Guilfoyle and W. Klingenberg, “From global to local: an index bound for umbilic points on smooth convex surfaces”, preprint, 2012. arXiv
- [Gutierrez et al. 1996] C. Gutierrez, F. Mercuri, and F. Sánchez-Bringas, “On a conjecture of Carathéodory: analyticity versus smoothness”, *Experiment. Math.* **5**:1 (1996), 33–37. MR Zbl
- [Hamburger 1940] H. Hamburger, “Beweis einer Caratheodoryschen Vermutung, I”, *Ann. of Math.* (2) **41** (1940), 63–86. MR Zbl
- [Hamburger 1941a] H. L. Hamburger, “Beweis einer Caratheodoryschen Vermutung, II”, *Acta Math.* **73** (1941), 175–228. MR Zbl
- [Hamburger 1941b] H. L. Hamburger, “Beweis einer Caratheodoryschen Vermutung, III”, *Acta Math.* **73** (1941), 229–332. MR Zbl
- [Klotz 1959] T. Klotz, “On G. Bol’s proof of Carathéodory’s conjecture”, *Comm. Pure Appl. Math.* **12** (1959), 277–311. MR Zbl
- [Scherbel 1993] H. Scherbel, *A new proof of Hamburger’s index theorem on umbilical points*, Ph.D. thesis, Eidgenössische Technische Hochschule Zürich, 1993, <http://search.proquest.com/docview/304065217>. MR
- [Smyth and Xavier 1992] B. Smyth and F. Xavier, “A sharp geometric estimate for the index of an umbilic on a smooth surface”, *Bull. London Math. Soc.* **24**:2 (1992), 176–180. MR Zbl

Received August 7, 2015. Revised September 25, 2016.

NAOYA ANDO  
GRADUATE SCHOOL OF SCIENCE AND TECHNOLOGY  
KUMAMOTO UNIVERSITY  
2-39-1 KUROKAMI  
KUMAMOTO 860-8555  
JAPAN  
ando@sci.kumamoto-u.ac.jp

TOSHIFUMI FUJIYAMA  
DEPARTMENT OF MATHEMATICAL INFORMATICS  
GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY  
UNIVERSITY OF TOKYO  
TOKYO 113-8656  
JAPAN  
toshifumi\_fujiyama@mist.i.u-tokyo.ac.jp

MASAACKI UMEHARA  
DEPARTMENT OF MATHEMATICAL AND COMPUTING SCIENCES  
TOKYO INSTITUTE OF TECHNOLOGY  
2-12-1-W8-34, O-OKAYAMA  
MEGURO-KU  
TOKYO 152-8552  
JAPAN  
umehara@is.titech.ac.jp

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Igor Pak  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pak.pjm@gmail.com](mailto:pak.pjm@gmail.com)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 288 No. 1 May 2017

---

$C^1$ -umbilics with arbitrarily high indices	1
NAOYA ANDO, TOSHIFUMI FUJIYAMA and MASAOKI UMEHARA	
Well-posedness of second-order degenerate differential equations with finite delay in vector-valued function spaces	27
SHANGQUAN BU and GANG CAI	
On cusp solutions to a prescribed mean curvature equation	47
ALEXANDRA K. ECHART and KIRK E. LANCASTER	
Radial limits of capillary surfaces at corners	55
MOZHGAN (NORA) ENTEKHABI and KIRK E. LANCASTER	
A new bicommutant theorem	69
ILIJAS FARAH	
Noncompact manifolds that are inward tame	87
CRAIG R. GUILBAULT and FREDERICK C. TINSLEY	
$p$ -adic variation of unit root $L$ -functions	129
C. DOUGLAS HAESSIG and STEVEN SPERBER	
Bavard's duality theorem on conjugation-invariant norms	157
MORIMICHI KAWASAKI	
Parabolic minimal surfaces in $\mathbb{M}^2 \times \mathbb{R}$	171
VANDERSON LIMA	
Regularity conditions for suitable weak solutions of the Navier–Stokes system from its rotation form	189
CHANGXING MIAO and YANQING WANG	
Geometric properties of level curves of harmonic functions and minimal graphs in 2-dimensional space forms	217
JINJU XU and WEI ZHANG	
Eigenvalue resolution of self-adjoint matrices	241
XUWEN ZHU	



0030-8730(201705)288:1;1-T