

*Pacific
Journal of
Mathematics*

BRAID GROUPS AND QUIVER MUTATION

JOSEPH GRANT AND BETHANY ROSE MARSH

BRAID GROUPS AND QUIVER MUTATION

JOSEPH GRANT AND BETHANY ROSE MARSH

We describe presentations of braid groups of type ADE and show how these presentations are compatible with mutation of quivers. In types A and D these presentations can be understood geometrically using triangulated surfaces. We then give a categorical interpretation of the presentations, with the new generators acting as spherical twists at simple modules on derived categories of Ginzburg dg-algebras of quivers with potential.

1. Introduction	77
2. Presentations of braid groups	80
3. Topological interpretation of the generators	87
4. Actions on categories	95
Acknowledgements	114
References	114

1. Introduction

Braid groups are fundamental objects in mathematics. Although they are of a topological and geometric nature, they have an algebraic interpretation: a simple presentation by generators and relations which is just based on adjacency of integers [Artin 1925]. This can be encoded in a line graph, and from there one can generalize to define a group from any finite graph, known as the Artin braid group.

The most well known groups defined from graphs are the Coxeter groups (we restrict to the simply laced cases, for simplicity). These are closely related to Artin braid groups: each Coxeter group is a quotient of a corresponding Artin braid group in a natural way. In particular, the symmetric group on n letters is a quotient of the classical braid group on n strands. Coxeter groups naturally split into two distinct classes: those of finite type, corresponding to the Dynkin diagrams of type ADE , and those of infinite type. Although all Artin braid groups are infinite, the Artin braid groups of Dynkin type have a different character to those not of Dynkin type, and are known as Artin groups of “finite type”.

This work was supported by the Engineering and Physical Sciences Research Council (grant number EP/G007497/1) and the Mathematical Sciences Research Institute, Berkeley.

MSC2010: primary 13F60, 16G20, 20F36; secondary 16E35, 16E45, 18E30.

Keywords: mutation, braid groups, cluster algebras, Ginzburg dg algebra, spherical twist.

This dichotomy also arises in another area of mathematics which has generated a lot of interest in the recent years: cluster algebras. In this theory, there is a notion of finite-type cluster algebras, which again correspond to the Dynkin diagrams [Fomin and Zelevinsky 2003]. Cluster algebras are specified by a directed graph, known as a quiver, together with other information. A key ingredient in the definition is the notion of mutation, which changes the arrows in a quiver in a nonobvious manner which generalizes reflection at a source or sink. Barot and Marsh [2015] have given new presentations of Coxeter groups of finite type based on quivers obtained from Dynkin diagrams under finite sequences of mutations. Our first result generalizes this to braid groups:

Theorem A. (Theorem 2.12) Let Q be a quiver, with vertices $1, 2, \dots, n$, obtained from a Dynkin quiver by a finite sequence of mutations. Let B_Q be the group with generators s_1, s_2, \dots, s_n , subject to the relations

- (a) $s_i s_j = s_j s_i$ if there is no arrow between i and j (in either direction);
- (b) $s_i s_j s_i = s_j s_i s_j$ if there is an arrow between i and j (in either direction);
- (c) $s_{i_1} s_{i_2} \cdots s_{i_n} s_{i_1} \cdots s_{i_{n-2}} = s_{i_2} s_{i_3} \cdots s_{i_n} s_{i_1} \cdots s_{i_{n-1}} = \cdots = s_{i_n} s_{i_1} s_{i_2} \cdots s_{i_n} s_{i_1} s_{i_2} \cdots s_{i_{n-3}}$, whenever $i_1 \rightarrow i_2 \rightarrow \cdots \rightarrow i_n \rightarrow i_1$ is a chordless cycle in Q .

Then B_Q is isomorphic to the Artin braid group of the same Dynkin type as Q .

We prove our result via isomorphisms between abstractly defined groups, which can be thought of as mutations of groups, even though the resulting groups are isomorphic. The Artin group presentations we obtain induce presentations of the corresponding Coxeter groups which are distinct from those in [Barot and Marsh 2015]; we also give a compatibility result which shows the relationship between the two presentations. Why have we chosen to use presentations which don't agree with the earlier work? This is explained in the following two sections of the paper, as we now detail.

Certain cluster algebras can be understood using pictures. A (tagged) triangulation of a Riemann surface with marked points on its boundary defines a quiver [Fomin and Zelevinsky 2002; Fomin et al. 2008]; see also [Caldero et al. 2006]. Then mutation of the quiver has a natural interpretation in terms of swapping one diagonal of a given quadrilateral for the other. So these cluster algebras have a topological interpretation. In particular, such descriptions are available for the infinite families of Dynkin type. It is natural to ask whether we can understand the generators above, and the isomorphisms corresponding to mutations, in terms of the geometry of the surface. The answer is yes, as shown by the following theorem:

Theorem B. (Theorem 3.6) Let Δ be a Dynkin diagram of type A_n or type D_n . In the former case, let (X, M) be a disk with $n + 3$ marked points on its boundary. In the latter case, let (X, M) be a disk with n marked points on its boundary and

one marked point in its interior, taken to be a cone point of order two (so X is an orbifold in this case).

Let \mathcal{T} be a tagged triangulation of (X, M) . Let $G_{\mathcal{T}}$ be the graph in (X, M) dual to \mathcal{T} . For each vertex i of the quiver $Q_{\mathcal{T}}$ associated to \mathcal{T} as in [Fomin and Zelevinsky 2002; Fomin et al. 2008], let σ_i be the braid of (X, M) associated to the edge of $G_{\mathcal{T}}$ crossing the tagged arc in \mathcal{T} corresponding to i (see Definition 3.3). Then there is an isomorphism between the subgroup $H_{\mathcal{T}}$ of the braid group generated by the σ_i and the group B_Q defined above, taking σ_i to s_i .

Furthermore, in type A_n , the subgroup $H_{\mathcal{T}}$ coincides with the braid group of (X, M) , while in type D_n , it is of index two in the braid group of (X, M) .

As well as the original combinatorial and commutative algebraic approach to cluster algebras, and the geometric approach described above, there is a third approach which has proved very powerful: the representation theoretic approach [Buan et al. 2006; Caldero et al. 2006]. This approach uses finite-dimensional (noncommutative) algebras and ideas from categorification to better understand cluster algebras, and has received intense study. Braid groups also appear in representation theory and categorification [Rouquier and Zimmermann 2003; Seidel and Thomas 2001]: in many important situations there are actions of braid groups on derived categories via spherical twists. One example of this is given by certain derived categories of differential graded algebras [Ginzburg 2007; Keller and Yang 2011] which are known to cover the categories appearing in the representation theoretic approach to cluster algebras [Amiot 2009]. One might hope that these categorical braid group actions are related to our presentations of braid groups, and we show that this is indeed the case.

First, we make a connection between the categorical and the geometric situations. The relevant differential graded algebras are defined by use of a quiver together with a formal sum of cycles in that quiver known as a potential [Ginzburg 2007]. Mutation of quivers of potential has been defined [Derksen et al. 2008] and, in the situations where our cluster algebra comes from a Riemann surface, the mutation of potentials also has a geometric interpretation [Labardini-Fragoso 2009]. Relying heavily on results of Labardini-Fragoso [2009; 2016], we observe that the potential defined on mutation-Dynkin quivers according to the geometric procedure is equivalent to the “obvious” potential that one might guess (Proposition 4.4). So, while the potential is important, it is in fact entirely determined by the quiver in types A and D . Note that this result could also be proved relatively easily via a direct calculation.

Next we show that we do indeed obtain an action of the groups B_Q (defined using mutation-Dynkin quivers) on derived categories of Ginzburg differential graded algebras in which the generators act via spherical twists. After setting up all the technical machinery correctly, the main difficulty in proving this is to check that the mutation procedure for the groups B_Q , which relates the group associated to a

quiver to the group associated to a mutated quiver, actually lifts to the categorical setting as a natural isomorphism of functors. We do this, using important results of Keller and Yang [2011]. From here, we can use the earlier theory developed here to show that the generators s_i of finite type Artin braid groups from Theorem A can be viewed as derived autoequivalences:

Theorem C. (Theorem 4.16) Let (Q, W) be a mutation-Dynkin quiver with potential of type ADE , and let $\Gamma_{Q,W}$ be the corresponding Ginzburg differential graded algebra. Let $\text{D}_{\text{fd}}(\Gamma_{Q,W})$ denote the full subcategory of the derived category $\text{D}(\Gamma_{Q,W})$ on objects with finite-dimensional total homology. Then there is a group homomorphism

$$B_Q \rightarrow \text{Aut D}_{\text{fd}}(\Gamma_{Q,W}), \quad s_i \mapsto F_i$$


sending the group generator associated to the vertex i of Q to the spherical twist F_i at the simple $\Gamma_{Q,W}$ -module S_i .

Since we started work on this project, we have become aware of independent work by other authors. A. King and Y. Qiu have a related project, and were aware of the new relations between spherical twists and a topological interpretation of the spherical twist group; see [Qiu 2016], especially Section 10.1. In particular, an independent proof of a version of Theorem 2.10 in types A and D was announced in [Qiu 2016]. A key difference in our approach is the use of an orbifold with cone point of degree two in type D . In [Nagao 2010, §2.2], K. Nagao refers to an action of the mapping class group of a marked surface on the derived category of a Ginzburg dg-algebra associated to a triangulation.

Since we released the first draft of this article, the preprint [Haley et al. 2014] has appeared, where the authors give a presentation (different from the one given here) of the Artin braid group for each diagram of finite type (in the cluster-theoretic sense). This includes the non-simply-laced cases (not considered here) but does not include a topological or categorical interpretation.

2. Presentations of braid groups


Braid groups. Let Δ be a graph of ADE Dynkin type, i.e., Δ is a graph of type A_n for $n \geq 1$, of type D_n for $n \geq 4$, or of type E_6, E_7 or E_8 .



In particular, Δ has no double edges or cycles. Let I be the set of vertices of Δ . We can associate a group B_Δ to Δ , which we call *the braid group of Δ* . It has a distinguished set of generators $S_\Delta = \{s_i\}_{i \in I}$, and the relations depend on whether or not two vertices are connected by an edge. They are

- (i) $s_i s_j = s_j s_i$ if i and j are not connected by an edge;
- (ii) $s_i s_j s_i = s_j s_i s_j$ if i and j are connected by an edge.

If Δ is of type A_n then we recover the ‘‘usual’’ braid group, sometimes denoted B_{n+1} . Its generators can be visualized as



and the relations of type (i) record the fact that crossings of adjacent pairs of strings which are far apart commute, while relations of type (ii) record a Reidemeister 3 move.

If we also impose the relation that $s_i^2 = 1$ for all $i \in I$ then we recover the Coxeter group of type Δ . More information on Coxeter groups and braid groups can be found in [Humphreys 1990; Kassel and Turaev 2008].

Mutation of quivers. A quiver is just a directed graph. Throughout this article we will only work with quivers with finitely many vertices and finitely many arrows that have no loops or oriented 2-cycles. For a given quiver Q , we again denote its set of vertices by I .

There is a procedure to obtain one quiver from another, called *quiver mutation*, due to Fomin and Zelevinsky [2002, §4]. Fix Q and let $k \in I$. Then we obtain the mutated quiver $\mu_k(Q)$ as follows:

- (i) for each pair of arrows $i \rightarrow k \rightarrow j$ through k , add a formal composite $i \rightarrow j$;
- (ii) reverse the orientation of all arrows incident with k ;
- (iii) remove a maximal set of 2-cycles (we may have created 2-cycles in the previous two steps).

It is a basic but important observation that quiver mutation does not change the set of vertices. One can also check that mutation is an involution.

We call a cycle in an unoriented graph (or in the underlying unoriented graph of a quiver) *chordless* if the full subgraph on the vertices of the cycle contains no edges which are not part of the cycle. We call a quiver *Dynkin* if its underlying unoriented graph is a Dynkin graph of type ADE , and *mutation-Dynkin* if it can be

obtained by mutating a Dynkin quiver finitely many times. By a theorem of Fomin and Zelevinsky [2003, Theorem 1.4], there are only finitely many quivers that can be obtained by mutating a given Dynkin quiver.

The following fact, due to Fomin and Zelevinsky, will be useful to us:

Proposition 2.1. *In any mutation-Dynkin quiver, there are no double arrows and all chordless cycles are oriented.*

Proof. By [Fomin and Zelevinsky 2003, Theorem 1.8], the entries in the corresponding exchange matrix B satisfy $|B_{xy}B_{yx}| \leq 3$ for all x, y (known as being 2-finite). Hence there cannot be any double arrows in the quiver.


Now let Q be a mutation-Dynkin quiver and C a chordless cycle in Q . Then, since Q is 2-finite, so is C . By Proposition 9.7 of the same paper, C must be an oriented cycle. □

Groups from quivers. Let Q be a mutation-Dynkin quiver.

Definition 2.2. Let B_Q be the group with generators $S_Q = \{s_i\}_{i \in I}$ subject to the following relations:

- (i) $s_i s_j = s_j s_i$ whenever i and j are vertices with no arrow between them,
- (ii) $s_i s_j s_i = s_j s_i s_j$ whenever i and j are vertices of Q and there is an arrow between them (in either direction);
- (iii) $s_1 s_2 \cdots s_n s_1 s_2 \cdots s_{n-2} = s_2 s_3 \cdots s_n s_1 \cdots s_{n-1}$
 \vdots
 $= s_n s_1 s_2 \cdots s_n s_1 s_2 \cdots s_{n-3}$

whenever Q contains an oriented chordless n -cycle



Remark 2.3. If Q is a Dynkin quiver, then B_Q is (isomorphic to) the Artin braid group of the corresponding Dynkin type.

This presentation is symmetric but not minimal:

Lemma 2.4. *For each single chordless n -cycle, in the presence of the relations of type (i) and (ii), any one of the relations of type (iii) implies all the others.*

Proof. It is enough to show that if the relation

$$(1) \quad s_1 s_2 \cdots s_n s_1 s_2 \cdots s_{n-2} = s_2 s_3 \cdots s_n s_1 s_2 \cdots s_{n-1}$$

holds then

$$s_1 s_2 \cdots s_n s_1 s_2 \cdots s_{n-2} = s_3 s_4 \cdots s_n s_1 s_2 \cdots s_n.$$

So, we assume that (1) holds. Then we have

$$s_2^{-1}s_1s_2 \cdots s_n s_1 s_2 \cdots s_{n-2} s_n = s_3 \cdots s_n s_1 s_2 \cdots s_{n-1} s_n.$$

The left-hand side can be rewritten, using relations of type (i) and (ii), as

$$\begin{aligned} s_2^{-1}s_1s_2 \cdots s_n s_1 s_2 \cdots s_{n-2} s_n &= s_1 s_2 s_1^{-1} s_3 \cdots s_n s_1 s_2 \cdots s_{n-2} s_n \\ &= s_1 s_2 s_3 \cdots s_{n-1} s_1^{-1} s_n s_1 s_2 \cdots s_{n-2} s_n \\ &= s_1 s_2 \cdots s_{n-1} s_n s_1 s_n^{-1} s_2 \cdots s_{n-2} s_n = s_1 s_2 \cdots s_n s_1 s_2 \cdots s_{n-2}, \end{aligned}$$

and the result follows. \square

Though the relations look different, by taking an appropriate quotient we can obtain the groups defined by Barot and Marsh [2015] directly:

Lemma 2.5. *If, along with the relations of types (i) and (ii), we also impose the relations $s_i^2 = 1$ for all $i \in I$, then the group B_Q becomes isomorphic to the group $\Gamma_{U(Q)}$ defined in [Barot and Marsh 2015, Section 3], where $U(Q)$ is the underlying graph of Q .*

Proof. As our definition is the usual definition of the braid group for a Dynkin quiver, this follows from results in [Barot and Marsh 2015] and the results below on how our groups change with quiver mutation, but since it is straightforward to give a direct proof, we do so.

We need to show that, in the presence of relations (i), (ii), and $s_i^2 = 1$ for all $i \in I$, our extra relation (iii) holds if and only if the relation

$$(s_1 s_2 \cdots s_{n-1} s_n s_{n-1} \cdots s_2)^2 = 1$$

and its rotations hold for each n -cycle $1 \rightarrow 2 \rightarrow \cdots \rightarrow n \rightarrow 1$. By symmetry, it is enough to check that the relation above is equivalent to $s_1 s_2 \cdots s_n s_1 s_2 \cdots s_{n-2} = s_2 s_3 \cdots s_n s_1 \cdots s_{n-1}$.

Using that $s_i = s_i^{-1}$, we see our relation is equivalent to

$$s_1 s_2 \cdots s_n s_1 s_2 \cdots s_{n-2} s_{n-1} s_{n-2} \cdots s_1 s_n \cdots s_3 s_2 = 1.$$

Multiplying out the relation from Barot and Marsh, we see that it is equivalent to

$$s_1 s_2 \cdots s_{n-1} s_n s_{n-1} \cdots s_2 s_1 s_2 \cdots s_{n-1} s_n s_{n-1} \cdots s_2 = 1.$$

Cancelling out n terms on the left and $n - 1$ terms on the right of these two expressions, it just remains to show

$$s_1 s_2 \cdots s_{n-2} s_{n-1} s_{n-2} \cdots s_2 s_1 = s_{n-1} s_{n-2} \cdots s_2 s_1 s_2 \cdots s_{n-2} s_{n-1}.$$

As there is an arrow $i \rightarrow i + 1$ for each i and the cycle is chordless, the symmetric group on n letters maps onto the subgroup generated by s_1, \dots, s_{n-1} with the



Figure 1. Mutation of a quiver of mutation Dynkin type.

transposition which swaps i and $i + 1$ being sent to s_i . It is easy to see that the corresponding relation holds in the symmetric group, with both sides of the equation representing the transposition which swaps 1 and n . \square

We will justify our choice of relations in [Remark 4.19](#).


Mutation of groups. Let B_Q be the group associated to the mutation-Dynkin quiver Q , as above, and let k be a vertex of Q . Denote $\mu_k(Q)$ by Q' . Our aim in this section is to show that B_Q is isomorphic to $B_{Q'}$. We will do this by using a group homomorphism $\varphi_k : B_Q \rightarrow B_{Q'}$ defined using a formula which lifts the formula used in [\[Barot and Marsh 2015, §5\]](#).


The following lemma follows from results in [\[Fomin and Zelevinsky 2003\]](#) (see [\[Barot and Marsh 2015, §2\]](#)).

Lemma 2.6. *Let Q be a quiver of mutation-Dynkin type, and fix a vertex k of Q . Suppose that k has two neighbouring vertices. Then the possibilities for the induced subquiver of Q containing vertex k and its neighbours are shown in [Figure 1](#). The effect of mutation is shown in each case.*

The following lemma follows from [\[Barot and Marsh 2015, Lemma 2.5\]](#).

Lemma 2.7. *Let Q be a quiver of mutation-Dynkin type, and fix a vertex k of Q . Let C be an oriented cycle in Q . Then C is one of the following. In each case we indicate what happens locally under mutation at k .*





- (d) An oriented cycle containing exactly one neighbour of k . Mutation at k reverses the arrow between k and its neighbour in C .
- (e) An oriented cycle containing no neighbours of k . Mutation at k does not affect C .

Recall that B_Q is defined using generators s_i for $i \in I$. We denote the corresponding generating set for $B_{Q'}$ by t_i , $i \in I$. Let F_Q be the free group on the generators s_i for $i \in I$.

Definition 2.8. Let $\varphi_k : F_Q \rightarrow B_{Q'}$ be the group homomorphism defined by

$$\varphi_k(s_i) = \begin{cases} t_k t_i t_k^{-1} & \text{if } i \rightarrow k \text{ in } Q; \\ t_i & \text{otherwise.} \end{cases}$$

Proposition 2.9. The group homomorphism φ_k induces a group homomorphism (which we also denote by φ_k) from B_Q to $B_{Q'}$.

Proof. Let us write $\tilde{s}_i = \varphi_k(s_i)$. We must show that the elements \tilde{s}_i in $B_{Q'}$ satisfy the defining relations of B_Q . Note that the t_i satisfy the defining relations for $B_{Q'}$.

Firstly, we check the relations (ii) for an arrow incident with k . Suppose that there is an arrow $i \rightarrow k$. Using the fact that $t_i t_k t_i = t_k t_i t_k$,

$$\begin{aligned} \tilde{s}_i \tilde{s}_k \tilde{s}_i &= t_k t_i t_k t_i t_k^{-1} = t_k^2 t_i t_k t_k^{-1} \\ &= t_k^2 t_i. \end{aligned}$$

Also,

$$\tilde{s}_k \tilde{s}_i \tilde{s}_k = t_k^2 t_k t_i t_k^{-1} t_k = t_k^2 t_i.$$

So

$$\tilde{s}_i \tilde{s}_k \tilde{s}_i = \tilde{s}_k \tilde{s}_i \tilde{s}_k,$$

as required.

If there is an arrow $i \leftarrow k$, then

$$\tilde{s}_i \tilde{s}_k \tilde{s}_i = t_i t_k t_i = t_k t_i t_k = \tilde{s}_k \tilde{s}_i \tilde{s}_k.$$

Next, we consider relations (i) and (ii) for all other arrows in Q . Relations of this kind involving pairs of vertices which are not neighbours of k follow immediately

from the corresponding relations in B_Q . If only one of the vertices in the relation is a neighbour of k , the relation again follows immediately since t_k commutes with any generator corresponding to a vertex not incident with k in Q' (or equivalently, in Q). So we only need to consider the case where both of the vertices in the pair are incident with k and we can use [Lemma 2.6](#).

Going in either direction in part (a) of [Lemma 2.6](#), the relation $\tilde{s}_i \tilde{s}_j = \tilde{s}_j \tilde{s}_i$ follows from the relation $t_i t_j = t_j t_i$ in $B_{Q'}$, so we consider part (b), firstly from left to right. The cycle in Q' gives the relation $t_k t_i = t_j t_k t_i t_j t_k^{-1} t_j^{-1}$. Also applying the relation $t_k^{-1} t_j^{-1} t_k^{-1} = t_j^{-1} t_k^{-1} t_j^{-1}$, we obtain

$$\begin{aligned} \tilde{s}_i \tilde{s}_j &= t_k t_i t_k^{-1} t_j = t_j t_k t_i t_j t_k^{-1} t_j^{-1} t_k^{-1} t_j \\ &= t_j t_k t_i t_k^{-1} = \tilde{s}_j \tilde{s}_i. \end{aligned}$$

Going from right to left in part (b), we have, using $t_j t_k t_j = t_k t_j t_k$, $t_i t_j = t_j t_i$ and $t_i t_k t_i = t_k t_i t_k$,

$$\begin{aligned} \tilde{s}_j \tilde{s}_i \tilde{s}_j &= t_k t_j t_k^{-1} t_i t_k t_j t_k^{-1} = t_j^{-1} t_k t_j t_i t_j^{-1} t_k t_j \\ &= t_j^{-1} t_k t_i t_k t_j = t_j^{-1} t_i t_k t_i t_j \\ &= t_i t_j^{-1} t_k t_j t_i = t_i t_k t_j t_k^{-1} t_i \\ &= \tilde{s}_i \tilde{s}_j \tilde{s}_i. \end{aligned}$$

Next, we have to check that the \tilde{s}_i satisfy the relations of type (iii) for Q , so we need to consider each type of cycle described in [Lemma 2.7](#). By [Lemma 2.4](#), it is enough to check that, for any given cycle in Q , one of the relations in (iii) holds.

For part (a),

$$\tilde{s}_k \tilde{s}_i \tilde{s}_j \tilde{s}_k = t_k t_i t_k t_j t_k^{-1} t_k = t_k t_i t_k t_j,$$

while

$$\begin{aligned} \tilde{s}_i \tilde{s}_j \tilde{s}_k \tilde{s}_i &= t_i t_k t_j t_k^{-1} t_k t_i = t_i t_k t_j t_i \\ &= t_i t_k t_i t_j, \end{aligned}$$

which is equal to $\tilde{s}_k \tilde{s}_i \tilde{s}_j \tilde{s}_k$ as required.

For part (b), applying a relation for the cycle in Q' in the fourth step,

$$\begin{aligned} \tilde{s}_{i_1} \tilde{s}_{i_2} \cdots \tilde{s}_{i_r} \tilde{s}_{i_1} \tilde{s}_{i_2} \cdots \tilde{s}_{i_{r-2}} &= t_k t_{i_1} t_k^{-1} t_{i_2} \cdots t_{i_r} t_k t_{i_1} t_k^{-1} t_{i_2} \cdots t_{i_{r-2}} \\ &= t_{i_1}^{-1} t_k t_{i_1} t_{i_2} \cdots t_{i_r} t_k t_{i_1} t_k^{-1} t_{i_2} \cdots t_{i_{r-2}} \\ &= t_{i_1}^{-1} t_k t_{i_1} t_{i_2} \cdots t_{i_r} t_k t_{i_1} t_{i_2} \cdots t_{i_{r-2}} t_k^{-1} \\ &= t_{i_1}^{-1} t_{i_1} t_{i_2} \cdots t_{i_r} t_k t_{i_1} t_{i_2} \cdots t_{i_{r-2}} t_{i_{r-1}} t_k^{-1} \\ &= t_{i_2} \cdots t_{i_r} t_k t_{i_1} t_k^{-1} t_{i_2} \cdots t_{i_{r-2}} t_{i_{r-1}} \\ &= \tilde{s}_{i_2} \cdots \tilde{s}_{i_r} \tilde{s}_{i_1} \tilde{s}_{i_2} \cdots \tilde{s}_{i_{r-2}} \tilde{s}_{i_{r-1}}. \end{aligned}$$

For part (c), applying a relation for the cycle in Q' in the fourth step,

$$\begin{aligned}
 \tilde{s}_{i_1} \tilde{s}_{i_2} \cdots \tilde{s}_{i_{r-1}} \tilde{s}_{i_r} \tilde{s}_k \tilde{s}_{i_1} \cdots \tilde{s}_{i_{r-1}} &= t_{i_1} t_{i_2} \cdots t_{i_{r-1}} t_k t_r t_k^{-1} t_k t_{i_1} \cdots t_{i_{r-1}} \\
 &= t_{i_1} t_{i_2} \cdots t_{i_{r-1}} t_k t_r t_{i_1} \cdots t_{i_{r-1}} \\
 &= t_{i_1} t_k t_{i_2} \cdots t_{i_{r-1}} t_r t_{i_1} \cdots t_{i_{r-1}} \\
 &= t_{i_1} t_k t_{i_1} t_{i_2} \cdots t_{i_{r-1}} t_r t_{i_1} \cdots t_{i_{r-2}} \\
 &= t_k t_{i_1} t_k t_{i_2} \cdots t_{i_{r-1}} t_r t_{i_1} \cdots t_{i_{r-2}} \\
 &= t_k t_{i_1} t_{i_2} \cdots t_{i_{r-1}} t_k t_r t_k^{-1} t_k t_{i_1} \cdots t_{i_{r-2}} \\
 &= \tilde{s}_k \tilde{s}_{i_1} \tilde{s}_{i_2} \cdots \tilde{s}_{i_{r-1}} \tilde{s}_{i_r} \tilde{s}_k \tilde{s}_{i_1} \cdots \tilde{s}_{i_{r-2}},
 \end{aligned}$$

and we are done. \square

Theorem 2.10. *The map $\varphi_k : B_Q \rightarrow B_{Q'}$ is a group isomorphism.*

Proof. As mutation is an involution, we can consider the composition

$$\varphi_k : B_Q \xrightarrow{\varphi_k} B_{Q'} \xrightarrow{\varphi_k} B_Q.$$

Fix some $i \in I$. Note that mutation at k does not change whether i and k are connected in the quiver; it just swaps the direction of any arrow that may exist between i and k . So if we have $i \rightarrow k$, then $s_i \mapsto t_k t_i t_k^{-1} \mapsto s_k s_i s_k^{-1}$. If we have $i \leftarrow k$, then $s_i \mapsto t_i \mapsto s_k s_i s_k^{-1}$. And if there is no arrow between i and k then $s_i \mapsto t_i \mapsto s_i$. But in this case s_i and s_k commute, so $s_i = s_k s_i s_k^{-1}$. Hence in every case $\varphi_k(s_i) = s_k s_i s_k^{-1}$, so φ_k is just a conjugation map and therefore $\varphi_k : B_Q \rightarrow B_{Q'}$ is an isomorphism. \square

Remark 2.11. The inverse of φ_k is the group isomorphism $\varphi_k^{-1} : B_{Q'} \xrightarrow{\sim} B_Q$ defined by

$$\varphi_k^{-1}(t_i) = \begin{cases} s_k^{-1} s_i s_k & \text{if } i \rightarrow k \text{ in } Q; \\ s_i & \text{otherwise.} \end{cases}$$

Noting [Remark 2.3](#), we have the following:

Theorem 2.12. *If Q is a mutation-Dynkin quiver of type Δ then $B_Q \cong B_\Delta$.*

3. Topological interpretation of the generators

Braid groups. In this section we consider quivers Q which are mutation-equivalent to an orientation of the Dynkin diagram of type Δ , where $\Delta = A_n$ or D_n . By [Theorem 2.12](#), B_Q is isomorphic to the Artin braid group B_Δ of the same Dynkin type. In other words, B_Q gives a presentation of B_Δ . In this section we give a geometric interpretation of this presentation.

We associate an oriented Riemann surface S (with boundary), together with marked points M , to Δ as follows. If $\Delta = A_n$, we take S to be a disk with $n - 3$




Figure 2. Type I (left) and type II (right) puzzle pieces for tagged triangulations in types A_n and D_n and the corresponding quivers.

marked points on its boundary, as in [Fomin and Zelevinsky 2002; 2003]. If $\Delta = D_n$, we take S to be a disk with one marked point in its interior and n marked points on its boundary, as in [Fomin et al. 2008; Schiffler 2008]. In each case, it was shown that every quiver of the corresponding mutation type arises from a triangulation of (S, M) (tagged, in the type D_n case) in the following way. We follow [Fomin et al. 2008], in a generality great enough to cover both cases (noting that there is at most one interior marked point).

A (simple) *arc* in (S, M) is a curve in S (considered up to isotopy) whose endpoints are marked points in M and which does not have any self-crossings, except possibly at its endpoints. Apart from these endpoints, it must be disjoint from M and the boundary of S , and it must not cut out an unpunctured one- or two-sided polygon.

Two arcs are said to be *compatible* if they are noncrossing in the interior of S . A maximal set of compatible arcs is a *triangulation*.

A *tagged arc* in (S, M) is an arc which does not cut out a once-punctured monogon; each of its ends is tagged, either plain or notched. Plain tags are omitted, while notched tags are displayed using the bow-tie symbol \bowtie . An end incident with a boundary marked point is always tagged plain. Two tagged arcs α, β are *compatible* if

- (i) the untagged arcs underlying α and β are compatible, and
- (ii) if the untagged versions of α and β are different but share an endpoint, then the corresponding ends of α and β are tagged in the same way.

A *tagged triangulation* \mathcal{T} of (S, M) is a maximal collection of tagged arcs in (S, M) . Note that if none of the marked points in M lies in the interior of S , every end of an arc in a tagged triangulation must be tagged plain, and tagged triangulations of S can be identified with triangulations of S .

The set M of marked points divides the boundary components of (S, M) into connected components, which we call *boundary arcs*. Note that the boundary arcs do not lie in a triangulation or tagged triangulation of (S, M) , by definition.

The tagged triangulation \mathcal{T} can be built up by gluing together puzzle pieces of the two types shown in Figure 2 (see [Fomin et al. 2008, Remark 4.2]) by gluing together along boundary arcs. Note that the puzzle piece of type II can only occur in the type D_n case, and then it occurs exactly once.

If α is an arc in a tagged triangulation \mathcal{T} , then the *flip* of \mathcal{T} at α is the unique tagged triangulation containing $\mathcal{T} \setminus \{\alpha\}$ but not containing α . By [Fomin et al. 2008], the set of tagged triangulations of (S, M) is connected under flips.

The *quiver* $Q_{\mathcal{T}}$ of a tagged triangulation \mathcal{T} has vertices corresponding to the arcs in \mathcal{T} . The quiver is built up by associating a quiver to each puzzle piece; see Figure 2. If a boundary arc in the puzzle piece is also a boundary arc of (S, M) , then the corresponding vertex in the quiver is omitted, together with all incident arrows. The quivers are then glued together by identifying vertices whenever the corresponding edges are glued together in the puzzle pieces.

In order to discuss braid groups, we need to consider more general curves in (S, M) . We define a *path* in (S, M) to be a (possibly nonsimple) curve whose endpoints lie in S (not necessarily in M).

Definition 3.1. Let \mathcal{T} be a tagged triangulation of (S, M) . We associate a graph to \mathcal{T} , which we call the *braid graph* $G_{\mathcal{T}}$ of \mathcal{T} , as follows. The vertices $V_{\mathcal{T}}$ of $G_{\mathcal{T}}$ are in bijection with the connected components of the complement of \mathcal{T} in (S, M) and, whenever two such connected components have a common tagged arc on their boundaries, there is an edge in $G_{\mathcal{T}}$ between the corresponding vertices. Thus the edges in $G_{\mathcal{T}}$ are in bijection with the arcs in \mathcal{T} .

We choose an embedding ι of $G_{\mathcal{T}}$ into (S, M) , mapping each vertex to an interior point of the corresponding connected component of the complement of \mathcal{T} in (S, M) and each edge to a path between the images of its endpoints transverse to the corresponding edge in \mathcal{T} . We identify $G_{\mathcal{T}}$ with its image under ι .

Note that in the type A case the braid graph is the tree from Section 3.1 of [Caldero et al. 2006].

We associate an orbifold X to S as follows. In the type A_n case, we just take $X = S$, and in the type D_n case we take X to be S with the interior marked point of S interpreted as a cone point of order two. In each case, the set M of marked points induces a corresponding set of marked points in X , which we also denote




Figure 3. Thickening of the path π (the middle path).

by M . Each arc or tagged arc α in (S, M) induces a corresponding arc or tagged arc in (X, M) which we also denote by α . Thus each (tagged) triangulation \mathcal{T} of (S, M) induces a corresponding set \mathcal{T} of (tagged) arcs in (X, M) .

Note also that orbifolds have been used to model cluster algebras in [Felikson et al. 2012]. In this approach, the model for B_n is an orbifold with a cone point of order two, regarded as a folding of D_n , where D_n is modelled by a disk with a single interior marked point (see also Lecture 15 of [Thurston 2012], which was given by A. Felikson).

We denote by X° the orbifold X with the cone point (if there is one) removed (so $X^\circ = X$ in type A_n). Given any set V of vertices in X° , we may consider the corresponding *braid group*, $\Gamma(X, V)$ following [Allcock 2002]. Each element of $\Gamma(X, V)$ (or *braid*) can be regarded as a permutation g of V together with a tuple $\gamma = (\gamma_v)_{v \in V}$ of paths $\gamma_v : [0, 1] \rightarrow X^\circ$ such that $\gamma_v(0) = v$ and $\gamma_v(1) = g(v)$ for each $v \in V$. In addition, for each $t \in [0, 1]$, the points $\gamma_v(t)$ for $v \in V$ must all be distinct for all $v \in V$. Braids are considered up to isotopy, and two braids can be multiplied by composing the paths in a natural way; we compose braids from right to left, as for functions.

Remark 3.2. Suppose V and V' are two sets of points in X° and there is a bijection $\rho : V \rightarrow V'$. Suppose also that there is a set of paths $\delta_v : [0, 1] \rightarrow X^\circ$, for $v \in V$, with $\delta_v(0) = v$ and $\delta_v(1) = \rho(v)$ for all $v \in V$. Suppose furthermore that the points $\gamma_v(t)$ for $v \in V$ and $t \in [0, 1]$ are all distinct. Then the maps δ_v induce a natural isomorphism between $\Gamma(X, V)$ and $\Gamma(X, V')$.

Definition 3.3. Each path π in X° with endpoints v_1, v_2 in V determines a braid σ_π in $\Gamma(X, V)$ as follows (see [Fox and Neuwirth 1962, §7]). We thicken the path π along its length (avoiding the other vertices), closing it off at the end points to form a (topological) disk. We give the boundary of the disk the clockwise orientation. The vertices v_1 and v_2 divide the boundary of the disk into two paths, one from v_1 to v_2 and the other from v_2 to v_1 . We define γ_{v_1} to be the former and γ_{v_2} to be the latter. See Figure 3. For $v \in V$ such that $v \neq v_1, v_2$, we define $\gamma_v(t)$ to be v for all $t \in [0, 1]$. Then σ_π is the braid $(\gamma_v)_{v \in V}$. Note that σ_π only depends on the isotopy class of the image of π in (X, V) . In particular, it is unchanged if π is reversed.




Figure 4. The braid σ_π .

An example of a braid σ_π is displayed as a picture (in the same way as in [Allcock 2002]) in Figure 4. In this figure only, we display π as a dashed line to distinguish it from the braid σ_π .

Interpretation of the generators. Let \mathcal{T} be a triangulation of (S, M) . Let $Q_\mathcal{T}$ be the quiver of \mathcal{T} . Then $Q_\mathcal{T}$ has vertices I corresponding to the arcs in \mathcal{T} . We denote the arc in \mathcal{T} associated to $i \in I$ by α_i . The corresponding edge in $G_\mathcal{T}$ is denoted π_i . Let $\sigma_i = \sigma_{\pi_i} \in \Gamma(X, V_\mathcal{T})$ be the corresponding braid. We define $H_\mathcal{T}$ to be the subgroup of $\Gamma(X, V_{\mathcal{T}_0})$ generated by the braids σ_i for $i \in I$.

Let \mathcal{T}_0 be an initial triangulation of (S, M) defined as follows. In the type A_n case, we choose a marked point P in M and take noncrossing arcs between P and each of the other marked points in M not incident with a boundary arc incident with P . In the type D_n case, we choose two marked points P, Q on the boundary of S . We take two arcs between the interior marked point and Q , one tagged plain at the interior marked point and the other one tagged notched, and an arc between P and Q (not homotopic to a boundary arc). We then take (noncrossing) arcs between P and every other marked point in M on the boundary of S not incident with a boundary arc incident with P . See Figure 5. Then the quiver $Q_{\mathcal{T}_0}$ associated to $Q_{\mathcal{T}_0}$ is a Dynkin quiver of type Δ . By Remark 2.3, $B_{Q_{\mathcal{T}_0}}$ is isomorphic to the Artin braid group of type Δ .

Proposition 3.4. *Let \mathcal{T}_0 be the triangulation of (X, M) defined as above. Then there is an isomorphism from $H_{\mathcal{T}_0}$ to $B_{Q_{\mathcal{T}_0}}$ taking the braid σ_i to the generator s_i of $B_{Q_{\mathcal{T}_0}}$. Furthermore, in type A_n , the subgroup $H_{\mathcal{T}_0}$ coincides with $\Gamma(X, V_{\mathcal{T}_0})$, while in type D_n , it is a subgroup of $\Gamma(X, V_{\mathcal{T}_0})$ of index two.*

Proof. For type A_n , see [Fox and Neuwirth 1962] and the explanation in [Allcock 2002, §4]. For type D_n , note that the elements σ_i for $i \in I$ coincide with the generators h_i defined in [Allcock 2002, §1] (via an isomorphism of the kind in Remark 3.2). The result then follows from [Allcock 2002, Theorem 1]. \square

The following lemma appears in [Sergiescu 1993, Théorème, part (iv)].

Lemma 3.5. *Let A, B, C be three distinct points in X° and suppose there is a topological disk in X° , with A, B and C lying in order clockwise around its boundary.*




Figure 5. Initial triangulations and the corresponding braid graphs and quivers.

Let AB denote the arc on this boundary between A and B . We define BC and CA similarly. Then $\sigma_{AB}\sigma_{BC} = \sigma_{BC}\sigma_{CA}$.

Theorem 3.6. *Let \mathcal{T} be an arbitrary tagged triangulation of (X, M) . Then there is an isomorphism from $H_{\mathcal{T}}$ to $B_{Q_{\mathcal{T}}}$ taking the braid σ_i to the generator s_i of $B_{Q_{\mathcal{T}}}$. Furthermore, in type A_n , the subgroup $H_{\mathcal{T}}$ coincides with $\Gamma(X, V)$, while in type D_n , it is a subgroup of $\Gamma(X, V)$ of index two.*

Proof. The result holds for $\mathcal{T} = \mathcal{T}_0$ by Proposition 3.4. Note that any triangulation can be obtained from \mathcal{T}_0 by applying a finite number of flips of tagged triangulations. We show that the theorem is true for an arbitrary tagged triangulation \mathcal{T} by induction on the number of flips required to obtain \mathcal{T} from \mathcal{T}_0 . To do this, it is enough to show that if the theorem holds for a tagged triangulation \mathcal{T} and α_i is a tagged arc in \mathcal{T} then the theorem also holds for the flip of \mathcal{T} at α_i .

So we assume the result holds for a tagged triangulation \mathcal{T} . Thus there is an isomorphism $\psi_{\mathcal{T}} : H_{\mathcal{T}} \rightarrow B_{Q_{\mathcal{T}}}$ sending σ_i to s_i . We denote the corresponding elements of $H_{\mathcal{T}}$ by τ_i and t_i . The tagged arcs in \mathcal{T} are denoted by α_i , for $i \in I$, and we denote the corresponding tagged arcs in \mathcal{T}' by β_i , for $i \in I$. The edges of $G_{\mathcal{T}}$ are denoted π_i , and we denote the edges of $G_{\mathcal{T}'}$ by ρ_i .

We define:

$$\tilde{\tau}_i = \begin{cases} \sigma_k^{-1} \sigma_i \sigma_k, & \text{if } i \rightarrow k \text{ in } Q; \\ \sigma_i, & \text{otherwise.} \end{cases}$$

Then it is easy to see that $H_{\mathcal{T}}$ is generated by the $\tilde{\tau}_i$ for $i \in I$.




Figure 6. Flip involving an arc (α_1) where two puzzle pieces of type I are glued.

We consider the possible types of flip that can occur, which are determined by the fact that \mathcal{T} can be constructed out of puzzle pieces. Suppose first that \mathcal{T}' is the flip of \mathcal{T} at an arc α where two puzzle pieces of type I are glued together. We label the corresponding vertices in I by 1, 2, 3, 4, 5, for simplicity, and suppose we are flipping at the edge in \mathcal{T} dual to α_1 . The braid graph local to the flip is shown in the left-hand diagram in Figure 6. Applying Lemma 3.5, we see that the middle figure shows paths $\tilde{\pi}_i$ with the property that $\tilde{\tau}_i = \sigma_{\tilde{\pi}_i}$ for $i = 1, 2, 3, 4, 5$.

Rotating vertices A and B clockwise, to get the right-hand diagram in Figure 6, we obtain, via Remark 3.2, an isomorphism from $H_{\mathcal{T}}$ to $H_{\mathcal{T}'}$ taking $\tilde{\tau}_i$ to τ_i for all $i \in I$. The inverse is an isomorphism from $H_{\mathcal{T}'}$ to $H_{\mathcal{T}}$ taking τ_i to $\sigma_k^{-1} \sigma_i \sigma_k$ if there is an arrow $i \rightarrow k$ in Q and to σ_i otherwise. Composing with the isomorphism $\varphi_k \circ \psi_{\mathcal{T}}$, where φ_k is the isomorphism in Proposition 2.9, we obtain an isomorphism from $H_{\mathcal{T}'}$ to $B_{Q_{\mathcal{T}'}}$ taking τ_i to t_i as required. This proves the required result in type A, so we are left with the type D case, where puzzle pieces of type II may occur.

We next consider a flip inside a puzzle piece of type II. We can apply essentially the same argument; see Figures 7 and 8. Here we draw the puzzle piece together with the two adjacent triangles, necessarily of type I (since there is only one cone point). To go from the middle diagram to the right-hand diagram in Figure 8, the vertex D should be moved anticlockwise around the cone point. We use the fact that in the right-hand diagram of Figure 8, the resulting path $\tilde{\pi}_1$ is isotopic to the path ρ_1 in $G_{\mathcal{T}'}$, using the fact that the cone point has order two.

Note that the adjoining type I puzzle pieces (in Figures 7 and 8) may not occur, but the argument is easily modified to cover these cases. We also need to consider




Figure 7. Flip (at α_1) inside a puzzle piece of type II, first case.




Figure 8. Flip (at α_2) inside a puzzle piece of type II, second case.

the flips from the right-hand diagram in each case to the corresponding left-hand one. We omit the details; a similar argument can be applied in these cases.

Finally, we need to consider a flip involving an arc where a puzzle piece of type I and a puzzle piece of type II have been glued together. These cases are shown in Figures 9 and 10: Figure 9 illustrates the case where the puzzle piece of type I is on the left of the puzzle piece of type II (when it is drawn as shown), while Figure 10 illustrates the case where it is on the right. Again, a similar argument applies in the case of flips from the right-hand diagram to the left-hand one in these cases. \square




Figure 9. Flip involving an arc (α_3) where puzzle pieces of type I and II are glued, first case.




Figure 10. Flip involving an arc (α_4) where puzzle pieces of type I and II are glued, second case.

4. Actions on categories

Quivers with potential. Fix an algebraically closed field \mathbb{F} . To any quiver Q we can associate the path algebra $\mathbb{F}Q$, which, as an \mathbb{F} -vector space, has basis given by all paths in Q of length ≥ 0 , and the multiplication of two paths p_1 and p_2 is

their concatenation $p_1 p_2$ if p_1 ends and p_2 starts at the same vertex, and is zero otherwise.

Let $\mathbb{F}Q_{\geq n}$ be the ideal of $\mathbb{F}Q$ generated by the paths in Q of length at least n . We can take the completion $\widehat{\mathbb{F}Q}$ of $\mathbb{F}Q$ with respect to $\mathbb{F}Q_{\geq 1}$, which is defined as:

$$\widehat{\mathbb{F}Q} = \varprojlim_n \frac{\mathbb{F}Q}{\mathbb{F}Q_{\geq n}} = \{(a_n + \mathbb{F}Q_{\geq n})_{n=1}^{\infty} \mid a_n \in \mathbb{F}Q, \varphi_n(a_n + \mathbb{F}Q_{\geq n}) = a_{n-1} + \mathbb{F}Q_{\geq n-1}\},$$

where the limit is taken along the chain of epimorphisms

$$\frac{\mathbb{F}Q}{\mathbb{F}Q_{\geq 1}} \xleftarrow{\varphi_2} \frac{\mathbb{F}Q}{\mathbb{F}Q_{\geq 2}} \xleftarrow{\varphi_3} \frac{\mathbb{F}Q}{\mathbb{F}Q_{\geq 3}} \xleftarrow{\dots}$$

Let $\widehat{\mathbb{F}Q}_{\text{cyc}}$ denote the subspace of (possibly infinite) linear combinations of cycles in Q . Recall that a *potential* for a quiver Q is an element W of $\widehat{\mathbb{F}Q}_{\text{cyc}}$, regarded up to cyclic equivalence (and for which no two cyclically equivalent paths in Q occur in the decomposition of W). The pair (Q, W) is called a *quiver with potential* [Derksen et al. 2008], which we occasionally abbreviate to QP. The following definition is due to Derksen, Weyman and Zelevinsky:

Definition 4.1 [Derksen et al. 2008, Definition 4.2]. Let Q_1 and Q_2 be two quivers with the same vertex set I , and (Q_1, W_1) and (Q_2, W_2) be two QPs. A *right equivalence* between (Q_1, W_1) and (Q_2, W_2) is an algebra isomorphism $\varphi: \widehat{\mathbb{F}Q}_1 \rightarrow \widehat{\mathbb{F}Q}_2$ such that $\varphi(W_1)$ is cyclically equivalent to W_2 and φ is the identity when restricted to the semisimple subalgebra $\mathbb{F}I$ of $\widehat{\mathbb{F}Q}_1$.

A quiver with potential (Q, W) with W containing paths of length two or more is *trivial* if Q is a disjoint union of 2-cycles and there is an algebra automorphism of $\widehat{\mathbb{F}Q}$ preserving the span of the arrows of Q (a *change of arrows*) which takes W to the sum of the 2-cycles in Q . A quiver with potential (Q, W) is said to be *reduced* if W is a linear combination of cycles in Q of length 3 or more.

The *splitting theorem* [Derksen et al. 2008, Theorem 4.6] states that every quiver with potential can be written as a direct sum of a reduced quiver with potential and a trivial quiver with potential which are unique up to right equivalence.

Let (Q, W) be a quiver with potential, and let k be a vertex of Q not involved in any 2-cycles. By replacing W with a cyclically equivalent potential on Q if necessary, we can assume that none of the cycles in the decomposition of W start or end at k . We denote by $\tilde{\mu}_k(Q, W)$ the nonreduced mutation of (Q, W) at k in Q , as defined in [Derksen et al. 2008, §5]. Then, by Theorem 5.2 of the same paper, the right equivalence class of $\tilde{\mu}_k(Q, W)$ is determined by the right equivalence class of (Q, W) . The *mutation* $\mu_k(Q, W)$ of (Q, W) at k is then defined to be the reduced component of $\tilde{\mu}_k(Q, W)$, and is uniquely determined up to right equivalence, given the right equivalence class of (Q, W) .




Figure 11. Terms in the potential W_T .

As before, we will say a quiver with potential (Q, W) is *Dynkin* if the underlying unoriented graph of Q is an orientation of a Dynkin quiver (and hence $W = 0$). We shall say that a quiver with potential (Q', W') is *mutation-Dynkin* if it can be obtained by repeatedly mutating a Dynkin quiver with potential in the above sense. *Note:* For the rest of this subsection we will restrict to Dynkin types A and D .

Let (S, M) be the Riemann surface with marked points associated to Δ as in Section 3. So, if $\Delta = A_n$, we take S to be a disk with $n - 3$ points on its boundary, and if $\Delta = D_n$, we take S to be a disk with one marked point in its interior and n marked points on its boundary.

Let Q be a mutation-Dynkin quiver. By [Fomin et al. 2008], $Q = Q_{\mathcal{T}}$ for some tagged triangulation \mathcal{T} of (S, M) . Let W, W' be the sum of the terms coming from local configurations in \mathcal{T} as shown in Figure 11 (where in (c) and (d) there are at least three arcs incident with the interior marked point).

Then $W_{\mathcal{T}}$ is the potential given by taking the sum of the induced cycles in $Q_{\mathcal{T}}$ (i.e., induced subgraphs of $Q_{\mathcal{T}}$ which are cycles), and $W'_{\mathcal{T}}$ is the potential associated to \mathcal{T} in [Labardini-Fragoso 2016, §3], taking the parameter associated to the internal marked point (if there is one) to be equal to -1 . Then we have the following:

Lemma 4.2. *The potentials $W_{\mathcal{T}}$ and $W'_{\mathcal{T}}$ are right equivalent.*

Proof. We assume we are in case D_n , since the two potentials coincide in case A_n . If the interior marked point is as in case (c) of Figure 11 (with at least 3 arcs incident with it), then there is a unique triangle in \mathcal{T} with sides 1 and 2. We label the arrows in the corresponding 3-cycle in $W_{\mathcal{T}}$ or $W'_{\mathcal{T}}$ by a, x, y , in order around the cycle. Then the automorphism φ of $\widehat{kQ_{\mathcal{T}}}$ negating a and x and taking each other arrow to itself gives a right equivalence between $W_{\mathcal{T}}$ and $W'_{\mathcal{T}}$, since a and x are not involved in any other terms in any of these potentials.

If the interior marked point is as in case (d), then $W_{\mathcal{T}}$ and $W'_{\mathcal{T}}$ coincide. \square

We recall the following special case of [Labardini-Fragoso 2016, Theorem 8.1].

Theorem 4.3 [Labardini-Fragoso 2016]. *Let $\mathcal{T}, \mathcal{T}'$ be triangulations of (S, M) . If \mathcal{T}' is obtained from \mathcal{T} by flipping at an arc α_k then $\mu_k(Q_{\mathcal{T}}, W'_{\mathcal{T}})$ is right equivalent to $(Q_{\mathcal{T}'}, W'_{\mathcal{T}'})$.*

By [Derksen et al. 2008, Theorem 7.1], it follows from this that the quiver of $\mu_k(Q_{\mathcal{T}}, W_{\mathcal{T}})$ coincides with the quiver obtained from $Q_{\mathcal{T}}$ by Fomin–Zelevinsky quiver mutation at k .

Hence we can effectively ignore potentials:

Proposition 4.4. *Any mutation-Dynkin quiver with potential (\tilde{Q}, \tilde{W}) of type A or D is right equivalent to $(\tilde{Q}, W_{\tilde{Q}})$, where $W_{\tilde{Q}}$ is the sum of all chordless cycles in \tilde{Q} .*

Proof. Note that a Dynkin quiver with zero potential is of the form $(Q_{\mathcal{T}}, W_{\mathcal{T}})$ for some triangulation \mathcal{T} ; see [Fomin et al. 2008]. Suppose that (\tilde{Q}, \tilde{W}) is obtained from a Dynkin quiver with zero potential by iterated mutation in the sense of [Derksen et al. 2008]. Then, by Theorem 4.3 and Lemma 4.2, (\tilde{Q}, \tilde{W}) is right equivalent to $(Q_{\mathcal{T}}, W_{\mathcal{T}})$ for some triangulation \mathcal{T} of (S, M) . \square

Note that an alternative proof of Proposition 4.4 would be to compute the mutation of a quiver with potential $(Q_{\mathcal{T}}, W'_{\mathcal{T}})$ directly, and show that it is right equivalent

to $(Q_{\mathcal{T}}, W'_{\mathcal{T}})$. This is not too difficult to do, but requires consideration of several cases and still requires arguments dealing with changes of sign as in [Lemma 4.2](#), so we instead refer to [\[Labardini-Fragoso 2016\]](#) above.

Differential graded algebras and modules. Let \mathbb{F} be an algebraically closed field. We think of \mathbb{F} as a graded \mathbb{F} -algebra concentrated in degree 0. If $V = \bigoplus V_i$ is a graded \mathbb{F} -module then let $V[j]$ be the graded \mathbb{F} -module with $(V[j])_i = V_{i+j}$. If $f : V \rightarrow W$ is a map of graded vector spaces with homogeneous components $f_i : V_i \rightarrow W_i$ then let $f[j] : V[j] \rightarrow W[j]$ be the map of graded vector spaces with homogeneous components $f[j]_i : V[j]_i \rightarrow W[j]_i$ defined by $f[j]_i(v) = (-1)^j f_{i+j}(v)$ for $v \in V[j]_i = V_{i+j}$. Thus $[1]$ is an endofunctor of the category of graded \mathbb{F} -modules, called the *shift functor*.

If V and W are graded vector spaces, we will refer to a map $f : V \rightarrow W[j]$ of graded vector spaces as a map from V to W of degree j . We use the Koszul sign rule for graded \mathbb{F} -algebras, so if $f : V \rightarrow V'$ and $g : W \rightarrow W'$ are maps of graded \mathbb{F} -modules of degree m and n then

$$(f \otimes g)(v \otimes w) = (-1)^{in} f(v) \otimes g(w)$$

for $v \in V_i$ and $w \in W$.

A unital differential graded algebra (or dg-algebra, or dga) over \mathbb{F} is a graded \mathbb{F} -algebra $A = \bigoplus_{i \in \mathbb{Z}} A_i$ with multiplication $m : A \otimes_{\mathbb{F}} A \rightarrow A$ of degree 0 together with a unit $\iota : \mathbb{F} \hookrightarrow A$ and an \mathbb{F} -linear differential $d : A \rightarrow A$ of degree +1. These should satisfy

- the associativity relation $m \circ (1 \otimes m) = m \circ (m \otimes 1)$;
- the boundary relation $d^2 = 0$;
- the Leibniz relation $d \circ m = m \circ (1 \otimes d + d \otimes 1)$;
- the unital relation $m \circ (\text{id}_A \otimes \iota) = m \circ (\iota \otimes \text{id}_A)$, which should agree with the \mathbb{F} -algebra structure of A .

We often denote our dga by (A, d) , or simply by A . Each dga (A, d) has an underlying unital graded algebra, obtained by simply forgetting the differential, which we denote $u(A)$.

A left module M for A is a graded left \mathbb{F} -module M which has a left action $m_M : A \otimes M \rightarrow M$ of $u(A)$ together with a map $d_M : M \rightarrow M$ of degree +1, called a *differential*, such that

$$d_M \circ m_M = m_M \circ (1 \otimes d_M + d \otimes 1).$$

We always have the *regular* module $M = A$ with $d_M = d$ and $m_M = m$. Similarly, a right module M for A is a graded right \mathbb{F} -module M which has a right action $m_M : M \otimes A \rightarrow M$ of $u(A)$ together with a differential d_M such that $d_M \circ m_M = m_M \circ (1 \otimes d + d_M \otimes 1)$. If (M, d_M) is an A -module, then $(M[1], d_M[1])$ is also an

A -module, which we sometimes just write as $M[1]$. Modules for A are modules for $u(A)$, simply by forgetting the differential.

A map $f : M \rightarrow N$ of left A -modules is a degree 0 map of $u(A)$ -modules such that f commutes with the differentials: $d_N \circ f = f \circ d_M$. We thus obtain a category $A\text{-Mod}$ of left A -modules, and we write the morphism spaces in this category as $\text{Hom}_{A\text{-Mod}}(M, N)$. $A\text{-Mod}$ is an \mathbb{F} -category: each morphism space is an \mathbb{F} -module.

Given two differential algebras (A, d_A) and (B, d_B) , an A - B -bimodule (M, d_M) is a graded \mathbb{F} -module which is a left (A, d_A) -module with left action m^ℓ and a right (B, d_B) -module with right action m^r where the two actions commute:

$$m^r \circ (m^\ell \otimes \text{id}_B) = m^\ell \circ (\text{id}_A \otimes m^r).$$

We will always assume that \mathbb{F} acts centrally. Under this assumption we can, and will, identify left A -modules with A - k -bimodules and A - B -bimodules with left $A \otimes_{\mathbb{F}} B^{\text{op}}$ -modules, where B^{op} denotes the algebra B with the order of multiplication reversed. A map of bimodules should commute with the differential on both the left and the right, and we obtain an \mathbb{F} -category $A\text{-Mod-}B$ of A - B -bimodules.

Given a map $f : M \rightarrow N$ of left A -modules, we can construct a new left A -module called the *cone* of f , denoted $\text{cone}(f)$. As a left module for $u(A)$, we have $\text{cone}(f) = N \oplus M[1]$. The differential is given by

$$\begin{pmatrix} d_N & 0 \\ f[1] & d_{M[1]} \end{pmatrix}.$$

If L is isomorphic to $\text{cone}(f)$ for some map $f : M \rightarrow N$, we say that L is an *extension* of M by $L[-1]$.

We will use the following lemma, whose proof follows immediately from the definitions, repeatedly.

Lemma 4.5. *Let $f : M \rightarrow N$ be a map in $A\text{-Mod}$.*

- (i) *Let $F : A\text{-Mod} \rightarrow B\text{-Mod}$ be an additive functor which commutes with the shift functor. Then we have an isomorphism $\text{cone}(Ff) \cong F \text{cone}(f)$ in $B\text{-Mod}$.*
- (ii) *For any commutative diagram*

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \sim \downarrow \varphi_M & & \sim \downarrow \varphi_N \\ M' & \xrightarrow{f'} & N' \end{array}$$

in $A\text{-Mod}$ where both φ_M and φ_N are isomorphisms, we have an isomorphism $\varphi_N \oplus \varphi_M[1] : \text{cone}(f) \rightarrow \text{cone}(f')$ of A -modules.

Let (A, d_A) , (B, d_B) , and (C, d_C) be dgas. If (M, d_M) is an A - B -bimodule and (N, d_N) is an A - C -bimodule then let $\text{Hom}_A^i(M, N)$ be the space of all graded left

$u(A)$ -module maps $f : M \rightarrow N$ of degree i . We do not require that these maps commute with the differential. We define

$$\text{Hom}_A(M, N) = \bigoplus_{i \in \mathbb{Z}} \text{Hom}_A^i(M, N),$$

and this is a graded $u(B)$ - $u(C)$ -bimodule. We also have a version for right modules, which we write as $\text{Hom}_{A^{\text{op}}}(M, N)$.

Note the distinction between $\text{Hom}_A(M, N)$ and the hom spaces in the category $A\text{-Mod}$. With the differential $d(f) = d_N \circ f - (-1)^i f \circ d_M$ for $f \in \text{Hom}_A^i(M, N)$, $\text{Hom}_A(M, N)$ becomes a B - C -bimodule. Similarly, if (M, d_M) is a B - A -bimodule and (N, d_N) is a C - A -bimodule, $\text{Hom}_{A^{\text{op}}}(M, N)$ is a C - B -bimodule. $\text{Hom}_A(-, -)$ is the internal hom in the bimodule category, and we can recover the hom spaces in $A\text{-Mod}$ as the 0-cycles of $\text{Hom}_A(M, N)$.

If (M, d_M) is an A - B -bimodule and (N, d_N) is a B - C -bimodule then let $M \otimes_B N$ denote the space $M \otimes_{u(B)} N$. It is a graded $u(A)$ - $u(C)$ -bimodule: if $m \in M_i$ and $n \in N_j$ then $m \otimes n$ has degree $i + j$. With the differential $d_M \otimes \text{id}_N + \text{id}_M \otimes d_N$, it becomes an A - C -bimodule.

For an A - B -bimodule (M, d_M) , we thus have functors

$$M \otimes_B - : B\text{-Mod} \rightarrow A\text{-Mod} \quad \text{and} \quad \text{Hom}_A(M, -) : A\text{-Mod} \rightarrow B\text{-Mod}.$$

The functor $M \otimes_B -$ is left adjoint to $\text{Hom}_A(M, -)$. For (N, d_N) a left A -module, the counit $\text{ev}_N : M \otimes_B \text{Hom}_A(M, N) \rightarrow N$ of the adjunction is the evaluation map, which acts as $x \otimes f \mapsto (-1)^{ij} f(m)$ for $x \in M_i$ and $f \in \text{Hom}_A^j(M, N)$.

Derived categories. Our references are [Keller 1994; 2006].

If A is a graded vector space and d is a differential, i.e., a degree $+1$ endomorphism of A which satisfies $d^2 = 0$, then the i -th homology of A , denoted $H_i(A)$, is the subquotient $\ker d_i / \text{im } d_{i-1}$, where $d_i : A_i \rightarrow A_{i+1}$ denotes the restriction of d to A_i . If (A, d) is a dga then the homology $H(A) = \bigoplus H_i(A)$ is a graded algebra, and if M is a left A -module then $H(M) = \bigoplus H_i(M)$ is a left $H(A)$ -module. In fact, taking homology is a functor from the category of A -modules to the category of graded $H(A)$ -modules. We say that a left A -module M is *acyclic* if $H(M) = 0$, and that a map $f : M \rightarrow N$ of A -modules is a *quasi-isomorphism* if $H(f)$ is an isomorphism.

The *category up to homotopy* of $A\text{-Mod}$, denoted $\mathbb{K}(A)$, is the \mathbb{F} -category whose objects are all left A -modules and whose morphism spaces, for $M, N \in A\text{-Mod}$, are $\text{Hom}_{\mathbb{K}(A)}(M, N) = H_0 \text{Hom}_A(M, N)$. The *derived category* of A , denoted $\mathbb{D}(A)$, is the \mathbb{F} -category obtained by localizing $\mathbb{K}(A)$ at the full subcategory of acyclic A -modules. As a map of modules is a quasi-isomorphism if and only if its cone is acyclic, this is equivalent to localizing $\mathbb{K}(A)$ at the class of all quasi-isomorphisms. So we have a canonical functor $\mathbb{K}(A) \rightarrow \mathbb{D}(A)$, which we call the projection functor.

The finite-dimensional derived category, denoted $D_{\text{fd}}(A)$, is the full subcategory of $D(A)$ on objects with finite-dimensional total homology, i.e., on A -modules M such that $H(M)$ is a finite-dimensional \mathbb{F} -vector space.

Let (A, d_A) be a dga. We say that

- $P \in A\text{-Mod}$ is *indecomposable projective* if it is an indecomposable direct summand of the regular module,
- $P \in A\text{-Mod}$ is *relatively projective* if it is a direct sum of shifts of indecomposable projective modules, and
- $P \in A\text{-Mod}$ is *cofibrant* if, for each surjective quasi-isomorphism $f : M \rightarrow N$, the map $\text{Hom}_{A\text{-Mod}}(P, f) : \text{Hom}_{A\text{-Mod}}(P, M) \rightarrow \text{Hom}_{A\text{-Mod}}(P, N)$ is surjective.

The following result characterizes cofibrant modules.

Proposition 4.6 [Keller 1994, Section 3; Keller and Yang 2011, Proposition 2.13]. *An A -module P is cofibrant if and only if it is an iterated extension of a relatively projective module by other relatively projective modules, possibly infinitely many times.*

Let $A\text{-cofib}$ denote the full subcategory of $K(A)$ on the cofibrant objects. The projection functor $K(A) \rightarrow D(A)$ induces an equivalence $A\text{-cofib} \xrightarrow{\sim} D(A)$. Each A -module M has a *cofibrant replacement*, defined up to quasi-isomorphism and denoted $\mathbf{p}M$, which can be realized as the image of M under the left adjoint $D(A) \rightarrow K(A)$ to the canonical projection functor [Keller 2006, Proposition 3.1].

Let (B, d_B) be another dga and let $F : A\text{-Mod} \rightarrow B\text{-Mod}$ be an additive functor. Then F preserves chain homotopies, and so induces a functor $K(F) : K(A) \rightarrow K(B)$. If $K(F)$ preserves quasi-isomorphisms then, by the universal property of localization, it induces a functor $D(F) : D(A) \rightarrow D(B)$. If $P \in A\text{-Mod-}B$ is cofibrant as a left A -module then, by [Keller 1994, Theorem 3.1(a)] and [Keller and Yang 2011, Proposition 2.13], $\text{Hom}_A(P, -)$ preserves acyclic modules, and so preserves quasi-isomorphisms. By imitating the proof of [Keller 1994, Theorem 3.1(a)] we see that if $P \in A\text{-Mod-}B$ is cofibrant as a right B -module then $P \otimes_B -$ also preserves acyclic modules. We often write $P \otimes_B -$ and $\text{Hom}_A(P, -)$, instead of $D(P \otimes_B -)$ and $D(\text{Hom}_A(P, -))$, for the induced functors $D(B) \rightarrow D(A)$ and $D(A) \rightarrow D(B)$.

For an arbitrary $M \in A\text{-Mod-}B$, we get a functor $M \otimes_B^L - : D(B) \rightarrow D(A)$, known as the *left derived functor* of $M \otimes_B -$, by composing the cofibrant replacement functor $D(B) \rightarrow K(B)$, the tensor functor $K(M \otimes_B -) : K(B) \rightarrow K(A)$, and the projection functor $K(A) \rightarrow D(B)$. By [Keller 1994, Lemma 6.3(a)], we have an isomorphism $M \otimes_B^L N \cong \mathbf{p}M \otimes_B N$ for all $N \in D(B)$. The following basic, but useful, lemma says that this isomorphism is natural.

Lemma 4.7. *Let $M \in \text{Mod-}B$.*

- (i) *We have a natural isomorphism of functors $\mathbf{p}M \otimes_B - \cong M \otimes_B^L -$.*

(ii) If M is cofibrant we have a natural isomorphism of functors $M \otimes_B - \cong M \otimes_B^L -$.

Proof. (i) We need to show that for each $N \in \mathcal{B}\text{-Mod}$ there is a quasi-isomorphism $\varphi_N : \mathbf{p} M \otimes_B N \rightarrow M \otimes_B \mathbf{p} N$ such that, for all maps $f : N \rightarrow N'$, the diagram

$$\begin{array}{ccc} \mathbf{p} M \otimes_B N & \xrightarrow{\varphi_N} & M \otimes_B \mathbf{p} N \\ \downarrow \mathbf{p} M \otimes f & & \downarrow M \otimes \mathbf{p} f \\ \mathbf{p} M \otimes_B N' & \xrightarrow{\varphi_{N'}} & M \otimes_B \mathbf{p} N' \end{array}$$

commutes. Consider the following diagram:

$$\begin{array}{ccccc} & \mathbf{p} M \otimes_B \mathbf{p} N & & & \\ & \swarrow \mathbf{p} M \otimes \pi_N & & \searrow \pi_M \otimes \mathbf{p} N & \\ \mathbf{p} M \otimes_B N & \xrightarrow{\varphi_N} & M \otimes_B \mathbf{p} N & & \\ & \searrow \mathbf{p} M \otimes f & & \swarrow M \otimes \mathbf{p} f & \\ & & \mathbf{p} M \otimes_B N' & \xrightarrow{\varphi_{N'}} & M \otimes_B \mathbf{p} N' \end{array}$$

As $\mathbf{p} M$ and $\mathbf{p} N$ are cofibrant and π_M and π_N are quasi-isomorphisms, both $\mathbf{p} M \otimes \pi_N$ and $\pi_M \otimes \mathbf{p} N$ are quasi-isomorphisms, therefore we can define $\varphi_N = (\pi_M \otimes \mathbf{p} N) \circ (\mathbf{p} M \otimes \pi_N)^{-1}$ and it is a quasi-isomorphism. Then to check naturality we need to show that

$$(M \otimes \mathbf{p} f) \circ (\pi_M \otimes \mathbf{p} N) \circ (\mathbf{p} M \otimes \pi_N)^{-1} = (\pi_M \otimes \mathbf{p} N') \circ (\mathbf{p} M \otimes \pi_{N'})^{-1} \circ (\mathbf{p} M \otimes f).$$

By the bifunctionality of the tensor product, the left-hand side is equal to

$$(\pi_M \otimes \mathbf{p} N') \circ (\mathbf{p} M \otimes \mathbf{p} f) \circ (\mathbf{p} M \otimes \pi_N)^{-1}$$

so we just need to show that

$$\mathbf{p} f \circ \pi_N^{-1} = \pi_{N'}^{-1} \circ f$$

but this follows from the functoriality $f \circ \pi_N = \pi_{N'} \circ \mathbf{p} f$ of the cofibrant replacement functor \mathbf{p} .

(ii) We just need to show that, for M cofibrant, there is a natural isomorphism $M \otimes_B - \cong \mathbf{p} M \otimes_B -$, and then the result will follow by part (i) of the lemma. This follows because $\pi_M : \mathbf{p} M \rightarrow M$ is a quasi-isomorphism and by the bifunctionality of the tensor product. \square

If the functor $M \otimes_B^L -$ is an equivalence $\mathcal{D}(B) \xrightarrow{\sim} \mathcal{D}(A)$, we say that M is a *tilting module*.

We say that a module M is of *finite projective dimension* if its cofibrant replacement is an iterated extension of finitely many shifted indecomposable projective modules.

The following basic lemma will also be useful. It can be found as [Keller 1994, Lemma 6.2(a)]. We include a proof for the convenience of the reader.

Lemma 4.8. *Let M be a left A -module of finite projective dimension and let P be its cofibrant replacement. Then we have a natural isomorphism of functors*

$$\mathrm{Hom}_A(P, A) \otimes_A - \xrightarrow{\sim} \mathrm{Hom}_A(P, -) : A\text{-Mod} \rightarrow \mathbb{F}\text{-Mod}.$$

Proof. First note that, for any $P \in A\text{-Mod}$, we always have a natural transformation

$$\mathrm{Hom}_A(P, A) \otimes_A - \rightarrow \mathrm{Hom}_A(P, -)$$

obtained by starting with the map

$$\mathrm{ev} \otimes 1 : (P \otimes_{\mathbb{F}} \mathrm{Hom}_A(P, A)) \otimes_A M \rightarrow A \otimes_A M,$$

using the associativity isomorphism to obtain a map

$$P \otimes_{\mathbb{F}} (\mathrm{Hom}_A(P, A) \otimes_A M) \rightarrow A \otimes_A M,$$

then using the adjunction

$$\begin{aligned} \mathrm{Hom}_{\mathbb{F}}(\mathrm{Hom}_A(P, A) \otimes_A M, \mathrm{Hom}_A(P, A \otimes_A M)) \\ \cong \mathrm{Hom}_A(P \otimes_{\mathbb{F}} (\mathrm{Hom}_A(P, A) \otimes_A M), A \otimes_A M), \end{aligned}$$

and finally using the natural isomorphism $A \otimes_A M \cong M$.

To show that our natural transformation is an isomorphism, we use induction on the number of times we need to extend a summand of the regular module to obtain P . We handle the base case as follows: the natural transformation is certainly an isomorphism when P is the regular module and so, as hom functors commute with finite direct sums, it is an isomorphism for all summands of the regular module. For our inductive step, suppose the lemma holds for P_1 and P_2 , and let $P = \mathrm{cone}(f)$ for some map $f : P_1 \rightarrow P_2$. Then, for $M \in A\text{-Mod}$, one can check that the map $\mathrm{Hom}_A(P, A) \otimes_A M \rightarrow \mathrm{Hom}_A(P, M)$ comes from the commutative diagram

$$\begin{array}{ccc} \mathrm{Hom}_A(P_1, A) \otimes_A M & \xleftarrow{\mathrm{Hom}(f, A) \otimes M} & \mathrm{Hom}_A(P_2, A) \otimes_A M \\ \downarrow & & \downarrow \\ \mathrm{Hom}_A(P_1, M) & \xleftarrow{\mathrm{Hom}(f, M)} & \mathrm{Hom}_A(P_2, M) \end{array}$$

as in the construction from the second half of Lemma 4.5, where the vertical maps come from the natural transformation described above. Therefore, as both vertical

maps are isomorphisms by induction, $\text{Hom}_A(P, A) \otimes_A M \rightarrow \text{Hom}_A(P, M)$ is an isomorphism. \square

Spherical twists. Our references are [Seidel and Thomas 2001; Rouquier and Zimmermann 2003; Grant 2013].

Let (A, d) be a dga and M be a left A -module with finite-dimensional total homology. Let $d \in \mathbb{Z}$. Following [Seidel and Thomas 2001], we say that M is d -spherical if

- M is a d -Calabi–Yau object, i.e., we have an isomorphism

$$\text{Hom}_{\text{D}_{\text{fd}}(A)}(M, N) \cong \text{Hom}_{\text{D}_{\text{fd}}(A)}(N, M[d])^*$$

which is functorial in N , and

- $\bigoplus_{i \in \mathbb{Z}} \text{Hom}_{\text{D}_{\text{fd}}(A)}(M, M[i])$ is isomorphic as a graded algebra to $\mathbb{F}[x]/\langle x^2 \rangle$, with x in degree d .

Associated to any spherical object M , we have a spherical twist functor $F_M : \text{D}_{\text{fd}}(A) \rightarrow \text{D}_{\text{fd}}(A)$ which is defined as follows. First, let $P = \mathbf{p} M$ be a cofibrant replacement of M . Then let X_M be the cone of the map of A - A -bimodules

$$P \otimes_{\mathbb{F}} \text{Hom}_A(P, A) \xrightarrow{\text{ev}} A,$$

where the nonzero map is the obvious evaluation map. As both $\text{Hom}_A(P, A)$ and A are cofibrant, X_M is cofibrant as a right A -module. Then we define the spherical twist at M by

$$F_M = X_M \otimes_A - : \text{D}_{\text{fd}}(A) \rightarrow \text{D}_{\text{fd}}(A).$$

The spherical twist is an autoequivalence of $\text{D}_{\text{fd}}(A)$ (so X_M is a tilting module).

Note that, by Lemmas 4.5 and 4.8, if M has finite projective dimension then

$$F_M(N) \cong P \otimes_{\mathbb{F}} \text{Hom}_A(P, N) \xrightarrow{\text{ev}} N.$$

We next need the fact that spherical twists are intertwined by derived equivalences. The following is a generalization of [Seidel and Thomas 2001, Lemma 2.11].

Proposition 4.9. *Let A, B be dgas. Let $T \in B\text{-Mod-}A$ be a tilting module and*

$$\Phi = T \otimes_A^L - : \text{D}_{\text{fd}}(A) \rightarrow \text{D}_{\text{fd}}(B)$$

be the associated derived equivalence. Let $M \in A\text{-Mod}$ have finite-dimensional total homology and suppose it is d -spherical, for some $d \in \mathbb{Z}$. Suppose that $\Phi(M) \in B\text{-Mod}$ has finite-dimensional total homology. Then $\Phi(M)$ is also d -spherical and we have an isomorphism of functors

$$\Phi \circ F_M \cong F_{\Phi(M)} \circ \Phi : \text{D}_{\text{fd}}(A) \xrightarrow{\sim} \text{D}_{\text{fd}}(B).$$

In particular, we have an isomorphism

$$F_{\Phi(M)} \cong \Phi \circ F_M \circ \Phi^{-1} : \mathbf{D}_{\text{fd}}(B) \xrightarrow{\sim} \mathbf{D}_{\text{fd}}(B)$$

where Φ^{-1} is the quasi-inverse functor of Φ .

Proof. As $\Phi : \mathbf{D}_{\text{fd}}(A) \rightarrow \mathbf{D}_{\text{fd}}(B)$ is a derived equivalence it has quasi-inverse $\Phi^{-1} : \mathbf{D}_{\text{fd}}(B) \rightarrow \mathbf{D}_{\text{fd}}(A)$, and so we have isomorphisms

$$\text{Hom}_{\mathbf{D}_{\text{fd}}(B)}(\Phi(M), \Phi(M)[i]) \cong \text{Hom}_{\mathbf{D}_{\text{fd}}(A)}(M, M[i])$$

and

$$\begin{aligned} \text{Hom}_{\mathbf{D}_{\text{fd}}(B)}(\Phi(M), N) &\cong \text{Hom}_{\mathbf{D}_{\text{fd}}(A)}(M, \Phi^{-1}(N)) \\ &\cong \text{Hom}_{\mathbf{D}_{\text{fd}}(A)}(\Phi^{-1}(N), M[d])^* \\ &\cong \text{Hom}_{\mathbf{D}_{\text{fd}}(B)}(N, \Phi(M)[d])^*, \end{aligned}$$

the second natural in $N \in \mathbf{D}_{\text{fd}}(B)$, using the facts that M is a d -Calabi–Yau object and the shift functor $[d]$ commutes with all triangulated functors. Thus $\Phi(M)$ is d -spherical.

By [Lemma 4.7](#) we may assume that T is cofibrant as a right B -module and that $\Phi = T \otimes_A -$. We want to show that

$$T \otimes_A X_M \otimes_A - \cong X_{\Phi(M)} \otimes_B T \otimes_A -,$$

so it is enough to check that we have an isomorphism

$$T \otimes_A X_M \cong X_{\Phi(M)} \otimes_B T$$

in $\mathbf{D}_{\text{fd}}(B \otimes_{\mathbb{F}} A^{\text{op}})$. To construct this isomorphism, we use the following extension of [Lemma 4.5](#), which follows from the triangulated Five Lemma: for any commutative diagram

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \sim \downarrow \varphi_M & & \sim \downarrow \varphi_N \\ M' & \xrightarrow{f'} & N' \end{array}$$

in $B\text{-Mod-}A$ where φ_M and φ_N are both quasi-isomorphisms, we have a quasi-isomorphism $\varphi_N \oplus \varphi_{M[1]} : \text{cone}(f) \rightarrow \text{cone}(f')$.

As above, write $P = \mathbf{p} M$. Then, by the first part of [Lemma 4.5](#), we just need to find two vertical maps which are quasi-isomorphisms and make the following diagram commute:

$$\begin{array}{ccc} T \otimes_A P \otimes_{\mathbb{F}} \text{Hom}_B(T \otimes_A P, B) \otimes_B T & \xrightarrow{\text{ev} \otimes 1_T} & B \otimes_B T \\ \sim \downarrow & & \sim \downarrow \\ T \otimes_A P \otimes_{\mathbb{F}} \text{Hom}_A(P, A) & \xrightarrow{1_T \otimes \text{ev}} & T \otimes_A A \end{array}$$

Our plan is to do this in stages: we will show that the vertical maps in the following diagram exist, and are quasi-isomorphisms, and that the diagram commutes.

$$\begin{array}{ccc}
 T \otimes_A P \otimes_F \text{Hom}_B(T \otimes_A P, B) \otimes_B T & \xrightarrow{\text{ev} \otimes 1_T} & B \otimes_B T \\
 \sim \downarrow & & = \downarrow \\
 T \otimes_A P \otimes_F \text{Hom}_B(T \otimes_A P, B \otimes_B T) & \xrightarrow{\text{ev}} & B \otimes_B T \\
 \sim \downarrow & & \sim \downarrow \\
 T \otimes_A P \otimes_F \text{Hom}_B(T \otimes_A P, T \otimes_A A) & \xrightarrow{\text{ev}} & T \otimes_A A \\
 \sim \downarrow & & = \downarrow \\
 T \otimes_A P \otimes_F \text{Hom}_A(P, A) & \xrightarrow{1_T \otimes \text{ev}} & T \otimes_A A
 \end{array}$$

Let us show that the first square commutes. We introduce some temporary notation for the rest of this proof. Let F and G denote the functors $F = T \otimes_A P \otimes_F -$ and $G = \text{Hom}_B(T \otimes_A P, -)$, so F is left adjoint to G , and let H denote the functor $- \otimes_B T$. Then we have unit and counit natural transformations $\varepsilon : FG \rightarrow 1$ and $\eta : 1 \rightarrow GF$, and a natural isomorphism $\zeta : HF \xrightarrow{\sim} FH$ coming from the associativity isomorphism for tensor products. We first need to define a map

$$\begin{aligned}
 HFGB &= T \otimes_A P \otimes_F \text{Hom}_B(T \otimes_A P, B) \otimes_B T \rightarrow T \otimes_A P \otimes_F \text{Hom}_B(T \otimes_A P, B \otimes_B T) \\
 &= FGHB.
 \end{aligned}$$

We define this as the composite

$$HFGB \xrightarrow{\zeta_{GB}} FHGB \xrightarrow{F\eta_{HGB}} FGFHGB \xrightarrow{FG\zeta^{-1}GB} FGHFGB \xrightarrow{FGH\varepsilon_B} FGHB.$$

One checks that this is an isomorphism using the same argument as in [Lemma 4.8](#). To see that the diagram commutes, we break it up into smaller diagrams as follows:

$$\begin{array}{ccc}
 HFGB & \xrightarrow{H\varepsilon_B} & HB \\
 \zeta_{GB} \downarrow \sim & & \uparrow H\varepsilon_B \\
 FHGB & \xrightarrow{1} & FHGB & \xrightarrow{\sim} & HFGB \\
 F\eta_{HGB} \downarrow & \nearrow \varepsilon_{FHGB} & & \searrow \zeta^{-1}GB & \uparrow \varepsilon_{HB} \\
 FGFHGB & & & & HFGB \\
 FG\zeta^{-1}GB \downarrow \sim & \nearrow \varepsilon_{HFGB} & & & \uparrow \varepsilon_{HB} \\
 FGHFGB & \xrightarrow{FGH\varepsilon_B} & FGHB & & \uparrow \varepsilon_{HB}
 \end{array}$$

Now we see that both squares commute by the naturality of ε , the triangle commutes by the triangle identity $\varepsilon_F \circ F\eta = 1_F$, and the pentagon commutes because the isomorphisms are defined by ζ and its inverse.

We use the obvious composite isomorphism $B \otimes_B T \xrightarrow{\sim} T \xrightarrow{\sim} T \otimes_A A$ to define the second square. This commutes because the evaluation map is a counit, and therefore a natural transformation.

To show that the third square commutes, we introduce some more notation. Let F' and G' denote the functors $F' = P \otimes_{\mathbb{F}} -$ and $G' = \text{Hom}_A(P, -)$, so F' is left adjoint to G' , and let H' and I' denote the functors $H' = T \otimes_A -$ and $I' = \text{Hom}_B(T, -)$, so H' is left adjoint (in fact, quasi-inverse) to I' . We denote the counit and unit maps of the first adjunction by $\varepsilon' : F'G' \rightarrow 1$ and $\eta' : 1 \rightarrow G'F'$, and of the second adjunction by $\varepsilon'' : H'I' \rightarrow 1$ and $\eta'' : 1 \rightarrow I'H'$. Note that, because H' induces an equivalence of derived categories, ε'' and η'' give quasi-isomorphisms when applied to any object.

Using the associativity isomorphism for tensor products we have a natural isomorphism of functors $F \cong H'F'$, and by the uniqueness of right adjoints (or by using the tensor-hom adjunctions directly) this gives another natural isomorphism $G \cong G'I'$.

We now redraw our final square, breaking it up into smaller diagrams:

$$\begin{array}{ccc}
 FGH'A & \xrightarrow{\varepsilon'_{H'A}} & H'A \\
 \downarrow \sim & & \nearrow \varepsilon''_{H'A} \\
 H'F'G'I'H'A & \xrightarrow{H'\varepsilon'_{I'H'A}} & H'I'H'A \\
 \uparrow H'F'G'\eta''_A & & \nwarrow H'\eta''_A \\
 H'F'G'A & \xrightarrow{H'\varepsilon'_A} & H'A \\
 & & \uparrow 1
 \end{array}$$

Here, the top square commutes by definition of the isomorphisms $F \cong H'F'$ and $G \cong G'I'$, the triangle commutes by the triangle identity $\varepsilon''H' \circ H'\eta'' = 1_{H'}$, and the bottom square commutes by the naturality of ε' . \square

We now describe the braid relations for spherical twists, as in Propositions 2.12 and 2.13 of [Seidel and Thomas 2001]; see also [Rouquier and Zimmermann 2003; Grant 2015].

Proposition 4.10. *Suppose that M and N are spherical objects of $\text{D}_{\text{fd}}(A)$ and let*

$$(M, N) = \dim_{\mathbb{F}} \bigoplus_{n \in \mathbb{Z}} \text{Hom}_{\text{D}_{\text{fd}}(A)}(M, N[n]).$$

Let $F_M, F_N : \text{D}_{\text{fd}}(A) \xrightarrow{\sim} \text{D}_{\text{fd}}(A)$ be the associated spherical twists.

- If $(M, N) = 0$ then $F_M \circ F_N \cong F_N \circ F_M$.
- If $(M, N) = 1$ then $F_M \circ F_N \circ F_M \cong F_N \circ F_M \circ F_N$.

Ginzburg dg-algebras. There is a well-known method to associate a differential graded algebra to a quiver with potential [Ginzburg 2007, Section 5; Keller and Yang 2011, Section 2.6].

Let (Q, W) be a quiver with potential. Construct a new quiver \bar{Q} by adding arrows to Q : for each arrow $a : i \rightarrow j$ in Q we add a new arrow $a^* : j \rightarrow i$, and for each vertex i in Q we add a new arrow $t_i : i \rightarrow i$. We view \bar{Q} as a graded quiver with the arrows of Q in degree 0, the arrows a^* in degree -1 , and the arrows t_i in degree -2 . This induces a grading on the path algebra $\mathbb{F}\bar{Q}$ of \bar{Q} such that the degree 0 part $(\mathbb{F}\bar{Q})_0$ is just the path algebra of $\mathbb{F}Q$ of Q . Let \bar{J} denote the ideal of $\mathbb{F}\bar{Q}$ generated by the arrows of \bar{Q} , and let $\widehat{\mathbb{F}\bar{Q}}$ denote the completion of the graded algebra $\mathbb{F}\bar{Q}$ with respect to \bar{J} , as on page 96.

We define a differential d on $\widehat{\mathbb{F}\bar{Q}}$ by requiring that d be zero on each idempotent e_i associated to a vertex i of Q , specifying how d acts on arrows of \bar{Q} , and then extending to the rest of $\widehat{\mathbb{F}\bar{Q}}$ using the Leibniz rule and continuity. For degree reasons, we must have $d(a) = 0$ for each arrow a of Q . For arrows a^* , we set $d(a^*) = \partial_a W$, where ∂_a denotes the cyclic derivative, and for arrows t_i we set $d(t_i) = e_i(\sum aa^* - a^*a)e_i$, where we sum over all arrows a of Q . Then $\Gamma_{Q,W} = (\widehat{\mathbb{F}\bar{Q}}, d)$ is called the Ginzburg dga of (Q, W) .

Note that if (Q_1, W_1) and (Q_2, W_2) are right equivalent, then we have an isomorphism of dgas $\Gamma_{Q_1, W_1} \cong \Gamma_{Q_2, W_2}$ [Keller and Yang 2011, Lemma 2.9]. Hence, if we are working with quivers with potential of mutation type A or D , by Proposition 4.4 we only need to consider the Ginzburg dgas Γ_{Q, W_Q} , and so can denote them Γ_Q .

Keller and Yang showed that QP-mutation lifts to equivalences of derived categories of Ginzburg dgas:

Theorem 4.11 [Keller and Yang 2011, Theorem 3.2]. *Suppose that (Q, W) is a QP and that $(Q', W') = \mu_k(Q, W)$ for some $k \in I$. There is a tilting complex T which gives an equivalence of triangulated categories*

$$\mu_k = \text{Hom}_{\Gamma_{Q', W'}}(T, -) : \text{D}(\Gamma_{Q', W'}) \rightarrow \text{D}(\Gamma_{Q, W}),$$

and it restricts to an equivalence of triangulated categories,

$$\mu_k = \text{Hom}_{\Gamma_{Q', W'}}(T, -) : \text{D}_{\text{fd}}(\Gamma_{Q', W'}) \rightarrow \text{D}_{\text{fd}}(\Gamma_{Q, W}).$$

Recall that, for a dga A , the finite-dimensional derived category $\text{D}_{\text{fd}}(A)$ is d -Calabi–Yau if there exists a bifunctorial isomorphism,

$$\text{Hom}_{\text{D}_{\text{fd}}(A)}(M, N) \cong \text{Hom}_{\text{D}_{\text{fd}}(A)}(N, M[d])^*,$$

where $(-)^*$ denotes the k -linear dual. We will need the following important result of Keller and Van den Bergh:

Theorem 4.12 [Keller 2011, Theorem 6.3 and Theorem A.12]. *The category $\text{D}_{\text{fd}}(\Gamma_{Q, W})$ is 3-Calabi–Yau.*

Let (Q, W) be a QP and $\Gamma = \Gamma_{Q, W}$. Associated to each vertex i of Q , we have the 1-dimensional simple left Γ -module, which we denote S_i . Keller and Yang

[2011, Section 2.14] explained how to construct the cofibrant replacement of S_i : as long as we remember the differential, we can proceed as if the Ginzburg dga were an ordinary hereditary algebra, and the underlying $u(\Gamma)$ -module of $\mathbf{p} S_i$ is the direct sum of one copy of the projective P_i and one copy of the shifted projective $P_j[1]$ for each arrow $j \rightarrow i$ in Q . Using this, they showed:

Lemma 4.13 [Keller and Yang 2011, Lemma 2.15]. *Let $i, j \in I$ and $n \in \mathbb{Z}$ and $\Gamma = \Gamma_{Q,W}$. Then $\mathrm{Hom}_{\mathrm{D}_{\mathrm{fd}}(\Gamma)}(S_i, S_j[n]) = 0$ if $n \neq 0, 1, 2, 3$, and*

$$\dim_{\mathbb{F}} \mathrm{Hom}_{\mathrm{D}_{\mathrm{fd}}(\Gamma)}(S_i, S_j[n]) = \begin{cases} \delta_{ij} & \text{if } n = 0, \\ \#\{\text{arrows } i \rightarrow j \text{ in } Q\} & \text{if } n = 1, \\ \#\{\text{arrows } j \rightarrow i \text{ in } Q\} & \text{if } n = 2, \\ \delta_{ij} & \text{if } n = 3, \end{cases}$$

where δ_{ij} is the Kronecker delta.

Relations between functors. By Theorem 4.12, every object of $\mathrm{D}_{\mathrm{fd}}(\Gamma_{Q,W})$ is a 3-Calabi–Yau object. By Lemma 4.13,

$$\bigoplus_{j \in \mathbb{Z}} \mathrm{Hom}_{\mathrm{D}_{\mathrm{fd}}(\Gamma_{Q,W})}(S_i, S_i[j]) \cong \mathbb{F}[x]/\langle x^2 \rangle$$

with x in degree 3. Hence S_i is 3-spherical, and we have a spherical twist F_{S_i} associated to S_i . We will sometimes write F_i instead of F_{S_i} .

Let k be a vertex of Q , and write $(Q', W') = \mu_k(Q, W)$. Then write $\Gamma = \Gamma_{Q,W}$ and $\Gamma' = \Gamma_{Q',W'}$ for the associated Ginzburg dgas. Write T_i for the left Γ' -module associated to the vertex i of Q' and G_i for the associated autoequivalence F_{T_i} of $\mathrm{D}_{\mathrm{fd}}(\Gamma')$. In this section we will study how the spherical twists $F_i : \mathrm{D}_{\mathrm{fd}}(\Gamma) \xrightarrow{\sim} \mathrm{D}_{\mathrm{fd}}(\Gamma)$ interact with the mutation functors $\mu_k : \mathrm{D}_{\mathrm{fd}}(\Gamma') \xrightarrow{\sim} \mathrm{D}_{\mathrm{fd}}(\Gamma)$. Our key tools will be Proposition 4.9 and the results on the images of the simple modules under the mutation functors [Keller and Yang 2011, Lemma 3.12(a)], which we will describe below.

If A is a dga and $M, N \in \mathrm{D}_{\mathrm{fd}}(A)$, we have a natural map,

$$M \otimes_{\mathbb{F}} \mathrm{Hom}_{\mathrm{D}_{\mathrm{fd}}(A)}(M, N) \rightarrow N,$$

in $\mathrm{D}_{\mathrm{fd}}(A)$ given by evaluation. For any graded vector space V , we have biadjoint functors $- \otimes_F V$ and $- \otimes_F V^*$, and these respect the left A -module structure, so we also obtain a natural map,

$$M \rightarrow N \otimes_{\mathbb{F}} \mathrm{Hom}_{\mathrm{D}_{\mathrm{fd}}(A)}(M, N)^*,$$

in $\mathrm{D}_{\mathrm{fd}}(A)$. Now let $L, N \in A\text{-Mod}$. The *universal extension of N by L* is the cone of the natural map

$$N[-1] \rightarrow L \otimes_{\mathbb{F}} \mathrm{Hom}_{\mathrm{D}_{\mathrm{fd}}(A)}(N[-1], L)^*$$

and the *universal coextension of L by N* is the cone of the natural map

$$N[-1] \otimes_{\mathbb{F}} \text{Hom}_{\text{D}_{\text{fd}}(A)}(N[-1], L) \rightarrow L.$$

The following result is due to Keller and Yang:

Lemma 4.14 [Keller and Yang 2011, Lemma 3.12(a)]. *We have $\mu_k(T_k) \cong S_k[1]$ and $\mu_k(T_i)$ is the universal extension of S_i by S_k , for $i \neq k$.*

The following result should be compared to Definition 2.8.

Proposition 4.15. *If Q has no double arrows then we have a natural isomorphism of functors*

$$F_{\mu_k^{-1}(S_i)} \cong \begin{cases} G_k G_i G_k^{-1} & \text{if } i \rightarrow k \text{ in } Q, \\ G_i & \text{otherwise.} \end{cases}$$

Proof. We first use Lemma 4.14 to calculate the images of the simple Γ' -modules under the inverse mutation functor μ^{-1} , where $\mu = \mu_k$. We know that $\mu(T_k) \cong S_k[1]$, so $\mu^{-1}(S_k) \cong T_k[-1]$. By assumption, there is at most one arrow between any two vertices in Q . If $i \neq k$ and there is no arrow $i \rightarrow k$ in Q then, by Lemma 4.14, $\text{Hom}_{\text{D}_{\text{fd}}(\Gamma)}(S_i[-1], S_k) = 0$ and so $\mu(T_i) \cong \text{cone}(S_i[-1] \rightarrow 0) \cong S_i$, so $\mu^{-1}(S_i) \cong T_i$.

If $i \neq k$ and there is an arrow $i \rightarrow k$ in Q then $\text{Hom}_{\text{D}_{\text{fd}}(\Gamma)}(S_i[-1], S_k)$ is 1-dimensional and so $\mu(T_i) \cong \text{cone}(S_i[-1] \rightarrow S_k)$, with the nonzero map determined up to a scalar. We can then use Lemma 4.5 to calculate $\mu(\text{cone}(T_k[-1] \rightarrow T_i))$: this is $\text{cone}(\mu(T_k)[-1] \rightarrow \mu(T_i))$ where, as μ is an equivalence, the map must again be nonzero and determined up to scalar. We know that $\mu(T_k)[-1] \cong S_k$ and $\mu(T_i)$ is $S_i \oplus S_k$ with appropriate differential. One can check that the injection $S_k \hookrightarrow S_i \oplus S_k$ respects the differentials, and so this must be our nonzero map. This is quasi-isomorphic to the map $0 \rightarrow S_i$, and so $\mu(\text{cone}(T_k[-1] \rightarrow T_i)) \cong S_i$ and hence $\mu^{-1}(S_i) \cong \text{cone}(T_k[-1] \rightarrow T_i)$. Note that this is the universal coextension of T_i by T_k .

Now we check that the formula holds. If $i = k$ then $F_{\mu^{-1}(S_i)} = F_{T_i[1]}$, and as the shift functor on $\text{D}_{\text{fd}}(\Gamma')$ is naturally isomorphic to $\Gamma'[1] \otimes_{\Gamma'} -$ we see that, by Proposition 4.9, $F_{T_i[1]} \cong [1] \circ G_i \circ [-1] \cong G_i$. If $i \neq k$ and there is no arrow $i \rightarrow k$ in Q then $\mu^{-1}(S_i) \cong T_i$ so $F_{\mu^{-1}(T_i)} = G_i$.

Finally, suppose $i \neq k$ and there is an arrow $i \rightarrow k$ in Q . As mutation at k reverses all arrows incident with k , and can never change the number of arrows incident with k , there must be exactly one arrow $k \rightarrow i$ in Q' . We first calculate $G_k(T_i)$: this is

$$\text{cone}(\mathbf{p} T_k \otimes_{\mathbb{F}} \text{Hom}_{\Gamma'}(\mathbf{p} T_k, T_i) \rightarrow T_i).$$

As $\text{Hom}_{\Gamma'}(\mathbf{p} T_k, T_i)$ is a differential graded \mathbb{F} -module, it is quasi-isomorphic to its homology, which is the direct sum $\bigoplus \text{Hom}_{K(\Gamma')}(T_k, T_i[n])$ with

$$\text{Hom}_{K(\Gamma')}(T_k, T_i[n]) \cong \text{Hom}_{\text{D}_{\text{fd}}(\Gamma')}(T_k, T_i[n])$$

in degree n . So by [Lemma 4.14](#) the homology is only nonzero in degree 1, where it is 1-dimensional, and thus

$$G_k(T_i) \cong \text{cone}(T_k \otimes_{\mathbb{F}} \mathbb{F}[-1] \rightarrow T_i).$$

So we see that $\mu^{-1}(S_i) \cong G_k(T_i)$ and therefore, using [Proposition 4.9](#) again, $F_{\mu^{-1}(S_i)} \cong G_k G_i G_k^{-1}$. \square

We are now able to show that our braid groups B_Q act via spherical twists on the category $\text{D}_{\text{fd}}(\Gamma)$.

Theorem 4.16. *Let (Q, W) be a mutation-Dynkin quiver with potential of type ADE. Then we have a group homomorphism*

$$B_Q \rightarrow \text{Aut D}_{\text{fd}}(\Gamma_{Q,W}), \quad s_i \mapsto F_i$$

sending the group generator associated to the vertex $i \in I$ to the spherical twist at the simple $\Gamma_{Q,W}$ -module S_i .

Proof. As (Q, W) is mutation-Dynkin, it is obtained by mutating a quiver with potential $(Q'', 0)$ finitely many times, where Q'' is a Dynkin quiver. Then we have a group homomorphism $B_{Q''} \rightarrow \text{Aut D}_{\text{fd}}(\Gamma_{Q'',0})$ by [Remark 2.3](#), [Proposition 4.10](#), and [Lemma 4.13](#). This gives the base case of an inductive argument, so we need to show that if the spherical twists F_i on $\Gamma = \Gamma_{Q,W}$ satisfy the relations of B_Q for a mutation-Dynkin quiver with potential (Q, W) then the spherical twists G_i on $\Gamma' = \Gamma_{Q',W'}$ satisfy the relations of $B_{Q'}$.

Assume the functors $F_i : \text{D}_{\text{fd}}(\Gamma) \rightarrow \text{D}_{\text{fd}}(\Gamma)$ satisfy the relations of B_Q and let $\mu = \mu_k : \text{D}_{\text{fd}}(\Gamma') \rightarrow \text{D}_{\text{fd}}(\Gamma)$ be the Keller–Yang derived equivalence. Then the functors $\mu^{-1} \circ F_i \circ \mu : \text{D}_{\text{fd}}(\Gamma') \rightarrow \text{D}_{\text{fd}}(\Gamma')$ also satisfy the relations of B_Q . By [Proposition 4.9](#) we have

$$\mu^{-1} \circ F_i \circ \mu \cong F_{\mu^{-1}(S_i)},$$

i.e., the following diagram commutes:

$$\begin{array}{ccc} \text{D}_{\text{fd}}(\Gamma') & \xrightarrow{\mu} & \text{D}_{\text{fd}}(\Gamma) \\ \downarrow F_{\mu^{-1}(S_i)} & & \downarrow F_{S_i} \\ \text{D}_{\text{fd}}(\Gamma') & \xrightarrow{\mu} & \text{D}_{\text{fd}}(\Gamma) \end{array}$$

So we have a group homomorphism $\rho : B_Q \xrightarrow{\rho} \text{Aut D}_{\text{fd}}(\Gamma')$ sending s_i to $F_{\mu^{-1}(S_i)}$. By [Proposition 2.1](#), Q has no double arrows, so we can use [Proposition 4.15](#) to

write ρ as

$$B_Q \xrightarrow{\rho} \text{Aut } D_{\text{fd}}(\Gamma'),$$

$$s_i \mapsto \begin{cases} G_k G_i G_k^{-1} & \text{if } i \rightarrow k \text{ in } Q; \\ G_i & \text{otherwise.} \end{cases}$$

Precomposing this with the group isomorphism $\varphi_k^{-1} : B'_Q \xrightarrow{\sim} B_Q$ of [Remark 2.11](#), we obtain the group homomorphism

$$B_{Q'} \xrightarrow{\varphi_k^{-1}} B_Q \xrightarrow{\rho} \text{Aut } D_{\text{fd}}(\Gamma')$$

$$t_i \mapsto \begin{cases} s_k^{-1} s_i s_k & \text{if } i \rightarrow k \text{ in } Q, \\ s_i & \text{otherwise} \end{cases} \mapsto \begin{cases} G_k^{-1} G_k G_i G_k^{-1} G_k & \text{if } i \rightarrow k \text{ in } Q, \\ G_i & \text{otherwise} \end{cases} \cong G_i$$

as required. □

Remark 4.17. Known results on the faithfulness of braid group actions can be transferred to our setting. Suppose Q'' is an orientation of an *ADE* graph and the usual action $B_{Q''} \rightarrow \text{Aut } D_{\text{fd}}(\Gamma_{Q'',0})$ is faithful. From the proof of [Theorem 4.16](#) we see that our actions of B_Q where Q is of mutation type *ADE* are just built by precomposing group isomorphisms with the action of $B_{Q''}$, and so these are also faithful under this assumption.

It was shown by Seidel and Thomas [[2001](#), Theorem 2.18], building on work of Khovanov and Seidel [[2002](#)], that given a collection of d -spherical objects, with $d \geq 2$, in a type A_n -configuration the action of the braid group by spherical twists is faithful. Thus the actions of [Theorem 4.16](#) are faithful in mutation type *A*. The faithfulness result of Seidel and Thomas was extended to all collections of 2-spherical objects in type *ADE* configurations by Brav and Thomas [[2011](#)], using the Garside structure of the braid monoid, but it is not immediately clear how to generalize their argument to the 3-Calabi–Yau situation.

Remark 4.18. Although we have shown that our braid groups of mutation-Dynkin quivers can be realized categorically, this is not a categorification of our earlier results because we cannot decategorify (see, for example, [[Baez and Dolan 1998](#)]): we cannot recover [Theorem 2.10](#) from [Theorem 4.16](#) because we use [Theorem 2.10](#) to prove [Theorem 4.16](#). The problem is that, for an arbitrary mutation-Dynkin quiver with potential (Q, W) , we do not in advance know the relations satisfied by the spherical twist functors F_i . This question will be addressed in a forthcoming paper.

Remark 4.19. The arguments of [[Grant 2015](#)] generalize to show that, if vertices i and j of Q are joined by an arrow, then $F_i F_j F_i \cong F_j F_i F_j$ can be realized as a single periodic twist. Similarly, one can show that if $i \rightarrow j \rightarrow k \rightarrow i$ is a 3-cycle in

Q then

$$F_1 F_2 F_3 F_1 \cong F_2 F_3 F_1 F_2 \cong F_3 F_1 F_2 F_3$$

can be realized as a single periodic twist. This exhausts the possibilities in type A ; we will study type D further in the future.

Acknowledgements

Marsh would like to thank K. Baur and A. King for useful discussions concerning orbifold interpretations of cluster algebras. Both authors would like to thank the referee, and I. Webster, for helpful comments on an earlier version of the paper. Some of the work for this paper was done while Marsh was Guest Professor in 2014 at the Department of Mathematical Sciences, Norwegian University of Science and Technology, Trondheim, N-7491, Norway.

References

- [Allcock 2002] D. Allcock, “Braid pictures for Artin groups”, *Trans. Amer. Math. Soc.* **354**:9 (2002), 3455–3474. [MR](#) [Zbl](#)
- [Amiot 2009] C. Amiot, “Cluster categories for algebras of global dimension 2 and quivers with potential”, *Ann. Inst. Fourier (Grenoble)* **59**:6 (2009), 2525–2590. [MR](#) [Zbl](#)
- [Artin 1925] E. Artin, “Theorie der Zöpfe”, *Abh. Math. Sem. Univ. Hamburg* **4**:1 (1925), 47–72. [MR](#) [JFM](#)
- [Baez and Dolan 1998] J. C. Baez and J. Dolan, “Categorification”, pp. 1–36 in *Higher category theory* (Evanston, IL, 1997), edited by E. Getzler and M. Kapranov, *Contemp. Math.* **230**, American Mathematical Society, Providence, RI, 1998. [MR](#) [Zbl](#)
- [Barot and Marsh 2015] M. Barot and B. R. Marsh, “Reflection group presentations arising from cluster algebras”, *Trans. Amer. Math. Soc.* **367**:3 (2015), 1945–1967. [MR](#) [Zbl](#)
- [Brav and Thomas 2011] C. Brav and H. Thomas, “Braid groups and Kleinian singularities”, *Math. Ann.* **351**:4 (2011), 1005–1017. [MR](#) [Zbl](#)
- [Buan et al. 2006] A. B. Buan, B. R. Marsh, M. Reineke, I. Reiten, and G. Todorov, “Tilting theory and cluster combinatorics”, *Adv. Math.* **204**:2 (2006), 572–618. [MR](#) [Zbl](#)
- [Caldero et al. 2006] P. Caldero, F. Chapoton, and R. Schiffler, “Quivers with relations arising from clusters (A_n case)”, *Trans. Amer. Math. Soc.* **358**:3 (2006), 1347–1364. [MR](#) [Zbl](#)
- [Derksen et al. 2008] H. Derksen, J. Weyman, and A. Zelevinsky, “Quivers with potentials and their representations, I: Mutations”, *Selecta Math. (N.S.)* **14**:1 (2008), 59–119. [MR](#) [Zbl](#)
- [Felikson et al. 2012] A. Felikson, M. Shapiro, and P. Tumarkin, “Cluster algebras and triangulated orbifolds”, *Adv. Math.* **231**:5 (2012), 2953–3002. [MR](#) [Zbl](#)
- [Fomin and Zelevinsky 2002] S. Fomin and A. Zelevinsky, “Cluster algebras, I: Foundations”, *J. Amer. Math. Soc.* **15**:2 (2002), 497–529. [MR](#) [Zbl](#)
- [Fomin and Zelevinsky 2003] S. Fomin and A. Zelevinsky, “Cluster algebras, II: Finite type classification”, *Invent. Math.* **154**:1 (2003), 63–121. [MR](#) [Zbl](#)
- [Fomin et al. 2008] S. Fomin, M. Shapiro, and D. Thurston, “Cluster algebras and triangulated surfaces, I: Cluster complexes”, *Acta Math.* **201**:1 (2008), 83–146. [MR](#) [Zbl](#)

- [Fox and Neuwirth 1962] R. Fox and L. Neuwirth, “The braid groups”, *Math. Scand.* **10** (1962), 119–126. [MR](#) [Zbl](#)
- [Ginzburg 2007] V. Ginzburg, “Calabi–Yau algebras”, preprint, 2007. [arXiv:math/0612139v3](#)
- [Grant 2013] J. Grant, “Derived autoequivalences from periodic algebras”, *Proc. Lond. Math. Soc.* (3) **106**:2 (2013), 375–409. [MR](#) [Zbl](#)
- [Grant 2015] J. Grant, “Lifts of longest elements to braid groups acting on derived categories”, *Trans. Amer. Math. Soc.* **367**:3 (2015), 1631–1669. [MR](#) [Zbl](#)
- [Haley et al. 2014] J. Haley, D. Hemminger, A. Landesman, and H. Peck, “Artin group presentations arising from cluster algebras”, preprint, 2014. [arXiv](#)
- [Humphreys 1990] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics **29**, Cambridge Univ. Press, 1990. [MR](#) [Zbl](#)
- [Kassel and Turaev 2008] C. Kassel and V. Turaev, *Braid groups*, Graduate Texts in Mathematics **247**, Springer, New York, 2008. [MR](#) [Zbl](#)
- [Keller 1994] B. Keller, “Deriving DG categories”, *Ann. Sci. École Norm. Sup.* (4) **27**:1 (1994), 63–102. [MR](#) [Zbl](#)
- [Keller 2006] B. Keller, “On differential graded categories”, pp. 151–190 in *International Congress of Mathematicians*, vol. II, edited by M. Sanz-Solé et al., Eur. Math. Soc., Zürich, 2006. [MR](#) [Zbl](#)
- [Keller 2011] B. Keller, “Deformed Calabi–Yau completions”, *J. Reine Angew. Math.* **654** (2011), 125–180. With an appendix by Michel Van den Bergh. [MR](#) [Zbl](#)
- [Keller and Yang 2011] B. Keller and D. Yang, “Derived equivalences from mutations of quivers with potential”, *Adv. Math.* **226**:3 (2011), 2118–2168. [MR](#) [Zbl](#)
- [Khovanov and Seidel 2002] M. Khovanov and P. Seidel, “Quivers, Floer cohomology, and braid group actions”, *J. Amer. Math. Soc.* **15**:1 (2002), 203–271. [MR](#) [Zbl](#)
- [Labardini-Fragoso 2009] D. Labardini-Fragoso, “Quivers with potentials associated to triangulated surfaces”, *Proc. Lond. Math. Soc.* (3) **98**:3 (2009), 797–839. [MR](#) [Zbl](#)
- [Labardini-Fragoso 2016] D. Labardini-Fragoso, “Quivers with potentials associated to triangulated surfaces, IV: Removing boundary assumptions”, *Selecta Math. (N.S.)* **22**:1 (2016), 145–189. [MR](#) [Zbl](#)
- [Nagao 2010] K. Nagao, “Triangulated surface, mapping class group and Donaldson–Thomas theory”, lecture notes, Graduate School of Mathematics, Nagoya University, 2010, available at <http://www.ms.u-tokyo.ac.jp/~kazushi/kinosaki2010/nagao.pdf>.
- [Qiu 2016] Y. Qiu, “Decorated marked surfaces: spherical twists versus braid twists”, *Math. Ann.* **365**:1–2 (2016), 595–633. [MR](#) [Zbl](#)
- [Rouquier and Zimmermann 2003] R. Rouquier and A. Zimmermann, “Picard groups for derived module categories”, *Proc. London Math. Soc.* (3) **87**:1 (2003), 197–225. [MR](#) [Zbl](#)
- [Schiffler 2008] R. Schiffler, “A geometric model for cluster categories of type D_n ”, *J. Algebraic Combin.* **27**:1 (2008), 1–21. [MR](#) [Zbl](#)
- [Seidel and Thomas 2001] P. Seidel and R. Thomas, “Braid group actions on derived categories of coherent sheaves”, *Duke Math. J.* **108**:1 (2001), 37–108. [MR](#) [Zbl](#)
- [Sergiescu 1993] V. Sergiescu, “Graphes planaires et présentations des groupes de tresses”, *Math. Z.* **214**:3 (1993), 477–490. [MR](#)
- [Thurston 2012] D. Thurston, “The geometry and algebra of curves on surfaces”, course notes, Department of Mathematics, University of California, Berkeley, 2012, available at <http://tinyurl.com/qiaochu>. Notes taken by Qiaochu Yuan.

Received March 28, 2016. Revised January 13, 2017.

JOSEPH GRANT
SCHOOL OF MATHEMATICS
UNIVERSITY OF EAST ANGLIA
NORWICH RESEARCH PARK
NORWICH
NR4 7TJ
UNITED KINGDOM
j.grant@uea.ac.uk

BETHANY ROSE MARSH
SCHOOL OF MATHEMATICS
UNIVERSITY OF LEEDS
LEEDS
LS2 9JT
UNITED KINGDOM
marsh@maths.leeds.ac.uk

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 290 No. 1 September 2017

The Vietoris–Rips complexes of a circle	1
MICHAŁ ADAMASZEK and HENRY ADAMS	
A tale of two Liouville closures	41
ALLEN GEHRET	
Braid groups and quiver mutation	77
JOSEPH GRANT and BETHANY R. MARSH	
Paley–Wiener theorem of the spectral projection for symmetric graphs	117
SHIN KOIZUMI	
Fundamental domains of arithmetic quotients of reductive groups over number fields	139
LEE TIM WENG	
Growth and distortion theorems for slice monogenic functions	169
GUANGBIN REN and XIEPING WANG	
Remarks on metaplectic tensor products for covers of GL_r	199
SHUICHIRO TAKEDA	
On relative rational chain connectedness of threefolds with anti-big canonical divisors in positive characteristics	231
YUAN WANG	
An orthogonality relation for spherical characters of supercuspidal representations	247
CHONG ZHANG	