

*Pacific  
Journal of  
Mathematics*

**A PALEY-WIENER THEOREM  
FOR THE SPECTRAL PROJECTION  
OF SYMMETRIC GRAPHS**

SHIN KOIZUMI

Volume 290 No. 1

September 2017



# A PALEY–WIENER THEOREM FOR THE SPECTRAL PROJECTION OF SYMMETRIC GRAPHS

SHIN KOIZUMI

**We prove a Paley–Wiener theorem for the spectral projection of symmetric graphs and, as a corollary, derive a Paley–Wiener theorem for the Helgason–Fourier transform. The proof is based on contour integration arguments similar to those used to prove the Paley–Wiener theorem for Euclidean spaces and symmetric spaces.**

## 1. Introduction

The theory of representations of free groups has been studied by many authors in analogy with the semisimple theory. This arises from the realization of a free group as a homogeneous tree and relies upon the use of the Poisson boundary and spherical function. Mantero and Zappa [1983] characterized the image of the Poisson transform of free groups and studied the uniform boundedness of the spherical representation. In [Cowling et al. 1998], Cowling, Meda and Setti studied the images of the Abel transform for various function spaces on homogeneous trees. Cowling and Setti [1999] gave the characterizations of the images of the spaces of compactly supported functions and rapidly decreasing functions.

The concept of tree has been extended in several aspects. For instance, Iozzi and Picardello [1983a; 1983b] extended the context of tree to symmetric graphs and gave an explicit expression of the spherical function. Later, the Plancherel measure on symmetric graphs was explicitly computed in [Kuhn and Soardi 1983; Faraut and Picardello 1984]. Recently Eddine [2013; 2015] investigated the characterization of the Abel transform for symmetric graphs and, as an application, solved the shifted wave equations on it.

In [Koizumi 2013], we studied the spectral projection on homogeneous trees and proved the Paley–Wiener theorem of the spectral projection, which is an analogue of that given by Bray [1996]. In this paper, we shall extend the works in [Koizumi 2013] to the case of symmetric graphs. Unlike the works of Cowling and Setti

---

*MSC2010:* primary 43A80; secondary 43A85.

*Keywords:* spectral projection, symmetric graph, Paley–Wiener theorem, generalized spherical function.

[1999], our proof is based on contour integration arguments, which are usually used to prove the Paley–Wiener theorem for the cases of the Euclidean spaces and the symmetric spaces [Johnson 1979; Campoli 1980].

A brief outline of this paper is as follows: Section 2 is devoted to the overview of the notation of symmetric graphs. In Section 3, we concretely write down the expressions of the Poisson transform on symmetric graphs. In Section 4, we construct the intertwining operators between the spherical representations and give the explicit expressions of the intertwining operators. In Section 5, we study the properties of the spectral projection for symmetric graphs. Finally in Section 6, we show the Paley–Wiener theorem of the spectral projection and prove the Paley–Wiener theorem of the Helgason–Fourier transform.

## 2. Notation and preliminaries

The standard symbols  $\mathbb{Z}$ ,  $\mathbb{R}$  and  $\mathbb{C}$  are used for the integers, the real numbers and the complex numbers, respectively. Let us set  $\mathbb{Z}_k = \mathbb{Z}/k\mathbb{Z}$ . Throughout this paper, the imaginary unit is denoted as  $i$ . If  $x \in \mathbb{C}$ ,  $\Re x$  and  $\Im x$  denote its real part and its imaginary part, respectively.

A graph  $\mathfrak{X}$  is symmetric of type  $k \geq 2$  and order  $r \geq 2$  if every vertex  $v$  belongs exactly to  $r$  polygons with  $k$  sides each with no sides and no vertex in common except  $v$ , and if every nontrivial loop in  $\mathfrak{X}$  runs through all edges of at least one polygon. If  $k = 2$ ,  $\mathfrak{X}$  reduces to a homogeneous tree of degree  $r$ . In what follows, we write  $q = (k - 1)(r - 1)$ ,  $\tau = 2\pi/\log q$  and  $\mathbb{T} = \mathbb{R}/\tau\mathbb{Z}$ . Different notions of length on a symmetric graph were introduced in [Iozzi and Picardello 1983a]. Here we use the definition of the length  $d(x, y)$  between two vertices  $x, y \in \mathfrak{X}$  to denote the minimal number of polygons crossed by a path connecting  $x$  and  $y$ . We fix a reference point  $o$  in  $\mathfrak{X}$  and write  $|x| = d(x, o)$ .

By the same arguments as in [Betori and Pagliacci 1984, Theorem 1], if  $k > 2$ , it is easy to see that every group acting simply transitively on  $\mathfrak{X}$  and isometrically with respect to the metric induced by this length is isometric to the free group  $G = \bigoplus_{i=1}^r \mathbb{Z}_k$ , while, for  $k = 2$ ,  $G$  is isometric to the free product of  $t$  copies of  $\mathbb{Z}$  and  $s$  copies of  $\mathbb{Z}_2$ , where  $2t + s = r$ . Hence every vertex of  $\mathfrak{X}$  is identified with an element of  $G$  and, under this identification, every polygon corresponds to an orbit under right translations by one of the factors  $\mathbb{Z}_k$ . For  $x \in \mathfrak{X}$  and  $n \leq |x|$ , we write  $x^{(n)}$  for the word of length  $n$  consisting of the first  $n$  blocks of  $x$  and simply write  $x'$  for  $x^{(|x|-1)}$ .

Let  $\mathfrak{S}_n$  be the set of words of length  $n$  in  $\mathfrak{X}$ . We write  $\Omega$  for the Poisson boundary of  $\mathfrak{X}$ . For  $\omega \in \Omega$  and  $n \in \mathbb{Z}_{\geq 0}$ , we denote by  $\omega_n$  the word of length  $n$  consisting of the first  $n$  blocks. Let  $E(x)$  denote the subset of  $\Omega$  of words that begin with the reduced word  $x \in \mathfrak{X}$ . We write  $\mathcal{M}$  and  $\mathcal{M}_n$  for the  $\sigma$ -algebra generated by  $\{E(x) : x \in \mathfrak{X}\}$  and  $\sigma$ -subalgebra generated by  $\{E(x) : |x| \leq n\}$  respectively. Then  $\mathcal{M}$  makes

$\Omega$  into a compact topological space and there exists a natural  $G$ -quasi-invariant probability measure  $\nu$  on  $(\Omega, \mathcal{M})$ . We write  $\mathcal{F}(\Omega)$  for the space comprised of the  $\mathcal{M}_n$ -measurable functions on  $\Omega$ . We denote by  $\mathcal{F}(\Omega)_c$  the linear span of the characteristic functions of  $E(x)$  for  $x \in \mathfrak{X}$ . The dual space  $\mathcal{F}'(\Omega)$  is identified with the space of the martingales on  $\Omega$  with respect to  $\{\mathcal{M}_n\}$ .

We write  $s_0 = (\frac{1}{2} - \log_q(k-1))i + \frac{1}{2}\tau$  and set

$$\Upsilon = \{s \in \mathbb{C} : s = \frac{1}{2}i + h\tau, s = s_0 + h\tau \ (h \in \mathbb{Z})\}.$$

We define the subsets  $b_x, c_x, d_x$  of  $\mathfrak{X}$  by the following: for  $x \in \mathfrak{X} \setminus \{o\}$

$$b_x = \{y \in \mathfrak{X} : d(y, x) = 1, |y| = |x|\},$$

$$c_x = \{y \in \mathfrak{X} : d(y, x) = 2, |y| = |x|\},$$

$$d_x = \{y \in \mathfrak{X} : d(y, x) \geq 3, |y| = |x|\},$$

and  $b_o = c_o = d_o = \emptyset$ . The subsets  $B(x)$  and  $C(x)$  of  $\Omega$  are defined by

$$B(x) = \bigcup_{y \in b_x} E(y), \quad C(x) = \bigcup_{y \in c_x} E(y).$$

For a function  $\eta$  on  $\Omega$  and  $n \in \mathbb{Z}_{\geq 0}$ , we define the averages  $E_n\eta$  and  $B_n\eta$  as follows:

$$E_n\eta(\omega) = \frac{1}{\nu(E(\omega_n))} \int_{E(\omega_n)} \eta(\omega') d\nu(\omega'), \quad B_n\eta(\omega) = \frac{1}{\nu(B(\omega_n))} \int_{B(\omega_n)} \eta(\omega') d\nu(\omega').$$

Then, as shown in [Mantero and Zappa 1983, p. 375], the set  $\{E_n\eta\}$  is a martingale associated to  $\eta \in L^1(\Omega)$  and the  $n$ -th martingale difference of  $\eta$  is given by  $D_n\eta = E_n\eta - E_{n-1}\eta$ . Here we set  $E_{-1} = 0$ . For  $x \in \mathfrak{X}$  and  $\omega \in \Omega$ , the Poisson kernel  $p(x, \omega)$  is defined to be the Radon-Nikodym derivative  $d\nu(x^{-1}\omega)/d\nu(\omega)$  and is computed as

$$p(x, \omega) = q^{\zeta(x, \omega)},$$

where  $\zeta(x, \omega) = \lim_{m \rightarrow \infty} (m - d(x, \omega_m))$  is the Busemann function. As shown in [Iozzi and Picardello 1983b, Proposition 2], for  $x \in \mathfrak{S}_n$ , we have

$$(2-1) \quad p(x, \omega) = q^n \chi_{E(x)}(\omega) + \sum_{j=1}^n q^{2j-n-1} \chi_{B(x^{(j)})}(\omega) + \sum_{j=1}^n q^{2j-n-2} \chi_{C(x^{(j)})}(\omega).$$

For  $\eta \in L^1(\Omega)$  and  $s \in \mathbb{T}$ , we define the Poisson transform  $P^s\eta$  by

$$(2-2) \quad P^s\eta(x) = \int_{\Omega} p(x, \omega)^{1/2+is} \eta(\omega) d\nu(\omega).$$

By duality, the Poisson transform is naturally extended to  $\mathcal{F}'(\Omega)$  and is denoted by the same symbol  $P^s$ .

Following [Mantero and Zappa 1983], we define the operators  $\varepsilon$  and  $\Delta$  on  $\mathfrak{X}$ , which are essentially the analogue of  $E_n$  and  $D_n$ . We set for  $n \in \mathbb{Z}_{\geq 0}$

$$S(n, x) = \begin{cases} \{x\}, & |x| \leq n, \\ \{y \in \mathfrak{X} : |y| = |x|, y^{(n)} = x^{(n)}\}, & |x| > n. \end{cases}$$

For a function  $\phi$  on  $\mathfrak{X}$  and  $n \in \mathbb{Z}_{\geq 0}$ , we define its average  $\varepsilon_n \phi$  by

$$(2-3) \quad \varepsilon_n \phi(x) = \frac{1}{\text{Card } S(n, x)} \sum_{y \in S(n, x)} \phi(y).$$

We also define  $\Delta_n \phi$  by

$$\Delta_n \phi(x) = \varepsilon_n \phi(x) - \varepsilon_{n-1} \phi(x).$$

Here we set  $\varepsilon_{-1} \phi = 0$ . We write  $\mu_1$  for the probability measure equidistributed on words of length 1. We also use the notation  $\kappa_1$  to denote the following:

$$(\phi * \kappa_1)(x) = \frac{1}{k-2} \sum_{y \in b_x} \phi(y).$$

We write  $C_c(\mathfrak{X})$  for the set of all compactly supported functions on  $\mathfrak{X}$ . For  $N \in \mathbb{Z}_{\geq 0}$ , we denote by  $C_N(\mathfrak{X})$  the subset of  $C_c(\mathfrak{X})$  consisting of all  $f \in C_c(\mathfrak{X})$  such that  $\text{supp } f \subseteq \mathfrak{B}_N$ . A function  $\phi$  on  $\mathfrak{X}$  is said to be radial if  $\varepsilon_0 \phi = \phi$  and cylindrical if  $\varepsilon_N \phi(x) = \phi(x)$  for some  $N \in \mathbb{Z}_{\geq 0}$ . For any function space  $E(\mathfrak{X})$ , we denote by  $E(\mathfrak{X})^\#$  and  $E(\mathfrak{X})_c$  the subspaces of  $E(\mathfrak{X})$  consisting of all radial functions and cylindrical functions, respectively. A function  $f$  on  $\mathbb{T}$  is said to be Weyl-invariant if  $f(s + \tau) = f(s)$  and  $f(-s) = f(s)$ .

Finally we pointed out that it is meaningful to study harmonic analysis for symmetric graphs using methods similar to that in symmetric spaces. For example, the explicit expressions of the intertwining operators obtained in Section 4 can be used to construct the composition series of the spherical representations and determine which parts of the subquotients are unitarizable. In Section 6, using this information, we can concretely characterize the image of the compactly supported functions under the Helgason–Fourier transform.

### 3. The Poisson transform on symmetric graphs

Iozzi and Picardello [1983b] studied the Poisson transform for symmetric graphs. They showed in their paper that the Poisson transform  $P^s$  is injective on  $\mathcal{F}(\Omega)_c$  if and only if  $s \notin \Upsilon$ . In this section, by carrying out similar arguments to that in [Mantero and Zappa 1983], we show that  $P^s$  is also surjective on  $\mathcal{F}'(\Omega)$  when  $s \notin \Upsilon$ .

As shown in [Iozzi and Picardello 1983b, Theorem 1], we have

$$(P^s \eta * \mu_1)(x) = \gamma(s) P^s \eta(x),$$

where

$$\gamma(s) = \frac{1}{r(k-1)}(q^{1/2+is} + q^{1/2-is} + k - 2).$$

By using the equation

$$\chi_{E(x^{(j)})}(\omega) - \chi_{E(x^{(j+1)})}(\omega) = \chi_{C(x^{(j+1)})}(\omega) + \chi_{B(x^{(j+1)})}(\omega),$$

(2-1) is expressed as

$$(3-1) \quad p(x, \omega) = q^{-|x|} \chi_{E(x^{(0)})}(\omega) + (1 - q^{-2}) \sum_{j=1}^{|x|} q^{2j-|x|} \chi_{E(x^{(j)})}(\omega) \\ + (1 - q^{-1}) \sum_{j=1}^{|x|} q^{2j-|x|-1} \chi_{B(x^{(j)})}(\omega).$$

Therefore for  $\omega \in \Omega$ , substituting (3-1) into (2-2), we have

$$P^s \eta(\omega_n) = \sum_{j=0}^n b_{j,n}(s) E_j \eta(\omega) + \frac{k-2}{q^{1/2+is} + 1} \sum_{j=1}^n b_{j,n}(s) B_j \eta(\omega),$$

where  $b_{0,n}(s) = q^{-n(1/2+is)}$  and

$$b_{j,n}(s) = \frac{q}{r(k-1)} (1 - q^{-1-2is}) q^{-n(1/2+is)+i2js}.$$

By the definitions of  $B_n$  and  $E_n$ , it is easy to verify that

$$E_m B_n = \begin{cases} B_n, & m \geq n, \\ E_m, & m < n, \end{cases} \quad B_m E_n = \begin{cases} E_n, & m > n, \\ B_m, & m \leq n. \end{cases}$$

And hence we obtain that

$$B_m D_n \eta = \begin{cases} D_n \eta, & m > n, \\ B_n \eta - E_{n-1} \eta, & m = n, \\ 0, & m < n. \end{cases}$$

Hereafter we suppose that  $D_M \eta = \eta$  for some  $M \geq 0$ . We first consider the case when  $M > 0$ . Since  $E_j \eta = 0$  and  $B_j \eta = 0$  for  $j < M$ , we have

$$P^s \eta(\omega_n) = 0 \quad \text{for } n < M,$$

and

$$(3-2) \quad P^s \eta(\omega_{M+\ell}) = \sum_{j=M}^{M+\ell} b_{j,M+\ell}(s) \eta(\omega) + \frac{k-2}{q^{1/2+is} + 1} \sum_{j=M+1}^{M+\ell} b_{j,M+\ell}(s) \eta(\omega) \\ + \frac{k-2}{q^{1/2+is} + 1} b_{M,M+\ell}(s) B_M \eta(\omega).$$

We define the function  $Q_M^0(s)$  on  $\mathbb{T}$  by  $Q_0^0(s) = 1$  and

$$Q_M^0(s) = \frac{\sqrt{q}}{r(k-1)} q^{-M/2} q^{i(M-1)s} (q^{1/2+is} - q^{-1/2-is})$$

for  $M > 0$ . By using this, (3-2) can be written

$$\begin{aligned} P^s \eta(\omega_{M+\ell}) &= q^{-\ell(1/2+is)} Q_M^0(s) \left\{ \sum_{j=M}^{M+\ell} q^{i2s(j-M)} \eta(\omega) + \frac{k-2}{q^{1/2+is} + 1} \sum_{j=M+1}^{M+\ell} q^{i2s(j-M)} \eta(\omega) \right. \\ &\quad \left. + \frac{k-2}{q^{1/2+is} + 1} B_M \eta(\omega) \right\}. \end{aligned}$$

Here we set  $Q_0(s) = R_0(s) = 1$  and

$$\begin{aligned} Q_M(s) &= \frac{\sqrt{q}}{r(k-1)} q^{-M/2} q^{i(M-1)s} (q^{1/2+is} - (k-1)q^{-1/2-is} + k-2), \\ R_M(s) &= \frac{\sqrt{q}}{r(k-1)} q^{-M/2} q^{i(M-1)s} (1 - q^{-1/2-is}) \end{aligned}$$

for  $M > 0$ . Then a direct computation yields that

$$(3-3) \quad P^s \eta(\omega_{M+\ell}) = q^{-\ell/2} \psi(\ell+1, s) Q_M(s) \eta(\omega) + (k-2) R_M(s) q^{-\ell(1/2+is)} (B_M \eta(\omega) - \eta(\omega)),$$

where

$$\psi(n, s) = \frac{\sin(ns \log q)}{\sin(s \log q)}.$$

As pointed out in [Cowling and Setti 1999, p. 242],  $D_M \eta = \eta$  if and only if  $\eta$  is constant on  $E(x)$  for every  $x \in \mathfrak{S}_M$  and the average of  $\eta$  with respect to  $E(y)$  for  $|y| < M$  is equal to 0. Therefore we can regard  $\eta$  as a function on  $\mathfrak{X}$  by setting  $\eta(x) = E_{|x|} \eta(\omega)$  for  $\omega \in E(x)$ . Under this identification, we have that  $\eta(x) = 0$  when  $|x| < M$  and  $\eta(x) = \eta(x^{(M)})$  when  $|x| \geq M$ . Moreover, for  $x \in \mathfrak{X}$  such that  $|x| > M$  and  $y \in b_x$ , it holds that  $\eta(x) = \eta(y)$  because  $|x'| = |y'| \geq M$ . We also remark that  $B_M \eta(\omega)$  corresponds to  $\eta * \kappa_1(x^{(M)})$ . For these reasons, (3-3) can be rewritten as follows:

$$(3-4) \quad P^s \eta(x) = q^{-(|x|-M)/2} \psi(|x|-M+1, s) Q_M(s) \eta(x^{(M)}) + (k-2) R_M(s) q^{-(|x|-M)(1/2+is)} (\eta * \kappa_1(x^{(M)}) - \eta(x^{(M)})).$$

In the case  $M = 0$ ,  $\eta$  is a constant function on  $\Omega$  and so  $P^s \eta$  is expressed in terms of the spherical function  $\phi_s$  given in [Iozzi and Picardello 1983b, Theorem 2] as:

$$P^s \eta(x) = \phi_s(x) \eta(o).$$



We summarize these results in the following proposition.

**Proposition 3.1.** *Let  $\eta \in L^1(\Omega)$  be such that  $D_M \eta = \eta$ . Then the Poisson transform  $P^s \eta$  has the following forms:*

(1) *If  $M > 0$ ,*

$$P^s \eta(x) = 0, \quad |x| < M,$$

$$P^s \eta(x) = q^{-(|x|-M)/2} \psi(|x| - M + 1, s) Q_M(s) \eta(x^{(M)}) \\ + (k - 2) R_M(s) q^{-(|x|-M)(1/2+is)} (\eta * \kappa_1(x^{(M)}) - \eta(x^{(M)})), \quad |x| \geq M.$$

(2) *If  $M = 0$ ,*

$$P^s \eta(x) = \phi_s(x) \eta(o).$$

Let  $\mathcal{L}_{\gamma(s)}(\mathfrak{X})$  denote the space consisting of the functions  $\phi$  on  $\mathfrak{X}$  satisfying the condition  $\phi * \mu_1 = \gamma(s)\phi$ . For  $M \in \mathbb{Z}_{\geq 0}$ , we denote by  $\mathcal{L}_{\gamma(s)}^M(\mathfrak{X})$  the subspace of  $\mathcal{L}_{\gamma(s)}(\mathfrak{X})$  consisting of the functions  $\phi$  which satisfy the following conditions:

(1)  $\Delta_M \phi = \phi$ ,

(2) for  $x \in \mathfrak{X}$  such that  $|x| > M$  and  $y \in b_x$ ,  $\phi(x) = \phi(y)$ .

Then  $\mathcal{L}_{\gamma(s)}^0(\mathfrak{X})$  is just the space of radial harmonic functions on  $\mathfrak{X}$ . We prove the following lemma, which is an analogue of [Mantero and Zappa 1983, Lemma 3.2].

**Lemma 3.2.** *Let  $\phi \in \mathcal{L}_{\gamma(s)}^M(\mathfrak{X})$  and  $\omega \in \Omega$ . Then we have the following:*

(1) *If  $M > 0$ ,*

$$\phi(\omega_n) = 0 \quad (n < M),$$

$$(3-5) \quad \phi(\omega_{M+\ell}) = q^{-\ell/2} \psi(\ell + 1, s) \phi(\omega_M) \\ + (k - 2) q^{-(\ell+1)/2} \psi(\ell, s) \times (\phi(\omega_M) - \phi * \kappa_1(\omega_M)).$$

(2) *If  $M = 0$ ,*

$$\phi(\omega_\ell) = \phi_s(\omega_\ell) \phi(o).$$

*Proof.* Since for  $\omega \in \Omega$  and  $n \geq M$ ,

$$\phi * \mu_1(\omega_n) = \frac{1}{r(k-1)} \left( \phi(\omega_{n-1}) + (k-2)\phi * \kappa_1(\omega_n) + \phi(\omega_{n+1}) + \sum_{y \in b_{\omega_{n+1}}} \phi(y) + \sum_{y \in c_{\omega_{n+1}}} \phi(y) \right),$$

and

$$\Delta_{n+1} \phi(\omega_{n+1}) = \phi(\omega_{n+1}) - \frac{1}{q} \left( \phi(\omega_{n+1}) + \sum_{y \in b_{\omega_{n+1}}} \phi(y) + \sum_{y \in c_{\omega_{n+1}}} \phi(y) \right) = 0,$$

we have

$$\phi * \mu_1(\omega_n) = \frac{1}{r(k-1)} \{ \phi(\omega_{n-1}) + (k-2)\phi * \kappa_1(\omega_n) + q\phi(\omega_{n+1}) \}.$$

Hence the condition  $\phi * \mu_1 = \gamma(s)\phi$  implies that

$$(q\beta(s) + k - 2)\phi(\omega_n) = \phi(\omega_{n-1}) + (k - 2)\phi * \kappa_1(\omega_n) + q\phi(\omega_{n+1}),$$

where  $\beta(s) = q^{-1/2+is} + q^{-1/2-is}$ . When  $n > M$ , we have that  $\phi * \kappa_1(\omega_n) = \phi(\omega_n)$  and therefore we have the recursion formulae:

$$(3-6) \quad \phi(\omega_n) = 0, \quad n < M,$$

$$(3-7) \quad \phi(\omega_{M+1}) = \beta(s)\phi(\omega_M) + (k - 2)q^{-1}(\phi(\omega_M) - \phi * \kappa_1(\omega_M)),$$

$$(3-8) \quad \phi(\omega_{M+\ell}) = \beta(s)\phi(\omega_{M+\ell-1}) - q^{-1}\phi(\omega_{M+\ell-2}), \quad \ell \geq 2.$$

In the case  $s \notin (\tau/2)\mathbb{Z}$ , the difference equation (3-8) has the fundamental solutions  $q^{-1/2+is}$  and  $q^{-1/2-is}$ . So using the initial condition (3-7) to determine the coefficients of the fundamental solutions, we obtain the following expression:

$$(3-9) \quad \phi(\omega_{M+\ell}) = C_1 q^{\ell(-1/2+is)} + C_2 q^{\ell(-1/2-is)},$$

where

$$C_1 = \frac{q^{is}\phi(\omega_M) + (k - 2)q^{-1/2}\{\phi(\omega_M) - \phi * \kappa_1(\omega_M)\}}{q^{is} - q^{-is}},$$

$$C_2 = \frac{-q^{-is}\phi(\omega_M) - (k - 2)q^{-1/2}\{\phi(\omega_M) - \phi * \kappa_1(\omega_M)\}}{q^{is} - q^{-is}}.$$

Similarly, when  $s = \frac{1}{2}m\tau$  ( $m \in \mathbb{Z}$ ), we also have

$$(3-10) \quad \phi(\omega_{M+\ell}) = (C'_1 + C'_2\ell)(-1)^{m\ell}q^{-\ell/2},$$

where

$$C'_1 = \phi(\omega_M), \quad C'_2 = \phi(\omega_M) + (k - 2)(-1)^m q^{-1/2}(\phi(\omega_M) - \phi * \kappa_1(\omega_M)).$$

Obviously both the expressions (3-9) and (3-10) agree with the equation (3-5). The case  $M = 0$  is analogous. This concludes the proof.  $\square$

Let  $x \in \mathfrak{S}_M$  and  $s \notin \Upsilon$ . Then we have from (3-4) that

$$(3-11) \quad P^s\eta(x) = Q_M(s)\eta(x) + (k - 2)R_M(s)(\eta * \kappa_1(x) - \eta(x)).$$

We put  $\phi(x) = P^s\eta(x)$  and write down  $\eta$  in terms of  $\phi$  and  $\phi * \kappa_1$ . Since

$$(3-12) \quad \phi * \kappa_1 * \kappa_1(x) = \frac{1}{k-2}\phi(x) + \frac{k-3}{k-2}\phi * \kappa_1(x),$$

we have from (3-11) that

$$(3-13) \quad \phi * \kappa_1(x) = Q_M(s)(\eta * \kappa_1)(x) + R_M(s)(\eta(x) - \eta * \kappa_1(x)).$$

Thus, solving the simultaneous equations (3-11) and (3-13), we get the expressions

$$\eta(x) = \frac{(q^{1/2+is} + k - 2)\phi(x) - (k - 2)\phi * \kappa_1(x)}{q^{1/2+is} Q_M(s)},$$

$$\eta * \kappa_1(x) = \frac{(q^{1/2+is} + 1)\phi * \kappa_1(x) - \phi(x)}{q^{1/2+is} Q_M(s)}.$$

The above expressions suggest the following proposition.

**Proposition 3.3** (cf., [Mantero and Zappa 1983, Proposition 3.4]). *Let  $s \notin \Upsilon$ . For  $\phi \in \mathcal{L}_{\gamma(s)}^M(\mathfrak{X})$ , there exists a function  $\eta$  on  $\Omega$  such that  $D_M \eta = \eta$  and  $P^s \eta = \phi$ .*

*Proof.* Suppose that  $M > 0$ . Indeed, define  $\eta(\omega)$  by

$$\eta(\omega) = \frac{(q^{1/2+is} + k - 2)\phi(\omega_M) - (k - 2)\phi * \kappa_1(\omega_M)}{q^{1/2+is} Q_M(s)}.$$

Then  $\phi(\omega_M) = P^s \eta(\omega_M)$  is trivial. Applying Lemma 3.2 to our case together with (3-11) and (3-13), we see that

$$\begin{aligned} \phi(\omega_{M+\ell}) &= q^{-\ell/2} \psi(\ell + 1, s) \phi(\omega_M) + (k - 2) q^{-(\ell+1)/2} \psi(\ell, s) \\ &\quad \times (\phi(\omega_M) - \phi * \kappa_1(\omega_M)) \\ &= q^{-\ell/2} \psi(\ell + 1, s) \{Q_M(s) \eta(\omega_M) + (k - 2) R_M(s) (\eta * \kappa_1(\omega_M) - \eta(\omega_M))\} \\ &\quad + (k - 2) q^{-(\ell+1)/2} \psi(\ell, s) q^{1/2+is} R_M(s) (\eta(\omega_M) - \eta * \kappa_1(\omega_M)) \\ &= q^{-\ell/2} \psi(\ell + 1, s) Q_M(s) \eta(\omega_M) + (k - 2) q^{-\ell/2} R_M(s) \\ &\quad \times \{\psi(\ell + 1, s) - q^{is} \psi(\ell, s)\} (\eta * \kappa_1(\omega_M) - \eta(\omega_M)) \\ &= P^s \eta(\omega_{M+\ell}). \end{aligned}$$

The case  $M = 0$  is analogous. This concludes the proof.  $\square$

The following proposition is proved in the same way as in [Mantero and Zappa 1983, Corollary 3.5] and hence we omit its proof.

**Proposition 3.4.** *Suppose that  $s \notin \Upsilon$ . Then the Poisson transform  $P^s$  is a bijective operator from  $\mathcal{F}'(\Omega)$  onto  $\mathcal{L}_{\gamma(s)}(\mathfrak{X})$ .*

#### 4. The construction of the intertwining operator

Mantero and Zappa [1983] defined the intertwining operator between the spherical representations for free groups and gave an explicit expression of the intertwining operator. In this section, we extend their results to the case of symmetric graphs.

Let  $s \in \mathbb{C}$  and define the action  $\pi_s$  of  $G$  on  $L^2(\Omega)$  by

$$(\pi_s(g)\eta)(\omega) = p(g \cdot o, \omega)^{1/2+is} \eta(g^{-1}\omega).$$

The representation  $(\pi_s, L^2(\Omega))$  is called the spherical representation. We denote by  $\lambda$  the left regular representation of  $G$  on  $\mathcal{L}_{\gamma(s)}(\mathfrak{X})$ . Then as indicated in [Iozzi and Picardello 1983b, p. 372], the Poisson transform  $P^{-s}$  intertwines  $\pi_s$  and  $\lambda$ . Therefore, in the case  $\pm s \notin \Upsilon$ , we see from Proposition 3.4 that the operator on  $\mathcal{F}'(\Omega)$  defined by  $I_s = (P^s)^{-1}P^{-s}$  is bijective and satisfies the following relation:

$$(4-1) \quad I_s \pi_s(g) = \pi_{-s}(g) I_s.$$

Let  $\eta \in L^1(\Omega)$  be such that  $D_M \eta = \eta$  for some  $M > 0$ . Under this assumption,  $I_s(B_M \eta) = B_M(I_s \eta)$  because

$$P^s(B_M I_s \eta)(x) = P^s I_s \eta * \kappa_1(x) = P^{-s} \eta * \kappa_1(x) = P^{-s}(B_M \eta)(x).$$

Since

$$P^s I_s \eta(\omega_{M+\ell}) = P^{-s} \eta(\omega_{M+\ell}),$$

we have from (3-3) that

$$\begin{aligned} & q^{-\ell/2} \psi(\ell+1, s) Q_M(s) I_s \eta(\omega) + (k-2) R_M(s) q^{-\ell(1/2+is)} (I_s B_M \eta(\omega) - I_s \eta(\omega)) \\ &= q^{-\ell/2} \psi(\ell+1, s) Q_M(-s) \eta(\omega) + (k-2) R_M(-s) q^{-\ell(1/2-is)} (B_M \eta(\omega) - \eta(\omega)). \end{aligned}$$

Taking  $\ell = 0$  and  $\ell = 1$  respectively, we obtain from the above equation that

$$(4-2) \quad \begin{aligned} Q_M^0(s) I_s \eta(\omega) + (k-2) R_M(s) I_s B_M \eta(\omega) \\ = Q_M^0(-s) \eta(\omega) + (k-2) R_M(-s) B_M \eta(\omega) \end{aligned}$$

and

$$(4-3) \quad \begin{aligned} \beta(s) Q_M(s) I_s \eta(\omega) + (k-2) q^{-(1/2+is)} R_M(s) (I_s B_M \eta(\omega) - I_s \eta(\omega)) \\ = \beta(s) Q_M(-s) \eta(\omega) + (k-2) q^{-(1/2-is)} R_M(-s) (B_M \eta(\omega) - \eta(\omega)). \end{aligned}$$

Solving the simultaneous equations (4-2) and (4-3), we have

$$(4-4) \quad \begin{aligned} I_s \eta(\omega) &= \frac{q^{-is} Q_M(-s) \eta(\omega) + (q^{is} - q^{-is}) Q_M^0(-s) \eta(\omega)}{q^{is} Q_M(s)} \\ &+ \frac{(k-2)(q^{is} - q^{-is}) R_M(-s) B_M \eta(\omega)}{q^{is} Q_M(s)}, \end{aligned}$$

$$(4-5) \quad \begin{aligned} I_s B_M \eta(\omega) &= \frac{(q^{is} - q^{-is}) R_M(-s) \{\eta(\omega) - B_M \eta(\omega)\}}{q^{is} Q_M(s)} \\ &+ \frac{q^{is} Q_M(-s) B_M \eta(\omega)}{q^{is} Q_M(s)}. \end{aligned}$$

For  $\eta \in L^1(\Omega)$  satisfying  $D_M\eta = \eta$ , we define  $\eta^+$  and  $\eta^-$  as

$$(4-6) \quad \eta^+(\omega) = \eta(\omega) + (k-2)B_M\eta(\omega),$$

$$(4-7) \quad \eta^-(\omega) = \eta(\omega) - B_M\eta(\omega).$$

Then we have from (4-4) and (4-5) that

$$(4-8) \quad I_s\eta^+(\omega) = \frac{Q_M(-s)}{Q_M(s)}\eta^+(\omega),$$

$$(4-9) \quad I_s\eta^-(\omega) = \frac{q^{-is}R_M(-s)}{q^{is}R_M(s)}\eta^-(\omega).$$

For  $n \in \mathbb{Z}_{\geq 0}$ , we denote by  $\mathcal{H}_n$  the subspace of  $\mathcal{F}(\Omega)_c$  consisting of  $\eta$  such that  $D_n\eta = \eta$ . We write  $\mathcal{H}_n^+$  and  $\mathcal{H}_n^-$  for the subspaces of  $\mathcal{H}_n$  generated by  $\{\eta^+ : \eta \in \mathcal{H}_n\}$  and  $\{\eta^- : \eta \in \mathcal{H}_n\}$ , respectively. Then it holds that  $\mathcal{H}_0 = \mathcal{H}_0^+$  and  $\mathcal{H}_n = \mathcal{H}_n^+ \oplus \mathcal{H}_n^-$  for  $n > 0$ . The expressions (4-8) and (4-9) give the explicit forms of the intertwining operator  $I_s$  when restricted to  $\mathcal{H}_n^+$  and  $\mathcal{H}_n^-$ , respectively.

Finally in this section, we list some properties of the Poisson transform. Analogously to (4-6) and (4-7), for a function  $\phi$  on  $\mathfrak{X}$ , we define  $\phi^+$  and  $\phi^-$  by

$$\phi^+(x) = \phi(x) + (k-2)\phi * \kappa_1(x), \quad \phi^-(x) = \phi(x) - \phi * \kappa_1(x).$$

Let  $\eta \in L^1(\Omega)$  be such that  $D_M\eta = \eta$ . Then taking into account (3-12), we see from Proposition 3.1 that

$$(4-10) \quad (P^s\eta)^+(x) = q^{-(|x|-M)/2}\psi(|x|-M+1, s)Q_M(s)\eta^+(\omega) = (P^s\eta^+)(x)$$

and

$$(4-11) \quad (P^s\eta)^-(x) = q^{-(|x|-M)/2}\psi(|x|-M+1, s)Q_M(s)\eta^-(\omega) \\ - (k-1)q^{-(|x|-M)(1/2+is)}R_M(s)\eta^-(\omega) \\ = (P^s\eta^-)(x).$$

We here set

$$\psi^-(\ell+1, s) = \sum_{j=0}^{\ell} q^{i(\ell-2j)s} + q^{-1/2}(k-1) \sum_{j=1}^{\ell} q^{i(\ell-2j+1)s}.$$

Then the expression (4-11) can be written

$$(4-12) \quad (P^s\eta)^-(x) = q^{-(|x|-M)/2}\psi^-(|x|-M+1, s)q^{1/2+is}R_M(s)\eta^-(\omega).$$

We summarize these in the following corollary.

**Corollary 4.1.** *Let  $\eta \in L^1(\Omega)$  be such that  $D_M\eta = \eta$ . Then*

$$(P^s\eta)^+(x) = (P^s\eta^+)(x), \quad (P^s\eta)^-(x) = (P^s\eta^-)(x).$$

In addition, for any  $x \in \mathfrak{X}$  satisfying  $|x| > M$ , we have

- (1)  $Q_M(s)^{-1}(P^s\eta)^+(x)$  is an even entire holomorphic function on  $\mathbb{C}$  with respect to the variable  $s$ ,
- (2)  $(q^{1/2+is}R_M(s))^{-1}(P^s\eta)^-(x)$  is an even entire holomorphic function on  $\mathbb{C}$  with respect to the variable  $s$ .

### 5. The spectral projection on symmetric graphs

We first review the Helgason–Fourier transform for symmetric graphs and its inversion formula, which were introduced by Eddine [2013; 2015].

Let  $(\pi_s, L^2(\Omega))$  be a spherical representation and let  $I_s$  be the intertwining operator defined in the previous section. The Helgason–Fourier transform  $\tilde{f}(s, \omega)$  of  $f \in C_c(\mathfrak{X})$  is defined by

$$(5-1) \quad \tilde{f}(s, \omega) = (\pi_s(f)1)(\omega) = \sum_{x \in \mathfrak{X}} f(x)p(x, \omega)^{1/2+is}.$$

Here 1 denotes the function identically one on  $\Omega$ . In [Jamal Eddine 2015, Lemma 3.10], Eddine proved the following inversion formula:

$$f(x) = \frac{(k-r)_+}{k} \int_{\Omega} \tilde{f}(s_0, \omega)p(x, \omega)^{1/2-is_0} d\nu(\omega) + c_G \int_{\Omega} \int_{\mathbb{T}} \tilde{f}(s, \omega)p(x, \omega)^{1/2-is} |c(s)|^{-2} ds d\nu(\omega),$$

where  $c_G = q/\{2\tau r(k-1)\}$  and

$$c(s) = \frac{\sqrt{q}}{q+1} \cdot \frac{q^{1/2+is} - (k-1)q^{-1/2-is} + k-2}{q^{is} - q^{-is}}$$

is a  $c$ -function. Here  $(k-r)_+$  stands for  $k-r$  when  $k > r$  and to 0 when  $k \leq r$ . As described in [Cowling and Setti 1999, p. 240], we see that  $I_s \tilde{f}(s, \omega) = \tilde{f}(-s, \omega)$  for almost all  $s \in \mathbb{T}$  and thus we obtain the following symmetry condition:

$$(5-2) \quad \int_{\Omega} \tilde{f}(s, \omega)p(x, \omega)^{1/2-is} d\nu(\omega) = \int_{\Omega} \tilde{f}(-s, \omega)p(x, \omega)^{1/2+is} d\nu(\omega).$$

Following Bray [1996], we define the spectral projection  $P_s f$  of  $f \in C_c(\mathfrak{X})$  by

$$(5-3) \quad P_s f(x) = (f * \phi_s)(x) = \int_G f(g_1)\phi_s(g_1^{-1}g) dg_1,$$

where  $x = g \cdot o$ . Applying the functional equation of the spherical function [Jamal Eddine 2015, Lemma 3.9] and using Fubini's theorem, we obtain

$$\begin{aligned} P_s f(x) &= \int_G f(g_1 \cdot o) \int_{\Omega} p(g_1 \cdot o, \omega)^{1/2+is} p(g \cdot o, \omega)^{1/2-is} d\nu(\omega) dg_1 \\ &= \int_{\Omega} \tilde{f}(s, \omega) p(x, \omega)^{1/2-is} d\nu(\omega). \end{aligned}$$

Thus the spectral projection  $P_s f(x)$  is Weyl-invariant with respect to the variable  $s$  and has the following inversion formula:

$$(5-4) \quad f(x) = \frac{(k-r)_+}{k} P_{s_0} f(x) + c_G \int_{\mathbb{T}} P_s f(x) |c(s)|^{-2} ds.$$

Let  $a \in \mathfrak{X}$  and define the function  $\xi_a$  on  $\Omega$  by  $\xi_o(\omega) = 1$  and for  $a \neq o$

$$\xi_a(\omega) = \nu(E(a))^{-1} \chi_{E(a)}(\omega) - \nu(E(a'))^{-1} \chi_{E(a')}(\omega).$$

Then it is easy to see that  $D_{|a|} \xi_a = \xi_a$  and  $B_{|a|} \xi_a = (\xi_* * \kappa_1)(a)$ . For  $a \in \mathfrak{X}$  and  $s \in \mathbb{C}$ , we define the generalized spherical function  $\Phi_{a,s}$  on  $\mathfrak{X}$  by

$$\Phi_{a,s}(x) = P^s \xi_a(x) = \int_{\Omega} p(x, \omega)^{1/2+is} \xi_a(\omega) d\nu(\omega).$$

By Proposition 3 in [Koizumi 2013] combined with Corollary 4.1, we have that

$$(\Delta_n P_s f)^{\pm}(x) = \int_{\Omega} \Phi_{\omega_n, -s}^{\pm}(x) \tilde{f}(s, \omega) d\nu(\omega).$$

The explicit expressions of  $\Phi_{\omega_n, -s}^{\pm}$  are given by (4-10) and (4-12). We see from these that

$$\begin{aligned} (\Delta_n P_s f)^+(x) &= 0 \quad \text{when } \pm s \in \Upsilon, \\ (\Delta_n P_s f)^-(x) &= 0 \quad \text{when } \pm s \in \frac{1}{2}i + \tau\mathbb{Z}. \end{aligned}$$

Furthermore, we have the following proposition.

**Proposition 5.1.** *Let  $f \in C_N(\mathfrak{X})$ . Then  $(\Delta_n P_{s_0} f)^-(x) = 0$  when  $n > N$ .*

*Proof.* By the definition of the Poisson transform, we have

$$\begin{aligned} (\Delta_n P_{s_0} f)^-(x) &= \sum_{y \in \mathfrak{X}} f(y) \left\{ \int_{\Omega} p(x, \omega)^{1/2-is_0} \Phi_{\omega_n, s_0}^-(y) d\nu(\omega) \right\} \\ &= \sum_{y \in \mathfrak{X}} f(y) \left\{ \int_{\Omega} (1-r)^{\zeta(x, \omega)} \left( \frac{1}{1-k} \right)^{|y|} \frac{k(k-1)^{n-1}}{r(r-1)^{n-1}} \xi_{\omega_n}^-(\omega') d\nu(\omega) \right\}, \end{aligned}$$

where we choose  $\omega' \in E(y)$ . Taking into account

$$\xi_{\omega_n}^-(\omega') = \begin{cases} r(k-1)q^{n-1}, & \omega \in E(y^{(n)}), \\ -r(k-1)q^{n-1}/(k-2), & \omega \in B(y^{(n)}), \\ 0, & \text{otherwise,} \end{cases}$$

we have

$$\begin{aligned} (\Delta_n P_{s_0} f)^-(x) &= k(k-1)^{2n-1} \sum_{\substack{y \in \mathfrak{X} \\ |y| \geq n}} f(y) \left(\frac{1}{1-k}\right)^{|y|} \int_{E(y^{(n)})} (1-r)^{\zeta(x,\omega)} d\nu(\omega) \\ &\quad - \frac{k(k-1)^{2n-1}}{k-2} \sum_{\substack{y \in \mathfrak{X} \\ |y| \geq n}} f(y) \left(\frac{1}{1-k}\right)^{|y|} \int_{B(y^{(n)})} (1-r)^{\zeta(x,\omega)} d\nu(\omega). \end{aligned}$$

Since  $f(y) = 0$  for  $|y| \geq n > N$ , we have  $(\Delta_n P_{s_0} f)^-(x) = 0$ .  $\square$

## 6. Paley–Wiener theorem for spectral projection

In this section, we shall characterize the image of  $C_c(\mathfrak{X})$  under the spectral projection on  $\mathfrak{X}$ . As an application, we shall prove the Paley–Wiener theorem of the Helgason–Fourier transform for symmetric graphs.

Throughout this section, for a function  $\phi$  on  $\mathfrak{X}$ , we denote  $\Delta_n \phi(x)$  by  $\phi_n(x)$ . Let  $N \in \mathbb{Z}_{\geq 0}$ . We denote by  $\mathcal{T}_N(\mathbb{T} \times \mathfrak{X})$  the set comprised of all functions  $F$  on  $\mathbb{T} \times \mathfrak{X}$  satisfying the following conditions:

- (N1)  $F(s, x)$  is a Weyl-invariant smooth function on  $\mathbb{R}$  with respect to the variable  $s$ .
- (N2) For each  $n \in \mathbb{Z}_{\geq 0}$  and  $s \in \mathbb{R}$ ,  $F_n(s, x) \in \mathcal{L}_{\gamma(s)}^n(\mathfrak{X})$ ,
- (N3) For each  $x \in \mathfrak{X}$ ,  $F(s, x)$  extends to a Weyl-invariant holomorphic function on  $\mathbb{C}$ .
- (N4) For each  $n \in \mathbb{Z}_{\geq 0}$ ,  $Q_n(-s)^{-1} F_n^+(s, x)$  is holomorphic on  $\mathbb{C}$  and there exists a constant  $C_N > 0$  which does not depend on the choice of  $n$  such that

$$|Q_n(-s)^{-1} F_n^+(s, x)| \leq C_N q^{(|x|-n+N)|\Im s|}.$$

- (N5) For each  $n \in \mathbb{Z}_{> 0}$ ,  $(q^{1/2-is} R_n(-s))^{-1} F_n^-(s, x)$  is holomorphic on  $\mathbb{C}$  and there exists a constant  $C_N > 0$  which does not depend on the choice of  $n$  such that

$$|(q^{1/2-is} R_n(-s))^{-1} F_n^-(s, x)| \leq C_N q^{(|x|-n+N)|\Im s|}.$$

- (N6)  $F_n^-(s_0, x) = 0$  when  $n > N$ .

We set

$$\mathcal{T}(\mathbb{T} \times \mathfrak{X}) = \bigcup_{N=0}^{\infty} \mathcal{T}_N(\mathbb{T} \times \mathfrak{X}).$$



The following proposition is obtained by the same arguments as in [Koizumi 2013, Proposition 4].

**Proposition 6.1.** *Let  $f \in C_N(\mathfrak{X})$ . Then  $F(s, x) = P_s f(x)$  belongs to  $\mathcal{T}_N(\mathbb{T} \times \mathfrak{X})$ .*

To prove the sufficient condition in the Paley–Wiener theorem, we need the following lemma.

**Lemma 6.2** (cf., [Koizumi 2013, Lemma 1]). *Let  $N \in \mathbb{Z}_{>0}$ ,  $F \in \mathcal{T}_N(\mathbb{T} \times \mathfrak{X})$  and  $a \in \mathfrak{S}_n$ . If  $n > N$  then  $F_n^+(s, a) = 0$  and  $F_n^-(s, a) = 0$  for all  $s \in \mathbb{T}$ .*

*Proof.* We shall first show that  $F_n^-(s, a) = 0$ . Let us set

$$\phi^-(s) = (q^{1/2-is} R_n(-s))^{-1} F_n^-(s, a).$$

Then we see

$$(6-1) \quad \phi^-(-s) = (q^{1/2+is} R_n(s))^{-1} F_n^-(s, a) = \frac{q^{1/2-is} R_n(-s)}{q^{1/2+is} R_n(s)} \phi^-(s).$$

We put

$$c^-(n, s) = \frac{q^{1/2-is} R_n(-s)}{q^{1/2+is} R_n(s)}.$$

Obviously we have

$$(6-2) \quad c^-(n, s) = -q^{-1/2-is(2n-1)} + (1 - q^{-1}) \sum_{\ell=0}^{\infty} q^{-\ell/2-is(2n+\ell)}.$$

The condition (N5) yields that  $\phi^-(s)$  is an entire function of exponential type  $N$ . We use the Paley–Wiener theorem on  $\mathbb{Z}$  to write

$$\phi^-(s) = \sum_{m \in \mathbb{Z}} \phi^-(m) q^{ims},$$

where  $\phi^-(m) = 0$  unless  $-N \leq m \leq N$ . Substituting (6-2) to (6-1), we have

$$\sum_{m \in \mathbb{Z}} \phi^-(m) q^{-ims} = \sum_{m \in \mathbb{Z}} \left[ -q^{-\frac{1}{2}-is(2n-1)} + (1 - q^{-1}) \sum_{\ell=0}^{\infty} q^{-\frac{\ell}{2}-is(2n+\ell)} \right] \times \phi^-(m) q^{ims},$$

and thus we have the following recursion formula:

$$(6-3) \quad \phi^-(m) = -q^{-1/2} \phi^-(2n - 1 - m) + (1 - q^{-1}) \sum_{\ell=0}^{\infty} q^{-\ell/2} \phi^-(2n + \ell - m).$$

From (6-3), when  $n > N + 1$ , it is easily verified that  $\phi^-(m) = 0$  for all  $m \in \mathbb{Z}$ . When  $n = N + 1$ , (6-3) implies

$$\phi^-(m) = -q^{-1/2} \phi^-(2N + 1 - m),$$

and so  $\phi^-(N) = 0$ . We consequently have that  $\phi^-(m) = 0$  for all  $m \in \mathbb{Z}$ .

We shall next show that  $F_n^+(s, a) = 0$ . We set  $\phi^+(s) = Q_n(-s)^{-1}F_n^+(s, a)$ . We use the Paley–Wiener theorem on  $\mathbb{Z}$  to write

$$\phi^+(s) = \sum_{m \in \mathbb{Z}} \phi^+(m)q^{ims},$$

where  $\phi^+(m) = 0$  unless  $-N \leq m \leq N$ . Putting

$$c^+(n, s) = \frac{Q_n(-s)}{Q_n(s)},$$

we have

$$\phi^+(-s) = c^+(n, s)\phi^+(s).$$

Because

$$c^+(n, s) = \frac{1 + (k-1)q^{-1/2+is}}{1 + (k-1)q^{-1/2-is}}c^-(n, s),$$

we have

$$\begin{aligned} (6-4) \quad \phi^+(m) &= -(k-1)q^{-1}\phi^+(2n-2-m) \\ &\quad + (k-1)(1-q^{-1})\sum_{\ell=0}^{\infty} q^{-(\ell+1)/2}\phi^+(2n+\ell-m-1) \\ &\quad - (1-(1-k)^2q^{-1})\sum_{\ell=0}^{\infty} (1-k)^\ell q^{-(\ell+1)/2}\phi^+(2n+\ell-1-m) \\ &\quad + \frac{(1-q^{-1})(1-(1-k)^2q^{-1})}{k}\sum_{\ell=0}^{\infty} (1-(1-k)^{\ell+1})q^{-\ell/2}\phi^+(2n+\ell-m). \end{aligned}$$

From (6-4), when  $n > N + 1$ , it is easily verified that  $\phi^+(m) = 0$  for all  $m \in \mathbb{Z}$ . In the case  $n = N + 1$ , (6-4) implies

$$\phi^+(m) = -(k-1)q^{-1}\phi^+(2N-m),$$

and so  $\phi^+(N) = 0$ . Therefore, in this case,  $\phi^+(m) = 0$  for all  $m \in \mathbb{Z}$ .  $\square$

The proof of the sufficient condition in the Paley–Wiener theorem is like the proof for the case of semisimple Lie groups given by Campoli [1980] and Johnson [1979]. We remark that  $\Im s_0 \geq 0$  when  $k \leq r$  and  $\Im s_0 < 0$  when  $k > r$ . We also remark that the residue of  $1/c(s)$  at  $s = s_0$  is equal to  $r(k-r)/\{ik(r-1)\log q\}$  and  $1/c(-s_0) = 1$ . We first show the following proposition.

**Proposition 6.3.** *Let  $N \in \mathbb{Z}_{>0}$ ,  $F \in \mathcal{F}_N(\mathbb{T} \times \mathfrak{X})$  and set*

$$f_0(x) = c_G \int_{\mathbb{T}} F_0(s, x) |c(s)|^{-2} ds.$$

Then there exists a function  $J_0 \in C_N(\mathfrak{X})^\#$  such that

$$f_0(x) - J_0(x) = \frac{(k-r)_+}{k} F_0(s_0, x).$$

*Proof.* We put  $F(s) = F_0(s, o)$ . Then it follows from Lemma 3.2 and the condition (N4) that  $F(s)$  is an even entire function of exponential order  $N$  and

$$F_0(s, x) = \phi_s(x) F(s).$$

We thus have

$$\begin{aligned} f(x) &= c_G \int_{\mathbb{T}} \phi_s(x) F(s) |c(s)|^{-2} ds \\ (6-5) \quad &= c_G \int_{\mathbb{T}} F(s) \frac{1}{c(-s)} q^{(is-1/2)|x|} ds + c_G \int_{\mathbb{T}} F(s) \frac{1}{c(s)} q^{(-is-1/2)|x|} ds. \end{aligned}$$

We write  $f_1(x)$  and  $f_2(x)$  for the first term and the second term of the last expression of (6-5), respectively. For a sufficiently large  $\eta > 0$ , let  $f_{1,\eta}$  denote the formula shifting the path of integral of  $f_1$  from  $\mathbb{T}$  to  $\mathbb{T} + i\eta$  and let  $f_{2,-\eta}$  denote the formula shifting the path of integral of  $f_2$  from  $\mathbb{T}$  to  $\mathbb{T} - i\eta$ .

Suppose that  $k \leq r$ . Because  $f_1$  is analytic inside the rectangle with corners  $\pm\tau/2$  and  $\pm\tau/2 + i\eta$ , we have by Cauchy's theorem that  $f_1 = f_{1,\eta}$ . Similarly we can also obtain that  $f_2 = f_{2,-\eta}$ . In case  $k > r$ , we have

$$\begin{aligned} f_1(x) - f_{1,\eta}(x) &= 2\pi i c_G \operatorname{Res}_{s=-s_0} \left\{ F(s) \frac{1}{c(-s)} q^{(is-1/2)|x|} \right\} \\ &= \frac{k-r}{2k} F(-s_0) \left( \frac{1}{1-k} \right)^{|x|}, \\ f_2(x) - f_{2,-\eta}(x) &= 2\pi i c_G \operatorname{Res}_{s=s_0} \left\{ F(s) \frac{1}{c(s)} q^{(-is-1/2)|x|} \right\} \\ &= \frac{k-r}{2k} F(s_0) \left( \frac{1}{1-k} \right)^{|x|}. \end{aligned}$$

Then the assertion follows immediately. □

In the nonradial case, we need slightly complicated calculations.

**Proposition 6.4.** *Let  $N \in \mathbb{Z}_{>0}$ ,  $F \in \mathcal{T}_N(\mathbb{T} \times \mathfrak{X})$  and set*

$$f_n(x) = c_G \int_{\mathbb{T}} F_n(s, x) |c(s)|^{-2} ds$$

*for  $n > 0$ . Then there exists a function  $J_n \in C_N(\mathfrak{X})$  such that*

$$f_n(x) - J_n(x) = \frac{(k-r)_+}{k} F_n(s_0, x).$$

*Proof.* We put  $a = x^{(n)}$  and choose  $\omega \in E(x)$ . Because  $F_n^\pm$  satisfies the condition (N2), we have from (4-10) and (4-12) that

$$(6-6) \quad F_n^+(s, x) = q^{-(|x|-|a|)/2} \psi(|x| - |a| + 1, s) Q_n(s) F_n^+(s, a),$$

$$(6-7) \quad F_n^-(s, x) = q^{-(|x|-|a|)/2} \psi^-(|x| - |a| + 1, s) q^{1/2+is} R_n(s) F_n^-(s, a).$$

In the case when  $n > N$ , Lemma 6.2 yields that  $F_n^+(s, a) = 0$  and  $F_n^-(s, a) = 0$ . On the other hand, since  $F_n^-(s_0, x) = 0$ , it suffices to set  $J_n(x) = 0$ .

In the following, we suppose that  $n \leq N$ . Substituting (6-6) and (6-7), we obtain

$$f_n^\pm(x) = \xi_a^\pm(\omega)^{-1} c_G \int_{\mathbb{T}} h_a^\pm(s) \Phi_{a,s}^\pm(x) |c(s)|^{-2} ds,$$

where  $h_a^+(s) = Q_{|a|}(s)^{-1} F_n^+(s, a)$ ,  $h_a^-(s) = (q^{1/2+is} R_{|a|}(s))^{-1} F_n^-(s, a)$  and  $\omega \in E(x)$ . We know that  $\Phi_{a,s}^\pm(x)$  has the following expansion:

$$(6-8) \quad \Phi_{a,s}^+(x) = \{c(s)q^{(is-1/2)|x|} - q^{i2(|a|-1)s} c(s)q^{(-is-1/2)|x|}\} \xi_a^+(\omega).$$

Hence we can show  $f_n^+ \in C_N(\mathfrak{X})$  for all  $k, r$  by the same arguments as in the proof of Proposition 6.3.

Hereafter we shall compute  $f_n^-(x)$ . It follows from (6-8) that

$$\begin{aligned} \Phi_{a,s}^-(x) &= \{c(s)q^{(is-1/2)|x|} - q^{i2(|a|-1)s} c(s)q^{(-is-1/2)|x|} \\ &\quad - (k-1)R_{|a|}(s)q^{-(|x|-|a|)(1/2+is)}\} \xi_a^-(\omega). \end{aligned}$$

Let us set

$$\begin{aligned} f_{n,1}^-(x) &= c_G \int_{\mathbb{T}} h_a^-(s) \frac{1}{c(-s)} q^{(is-1/2)|x|} ds, \\ f_{n,2}^-(x) &= c_G \int_{\mathbb{T}} h_a^-(s) \frac{1}{c(-s)} q^{i2(|a|-1)s} q^{(-is-1/2)|x|} ds, \\ f_{n,3}^-(x) &= c_G \int_{\mathbb{T}} h_a^-(s) R_{|a|}(s) q^{-(|x|-|a|)(1/2+is)|x|} |c(s)|^{-2} ds. \end{aligned}$$

Suppose first that  $k \leq r$ . Then as shown in Proposition 6.3, we see that  $f_{n,1}^- \in C_N(\mathfrak{X})$ . Keeping the notation in the proof of Proposition 6.3, we have

$$\begin{aligned} f_{n,2}^-(x) - f_{n,2,-\eta}^-(x) &= 2\pi i c_G \operatorname{Res}_{s=-s_0} \left\{ h_a^-(s) \frac{1}{c(-s)} q^{i2(|a|-1)s} q^{(-is-1/2)|x|} \right\} \\ &= \frac{r(k-r)}{2k} h_a^-(-s_0) (1-k)^{-|a|+1} (1-r)^{-|x|+|a|-1}, \\ f_{n,3}^-(x) - f_{n,3,-\eta}^-(x) &= 2\pi i c_G \operatorname{Res}_{s=-s_0} \{ h_a^-(s) R_{|a|}(s) q^{-(is+1/2)(|x|-|a|)} |c(s)|^{-2} \} \\ &= \frac{r(k-r)}{2k} h_a^-(-s_0) (1-k)^{-|a|} (1-r)^{-|x|+|a|-1}. \end{aligned}$$

Setting

$$f_{n,\eta}^-(x) = f_{n,1}^-(x) - f_{n,2,-\eta}^-(x) - (k-1)f_{n,3,-\eta}^-(x),$$

we obtain that

$$f_n^-(x) - f_{n,\eta}^-(x) = 0.$$

Thus we have by contour integration arguments that  $f_n^- \in C_N(\mathfrak{X})$ .

Suppose next that  $k > r$ . In this case, by the same discussion as in the proof of Proposition 6.3, we see that  $f_{n,2}^- \in C_N(\mathfrak{X})$ . Moreover we have

$$\begin{aligned} f_{n,1}^-(x) - f_{n,1,\eta}^-(x) &= 2\pi i c_G \operatorname{Res}_{s=-s_0} \left\{ h_a^-(s) \frac{1}{c(-s)} q^{(is-1/2)|x|} \right\} \\ &= \frac{k-r}{2k} \left( \frac{1}{1-k} \right)^{|x|} h_a^-(-s_0), \\ f_{n,3}^-(x) - f_{n,3,-\eta}^-(x) &= 2\pi i c_G \operatorname{Res}_{s=s_0} \{ h_a^-(s) R_{|a|}(s) q^{-(|x|-|a|)(1/2+is)} |c(s)|^{-2} \} \\ &= \frac{k-r}{2r} h_a^-(s_0) (1-k)^{-|x|+|a|-2} (1-r)^{-|a|+1}. \end{aligned}$$

Let us set

$$f_{n,\eta}^-(x) = f_{n,1,\eta}^-(x) - f_{n,2}^-(x) - (k-1)f_{n,3,-\eta}^-(x).$$

Then we have

$$f_n^-(x) - f_{n,\eta}^-(x) = \left( \frac{1}{1-k} \right)^{|x|} \left\{ \frac{k-r}{2k} h_a^-(-s_0) + \frac{k-r}{2r} h_a^-(s_0) (1-k)^{|a|-1} (1-r)^{-|a|+1} \right\}.$$

We see from the definition of  $h_a^-(s)$  that

$$h_a^-(-s_0) = (1-k)^{|a|} F_n^-(s_0, a), \quad h_a^-(s_0) = \frac{r(1-k)(1-r)^{|a|-1}}{k} F_n^-(s_0, a).$$

We therefore obtain

$$f_n^-(x) - f_{n,\eta}^-(x) = \left( \frac{1}{1-k} \right)^{|x|} \frac{k-r}{k} (1-k)^{|a|} F_n^-(s_0, a) = \frac{k-r}{k} F_n^-(s_0, x).$$

By contour integration arguments, we see that there exists  $J_n^- \in C_N(\mathfrak{X})$  such that

$$f_n^-(x) - J_n^-(x) = \frac{(k-r)_+}{k} F_n^-(s_0, x),$$

concluding the proof. □

Summarizing the arguments in this section, we arrive at the following theorem.

**Theorem 6.5.** *Let  $N \in \mathbb{Z}_{>0}$  and  $F \in \mathcal{T}_N(\mathbb{T} \times \mathfrak{X})$ . We set*

$$f(x) = c_G \int_{\mathbb{T}} F(s, x) |c(s)|^{-2} ds.$$

*Then there exists a function  $J \in C_N(\mathfrak{X})$  such that*

$$f(x) - J(x) = \frac{(k-r)_+}{k} F(s_0, x).$$

*Proof.* Let  $x \in \mathfrak{X}$  be such that  $|x| > N$ . We choose a positive integer  $M$  so that  $|x| \leq M$ . Then  $f(x)$  can be written as the following finite sum:

$$f(x) = \varepsilon_M f(x) = f_0(x) + f_1(x) + \cdots + f_M(x).$$

Applying Propositions 6.3 and 6.4 to each  $F_n$ , we have

$$\sum_{n=0}^M f_n(x) - \sum_{n=0}^{\min(M,N)} J_n(x) = \frac{(k-r)_+}{k} \sum_{n=0}^{\min(M,N)} F_n(s_0, x).$$

We thus have the required result. □

In the remainder of this section, as a corollary of Theorem 6.5, we shall prove the Paley–Wiener theorem of the Helgason–Fourier transform.

Let  $N \in \mathbb{Z}_{\geq 0}$ . Denote by  $\mathcal{Z}_N(\mathbb{T} \times \Omega)$  the set of all functions  $F$  on  $\mathbb{T} \times \Omega$  satisfying the following conditions:

- (H1)  $F(s, \omega)$  is a smooth function on  $\mathbb{T}$  with respect to the variable  $s$ ,
- (H2)  $F(s + \tau, \omega) = F(s, \omega)$ ,
- (H3)  $F(s, \omega)$  extends to a  $\tau$ -periodic entire function of exponential type  $N$ ,
- (H4)  $F$  satisfies the symmetry condition (5-2),
- (H5)  $(D_n F)^-(s_0, \omega) = 0$  when  $n > N$ .

With the notation above, we show the following theorem.

**Theorem 6.6.** *Let  $N \in \mathbb{Z}_{\geq 0}$ ,  $F \in \mathcal{Z}_N(\mathbb{T} \times \Omega)$  and set*

$$f(x) = c_G \int_{\mathbb{T}} \int_{\Omega} F(s, \omega) p(x, \omega)^{1/2-is} ds dv(\omega).$$

*Then there exists a function  $J \in C_N(\mathfrak{X})$  such that*

$$f(x) - J(x) = \frac{(k-r)_+}{k} \int_{\Omega} F(s_0, \omega) p(x, \omega)^{1/2-is_0} dv(\omega).$$

*Proof.* Let  $F \in \mathcal{Z}_N(\mathbb{T} \times \Omega)$ . It suffices to show  $P^{-s} F \in \mathcal{T}_N(\mathbb{T} \times \mathfrak{X})$ . The conditions (H1), (H2) and (H6) are immediate from the conditions (H1), (H2) and (H5). Noting

$$|q^{-n/2} \psi(n+1, s)| \leq \frac{q+1}{q-1} q^{n|\Re s|}, \quad |q^{-n/2} \psi^-(n+1, s)| \leq \frac{q+1}{q-1} q^{n|\Im s|},$$

we have from (4-10) that

$$\begin{aligned} |Q_n(-s)^{-1} (P^{-s} F)_n^+(s, x)| &= |q^{-(|x|-n)/2} \psi(|x|-n+1, s) (D_n F)^+(s, \omega)| \\ &\leq C'_N q^{(|x|-n+N)|\Im s|} \end{aligned}$$

for some constant  $C'_N$  which does not depend on the choice of  $n$ . The condition (N5) is obtained in the same fashion as above. This concludes the proof. □

## References

- [Betori and Pagliacci 1984] W. Betori and M. Pagliacci, “Harmonic analysis for groups acting on trees”, *Boll. Un. Mat. Ital. B* (6) **3**:2 (1984), 333–349. MR Zbl
- [Bray 1996] W. O. Bray, “Generalized spectral projections on symmetric spaces of noncompact type: Paley–Wiener theorems”, *J. Funct. Anal.* **135**:1 (1996), 206–232. MR Zbl
- [Campoli 1980] O. A. Campoli, “Paley–Wiener type theorems for rank-1 semisimple Lie groups”, *Rev. Un. Mat. Argentina* **29**:3 (1980), 197–221. MR Zbl
- [Cowling and Setti 1999] M. Cowling and A. G. Setti, “The range of the Helgason–Fourier transformation on homogeneous trees”, *Bull. Austral. Math. Soc.* **59**:2 (1999), 237–246. MR Zbl
- [Cowling et al. 1998] M. Cowling, S. Meda, and A. G. Setti, “An overview of harmonic analysis on the group of isometries of a homogeneous tree”, *Exposition. Math.* **16**:5 (1998), 385–423. MR Zbl
- [Faraut and Picardello 1984] J. Faraut and M. A. Picardello, “The Plancherel measure for symmetric graphs”, *Ann. Mat. Pura Appl.* (4) **138** (1984), 151–155. MR Zbl
- [Iozzi and Picardello 1983a] A. Iozzi and M. A. Picardello, “Graphs and convolution operators”, pp. 187–208 in *Topics in modern harmonic analysis, I* (Turin/Milan, 1982), edited by F. Ricci, Istituto Nazionale di Alta Matematica, Rome, 1983. MR
- [Iozzi and Picardello 1983b] A. Iozzi and M. A. Picardello, “Spherical functions on symmetric graphs”, pp. 344–386 in *Harmonic analysis* (Cortona, 1982), edited by F. Ricci and G. Weiss, Lecture Notes in Math. **992**, Springer, Berlin, 1983. MR
- [Jamal Eddine 2013] A. Jamal Eddine, *Equations d’évolution sur certains groupes hyperboliques*, Thèse de doctorat, Université d’Orléans, 2013, <http://tinyurl.com/ajeddine>.
- [Jamal Eddine 2015] A. Jamal Eddine, “An overview of harmonic analysis and the shifted wave equation on symmetric graphs”, preprint, 2015. arXiv
- [Johnson 1979] K. D. Johnson, “Paley–Wiener theorems on groups of split rank one”, *J. Funct. Anal.* **34**:1 (1979), 54–71. MR Zbl
- [Koizumi 2013] S. Koizumi, “An analogue of the spectral projection for homogeneous trees”, *Hiroshima Math. J.* **43**:2 (2013), 207–221. MR Zbl
- [Kuhn and Soardi 1983] G. Kuhn and P. M. Soardi, “The Plancherel measure for polygonal graphs”, *Ann. Mat. Pura Appl.* (4) **134** (1983), 393–401. MR Zbl
- [Mantero and Zappa 1983] A. M. Mantero and A. Zappa, “The Poisson transform and representations of a free group”, *J. Funct. Anal.* **51**:3 (1983), 372–399. MR Zbl

Received August 28, 2016. Revised January 29, 2017.

SHIN KOIZUMI

FACULTY OF ECONOMY, MANAGEMENT AND INFORMATION SCIENCE

ONOMICHI CITY UNIVERSITY

1600 HISAYAMADA-CHO

ONOMICHI-SHI 722-8506

JAPAN

koizumi\_qup\_shin@live.jp





# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Igor Pak  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pak.pjm@gmail.com](mailto:pak.pjm@gmail.com)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 290    No. 1    September 2017

---

The Vietoris–Rips complexes of a circle	1
MICHAŁ ADAMASZEK and HENRY ADAMS	
A tale of two Liouville closures	41
ALLEN GEHRET	
Braid groups and quiver mutation	77
JOSEPH GRANT and BETHANY R. MARSH	
Paley–Wiener theorem of the spectral projection for symmetric graphs	117
SHIN KOIZUMI	
Fundamental domains of arithmetic quotients of reductive groups over number fields	139
LEE TIM WENG	
Growth and distortion theorems for slice monogenic functions	169
GUANGBIN REN and XIEPING WANG	
Remarks on metaplectic tensor products for covers of $GL_r$	199
SHUICHIRO TAKEDA	
On relative rational chain connectedness of threefolds with anti-big canonical divisors in positive characteristics	231
YUAN WANG	
An orthogonality relation for spherical characters of supercuspidal representations	247
CHONG ZHANG	