

*Pacific  
Journal of  
Mathematics*

ON HOMOLOGY THEORIES OF CUBICAL DIGRAPHS

ALEXANDER GRIGOR'YAN AND YURI MURANOV

Volume 322 No. 1

January 2023



## ON HOMOLOGY THEORIES OF CUBICAL DIGRAPHS

ALEXANDER GRIGOR'YAN AND YURI MURANOV

**We prove the equivalence of the singular cubical homology and the path homology on the category of cubical digraphs. As a corollary we obtain a new relation between the singular cubical homology of digraphs and simplicial homology.**

### 1. Introduction

The path homology theory and the singular cubical homology theory for the category of digraphs were introduced in [1; 2; 3; 4; 5]. In this category, there is a natural mapping of the cubical homology theory to the path homology theory, that induces an isomorphism of homology groups in dimensions 0 and 1. However, in [5] an example of a digraph was constructed, for which the path homology is trivial in dimension 2 while the singular cubical homology is nontrivial in this dimension. Hence, in general, these two theories give different homologies in dimensions  $\geq 2$ . A natural question arises whether these two theories are equivalent on some subclass of digraphs.

In this paper we present a class of cubical digraphs and prove the equivalence of the singular cubical homology and the path homology theories on this class. As the main technical tool for that, we prove that the image of every map of a digraph cube to a cubical digraph is contractible.

The paper is organized as follows. In Section 2, we recall the basic definitions from graph theory and describe some properties of singular cubical homology  $H_*^c$  and the path homology  $H_*$  on the category of digraphs using [1; 2; 3; 5]. In Section 3, we recall the definition of cubical digraph from [2] and prove the contractibility of the image of a digraph cube in a cubical digraph for any digraph map. In Section 4, we prove the main result of the paper:

**Theorem 1.1.** *On the category of cubical digraphs, the singular cubical homology theory is equivalent to the path homology theory.*

---

The authors were supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - SFB 1283/2 2021 - 317210226.

*MSC2020:* primary 18G85, 55N10, 55N35, 57M15; secondary 05C20, 05C25, 05C38.

*Keywords:* homology of digraphs, equivalence of homology theories, path homology theory, singular cubical homology theory, cubical digraph, paths in digraphs.

In Corollary 4.6 we obtain a consequence about the relation between the singular cubical homology theory of digraphs and simplicial homology.

## 2. Singular cubical and path homology theories

In this section we give necessary preliminary material about digraphs and homology theories on the category of finite digraphs.

**Definition 2.1.** A *digraph*  $G$  is a pair  $(V_G, E_G)$  of a set  $V = V_G$  of *vertices* and a subset  $E_G \subset \{V_G \times V_G \setminus \text{diagonal}\}$  of ordered pairs  $(v, w)$  of vertices which are called *arrows* and are denoted  $v \rightarrow w$ . The vertex  $v = \text{orig}(v \rightarrow w)$  is called the *origin* of the arrow and the vertex  $w = \text{end}(v \rightarrow w)$  is called the *end* of the arrow.

For two vertices  $v, w \in V_G$ , we write  $v \cong w$  if either  $v = w$  or  $v \rightarrow w$ .

A *subgraph*  $H$  of a digraph  $G$  is a digraph whose set of vertices is a subset of that of  $G$  and the set edges of  $H$  is a subset of the set of edges of  $G$ . In this case we write  $G \supset H$ .

A subgraph  $H$  of  $G$  is called *induced* if the edges of  $H$  are all those edges of  $G$  whose adjacent vertices belong to  $H$ . In this case we write  $G \sqsupset H$ .

A *directed path*  $p = (a_1, \alpha_1, a_2, \alpha_2, \dots, \alpha_n, a_{n+1})$  in a digraph  $G$  is a sequence of vertices  $a_i$  and arrows  $\alpha_i$  such that  $\alpha_i = (a_i \rightarrow a_{i+1})$ . The number  $n$  of arrows in path is called *length* of the path and is denoted by  $|p|$ . The vertex  $a_1$  is called the *origin* of the path and the vertex  $a_{n+1}$  is called the *end* of the path.

**Definition 2.2.** A *digraph map* (or simply *map*) from a digraph  $G$  to a digraph  $H$  is a map  $f : V_G \rightarrow V_H$  such that  $v \cong w$  in  $G$  implies  $f(v) \cong f(w)$  in  $H$ .

A digraph map  $f$  is *nondegenerate* if  $v \rightarrow w$  in  $G$  implies  $f(v) \rightarrow f(w)$  in  $H$ .

The set of all digraphs with digraph maps form the *category* of digraphs that will be denoted by  $\mathcal{D}$ .

**Definition 2.3.** For two digraphs  $G$  and  $H$ , the *box product*  $\Pi = G \square H$  is defined as a digraph with a set of vertices  $V_\Pi = V_G \times V_H$  and a set of arrows  $E_\Pi$  given by the rule

$$(x, y) \rightarrow (x', y') \quad \text{if } x = x' \text{ and } y \rightarrow y', \text{ or } x \rightarrow x' \text{ and } y = y',$$

where  $x, x' \in V_G$  and  $y, y' \in V_H$ .

Fix  $n \geq 0$ . Denote by  $I_n$  any digraph with the set of vertices  $V = \{0, 1, \dots, n\}$  such that, for  $i = 0, 1, \dots, n-1$ , there is exactly one arrow  $i \rightarrow i+1$  or  $i+1 \rightarrow i$  and there are no other arrows. Such a digraph is called a *line* digraph. It is called a *direct line* digraph if, additionally, all arrows have the form  $i \rightarrow i+1$ . We denote the digraph  $0 \rightarrow 1$  by  $I$ .

For any  $n \geq 0$ , define a *standard  $n$ -cube digraph*  $I^n$  as follows. For  $n = 0$  we put  $I^0 = \{0\}$  which is an one-vertex digraph. For  $n \geq 1$ , the set of vertices of  $I^n$

consists of all  $2^n$  binary sequences  $a = (a_1, \dots, a_n)$ , and there is an arrow  $a \rightarrow b$  between two such vertices if and only if the sequence  $b = (b_1, \dots, b_n)$  is obtained from  $a = (a_1, \dots, a_n)$  by replacing a digit 0 by 1 at exactly one position. It is easy to see that

$$I^n = \underbrace{I \square I \square I \square \dots \square I}_{n \text{ times}}$$

For example, the digraph  $0 \rightarrow 1$  is an 1-cube. Any digraph that is isomorphic to  $I^2$  will be referred to as a *square*. Any digraph that is isomorphic to  $I_n$  and isomorphic to the standard  $n$ -cube will be referred to as an  *$n$ -cube digraph*.

Let us recall the notion of homotopy in the category of digraphs that was introduced in [1].

**Definition 2.4.** Two digraph maps  $f, g : G \rightarrow H$  are called *homotopic* if there exists a line digraph  $I_n$  with  $n \geq 1$  and a digraph map

$$F : G \square I_n \rightarrow H$$

such that

$$F|_{G \square \{0\}} = f \quad \text{and} \quad F|_{G \square \{n\}} = g,$$

where we identify  $G \square \{0\}$  and  $G \square \{n\}$  with  $G$  in a natural way. In this case we shall write  $f \simeq g$ . The map  $F$  is called a *homotopy* between  $f$  and  $g$ .

In the case  $n = 1$  we refer to the map  $F$  as an *one-step homotopy*.

**Definition 2.5.** Digraphs  $G$  and  $H$  are called *homotopy equivalent* if there exist digraph maps

$$f : G \rightarrow H, \quad g : H \rightarrow G$$

such that

$$f \circ g \simeq \text{id}_H, \quad g \circ f \simeq \text{id}_G.$$

In this case we shall write  $H \simeq G$  and the maps  $f$  and  $g$  are called *homotopy inverses* of each other.

A digraph  $G$  is called *contractible* if  $G \simeq \{*\}$  where  $\{*\}$  is an one-vertex digraph.

**Definition 2.6** [1, Definition 3.4]. Let  $G$  be a digraph and  $H$  be its subgraph.

- (i) A *retraction* of  $G$  onto  $H$  is a map  $r : G \rightarrow H$  such that  $r|_H = \text{id}_H$ .
- (ii) A retraction  $r : G \rightarrow H$  is called a *deformation retraction* if  $i \circ r \simeq \text{id}_G$ , where  $i : H \rightarrow G$  is the natural inclusion.

**Proposition 2.7** [1, Corollary 3.7]. *Let  $r : G \rightarrow H$  be a retraction of a digraph  $G$  onto a subdigraph  $H$  and*

$$(2-1) \quad x \rightrightarrows r(x) \quad \text{for all } x \in V_G \quad \text{or} \quad r(x) \rightrightarrows x \quad \text{for all } x \in V_G.$$

*Then  $r$  is a deformation retraction, the digraphs  $G$  and  $H$  are homotopy equivalent, and  $i, r$  are the homotopy inverses of each other.*

Now we recall the definitions of path homology groups from [2] with the group of coefficients  $\mathbb{Z}$ . An *elementary  $p$ -path* on a finite set  $V$  is any (ordered) sequence  $i_0, \dots, i_p$  of  $p + 1$  vertices of  $V$  that will be denoted by  $e_{i_0 \dots i_p}$ . By  $\Lambda_p = \Lambda_p(V)$  we denote the free abelian group generated by all elementary  $p$ -paths  $e_{i_0 \dots i_p}$ . The elements of  $\Lambda_p$  are called  *$p$ -paths*. Thus, each  $p$ -path  $v \in \Lambda_p$  has the form

$$v = \sum_{i_0, \dots, i_p \in V} v^{i_0 i_1 \dots i_p} e_{i_0 i_1 \dots i_p},$$

where  $v^{i_0 i_1 \dots i_p} \in \mathbb{Z}$  are the coefficients of  $v$ .

For  $p \geq 0$ , define the *boundary operator*  $\partial : \Lambda_{p+1} \rightarrow \Lambda_p$  on basic elements by

$$(2-2) \quad \partial e_{i_0 \dots i_{p+1}} = \sum_{q=0}^{p+1} (-1)^q e_{i_0 \dots \hat{i}_q \dots i_{p+1}},$$

where  $\hat{k}$  means omission of the corresponding index, and extend  $\partial$  to  $\Lambda_{p+1}$  by linearity. Set also  $\Lambda_{-1} = \{0\}$  and define  $\partial : \Lambda_0 \rightarrow \Lambda_{-1}$  by  $\partial v = 0$  for all  $v \in \Lambda_0$ . It follows from this definition that  $\partial^2 v = 0$  for any  $p$ -path  $v$ .

An elementary  $p$ -path  $e_{i_0 \dots i_p}$  for  $p \geq 1$  is called *regular* if  $i_k \neq i_{k+1}$  for all  $k$ . For  $p \geq 1$ , let  $\mathcal{I}_p$  be the subgroup of  $\Lambda_p$  that is spanned by all irregular  $e_{i_0 \dots i_p}$  and we set  $\mathcal{I}_0 = \mathcal{I}_{-1} = 0$ . Then  $\partial \mathcal{I}_{p+1} \subset \mathcal{I}_p$  for  $p \geq -1$ . Consider the chain complex  $\mathcal{R}_*$  with

$$\mathcal{R}_p = \mathcal{R}_p(V) = \Lambda_p / \mathcal{I}_p$$

and with the chain map that is induced by  $\partial$ .

Now we define allowed paths on a digraph  $G = (V, E)$ . A regular elementary path  $e_{i_0 \dots i_p}$  in  $V$  is called *allowed* if  $i_{k-1} \rightarrow i_k$  for any  $k = 1, \dots, p$ , and *nonallowed* otherwise. For  $p \geq 1$ , denote by  $\mathcal{A}_p = \mathcal{A}_p(G)$  the subgroup of  $\mathcal{R}_p$  spanned by the allowed elementary  $p$ -paths, that is,

$$\mathcal{A}_p = \text{span}\{e_{i_0 \dots i_p} : i_0 \dots i_p \text{ is allowed}\}$$

and set  $\mathcal{A}_{-1} = 0$ . The elements of  $\mathcal{A}_p$  are called *allowed  $p$ -paths*.

Consider the following subgroup of  $\mathcal{A}_p$  for  $p \geq 0$ :

$$(2-3) \quad \Omega_p = \Omega_p(G) = \{v \in \mathcal{A}_p : \partial v \in \mathcal{A}_{p-1}\}.$$

The elements of  $\Omega_p$  are called  *$\partial$ -invariant  $p$ -paths*. It is easy to see that  $\partial \Omega_{p+1} \subset \Omega_p$  so that we obtain a chain complex

$$(2-4) \quad 0 \longleftarrow \Omega_0 \xleftarrow{\partial} \Omega_1 \xleftarrow{\partial} \dots \xleftarrow{\partial} \Omega_{p-1} \xleftarrow{\partial} \Omega_p \xleftarrow{\partial} \dots$$

The *path homology groups*  $H_*(G)$  of the digraph  $G$  are defined as the homology groups of the chain complex (2-4), that is,

$$H_p(G) := \ker \partial|_{\Omega_p} / \text{Im } \partial|_{\Omega_{p+1}}.$$

In what follows we will also need a natural augmentation  $\varepsilon : \Omega_0 \rightarrow \mathbb{Z}$  defined by

$$\varepsilon\left(\sum k_i e_i\right) = \sum k_i, \quad k_i \in \mathbb{Z}.$$

Clearly,  $\varepsilon$  is an epimorphism and  $\varepsilon \circ \partial = 0$ .

Now we recall from [5] the construction of the cubical singular homology theory of digraphs.

**Definition 2.8.** A singular  $n$ -cube in a digraph  $G$  is a digraph map  $\phi : I^n \rightarrow G$ .

Fix  $n \geq 1$ . For any  $1 \leq j \leq n$  and  $\epsilon = 0, 1$ , define the inclusion  $F_{j\epsilon}^{n-1} : I^{n-1} \rightarrow I^n$  of digraphs as follows: if  $n \geq 2$ , then

$$(2-5) \quad F_{j\epsilon}^{n-1}(c_1, \dots, c_{n-1}) = \begin{cases} (\epsilon, c_1, \dots, c_{n-1}) & \text{for } j = 1, \\ (c_1, \dots, c_{j-1}, \epsilon, c_j, \dots, c_{n-1}) & \text{for } 1 < j < n, \\ (c_1, \dots, c_{n-1}, \epsilon) & \text{for } j = n \end{cases}$$

and if  $n = 1$ , then  $F_{1\epsilon}^{n-1}(0) = (\epsilon)$ . We shall write shortly  $F_{j\epsilon}$  instead of  $F_{j\epsilon}^{n-1}$  if the dimension  $n - 1$  is clear from the context. Denote by  $I_{j\epsilon}^{n-1}$  the image of  $F_{j\epsilon}^{n-1}$ . We shall write  $I_{j\epsilon}$  instead of  $I_{j\epsilon}^{n-1}$  if the dimension is clear from the context.

Let  $Q_{-1} = 0$ . For  $n \geq 0$ , denote  $Q_n = Q_n(G)$  the free abelian group generated by all singular  $n$ -cubes in  $G$ , and denote by  $\phi^\square$  the singular  $n$ -cube  $\phi$  as the element of the group  $Q_n$ . For  $n \geq 1$  and  $1 \leq p \leq n$ , denote

$$(2-6) \quad \phi_{p\epsilon}^\square = (\phi \circ F_{p\epsilon})^\square \in Q_{n-1}.$$

For any  $n \geq 1$ , define a homomorphism  $\partial^c : Q_n \rightarrow Q_{n-1}$  on the basis elements  $\phi^\square$  by the rule

$$(2-7) \quad \partial^c \phi^\square = \sum_{p=1}^n (-1)^p (\phi_{p0}^\square - \phi_{p1}^\square),$$

and  $\partial^c = 0$  for  $n = 0$ . Then  $(\partial^c)^2 = 0$  and the groups  $Q_n(G)$  form a chain complex that we denote  $Q_* = Q_*(G)$ .

For  $n \geq 1$  and  $1 \leq p \leq n$ , consider the natural projection  $T^p : I^n \rightarrow I^{n-1}$  on the  $p$ -face  $I^{n-1}$  defined as follows. For  $n = 1$ ,  $T^1$  is the unique digraph map  $I^1 \rightarrow I^0$ . For  $n \geq 2$ , we have on the set of vertices

$$T^p(i_1, \dots, i_n) = (i_1, \dots, i_{p-1}, i_{p+1}, \dots, i_n).$$

The singular  $n$ -cube  $\phi : I^n \rightarrow G$  is degenerate if there is  $1 \leq p \leq n$  such that  $\phi = \psi \circ T^p$  where  $\psi : I^{n-1} \rightarrow G$  is a singular  $(n - 1)$ -cube. Then an abelian group  $B_n = B_n(G)$  that is generated by all degenerated  $n$ -cubes is a subgroup  $Q_n$  for  $n \geq 1$ . We put also  $B_0 = 0$ ,  $B_{-1} = 0$ . Then the quotient group

$$(2-8) \quad \Omega_p^c(G) = Q_p(G)/B_p(G)$$

is defined for  $p \geq 0$ . We have  $\partial(B_n) \subset B_{n-1}$ , and hence  $B_*(G) \subset Q_*(G)$ . Hence the quotient complex  $\Omega_*^c(G) = Q_*(G)/B_*(G)$  is defined. We continue to denote the boundary operator in this complex  $\partial^c$ . The homology group  $H_k(\Omega_*^c(G))$  is called the *singular cubical homology group of digraph  $G$  in dimension  $k$*  and is denoted  $H_k^c(G)$ .

We have a natural augmentation homomorphism  $\varepsilon : \Omega_0^c(G) \rightarrow \mathbb{Z}$ , defined by

$$\varepsilon\left(\sum k_i \phi_i\right) = \sum k_i, \quad k_i \in \mathbb{Z}.$$

Then  $\varepsilon$  is an epimorphism and  $\varepsilon \circ \partial^c = 0$ .

Here are some basic properties of the path and the singular cubical homology groups from [2] and [5].

- The groups  $H_*^c(X)$  and  $H_*(X)$  are functors from the category  $\mathcal{D}$  to the category of abelian groups.
- Let  $f \simeq g : X \rightarrow Y$  be two homotopic digraph maps. Then the induced homomorphisms  $f_*$ ,  $g_*$  of homology groups are equal for  $k \geq 0$  for the both theories.

### 3. Maps from cube to cubical digraph

In this section we slightly reformulate the definition of a cubical digraph from [2] and prove Theorem 3.6 saying that an image of a cube in a cubical digraph is contractible.

Recall that any vertex of a cube  $I^n$  is given by a sequence of binary numbers  $(a_1, \dots, a_n)$ . For any arrow  $a \rightarrow b$  in a digraph cube  $I^n$  we have also the arrow

$$(3-1) \quad \gamma_i = (0, \dots, 0) \rightarrow (b_1 - a_1, \dots, b_n - a_n)$$

in  $I^n$  where the right sequence represents a vertex in  $I^n$  that has only one nontrivial element 1 at some position  $i$ . We say that two arrows  $\alpha = (a \rightarrow b)$  and  $\beta = (c \rightarrow d)$  of  $I^n$  are *parallel* and write  $\alpha \parallel \beta$  if


$$(b_1 - a_1, \dots, b_n - a_n) = (d_1 - c_1, \dots, d_n - c_n).$$

In the opposite case we say that the arrows  $\alpha$  and  $\beta$  are *orthogonal*.

An arrow  $\alpha \in E_{I^n}$  defines two  $(n-1)$ -faces of  $I^n$ : the face  $I_0 = I_0^\alpha$  which contains the origin vertices of the arrows that are parallel to  $\alpha$  and the face  $I_1 = I_1^\alpha$  which contains the end vertices of the arrows that are parallel to  $\alpha$ . Note that any arrow that is orthogonal to  $\alpha$  lies in  $I_0$  or in  $I_1$ .

For the digraph cube  $I^n$ , there is a natural partial order on the set of its vertices  $V_{I^n}$  that is defined as follows: we write  $a \leq b$  if there exists a path along the arrows with the origin vertex  $a$  and the end vertex  $b$ . Now we introduce the *distance*  $\Delta(a, b)$  for a pair of vertex  $a, b \in I^n$  that is defined only for comparable pair of vertices.





**Figure 1.** The map  $f : I^3 \rightarrow G$  with noncontractible image.

Let  $a, b$  be two vertices of  $I^n$  such that  $a \leq b$ . As it follows from the definition of  $I^n$ , the length of the path  $p$  from  $a$  to  $b$  does not depend on the choice of the path, and we set

$$\Delta(a, b) = \Delta(b, a) := |p|.$$

We shall refer to the vertex  $a = (0, \dots, 0)$  of a cube as the *origin vertex* and to the vertex  $d = (1, \dots, 1)$  as the *end vertex*.

It follows immediately from the definition of  $I^n$  that, for any vertex  $x$ , the distances  $\Delta(a, x)$  and  $\Delta(x, d)$  are well defined. For an arrow  $\alpha = (x \rightarrow y)$  we define  $\Delta(\alpha, d) := \Delta(y, d)$ .

Let  $a \leq b$  be a pair of comparable vertices of  $I^n$ . Denote by  $I_{a,b}$  the induced subgraph of  $I^n$  with the set of vertices  $\{c \in V_{I^n} \mid a \leq c \leq b\}$ . Clearly,  $I_{a,b}$  is isomorphic to a digraph cube  $I^k$ , where  $k = |p| = \Delta(a, b)$ .

**Definition 3.1.** A subgraph  $G$  of  $I^n$  is called *cubical* if, for any two vertices  $a, b \in V_G \subset V_{I^n}$  with  $a \leq b$ , we have  $I_{a,b} \sqsubset G$ .

Note that the set of all paths from  $a$  to  $b$  in  $I_{a,b}$  coincides with the set of all paths from  $a$  to  $b$  in  $G$ . It is easy to see that cubical digraphs with digraph maps form a category. Now we prove that the image of a cube  $I^n$  in any cubical digraph is contractible. Note that this statement is not true for general digraphs.

**Example 3.2.** Consider a digraph map  $f$  (see Figure 1) that maps the cube  $I^3$  onto the cycle digraph  $G$  and that is defined by  $f(1) = f(8) = x$ ,  $f(2) = f(3) = f(5) = y$ ,  $f(4) = f(6) = f(7) = z$ . Then the images of this map  $G$  is noncontractible.

Now consider a digraph map  $f : I^n \rightarrow G$  where  $G$  is a cubical digraph. The image  $f(I^n)$  is connected as the image of a connected digraph. Let  $s = (0, \dots, 0) \in V_{I^n}$  be the origin vertex and  $z = (1, \dots, 1) \in V_{I^n}$  be the end vertex of  $I^n$ . Then  $f(s) \in V_G$ ,  $f(z) \in V_G$  and  $f(I^n) \subset I_{f(s), f(z)} \subset G$  where  $I_{f(s), f(z)}$  is isomorphic to

an  $m$ -dimensional cube which we denote  $J = J^m \cong I^m$  where  $m = \Delta(f(s), f(z))$ . Hence, without loss of generality, we can assume that  $G = I_{f(s), f(z)} = J$ , that is,

$$f(s) = (0, \dots, 0) \in V_J, \quad f(z) = d = (1, \dots, 1) \in V_J.$$

For  $m = 0, 1, 2$  the image  $f(I^n) \subset G$  is contractible since all connected subgraphs of the digraphs  $J^0, J^1$ , and  $J^2$  are contractible.

Consider the case  $J = J^m$  where  $m \geq 3$  and  $d = (1, \dots, 1) \in V_J$  is the end vertex of the cube  $J$ . Since  $d = f(z) \in \text{Image}(f)$ , there exists a nonempty set of arrows  $\Gamma \subset E_J$  defined by

$$[\tau \in \Gamma] \Leftrightarrow [\text{end}(\tau) = d \text{ and } \tau = f(\alpha), \alpha \in E_{I^n}].$$

The set  $\Gamma$  consists of arrows in  $E_J$  with the end vertex  $d$  that are lying in the image of the map  $f$ . Let  $\gamma = (c \rightarrow d) \in \Gamma$  be an arrow satisfying

$$(3-2) \quad \begin{cases} f(\alpha) = f(x \rightarrow y) = (c \rightarrow d) = \gamma, \\ \Delta(\alpha, z) = \Delta(y, z) = k \geq 0 \text{ is minimal.} \end{cases}$$

Note that  $\alpha$  is not uniquely defined.

**Lemma 3.3.** *For every vertex  $v \in V_{I^n}$  with  $\Delta(v, z) \leq k$  we have  $f(v) = d$ . Hence the cube  $I_{y,z} \sqsubset I^n$  is mapped by  $f$  into the vertex  $d$ .*

*Proof.* It follows immediately from the definition of  $k$  in (3-2).  $\square$

The arrow  $\gamma$  defines two  $(m-1)$ -dimensional faces  $J_0$  and  $J_1$  of the cube  $J$  with  $c \in V_{J_0}$ ,  $d \in V_{J_1}$  and we have the natural projection  $\pi : J \rightarrow J_0$  along the arrow  $\gamma$ . Let  $H$  be a subgraph of  $I^n$ . We define subgraphs  $K_0, K_1, K \subset J$  that depend on the map  $f : I^n \rightarrow J$  and  $H \subset I^n$  such that

$$(3-3) \quad K := f(H) \subset J, \quad K_0 := f(H) \cap J_0 \subset J_0, \quad \text{and} \quad K_1 := f(H) \cap J_1 \subset J_1.$$

It is easy to see that for an arrow  $(v \rightarrow w) \in E_J$  we have

$$(3-4) \quad [(v \rightarrow w) \parallel \gamma] \Leftrightarrow [(v \in J_0) \text{ and } (w \in J_1)].$$

For technical reasons we introduce the following definition.

**Definition 3.4.** Let  $H$  be a subgraph of  $I^n$  and  $f : I^n \rightarrow J$  be a digraph map. Let the digraphs  $K_0, K_1, K \subset J$  be defined as above using (3-2) and (3-3). We say that the subgraph  $H$  satisfies the  $\Pi$ -condition if the following conditions are satisfied:

- (1) For all  $w \in V_{K_1}$  there is a vertex  $v \in V_{K_0}$  such that  $(v \rightarrow w) \in E_K$ .
- (2) For all  $(w \rightarrow w') \in E_{K_1}$  we have  $\pi(w \rightarrow w') \in E_{K_0}$ .

The next statement is our key technical result.

**Proposition 3.5.** *Consider the map  $f : I^n \rightarrow J = J^m$  with  $m \geq 3$ . Let  $k$  and  $\gamma$  are defined in (3-2). Then the cube  $I^n$  satisfies the  $\Pi$ -condition.*

*Proof.* Using induction on  $k \geq 0$ .

The base of induction,  $k = 0$ . Hence  $y = z = (1, \dots, 1) \in V_{I^n}$  is the end vertex of  $I^n$  and  $n \geq m \geq 3$ . The arrow  $\alpha = (x \rightarrow z) \in E_{I^n}$  with

$$(3-5) \quad f(\alpha) = f(x \rightarrow z) = \gamma = (c \rightarrow d)$$

defines  $(n-1)$ -face  $I_0 = I_{s,x}$  and opposite  $(n-1)$ -face  $I_1$  of the cube  $I^n$ . Let  $a = (0, \dots, 0)$  be the origin vertex of  $J$  (and hence the origin vertex of  $J_0$ ) and  $b$  the origin vertex of  $J_1$ . Then  $a \rightarrow b$  is parallel to  $\gamma = (c \rightarrow d)$ . We have

$$(3-6) \quad f(I_0) = f(I_{s,x}) \subset I_{f(s),f(x)} = I_{a,c} = J_0$$

and hence, by (3-3) for  $H = I^n$ , we have  $f(I_0) \subset K_0$ . Let  $t$  be a vertex of  $I_1$  such that  $w = f(t) \notin V_{K_0}$ , that is,  $w \in V_{K_1} \subset V_{J_1}$ . There exists an unique vertex  $r \in V_{I_0}$  such that  $(r \rightarrow t) \in E_{I^n}$  is parallel to  $\alpha$  and  $f(r) = v \in K_0 \subset J_0$  by (3-6). Thus  $f(r \rightarrow t) = v \rightarrow w$  with  $v \in V_{K_0}$  and condition (1) of Definition 3.4 is satisfied.

Now let  $\tau = (w \rightarrow w') \in E_{K_1}$  be an arrow such that  $f(t \rightarrow t') = \tau$ , that is,

$$f(t) = w, \quad f(t') = w', \quad t, t' \in V_{I_1}.$$

The same line of arguments as above gives the vertices  $r, r' \in V_{I_0}$  such that  $(r \rightarrow t)$  and  $(r' \rightarrow t')$  are parallel to  $\alpha$  and hence,  $\pi(\tau) = f(r \rightarrow r')$  since  $f(r), f(r') \in V_{K_0}$ . This proves condition (2) of Definition 3.4. Thus  $\Pi$ -condition is satisfied for the cube  $I^n$  and  $k = 0$ .

We now consider the induction step. By inductive assumption we have that any map  $f : I^n \rightarrow J$  satisfies the  $\Pi$ -condition if  $\Delta(y, z) \leq k - 1 \geq 0$ . Consider the case  $\Delta(y, z) = k \geq 1$  and hence

$$\Delta(x, z) = \Delta(y, z) + 1 = k + 1 \geq 2$$

where

$$z = \underbrace{(1, \dots, 1)}_n \in V_{I^n}.$$

Thus, without loss of generality, we can suppose that

$$(3-7) \quad x = \underbrace{(1, \dots, 1)}_{n-k-1} \underbrace{(0, 0, \dots, 0)}_{k+1}, \quad y = \underbrace{(1, \dots, 1)}_{n-k-1} \underbrace{(1, 0, \dots, 0)}_k.$$

From now we put  $y_0 = y \in V_{I^n}$  and let the vertex  $y_i$  is obtained from  $y$  by replacing the last coordinate 1 in  $y$  by 0, and  $i$ -th coordinate 0 of  $y$  by 1 for  $1 \leq i \leq k$ . For example,

$$y_2 = \underbrace{(1, \dots, 1)}_{n-k-1} \underbrace{(0, 0, 1, 0, \dots, 0)}_k, \quad y_k = \underbrace{(1, \dots, 1)}_{n-k-1} \underbrace{(0, 0, 0, \dots, 0, 1)}_k.$$

We also define

$$\alpha_i = (x \rightarrow y_i) \in E_{I^n} \quad \text{for } 0 \leq i \leq k.$$

By Lemma 3.3 we have

$$f(\alpha_i) = f(x \rightarrow y_i) = (c \rightarrow d) = \gamma \quad \text{for } 0 \leq i \leq k.$$

Let  $I_0 = I_{s,x}$  be  $(n - k - 1)$ -dimensional subcube of  $I^n$ . Then, as before, we have

$$f(I_0) \subset K_0 \subset J_0.$$

Consider a vertex  $t \in V_{I^n}$  and  $t \notin V_{I_0}$  that has the form

$$t = (a_1, \dots, a_{n-k-1}, b_0, \dots, b_k) \notin I_0, \quad a_i, b_j \in \{0, 1\},$$

where at least one coordinate  $b_j$  is 1. If at least one coordinate  $b_j$  is 0 we obtain that  $t \in I_{s,z_j} \sqsubset I^n$  where

$$z_j = (\underbrace{1, \dots, 1}_{n-k-1}, \underbrace{1, \dots, \overset{j}{0}, \dots, 1}_{k+1}).$$

The  $(n - 1)$ -dimensional subcube  $I_{s,z_j} \subset I^n$  contains the vertices  $x$  and  $t$ . Moreover,  $\Delta(x, z_j) = k$  and there is an arrow

$$\alpha_i = (x \rightarrow y_i) \in E_{I_{s,z_j}}$$

with

$$f(\alpha_i) = \gamma \quad \text{and} \quad \Delta(\alpha_i, z_j) = k - 1.$$

Hence, by the inductive assumption, the map

$$f|_{I_{s,z_j}} : I_{s,z_j} \rightarrow J$$

satisfies the  $\Pi$ -condition. Hence the conditions (1) and (2) of Definition 3.4 are satisfied for every  $(n - 1)$ -dimensional subcube  $I_{s,z_j} \subset I^n$ .

Now consider a vertex  $t$  for which all  $(k + 1)$ -coordinates  $b_j$  are equal to 1 such that  $t \notin I_{x,z}$ . This means that at least one of the first  $(n - k - 1)$ -coordinates  $a_i$  is 0. Recall that  $(k + 1) \geq 2$ . Thus, consider the vertices

$$(3-8) \quad t = (a_1, \dots, a_{n-k-1}, \underbrace{1, \dots, 1}_{k+1}) \notin I_0, \quad r = (a_1, \dots, a_{n-k-1}, \underbrace{0, \dots, 0}_{k+1}) \in I_0,$$

where  $a_i \in \{0, 1\}$ . Consider a directed path  $p$  in the digraph  $I_0$  from the vertex  $r \in V_{I_0}$  to the vertex  $x \in V_{I_0}$  of the length  $l = |p| \geq 1$  (since  $t \notin I_{x,z}$ ). Write this path in the form

$$p = (r \rightarrow x_1 \rightarrow x_2 \rightarrow \dots \rightarrow x_{l-1} \rightarrow x_l = x) \subset I_{r,x} \subset I_0.$$

Consider a directed path  $q$  from the vertex  $r \in V_{I_0}$  to the vertex  $t$  of the length

$$k + 1 = |q| \geq 2.$$

Note that  $q$  lies in the digraph  $I_{r,t}$  of dimension  $k + 1$ . Write this path in the form

$$q = (r \rightarrow r^1 \rightarrow r^2 \rightarrow \dots \rightarrow r^k \rightarrow r^{k+1} = t) \subset I_{r,t}.$$

Any such two paths  $p$  and  $q$  define a unique subgraph of the digraph  $I^n$  that has the form

$$(3-9) \quad \begin{array}{ccccccccc} t = r^{k+1} & \longrightarrow & r_1^{k+1} & \longrightarrow & r_2^{k+1} & \longrightarrow & \dots & \longrightarrow & r_l^{k+1} = z \\ \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ r^k & \longrightarrow & r_1^k & \longrightarrow & r_2^k & \longrightarrow & \dots & \longrightarrow & r_l^k \\ \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \dots & \longrightarrow & \dots & \longrightarrow & \dots & \longrightarrow & \dots & \longrightarrow & \dots \\ \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ r^1 & \longrightarrow & r_1^1 & \longrightarrow & r_2^1 & \longrightarrow & \dots & \longrightarrow & r_l^1 \\ \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ r & \longrightarrow & x_1 & \longrightarrow & x_2 & \longrightarrow & \dots & \longrightarrow & x_l = x \end{array}$$

Now using induction in the length  $l = |p| \geq 1$  we prove the following statement.

**Statement A.** For every path  $q$  and every path  $p$ , as above, there is a path

$$p' = (r \rightarrow x'_1 \rightarrow x'_2 \rightarrow \dots \rightarrow x'_{l-1} \rightarrow x'_l = x) \subset I_{r,x} \subset I_0$$

(that may be equal to  $p$ ) such that  $q$  and  $p'$  defines the subgraph (similarly as above)

$$(3-10) \quad \begin{array}{ccccccccc} t = r^{k+1} & \longrightarrow & r_1^{k+1'} & \longrightarrow & r_2^{k+1'} & \longrightarrow & \dots & \longrightarrow & r_l^{k+1'} = z \\ \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ r^k & \longrightarrow & r_1^{k'} & \longrightarrow & r_2^{k'} & \longrightarrow & \dots & \longrightarrow & r_l^{k'} \\ \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \dots & \longrightarrow & \dots & \longrightarrow & \dots & \longrightarrow & \dots & \longrightarrow & \dots \\ \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ r^1 & \longrightarrow & r_1^{1'} & \longrightarrow & r_2^{1'} & \longrightarrow & \dots & \longrightarrow & r_l^{1'} \\ \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ r & \longrightarrow & x'_1 & \longrightarrow & x'_2 & \longrightarrow & \dots & \longrightarrow & x'_l = x \end{array}$$

and at least one of the following conditions is satisfied:

$$(3-11) \quad (i) f(t) = f(r^k), \quad (ii) f(t) = f(r_1^k), \quad (iii) f(t) = f(r_1^{k'}).$$

The base of induction for Statement A, the case  $l = 1$ . Consider the unique path  $p = (r \rightarrow x) \subset I_0$  of the length  $l = 1$  and a path  $q$  as above. We have the following subgraph of the digraph  $I^n$ :

$$(3-12) \quad \begin{array}{ccc} t = r^{k+1} & \longrightarrow & r_1^{k+1} = z \\ \uparrow & & \uparrow \\ r^k & \longrightarrow & r_1^k \\ \uparrow & & \uparrow \\ \vdots & \longrightarrow & \vdots \\ \uparrow & & \uparrow \\ r^1 & \longrightarrow & r_1^1 \\ \uparrow & & \uparrow \\ r & \longrightarrow & x_1 = x \end{array}$$

where

$$r, x \in V_{I_0} \quad \text{and} \quad f(r), f(x) \in V_{K_0},$$

and

$$f(r_i^i) = d \quad \text{for } 1 \leq i \leq k + 1$$

since  $k \geq 1$ . Hence,

$$f(r^k) = f(r_1^{k+1}) = d$$

and thus at least one of the conditions (i) or (ii) in (3-11) is satisfied because there are no triangles in the digraph  $J$ . We put in this case  $p' = p$ , and hence the base of induction  $l = 1$  is proved.

Inductive step of induction for Statement A. Consider vertices  $t, r \in V_J$  given in (3-8) where

$$\Delta(t, r) = k + 1 \geq 2 \quad \text{and} \quad \Delta(r, x) \geq 2.$$

Let  $p$  be a path from  $r$  to  $x$  and  $q$  be a path from  $r$  to  $t$  as the above. Recall that

$$|q| = k + 1 \geq 2 \quad \text{and} \quad |p| = l \geq 2.$$

These paths define the subgraph of  $I^n$  given in (3-9). By the inductive assumption, for the vertex  $r_1^{k+1}$  at least one of the conditions

$$(3-13) \quad (i) f(r_1^{k+1}) = f(r_1^k), \quad (ii) f(r_1^{k+1}) = f(r_2^k), \quad (iii) f(r_1^{k+1}) = f(r_2^{k''}),$$

that is similar to (3-11) is realized. In (3-13) we have a path

$$r^k \rightarrow r_1^k \rightarrow r_2^{k''} \rightarrow \dots \rightarrow r_l^k,$$

that is, from (3-9), similar to the path

$$r^k \rightarrow r_1^k \rightarrow r_2^k \rightarrow \dots \rightarrow r_l^k.$$

If condition (i) is realized, that is,  $f(r_1^{k+1}) = f(r_1^k)$ , then for  $f(t)$  at least one of the conditions (i) or (ii) in (3-11) is satisfied since there are no triangles in the digraph  $J$  (similar to the case  $l = 1$ ).

If condition (ii) is realized and condition (i) is not realized, that is,

$$f(r_1^{k+1}) = f(r_2^k) \quad \text{and} \quad f(r_1^k) \neq f(r_2^k),$$

we can consider the subcube of  $I^n$  given in Figure 2 that is defined by the subgraph of (3-9) given by

$$(3-14) \quad \begin{array}{ccccc} t = r^{k+1} & \longrightarrow & r_1^{k+1} & \longrightarrow & r_2^{k+1} \\ \uparrow & & \uparrow & & \uparrow \\ r^k & \longrightarrow & r_1^k & \longrightarrow & r_2^k \end{array}$$

We have


$$f(r_1^{k+1}) = f(r_2^k) \quad \text{and} \quad f(r_1^k) \neq f(r_2^k),$$

that is,

$$f(r_1^k \rightarrow r_1^{k+1}) = f(r_1^k \rightarrow r_2^k) \in E_J$$

is an arrow. If  $f(r^k) = f(r_1^k)$ , then the same line of above gives that

$$f(t) = f(r_1^k) \quad \text{or} \quad f(t) = f(r_2^k)$$



**Figure 2.** The subcube of  $I^n$  that is defined by the digraph in (3-13).

and the step of induction is proved. Let  $f(r_k) \neq f(r_1^k)$ , then

$$f(I_{r^k, r_2^k}) \subset f(I_{f(r^k), f(r_2^k)}) \quad \text{and} \quad f(I_{r^k, r_1^{k+1}}) \subset f(I_{f(r^k), f(r_2^k)}),$$

where  $I_{f(r^k), f(r_2^k)}$  is the digraph square. Hence at least one of conditions

$$f(r^{k+1}) = f(r_1^k) \quad \text{or} \quad f(r^{k+1}) = f(r_1^{k'})$$

is satisfied and the inductive assumption is proved.

Consider the case when condition (iii) is realized and conditions (i) and (ii) are not realized. This case is the same as the case (ii). We must to start the consideration from the path

$$r^k \rightarrow r_1^k \rightarrow r_2^{k''} \rightarrow \dots \rightarrow r_l^k$$

on the place of the path

$$r^k \rightarrow r_1^k \rightarrow r_2^k \rightarrow \dots \rightarrow r_l^k$$

from (3-9). This finishes the proof of the inductive step as well as Statement A.

Since each of the vertices  $r^k, r_1^k, r_1^{k'}$  lies in the image of a subcube  $I_{r, z_j}$  it follows from Statement A that image  $w = f(t)$  lies in the image of a subcube  $I_{r, z_j}$  with

$$\Delta(x, z_j) = \Delta(r, z_j) = k,$$

which satisfies  $\Pi$ -condition by the inductive assumption in  $k$ . Hence condition (1) of Definition 3.4 is satisfied for every subcube  $I_{r, t} \subset I^n$ . By a similar way, it follows from Statement A that the image of every arrow with end or origin  $t$  lies in the image of a subcube  $I_{r, z_j}$  which satisfies  $\Pi$ -condition by the inductive assumption in  $k$ . Hence condition (2) of Definition 3.4 is satisfied for every subcube  $I_{r, t} \subset I^n$ . Hence every cube  $I_{r, t}$  satisfies the  $\Pi$ -condition, and hence the cube  $I^n$  satisfies the  $\Pi$ -condition. This completes the proof of Proposition 3.5.  $\square$

**Theorem 3.6.** *Let  $f : I^n \rightarrow G$  be a digraph map to a cubical digraph. Then the image  $f(I^n) \subset G$  is contractible.*

*Proof.* The image  $f(I^n)$  lies in the digraph  $J = J^m$ . Now we use the induction in  $m$ . For  $m = 0, 1, 2$  the image  $f(I^n)$  is contractible since all connected subgraphs of  $J$  are contractible. For  $m \geq 3$  the digraph  $I^n$  satisfies the  $\Pi$ -condition, then (2-7) and conditions (1) and (2) of Definition 3.4 imply that restriction  $\pi|_K$  of the projection  $\pi : J^m \rightarrow J_0^{m-1}$  to the image  $K$  of the map  $f$  is well defined deformation retraction to  $K_0$ . But  $K_0$  is contractible by the inductive assumption in  $m$ .  $\square$

#### 4. Equivalence to homology theories on cubical digraphs

In this section we prove our main result, Theorem 1.1, stated below as Theorem 4.5. For that we use the acyclic carrier theorem from homology theory (see, for example, [6, Section 3.4] and [7, Section 1.2.1]). Recall that a chain complex  $C_*$  is called



*nonnegative* if  $C_p = 0$  for  $p < 0$  and is called *free* if  $C_p$  are finitely generated free abelian groups for all  $p$ . We say that  $C_*$  is a *geometric chain complex* if it is nonnegative, free, and if a basis  $\mathcal{B}_p$  is chosen in the group  $C_p$  for any  $p \geq 0$ . For example, any finite simplicial complex gives rise to a geometric chain complex, where  $\mathcal{B}_p$  consists of all  $p$ -simplexes.

Let  $C_*$  be a geometric chain complex with fixed bases  $\mathcal{B}_p$ . For  $b \in \mathcal{B}_{p-1}$  and  $b' \in \mathcal{B}_p$ , we write  $b < b'$  if  $b$  enters with a nonzero coefficient into the expansion of  $\partial b'$  in the basis  $\mathcal{B}_{p-1}$ . The *augmentation homomorphism*  $\varepsilon : C_0 \rightarrow \mathbb{Z}$  is defined by

$$\varepsilon\left(\sum_i k_i b_i\right) = \sum_i k_i, \quad k_i \in \mathbb{Z}, b_i \in \mathcal{B}_0,$$

and by  $\tilde{C}_*$  we denote the augmented complex

$$0 \longleftarrow \mathbb{Z} \xleftarrow{\varepsilon} C_0 \xleftarrow{\partial} C_1 \xleftarrow{\partial} \dots$$

A geometric chain complex  $C_*$  is called *acyclic* if all homology groups of the augmented complex  $\tilde{C}_*$  are trivial.

Let  $C_*$  and  $D_*$  be two geometric complexes with augmentation homomorphism  $\varepsilon$  and  $\varepsilon'$ , respectively. A chain map  $\phi_* : C_* \rightarrow D_*$  is called *augmentation preserving* if  $\varepsilon' \phi_0(c) = \varepsilon(c)$  for any  $c \in C_0$ .

**Definition 4.1.** Let  $C_*$  and  $D_*$  be two geometric chain complexes.

(i) An *algebraic carrier function* from  $C_*$  to  $D_*$  is a mapping  $E$  that assigns to any basis element  $b$  in  $C_*$  a subcomplex  $E_*(b) := E(b)$  of  $D_*$ , such that  $b < b'$  implies  $E_*(b) \subset E_*(b')$ .

(ii) An algebraic carrier function  $E$  is called *acyclic* if each complex  $E_*(b)$  is nonempty and acyclic.

(iii) A chain map  $f_* : C_* \rightarrow D_*$  is *carried by*  $E$  if  $f_n(b) \in E_*(b)$  for any basis element  $b$  in  $C_n$ .

We state the acyclic carrier theorem in the following form.

**Theorem 4.2** [6, Section 3.4; 7, Section 1.2.1]. *Let  $C_*$  and  $D_*$  be two geometric chain complexes and  $E$  be an acyclic carrier function from  $C_*$  to  $D_*$ . If  $f_*, g_* : C_* \rightarrow D_*$  are augmentation preserving chain maps that are carried by  $E$ , then  $f_*$  and  $g_*$  are chain homotopic.*

Before the proof of Theorem 1.1, we state and prove some technical results. We use the notations of [2; 5]. Let  $G$  be a cubical digraph. The free abelian groups  $\Omega_p^c = \Omega_p^c(G)$  and  $\Omega_p = \Omega_p(G)$  defined in (2-3) and (2-8) are finitely generated.

Let  $I^0 = \{*\}$  be the one-vertex digraph. Any 0-dimensional singular cube  $\phi : I^0 = \{*\} \rightarrow G$  is given by the vertex  $\phi(*) \in V_G$  and thus we obtain the map  $\tau_0 : \Omega_0^c(G) \rightarrow \Omega_0(G)$  which preserve augmentation.

For any digraph cube  $I^n$  ( $n \geq 1$ ) we denote by  $P$  the set of all directed paths of the length  $n$  going from the origin vertex

$$\underbrace{(0, \dots, 0)}_n$$

of the cube to the end vertex

$$\underbrace{(1, \dots, 1)}_n.$$

Every path  $p \in P$  has the form

$$(4-1) \quad p = (a_0 \rightarrow a_1 \rightarrow a_2 \rightarrow \dots \rightarrow a_n), \quad a_i \in V_{I^n}.$$

In (4-1) for  $1 \leq i \leq n$  the vertex  $a_i$  differs from  $a_{i-1}$  only by one coordinate  $1 \leq \pi(i) \leq n$  that equals 0 for  $a_{i-1}$  and 1 for  $a_i$ . Let  $\sigma(p)$  be a sign of the permutation

$$\pi(p) = \begin{pmatrix} 1 & 2 & \dots & n \\ \pi(1) & \pi(2) & \dots & \pi(n) \end{pmatrix}.$$

Consider the path  $w_n \in \Omega_n(I^n)$  given by

$$(4-2) \quad w_n = \sum_{p \in P} (-1)^{\sigma(p)} p,$$

which is the generator of the group  $\Omega_n(I^n)$  (see [5] and [2]). For any singular  $n$ -dimensional cube  $\phi : I^n \rightarrow G$ , which gives a basic element  $\phi^\square \in \Omega_n^c(G)$ , we have a morphism of chain complexes defined in [5]

$$(4-3) \quad \tau_* : \Omega_*^c(G) \rightarrow \Omega_*(G), \quad \tau_n(\phi^\square) := \phi_*(w_n),$$

where  $\phi_* : \Omega_*(I^n) \rightarrow \Omega_*(G)$  is the induced of  $\phi$  morphism of chain complexes.

For  $n \geq 0$  consider the set  $K_n$  of all subcubes  $G$  of dimension  $n$  that have the form  $I_{s,t}$  with  $s, t \in V_G$ . By [2; 5], for every cube  $I_{s,t} \in K_n$  there is an isomorphism  $\chi_{s,t} : I^n \rightarrow I_{s,t}$  such that the set of elements

$$\{(\chi_{s,t})_*(w_n) : I_{s,t} \in K_n\}$$

gives the basis of  $\Omega_n(G)$ . For  $n \geq 1$ , define homomorphisms  $\theta_n : \Omega_n(G) \rightarrow \Omega_n^c(G)$  on basic elements by

$$(4-4) \quad \theta_n((\chi_{s,t})_*(w_n)) = \chi_{s,t}^\square,$$

and then extend it by linearity. It is clear that  $\theta_0$  preserves the augmentation.

**Proposition 4.3.** *The homomorphisms  $\theta_n$  define a morphism of chain complexes*

$$(4-5) \quad \theta_* : \Omega_*(G) \rightarrow \Omega_*^c(G),$$

which is a right inverse morphism to  $\tau_*$ , that is,

$$\tau_*\theta_* = \text{Id} : \Omega_*(G) \rightarrow \Omega_*(G).$$

*Proof.* Let us first prove that  $\tau_n\theta_n = \text{Id}$ . For  $n = 0, 1$  this is trivial. Let  $n \geq 2$  and  $(\chi_{s,t})_*(w_n) \in \Omega_n(G)$  be a basic element. By (4-4) and (4-3) we have

$$(4-6) \quad \tau_n\theta_n((\chi_{s,t})_*(w_n)) = \tau_n(\chi_{s,t}^\square) = \chi_{s,t}_*(w_n).$$

Consider the diagram

$$(4-7) \quad \begin{array}{ccccc} \Omega_n(G) & \xrightarrow{\theta_n} & \Omega_n^c(G) & \xrightarrow{\tau_n} & \Omega_n(G) \\ \downarrow \partial & & \downarrow \partial^c & & \downarrow \partial \\ \Omega_{n-1}(G) & \xrightarrow{\theta_{n-1}} & \Omega_{n-1}^c(G) & \xrightarrow{\tau_{n-1}} & \Omega_{n-1}(G) \end{array}$$

where the horizontal compositions are identity homomorphisms by (4-6), the right square is commutative and the large square is evidently commutative. Now we prove that the left square is commutative. It follows from [2, Lemma 4] that, for

$$(\phi_{s,t})_*(w_n) \in \Omega_n(G),$$

we have

$$(4-8) \quad \begin{aligned} \theta_{n-1}(\partial((\phi_{s,t})_*(w_n))) &= \theta_{n-1}\left(\sum_{I_{s',t'} \subset I_{s,t}} (-1)^{\sigma(I,I')} (\phi_{s',t'})_*(w_{n-1})\right) \\ &= \sum (-1)^{\sigma(I,I')} \phi_{s',t'}^\square, \end{aligned}$$

where the sum is taken over all  $(n-1)$ -cubes  $I' = I_{s',t'} \subset I_{s,t} = I$ . By (2-7) and (4-4) for

$$(\phi_{s,t})_*(w_n) \in \Omega_n(G),$$

we have

$$(4-9) \quad \partial^c(\theta((\phi_{s,t})_*(w_n))) = \partial^c(\phi_{s,t}^\square) = \sum_{p=1}^n (-1)^p ((\phi_{s,t}^\square)_{p,0} - (\phi_{s,t}^\square)_{p,1}),$$

where the sum consists of all singular  $(n-1)$ -subcubes of the cube  $I^n$  with coefficients. Since bottom row in (4-7) is the identity homomorphism we conclude from (4-3), (4-8) and (4-9) that the left square in (4-7) is commutative, which finishes the proof.  $\square$

**Proposition 4.4.** *There is a chain homotopy between  $\theta_*\tau_* : \Omega_*^c(G) \rightarrow \Omega_*^c(G)$  and the identity map  $\text{Id} : \Omega_*^c(G) \rightarrow \Omega_*^c(G)$ .*

*Proof.* The chain complex  $\Omega_*^c(G)$  is geometric and the chain maps  $\theta_*\tau_*$  and  $\text{Id}$  evidently preserve augmentation. For a singular cube  $\phi : I^n \rightarrow G$  consider the

subgraph  $G_\phi \subset G$ , that is, image of  $\phi$ . This is a contractible cubical digraph by Theorem 3.6. Thus we assign to every basic element  $\phi^\square \in \Omega_*^c(G)$  the subcomplex

$$(4-10) \quad E_*(\phi^\square) \stackrel{\text{def}}{=} \Omega_*^c(G_\phi) \subset \Omega_*^c(G),$$

which is acyclic since  $G_\phi$  is contractible.

Now we check that  $E$  is an algebraic carrier function, that is, condition (i) of (4-1) is satisfied. Let  $\phi^\square \in \Omega_*^c(G)$  be a basic element given by a singular cube  $\phi : I^n \rightarrow G$  with  $n \geq 0$ . By (2-6) and (2-7), the element  $\partial(\phi^\square)$  is given by the sum of the basic elements  $(\phi \circ V_{p\epsilon})^\square$  with coefficients  $(\pm 1)$  where the maps  $V_{p\epsilon} : I^{n-1} \rightarrow I^n$  are the inclusions. Hence the digraph  $G_{\phi \circ V_{p\epsilon}}$  is a subgraph of  $G_\phi$ , and hence the chain complex

$$E_*((\phi \circ V_{p\epsilon})^\square) = \Omega_*^c(G_{\phi \circ V_{p\epsilon}})$$

is a subcomplex of  $E_*(\phi^\square)$ . Thus for the basic singular cube  $b \in \Omega_{n-1}^c(G)$  and  $b < \phi^\square$  we obtain that  $b = (\phi \circ V_{p\epsilon})^\square$  and

$$E_*(b) = E_*((\phi \circ V_{p\epsilon})^\square) < E_*(\phi^\square).$$

Hence we have the algebraic acyclic carrier function  $E$  from  $\Omega_*^c(G)$  to itself.

Now we prove that the chain maps  $\theta_* \tau_*$  and  $\text{Id}$  from  $\Omega_*^c(G)$  to itself are carried by the function  $E$ . Consider a basic element  $\phi^\square \in \Omega_n^c(G)$ . Then

$$(4-11) \quad \text{Id}(\phi^\square) = \phi^\square \in \Omega_*^c(G_\phi) = E_*(\phi^\square)$$

since image of  $\phi$  is the digraph  $G_\phi$ . Hence the chain map

$$\text{Id} : \Omega_n^c(G) \rightarrow \Omega_n^c(G)$$

is carried by the algebraic carrier function  $E$ .

By (4-3) and (4-4), we have

$$(4-12) \quad \theta_n \tau_n(\phi^\square) = \theta_n(\phi_*(w_n)), \quad \phi : I^n \rightarrow G.$$

We have only two different possibilities for the  $\phi_*(w_n)$ . In the first case,  $\phi$  is an isomorphism on its image  $G_\phi = I_{s,t} \cong I^n$  with

$$s = \phi(0, \dots, 0), \quad t = \phi(1, \dots, 1),$$

where  $(0, \dots, 0) \in V_{I^n}$  is the origin vertex and  $(1, \dots, 1) \in V_{I^n}$  is the end vertex of the cube  $I^n$ . Note that for any isomorphism  $\psi : I^n \rightarrow I^n$  we have  $\psi_*(w_n) = \pm w_n$ . Hence in this case subgraph  $G_\phi \subset G$  coincides with the subgraph cube  $G_{\chi_{s,t}} \subset G$  and by (4-4) we have

$$(4-13) \quad \theta_n \tau_n(\phi^\square) = \theta_n(\phi_*(w_n)) = \theta_n(\pm(\chi_{s,t})_*(w_n)) = \pm \chi_{s,t}^\square,$$

where

$$\chi_{s,t} : I^n \rightarrow D_{s,t} = G_\phi.$$

That is,

$$\theta_n \tau_n(\phi^\square) \in \Omega_n^c(G_{\chi_{s,t}}) = \Omega_n^c(G_\phi) = E_n(\phi^\square).$$

In the second case, the image of  $\phi$  does not contain any cube of dimension  $n$ , and hence  $\phi_*(w_n) = 0$ . Consequently, we have

$$\theta_n \phi_*(w_n) = 0 \in E_*(\phi^\square).$$

Then the claim follows from the acyclic carriers Theorem 4.2.  $\square$

**Theorem 4.5.** *For any finite cubical digraph  $G$ , the chain maps  $\tau_*$  and  $\theta_*$  are homotopy inverses, and hence induce isomorphisms of homology groups*

$$H_*^c(G) \cong H_*(G).$$

*Proof.* Indeed, it follows from Propositions 4.3 and 4.4 that the chain maps  $\tau_*$  and  $\theta_*$  are homotopy inverses.  $\square$

**Corollary 4.6.** *Let  $\Delta$  be a finite simplicial complex. Consider a digraph  $G_\Delta$  (see [2]) with the set of vertices given by the set of all simplexes from  $\Delta$ , and*

$$s \rightarrow t (t, s \in \Delta) \quad \text{if and only if } s \supset t \text{ and } \dim s = \dim t + 1.$$

*Then the graph  $G_\Delta$  is a cubical digraph and*

$$H_*^c(G_\Delta) \cong H_*(\Delta),$$

*where  $H_*(\Delta)$  are the simplicial homology groups of  $\Delta$ .*

*Proof.* Indeed, it is proved in [2] that path homology groups  $H_*(G_\Delta)$  are isomorphic to simplicial homology groups  $H_*(\Delta)$ .  $\square$

## References

- [1] A. Grigor'yan, Y. Lin, Y. Muranov, and S.-T. Yau, "Homotopy theory for digraphs", *Pure Appl. Math. Q.* **10**:4 (2014), 619–674. MR
- [2] A. Grigor'yan, Y. V. Muranov, and S.-T. Yau, "Graphs associated with simplicial complexes", *Homology Homotopy Appl.* **16**:1 (2014), 295–311. MR Zbl
- [3] A. Grigor'yan, Y. Lin, Y. Muranov, and S.-T. Yau, "Cohomology of digraphs and (undirected) graphs", *Asian J. Math.* **19**:5 (2015), 887–931. MR
- [4] A. Grigor'yan, Y. Muranov, and S.-T. Yau, "Homologies of digraphs and Künneth formulas", *Comm. Anal. Geom.* **25**:5 (2017), 969–1018. MR Zbl
- [5] A. A. Grigor'yan, Y. V. Muranov, and R. B. Jimenez, "Homology of digraphs", *Mat. Zametki* **109**:5 (2021), 705–722. MR Zbl
- [6] P. J. Hilton and S. Wylie, *Homology theory: an introduction to algebraic topology*, Cambridge University Press, New York, 1960. MR Zbl
- [7] V. V. Prasolov, *Elements of homology theory*, Graduate Studies in Mathematics **81**, American Mathematical Society, Providence, RI, 2007. MR Zbl

Received January 22, 2021. Revised November 23, 2022.

ALEXANDER GRIGOR'YAN  
DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF BIELEFELD  
BIELEFELD  
GERMANY  
[grigor@math.uni-bielefeld.de](mailto:grigor@math.uni-bielefeld.de)

YURI MURANOV  
FACULTY OF MATHEMATICS AND COMPUTER SCIENCE  
UNIVERSITY OF WARMIA AND MAZURY IN OLSZTYN  
OLSZTYN  
POLAND  
[muranov@matman.uwm.edu.pl](mailto:muranov@matman.uwm.edu.pl)

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Matthias Aschenbrenner  
Fakultät für Mathematik  
Universität Wien  
Vienna, Austria  
[matthias.aschenbrenner@univie.ac.at](mailto:matthias.aschenbrenner@univie.ac.at)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Robert Lipshitz  
Department of Mathematics  
University of Oregon  
Eugene, OR 97403  
[lipshitz@uoregon.edu](mailto:lipshitz@uoregon.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2023 is US \$605/year for the electronic version, and \$820/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 322    No. 1    January 2023

---

Rips construction without unique product	1
GOULNARA ARZHANTSEVA and MARKUS STEENBOCK	
No periodic geodesics in jet space	11
ALEJANDRO BRAVO-DODDOLI	
Coarse geometry of Hecke pairs and the Baum–Connes conjecture	21
CLÉMENT DELL’AIERA	
On homology theories of cubical digraphs	39
ALEXANDER GRIGOR’YAN and YURI MURANOV	
The geometry and topology of stationary multiaxisymmetric vacuum black holes in higher dimensions	59
VISHNU KAKKAT, MARCUS KHURI, JORDAN RAINONE and GILBERT WEINSTEIN	
Quasilinear Schrödinger equations: ground state and infinitely many normalized solutions	99
HOUWANG LI and WENMING ZOU	
Thomae’s function on a Lie group	139
MARK REEDER	
On the potential function of the colored Jones polynomial with arbitrary colors	171
SHUN SAWABE	
Pushforward and smooth vector pseudo-bundles	195
ENXIN WU	