

*Pacific
Journal of
Mathematics*

**ON THE POTENTIAL FUNCTION OF THE
COLORED JONES POLYNOMIAL WITH ARBITRARY COLORS**

SHUN SAWABE

ON THE POTENTIAL FUNCTION OF THE COLORED JONES POLYNOMIAL WITH ARBITRARY COLORS

SHUN SAWABE

We consider the potential function of the colored Jones polynomial for a link with arbitrary colors and obtain the cone-manifold structure for the link complement. In addition, we establish a relationship between a saddle point equation and hyperbolicity of the link complement. This provides evidence for the Chen–Yang conjecture on the link complement.

1. Introduction

The volume conjecture is one of the most important problems in low-dimensional topology. Kashaev [1997] discovered that a certain limit of the Kashaev invariant of specific hyperbolic knots such as the figure-eight knot is equal to the hyperbolic volume of their complements. Murakami and Murakami [2001] proved that the Kashaev invariant is a specialization of the colored Jones polynomial and conjectured that a similar limit of the colored Jones polynomial for an arbitrary knot is equal to the simplicial volume of its complement. In addition, Chen and Yang [2018] considered the volume conjectures for 3-manifold invariants such as the Reshetikhin–Turaev invariant and the Turaev–Viro invariant, and provided numerical evidence for them for specific 3-manifolds. Detcherry, Kalfagianni, and Yang [Detcherry et al. 2018] showed the relationship between the colored Jones polynomial for a link and the Turaev–Viro invariant of its complement. By using this relation, they mathematically verified the Chen–Yang conjecture for complements of the figure-eight knot and Borromean rings. In addition, Belletti, Detcherry, Kalfagianni, and Yang verified the Chen–Yang conjecture for fundamental shadow links in [Belletti et al. 2022].

Meanwhile, theoretical evidence of the original volume conjecture has been considered. Kashaev and Tirkkonen [2000] proved the volume conjecture for torus knots. On the other hand, Yokota [2000] found a correspondence between quantum factorials in the Kashaev invariant and an ideal triangulation of a hyperbolic knot complement. He showed that a saddle point equation for the potential function (see Section 3 for the definition) of the invariant is equivalent to a hyperbolicity equation.

MSC2020: 57K14, 57K31, 57K32.

Keywords: Chen–Yang conjecture, cone-manifold, potential function, volume conjecture.

Also, the potential function of the colored Jones polynomial $J_N(K; e^{2\pi\sqrt{-1}/N})$ for a hyperbolic knot K is considered in [Cho 2016a; 2016b; Cho and Murakami 2013].

In this study, we consider the potential function of the colored Jones polynomial for a link L with arbitrary colors. We establish a relationship between a saddle point equation and a hyperbolicity equation of the link complement. More precisely, for a fixed diagram D of the link L , we introduce a potential function $\Phi_D(a_1, \dots, a_n, w_1, \dots, w_\nu)$ of the colored Jones polynomial $J_{\mathbf{a}(N)}(L; e^{2\pi\sqrt{-1}/N})$ with new parameters corresponding to the colors $\mathbf{a}(N)$ of link components. When we fix the new parameters $\mathbf{a} = (a_1, \dots, a_n)$, the saddle point $(\sigma_1(\mathbf{a}), \dots, \sigma_\nu(\mathbf{a}))$ of $\Phi_D(\mathbf{a}, -)$ gives a noncomplete hyperbolic structure to the link complement. In fact, the manifold M_{a_1, \dots, a_n} with the hyperbolic structure is a cone-manifold. Specifically, we prove the following statement:

Theorem 4.1. *The hyperbolic volume of the cone-manifold M_{a_1, \dots, a_n} is equal to the imaginary part of*

$$\tilde{\Phi}_D = \Phi_D - \sum_{j=1}^{\nu} w_j \frac{\partial \Phi_D}{\partial w_j} \log w_j$$

evaluated at $w_j = \sigma_j(\mathbf{a})$ for $j = 1, \dots, \nu$.

Here, the function $\Phi_D(\mathbf{a}, \sigma_1(\mathbf{a}), \dots, \sigma_\nu(\mathbf{a}))$ determines the Neumann–Zagier potential function [Neumann and Zagier 1985]. Furthermore, we prove that the derivatives of the potential function with respect to the new parameters correspond to the completeness of the hyperbolic structure of the link complement. Note that similar arguments for the Kashaev invariant of the 5_2 knot are indicated in [Yokota 2003]. As an application, we prove the following theorem:

Theorem 5.3. *Let D be a diagram of a hyperbolic link with n components, and let $\mathbf{1}$ be $(1, \dots, 1) \in \mathbb{Z}^n$. The point $(\mathbf{1}, \sigma_1(\mathbf{1}), \dots, \sigma_\nu(\mathbf{1}))$ is a saddle point of the function $\Phi_D(a_1, \dots, a_n, w_1, \dots, w_\nu)$ and gives a complete hyperbolic structure to the link complement.*

The paper is organized as follows: In Section 2, we recall the facts on the colored Jones polynomial and the Turaev–Viro invariant. In Section 3, we give the potential function of the colored Jones polynomial. In Section 4, we consider the case where the new parameters are fixed and prove Theorem 4.1. In Section 5, we regard the new parameters as variables and prove Theorem 5.3. In Section 6, we briefly mention the Witten–Reshetikhin–Turaev invariant.

2. Preliminaries

In this section, we review some facts on the invariants for a link and a 3-manifold.

The colored Jones polynomial and the Turaev–Viro invariant. Let L be an oriented n -component link, let \mathbf{i} be a multiinteger, and let t be an indeterminate. The colored Jones polynomial $J_{\mathbf{i}}(L; t)$ is defined skein-theoretically by using the Kauffman bracket, which is a map $\langle \cdot \rangle$ from the set of all unoriented diagrams of links to the ring of Laurent polynomials $\mathbb{Z}[A, A^{-1}]$ in an indeterminate A given by the following axioms:

- (1) For the trivial diagram \bigcirc ,

$$\langle \bigcirc \rangle = 1.$$

- (2) For an unoriented diagram D with the trivial component added,

$$\langle D \sqcup \bigcirc \rangle = (-A^2 - A^{-2})\langle D \rangle.$$

- (3) For each crossing,

$$\langle \begin{array}{c} \diagup \\ \diagdown \end{array} \rangle = A\langle \begin{array}{c} \diagdown \\ \diagup \end{array} \rangle + A^{-1}\langle \begin{array}{c} \diagdown \\ \diagdown \end{array} \rangle$$

Let D_0 be an unoriented diagram of the link L . The colored Jones polynomial $J_{\mathbf{i}}(L; t)$ for the link L is a certain normalization of the Kauffman bracket of the parallelized diagram of D_0 in which the Jones–Wenzl idempotent is inserted, where $t = A^{-4}$ [Detcherry et al. 2018].

Remark 2.1. In this paper, we normalize the colored Jones polynomial so that the one for the trivial knot is equal to 1.

From the perspective of skein theory, we can define the 3-manifold invariants such as the Reshetikhin–Turaev invariant or the Turaev–Viro invariant. Detcherry, Kalfagianni, and Yang [Detcherry et al. 2018] presented the relationship between the Turaev–Viro invariant for the link complement and the colored Jones polynomial.

Theorem 2.2 [Detcherry et al. 2018]. *Let $L \subset S^3$ be a link with n components and $\bar{t} = q^2$. Namely, $\bar{t} = q^2 = A^4$.*

- (1) *For an integer $r \geq 3$ and a primitive $4r$ -th root of unity A ,*

$$TV_r(S^3 \setminus L, q) = \eta_r^2 \sum_{1 \leq i \leq r-1} |J'_i(L; \bar{t})|^2.$$

- (2) *For an odd integer $r = 2m + 1 \geq 3$ and a primitive $2r$ -th root of unity A ,*

$$TV_r(S^3 \setminus L, q) = 2^{n-1}(\eta'_r)^2 \sum_{1 \leq i \leq m} |J'_i(L; \bar{t})|^2.$$

Here, η_r and η'_r are

$$\eta_r = \frac{A^2 - A^{-2}}{\sqrt{-2r}} \quad \text{and} \quad \eta'_r = \frac{A^2 - A^{-2}}{\sqrt{-r}}.$$

In addition, for a multiinteger $\mathbf{i} = (i_1, \dots, i_n)$, we let $1 \leq \mathbf{i} \leq m$ denote that $1 \leq i_k \leq m$ for all integers $k = 1, \dots, n$.

Remark 2.3. In [Detcherry et al. 2018], the normalization of the colored Jones polynomial and conventions on parameters are slightly different from the ones in this paper. Therefore, we use the notation $J'_i(L; \bar{t})$ in Theorem 2.2.

These invariants are conjectured to relate to the geometry of the 3-manifold. Murakami and Murakami [2001] conjectured that a certain limit of the colored Jones polynomial for a knot is equal to the volume of the complement of the knot.

Conjecture 2.4 (volume conjecture [Murakami and Murakami 2001]). *For any knot K ,*

$$2\pi \lim_{N \rightarrow \infty} \frac{\log |J_N(K; t = e^{2\pi\sqrt{-1}/N})|}{N} = v_3 \|K\|,$$

where v_3 is the volume of the ideal regular tetrahedron in the three-dimensional hyperbolic space and $\|\cdot\|$ is the simplicial volume for the complement of K .

This conjecture was generalized to the one for 3-manifold invariants.

Conjecture 2.5 (Chen–Yang conjecture [2018]). *For any 3-manifold M with a complete hyperbolic structure of the finite volume,*

$$2\pi \lim_{r \rightarrow \infty} \frac{\log TV_r(M, q = e^{2\pi\sqrt{-1}/r})}{r} = \text{Vol}(M),$$

where r runs over all odd integers, $TV(M)$ is a Turaev–Viro invariant of M and $\text{Vol}(M)$ is a hyperbolic volume of M .

Moreover, Detcherry, Kalfagianni, and Yang proved the following theorem by using Theorem 2.2:

Theorem 2.6 [Detcherry et al. 2018]. *Let L be either the figure-eight knot or the Borromean rings, and let M be the complement of L in S^3 . Then,*

$$\begin{aligned} 2\pi \lim_{r \rightarrow \infty} \frac{\log TV_r(M, q = e^{2\pi\sqrt{-1}/r})}{r} \\ = 4\pi \lim_{m \rightarrow \infty} \frac{\log |J'_m(L; \bar{t} = e^{4\pi\sqrt{-1}/(2m+1)})|}{2m+1} = \text{Vol}(M), \end{aligned}$$

where $r = 2m + 1$ runs over all odd integers.

Remark 2.7. If t is a root of unity, \bar{t} is the complex conjugate of t . Therefore,

$$\lim_{m \rightarrow \infty} \frac{\log |J'_m(L; \bar{t} = e^{4\pi\sqrt{-1}/(2m+1)})|}{2m+1} = \lim_{m \rightarrow \infty} \frac{\log |J_m(L; t = e^{4\pi\sqrt{-1}/(2m+1)})|}{2m+1}.$$

Meanwhile, the evidence of the volume conjecture was established in [Yokota 2000]. What is important is that a saddle point equation of a potential function of the colored Jones polynomial for a knot coincides with a gluing condition of the ideal triangulation of the knot complement. This and Theorem 2.2 indicate that if we can establish a similar relationship between a hyperbolicity equation

and a potential function of the colored Jones polynomial with arbitrary colors, the relationship is evidence of the Chen–Yang conjecture for a link complement.

The R -matrix of the colored Jones polynomial. In this subsection, we give the R -matrix of the colored Jones polynomial by following [Kirby and Melvin 1991]. For an integer $r > 1$, let \mathcal{A}_r be the algebra generated by X, Y, K , and \bar{K} with the following relations:

$$\begin{aligned} \bar{K} &= K^{-1}, & KX &= sXK, & KY &= s^{-1}YK, \\ XY - YX &= \frac{K^2 - \bar{K}^2}{s - s^{-1}}, & X^r &= Y^r = 0, & K^{4r} &= 1, \end{aligned}$$

where $s = e^{\pi\sqrt{-1}/r}$. Namely, \mathcal{A}_r is $\mathcal{U}_q(\mathfrak{sl}_2)$ with the last 3 relations. The universal R -matrix $\mathcal{R} \in \mathcal{A}_r \otimes \mathcal{A}_r$ is given by

$$\mathcal{R} = \frac{1}{4r} \sum_{\substack{0 \leq k < r \\ 0 \leq a, b < 4r}} \frac{(s - s^{-1})^k}{[k]_s!} s^{-(ab + (b-a)k + k)/2} X^k K^a \otimes Y^k K^b.$$

Here, we put

$$[k]_s = \frac{s^k - s^{-k}}{s - s^{-1}}, \quad [k]_s! = [k]_s \cdots [1]_s, \quad [0]_s! = 1.$$

Let N be a positive integer and m be the half-integer satisfying $N = 2m + 1$. We define the action of \mathcal{A}_r on an N -dimensional complex vector space V with a basis $\{e_{-m}, e_{-m+1}, \dots, e_m\}$ by

$$Xe_i = [m - i + 1]_s e_{i-1}, \quad Ye_i = [m + i + 1]_s e_{i+1}, \quad Ke_i = s^{-i} e_i.$$

Here, e_i in this paper corresponds to e_{-i} in [Kirby and Melvin 1991]. Let V' be an $(N' = 2m' + 1)$ -dimensional complex vector space with basis $\{e'_{-m'}, \dots, e'_{m'}\}$. Then, the quantum R -matrix $R_{VV'} : V \otimes V' \rightarrow V' \otimes V$ is given by

$$\begin{aligned} &R_{VV'}(e_i \otimes e'_j) \\ &= \sum_{k=0}^{\min\{m+i, m'-j\}} \frac{\{m - i + k\}_s! \{m' + j + k\}_s!}{\{k\}_s! \{m - i\}_s! \{m' + j\}_s!} s^{2ij + k(i-j) - k(k+1)/2} e'_{j+k} \otimes e_{i-k}, \end{aligned}$$

where $\{k\}_s = s^k - s^{-k}$, $\{k\}_s! = \{k\}_s \cdots \{1\}_s$, and $\{0\}_s! = 1$.

Also, its inverse is

$$\begin{aligned} &R_{VV'}^{-1}(e'_i \otimes e_j) \\ &= \sum_{k=0}^{\min\{m-i, m'+j\}} (-1)^k \frac{\{m - j + k\}_s! \{m' + i + k\}_s!}{\{k\}_s! \{m - j\}_s! \{m' + i\}_s!} s^{-2ij + k(i-j)/2 + k(k+1)/2} e_{j-k} \otimes e'_{i+k}. \end{aligned}$$




Figure 2.1. The links that are identical except for these regions.

These matrices and the isomorphism $\mu : V \rightarrow V$, where

$$\mu(e_i) = s^{-2i} e_i, \quad i = -m, \dots, m,$$

defines a link invariant \tilde{J} . If $V = V'$ and $\dim V = 2$, then

$$R_{VV} = \begin{pmatrix} s^{1/2} & 0 & 0 & 0 \\ 0 & 0 & s^{-1/2} & 0 \\ 0 & s^{-1/2} & s^{1/2} & -s^{-3/2} \\ 0 & 0 & 0 & s^{1/2} \end{pmatrix}$$

and satisfies

$$s^{1/2} R_{VV} - s^{-1/2} R_{VV}^{-1} = (s - s^{-1}) I_4,$$

where I_4 is the 4×4 identity matrix. Considering the writhes, this implies

$$(2.1) \quad s^2 \tilde{J}(L_+) - s^{-2} \tilde{J}(L_-) = (s - s^{-1}) \tilde{J}(L_0),$$

where L_+ , L_- , and L_0 are the links in [Figure 2.1](#).

Under the substitution $s = -t^{-1/2}$, the relation (2.1) coincides with the skein relation of the Jones polynomial. Therefore, under this substitution the R -matrix of the colored Jones polynomial $J_i(L; t)$ for L with colors $\mathbf{i} = (i_1, \dots, i_n) \in \mathbb{Z}_{>0}^n$, where i_j , with $j = 1, \dots, n$, is the dimension of the assigned representation, is

$$(2.2) \quad R_{VV'}(e_i \otimes e'_j) = \sum_{k=0}^{\min\{m+i, m'-j\}} (-1)^{k+k(m+m')+2ij} \frac{\{m-i+k\}! \{m'+j+k\}!}{\{k\}! \{m-i\}! \{m'+j\}!} \\ \times t^{-ij-k(i-j)/2+k(k+1)/4} e'_{j+k} \otimes e_{i-k},$$

and its inverse is

$$R_{VV'}^{-1}(e'_i \otimes e_j) = \sum_{k=0}^{\min\{m-i, m'+j\}} (-1)^{-k(m+m')-2ij} \frac{\{m-j+k\}! \{m'+i+k\}!}{\{k\}! \{m-j\}! \{m'+i\}!} \\ \times t^{ij-k(i-j)/2-k(k+1)/4} e_{j-k} \otimes e'_{i+k},$$

where

$$\{k\} = t^{k/2} - t^{-k/2}, \quad \{k\}! = \{k\} \{k-1\} \cdots \{1\}, \quad \{0\}! = 1.$$

3. Potential function

Let $L = L_1 \cup \dots \cup L_n$ be an oriented n -component link. We deform L so that L is a closure of a braid. Let D be its oriented diagram, and $\xi_N = e^{2\pi\sqrt{-1}/N}$ be the primitive N -th root of unity. For each link component L_i , with $i = 1, \dots, n$, we assign its color $a_i(N) \in \mathbb{Z}_{>0}$. We put $\mathbf{a}(N) = (a_1(N), \dots, a_n(N))$. In this section, we determine a potential function of the colored Jones polynomial $J_{\mathbf{a}(N)}(L; \xi_N^p)$ for L , where p is a nonzero integer. See [Cho 2016b] for details.

Definition 3.1. Suppose that the asymptotic behavior of a certain quantity Q_N for a sufficiently large N is

$$Q_N \sim \int \dots \int_{\Omega} P_N e^{N/(2\pi\sqrt{-1})\Phi(z_1, \dots, z_\nu)} dz_1 \dots dz_\nu,$$

where P_N grows at most polynomially and Ω is a region in \mathbb{C}^ν . We call this function $\Phi(z_1, \dots, z_\nu)$ a potential function of Q_N .

We can easily verify that

$$(3.1) \quad \{k\}! = (-1)^k t^{-k(k+1)/4} (t)_k,$$

where $(t)_k = (1-t)(1-t^2) \dots (1-t^k)$. Thus, we approximate $(\xi_N^p)_k$ by continuous functions.

Proposition 3.2. For a sufficiently large integer N ,

$$\log(\xi_N^p)_k = \frac{N}{2p\pi\sqrt{-1}} \left(-\text{Li}_2(\xi_N^{pk}) + \frac{\pi^2}{6} + o(1) \right),$$

where Li_2 is a dilogarithm function

$$\text{Li}_2(z) = - \int_0^z \frac{\log(1-x)}{x} dx.$$

Remark 3.3. The dilogarithm function satisfies

$$\text{Li}_2(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^2}, \quad \text{for } |z| < 1, \quad \text{and} \quad \text{Li}_2(1) = \sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}.$$

Proof. By the direct calculation, we have

$$\begin{aligned} \log(\xi_N^p)_k &= \sum_{j=1}^k \log(1 - e^{2p\pi j\sqrt{-1}/N}) = N \left(\int_0^{k/N} \log(1 - e^{2p\pi\sqrt{-1}\theta}) d\theta + o(1) \right) \\ &= \frac{N}{2p\pi\sqrt{-1}} \left(\int_1^{\xi_N^{pk}} \frac{\log(1-x)}{x} dx + o(1) \right) \\ &= \frac{N}{2p\pi\sqrt{-1}} \left(-\text{Li}_2(\xi_N^{pk}) + \frac{\pi^2}{6} + o(1) \right). \quad \square \end{aligned}$$

First, we consider the case where the strings at a crossing are in the different components. Let $\{a(N)\}_{N=1,2,\dots}$ and $\{b(N)\}_{N=1,2,\dots}$ be sequences of natural numbers. We can approximate the R -matrix by Proposition 3.2. For a positive crossing of the link diagram, the R -matrix $R_{V'V'}$ of (2.2) is labeled. For convenience, we recall the summand of the R -matrix:

$$(-1)^{k+k(m_N+m'_N)+2ij} t^{-ij-k(i-j)/2+k(k+1)/4} \frac{\{m_N - i + k\}! \{m'_N + j + k\}!}{\{k\}! \{m_N - i\}! \{m'_N + j\}!}.$$

Here, m_N and m'_N are the half-integers satisfying $a(N) = 2m_N + 1$ and $b(N) = 2m'_N + 1$. If we assume that $a(N)$ and $b(N)$ are odd numbers, indices i and j are integers. Moreover, by adding 2 to $a(N)$ or $b(N)$ if necessary, we can assume that $m_N + m'_N$ is an even integer without changing the values of the limit $a(N)/N$ and $b(N)/N$. Therefore, under these assumptions the summand is

$$(-1)^k t^{-ij-k(i-j)/2+k(k+1)/4} \frac{\{m_N - i + k\}! \{m'_N + j + k\}!}{\{k\}! \{m_N - i\}! \{m'_N + j\}!}.$$

From (3.1), we have

$$t^{-ij-((m_N+m'_N)/2)k} \frac{(t)_{m_N-i+k} (t)_{m'_N+j+k}}{(t)_k (t)_{m_N-i} (t)_{m'_N+j}}.$$

Under substitution $x = \xi_N^i$, $y = \xi_N^j$, and $z = \xi_N^k$, the potential function for a positive crossing is

$$\frac{1}{p} \left\{ -\pi \sqrt{-1} p \frac{a+b}{2} \log(z^p) - \log(x^p) \log(y^p) - \frac{\pi^2}{6} - \text{Li}_2\left(e_a^p \frac{z^p}{x^p}\right) - \text{Li}_2(e_b^p y^p z^p) + \text{Li}_2\left(\frac{e_a^p}{x^p}\right) + \text{Li}_2(e_b^p y^p) + \text{Li}_2(z^p) \right\},$$

where $a(N)/N \rightarrow a$, $b(N)/b \rightarrow b$, and $e_a = e^{\pi\sqrt{-1}a}$. Note that the indices of the summand are labeled to the edges of the link diagram. We change these indices to the ones corresponding to regions of the link diagram as shown in Figure 3.1.




Figure 3.1. Indices corresponding to an edge E_i and regions R_l and R_r .




Figure 3.2. Indices corresponding to regions around a crossing.

If k_{j_1}, \dots, k_{j_4} are indices around a crossing as shown in Figure 3.2, we have

$$i = k_{j_2} - k_{j_1}, \quad j = k_{j_3} - k_{j_2}, \quad j + k = k_{j_4} - k_{j_1}, \quad i - k = k_{j_3} - k_{j_4}.$$

From the above equations, we have $k = k_{j_2} + k_{j_4} - k_{j_1} - k_{j_3}$. Therefore, by putting $w_{j_i} = \xi_N^{k_{j_i}}$ and substituting

$$x = \frac{w_{j_2}}{w_{j_1}}, \quad y = \frac{w_{j_3}}{w_{j_2}}, \quad z = \frac{w_{j_2} w_{j_4}}{w_{j_1} w_{j_3}},$$

the potential function for a positive crossing c is

$$\begin{aligned} \Phi_{c,p}^+ = \frac{1}{p} \left\{ \pi \sqrt{-1} p^2 \frac{a+b}{2} \log \frac{w_{j_1} w_{j_3}}{w_{j_2} w_{j_4}} - p^2 \log \frac{w_{j_2}}{w_{j_1}} \log \frac{w_{j_3}}{w_{j_2}} \right. \\ \left. - \text{Li}_2 \left(e_a^p \frac{w_{j_4}^p}{w_{j_3}^p} \right) - \text{Li}_2 \left(e_b^p \frac{w_{j_4}^p}{w_{j_1}^p} \right) + \text{Li}_2 \left(\frac{w_{j_2}^p w_{j_4}^p}{w_{j_1}^p w_{j_3}^p} \right) \right. \\ \left. + \text{Li}_2 \left(e_a^p \frac{w_{j_1}^p}{w_{j_2}^p} \right) + \text{Li}_2 \left(e_b^p \frac{w_{j_3}^p}{w_{j_2}^p} \right) - \frac{\pi^2}{6} \right\}. \end{aligned}$$

If the strings at a crossing are in the same component, we have to consider the modification on the Reidemeister move I. The Reidemeister move I on the component with a color $a(N)$ leads to the multiplication by $s^{2m_N+2m_N} = (-1)^{2m_N+2m_N} t^{-m_N-m_N}$. Therefore, we have to multiply $(-1)^{-2m_N-2m_N} t^{m_N+m_N}$ to cancel it. Under the assumption that $a(N)$ is an odd integer, this corresponds to the addition of the function $(\pi \sqrt{-1} pa)^2/p$. Therefore, the potential function is

$$\begin{aligned} \Phi_{c,p}^+ = \frac{1}{p} \left\{ (\pi \sqrt{-1} pa)^2 + \pi \sqrt{-1} p^2 a \log \frac{w_{j_1} w_{j_3}}{w_{j_2} w_{j_4}} - p^2 \log \frac{w_{j_2}}{w_{j_1}} \log \frac{w_{j_3}}{w_{j_2}} - \frac{\pi^2}{6} \right. \\ \left. - \text{Li}_2 \left(e_a^p \frac{w_{j_4}^p}{w_{j_3}^p} \right) - \text{Li}_2 \left(e_a^p \frac{w_{j_4}^p}{w_{j_1}^p} \right) + \text{Li}_2 \left(\frac{w_{j_2}^p w_{j_4}^p}{w_{j_1}^p w_{j_3}^p} \right) + \text{Li}_2 \left(e_a^p \frac{w_{j_1}^p}{w_{j_2}^p} \right) + \text{Li}_2 \left(e_a^p \frac{w_{j_3}^p}{w_{j_2}^p} \right) \right\}. \end{aligned}$$

Similarly, we obtain

$$\begin{aligned} \Phi_{c,p}^- = \frac{1}{p} & \left\{ -\pi\sqrt{-1}p^2\frac{a+b}{2} \log \frac{w_{j_1}w_{j_3}}{w_{j_2}w_{j_4}} + p^2 \log \frac{w_{j_3}}{w_{j_4}} \log \frac{w_{j_4}}{w_{j_1}} \right. \\ & - \text{Li}_2\left(e_a^p \frac{w_{j_1}^p}{w_{j_4}^p}\right) - \text{Li}_2\left(e_b^p \frac{w_{j_3}^p}{w_{j_4}^p}\right) - \text{Li}_2\left(\frac{w_{j_2}^p w_{j_4}^p}{w_{j_1}^p w_{j_3}^p}\right) \\ & \left. + \text{Li}_2\left(e_a^p \frac{w_{j_2}^p}{w_{j_3}^p}\right) + \text{Li}_2\left(e_b^p \frac{w_{j_2}^p}{w_{j_1}^p}\right) + \frac{\pi^2}{6} \right\} \end{aligned}$$

for a negative crossing c between different components, and

$$\begin{aligned} \Phi_{c,p}^- = \frac{1}{p} & \left\{ -(\pi\sqrt{-1}pa)^2 - \pi\sqrt{-1}p^2a \log \frac{w_{j_1}w_{j_3}}{w_{j_2}w_{j_4}} + p^2 \log \frac{w_{j_3}}{w_{j_4}} \log \frac{w_{j_4}}{w_{j_1}} + \frac{\pi^2}{6} \right. \\ & \left. - \text{Li}_2\left(e_a^p \frac{w_{j_1}^p}{w_{j_4}^p}\right) - \text{Li}_2\left(e_a^p \frac{w_{j_3}^p}{w_{j_4}^p}\right) - \text{Li}_2\left(\frac{w_{j_2}^p w_{j_4}^p}{w_{j_1}^p w_{j_3}^p}\right) + \text{Li}_2\left(e_a^p \frac{w_{j_2}^p}{w_{j_3}^p}\right) + \text{Li}_2\left(e_a^p \frac{w_{j_2}^p}{w_{j_1}^p}\right) \right\} \end{aligned}$$

for a negative crossing c between the same component. The potential function $\Phi_{D,p}$ of $J_{a(N)}(L, \xi_N^p)$ is a summation of these potential functions with respect to all crossings of D . That is,

$$\Phi_{D,p}(\mathbf{a}, w_1, \dots, w_v) = \sum_{c \text{ is a crossing}} \Phi_{c,p}^{\text{sgn}(c)},$$

where

$$\mathbf{a} = (a_1, \dots, a_n), \quad a_i = \lim_{N \rightarrow \infty} \frac{a_i(N)}{N}$$

and $\text{sgn}(c)$ is a signature of a crossing c . This potential function essentially coincides with Yoon's generalized potential function [Yoon 2021]. We can easily verify the following property by the definition of $\Phi_{D,p}$:

Proposition 3.4. $\Phi_{D,p}(\mathbf{a}, w_1, \dots, w_v)$ satisfies

$$\Phi_{D,p}(\mathbf{a}, w_1, \dots, w_v) = \frac{1}{p} \Phi_{D,1}(p\mathbf{a}, w_1^p, \dots, w_v^p).$$

Therefore, We mainly consider the case where $p = 1$ and write $\Phi_D = \Phi_{D,1}$.

4. A noncomplete hyperbolic structure

In this section, we provide geometric meanings of the potential function. In the rest of this paper, we assume that L is a hyperbolic link with n components. In this section, we also assume that $a_i \in [1 - \varepsilon, 1]$ for all $i = 1, \dots, n$, where ε is a sufficiently small positive real number. First, we consider derivatives of the potential functions with respect to the parameters corresponding to the regions of the link

diagram [Cho and Murakami 2013]. For a positive crossing c between different components, we have:

$$(4.1) \quad \begin{aligned} w_{j_1} \frac{\partial \Phi_c^+}{\partial w_{j_1}} &= \pi \sqrt{-1} \frac{a-b}{2} + \log \left(1 - e_a \frac{w_{j_1}}{w_{j_2}} \right)^{-1} \left(1 - e_b^{-1} \frac{w_{j_1}}{w_{j_4}} \right)^{-1} \left(1 - \frac{w_{j_1} w_{j_3}}{w_{j_2} w_{j_4}} \right), \\ w_{j_2} \frac{\partial \Phi_c^+}{\partial w_{j_2}} &= \pi \sqrt{-1} \frac{a+b}{2} + \log \left(1 - e_a^{-1} \frac{w_{j_2}}{w_{j_1}} \right) \left(1 - e_b^{-1} \frac{w_{j_2}}{w_{j_3}} \right) \left(1 - \frac{w_{j_2} w_{j_4}}{w_{j_1} w_{j_3}} \right)^{-1}, \\ w_{j_3} \frac{\partial \Phi_c^+}{\partial w_{j_3}} &= \pi \sqrt{-1} \frac{-a+b}{2} + \log \left(1 - e_a^{-1} \frac{w_{j_3}}{w_{j_4}} \right)^{-1} \left(1 - e_b \frac{w_{j_3}}{w_{j_2}} \right)^{-1} \left(1 - \frac{w_{j_1} w_{j_3}}{w_{j_2} w_{j_4}} \right), \\ w_{j_4} \frac{\partial \Phi_c^+}{\partial w_{j_4}} &= -\pi \sqrt{-1} \frac{a+b}{2} + \log \left(1 - e_a \frac{w_{j_4}}{w_{j_3}} \right) \left(1 - e_b \frac{w_{j_4}}{w_{j_1}} \right) \left(1 - \frac{w_{j_2} w_{j_4}}{w_{j_1} w_{j_3}} \right)^{-1}. \end{aligned}$$

Similarly, for a negative crossing c between different components, we have

$$(4.2) \quad \begin{aligned} w_{j_1} \frac{\partial \Phi_c^-}{\partial w_{j_1}} &= \pi \sqrt{-1} \frac{-a+b}{2} + \log \left(1 - e_a \frac{w_{j_1}}{w_{j_4}} \right) \left(1 - e_b^{-1} \frac{w_{j_1}}{w_{j_2}} \right) \left(1 - \frac{w_{j_1} w_{j_3}}{w_{j_2} w_{j_4}} \right)^{-1}, \\ w_{j_2} \frac{\partial \Phi_c^-}{\partial w_{j_2}} &= \pi \sqrt{-1} \frac{a+b}{2} + \log \left(1 - e_a \frac{w_{j_2}}{w_{j_3}} \right)^{-1} \left(1 - e_b \frac{w_{j_2}}{w_{j_1}} \right)^{-1} \left(1 - \frac{w_{j_2} w_{j_4}}{w_{j_1} w_{j_3}} \right), \\ w_{j_3} \frac{\partial \Phi_c^-}{\partial w_{j_3}} &= \pi \sqrt{-1} \frac{a-b}{2} + \log \left(1 - e_a^{-1} \frac{w_{j_3}}{w_{j_2}} \right) \left(1 - e_b \frac{w_{j_3}}{w_{j_4}} \right) \left(1 - \frac{w_{j_1} w_{j_3}}{w_{j_2} w_{j_4}} \right)^{-1}, \\ w_{j_4} \frac{\partial \Phi_c^-}{\partial w_{j_4}} &= -\pi \sqrt{-1} \frac{a+b}{2} + \log \left(1 - e_a^{-1} \frac{w_{j_4}}{w_{j_1}} \right)^{-1} \left(1 - e_b^{-1} \frac{w_{j_4}}{w_{j_3}} \right)^{-1} \left(1 - \frac{w_{j_2} w_{j_4}}{w_{j_1} w_{j_3}} \right). \end{aligned}$$

If a crossing is between the same component, the derivatives are (4.1) and (4.2) with $a = b$. These correspond to Thurston's triangulation [1999] of the link complement (see Figure 4.1).

Here, we put

$$\begin{aligned} u_1 &= e_a \frac{w_{j_1}}{w_{j_2}}, & u_2 &= e_a^{-1} \frac{w_{j_3}}{w_{j_4}}, & u_3 &= \frac{w_{j_2} w_{j_4}}{w_{j_1} w_{j_3}}, & u_4 &= e_b^{-1} \frac{w_{j_1}}{w_{j_4}}, & u_5 &= e_b \frac{w_{j_3}}{w_{j_2}}, \\ v_1 &= e_a^{-1} \frac{w_{j_4}}{w_{j_1}}, & v_2 &= e_a \frac{w_{j_2}}{w_{j_3}}, & v_3 &= \frac{w_{j_1} w_{j_3}}{w_{j_2} w_{j_4}}, & v_4 &= e_b \frac{w_{j_2}}{w_{j_1}}, & v_5 &= e_b^{-1} \frac{w_{j_4}}{w_{j_3}} \end{aligned}$$

in Figure 4.1. Furthermore, for a complex number z , denote

$$z' = \frac{1}{1-z} \quad \text{and} \quad z'' = 1 - \frac{1}{z}.$$

Note that if there exists a nonalternating part, the ideal tetrahedron abuts the one with the inverse complex number labeled. Thus we can ignore the contribution of such a part. Let G_i be a product of the parameters of ideal tetrahedra around the




Figure 4.1. Ideal tetrahedra on a positive crossing (left), and ideal tetrahedra on a negative crossing (right).

region R_i corresponding to the parameter w_i . Then, we have

$$w_i \frac{\partial \Phi_D}{\partial w_i} = \frac{\pi \sqrt{-1}}{2} r(a_1, \dots, a_n) + \log G_i,$$

where $\pi \sqrt{-1} r(a_1, \dots, a_n) / 2$ is the summation of first terms of $w_i \partial \Phi_c^\pm / \partial w_i$ with c running over all crossings around R_i . However, this is equal to 0 because the contribution of each parameter a to $r(a_1, \dots, a_n)$ is canceled as in Figure 4.2.

Therefore, the equations

$$(4.3) \quad \exp\left(w_i \frac{\partial \Phi_D}{\partial w_i}\right) = 1, \quad i = 1, 2, \dots, \nu,$$

coincide with the gluing condition of the ideal tetrahedra. Hence, we can obtain a hyperbolic structure from a saddle point $(\sigma_1(\mathbf{a}), \dots, \sigma_\nu(\mathbf{a}))$ of $\Phi_D(\mathbf{a}, -)$,




Figure 4.2. Signatures of parameters corresponding to edges (left). Note that the pattern of signatures is independent of the signature of a crossing. Contributions of each parameter (right). White circles represent either positive or negative crossings.




Figure 4.3. The dilation component of the meridian of the link component with the parameter a .

where $\mathbf{a} = (a_1, \dots, a_n)$. In addition, this hyperbolic structure is not complete in general because the dilation component of the meridian of the link component with the color a is equal to e_a^{-2} (see Figure 4.3).

Note that $\mathbf{a} = (1, \dots, 1)$ is the case of the original volume conjecture. So we suppose that $(\sigma_1(\mathbf{a}), \dots, \sigma_v(\mathbf{a}))$ gives $S^3 \setminus L$ the hyperbolic structure with the finite volume $\text{Vol}(S^3 \setminus L)$ when $\mathbf{a} = (1, \dots, 1)$ [Cho 2016a]. Let M_{a_1, \dots, a_n} be a manifold with the hyperbolic structure given by $(\sigma_1(\mathbf{a}), \dots, \sigma_v(\mathbf{a}))$. We will determine the detail of this noncomplete hyperbolic manifold M_{a_1, \dots, a_n} . Let a be a real number slightly less than 1. Note that the action derived from each meridian does not change a length because $|e_a^{-2}| = 1$. Therefore, the action derived from each longitude changes a length, since otherwise, both meridians and longitudes do not change a length and this results in the complete hyperbolic structure [Benedetti and Petronio 1992]. Therefore, the developing image in the upper half-space \mathbb{H}^3 of the link complement around the edge corresponding to parameter a should be as shown in Figure 4.4. If we glue faces by the action of meridians in Figure 4.4, each face is glued with the face rotated $2\pi(1 - a)$ around the singular set. Therefore, M_{a_1, \dots, a_n} is a cone-manifold of L with cone-angle $2\pi(1 - a_i)$ around the component corresponding to a_i . Specifically, we can prove the following proposition:

Theorem 4.1. *The hyperbolic volume of the cone-manifold M_{a_1, \dots, a_n} is equal to the imaginary part of the value¹ of a function*

$$\tilde{\Phi}_D = \Phi_D - \sum_{j=1}^v w_j \frac{\partial \Phi_D}{\partial w_j} \log w_j$$

evaluated at $w_j = \sigma_j(\mathbf{a})$, with $j = 1, \dots, v$.

¹In [Murakami 2000], this value is called the optimistic limit.




Figure 4.4. The developing image of the link complement with the noncomplete hyperbolic structure.

Proof. The hyperbolic volume $V(z)$ of the ideal tetrahedron with modulus z is given by the Bloch–Wigner function [Zagier 2007]

$$(4.4) \quad V(z) = \operatorname{Im} \operatorname{Li}_2(z) + \log |z| \arg(1 - z).$$

We only consider the case where a crossing is between different components. Let $V_c^\pm(a, b)$ be the sum of hyperbolic volumes of five ideal tetrahedra at a positive or negative crossing c , respectively. By using (4.4), we can show that

$$\operatorname{Im} \Phi_c^+ - V_c^+(a, b) = A_{j_1}^+ \log |w_{j_1}| + A_{j_2}^+ \log |w_{j_2}| + A_{j_3}^+ \log |w_{j_3}| + A_{j_4}^+ \log |w_{j_4}|,$$

where $A_{j_i}^+$, with $i = 1, 2, 3, 4$, are:

$$\begin{aligned} A_{j_1}^+ &= \frac{\pi}{2}(a - b) + \arg\left(1 - e_a \frac{w_{j_1}}{w_{j_2}}\right)^{-1} \left(1 - e_b^{-1} \frac{w_{j_1}}{w_{j_4}}\right)^{-1} \left(1 - \frac{w_{j_1} w_{j_3}}{w_{j_2} w_{j_4}}\right), \\ A_{j_2}^+ &= \frac{\pi}{2}(a + b) + \arg\left(1 - e_a^{-1} \frac{w_{j_2}}{w_{j_1}}\right) \left(1 - e_b^{-1} \frac{w_{j_2}}{w_{j_3}}\right) \left(1 - \frac{w_{j_2} w_{j_4}}{w_{j_1} w_{j_3}}\right)^{-1}, \\ A_{j_3}^+ &= \frac{\pi}{2}(-a + b) + \arg\left(1 - e_a^{-1} \frac{w_{j_3}}{w_{j_4}}\right)^{-1} \left(1 - e_b \frac{w_{j_3}}{w_{j_2}}\right)^{-1} \left(1 - \frac{w_{j_1} w_{j_3}}{w_{j_2} w_{j_4}}\right), \\ A_{j_4}^+ &= -\frac{\pi}{2}(a + b) + \arg\left(1 - e_a \frac{w_{j_4}}{w_{j_3}}\right) \left(1 - e_b \frac{w_{j_4}}{w_{j_1}}\right) \left(1 - \frac{w_{j_2} w_{j_4}}{w_{j_1} w_{j_3}}\right)^{-1}. \end{aligned}$$

Similarly, we can show that

$$\operatorname{Im} \Phi_c^- - V_c^-(a, b) = A_{j_1}^- \log |w_{j_1}| + A_{j_2}^- \log |w_{j_2}| + A_{j_3}^- \log |w_{j_3}| + A_{j_4}^- \log |w_{j_4}|,$$

where $A_{j_i}^-$, with $i = 1, 2, 3, 4$, are:

$$A_{j_1}^- = \frac{\pi}{2}(-a + b) + \arg\left(1 - e_a \frac{w_{j_1}}{w_{j_4}}\right) \left(1 - e_b^{-1} \frac{w_{j_1}}{w_{j_2}}\right) \left(1 - \frac{w_{j_1} w_{j_3}}{w_{j_2} w_{j_4}}\right)^{-1},$$

$$A_{j_2}^- = \frac{\pi}{2}(a + b) + \arg\left(1 - e_a \frac{w_{j_2}}{w_{j_3}}\right)^{-1} \left(1 - e_b \frac{w_{j_2}}{w_{j_1}}\right)^{-1} \left(1 - \frac{w_{j_2} w_{j_4}}{w_{j_1} w_{j_3}}\right),$$

$$A_{j_3}^- = \frac{\pi}{2}(a - b) + \arg\left(1 - e_a^{-1} \frac{w_{j_3}}{w_{j_2}}\right) \left(1 - e_b \frac{w_{j_3}}{w_{j_4}}\right) \left(1 - \frac{w_{j_1} w_{j_3}}{w_{j_2} w_{j_4}}\right)^{-1},$$

$$A_{j_4}^- = -\frac{\pi}{2}(a + b) + \arg\left(1 - e_a^{-1} \frac{w_{j_4}}{w_{j_1}}\right)^{-1} \left(1 - e_b^{-1} \frac{w_{j_4}}{w_{j_3}}\right)^{-1} \left(1 - \frac{w_{j_2} w_{j_4}}{w_{j_1} w_{j_3}}\right).$$

By summing up over all crossings, we verify the proposition. \square

Example 4.2 (figure-eight knot). Let θ be a real number in $[0, \frac{\pi}{3}]$. The volume $V(\theta)$ of the cone-manifold of the figure-eight knot with a cone-angle θ is given by the formula [Mednykh 2003; Mednykh and Rasskazov 2006]

$$V(\theta) = \int_{\theta}^{2\pi/3} \operatorname{arccosh}(1 + \cos \theta - \cos 2\theta) d\theta.$$

In this case, the cone-manifold admits a hyperbolic structure. On the other hand, the colored Jones polynomial for the figure-eight knot is given by Habiro and Le's formula [Habiro 2000]

$$J_N(4_1; t) = \frac{1}{\{N\}} \sum_{p=0}^{N-1} \frac{\{N+p\}!}{\{N-p-1\}!}.$$

We assume that a is in $(\frac{5}{6}, 1)$ so that $0 < 2\pi(1-a) < \frac{\pi}{3}$. The potential function of $J_{a(N)}(4_1, \xi_N)$ is

$$\Phi(a, x) = -2\pi \sqrt{-1} a \log x - \operatorname{Li}_2(e_a^2 x) + \operatorname{Li}_2(e_a^2 x^{-1}),$$

and the derivative of this function with respect to x is

$$\frac{\partial \Phi}{\partial x} = \frac{1}{x} \log(-x + e_a^2 + e_a^{-2} - x^{-1}).$$

As a solution of the equation $\partial \Phi / \partial x = 0$, we obtain

$$x_0(a) = \left(\cos 2\pi a - \frac{1}{2}\right) - \sqrt{\left(\cos 2\pi a - \frac{3}{2}\right) \left(\cos 2\pi a + \frac{1}{2}\right)}.$$

Since $\frac{5}{6} < a < 1$, the absolute value of $x_0(a)$ is equal to 1. So we put $x_0(a) = e^{\sqrt{-1}\varphi(a)}$, where $\varphi(a) \in (-\pi, \pi]$. Then, the imaginary part of $\Phi(a, x_0(a))$ is

$$\operatorname{Im} \Phi(a, x_0(a)) = -2\Lambda \left(\pi a + \frac{\varphi(a)}{2}\right) + 2\Lambda \left(\pi a - \frac{\varphi(a)}{2}\right).$$

We will show that $\text{Im } \Phi(a, x_0(a)) = V(2\pi(1-a))$ as a function on the closed interval $[\frac{2}{3}, 1]$. If $a = \frac{2}{3}$, they are both 0. The derivative with respect to a is

$$\begin{aligned} \frac{d\Phi(a, x_0(a))}{da} &= \frac{\partial\Phi}{\partial a}(a, x_0(a)) + \frac{\partial\Phi}{\partial x}(a, x_0(a)) \frac{dx_0(a)}{da} \\ &= 2\pi\sqrt{-1} \log \frac{1 - e_a^2 x_0(a)}{x_0(a) - e_a^2} \\ &= -2\pi^2 + 2\pi\sqrt{-1} \log \left(\frac{e_a^2 x_0(a) - 1}{e_a^{-2} x_0(a) - 1} e_a^{-2} \right). \end{aligned}$$

Since $e^{\sqrt{-1}\theta} - 1 = 2 \sin(\theta/2)e^{\sqrt{-1}(\pi+\theta)/2}$, we obtain

$$\frac{d\Phi(a, x_0(a))}{da} = -2\pi^2 + 2\pi\sqrt{-1} \log \frac{\sin((\varphi(a) + 2\pi a)/2)}{\sin((\varphi(a) - 2\pi a)/2)}.$$

Let $f(a)$ be the function inside the log, then

$$\cosh \log f(a) = \frac{\sin^2((\varphi(a) + 2\pi a)/2) + \sin^2((\varphi(a) - 2\pi a)/2)}{2 \sin((\varphi(a) + 2\pi a)/2) \sin((\varphi(a) - 2\pi a)/2)}.$$

Note that the denominator of the right-hand side is $\cos(2\pi a) - \cos \varphi(a) = \frac{1}{2}$. Then,

$$\begin{aligned} \cosh \log f(a) &= 2 \left(\sin^2 \frac{\varphi(a) + 2\pi a}{2} + \sin^2 \frac{\varphi(a) - 2\pi a}{2} \right) \\ &= 2 - \cos(\varphi(a) + 2\pi a) - \cos(\varphi(a) - 2\pi a) \\ &= 2 - 2 \cos \varphi(a) \cos 2\pi a \\ &= 1 + \cos 2\pi a - \cos 4\pi a. \end{aligned}$$

Therefore, we obtain

$$\frac{d\Phi(a, x_0(a))}{da} = -2\pi^2 + 2\pi\sqrt{-1} \operatorname{arccosh}(1 + \cos 2\pi a - \cos 4\pi a).$$

Clearly, the imaginary part of this function is $2\pi \operatorname{arccosh}(1 + \cos 2\pi a - \cos 4\pi a)$ which is equal to $dV(2\pi(1-a))/da$. This shows that $V(2\pi(1-a)) = \text{Im } \Phi(a, x_0(a))$.

Remark 4.3. We can show the following statement by the same procedure that appeared in [Murakami 2004]²: Let $a \in (\frac{5}{12}, \frac{1}{2})$ be the limit of $a(N)/N$, where $N \rightarrow \infty$. Then, the limit

$$4\pi \lim_{N \rightarrow \infty} \frac{\log |J_{a(N)}(4_1; \xi_N^2)|}{N}$$

is equal to the volume of the cone-manifold of the figure-eight knot with a cone-angle $2\pi - 4\pi a$, where N runs over all odd integers.

²In [Murakami 2004], the value substituted for t is slightly changed from the N -th root of unity.

Example 4.4 (Borromean rings). Let K_B be the Borromean rings, $K_B(\alpha, \beta, \gamma)$ be the cone manifold of K_B with cone-angles α, β, γ , and $\Delta(\alpha, \theta) = \Lambda(\alpha + \theta) - \Lambda(\alpha - \theta)$, where $\Lambda(x)$ is the Lobachevsky function. If $0 < \alpha, \beta, \gamma < \pi$, then $K_B(\alpha, \beta, \gamma)$ admits a hyperbolic structure, and its volume is given by

$$\text{Vol } K_B(\alpha, \beta, \gamma) = 2\left(\Delta\left(\frac{\alpha}{2}, \theta\right) + \Delta\left(\frac{\beta}{2}, \theta\right) + \Delta\left(\frac{\gamma}{2}, \theta\right) - 2\Delta\left(\frac{\pi}{2}, \theta\right) - \Delta(0, \theta)\right),$$

where $\theta \in (0, \frac{\pi}{2})$ is defined by the following conditions [Mednykh 2003]:

$$\begin{aligned} T = \tan \theta, \quad L = \tan \frac{\alpha}{2}, \quad M = \tan \frac{\beta}{2}, \quad N = \tan \frac{\gamma}{2}, \\ T^4 - (L^2 + M^2 + N^2 + 1)T^2 - L^2M^2N^2 = 0. \end{aligned}$$

We define the function $\tilde{\Delta}(x, y, z, \theta)$ by

$$\tilde{\Delta}(x, y, z, \theta) = 2\left(\Delta(x, \theta) + \Delta(y, \theta) + \Delta(z, \theta) - 2\Delta\left(\frac{\pi}{2}, \theta\right) - \Delta(0, \theta)\right)$$

for convenience. On the other hand, the colored Jones polynomial for K_B is given by [Habiro 2000]

$$J_{(l,m,n)}(K_B; t) = \sum_{i=1}^{\min(l,m,n)-1} \frac{\{l+i\}! \{m+i\}! \{n+i\}! (\{i\}!)^2}{\{1\}! \{l-i-1\}! \{m-i-1\}! \{n-i-1\}! (\{2i+1\}!)^2}.$$

Let a, b , and c be the limit of $l/N, m/N$, and n/N , respectively. The potential function $\Phi_{K_B}(x)$ of $J_{(l,m,n)}(K_B; \xi_N)$ is

$$\begin{aligned} \Phi_{K_B}(a, b, c, x) = -2\pi\sqrt{-1}(a+b+c)\log x + \frac{3}{2}(\log x)^2 \\ - \text{Li}_2(e_a^2x) - \text{Li}_2(e_b^2x) - \text{Li}_2(e_c^2x) - 2\text{Li}_2(x) \\ + \text{Li}_2\left(\frac{e_a^2}{x}\right) + \text{Li}_2\left(\frac{e_b^2}{x}\right) + \text{Li}_2\left(\frac{e_c^2}{x}\right) + 2\text{Li}_2(x^2). \end{aligned}$$

The derivative of $\Phi_{K_B}(x)$ with respect to x is

$$x \frac{\partial \Phi_{K_B}}{\partial x} = \log\left(e_a^{-2}e_b^{-2}e_c^{-2}F(a, x)F(b, x)F(c, x) \frac{x^3(1-x)^2}{(1-x^2)^4}\right),$$

where $F(a, x) = (1 - e_a^2x)(1 - e_a^2/x)$. Under the substitution $x = e^{2\pi\sqrt{-1}\zeta}$, we obtain

$$\begin{aligned} \frac{1}{2\pi\sqrt{-1}} \frac{\partial \Phi_{K_B}}{\partial \zeta} \\ = \log \frac{\sin \pi(\zeta+a) \sin \pi(\zeta-a) \sin \pi(\zeta+b) \sin \pi(\zeta-b) \sin \pi(\zeta+c) \sin \pi(\zeta-c)}{\sin^2 \pi \zeta \cos^4 \pi \zeta} \\ = \log \frac{\tan^2 \pi \zeta - A^2}{1+A^2} \frac{\tan^2 \pi \zeta - B^2}{1+B^2} \frac{\tan^2 \pi \zeta - C^2}{1+C^2} \frac{1}{\tan^2 \pi \zeta}, \end{aligned}$$

where $A = \tan \pi(1 - a)$, $B = \tan \pi(1 - b)$, and $C = \tan \pi(1 - c)$. Therefore, if $\pm \tan \pi \zeta$ are solutions of the equation

$$\frac{t^2 - A^2}{1 + A^2} \frac{t^2 - B^2}{1 + B^2} \frac{t^2 - C^2}{1 + C^2} \frac{1}{t^2} = 1,$$

which is equivalent to the equation

$$(t^2 + 1)(t^4 - (A^2 + B^2 + C^2 + 1)t^2 - A^2 B^2 C^2) = 0,$$

then $x = e^{2\pi\sqrt{-1}\zeta}$ is a saddle point of $\Phi_{K_B}(a, b, c, x)$. By using the properties of the Lobachevsky function, such as

$$\begin{aligned} \text{Li}_2(e^{2\sqrt{-1}\theta}) &= \frac{\pi^2}{6} - \theta(\pi - \theta) + 2\sqrt{-1}\Lambda(\theta), \\ \Lambda(2\theta) &= 2\Lambda(\theta) + 2\Lambda\left(\theta + \frac{\pi}{2}\right), \end{aligned}$$

we obtain

$$\begin{aligned} \text{Im } \Phi_{K_B}(a, b, c, e^{2\pi\sqrt{-1}\zeta}) &= \tilde{\Delta}(\pi(1 - a), \pi(1 - b), \pi(1 - c), \pi(1 - \zeta)) \\ &= \text{Vol } K_B(2\pi(1 - a), 2\pi(1 - b), 2\pi(1 - c)). \end{aligned}$$

5. The completeness condition

In the previous section, we fixed a_1, \dots, a_n . In this section, we regard them as variables and find a geometric meaning. First, we consider the case where a crossing is between different components. The derivatives of the potential function with respect to the parameters corresponding to the colors are:

$$\begin{aligned} \frac{\partial \Phi_c^+}{\partial a} &= \frac{\pi\sqrt{-1}}{2} \log\left(1 - e_a \frac{w_{j_4}}{w_{j_3}}\right) \left(1 - e_a \frac{w_{j_1}}{w_{j_2}}\right)^{-1} \left(1 - e_a^{-1} \frac{w_{j_3}}{w_{j_4}}\right) \left(1 - e_a^{-1} \frac{w_{j_2}}{w_{j_1}}\right)^{-1}, \\ \frac{\partial \Phi_c^+}{\partial b} &= \frac{\pi\sqrt{-1}}{2} \log\left(1 - e_b \frac{w_{j_4}}{w_{j_1}}\right) \left(1 - e_b \frac{w_{j_3}}{w_{j_2}}\right)^{-1} \left(1 - e_b^{-1} \frac{w_{j_1}}{w_{j_4}}\right) \left(1 - e_b^{-1} \frac{w_{j_2}}{w_{j_3}}\right)^{-1}, \\ \frac{\partial \Phi_c^-}{\partial a} &= \frac{\pi\sqrt{-1}}{2} \log\left(1 - e_a \frac{w_{j_1}}{w_{j_4}}\right) \left(1 - e_a \frac{w_{j_2}}{w_{j_3}}\right)^{-1} \left(1 - e_a^{-1} \frac{w_{j_4}}{w_{j_1}}\right) \left(1 - e_a^{-1} \frac{w_{j_3}}{w_{j_2}}\right)^{-1}, \\ \frac{\partial \Phi_c^-}{\partial b} &= \frac{\pi\sqrt{-1}}{2} \log\left(1 - e_b \frac{w_{j_3}}{w_{j_4}}\right) \left(1 - e_b \frac{w_{j_2}}{w_{j_1}}\right)^{-1} \left(1 - e_b^{-1} \frac{w_{j_4}}{w_{j_3}}\right) \left(1 - e_b^{-1} \frac{w_{j_1}}{w_{j_2}}\right)^{-1}. \end{aligned}$$

We can observe the correspondence between these derivatives and dilation components by cusp diagrams (Figure 5.1). In Figure 5.1, $\partial \Phi_c^+ / \partial a$ corresponds to the upper side of a positive crossing (top left), $\partial \Phi_c^+ / \partial b$ to the lower side of a positive crossing (top right), $\partial \Phi_c^- / \partial a$ to the upper side of a negative crossing (bottom left), and $\partial \Phi_c^- / \partial b$ to the upper side of a negative crossing (bottom right). A similar correspondence holds in the case where a crossing is between the same component.




Figure 5.1. Cusp diagrams: upper side of a positive crossing (top left), lower side of a positive crossing (top right), upper side of a negative crossing (bottom left), and lower side of a negative crossing (bottom right).

Let l_i be the longitude that is parallel to the component L_i , and let \tilde{l}_i be the longitude of the component L_i with $\text{lk}(\tilde{l}_i, L_i) = 0$. For a curve γ on the cusp diagram, we define $\delta(\gamma)$ as the dilation component of γ . Then, by the above observation

$$\exp\left(\frac{1}{\pi\sqrt{-1}} \frac{\partial \Phi'_D}{\partial a_i}\right) = \exp\left(\frac{1}{2} \log \delta(l_i)^2\right) = \delta(l_i),$$

where Φ'_D is a potential function of the colored Jones polynomial without the modification for the Reidemeister move I. Next, we consider the contribution of the modification. For a positive crossing between the same component with a parameter a , the modification corresponds to the addition of $(\pi\sqrt{-1}a)^2$, and its derivative is

$$\frac{1}{\pi\sqrt{-1}} \frac{d}{da} (\pi\sqrt{-1}a)^2 = 2\pi\sqrt{-1}a = \log e_a^2.$$

Here, e_a^2 is equal to the dilation component of the meridian with the inverse orientation. Similarly, for a negative crossing, the derivative of $-(\pi\sqrt{-1}a)^2$ corresponds to the dilation component of the meridian. Therefore,

$$(5.1) \quad \exp\left(\frac{1}{\pi\sqrt{-1}} \frac{\partial \Phi_D}{\partial a_i}\right) = \delta(\tilde{l}_i).$$




Figure 5.2. Cusp diagrams of a knot complement: upper side of a positive crossing (top left), lower side of a positive crossing (top right), upper side of a negative crossing (bottom left), lower side of a negative crossing (bottom right).

Remark 5.1. If K is a knot, we have a more simple correspondence. The derivatives of Φ_c^\pm with respect to a are:

$$(5.2) \quad \frac{1}{\pi\sqrt{-1}} \frac{\partial \Phi_c^+}{\partial a} = \log e_a^2 + \log \left(1 - e_a^{-1} \frac{w_{j_3}}{w_{j_4}} \right) \left(1 - e_a \frac{w_{j_4}}{w_{j_1}} \right) \left(1 - e_a^{-1} \frac{w_{j_2}}{w_{j_1}} \right)^{-1} \left(1 - e_a \frac{w_{j_3}}{w_{j_2}} \right)^{-1},$$

$$(5.3) \quad \frac{1}{\pi\sqrt{-1}} \frac{\partial \Phi_c^-}{\partial a} = \log e_a^{-2} + \log \left(1 - e_a^{-1} \frac{w_{j_4}}{w_{j_1}} \right) \left(1 - e_a \frac{w_{j_3}}{w_{j_4}} \right) \left(1 - e_a^{-1} \frac{w_{j_3}}{w_{j_2}} \right)^{-1} \left(1 - e_a \frac{w_{j_2}}{w_{j_1}} \right)^{-1}.$$

The second term of (5.2) corresponds to the upper side and the lower side of a positive crossing (Figure 5.2, top left and right), and the second term of (5.3) corresponds to the upper side and the lower side of a negative crossing (Figure 5.2, bottom left and right)

Remark 5.2. Changing the variable a_i to $u_i = 2\pi\sqrt{-1}a_i$, we have

$$2 \frac{\partial \Phi_D}{\partial u_i} = \frac{1}{\pi\sqrt{-1}} \frac{\partial \Phi_D}{\partial a_i}.$$

Then,

$$\Psi(\mathbf{u}) = 4(\Phi_D(\mathbf{u}, \sigma_1(\mathbf{u}), \dots, \sigma_\nu(\mathbf{u})) - \Phi_D(\mathbf{0}, \sigma_1(\mathbf{0}), \dots, \sigma_\nu(\mathbf{0}))),$$

where $\mathbf{u} = (u_1, \dots, u_n)$ and $\mathbf{0} = (0, \dots, 0)$ satisfy the conditions of the Neumann–Zagier potential function [1985]. Namely, Ψ satisfies $\Psi(\mathbf{0}) = 0$ and

$$\frac{1}{2} \frac{\partial \Psi}{\partial u_i} = \log \delta(\tilde{l}_i).$$

If $a_i = 1$, with $i = 1, \dots, n$, all the dilation components of meridians are 1. Furthermore, the contributions of parts, as shown in Figure 4.3, to the dilation component of the longitude is 1, hence $\delta(\tilde{l}_i) = 1$, $(1, \dots, n)$. Therefore, the point $(\mathbf{1}, \sigma_1(\mathbf{1}), \dots, \sigma_\nu(\mathbf{1}))$ gives a complete hyperbolic structure to the link complement [Benedetti and Petronio 1992], where $\mathbf{1} = (1, \dots, 1)$. Moreover, by (4.3) and (5.1) the point is a solution of the following system of equations:

$$\begin{cases} \exp\left(w_i \frac{\partial \Phi_D}{\partial w_i}\right) = 1, & i = 1, \dots, \nu, \\ \exp\left(\frac{1}{\pi\sqrt{-1}} \frac{\partial \Phi_D}{\partial a_j}\right) = 1, & j = 1, \dots, n. \end{cases}$$

Hence, we obtain the following theorem:

Theorem 5.3. *Let D be a diagram of a hyperbolic link with n components, and let $\mathbf{1}$ be $(1, \dots, 1) \in \mathbb{Z}^n$. The point $(\mathbf{1}, \sigma_1(\mathbf{1}), \dots, \sigma_\nu(\mathbf{1}))$ is a saddle point of the function $\Phi_D(a_1, \dots, a_n, w_1, \dots, w_\nu)$ and gives a complete hyperbolic structure to the link complement.*

6. The Witten–Reshetikhin–Turaev invariant

In [Kirby and Melvin 1991], the Witten–Reshetikhin–Turaev invariant for the manifold obtained by Dehn surgery on a link is stated. Furthermore, Murakami [2000] calculated the optimistic limit of the Witten–Reshetikhin–Turaev invariant for the manifold obtained by integer surgery on the figure-eight knot. By a similar argument as in Section 4, we would be able to explain the correspondence of the Witten–Reshetikhin–Turaev invariant and the geometry of the manifold obtained by Dehn surgery on a link. The procedure might be as follows: The Witten–Reshetikhin–Turaev invariant for the manifold M_{f_1, \dots, f_n} obtained by Dehn surgery on a link $L = L_1 \cup \dots \cup L_n$ with a framing f_i on L_i , where $i = 1, \dots, n$, can be written as a summation of the colored Jones polynomial $J_{\mathbf{k}}(L; \xi_N)$ multiplied by $t^{-(1/4)\sum f_j k_j^2}$, where $\mathbf{k} = (k_1, \dots, k_n)$ are colors of L . See [Kirby and Melvin 1991] for details, but note that t in [Kirby and Melvin 1991] and t in this paper are different. We suppose that M_{f_1, \dots, f_n} admits a hyperbolic structure. Let α_i be $e^{\pi\sqrt{-1}a_i}$ and regard




Figure 6.1. The schematic diagram of the developing image in the case of $f_i = 6$.

it as a complex parameter that is not necessarily in the unit circle. Then, we have

$$\frac{1}{\pi\sqrt{-1}} \frac{\partial\Phi}{\partial\alpha_i} = \alpha_i \frac{\partial\Phi}{\partial\alpha_i}.$$

Multiplying $t^{-(1/4)\sum f_j k_j^2}$ leads to the addition of $-\sum f_j (\log \alpha_j)^2$ to the potential function. The derivative of it with respect to α_i is

$$\alpha_i \frac{\partial}{\partial\alpha_i} \left(-\sum_{j=1}^n f_j (\log \alpha_j)^2 \right) = -2f_i \log \alpha_i = \log \alpha_i^{-2f_i}.$$

Then, the saddle point equation is equivalent to the system of equations consisting of the gluing condition and

$$\delta(\tilde{l}_i) = \alpha_i^{2f_i}, \quad i = 1, \dots, n.$$

Recall that the dilation component of the meridian m_i of L_i is α_i^{-2} , which implies that $\delta(m_i)^{-f_i} = \delta(\tilde{l}_i)$. If we suppose that $|\alpha_i|$ is less than 1 and f_i is a positive integer, the developing image would be as shown in Figure 6.1. By filling in the singular set, the developing image becomes complete.

Acknowledgment

The author is grateful to Jun Murakami for his helpful comments.

References

- [Belletti et al. 2022] G. Belletti, R. Detcherry, E. Kalfagianni, and T. Yang, “Growth of quantum $6j$ -symbols and applications to the volume conjecture”, *J. Differential Geom.* **120**:2 (2022), 199–229. [MR Zbl](#)
- [Benedetti and Petronio 1992] R. Benedetti and C. Petronio, *Lectures on hyperbolic geometry*, Springer, Berlin, 1992. [MR Zbl](#)

- [Chen and Yang 2018] Q. Chen and T. Yang, “Volume conjectures for the Reshetikhin–Turaev and the Turaev–Viro invariants”, *Quantum Topol.* **9**:3 (2018), 419–460. [MR](#) [Zbl](#)
- [Cho 2016a] J. Cho, “Optimistic limit of the colored Jones polynomial and the existence of a solution”, *Proc. Amer. Math. Soc.* **144**:4 (2016), 1803–1814. [MR](#) [Zbl](#)
- [Cho 2016b] J. Cho, “Optimistic limits of the colored Jones polynomials and the complex volumes of hyperbolic links”, *J. Aust. Math. Soc.* **100**:3 (2016), 303–337. [MR](#) [Zbl](#)
- [Cho and Murakami 2013] J. Cho and J. Murakami, “Optimistic limits of the colored Jones polynomials”, *J. Korean Math. Soc.* **50**:3 (2013), 641–693. [MR](#) [Zbl](#)
- [Detcherry et al. 2018] R. Detcherry, E. Kalfagianni, and T. Yang, “Turaev–Viro invariants, colored Jones polynomials, and volume”, *Quantum Topol.* **9**:4 (2018), 775–813. [MR](#) [Zbl](#)
- [Habiro 2000] K. Habiro, “On the colored Jones polynomials of some simple links”, *RIMS Kōkyūroku* **1172** (2000), 34–43. In Japanese. [MR](#) [Zbl](#)
- [Kashaev 1997] R. M. Kashaev, “The hyperbolic volume of knots from the quantum dilogarithm”, *Lett. Math. Phys.* **39**:3 (1997), 269–275. [MR](#) [Zbl](#)
- [Kashaev and Tirkkonen 2000] R. M. Kashaev and O. Tirkkonen, “Proof of the volume conjecture for torus knots”, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **269**:Vopr. Kvant. Teor. Polya i Stat. Fiz. 16 (2000), 262–268, 370. In Japanese; translated in *J. Math. Sci.* **115**:1 (2003), 2033–2036. [MR](#) [Zbl](#)
- [Kirby and Melvin 1991] R. Kirby and P. Melvin, “The 3-manifold invariants of Witten and Reshetikhin–Turaev for $sl(2, \mathbb{C})$ ”, *Invent. Math.* **105**:3 (1991), 473–545. [MR](#)
- [Mednykh 2003] A. D. Mednykh, “On hyperbolic and spherical volumes for knot and link cone-manifolds”, pp. 145–163 in *Kleinian groups and hyperbolic 3-manifolds* (Warwick, 2001), edited by Y. Komori et al., London Math. Soc. Lecture Note Ser. **299**, Cambridge Univ. Press, 2003. [MR](#) [Zbl](#)
- [Mednykh and Rasskazov 2006] A. Mednykh and A. Rasskazov, “Volumes and degeneration of cone-structures on the figure-eight knot”, *Tokyo J. Math.* **29**:2 (2006), 445–464. [MR](#) [Zbl](#)
- [Murakami 2000] H. Murakami, “Optimistic calculations about the Witten–Reshetikhin–Turaev invariants of closed three-manifolds obtained from the figure-eight knot by integral Dehn surgeries”, *RIMS Kōkyūroku* **1172** (2000), 70–79. In Japanese. [MR](#) [Zbl](#)
- [Murakami 2004] H. Murakami, “Some limits of the colored Jones polynomials of the figure-eight knot”, *Kyungpook Math. J.* **44**:3 (2004), 369–383. [MR](#) [Zbl](#)
- [Murakami and Murakami 2001] H. Murakami and J. Murakami, “The colored Jones polynomials and the simplicial volume of a knot”, *Acta Math.* **186**:1 (2001), 85–104. [MR](#) [Zbl](#)
- [Neumann and Zagier 1985] W. D. Neumann and D. Zagier, “Volumes of hyperbolic three-manifolds”, *Topology* **24**:3 (1985), 307–332. [MR](#) [Zbl](#)
- [Thurston 1999] D. Thurston, “Hyperbolic volume and the Jones polynomial”, lecture notes, 1999, available at <https://dpthurst.pages.iu.edu/speaking/Grenoble.pdf>.
- [Yokota 2000] Y. Yokota, “On the volume conjecture for hyperbolic knots”, preprint, 2000. [arXiv math/0009165](https://arxiv.org/abs/math/0009165)
- [Yokota 2003] Y. Yokota, “From the Jones polynomial to the A-polynomial of hyperbolic knots”, *Interdiscip. Inform. Sci.* **9**:1 (2003), 11–21. [MR](#) [Zbl](#)
- [Yoon 2021] S. Yoon, “On the potential functions for a link diagram”, *J. Knot Theory Ramifications* **30**:7 (2021), art. id. 2150056. [MR](#)
- [Zagier 2007] D. Zagier, “The dilogarithm function”, pp. 3–65 in *Frontiers in number theory, physics, and geometry, II* (Les Houches, 2003), edited by P. Cartier et al., Springer, Berlin, 2007. [MR](#) [Zbl](#)

SHUN SAWABE
DEPARTMENT OF PURE AND APPLIED MATHEMATICS
SCHOOL OF FUNDAMENTAL SCIENCE AND ENGINEERING
WASEDA UNIVERSITY
TOKYO
JAPAN
sa-shun.1729ttw@asagi.waseda.jp

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2023 is US \$605/year for the electronic version, and \$820/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 322 No. 1 January 2023

Rips construction without unique product	1
GOULNARA ARZHANTSEVA and MARKUS STEENBOCK	
No periodic geodesics in jet space	11
ALEJANDRO BRAVO-DODDOLI	
Coarse geometry of Hecke pairs and the Baum–Connes conjecture	21
CLÉMENT DELL’AIERA	
On homology theories of cubical digraphs	39
ALEXANDER GRIGOR’YAN and YURI MURANOV	
The geometry and topology of stationary multiaxisymmetric vacuum black holes in higher dimensions	59
VISHNU KAKKAT, MARCUS KHURI, JORDAN RAINONE and GILBERT WEINSTEIN	
Quasilinear Schrödinger equations: ground state and infinitely many normalized solutions	99
HOUWANG LI and WENMING ZOU	
Thomae’s function on a Lie group	139
MARK REEDER	
On the potential function of the colored Jones polynomial with arbitrary colors	171
SHUN SAWABE	
Pushforward and smooth vector pseudo-bundles	195
ENXIN WU	