

*Pacific
Journal of
Mathematics*

**A LIFT OF WEST'S STACK-SORTING MAP
TO PARTITION DIAGRAMS**

JOHN M. CAMPBELL

Volume 324 No. 2

June 2023

A LIFT OF WEST'S STACK-SORTING MAP TO PARTITION DIAGRAMS

JOHN M. CAMPBELL

We introduce a lifting of West's stack-sorting map s to partition diagrams, which are combinatorial objects indexing bases of partition algebras. Our lifting \mathcal{S} of s is such that \mathcal{S} behaves in the same way as s when restricted to diagram basis elements in the order- n symmetric group algebra as a diagram subalgebra of the partition algebra \mathcal{P}_n^ξ . We then introduce a lifting of the notion of 1-stack-sortability, using our lifting of s . By direct analogy with Knuth's famous result that a permutation is 1-stack-sortable if and only if it avoids the pattern 231, we prove a related pattern-avoidance property for partition diagrams, as opposed to permutations, according to what we refer to as *stretch-stack-sortability*.

1. Introduction

For a permutation p in the symmetric group S_n , we write

$$(1) \quad p = LnR,$$

letting p be denoted as a string or tuple given by the entries of the bottom row of the two-line notation for p . We then let *West's stack-sorting map* s [6; 7; 8; 9; 10; 11; 12; 13; 14] (compare [27]) be defined recursively so that

$$(2) \quad s(p) = s(L)s(R)n$$

and so that s maps permutations to permutations and sends the empty permutation to itself. Our notation and terminology concerning this mapping are mainly based on references such as [6; 7; 8; 9; 10; 11; 12; 13; 14]. In this article, we introduce a lifting of s so as to allow combinatorial objects known as *partition diagrams* as input.

The problem of generalizing West's stack-sorting map has been considered in a number of different contexts. Notably, the stack-sorting map s has been

MSC2020: primary 05A05; secondary 05E16.

Keywords: stack-sorting, partition diagram, permutation, permutation pattern, partition monoid.

generalized to Coxeter groups [12] and to words [13], and Cerbai, Claesson, and Ferrari generalized

$$(3) \quad s : S_n \rightarrow S_n$$

to a function $s_\sigma : S_n \rightarrow S_n$ for a permutation pattern σ [5], so that $s = s_{21}$, with a related generalization of s having been given by Defant and Zheng in [14]. Since s is defined on permutations, it is natural to consider generalizing this mapping using “permutation-like” combinatorial objects. The bases of the partition algebra \mathcal{P}_n^ξ generalize the bases of the order- n symmetric group algebra in ways that are of interest from both a combinatorial and an algebraic perspective, and this leads us to consider how partition diagrams, which index the bases of \mathcal{P}_n^ξ , may be used to generalize West’s stack-sorting map. We generalize, in this article, $s : S_n \rightarrow S_n$ to a mapping $\mathcal{S} : \mathcal{P}_n \rightarrow \mathcal{P}_n$ from the partition monoid \mathcal{P}_n to itself such that \mathcal{S} restricted to permuting diagrams behaves in the same way as s . We then apply our lifting \mathcal{S} to determine an analogue of a famous result due to Knuth [20, Section 2.2.1] on 1-stack-sortable permutations.


The representation theory of \mathcal{P}_n^ξ is intimately linked with that for the $n!$ -dimensional symmetric group algebra. Indeed, there is so much about the representation theory of \mathcal{P}_n^ξ and associated combinatorial properties of \mathcal{P}_n^ξ that are directly derived from or otherwise based on the representation theory of the algebra $\text{span}(S_n)$ and associated combinatorics [2; 3; 17; 18; 22; 23; 25]. The Schur–Weyl duality given by how the bases of \mathcal{P}_n^ξ generalize the bases of $\text{span}(S_n)$ is of importance in both combinatorial representation theory and in the areas of statistical mechanics in which partition algebras had originally been defined by Martin [21; 22; 23; 24] and Jones [19]. The foregoing considerations are representative of the extent to which partition diagrams generalize permutations in a way that is of much significance in both mathematics and physics. This motivates our lifting West’s stack-sorting map so as to allow partition diagrams, in addition to permutations, as input.

Preliminaries. To be consistent with the notation in references as in [6; 7; 8; 9; 10; 11; 12; 13; 14] for indexing symmetric groups and maximal elements in permutations, as in (2) and (3), we let the order of a given partition algebra/monoid be denoted as in the order of the symmetric group shown in (3). In particular, following [16], we let the partition monoid of a given order be denoted as \mathcal{P}_n , and we let the corresponding partition algebra with a complex parameter ξ be denoted as \mathcal{P}_n^ξ . These structures are defined as follows.


We let \mathcal{P}_n consist of set-partitions μ of $\{1, 2, \dots, n, 1', 2', \dots, n'\}$ that we denote with a two-line notation by analogy with permutations, by aligning nodes labeled with $1, 2, \dots, n$ into an upper row and nodes labeled with $1', 2', \dots, n'$ into a bottom row, and by forming any graph G such that the set of components of G

equals μ . Graphs of this form, denoted in the manner we have specified, are referred to as *partition diagrams*, and two such partition diagrams are considered to be the same if the connected components are the same in both cases. The components of the graph G given as before are referred to as *blocks*.

Example 1. The set-partition $\{\{1, 4\}, \{2, 3, 4', 5'\}, \{5\}, \{1', 3'\}, \{2'\}\}$ may be denoted as




or, equivalently, as




Following [18], we let the underlying field for partition algebras and symmetric group algebras be \mathbb{C} , as \mathbb{C} being algebraically closed is of direct relevance in terms of the study of the semisimple structure for partition algebras [18]. We may let the symmetric group algebra of order n be denoted by taking the linear span $\text{span}(S_n)$, or, more explicitly, $\text{span}_{\mathbb{C}}\{\sigma : \sigma \in S_n\}$, of the symmetric group S_n .

For two partition diagrams d_1 and d_2 , we place d_1 on top of d_2 so that the bottom nodes of d_1 overlap with the top nodes of d_2 , then we remove the central row in this concatenation $d_1 * d_2$ in such a way so as to preserve the relation given by topmost nodes being in the same component as bottommost nodes in $d_1 * d_2$. We then let $d_1 \circ d_2$ denote the graph thus obtained from this concatenation. This is the underlying multiplicative operation for partition monoids.

Example 2. Borrowing an example from [16], we let d_1 be as in Example 1, and we let d_2 be as below:



We may verify that the monoid product $d_1 \circ d_2$, in this case, is as below [16]:



For a complex parameter ξ , we endow the \mathbb{C} -span of \mathcal{P}_n with a multiplicative binary operation as $d_1 d_2 = \xi^\ell d_1 \circ d_2$, where ℓ denotes the number of components contained entirely in the middle row of $d_1 * d_2$. By extending this binary operation linearly, this gives us the underlying multiplicative operation for the structure known as the partition algebra, which is denoted as \mathcal{P}_n^ξ . The set of all elements in \mathcal{P}_n , as elements in \mathcal{P}_n^ξ , is referred to as the *diagram basis* of \mathcal{P}_n^ξ .

Although the algebraic structure of \mathcal{P}_n^ξ will not be used directly in this article, it is useful to define \mathcal{P}_n^ξ explicitly, since we are to heavily make use of an algebra homomorphism defined on partition algebras and introduced in [4], and it is often convenient to use notation and terminology associated with \mathcal{P}_n^ξ or its algebraic structure, e.g., by referring to the diagram basis of \mathcal{P}_n^ξ .

For a partition diagram π , the *propagation number* of π refers to the number of blocks in π containing at least one vertex in the top row and at least one vertex in the bottom row. So, there is a clear bijection between the set of all permutations in S_n and the set of all diagrams in \mathcal{P}_n that are of propagation number n . In our lifting West’s stack-sorting map to the partition monoid, to be consistent with two-line notation for permutations and with (1), we let a permutation

$$(4) \quad p: \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$$

be in correspondence with the partition diagram π given by the set-partition

$$(5) \quad \{\{1, p(1)'\}, \{2, p(2)'\}, \dots, \{n, p(n)'\}\},$$

although it is common to instead let a permutation p be mapped to the partition diagram obtained by reflecting π vertically.

2. A lift of West’s stack-sorting map


As Defant and Kravitz explain in [13], there is a matter of ambiguity in the problem of lifting the mapping s so as to allow words involving repeated characters. We encounter similar kinds of problems in terms of the problem of lifting s so as to allow partition diagrams, as opposed to permutations written as permutation diagrams, as the argument. This is illustrated below.

Example 3. According to the embedding indicated in (5), we let the permutation $\begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}$ be written as



If we visualize the block $\{2, 3'\}$ being removed and placed on the right of a resultant configuration, by analogy with how the permutation (1) gets mapped to the right-hand side of (2) under the application of West’s stack-sorting map, there is a clear way how this removal “splits” the permutation diagram in (6) into a left and a right configuration by direct analogy with (1), and part of the purpose of our procedure in the next subsection is to formalize this idea in a way that may be extended to all

partition diagrams. In contrast, if we remove, for example, the rightmost block of



where the above coloring is to distinguish the blocks in the partition diagram depicted, then the block highlighted in green is not strictly to the left of the rightmost block, since there are green nodes to the left and right of a red node in the upper row.

To lift the recursive definition for s indicated in (2) so as to allow members of \mathcal{P}_n as the input for an extension or lifting of s , and to deal with the matter of ambiguity explained in Example 3, we would want to mimic (2) by allowing the possibility of “middle” configurations, as opposed to the left-right dichotomy indicated in (1) and (2). We formalize this idea below.

A procedure for stack-sorting partition diagrams. Define $\mathcal{S}_n: \mathcal{P}_n \rightarrow \mathcal{P}_n$ according to the following procedure, for an arbitrary partition diagram π in \mathcal{P}_n . We order the bottom nodes of a given partition diagram in \mathcal{P}_n in the natural way, with $1' < 2' < \dots < n'$. For the sake of convenience, we may write $\mathcal{S} = \mathcal{S}_n$.

- (1) If there are no propagating blocks in π , skip the below steps involving propagating blocks.
- (2) Take the largest bottom node of π that is part of a propagating block B . This propagating block separates π into three (possibly empty) classes of configurations according to how the top nodes of π are separated by removing B . Explicitly, we define $\mathcal{L}, \mathcal{M}_1, \mathcal{M}_2, \dots, \mathcal{M}_\mu$, and \mathcal{R} as follows, by direct analogy with (1). We may denote $\mathcal{L}, \mathcal{M}_i$, and \mathcal{R} as diagrams in \mathcal{P}_n . The blocks of \mathcal{L} (resp. \mathcal{R}) consist of one of the following:

- Any blocks of π with upper nodes that are all strictly to the left (resp. right) of all of the upper nodes of B .
- Any nonpropagating blocks of π on the bottom row of π with nodes that are all strictly to the left (resp. right) of all of the lower nodes of B .
- Singleton blocks.

The blocks of expressions of the form \mathcal{M}_i consist either of

- any blocks of π that do not satisfy either of the first two bullet points listed above, or
- singleton blocks.

We order $\mathcal{M}_1 < \mathcal{M}_2 < \dots < \mathcal{M}_\mu$ according to the ordering of the minimal elements, subject the ordering whereby $1' < 2' < \dots < n' < 1 < 2 < \dots < n$. We

then let B' denote the partition diagram obtained from B by adding any singleton nodes so as to form a partition diagram in \mathcal{P}_n . With this setup, we write

$$(8) \quad \mathcal{S}(\pi) = \mathcal{S}(\mathcal{L}) \odot \mathcal{S}(\mathcal{M}_1) \odot \cdots \odot \mathcal{S}(\mathcal{M}_\mu) \odot \mathcal{S}(\mathcal{R}) \odot B'$$

where the associative binary relation \odot is to later be defined.

(3) Repeatedly apply the above step wherever possible, i.e., to the expressions in or derived from (8) given by \mathcal{S} evaluated at a partition diagram.


(4) The above steps yield a \odot -product of expressions of the forms

$$(9) \quad \mathcal{S}(N_1), \dots, \mathcal{S}(N_{m_1}) \quad \text{and} \quad B'_1, \dots, B'_{m_2},$$

not necessarily in this order, for nonpropagating diagrams N_i , and where each expression of the form B'_j is a partition diagram with exactly one propagating block and with singleton blocks anywhere else. The factors of the aforementioned \odot -product indicated in (9) are ordered in the following way: B'_j is the j -th factor of the form B'_κ appearing in this \odot -product, and $\mathcal{S}(N_i)$ is the i -th factor of the form $\mathcal{S}(N_\kappa)$ appearing in this \odot -product. The operation \odot indicates that the following is to be applied. We label the top nodes in the propagating block in B'_1 from left to right with consecutive integers starting with 1, and we then label the top nodes in the propagating block in B'_2 with consecutive integers (starting with 1 plus the number of top nodes in the propagating block in B'_1), and we continue in this manner. We then continue with this labeling, by labeling any nonpropagating blocks of size greater than 1 in the top row of the N_i -expressions, in order of the nodes as they appear among consecutive N_i -diagrams. If there are any unused labels for the top row, label singleton blocks in the upper row with these leftover labels. However, for the bottom nodes of any nonsingleton blocks from π , we let these bottom nodes keep their original labelings (and if necessary we add in singleton blocks in a bottom row to form a partition diagram based on the preceding steps).

(5) As indicated above, the above steps produce an element in \mathcal{P}_n . We set $\mathcal{S}(\pi)$ to be this element.

Example 4. Let π denote the partition diagram




in \mathcal{P}_8 corresponding to the set-partition


$$\{\{1, 2\}, \{3, 5, 7, 2', 4', 6'\}, \{4, 3'\}, \{6, 7'\}, \{8\}, \{1'\}, \{5', 8'\}\}.$$

The largest bottom node of π that is part of a propagating block B , in this case, is $7'$, and B is $\{6, 7'\}$. For the sake of clarity, we highlight this block in the following


manner:




The blocks of \mathcal{L} are highlighted in cyan as below (with the propagating block B again highlighted in a different color for the sake of clarity).



We find that there is only one expressions of the form \mathcal{M}_i , and we may denote this expression as \mathcal{M} . The blocks of \mathcal{M} are highlighted in green as below.



We see that \mathcal{R} consists of only one block, which is highlighted in orange, as below.



So, according to our lifting of West's stack-sorting map, we obtain the following, where singleton nodes colored black indicate that these singleton nodes have been "added in" according to the above given procedure.

$$\mathcal{S}(\pi) =$$

$$\mathcal{S} \left(\begin{array}{cccccccc} \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \right) \odot$$

$$\mathcal{S} \left(\begin{array}{cccccccc} \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \right) \odot$$

$$\mathcal{S} \left(\begin{array}{cccccccc} \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \right) \odot$$

$$\left(\begin{array}{cccccccc} \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \right)$$

Now, let us repeat the given procedure to the first \mathcal{S} -factor on the right-hand side of the above equality, and then to the resultant \mathcal{S} -factor involving an argument


with a block of size 6. This results in the following equality:

$$\mathcal{S}(\pi) =$$


The diagram shows ten 8x2 grids of dots, each preceded by a script \mathcal{S} and followed by a circled dot. The grids contain various colored dots and connecting lines:

- Grid 1: Two blue dots in the top row, connected by a curved line. A blue dot in the bottom row.
- Grid 2: All dots are black.
- Grid 3: All dots are black.
- Grid 4: A blue dot in the top row, connected by a straight line to a blue dot in the bottom row.
- Grid 5: All dots are black.
- Grid 6: Two green dots in the bottom row, connected by a curved line.
- Grid 7: All dots are black.
- Grid 8: Six green dots in the bottom row, connected by a series of arcs and lines.
- Grid 9: One orange dot in the top row.
- Grid 10: Two red dots in the bottom row, connected by a straight line.

So, we have determined a \odot -product of expressions of the forms indicated in (9). Now, apply the labeling indicated as follows, according to our procedure.



So, this shows us how the partition diagram in (10) gets mapped to



according to our sorting map.

Sorting permuting diagrams. Since we claim that our sorting map $\mathcal{S} : \mathcal{P}_n \rightarrow \mathcal{P}_n$ lifts (3), it would be appropriate to formalize and prove this claim, as in Theorem 5 below and our proof of this theorem. We proceed to briefly review some preliminaries concerning Theorem 5.

Our convention for mapping permutations to permutation diagrams, as indicated in (5), is to be used consistently throughout our article. Again, this convention is consistent with the notation for West’s stack-sorting map indicated in (1), since it mimics two-line notation for permutations. We are to extend, as below, this embedding so as to be applicable to expressions as in L and R in the permutation decomposition shown in (1).

The *rook algebra* is a subalgebra of \mathcal{P}_n^ξ that is spanned by partial permutations. Partial permutations are diagrams that consist of blocks of size 1 and blocks of size 2 that consist of a vertex in the top row and a vertex in the bottom row [17]. Following [28], the expression Rd_k denotes the set of all rook k -diagrams. Similarly, for each $r \in \mathbb{Z}$ satisfying $0 \leq r \leq k$, the expression $\text{Rd}_k[r]$ denotes the set of rook k -diagrams with precisely r singleton vertices in each row. Given a permutation decomposition of the form indicated in (1), we can identify L (resp. R) with a partial permutation such that the primed elements in any 2-blocks in this partial permutation are the primed versions of any numbers in L (resp. R) and in such a way so as to agree with the two-line notation indicated in (1). Explicitly, if L is empty, we let it be mapped to the partition diagram in \mathcal{P}_n consisting of singleton blocks, and if we write

$$L = l_1 l_2 \cdots l_{\ell(L)},$$

we may identify L with the partition diagram with 2-blocks of the forms

$$\{p^{-1}(l_1), l'_1\}, \{p^{-1}(l_2), l'_2\}, \dots, \{p^{-1}(l_{\ell(L)}), l'_{\ell(L)}\}$$

and with singleton blocks everywhere else, and similarly for R . This agrees with our convention indicated in (5) for embedding S_n into \mathcal{P}_n .

Again with reference to the permutation decomposition in (1) we may identify the expression n with a partial permutation and in a similar fashion as above, with the understanding that this expression is part of the concatenation in (1). Explicitly,

we would identify n with the partition diagram given by the set-partition

$$(11) \quad \{\{p^{-1}(n), n'\}\} \cup$$

$$(12) \quad \{\{x\} : x \in \mathbb{N}, 1 \leq x \leq n, x \neq p^{-1}(n)\} \cup$$

$$(13) \quad \{\{x\} : x \in \mathbb{N}, 1 \leq x \leq n, x \neq n\}.$$

As illustrated in Example 4, for permutation diagrams π_1 and π_2 , if π_2 consists entirely of singleton nodes then $\mathcal{S}(\pi_1) \odot \mathcal{S}(\pi_2) = \mathcal{S}(\pi_2) \odot \mathcal{S}(\pi_1) = \mathcal{S}(\pi_1)$. This algebraic property is to be used in our proof of Theorem 5.

Theorem 5. *Let p be a permutation in S_n , and let π denote the corresponding partition diagram according to (5). Then $\mathcal{S}(\pi)$ equals the partition diagram corresponding to $s(p)$.*

Proof. As above, we let $p \in S_n$, writing $p: \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$. As above, we let π denote the partition diagram corresponding to (5), i.e., so that the set of blocks of π is (5). With respect to the notation in (8), the expression $\mathcal{S}(\mathcal{L})$ (resp. $\mathcal{S}(\mathcal{R})$) is equal to \mathcal{S} evaluated at a diagram obtained from π by taking any blocks consisting of a bottom node that is labeled the primed version of a number to the left (resp. right) of n in the sense indicated in (1), i.e., a number in L (resp. R) according to the notation in (1) for a permutation $p \in S_n$. Since π is a permutation diagram, any \mathcal{M} -expression consists entirely of singleton nodes. So, the product in (8) reduces to

$$(14) \quad \mathcal{S}(\pi) = \mathcal{S}(\mathcal{L}) \odot \mathcal{S}(\mathcal{R}) \odot B'.$$

As indicated above, \mathcal{L} (resp. \mathcal{R}) is precisely the partition diagram given by embedding L (resp. R) into \mathcal{P}_n . Similarly, B' is precisely the partition diagram given by embedding the factor n in (1), in the manner indicated in (11)–(13). So, letting it be understood that the factors in (1) and the left-hand side of (1) may be identified with their respective embeddings, we find that the \odot -product in (14) may be written as

$$\mathcal{S}(p) = \mathcal{S}(L) \odot \mathcal{S}(R) \odot n.$$

By comparing both sides of the above decomposition with both sides of the decomposition in (2), an inductive argument provides us with the desired result. \square


Since we have lifted West's stack-sorting map so as to allow partition diagrams in addition to permuting diagrams as input, this leads us to consider how fundamental properties concerning the s -map (3) may be "translated" according to our lifting. In particular, we are led to desire to generalize Knuth's classic result that a permutation is 1-stack-sortable if and only if it avoids the pattern 231.

The intricate combinatorial structures and behaviors associated with our lifting \mathcal{S} of West's stack-sorting map are investigated in Section 3 below. Such investigations

are inspired by the extent of mathematical interest concerning the beautiful combinatorics, at both a structural and enumerative level, associated with an important variant of the s -map referred to as *pop-stack sorting* for permutations, with reference to the work of Asinowski et. al. [1].

3. Pattern avoidance

We again let $p \in S_n$ be written as a mapping in the manner indicated in (4). In the context of the study of pattern avoidance in permutations, it is often desirable to denote p using an $n \times n$ grid in the Cartesian plane. For example, if we begin by letting a permutation $p \in S_3$ be denoted in the manner indicated in (1), writing $p = 312$, we may let this be denoted using the $n \times n$ grid below.



More generally, we may denote a permutation

$$p = p(1)p(2) \cdots p(n)$$


with an $n \times n$ array such that an (i, j) -entry is nonempty if and only if (i, j) is of the form $(k, n - p^{-1}(k) + 1)$ for some index k , and where an $(k, n - p^{-1}(k) + 1)$ -entry is highlighted in some way, as above. We say that a permutation $p = p(1)p(2) \cdots p(n)$ contains the pattern 231 if there exist indices k_1, k_2 , and k_3 such that $k_1 < k_2 < k_3$ and such that the entries

$$(15) \quad (k_1, n - p^{-1}(k_1) + 1),$$

$$(16) \quad (k_2, n - p^{-1}(k_2) + 1),$$


$$(17) \quad (k_3, n - p^{-1}(k_3) + 1),$$

satisfy the following in the $n \times n$ array we use to denote p : Point (15) is strictly below (16) and is strictly above point (17). In other words, the $n \times n$ array corresponding to p contains a pattern that is equivalent, in the sense that we have specified, to the following configuration:




A permutation is said to *avoid* the pattern 231 if it does not contain this pattern. In a similar fashion, we may characterize or define a permutation that avoids the

pattern 12 as a *decreasing* permutation, i.e., a permutation of the following form:



Correspondingly, an *increasing* permutation is of the form shown below, i.e., it is an identity permutation:




A permutation p is said to be t -stack-sortable if

$$\underbrace{s \circ s \circ \dots \circ s}_t(p)$$

is increasing; see [9; 13] for related research that has inspired much about this article.

The concept of a 231-avoiding permutation is of considerable interest for the purposes of this article, so it is worthwhile to illustrate a permutation of this form. In this regard, we see that the permutation corresponding to

(18)



avoids the pattern 231. So, according to Knuth's characterization of 1-stack-sortable permutations, we would expect the permutation illustrated in (18) to be 1-stack-sortable. It is worthwhile for our purposes to illustrate this.

Example 6. Being consistent with the notation in (1), the permutation $p \in S_6$ depicted in (18) may be written as 543216. According to the recursion shown in (2) for West's stack-sorting map, we obtain

$$\begin{aligned} s(p) &= s(543216) = s(54321)6 = s(4321)56 \\ &= s(321)456 \\ &= s(21)3456 = s(1)23456 = 123456. \end{aligned}$$

So, since $s(p)$ is the identity permutation in S_6 , we have that p is 1-stack-sortable.

The main goal of our current section is to generalize the following groundbreaking result due to Knuth, which, as indicated in [9], was the starting point for the study of both stack-sorting and pattern avoidance in permutations.

Theorem 7 [20]. *A permutation is 1-stack sortable if and only if it avoids 231 (compare [9]).*


So, in view of our lifting \mathcal{S} of the mapping s , what would be an appropriate way of generalizing Theorem 7 according to the mapping \mathcal{S} ? In particular, how can the concept of “1-stack-sortability” be translated in a meaningful way so as to be applicable with respect to \mathcal{S} ? To answer these questions, we are to make use of a morphism on partition algebras that we had previously applied in a representation-theoretic context [4].

Stretch morphisms. To illustrate the problem of determining a suitable analogue of Theorem 7 according to our lifting \mathcal{S} of West’s stack-sorting map, let us consider \mathcal{S} evaluated at a permuting diagram that is 1-stack-sortable.

Example 8. We see that the permutation 312 avoids the pattern 231. So, let us consider \mathcal{S} evaluated at the permutation diagram corresponding to 312, as below.

$$(19) \quad \mathcal{S} \left(\begin{array}{ccc} \circ & \circ & \circ \\ & \diagdown & / \\ \circ & \circ & \circ \end{array} \right) = \mathcal{S} \left(\begin{array}{ccc} \bullet & \circ & \circ \\ & \diagdown & / \\ \circ & \circ & \bullet \end{array} \right) \left(\begin{array}{ccc} \circ & \bullet & \bullet \\ & \diagdown & / \\ \bullet & \bullet & \circ \end{array} \right) = \mathcal{S} \left(\begin{array}{ccc} \bullet & \circ & \bullet \\ & \diagdown & / \\ \circ & \bullet & \bullet \end{array} \right) \left(\begin{array}{ccc} \bullet & \bullet & \circ \\ & \diagdown & / \\ \bullet & \circ & \bullet \end{array} \right) \left(\begin{array}{ccc} \circ & \bullet & \bullet \\ & \diagdown & / \\ \bullet & \bullet & \circ \end{array} \right)$$

Following through with the procedure introduced in Section 2, we obtain that identity permutation diagram in \mathcal{P}_3 shown below.



From Theorem 5 together with Example 8, given a partition diagram $\pi \in \mathcal{P}_n$, to generalize the notion of 1-stack-sortability in an applicable and meaningful way, it would be appropriate to use a condition whereby $\mathcal{S}(\pi)$ belongs to a fixed class of

generalizations of the multiplicative identity element

$$(20) \quad \begin{array}{c} \circ \quad \circ \quad \circ \\ | \quad | \quad \dots \quad | \\ \circ \quad \circ \quad \circ \\ \underbrace{\hspace{1.5cm}}_n \end{array}$$

in \mathcal{P}_n . This leads us to apply, as below, a morphism on partition algebras introduced in a representation-theoretic context in [4].

Since our lifting of West's stack-sorting map applies to partition diagrams in general, as opposed to, say, rook diagrams, it would be appropriate to allow nonrook diagrams in our lifting of the concept of 1-stack sortability. However, our lifting of $s: S_n \rightarrow S_n$ is such that it preserves the sizes of blocks in a partition diagram (but not necessarily the arrangement or ordering of the blocks). So, it would be appropriate to let a partition diagram be mapped, under \mathcal{S} , to a generalization of (20) allowing for the possibility of blocks other than 2-blocks. In this regard, we are to employ what is referred to as the *Stretch* morphism for partition algebras, as introduced in [4].

Let S be a finite set of natural numbers. Let π be a set-partition of $S \cup S'$. Let k be a natural number such that $k \geq \max(S)$. Following [4], we write $\delta_k(\pi)$ to denote the diagram basis element in \mathcal{P}_k^ξ corresponding to the set-partition given by adding blocks of the form $\{i, i'\}$ to π , where i is a natural number such that $i \notin S$ and $i \leq k$. Again, following [4], we let $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{\ell(\alpha)})$ be a set-composition of a finite set of natural numbers (referring to [4] for details on combinatorial terms related to Stretch morphisms), and we write $m = \ell(\alpha)$, and set $k \geq \max(\bigcup \alpha)$. As in [4], we define

$$(21) \quad \text{Stretch}_{\alpha,k}: \mathcal{P}_m^\xi \rightarrow \mathcal{P}_k^\xi$$

in the following manner.

Let d_π be a member of the diagram basis of \mathcal{P}_m^ξ . Let $\pi = \{\pi_1, \pi_2, \dots, \pi_{\ell(\pi)}\}$. Then

$$(22) \quad \text{Stretch}_{\alpha,k}(d_\pi) = \delta_k \left(\left\{ \bigcup_{\substack{i \in \pi_j \\ i \text{ is unprimed}}} \alpha_i \cup \bigcup_{i' \in \pi_j} \alpha'_i : 1 \leq j \leq \ell(\pi) \right\} \right).$$

We may extend this definition linearly as in [4] so as to obtain an algebra homomorphism, but this is not important for our purposes.

An interesting feature concerning the problem of generalizing 1-stack-sortability by generalizing (3) so as to allow partition diagrams as the arguments of a lifting/extension of the s -map may be explained as follows. By enforcing different conditions on how (20) should be generalized in order to generalize 1-stack-sortability, say, in the context of a given application, this gives rise to different families of

combinatorial objects in the study of the classification of partition diagrams that reduce to a specified generalization of (20), under the application of \mathcal{S} .

Definition 9. We say that a partition diagram π is *stretch-stack-sortable* if $\mathcal{S}(\pi)$ is the evaluation of a Stretch map on an identity permutation diagram.

To begin with, we illustrate the effect of applying (22) to the multiplicative identity elements in a partition algebra.


Example 10. With regard to the notation in (22), we let d_π be the partition diagram corresponding to

$$(23) \quad \pi = \{\{1, 1'\}, \{2, 2'\}, \{3, 3'\}, \{4, 4'\}\}.$$

Define α as the set-composition $(\{1, 2\}, \{3\}, \{5, 6, 7\}, \{4\})$. Observe that the length of α equals the order of d_π , in accordance with the definition in (22). Let us consider the $j = 3$ case within the family of the argument of δ_k indicated in (22), letting the elements in π be ordered in the manner we have written these elements in (23). So, the element corresponding to this $j = 3$ case is

$$\bigcup_{\substack{i \in \pi_3 \\ i \text{ is unprimed}}} \alpha_i \cup \bigcup_{i' \in \pi_3} \alpha'_{i'} = \alpha_3 \cup \alpha'_3 = \{5, 6, 7, 5', 6', 7'\}.$$

Continuing similarly with respect to the other possible values for the j -index, we obtain that $\text{Stretch}_{\alpha,7}$ evaluated at d_π is equal to




Now, let us consider an illustration of a partition diagram satisfying the conditions of Definition 9.

Example 11. According to the procedure in Section 2, we may verify the evaluation

$$\mathcal{S} \left(\begin{array}{c} \text{Diagram with 4 nodes and 4 edges} \end{array} \right) = \begin{array}{c} \text{Diagram with 4 nodes and 3 edges} \end{array}$$

So, the argument of \mathcal{S} , in this case, is stretch-stack-sortable.

Example 12. We also find that



it stretch-stack-sortable.

Examples 11 and 12 illustrate the problem of classifying stretch-stack-sortable partition diagrams. The importance of diagram algebras in both the study of algebraic groups and in the field of knot theory provides a source of further motivation concerning our interest in lifting and generalizing the notions of stack-sortability and pattern avoidance in permutations to basis elements in diagram algebras. The following theorem appears to be the first direct step in this direction. The situation indicated via the bullet points given in Theorem 13 formalizes how the notion of avoiding the pattern 231 may be lifted from permutations to partition diagrams, in a way that is directly applicable to lifting West's stack-sorting map.

Theorem 13. *A partition diagram π is stretch-stack-sortable if and only if:*

- (1) *The only blocks of π are propagating.*
- (2) *For each block of π , it has the same number of upper and lower vertices.*
- (3) *For each block of π , all of its lower nodes are consecutive as primed integers.*
- (4) *Letting the blocks of π be denoted in the form $C_i \cup D'_i$ for indices i , and where C_i and D'_i respectively denote the set of upper and lower nodes of a given block, and writing*

$$(24) \quad D'_1 < D'_2 < \dots,$$

according to the consecutive primed integers labeling D'_i , the situation indicated via the following bullet points cannot occur.

- $D'_{i_1} < D'_{i_2} < D'_{i_3}$.
- *By applying the procedure used to define \mathcal{S} , at the stage in this application when D'_{i_3} contains the greatest nonsingleton primed nodes, one of the following situations occurs:*
 - *either D'_{i_2} is part of an \mathcal{L} -configuration and D'_{i_1} is part of an \mathcal{R} -configuration, or*
 - *D'_{i_2} is part of an \mathcal{L} -configuration and D'_{i_1} is part of an \mathcal{M} -configuration, or*
 - *D'_{i_2} is part of an \mathcal{M} -configuration and D'_{i_1} is part of an \mathcal{R} -configuration, or*
 - *D'_{i_2} is part of an \mathcal{M}_j -configuration and D'_{i_1} is part of an \mathcal{M}_k -configuration, with $j < k$.*

Proof. (\Rightarrow) Let π be a stretch-stack-sortable partition diagram. The mapping \mathcal{S} is such that the number of blocks in a partition diagram μ with u upper nodes and l lower nodes is equal to the number of blocks in $\mathcal{S}(\mu)$ with u upper nodes and l lower nodes. So, π cannot have any nonpropagating blocks, because, otherwise, π would have a nonpropagating block, contradicting that π is stretch-stack-sortable. By using the same property of \mathcal{S} that we have previously indicated, we have that each block of π is such that it has the same number of upper and lower vertices.

Let B be a block of π , and, by way of contradiction, suppose that it is not the case that all of its lower nodes are consecutive as primed integers. According to the procedure used to define \mathcal{S} , in $\mathcal{S}(\pi)$, there would have to be a new block N such that the labels for the lower nodes of N are the same as the labels of the lower nodes of B and such that the upper nodes of N are consecutive integers. So, there would be a block N in $\mathcal{S}(\pi)$ with consecutive integers labeling the upper nodes of N and such that the lower nodes of N do not form a set of consecutive primed integers. However, this is impossible for the stretch of an identity diagram. Now, by way of contradiction, suppose that the situation indicated via the above six bullet points occurs. In any out of the four possibilities corresponding to the last four bullet points, the removal of the block containing D'_{i_3} has the effect of separating the partition diagram containing D'_{i_1} and D'_{i_2} in such a way so that in the \odot -product shown in (8), the block containing D'_{i_1} will be in an argument of \mathcal{S} strictly to the left of the factor in the \odot -product given by \mathcal{S} evaluated at an expression involving D'_{i_2} . Consequently, the ordering of the positions of the bottom nodes of D'_{i_1} and D'_{i_2} will be reversed, but then the top nodes corresponding to D'_{i_2} will be labeled with integers strictly smaller than the top nodes corresponding to D'_{i_1} in the evaluation of $\mathcal{S}(\pi)$, giving us a crossing in the partition diagram $\mathcal{S}(\pi)$ that would be impossible in the stretch of an identity partition diagram.

(\Leftarrow) Conversely, suppose that a partition diagram $\pi \in \mathcal{P}_n$ satisfies all four of the conditions listed above. We apply \mathcal{S} to π , so as to obtain a decomposition of the form indicated in (8). Adopting notation from (8), we may deduce that the following properties hold, again working under the assumption that the four conditions in the theorem under consideration hold:

- (i) The expression B' consists, apart from singleton blocks, of a single propagating block with the same number of upper and lower nodes, and such that the lower nodes are labeled with consecutive primed integers ending with n' .
- (ii) If we were to keep the labelings for the lower nodes as in π' , then the labeling for the lower nonsingleton blocks in the \odot -decomposition in (8) would be in the same order.

This latter property may be verified by a case analysis of the form suggested by the last four out of the six bullet points given above. Repeating this argument inductively, and mimicking notation from (9), we find that $\mathcal{S}(\pi)$ may be written as an \odot -product of the form

$$(25) \quad \mathcal{B}'_1 \odot \mathcal{B}'_2 \odot \cdots \odot \mathcal{B}'_{m_2},$$


where each expression of the form B'_i consists of, apart from singleton blocks, only one propagating block, and this propagating block has the same number of top

nodes as bottom nodes and has consecutive primed integers as its bottom nodes; moreover, the bottom nodes according to the ordering in (25) form the sequence

$$1' < 2' < \dots < n'.$$

So, a direct application of the labeling according to the definition for the \odot -product gives us a stretch of an identity partition diagram. \square

Example 14. If we return to the partition diagram in \mathcal{P}_9 that is illustrated in (7) within Example 3, we find that there is a propagating block containing the largest bottom node $9'$. Being consistent with our notation in (24), we write $D'_1 = \{1', 2', 3'\}$, $D'_2 = \{4', 5', 6'\}$, and $D'_3 = \{7', 8', 9'\}$, and we let the corresponding blocks in (7) be denoted as $C_1 \cup D'_1$, $C_2 \cup D'_2$, and $C_3 \cup D'_3$. We have that $D'_1 < D'_2 < D'_3$, but when we apply the procedure used to define \mathcal{S} to the partition diagram in (7), as we remove the block $C_3 \cup D'_3$, we find that D'_2 is part of an \mathcal{L} -configuration and D'_1 is part of an \mathcal{M} -configuration, which is one of the forbidden patterns indicated in Theorem 13. So, according to Theorem 13, the partition diagram in (7) should not be stretch-stack-sortable, and we may verify that \mathcal{S} evaluated at this same diagram in (7) is equal to



being consistent with the coloring in (7). We see that the partition diagram in (26) is not a stretch of any identity partition diagram.

4. Future work and applications

In regard to Knuth's famous result that the number of 1-stack-sortable permutations in S_n is equal to the n -th Catalan number $\frac{1}{n+1} \binom{2n}{n}$, it would be desirable to obtain a meaningfully similar result for counting the elements in \mathcal{P}_n satisfying the conditions in Theorem 13, i.e., the number of stretch-stack-sortable elements in \mathcal{P}_n . We leave this as an open problem.

An advantage of our lifting of West's stack-sorting map may be described as follows. Since our mapping \mathcal{S} may be evaluated at an arbitrary partition diagram, this allows us to apply and experiment with different ways of generalizing stack-sortability, based on different kinds of partition diagram generalizations of identity permutations. We may obtain many further combinatorial results, relative to our main results, through the use of variants and generalizations of Definition 9. For example, what kinds of bijective and enumerative results can we obtain, relative to the theorems introduced in this article, if we instead consider partition diagrams that get mapped, under the application of \mathcal{S} , to stretches of rook diagrams with

2-blocks of the form $\{i, i'\}$? What if nonrectangular and nonsingleton blocks are allowed?

Many remarkable results and applications have concerned preimages under West's stack-sorting map, as in the work of Defant, Engen, and Miller in [15], for example. What kinds of applications and combinatorial results can we obtain concerning preimages under our lifting of s ?

Since our definition for stretch-stack-sortability is a lifting of 1-stack-sortability, this raises the question as to how the notion of t -stack-sortability may be generalized, according to our procedure from Section 2. The study of t -stack-sortability is widely known to be much more difficult even for the $t = 2$ case, relative to the $t = 1$ case; see [26, Section 1.2.3], for example. In consideration as to Zeilberger's renowned proof [29] of West's conjecture that the number of 2-stack-sortable permutations in S_n is

$$\frac{2(3n)!}{(n+1)!(2n+1)!},$$

this inspires the determination of an analogue of the above indicated enumerative result, using a lifting of 2-stack-sortability via our mapping \mathcal{S} .

Acknowledgements

The author wants to thank Mike Zabrocki for having described to the author (and introduced to the author) the partition algebra morphisms of the form indicated in (22) and how such morphisms may be applied in the representation theory for partition algebras. The author is very thankful for the referee feedback that has been provided.

References

- [1] A. Asinowski, C. Banderier, S. Billey, B. Hackl, and S. Linusson, "Pop-stack sorting and its image: permutations with overlapping runs", *Acta Math. Univ. Comenian. (N.S.)* **88**:3 (2019), 395–402. MR
- [2] G. Benkart and T. Halverson, "Partition algebras and the invariant theory of the symmetric group", pp. 1–41 in *Recent trends in algebraic combinatorics*, Assoc. Women Math. Ser. **16**, Springer, Cham, 2019. MR Zbl
- [3] G. Benkart and T. Halverson, "Partition algebras $P_k(n)$ with $2k > n$ and the fundamental theorems of invariant theory for the symmetric group S_n ", *J. Lond. Math. Soc. (2)* **99**:1 (2019), 194–224. MR Zbl
- [4] J. M. Campbell, "Alternating submodules for partition algebras, rook algebras, and rook-Brauer algebras", preprint, 2022, available at <https://hal.science/hal-03850236>. To appear in *J. Pure Appl. Algebra*.
- [5] G. Cerbai, A. Claesson, and L. Ferrari, "Stack sorting with restricted stacks", *J. Combin. Theory Ser. A* **173** (2020), art. id. 105230. MR Zbl
- [6] C. Defant, "Descents in t -sorted permutations", *J. Comb.* **11**:3 (2020), 511–526. MR Zbl

- [7] C. Defant, “Polyurethane toggles”, *Electron. J. Combin.* **27**:2 (2020), art. id. P2.46. MR Zbl
- [8] C. Defant, “Stack-sorting preimages of permutation classes”, *Sém. Lothar. Combin.* **82** (2020), art. id. B82b. MR Zbl
- [9] C. Defant, “Enumeration of stack-sorting preimages via a decomposition lemma”, *Discrete Math. Theor. Comput. Sci.* **22**:2 (2021), art. id. 3. MR Zbl
- [10] C. Defant, “Fertility monotonicity and average complexity of the stack-sorting map”, *European J. Combin.* **93** (2021), art. id. 103276. MR Zbl
- [11] C. Defant, “Highly sorted permutations and Bell numbers”, *Enumer. Comb. Appl.* **1**:1 (2021), art. id. S2R6. MR Zbl
- [12] C. Defant, “Stack-sorting for Coxeter groups”, *Comb. Theory* **2**:1 (2022), art. id. 18. MR Zbl
- [13] C. Defant and N. Kravitz, “Stack-sorting for words”, *Australas. J. Comb.* **77** (2020), 51–68. Zbl
- [14] C. Defant and K. Zheng, “Stack-sorting with consecutive-pattern-avoiding stacks”, *Adv. Appl. Math.* **128** (2021), art. id. 102192. Zbl
- [15] C. Defant, M. Engen, and J. A. Miller, “Stack-sorting, set partitions, and Lassalle’s sequence”, *J. Combin. Theory Ser. A* **175** (2020), art. id. 105275. MR Zbl
- [16] J. East, “Generators and relations for partition monoids and algebras”, *J. Algebra* **339** (2011), 1–26. MR Zbl
- [17] T. Halverson and T. N. Jacobson, “Set-partition tableaux and representations of diagram algebras”, *Algebr. Comb.* **3**:2 (2020), 509–538. MR Zbl
- [18] T. Halverson and A. Ram, “Partition algebras”, *European J. Combin.* **26**:6 (2005), 869–921. MR Zbl
- [19] V. F. R. Jones, “The Potts model and the symmetric group”, pp. 259–267 in *Subfactors* (Kyuzeso, 1993), World Sci. Publ., River Edge, NJ, 1994. MR Zbl
- [20] D. E. Knuth, *The art of computer programming, I: Fundamental algorithms*, vol. 1, Addison-Wesley, Reading, MA, 1973. MR
- [21] P. Martin, *Potts models and related problems in statistical mechanics*, Series on Advances in Statistical Mechanics **5**, World Scientific Publishing Co., Teaneck, NJ, 1991. MR Zbl
- [22] P. Martin, “Temperley–Lieb algebras for nonplanar statistical mechanics — the partition algebra construction”, *J. Knot Theory Ramifications* **3**:1 (1994), 51–82. MR
- [23] P. Martin, “The structure of the partition algebras”, *J. Algebra* **183**:2 (1996), 319–358. MR Zbl
- [24] P. P. Martin, “The partition algebra and the Potts model transfer matrix spectrum in high dimensions”, *J. Phys. A* **33**:19 (2000), 3669–3695. MR Zbl
- [25] P. Martin and D. Woodcock, “On central idempotents in the partition algebra”, *J. Algebra* **217**:1 (1999), 156–169. MR Zbl
- [26] V. Vatter, *The enumeration and structure of permutation classes*, Ph.D. thesis, Rutgers University, New Brunswick, NJ, 2006, available at <https://sites.math.rutgers.edu/~zeilberg/Theses/VinceVatterThesis.pdf>.
- [27] J. West, *Permutations with forbidden subsequences and stack-sortable permutations*, Ph.D. thesis, Massachusetts Institute of Technology, Cambridge, MA, 1990, available at <https://dspace.mit.edu/handle/1721.1/13641>. MR
- [28] Z. K. Xiao, “On tensor spaces for rook monoid algebras”, *Acta Math. Sin. (Engl. Ser.)* **32**:5 (2016), 607–620. MR Zbl
- [29] D. Zeilberger, “A proof of Julian West’s conjecture that the number of two-stack-sortable permutations of length n is $2(3n)!/((n+1)!(2n+1)!)$ ”, *Discrete Math.* **102**:1 (1992), 85–93. MR Zbl

Received December 11, 2022. Revised April 15, 2023.

JOHN M. CAMPBELL
DEPARTMENT OF MATHEMATICS
TORONTO METROPOLITAN UNIVERSITY
TORONTO, ON
CANADA
jmaxwellcampbell@gmail.com

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2023 is US \$605/year for the electronic version, and \$820/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 324 No. 2 June 2023

Polynomial conditions and homology of FI-modules CIHAN BAHRAN	207
A lift of West's stack-sorting map to partition diagrams JOHN M. CAMPBELL	227
Limit cycles of linear vector fields on $(\mathbb{S}^2)^m \times \mathbb{R}^n$ CLARA CUFÍ-CABRÉ and JAUME LLIBRE	249
Horospherical coordinates of lattice points in hyperbolic spaces: effective counting and equidistribution TAL HORESH and AMOS NEVO	265
Bounded Ricci curvature and positive scalar curvature under Ricci flow KLAUS KRÖNCKE, TOBIAS MARXEN and BORIS VERTMAN	295
Polynomial Dedekind domains with finite residue fields of prime characteristic GIULIO PERUGINELLI	333
The cohomological Brauer group of weighted projective spaces and stacks MINSEON SHIN	353
Pochette surgery of 4-sphere TATSUMASA SUZUKI and MOTOO TANGE	371