

*Pacific
Journal of
Mathematics*

**THE FUNDAMENTAL GROUP OF AN EXTENSION IN
A TANNAKIAN CATEGORY AND THE UNIPOTENT RADICAL
OF THE MUMFORD–TATE GROUP OF AN OPEN CURVE**

PAYMAN ESKANDARI AND V. KUMAR MURTY

THE FUNDAMENTAL GROUP OF AN EXTENSION IN A TANNAKIAN CATEGORY AND THE UNIPOTENT RADICAL OF THE MUMFORD–TATE GROUP OF AN OPEN CURVE

PAYMAN ESKANDARI AND V. KUMAR MURTY

In the first part, we give a self-contained account of Tannakian fundamental groups of extensions, generalizing a result of Hardouin (2008; 2011). In the second part, we use Hardouin’s characterization of Tannakian groups of extensions to give a characterization of the unipotent radical of the Mumford–Tate group of an open complex curve. Consequently, we prove a formula that relates the dimension of the unipotent radical of the Mumford–Tate group of an open complex curve $X \setminus S$ with X smooth and projective and S a finite set of points to the rank of the subgroup of the Jacobian of X supported on S .

1. Introduction

Let X be a smooth complex projective curve and $S \subset X(\mathbb{C})$ a finite nonempty set of points. There is an exact sequence

$$(1) \quad 0 \rightarrow H^1(X) \rightarrow H^1(X \setminus S) \xrightarrow{\text{residue}} \mathbb{Q}(-1)^{|S|-1} \rightarrow 0$$

of (rational) mixed Hodge structures, where the first arrow is induced by the inclusion $X \setminus S \subset X$. In connection to a new proof of the Manin–Drinfeld theorem for modular curves, Deligne proved in the 1970s that this sequence splits (or equivalently, $H^1(X \setminus S)$ is semisimple) if and only if the rank of the subgroup of the Jacobian of X supported on S is zero (see [7, Section 10.3] and [8, Remarque 7.5], and also [11] for another argument).

To any mixed Hodge structure H , one associates an algebraic group called the Mumford–Tate group of H , which we denote by $\mathcal{MT}(H)$. This group can be defined in at least two equivalent ways: In the original definition, due to Mumford (and then refined by Serre) in the pure case, $\mathcal{MT}(H)$ is the subgroup of $\mathrm{GL}(H_{\mathbb{Q}})$ (where as usual, $H_{\mathbb{Q}}$ denotes the underlying rational vector space of H) which fixes all Hodge classes of weight zero in finite direct sums of objects of the form

$$H^{\otimes m} \otimes (H^{\vee})^{\otimes n} \quad (m, n \in \mathbb{Z}_{\geq 0}).$$

MSC2020: primary 11G30, 14C30; secondary 18D30.

Keywords: Tannakian categories, Mumford–Tate groups, mixed Hodge structures.

The second definition, which is somewhat more natural and more conceptual, is in terms of Tannakian formalism: $\mathcal{MT}(H)$ is the fundamental group of the Tannakian subcategory $\langle H \rangle$ of the category of mixed Hodge structures generated by H (see Section 2 for a brief reminder on Tannakian fundamental groups; see [1] for the equivalence of the two definitions). This means that one has a canonical equivalence of categories between $\langle H \rangle$ and the category of finite-dimensional representations of $\mathcal{MT}(H)$.

The unipotent radical of $\mathcal{MT}(H)$ measures how far H is from being semisimple. In particular, H is semisimple if and only if the unipotent radical of $\mathcal{MT}(H)$ is trivial. Thus Deligne’s result about $H^1(X \setminus S)$ can be paraphrased as follows: the unipotent radical of $\mathcal{MT}(H^1(X \setminus S))$ is trivial if and only if the rank of the subgroup of the Jacobian of X supported on S is zero.

The unipotent radical of the Mumford–Tate group of a 1-motive (of which the Mumford–Tate group of $H^1(X \setminus S)$ is an example) has been studied in great generality by Bertolin [3; 4] and Jossen [18]. On his path to prove the main theorem of [18], Jossen gives a characterization of this unipotent radical in Theorem 6.2 of the same article.

In the case of $H^1(X \setminus S)$, Jossen’s characterization is the following: Suppose $S = \{p_0, \dots, p_n\}$. Let P be the identity connected component of the Zariski closure of the subgroup generated by

$$(p_1 - p_0, \dots, p_n - p_0)$$

in $\text{Jac}(X)^n$, where $\text{Jac}(X)$ is the Jacobian of X . Then P itself is an abelian subvariety of $\text{Jac}(X)^n$. Jossen’s theorem asserts that the Lie algebra of the unipotent radical of $\mathcal{MT}(H^1(X \setminus S))$ is canonically isomorphic to $H_1(P)$. In particular, the dimension of the unipotent radical of $\mathcal{MT}(H^1(X \setminus S))$ is twice the dimension of P .

To get a more concrete description (one that does not involve the Zariski closure) of the dimension of the unipotent radical of $\mathcal{MT}(H^1(X \setminus S))$, one can note that linear relations between the points $p_1 - p_0, \dots, p_n - p_0$ with coefficients in the endomorphism algebra of $\text{Jac}(X)$ cut down the dimension of P .

One of the main results of this paper gives a more explicit description of the unipotent radical of $\mathcal{MT}(H^1(X \setminus S))$ that avoids the Zariski closure (see Theorem 4.9.1). As a consequence, in the case where $\text{Jac}(X)$ is simple, we get the following clean formula for the dimension of the unipotent radical (see Theorem 4.9.2(b)):

Theorem A. *Let X , S , and $\text{Jac}(X)$ be as above. Let g , E , and $\mathcal{U}(H^1(X \setminus S))$ be respectively the genus of X , the endomorphism algebra $\text{End}(\text{Jac}(X)) \otimes \mathbb{Q}$ of $\text{Jac}(X)$, and the unipotent radical of the Mumford–Tate group of $H^1(X \setminus S)$. Suppose that $\text{Jac}(X)$ is simple. Then the dimension of $\mathcal{U}(H^1(X \setminus S))$ is equal to $2g$ times the E -rank of the E -submodule of $\text{Jac}(X)(\mathbb{C}) \otimes \mathbb{Q}$ generated by the subgroup supported on S .*

In the general case where $\text{Jac}(X)$ is not necessarily simple, for every division algebra in $\text{End}(\text{Jac}(X)) \otimes \mathbb{Q}$ we get an upper bound for the dimension of $\mathcal{U}(H^1(X \setminus S))$ (see [Theorem 4.9.2\(a\)](#)).

Let us put this discussion on hold for the moment and go to the abstract setting of (neutral) Tannakian categories. Let \mathbf{T} be a Tannakian category over a field K of characteristic zero, and ω a fiber functor over K (the example relevant to the earlier discussion being the category of mixed Hodge structures and the forgetful functor $H \mapsto H_{\mathbb{Q}}$). Suppose we have an extension

$$(2) \quad 0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$$

in \mathbf{T} . Denoting the Tannakian fundamental groups of objects with respect to ω by $\mathcal{G}(-)$, we have a natural surjection

$$\mathcal{G}(M) \rightarrow \mathcal{G}(L \oplus N).$$

Let $\mathcal{U}(M)$ be the kernel of this map (if N and L are semisimple, then $\mathcal{U}(M)$ is the unipotent radical of $\mathcal{G}(M)$). By Tannakian formalism, there is an object

$$\underline{\text{Lie}}(\mathcal{U}(M)) \subset \underline{\text{Hom}}(N, L),$$

whose image under ω is the Lie algebra of $\mathcal{U}(M)$. The question of characterization of $\underline{\text{Lie}}(\mathcal{U}(M))$ has been studied and answered earlier by Hardouin and Bertrand in the case where $N = \mathbb{1}$ and L is semisimple: A theorem of Hardouin [[15](#), Theorem 2] (see also [[14](#)]) asserts that in this case, $\underline{\text{Lie}}(\mathcal{U}(M))$ is the smallest subobject of

$$\underline{\text{Hom}}(\mathbb{1}, L) \cong L$$

such that the pushforward of (2) along the quotient map

$$L \rightarrow L/\underline{\text{Lie}}(\mathcal{U}(M))$$

splits. The result was earlier proved by Bertrand [[5](#), Theorem 1.1] in the setting of D -modules.

The case of arbitrary semisimple N (with L continued to be semisimple as well) can be deduced from Hardouin’s result. In this case, the characterization becomes as follows: If ν is the extension of $\mathbb{1}$ by $\underline{\text{Hom}}(N, L)$ corresponding to (2) under the canonical isomorphism

$$\text{Ext}(N, L) \cong \text{Ext}(\mathbb{1}, \underline{\text{Hom}}(N, L))$$

(where Ext means the Yoneda Ext^1 group in \mathbf{T}), then $\underline{\text{Lie}}(\mathcal{U}(M))$ is the smallest subobject of $\underline{\text{Hom}}(N, L)$ such that the pushforward of ν under the quotient map

$$\underline{\text{Hom}}(N, L) \rightarrow \underline{\text{Hom}}(N, L)/\underline{\text{Lie}}(\mathcal{U}(M))$$

splits.

The goal of this paper is twofold. Our first goal, to which the first part of the paper is devoted, is to give a self-contained and general treatment of Tannakian groups of extensions in characteristic zero. More precisely, in the general setting of the extension (2) in a Tannakian category, in [Theorem 3.3.1](#) we give a characterization of $\underline{\text{Lie}}(\mathcal{U}(M))$ as a subobject of $\underline{\text{Hom}}(N, L)$, without assuming that N or L is semisimple. In the semisimple case, the result simplifies to Hardouin’s characterization (see [Corollary 3.4.1](#)). We also discuss a dual variant of the characterization of $\underline{\text{Lie}}(\mathcal{U}(M))$ ([Theorem 3.5.1](#) and in the semisimple case, [Corollary 3.5.2](#)), which is more convenient in some settings.

We should point out that the generalization to the nonsemisimple situation is indeed useful in practical applications: extensions as in (2) with nonsemisimple L and N arise naturally, for example, in a nonsemisimple Tannakian category with a weight filtration, e.g., the category of mixed motives. In fact, in [\[12\]](#) we build on [Theorem 3.3.1](#) to refine a result of Deligne from [\[18, Appendix\]](#) on unipotent radicals of Tannakian fundamental groups in a Tannakian category with a weight filtration, and then give applications to mixed motives which have “large” unipotent radicals of motivic Galois groups (see the aforementioned paper for more details).

The second goal of the paper, to which the second part of the paper is devoted, is to apply the method of the first part to study the unipotent radical of the Mumford–Tate group of an open curve. Here we take \mathbf{T} to be the category of mixed Hodge structures and apply results about Tannakian groups of extensions to the extension (1). This approach leads to a characterization of the unipotent radical of the Mumford–Tate group of an open curve (see [Theorem 4.9.1](#)). The dimension formula and upper bounds mentioned above follow from this characterization of the unipotent radical (see [Theorem 4.9.2](#)).

The proof of [Theorem 4.9.1](#) has two ingredients: The first ingredient is the semisimple case of [Theorem 3.3.1](#) due to Hardouin (or more precisely, its dual variant given in [Corollary 3.5.2](#)). This gives a characterization of $\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))$ as follows: if μ is the element of

$$\text{Ext}(H^1(X)^{|S|-1}, \mathbb{1})$$

corresponding to (1) under the canonical isomorphisms

$$\begin{aligned} \text{Ext}(\mathbb{Q}(-1)^{|S|-1}, H^1(X)) \\ \cong \text{Ext}(H_1(X) \otimes \mathbb{Q}(-1)^{|S|-1}, \mathbb{1}) \end{aligned} \quad \begin{array}{c} \text{Poincaré duality} \\ \cong \end{array} \quad \text{Ext}(H^1(X)^{|S|-1}, \mathbb{1}),$$

then the orthogonal complement (see [Section 3.5](#)) of $\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))$ is the largest subobject of $H^1(X)^{|S|-1}$ on which μ restricts to a split extension. The second ingredient of the argument is now the calculation of the restrictions of the extension μ along different maps $H^1(X) \rightarrow H^1(X)^{|S|-1}$.

[Theorem 4.9.1](#) can be deduced alternatively from Jossen’s general characterization of the unipotent radical of the Mumford–Tate group of an arbitrary 1-motive given in [[18](#), [Theorem 6.2](#)]. Although [Theorem 4.9.1](#) is weaker than Jossen’s [[18](#), [Theorem 6.2](#)], we hope that the reader might find some value in the simplicity of our approach and exposition, which solely rely on the general material on Tannakian groups and the calculation of the relevant extensions in the category of mixed Hodge structures. This approach can be applied to any situation where the relevant extensions can be calculated and described nicely. It is also hopefully more accessible to some audiences.

The paper is organized as follows. In the next section, we recall some basic generalities about Tannakian categories. In [Section 3](#) we prove the characterizations of $\underline{\text{Lie}}(\mathcal{U}(M))$ in a general Tannakian category and for general L and N (with notation as above). A reader not familiar with the language of Tannakian categories but familiar with properties of the category of mixed Hodge structures may assume in [Sections 2](#) and [3](#) that \mathbf{T} is the latter category and ω is the forgetful functor. In [Section 4](#), we come back to the problem of studying the unipotent radical of the Mumford–Tate group of an open curve, and prove [Theorems 4.9.1](#) and [4.9.2](#).

2. Preliminaries

In this section we briefly recall a few facts and constructions about Tannakian categories. For any commutative ring R , let \mathbf{Mod}_R denote the category of R -modules. Throughout, K is a field of characteristic zero. The categories of groups and commutative K -algebras are respectively denoted by **Groups** and \mathbf{Alg}_K . For an affine group scheme \mathcal{G} over K , let $\mathbf{Rep}(\mathcal{G})$ be the category of finite-dimensional representations of \mathcal{G} over K . We use the language of [[10](#)] for the theory of Tannakian categories. Our Tannakian categories are all neutral.

2.1. Let \mathbf{T} be a Tannakian category over K with unit object $\mathbb{1}$; thus \mathbf{T} is a K -linear rigid abelian tensor category with the identity $\mathbb{1}$ of the tensor structure satisfying $\text{End}(\mathbb{1}) = K$, for which there exists a fiber functor, i.e., a K -linear exact faithful tensor functor

$$\mathbf{T} \rightarrow \mathbf{Mod}_K.$$

Let ω be such a functor. Let

$$\underline{\text{Aut}}^{\otimes}(\omega) : \mathbf{Alg}_K \rightarrow \mathbf{Groups}$$

be the functor that sends a commutative K -algebra R to

$$\begin{aligned} \text{Aut}^{\otimes}(\omega \otimes 1_R) &:= \text{the group of automorphisms of the functor} \\ \omega \otimes 1_R : \mathbf{T} &\rightarrow \mathbf{Mod}_R \\ &\text{respecting the tensor structures.} \end{aligned}$$

The fundamental theorem of the theory of Tannakian categories [10, Theorem 2.11] asserts that $\underline{\text{Aut}}^{\otimes}(\omega)$ is representable by an affine group scheme $\mathcal{G}(\mathbf{T}, \omega)$ over K (so that $\underline{\text{Aut}}^{\otimes}(\omega)$ is the functor of points of $\mathcal{G}(\mathbf{T}, \omega)$), and that the functor

$$\mathbf{T} \rightarrow \mathbf{Rep}(\mathcal{G}(\mathbf{T}, \omega))$$

sending

$$M \mapsto \omega M$$

(with the natural action of $\mathcal{G}(\mathbf{T}, \omega)$ on ωM) is an equivalence of tensor categories. We call $\mathcal{G}(\mathbf{T}, \omega)$ the fundamental (or the Tannakian) group of \mathbf{T} with respect to ω .

If \mathbf{T}' is also a Tannakian category over K , a tensor functor $\phi : \mathbf{T}' \rightarrow \mathbf{T}$ gives rise to a morphism

$$\phi^{\#} : \mathcal{G}(\mathbf{T}, \omega) \rightarrow \mathcal{G}(\mathbf{T}', \omega \circ \phi)$$

of group schemes over K , sending an automorphism of $\omega \otimes 1_R$ for any K -algebra R to the obvious automorphism induced on $(\omega \otimes 1_R) \circ \phi = (\omega \circ \phi) \otimes 1_R$. The morphism $\phi^{\#}$ is surjective (or faithfully flat) if and only if ϕ is fully faithful and moreover, satisfies the following property: for every $M \in \mathbf{T}'$, every subobject of $\phi(M)$ is isomorphic to $\phi(L)$ for some subobject L of M (see [10, Proposition 2.21], for instance). In particular, if \mathbf{T}' is a full Tannakian subcategory of \mathbf{T} which is closed under taking subobjects, then the inclusion $\mathbf{T}' \subset \mathbf{T}$ gives rise to a surjective morphism $\mathcal{G}(\mathbf{T}, \omega) \rightarrow \mathcal{G}(\mathbf{T}', \omega|_{\mathbf{T}'})$, where $\omega|_{\mathbf{T}'}$ is the restriction of ω to \mathbf{T}' .

2.2. Let M be an object of \mathbf{T} . Let $\langle M \rangle$ denote the full Tannakian subcategory of \mathbf{T} generated by M , that is, the smallest full Tannakian subcategory of \mathbf{T} that contains M , and is closed under taking subobjects (or subquotients). Set

$$\mathcal{G}(M, \omega) := \mathcal{G}(\langle M \rangle, \omega|_{\langle M \rangle}) = \underline{\text{Aut}}^{\otimes}(\omega|_{\langle M \rangle});$$

we refer to this group as the fundamental (or the Tannakian) group of M with respect to ω . Starting with M and $\mathbb{1}$, we can obtain every object of $\langle M \rangle$ by finitely many iterations of taking direct sums, duals, tensor products, and subquotients. It follows that the natural map

$$\mathcal{G}(M, \omega) \rightarrow \text{GL}_{\omega M}, \quad \sigma \mapsto \sigma_M$$

(restricting to the action on ωM) is injective, so that, indeed, $\mathcal{G}(M, \omega)$ is an algebraic group over K . (Here, complying with the standard notation for natural transformations, $\sigma_M : \omega M \rightarrow \omega M$ is how σ acts on ωM .) Often we will identify $\mathcal{G}(M, \omega)$ as a subgroup of $\text{GL}_{\omega M}$ via the injection above.

Since $\langle M \rangle$ is closed under taking subobjects, the natural map $\mathcal{G}(\mathbf{T}, \omega) \rightarrow \mathcal{G}(M, \omega)$ (induced by the inclusion $\langle M \rangle \subset \mathbf{T}$) is surjective. The kernel of this map consists of all $\sigma \in \mathcal{G}(\mathbf{T}, \omega)$ such that σ_M is identity (then by functoriality, σ_N is also identity for every $N \in \langle M \rangle$).

2.3. For any algebraic group G , let $\text{Lie}(G)$ be the Lie algebra of G . Let \mathcal{N} be a normal subgroup of $\mathcal{G}(M, \omega)$. Consider the adjoint representation

$$(3) \quad \text{Ad} : \mathcal{G}(M, \omega) \rightarrow \text{GL}_{\text{Lie}(\mathcal{N})}.$$

In view of the equivalence of categories

$$(4) \quad \langle M \rangle \rightarrow \mathbf{Rep}(\mathcal{G}(M, \omega)), \quad A \mapsto \omega A,$$

there is a canonical object $\underline{\text{Lie}}(\mathcal{N})$ in $\langle M \rangle$ with

$$\omega \underline{\text{Lie}}(\mathcal{N}) = \text{Lie}(\mathcal{N}),$$

such that the natural action of $\mathcal{G}(M, \omega)$ on $\omega \underline{\text{Lie}}(\mathcal{N})$ (through the definition of $\mathcal{G}(M, \omega)$ as the group of tensor automorphisms of the functor ω) coincides with the adjoint representation (3).

3. The fundamental group of an extension

The goal of this section is to study the fundamental group of an extension in a Tannakian category. As before, let \mathbf{T} be a Tannakian category over a field K of characteristic zero. Fix a fiber functor $\omega : \mathbf{T} \rightarrow \mathbf{Mod}_K$. We shall drop ω from the notation for fundamental groups, and simply write $\mathcal{G}(M)$ (for M an object of \mathbf{T}). We use the notation I_A for the identity map on an object A of a given category. We use an unadorned Hom to denote a Hom group in a category of modules, with the coefficient ring understood from the context. In \mathbf{T} or any category of modules, the dual of an object A is denoted by A^\vee .

3.1. Let L, M and N be objects of \mathbf{T} given in an exact sequence

$$(5) \quad 0 \rightarrow L \xrightarrow{i} M \xrightarrow{q} N \rightarrow 0,$$

where (as indicated in the diagram) the morphisms $L \rightarrow M$ and $M \rightarrow N$ are respectively denoted by i and q .

The inclusion $\iota : \langle L \oplus N \rangle \subset \langle M \rangle$ induces a surjective morphism

$$\iota^\# : \mathcal{G}(M) \rightarrow \mathcal{G}(L \oplus N).$$

Let $\mathcal{U}(M)$ be the kernel of this map; it consists of those $\sigma \in \mathcal{G}(M)$ which act trivially on $\omega L \oplus \omega N$, or equivalently, on both ωL and ωN (i.e., $\sigma_L = I_{\omega L}$ and $\sigma_N = I_{\omega N}$). Note that while for simplicity we did not incorporate L and N in the notation for $\mathcal{U}(M)$, in general, $\mathcal{U}(M)$ will also depend on L and N . Our goal in this section is to study the group $\mathcal{U}(M)$.

First, let us describe the map $\iota^\#$ more concretely. Use the map i (see (5)) to identify ωL as a subspace of ωM . Moreover, once and for all, choose a section of

the surjection $\omega q : \omega M \rightarrow \omega N$ to identify

$$\omega M = \omega L \oplus \omega N$$

(as vector spaces). Then the functor ω applied to the sequence (5) gives

$$(6) \quad 0 \rightarrow \omega L \rightarrow \omega L \oplus \omega N \rightarrow \omega N \rightarrow 0,$$

where the second and third arrows are the inclusion and projection maps.

Let σ be an element of $\mathcal{G}(M)$. Since σ is an automorphism of the functor ω , we have a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \omega L & \longrightarrow & \omega L \oplus \omega N & \longrightarrow & \omega N & \longrightarrow & 0 \\ & & \downarrow \sigma_L & & \downarrow \sigma_M & & \downarrow \sigma_N & & \\ 0 & \longrightarrow & \omega L & \longrightarrow & \omega L \oplus \omega N & \longrightarrow & \omega N & \longrightarrow & 0 \end{array}$$

It follows that

$$\sigma_M = \begin{pmatrix} \sigma_L & f \\ 0 & \sigma_N \end{pmatrix} \in \text{GL}_{\omega L \oplus \omega N}$$

for some $f \in \text{Hom}(\omega N, \omega L)$. Let

$$G(M) \subset \text{GL}_{\omega L \oplus \omega N}$$

be the subgroup consisting of the elements which stabilize ωL . Regarding $\mathcal{G}(M)$ as a subgroup of $\text{GL}_{\omega M} = \text{GL}_{\omega L \oplus \omega N}$ (via $\sigma \mapsto \sigma_M$), we have

$$\mathcal{G}(M) \subset G(M).$$

Similarly, for any σ in $\mathcal{G}(L \oplus N)$,

$$\sigma_{L \oplus N} = \begin{pmatrix} \sigma_L & 0 \\ 0 & \sigma_N \end{pmatrix} \in \text{GL}_{\omega L \oplus \omega N}.$$

Thinking of $\mathcal{G}(L \oplus N)$ (resp. $\text{GL}_{\omega L} \times \text{GL}_{\omega N}$) as a subgroup of $\text{GL}_{\omega L \oplus \omega N}$ via $\sigma \mapsto \sigma_{L \oplus N}$ (resp. the diagonal embedding), we have

$$\mathcal{G}(L \oplus N) \subset \text{GL}_{\omega L} \times \text{GL}_{\omega N}.$$

The map $\iota^\#$ is then the restriction of

$$\begin{aligned} \varphi : G(M) &\rightarrow \text{GL}_{\omega L} \times \text{GL}_{\omega N} \\ \begin{pmatrix} g & * \\ 0 & g' \end{pmatrix} &\mapsto \begin{pmatrix} g & 0 \\ 0 & g' \end{pmatrix} \quad (g \in \text{GL}_{\omega L}, g' \in \text{GL}_{\omega N}). \end{aligned}$$

Let

$$U(M) := \ker(\varphi).$$

Thus $U(M)$ is the subgroup of $\mathrm{GL}_{\omega L \oplus \omega N}$ consisting of the elements of the form

$$\begin{pmatrix} I_{\omega L} & * \\ 0 & I_{\omega N} \end{pmatrix},$$

and in particular, is an abelian unipotent group. We have a commutative diagram

$$(7) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \mathcal{U}(M) & \longrightarrow & \mathcal{G}(M) & \xrightarrow{i^\#} & \mathcal{G}(L \oplus N) & \longrightarrow & 1 \\ & & & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & U(M) & \longrightarrow & G(M) & \xrightarrow{\varphi} & \mathrm{GL}_{\omega L} \times \mathrm{GL}_{\omega N} & \longrightarrow & 1 \end{array}$$

where the injective arrows are inclusion maps and the rows are exact. Thus

$$U(M) \subset \mathcal{U}(M).$$

Being a subgroup of an abelian unipotent group, $\mathcal{U}(M)$ is abelian and unipotent.

As discussed in Section 2.3, the adjoint representation of $\mathcal{G}(M)$ gives a canonical object $\underline{\mathrm{Lie}}(\mathcal{U}(M))$ of $\langle M \rangle$ whose image under ω is $\mathrm{Lie}(\mathcal{U}(M))$. Since $\mathcal{U}(M)$ is abelian, the action of $\mathcal{G}(M)$ on $\mathrm{Lie}(\mathcal{U}(M))$ factors through an action of $\mathcal{G}(L \oplus N)$, so that indeed, the object $\underline{\mathrm{Lie}}(\mathcal{U}(M))$ belongs to the subcategory $\langle L \oplus N \rangle$.

The Lie algebra of $U(M)$ can be identified with

$$\mathrm{Hom}(\omega N, \omega L)$$

(with trivial Lie bracket). The exponential map

$$\exp : \mathrm{Lie}(U(M)) = \mathrm{Hom}(\omega N, \omega L) \rightarrow U(M)(K)$$

(with its inverse denoted by \log) is given by

$$(8) \quad \exp(f) = \begin{pmatrix} I_{\omega L} & f \\ 0 & I_{\omega N} \end{pmatrix}.$$

Let $\underline{\mathrm{Hom}}(N, L)$ denote the internal hom object in the category \mathbf{T} . We identify

$$\omega(\underline{\mathrm{Hom}}(N, L)) = \mathrm{Hom}(\omega N, \omega L)$$

via the canonical isomorphism between the two.

The following observation is standard.

Proposition 3.1.1. *The inclusion map*

$$\mathrm{Lie}(\mathcal{U}(M)) \rightarrow \mathrm{Hom}(\omega N, \omega L)$$

is ω of a morphism

$$\underline{\mathrm{Lie}}(\mathcal{U}(M)) \rightarrow \underline{\mathrm{Hom}}(N, L).$$

(In other words, $\underline{\mathrm{Lie}}(\mathcal{U}(M))$ can be identified as a subobject of $\underline{\mathrm{Hom}}(N, L)$.)

Proof. In view of the equivalence of categories (4), it is enough to show that the natural actions of $\mathcal{G}(M)$ on $\text{Lie}(\mathcal{U}(M))$ and $\text{Hom}(\omega N, \omega L)$ are compatible. In other words, we need to show that for any commutative K -algebra R and $\sigma \in \mathcal{G}(M)(R)$, we have

$$\sigma_{\underline{\text{Lie}}(\mathcal{U}(M))} = \sigma_{\underline{\text{Hom}}(N, L)}|_{\text{Lie}(\mathcal{U}(M))_R},$$

where for any vector space V over K , we denote $V_R := V \otimes R$. We may identify

$$(\omega \underline{\text{Hom}}(N, L))_R = \text{Hom}((\omega N)_R, (\omega L)_R)$$

(Hom in R -modules). Considering the evaluation map $N \otimes N^\vee \rightarrow \mathbb{1}$ and the canonical isomorphism $\underline{\text{Hom}}(N, L) \cong N^\vee \otimes L$ (which after applying ω , are the corresponding maps in linear algebra), one easily sees that the map $\sigma_{\underline{\text{Hom}}(N, L)}$ is given by

$$f \mapsto \sigma_L \circ f \circ \sigma_N^{-1} \quad (f \in \text{Hom}((\omega N)_R, (\omega L)_R)).$$

We now calculate the map $\sigma_{\underline{\text{Lie}}(\mathcal{U}(M))}$. By definition, the action of $\mathcal{G}(M)$ on $\text{Lie}(\mathcal{U}(M))$ is the restriction of the adjoint representation of $\mathcal{G}(M)$ to $\text{Lie}(\mathcal{U}(M))$. Let

$$f \in \text{Lie}(\mathcal{U}(M))_R \subset \text{Lie}(U(M))_R = \text{Hom}((\omega N)_R, (\omega L)_R).$$

Then $\sigma_{\underline{\text{Lie}}(\mathcal{U}(M))}(f)$ is characterized by

$$\exp(\sigma_{\underline{\text{Lie}}(\mathcal{U}(M))}(f)) = \sigma_M \exp(f) \sigma_M^{-1},$$

where \exp is the isomorphism between $\text{Lie}(\mathcal{U}(M))$ and $\mathcal{U}(M)$ as varieties over K , and via the inclusion $\mathcal{U}(M) \subset U(M)$, is given by (8) (with coefficients extended to R). Writing

$$\sigma_M = \begin{pmatrix} \sigma_L & h \\ 0 & \sigma_N \end{pmatrix},$$

where $h \in \text{Hom}((\omega N)_R, (\omega L)_R)$, we have

$$\begin{aligned} \sigma_M \exp(f) \sigma_M^{-1} &= \begin{pmatrix} \sigma_L & h \\ 0 & \sigma_N \end{pmatrix} \begin{pmatrix} I_{(\omega L)_R} & f \\ 0 & I_{(\omega N)_R} \end{pmatrix} \begin{pmatrix} \sigma_L^{-1} & -\sigma_L^{-1} \circ h \circ \sigma_N^{-1} \\ 0 & \sigma_N^{-1} \end{pmatrix} \\ &= \begin{pmatrix} I_{(\omega L)_R} & \sigma_L \circ f \circ \sigma_N^{-1} \\ 0 & I_{(\omega N)_R} \end{pmatrix} = \exp(\sigma_L \circ f \circ \sigma_N^{-1}). \end{aligned}$$

Thus

$$\sigma_{\underline{\text{Lie}}(\mathcal{U}(M))}(f) = \sigma_L \circ f \circ \sigma_N^{-1},$$

as desired. \square

Remark 3.1.2. (1) The embedding

$$\text{Lie}(\mathcal{U}(M)) \subset \text{Hom}(\omega N, \omega L)$$

is independent of the section of ωq used to identify $\omega M = \omega L \oplus \omega N$. Indeed, if we had chosen a different section of ωq and hence a different identification of ωM

as $\omega L \oplus \omega N$, then the resulting embedding $\mathcal{G}(M) \hookrightarrow \mathrm{GL}_{\omega L \oplus \omega N}$ would differ from the previous one by conjugation by an element of $U(M)$. Since $U(M)$ is abelian, the two embeddings agree on $\mathcal{U}(M)$. Thus our identification of $\underline{\mathrm{Lie}}(\mathcal{U}(M))$ as a subobject of $\underline{\mathrm{Hom}}(N, L)$ is independent of the choice of the section of ωq .

(2) If L and N are semisimple, then $\mathcal{U}(M)$ is the unipotent radical of $\mathcal{G}(M)$, and in particular will only depend on M (and not on the choices of L or N). (Recall that L and N are semisimple if and only if the category $\langle L \oplus N \rangle$ is semisimple if and only if $\mathcal{G}(L \oplus N)$ is reductive.)

3.2. Before we proceed any further, let us recall a categorical construction. The extension (5) gives an element of

$$\mathrm{Ext}(N, L),$$

where Ext denotes the Yoneda Ext^1 group in \mathbf{T} . Recall that one has a canonical isomorphism

$$(9) \quad \mathrm{Ext}(N, L) \cong \mathrm{Ext}(\mathbb{1}, \underline{\mathrm{Hom}}(N, L)).$$

Let

$$v \in \mathrm{Ext}(\mathbb{1}, \underline{\mathrm{Hom}}(N, L))$$

be the extension class corresponding to (5) under the canonical isomorphism (9). Then v is the class of the extension obtained by first applying $\underline{\mathrm{Hom}}(N, -)$ to the sequence (5):

$$0 \rightarrow \underline{\mathrm{Hom}}(N, L) \rightarrow \underline{\mathrm{Hom}}(N, M) \rightarrow \underline{\mathrm{Hom}}(N, N) \rightarrow 0,$$

and then pulling back along the canonical morphism

$$e : \mathbb{1} \rightarrow \underline{\mathrm{Hom}}(N, N)$$

characterized by the fact that

$$\omega e(1) \in \omega \underline{\mathrm{Hom}}(N, N) = \mathrm{Hom}(\omega N, \omega N)$$

is the identity map. Going through this procedure, assuming $N \neq 0$, we see that v is the class of the extension

$$(10) \quad 0 \rightarrow \underline{\mathrm{Hom}}(N, L) \rightarrow \underline{\mathrm{Hom}}(N, M)^\dagger \rightarrow \mathbb{1} \rightarrow 0,$$

where

- $\underline{\mathrm{Hom}}(N, M)^\dagger$ is the subobject of $\underline{\mathrm{Hom}}(N, M)$ characterized by

$$\begin{aligned} & \omega \underline{\mathrm{Hom}}(N, M)^\dagger \\ &= \mathrm{Hom}(\omega N, \omega M)^\dagger \\ &:= \{ f \in \mathrm{Hom}(\omega N, \omega M) : (\omega q) \circ f = \lambda(f) \mathrm{Id}_{\omega N} \text{ for some } \lambda(f) \in K \}, \end{aligned}$$

- after applying ω , the injective arrow is $f \mapsto (\omega i) \circ f$, and

- after applying ω , the surjective arrow is the map $f \mapsto \lambda(f)$, where $\lambda(f) \in K$ is as in the definition of $\underline{\text{Hom}}(N, M)^\dagger$ above.

If N (and hence $\underline{\text{Hom}}(N, L)$) is zero, then ν is the trivial extension

$$0 \rightarrow 0 \rightarrow \mathbb{1} \rightarrow \mathbb{1} \rightarrow 0.$$

For convenience, we set $\underline{\text{Hom}}(N, M)^\dagger := \mathbb{1}$ in this case.

3.3. We are ready to give the characterization of the subobject $\underline{\text{Lie}}(\mathcal{U}(M))$ of $\underline{\text{Hom}}(N, L)$. To simplify the notation, we identify $\underline{\text{Hom}}(N, L)$ with its image under the injection $\underline{\text{Hom}}(N, L) \rightarrow \underline{\text{Hom}}(N, M)^\dagger$.

Theorem 3.3.1. *Let A be a subobject of $\underline{\text{Hom}}(N, L)$. Then A contains $\underline{\text{Lie}}(\mathcal{U}(M))$ if and only if the quotient*

$$\underline{\text{Hom}}(N, M)^\dagger / A$$

belongs to the subcategory $\langle L \oplus N \rangle$. (Thus $\underline{\text{Lie}}(\mathcal{U}(M))$ is the smallest subobject of $\underline{\text{Hom}}(N, L)$ with this property.)

Proof. The theorem is trivial if $N = 0$, so we may assume $N \neq 0$. An object X of $\langle M \rangle$ belongs to the subcategory $\langle L \oplus N \rangle$ if and only if the subgroup $\mathcal{U}(M)$ of $\mathcal{G}(M)$ acts trivially on ωX . Thus the assertion in the theorem can be paraphrased as that A contains $\underline{\text{Lie}}(\mathcal{U}(M))$ if and only if the action of $\mathcal{U}(M)$ on $\omega(\underline{\text{Hom}}(N, M)^\dagger / A)$ is trivial.

Let $\sigma \in \mathcal{G}(M)(K)$. Let $A \subset \underline{\text{Hom}}(N, L)$. The morphism

$$\underline{\text{Hom}}(N, M)^\dagger \rightarrow \underline{\text{Hom}}(N, M)^\dagger / A$$

gives rise to a commutative diagram

$$\begin{array}{ccc} \text{Hom}(\omega N, \omega M)^\dagger & \longrightarrow & \text{Hom}(\omega N, \omega M)^\dagger / \omega A \\ \downarrow \sigma_{\underline{\text{Hom}}(N, M)^\dagger} & & \downarrow \sigma_{\underline{\text{Hom}}(N, M)^\dagger / A} \\ \text{Hom}(\omega N, \omega M)^\dagger & \longrightarrow & \text{Hom}(\omega N, \omega M)^\dagger / \omega A \end{array}$$

Thus

$$\sigma_{\underline{\text{Hom}}(N, M)^\dagger / A}(f + \omega A) = \sigma_{\underline{\text{Hom}}(N, M)^\dagger}(f) + \omega A$$

for every $f \in \text{Hom}(\omega N, \omega M)^\dagger$.

As before, we use our fixed section of $\omega q : \omega M \rightarrow \omega N$ to identify $\omega M = \omega L \oplus \omega N$. Then we have

$$(11) \quad \begin{array}{ccc} \text{Hom}(\omega N, \omega M) & \xlongequal{\quad} & \text{Hom}(\omega N, \omega L) \oplus \text{Hom}(\omega N, \omega N) \\ \uparrow & & \uparrow \\ \text{Hom}(\omega N, \omega M)^\dagger & \xlongequal{\quad} & \text{Hom}(\omega N, \omega L) \oplus K \cdot I_{\omega N} \end{array}$$

Suppose $\sigma \in \mathcal{U}(M)(K)$. Then σ_L and σ_N are both identity, and the action of σ on the $\mathcal{G}(M)$ -invariant subspace $\underline{\mathrm{Hom}}(\omega N, \omega L)$ of $\underline{\mathrm{Hom}}(\omega N, \omega M)^\dagger$ is trivial. Thus

$$\sigma_{\underline{\mathrm{Hom}}(N, M)^\dagger/A} = I_{\underline{\mathrm{Hom}}(\omega N, \omega M)^\dagger/\omega A}$$

if and only if

$$\sigma_{\underline{\mathrm{Hom}}(N, M)^\dagger/A}(I_{\omega N} + \omega A) = I_{\omega N} + \omega A,$$

where here, as well as in the rest of this argument except in (12) below, $I_{\omega N}$ is considered as an element of $\underline{\mathrm{Hom}}(\omega N, \omega M)^\dagger$ via the decomposition (11). This is equivalent to

$$\sigma_{\underline{\mathrm{Hom}}(N, M)^\dagger}(I_{\omega N}) - I_{\omega N} \in \omega A.$$

Note that $\sigma_{\underline{\mathrm{Hom}}(N, M)}$ (and hence $\sigma_{\underline{\mathrm{Hom}}(N, M)^\dagger}$) is given by

$$f \mapsto \sigma_M \circ f \circ \sigma_N^{-1} = \sigma_M \circ f \quad (f \in \underline{\mathrm{Hom}}(\omega N, \omega M)).$$

We have

$$(12) \quad \sigma_M = \begin{pmatrix} I_{\omega L} & \log(\sigma_M) \\ 0 & I_{\omega N} \end{pmatrix} \in \mathrm{GL}_{\omega L \oplus \omega N}(K),$$

where $\log(\sigma_M) \in \underline{\mathrm{Hom}}(\omega N, \omega L)$. Then

$$\sigma_{\underline{\mathrm{Hom}}(N, M)^\dagger}(I_{\omega N}) = \sigma_M \circ I_{\omega N} = \log(\sigma_M) + I_{\omega N},$$

so that

$$\sigma_{\underline{\mathrm{Hom}}(N, M)^\dagger}(I_{\omega N}) - I_{\omega N} = \log(\sigma_M).$$

We have shown that any element $\sigma \in \mathcal{U}(M)(K)$ acts trivially on $\omega(\underline{\mathrm{Hom}}(N, M)^\dagger/A)$ if and only if $\log(\sigma_M)$ is in ωA . The group $\mathcal{U}(M)$ is unipotent and hence $\mathcal{U}(M)(K)$ is dense in $\mathcal{U}(M)$. It follows that $\mathcal{U}(M)$ acts trivially on $\omega(\underline{\mathrm{Hom}}(N, M)^\dagger/A)$ if and only if for every $\sigma \in \mathcal{U}(M)(K)$, we have $\log(\sigma_M) \in \omega A$, i.e., if and only if $\mathrm{Lie}(\mathcal{U}(M)) \subset \omega A$. This completes the proof. \square

3.4. For every subobject A of $\underline{\mathrm{Hom}}(N, L)$, pushing extensions forward along the natural map $\underline{\mathrm{Hom}}(N, L) \rightarrow \underline{\mathrm{Hom}}(N, L)/A$ we have a map

$$\mathrm{Ext}(\mathbb{1}, \underline{\mathrm{Hom}}(N, L)) \rightarrow \mathrm{Ext}(\mathbb{1}, \underline{\mathrm{Hom}}(N, L)/A).$$

We denote the image of ν under this map by ν/A . [Theorem 3.3.1](#) has the following corollary:

Corollary 3.4.1. (a) *If A is a subobject of $\underline{\mathrm{Hom}}(N, L)$ such that ν/A is trivial, then $\underline{\mathrm{Lie}}(\mathcal{U}(M)) \subset A$.*

(b) *Suppose L and N are semisimple. Then $\nu/\underline{\mathrm{Lie}}(\mathcal{U}(M))$ is trivial (and hence $\underline{\mathrm{Lie}}(\mathcal{U}(M))$ is the smallest subobject of $\underline{\mathrm{Hom}}(N, L)$ with this property).*

Proof. We may assume $N \neq 0$. Suppose ν/A is trivial. Then

$$\underline{\text{Hom}}(N, M)^\dagger/A \simeq \underline{\text{Hom}}(N, L)/A \oplus \mathbb{1},$$

and hence $\underline{\text{Hom}}(N, M)^\dagger/A$ belongs to the subcategory generated by L and N . Thus (a) follows from [Theorem 3.3.1](#).

As for (b), the theorem implies that $\underline{\text{Hom}}(N, M)^\dagger/\underline{\text{Lie}}(\mathcal{U}(M))$ is in $\langle L \oplus N \rangle$, which is a semisimple category by the hypothesis of semisimplicity of L and N . Thus $\nu/\underline{\text{Lie}}(\mathcal{U}(M))$ splits. \square

Remark 3.4.2. The semisimple case of [Corollary 3.4.1](#) is originally due to Hardouin (see Theorem 2 of [\[15\]](#) as well as Théorème 2.1 of [\[14\]](#)). Hardouin shows that when L and N are semisimple, $\underline{\text{Lie}}(\mathcal{U}(M))$ is the smallest subobject of $\underline{\text{Hom}}(N, L)$ such that $\nu/\underline{\text{Lie}}(\mathcal{U}(M))$ is trivial. The same statement was earlier proved by Bertrand [\[5, Theorem 1.1\]](#) in the special case where \mathbf{T} is the category of D -modules over a differential field of characteristic zero. (Both Hardouin and Bertrand take $N = \mathbb{1}$, but one can deduce the case of arbitrary (semisimple) N from that.)

3.5. Consider the canonical nondegenerate pairing

$$(13) \quad (L^\vee \otimes N) \otimes \underline{\text{Hom}}(N, L) \rightarrow \mathbb{1}$$

given (after applying ω) by

$$(\gamma \otimes x) \otimes f \mapsto \gamma(f(x)).$$

For any subobject A of $\underline{\text{Hom}}(N, L)$ (resp. $L^\vee \otimes N$), we denote by A^\perp the subobject of $L^\vee \otimes N$ (resp. $\underline{\text{Hom}}(N, L)$) orthogonal to A with respect to the above pairing. It is clear that A can be recovered from A^\perp by $A^{\perp\perp} = A$.

In particular, we have a subobject

$$\underline{\text{Lie}}(\mathcal{U}(M))^\perp \subset L^\vee \otimes N.$$

In this subsection we shall give a dual variant of [Theorem 3.3.1](#) which characterizes this object. In some situations (such as the application in [Section 4](#)), this variant might be more convenient to use than the original version.

Let

$$\mu \in \text{Ext}(L^\vee \otimes N, \mathbb{1})$$

be the extension class corresponding to the defining extension of M (i.e., [\(5\)](#)) under the canonical isomorphism

$$(14) \quad \text{Ext}(N, L) \cong \text{Ext}(L^\vee \otimes N, \mathbb{1}).$$

The extension class μ is obtained as follows. Let

$$ev : L^\vee \otimes L \rightarrow \mathbb{1}$$

be the evaluation pairing between L and its dual. Then μ is the class of the pushforward of the extension

$$0 \rightarrow L^\vee \otimes L \xrightarrow{I_{L^\vee} \otimes i} L^\vee \otimes M \xrightarrow{I_{L^\vee} \otimes q} L^\vee \otimes N \rightarrow 0$$

(obtained by tensoring (5) by L^\vee) through the morphism ev . More explicitly, when L is not zero, μ is given by the extension

$$(15) \quad 0 \rightarrow \mathbb{1} \rightarrow (L^\vee \otimes M)^\dagger \rightarrow L^\vee \otimes N \rightarrow 0,$$

where

- $(L^\vee \otimes M)^\dagger$ is the quotient of $L^\vee \otimes M$ by $(I_{L^\vee} \otimes i)(\ker(ev))$,
- the injective arrow is the composition

$$\mathbb{1} \xrightarrow{\simeq, \text{induced by } ev} (L^\vee \otimes L) / \ker(ev) \xrightarrow{\text{induced by } I_{L^\vee} \otimes i} (L^\vee \otimes M)^\dagger, \text{ and}$$

- the surjective arrow is induced by $I_{L^\vee} \otimes q$.

If $L = 0$, then μ is given by the extension

$$0 \rightarrow \mathbb{1} \rightarrow \mathbb{1} \rightarrow 0 \rightarrow 0.$$

For convenience, in this case we set $(L^\vee \otimes M)^\dagger := \mathbb{1}$.

We shall use the following notation for restrictions of extensions. For every subobject B of $L^\vee \otimes N$, let $\mu|_B$ be the restriction of μ to B (i.e., the pullback of μ along the inclusion map $B \rightarrow L^\vee \otimes N$).

We can now state the dual variants of [Theorem 3.3.1](#) and [Corollary 3.4.1](#).

Theorem 3.5.1. *Let B be a subobject of $L^\vee \otimes N$. Then*

$$B \subset \underline{\text{Lie}}(\mathcal{U}(M))^\perp$$

if and only if the preimage of B under the surjective arrow in (15) belongs to the subcategory $\langle L \oplus N \rangle$.

Proof. One can prove this directly, similar to the proof of [Theorem 3.3.1](#), by calculating the action of $\mathcal{U}(M)$ on $(L^\vee \otimes M)^\dagger$ (and its subobjects) explicitly. We shall instead use a few categorical considerations to show that the statement is equivalent to [Theorem 3.3.1](#). Let T be an object of \mathbf{T} . For any subobject A of T , denote by A^\perp the orthogonal complement of A with respect to the evaluation pairing

$$T^\vee \otimes T \rightarrow \mathbb{1}.$$

Dualizing the exact sequence

$$0 \rightarrow A \rightarrow T \rightarrow T/A \rightarrow 0,$$

we get

$$0 \rightarrow (T/A)^\vee \rightarrow T^\vee \rightarrow A^\vee \rightarrow 0.$$

Use this to identify

$$A^\perp \stackrel{\text{(by definition)}}{=} \ker(T^\vee \rightarrow A^\vee) \cong (T/A)^\vee.$$

There is a commutative diagram

$$\begin{array}{ccc} \mathrm{Ext}(\mathbb{1}, T) & \xrightarrow{\text{dualizing, } \cong} & \mathrm{Ext}(T^\vee, \mathbb{1}) \\ \text{pushforward} \downarrow & & \downarrow \text{pullback} \\ \mathrm{Ext}(\mathbb{1}, T/A) & \xrightarrow{\text{dualizing, } \cong} & \mathrm{Ext}(A^\perp, \mathbb{1}) \end{array}$$

where the horizontal maps dualize extensions. Apply this with $T = \underline{\mathrm{Hom}}(N, L)$, and use the pairing (13) to identify $L^\vee \otimes N$ as T^\vee (so that (13) becomes simply the evaluation pairing between T and T^\vee). It is easy to see that ν and μ are duals of one another, with the isomorphism between $(L^\vee \otimes M)^\dagger$ and the dual of $\underline{\mathrm{Hom}}(N, M)^\dagger$ defined by the pairing

$$(L^\vee \otimes M)^\dagger \otimes \underline{\mathrm{Hom}}(N, M)^\dagger \rightarrow \mathbb{1},$$

which after applying ω is given by

$$\overline{g \otimes x} \otimes f \mapsto g(\lambda(f)x - f((\omega q)(x))).$$

(Here $\overline{g \otimes x}$ is the image of $g \otimes x \in \omega(L^\vee) \otimes \omega M$ in $\omega(L^\vee \otimes M)^\dagger$, and f is in $\mathrm{Hom}(\omega N, \omega M)^\dagger$.) Thus by the above diagram, for any subobject A of $\underline{\mathrm{Hom}}(N, L)$, we have an isomorphism between

$$(\underline{\mathrm{Hom}}(N, M)^\dagger/A)^\vee$$

and the preimage of A^\perp under the surjective arrow in (15). The equivalence of Theorems 3.3.1 and 3.5.1 is clear from this. \square

The argument also gives the following dual variant of Corollary 3.4.1:

Corollary 3.5.2. (a) *If B is a subobject of $L^\vee \otimes N$ such that $\mu|_B$ is trivial, then $B \subset \underline{\mathrm{Lie}}(\mathcal{U}(M))^\perp$.*

(b) *Suppose L and N are semisimple. Then the restriction of μ to $\underline{\mathrm{Lie}}(\mathcal{U}(M))^\perp$ is trivial. (Hence $\underline{\mathrm{Lie}}(\mathcal{U}(M))^\perp$ is the largest subobject of $L^\vee \otimes N$ with this property.)*

4. The unipotent radical of the Mumford–Tate group of H^1 of an algebraic curve

Let \mathbf{MHS} be the category of rational mixed Hodge structures. The category \mathbf{MHS} is a neutral Tannakian category over \mathbb{Q} . The forgetful functor $\omega_B : \mathbf{MHS} \rightarrow \mathbf{Mod}_{\mathbb{Q}}$ sending an object to its underlying rational vector space is a fiber functor. For any

rational mixed Hodge structure M , the group $\mathcal{G}(M)$ with $(\mathbf{T}, \omega) = (\mathbf{MHS}, \omega_B)$ is called the Mumford–Tate group of M . In this section, we will use the results of the previous section to study the unipotent radical of the Mumford–Tate group of the degree one cohomology of a smooth complex projective curve minus a finite set of points.

4.1. Notation. By a mixed Hodge structure we always mean a rational one. As usual, $\mathbb{Q}(-n)$ denotes the Hodge structure of weight $2n$ with underlying rational vector space $(2\pi i)^{-n}\mathbb{Q}$, with its complexification identified with \mathbb{C} via

$$(2\pi i)^{-n} \otimes 1 \mapsto (2\pi i)^{-n}.$$

The unit object $\mathbb{1}$ is $\mathbb{Q}(0)$. For any object M of \mathbf{MHS} , we denote by $M_{\mathbb{Q}}$ the underlying rational vector space of M . If R is a commutative \mathbb{Q} -algebra, M_R denotes $M_{\mathbb{Q}} \otimes R$.

Given a pure Hodge structure H of weight -1 , we denote by JH the intermediate Jacobian

$$JH := \frac{H_{\mathbb{C}}}{F^0 H_{\mathbb{C}} + H_{\mathbb{Q}}},$$

where F^\cdot is the Hodge filtration.

Given any smooth complex variety X , by $H^i(X)$ we mean the mixed Hodge structure on the degree i Betti cohomology of X (with underlying rational vector space $H^i(X, \mathbb{Q})$). We shall identify $H^i(X)_{\mathbb{C}} = H^i(X, \mathbb{C})$ with $H^i_{dR}(X)$ (= smooth complex de Rham cohomology) via the isomorphism of de Rham. By $H_i(X)$ we mean the dual of $H^i(X)$; it is a mixed Hodge structure with underlying rational vector space $H_i(X, \mathbb{Q})$.

All the Ext (= Yoneda Ext^1) groups in this section are in \mathbf{MHS} .

4.2. Carlson [6] gives an explicit description of Ext groups in \mathbf{MHS} . We briefly recall this description here in a special case that is of interest to us.

Let A be a pure Hodge structure of weight 1. Carlson gives a canonical isomorphism

$$\text{Ext}(A, \mathbb{1}) \rightarrow J(A^\vee),$$

where A^\vee is the dual Hodge structure to A . The isomorphism is functorial in A . See [6] for details. (Carlson [6] proves the analogous result for integral mixed Hodge structures. The proof of the rational case is identical.)

4.3. From this point on, let X be a smooth complex projective curve. We denote the Jacobian variety of X by $\text{Jac}(X)$. Let $CH_0^{\text{hom}}(X)$ be the group of divisors of degree 0 on X modulo the subgroup of principal divisors. (In other words, $CH_0^{\text{hom}}(X)$ is the homologically trivial subgroup of the Chow group $CH_0(X)$.) The group $CH_0^{\text{hom}}(X)$ is the group of complex points of $\text{Jac}(X)$.

Set $CH_0^{\text{hom}}(X)_{\mathbb{Q}} := CH_0^{\text{hom}}(X) \otimes \mathbb{Q}$. The Abel–Jacobi map on X gives an isomorphism

$$AJ_X : CH_0^{\text{hom}}(X)_{\mathbb{Q}} \rightarrow JH_1(X),$$

sending the class of $p - q$, with $p, q \in X$, to the class of the functional \int_q^p on the space of harmonic 1-forms on X . (See, for instance, [2, Chapter 1]. Note that here, said integral means the integral over any path from q to p . The choice of the path will not matter in $JH_1(X)$.)

Composing AJ_X with Carlson’s isomorphism we get an isomorphism

$$(16) \quad \text{Ext}(H^1(X), \mathbb{1}) \cong CH_0^{\text{hom}}(X)_{\mathbb{Q}}.$$

We shall identify these two groups to simplify the notation.

4.4. Let S be a finite nonempty set of (complex) points of X . We identify $H^1(X)$ as a subobject of $H^1(X \setminus S)$ via the map induced by the inclusion $(X \setminus S) \subset X$. The reader can refer to Deligne’s [7, Section 10.3] for a thorough study of the mixed Hodge structure $H^1(X \setminus S)$.

Since $X \setminus S$ is affine, every element of $H^1(X \setminus S)_{\mathbb{C}}$ can be represented by a meromorphic differential form on X with possible singularities only along S , and has a well-defined residue at every $p \in X$. Indeed, if $c = [\omega]$ with ω a meromorphic form, set $\text{res}_p(c) := \text{res}_p(\omega)$ (= the residue of ω at p , which is $1/(2\pi i)$ times the integral of ω along a small positively oriented loop around p). The subspace $H^1(X)_{\mathbb{C}}$ of $H^1(X \setminus S)_{\mathbb{C}}$ consists of the cohomology classes with zero residue everywhere (in other words, classes of differentials of the second kind).

For any vector space or mixed Hodge structure V , we denote by $(V^S)'$ the kernel of the map

$$V^S \rightarrow V, \quad (v_p)_{p \in S} \mapsto \sum_{p \in S} v_p$$

(where the v_p are in V).

One has a short exact sequence of mixed Hodge structures

$$(17) \quad 0 \rightarrow H^1(X) \rightarrow H^1(X \setminus S) \xrightarrow{\text{res}_S} (\mathbb{Q}(-1)^S)' \rightarrow 0,$$

where the injective arrow is inclusion and $\text{res}_S : H^1(X \setminus S)_{\mathbb{C}} \rightarrow (\mathbb{C}^S)'$ is the map $c \mapsto (\text{res}_p(c))_{p \in S}$.

4.5. We shall apply the results of Section 3.3 to the exact sequence (17). The Hodge structure $H^1(X)$ is polarizable and hence semisimple (see, for instance, [19, Section 7.1.2] and [9, Proposition 3.6]). Thus the group

$$U(H^1(X \setminus S)) := \ker(\mathcal{G}(H^1(X \setminus S)) \rightarrow \mathcal{G}(H^1(X) \oplus \mathbb{Q}(-1)))$$

is the unipotent radical of the Mumford–Tate group of $H^1(X \setminus S)$. In view of Section 3.5, the determination of the group $\mathcal{U}(H^1(X \setminus S))$ amounts to finding

$$\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^\perp \subset H^1(X)^\vee \otimes (\mathbb{Q}(-1)^S)'$$

We use the Poincaré duality isomorphism

$$PD : H^1(X)(1) \rightarrow H^1(X)^\vee, \quad [\eta] \mapsto \frac{1}{2\pi i} \int_X \eta \wedge -,$$

where η is a closed smooth 1-form on X and the isomorphism

$$H^1(X)(1) \otimes (\mathbb{Q}(-1)^S)' \rightarrow (H^1(X)^S)', \quad c \otimes (a_p)_{p \in S} \mapsto (a_p c)_{p \in S}$$

to identify

$$H^1(X)^\vee \otimes (\mathbb{Q}(-1)^S)' \cong (H^1(X)^S)'$$

Following the notation of Section 3.5, we let

$$\mu \in \text{Ext}((H^1(X)^S)', \mathbb{1})$$

be the element corresponding to the sequence (17) under the canonical isomorphism

$$(18) \quad \begin{aligned} \text{Ext}((\mathbb{Q}(-1)^S)', H^1(X)) &\cong \text{Ext}(H^1(X)^\vee \otimes (\mathbb{Q}(-1)^S)', \mathbb{1}) \\ &= \text{Ext}((H^1(X)^S)', \mathbb{1}). \end{aligned}$$

By Corollary 3.5.2 (and on recalling that $H^1(X)$ is semisimple), we have that $\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^\perp$ is the largest subobject of $(H^1(X)^S)'$ with the property that the restriction of μ to it is trivial.

4.6. Let us consider the restrictions of μ to some obvious subobjects of $(H^1(X)^S)'$. For each $p \in S$, let $\iota_p : H^1(X) \rightarrow H^1(X)^S$ be the embedding into the p -coordinate. Given $p, q \in S$, we have a morphism

$$\iota_p - \iota_q : H^1(X) \rightarrow (H^1(X)^S)'$$

(which is an embedding if $p \neq q$).

Proposition 4.6.1. *Let $p, q \in S$. Via the identification (16), we have*

$$(\iota_p - \iota_q)^*(\mu) = p - q$$

(where $(\iota_p - \iota_q)^*(\mu)$ is the pullback of μ along $\iota_p - \iota_q$, and with abuse of notation the class of $p - q$ in $CH_0^{\text{hom}}(X)_{\mathbb{Q}}$ is also denoted by $p - q$).

Proof. This is a reformulation of a well-known result about Hodge theory of open curves, which in turn is a special case of general results about equivalence of various definitions of the Abel–Jacobi map (see the remark below). With abuse of notation,

let ι_p also denote the embedding of $\mathbb{Q}(-1)$ as the p -coordinate of $\mathbb{Q}(-1)^S$. Then we have a commutative diagram

$$\begin{CD} \text{Ext}((\mathbb{Q}(-1)^S)', H^1(X)) @>{(\iota_p - \iota_q)^*}>> \text{Ext}(\mathbb{Q}(-1), H^1(X)) \\ @A{\cong}A\downarrow @A{\cong}A\downarrow \\ \text{Ext}((H^1(X)^S)', \mathbb{1}) @>{(\iota_p - \iota_q)^*}>> \text{Ext}(H^1(X), \mathbb{1}) \end{CD}$$

where the vertical isomorphisms are given by (14) and Poincaré duality. Under the isomorphism on the left (i.e., (18)), μ and (17) correspond to each other. The pullback of the extension (17) along $\iota_p - \iota_q : \mathbb{Q}(-1) \rightarrow (\mathbb{Q}(-1)^S)'$ is the extension

$$(19) \quad \begin{array}{ccccccc} 0 & \rightarrow & H^1(X) & \rightarrow & (\text{res}_S)^{-1}((\iota_p - \iota_q)(\mathbb{Q}(-1))) & \xrightarrow{\text{res}_p} & \mathbb{Q}(-1) \rightarrow 0 \\ & & & & \parallel & & \\ & & & & H^1(X \setminus \{p, q\}) & & \end{array}$$

This extension corresponds to $p - q$ under

$$(20) \quad \text{Ext}(\mathbb{Q}(-1), H^1(X)) \cong \text{Ext}(H^1(X), \mathbb{1}) \cong JH_1(X) \cong CH_0^{\text{hom}}(X)_{\mathbb{Q}}.$$

See, for example, Sections 9.0–9.2 of Jannsen [17]. □

Remark 4.6.2. The fact that the extension (19) corresponds to $p - q$ under (20) is already stated in Section 4.3 of Deligne’s [8]. The same paragraph outlines a motivically inspired definition of the Abel–Jacobi map, which naturally takes values in Ext groups in any suitable cohomology theory. Via this approach and in the case of Hodge theory (or more precisely, cohomology with values in **MHS**), the Abel–Jacobi image of $p - q$ is by definition the extension (19) (in other cohomology theories, by definition the Abel–Jacobi image is the analogous extension). A detailed description of this motivic approach towards the Abel–Jacobi map for any smooth complex variety can be found in Sections 9.0 and 9.1 of Jannsen’s book [17]. The fact that for Hodge theory the Abel–Jacobi map defined in terms of extensions coincides with the classical (Griffiths) Abel–Jacobi map with values in intermediate Jacobians is asserted in Lemma 9.2 of [17] and follows from the works [16] and [13] of Jannsen and Esnault–Viehweg. (See Section 9.2 of [17] for more details.)

4.7. We now calculate the slightly more complicated restrictions of μ . Let

$$E = \text{End}^0(\text{Jac}(X)) := \text{End}(\text{Jac}(X)) \otimes \mathbb{Q}$$

be the endomorphism algebra of the Jacobian of X . We have an (anti-) isomorphism

$$E \rightarrow \text{End}(H^1(X)), \quad f \mapsto f^*,$$

where for any element f of the endomorphism algebra of $\text{Jac}(X)$, by f^* we mean the pullback map on cohomology.¹ This induces an isomorphism

$$(21) \quad (E^S)' \rightarrow \text{Hom}(H^1(X), (H^1(X)^S)'), \quad (f_p)_{p \in S} \mapsto \sum_{p \in S} \iota_p f_p^*.$$

Consider the composition

$$(22) \quad \text{Hom}(H^1(X), (H^1(X)^S)') \xrightarrow{\phi \mapsto \phi^* \mu} \text{Ext}(H^1(X), \mathbb{1}) \stackrel{(16)}{\cong} CH_0^{\text{hom}}(X)_{\mathbb{Q}} = \text{Jac}(X)(\mathbb{C}) \otimes \mathbb{Q}.$$

Since every simple subobject of $(H^1(X)^S)'$ is the image of a morphism

$$H^1(X) \rightarrow (H^1(X)^S)'$$

(because $\langle H^1(X) \rangle$ is semisimple), the following corollary of [Proposition 4.6.1](#) can be used to describe all restrictions of μ .

Corollary 4.7.1. *Let $(f_p)_{p \in S} \in (E^S)'$. Then*

$$\left(\sum_{p \in S} \iota_p f_p^* \right)^* \mu = \sum_{p \in S} f_p(p - e),$$

where e is any point in X .

Proof. Let $(f_p)_{p \in S} \in (E^S)'$ and $e \in S$. Since $\sum_{p \in S} f_p = 0$, we have

$$\sum_{p \in S} \iota_p f_p^* = \sum_{p \in S} (\iota_p - \iota_e) f_p^*.$$

Thus

$$\left(\sum_{p \in S} \iota_p f_p^* \right)^* \mu = \sum_{p \in S} ((\iota_p - \iota_e) f_p^*)^* \mu = \sum_{p \in S} (f_p^*)^* (\iota_p - \iota_e)^* \mu = \sum_{p \in S} f_p(p - e),$$

where in the last line we used [Proposition 4.6.1](#) together with the commutativity of the diagram

$$\begin{array}{ccc} \text{Ext}(H^1(X), \mathbb{1}) \cong JH_1(X) & \xleftarrow{\simeq, AJ_X} & CH_0^{\text{hom}}(X)_{\mathbb{Q}} = \text{Jac}(X)(\mathbb{C}) \otimes \mathbb{Q} \\ \downarrow (f^*)^* & & \downarrow f \\ \text{Ext}(H^1(X), \mathbb{1}) \cong JH_1(X) & \xleftarrow{\simeq, AJ_X} & CH_0^{\text{hom}}(X)_{\mathbb{Q}} = \text{Jac}(X)(\mathbb{C}) \otimes \mathbb{Q} \quad \square \end{array}$$

For any $(f_p)_{p \in S} \in (E^S)'$, the value of $\sum_{p \in S} f_p(p - e)$ does not depend on the choice of $e \in X$. To simplify the notation, let us denote this common value by

$$\sum_{p \in S} f_p(p).$$

¹We use the symbol $*$ for pullback of extensions and the symbol \star for pullback of cohomology induced by morphisms of varieties.

Note that if

$$(f_p)_{p \in S} \in (\mathbb{Q}^S)' \subset (E^S)',$$

then $\sum_{p \in S} f_p(p)$ defined above agrees with the other possible interpretation of the notation (i.e., the image of the divisor $\sum_{p \in S} f_p p$ of degree zero with coefficients in \mathbb{Q} in $CH_0^{\text{hom}}(X)_{\mathbb{Q}}$).

4.8. Having computed the restrictions of μ , we return to the problem of determination of $\mathcal{U}(H^1(X \setminus S))$.

Proposition 4.8.1. *Let $(f_p)_{p \in S} \in (E^S)'$. The following statements are equivalent:*

- (i) *The restriction of μ to the image of $\sum_{p \in S} \iota_p f_p^*$ splits.*
- (ii) *$\sum_{p \in S} f_p(p)$ is zero in $CH_0^{\text{hom}}(X)_{\mathbb{Q}}$.*
- (iii) *The image of $\sum_{p \in S} \iota_p f_p^*$ is contained in $\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^{\perp}$.*

Proof. Recall that by [Corollary 3.5.2](#), $\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^{\perp}$ is the largest subobject of $(H^1(X)^S)'$ with the property that the restriction of μ to it splits (see [Section 4.5](#)). This gives the equivalence of statements (i) and (iii) of the proposition. The equivalence of (i) and (ii) follows from [Corollary 4.7.1](#), on noting (by weight considerations) that for any quotient B of $H^1(X)$, the canonical map

$$\text{Ext}(B, \mathbb{1}) \rightarrow \text{Ext}(H^1(X), \mathbb{1})$$

is injective. □

In particular, the proposition recovers the following well-known result, originally due to Deligne (see the remark below), which gives an arithmetic criterion for when $\mathcal{U}(H^1(X \setminus S))$ is trivial (or equivalently, for when the sequence [\(17\)](#) splits):

Corollary 4.8.2. *The group $\mathcal{U}(H^1(X \setminus S))$ is trivial if and only if the subgroup of the Jacobian of X supported on S has zero rank.*

Proof. Note that $\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^{\perp} = (H^1(X)^S)'$ if and only if $\text{Im}(\iota_p - \iota_q)$ is contained in $\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^{\perp}$ for every $p, q \in S$, which in turn is equivalent to $p - q$ being zero in $CH_0^{\text{hom}}(X)_{\mathbb{Q}}$ for every $p, q \in S$. □

Remark 4.8.3. [Corollary 4.8.2](#) is originally due to Deligne, implicit in [\[7\]](#) and announced explicitly in [\[8, Remarque 7.5\]](#), in relation to a new proof of the Manin–Drinfeld theorem on modular curves. See [\[11\]](#) for a more detailed discussion of this.

4.9. We are ready to give the main result of this part of the paper. The results gives a characterization of $\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^{\perp}$ (and hence $\mathcal{U}(H^1(X \setminus S))$).

Theorem 4.9.1. *Let A be the subobject of $(H^1(X)^S)'$ which is the sum of the images of all the maps of the form*

$$\sum_{p \in S} \iota_p f_p^* \in \text{Hom}(H^1(X), (H^1(X)^S)'),$$

with $(f_p)_{p \in S} \in (E^S)'$ and $\sum_{p \in S} f_p(p) = 0$ (see Section 4.7). Then

$$A = \underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^\perp.$$

Proof. The inclusion

$$A \subset \underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^\perp$$

is immediate from Proposition 4.8.1. To see the reverse inclusion, first note that since $\langle H^1(X) \rangle$ is semisimple, $\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^\perp$ is a direct sum of simple subobjects. Let B be a simple subobject of $\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^\perp$. Then B is the image of a map $H^1(X) \rightarrow (H^1(X)^S)'$. Any such map is of the form $\sum_{p \in S} \iota_p f_p^*$ for some $(f_p)_{p \in S} \in (E^S)'$. By Proposition 4.8.1, for the image of such a map to be in $\underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^\perp$ we must have

$$\sum_{p \in S} f_p(p) = 0$$

in $CH_0^{\text{hom}}(X)_{\mathbb{Q}}$. Thus $B \subset A$. □

We end the paper by deducing the following result about the dimension of $\mathcal{U}(H^1(X \setminus S))$ (note that part (b) is Theorem A of the introduction).

Theorem 4.9.2. *Let g be the genus of X . Recall that E is the endomorphism algebra of the Jacobian $\text{Jac}(X)$.*

- (a) *Suppose D is any division algebra contained in E . Then the dimension of $\mathcal{U}(H^1(X \setminus S))$ is at most $2g$ times the D -rank of the D -submodule of $\text{Jac}(X)(\mathbb{C}) \otimes \mathbb{Q}$ generated by the subgroup supported on S .*
- (b) *Suppose $H^1(X)$ is simple. Then the dimension of $\mathcal{U}(H^1(X \setminus S))$ is equal to $2g$ times the E -rank of the E -submodule of $\text{Jac}(X)(\mathbb{C}) \otimes \mathbb{Q}$ generated by the subgroup supported on S .*

Proof. Let A be as in Theorem 4.9.1.

- (a) For any subalgebra R of E , let Λ_R be the composition

$$(R^S)' \hookrightarrow (E^S)' \xrightarrow{(21)} \text{Hom}(H^1(X), (H^1(X)^S)') \xrightarrow{(22)} \text{Jac}(X)(\mathbb{C}) \otimes \mathbb{Q}.$$

This is R -linear by Corollary 4.7.1. The image of Λ_R is the R -submodule of $\text{Jac}(X)(\mathbb{C}) \otimes \mathbb{Q}$ generated by the subgroup supported on S . Let A_R be the subobject of $(H^1(X)^S)'$ which is the sum of the images of the maps $\sum_{p \in S} \iota_p f_p^*$ with $(f_p)_{p \in S}$ in $\ker(\Lambda_R)$, so that $A_R \subset A$ and $A_E = A$. If $\beta = \{(f_p^{(r)})_{p \in S}\}_{1 \leq r \leq d}$ is an R -spanning set for $\ker(\Lambda_R)$, then A_R is the sum of the images of $\sum_{p \in S} \iota_p (f_p^{(r)})^*$ for $1 \leq r \leq d$. Moreover, if $R = D$ is a division algebra and β is D -linearly independent, then A_D is the direct sum of the images of the previous d maps. Since each of these images is then a copy of $H^1(X)$ (because D is a division algebra), we have

$$\begin{aligned} \dim \underline{\text{Lie}}(\mathcal{U}(H^1(X \setminus S)))^\perp &= \dim(A) \geq \dim(A_D) = 2g \cdot \dim_D(\ker(\Lambda_D)) \\ &= 2g(|S| - 1 - \dim_D \text{Im}(\Lambda_D)). \end{aligned}$$

Taking orthogonal complements we get the desired bound.

(b) Since $H^1(X)$ is simple, E is a division algebra. Taking $D = E$, by the proof of part (b) we have

$$\dim(A) = \dim(A_E) = 2g(|S| - 1 - \dim_E \operatorname{Im}(\Lambda_E)).$$

The claimed formula follows. □

Acknowledgements

We thank Madhav Nori for his comments on a draft of the paper. When we wrote the first draft of this paper, we were not aware of the related works of Bertrand [5], Hardouin [14; 15] and Jossen [18]. We thank the referees of the earlier versions of the article for bringing these works to our attention, as well as for providing comments that helped us improve the results and exposition of the article. We also thank a referee for the *Pacific Journal of Mathematics* for providing us with many comments that helped us significantly improve the exposition of the second part of the paper.

Eskandari was at the University of Toronto and the Fields Institute for Research in Mathematical Sciences when this work was done. He wishes to thank both institutions for providing a pleasant working environment.

This work was a prequel to the work [12].

References

- [1] Y. André, “Mumford–Tate groups of mixed Hodge structures and the theorem of the fixed part”, *Compositio Math.* **82**:1 (1992), 1–24. [MR](#) [Zbl](#)
- [2] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris, *Geometry of algebraic curves, I*, Grundle Math. Wissen. **267**, Springer, New York, 1985. [MR](#) [Zbl](#)
- [3] C. Bertolin, “The Mumford–Tate group of 1-motives”, *Ann. Inst. Fourier (Grenoble)* **52**:4 (2002), 1041–1059. [MR](#) [Zbl](#)
- [4] C. Bertolin, “Le radical unipotent du groupe de Galois motivique d’un 1-motif”, *Math. Ann.* **327**:3 (2003), 585–607. [MR](#) [Zbl](#)
- [5] D. Bertrand, “Unipotent radicals of differential Galois group and integrals of solutions of inhomogeneous equations”, *Math. Ann.* **321**:3 (2001), 645–666. [MR](#)
- [6] J. A. Carlson, “Extensions of mixed Hodge structures”, pp. 107–127 in *Journées de Géométrie Algébrique d’Angers* (Juillet, 1979), Sijthoff & Noordhoff, Alphen aan den Rijn, 1980. [MR](#) [Zbl](#)
- [7] P. Deligne, “Théorie de Hodge, III”, *Inst. Hautes Études Sci. Publ. Math.* **44** (1974), 5–77. [MR](#) [Zbl](#)
- [8] P. Deligne, “Valeurs de fonctions L et périodes d’intégrales”, pp. 313–346 in *Proc. Sympos. Pure Math., part 2*, vol. 33, Amer. Math. Soc., Providence, R.I., 1979. [MR](#) [Zbl](#)
- [9] P. Deligne, “Hodge cycles on abelian varieties”, pp. 9–100 in *Hodge cycles, motives, and Shimura varieties*, Lecture Notes in Mathematics **900**, Springer, Berlin, 1982. [Zbl](#)

- [10] P. Deligne and J. S. Milne, “Tannakian categories”, pp. 101–228 in *Hodge cycles, motives, and Shimura varieties*, Lecture Notes in Mathematics **900**, Springer, Berlin, 1982. [Zbl](#)
- [11] R. Elkik, “Le théorème de Manin–Drinfeld”, pp. 59–67 in *Séminaire sur les Pinceaux de Courbes Elliptiques*, Astérisque **183**, Société Mathématique de France, Paris, 1990. [MR](#) [Zbl](#)
- [12] P. Eskandari and V. K. Murty, “On unipotent radicals of motivic Galois groups”, *Algebra Number Theory* **17**:1 (2023), 165–215. [MR](#)
- [13] H. Esnault and E. Viehweg, “Deligne–Beilinson cohomology”, pp. 43–91 in *Beilinson’s conjectures on special values of L-functions*, *Perspect. Math.* **4**, Academic Press, Boston, MA, 1988. [MR](#) [Zbl](#)
- [14] C. Hardouin, “Hypertranscendance et Groupes de Galois aux différences”, 2006. [arXiv 0609646](#)
- [15] C. Hardouin, “Unipotent radicals of Tannakian Galois groups in positive characteristic”, pp. 223–239 in *Arithmetic and Galois theories of differential equations*, *Sémin. Congr.* **23**, Soc. Math. France, Paris, 2011. [MR](#) [Zbl](#)
- [16] U. Jannsen, “Deligne homology, Hodge- \mathcal{D} -conjecture, and motives”, pp. 305–372 in *Beilinson’s conjectures on special values of L-functions*, *Perspect. Math.* **4**, Academic Press, Boston, MA, 1988. [MR](#) [Zbl](#)
- [17] U. Jannsen, *Mixed motives and algebraic K-theory*, Lecture Notes in Mathematics **1400**, Springer, Berlin, 1990. [MR](#) [Zbl](#)
- [18] P. Jossen, “On the Mumford–Tate conjecture for 1-motives”, *Invent. Math.* **195**:2 (2014), 393–439. [MR](#) [Zbl](#)
- [19] C. Voisin, *Hodge theory and complex algebraic geometry, I*, Cambridge Studies in Advanced Mathematics **76**, Cambridge University Press, Cambridge, 2002. [MR](#) [Zbl](#)

Received August 20, 2021. Revised March 23, 2023.

PAYMAN ESKANDARI
 DEPARTMENT OF MATHEMATICS AND STATISTICS
 UNIVERSITY OF WINNIPEG
 WINNIPEG, MB
 CANADA
p.eskandari@uwinnipeg.ca

V. KUMAR MURTY
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF TORONTO
 TORONTO, ON
 CANADA
murty@math.toronto.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2023 is US \$605/year for the electronic version, and \$820/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 325 No. 2 August 2023

Totally geodesic hyperbolic 3-manifolds in hyperbolic link complements of tori in S^4	191
MICHELLE CHU and ALAN W. REID	
Infinite homotopy stable class for 4-manifolds with boundary	209
ANTHONY CONWAY, DIARMUID CROWLEY and MARK POWELL	
Counterexamples to the nonsimply connected double soul conjecture	239
JASON DEVITO	
The fundamental group of an extension in a Tannakian category and the unipotent radical of the Mumford–Tate group of an open curve	255
PAYMAN ESKANDARI and V. KUMAR MURTY	
The size of semigroup orbits modulo primes	281
WADE HINDES and JOSEPH H. SILVERMAN	
Tropical Lagrangian multisections and toric vector bundles	299
YAT-HIN SUEN	
Some effectivity results for primitive divisors of elliptic divisibility sequences	331
MATTEO VERZOBIO	
Poisson manifolds of strong compact type over 2-tori	353
LUKA ZWAAN	