

*Pacific
Journal of
Mathematics*

**LOCAL GALOIS REPRESENTATIONS
OF SWAN CONDUCTOR ONE**

NAOKI IMAI AND TAKAHIRO TSUSHIMA

LOCAL GALOIS REPRESENTATIONS OF SWAN CONDUCTOR ONE

NAOKI IMAI AND TAKAHIRO TSUSHIMA

We construct the local Galois representations over the complex field whose Swan conductors are one by using étale cohomology of Artin–Schreier sheaves on affine lines over finite fields. Then, we study the Galois representations, and give an explicit description of the local Langlands correspondences for simple supercuspidal representations. We discuss also a more natural realization of the Galois representations in the étale cohomology of Artin–Schreier varieties.

Introduction

Let K be a nonarchimedean local field. Let n be a positive integer. The existence of the local Langlands correspondence for $\mathrm{GL}_n(K)$, proved in [Laumon et al. 1993] and [Harris and Taylor 2001], is one of the fundamental results in the Langlands program. However, even in this fundamental case, an explicit construction of the local Langlands correspondence has not yet been obtained. One of the most striking results in this direction is the result of Bushnell and Henniart [2005a; 2005b; 2010] for essentially tame representations. On the other hand, we don't know much about the explicit construction outside essentially tame representations.

We discuss this problem for representations of Swan conductor 1. The irreducible supercuspidal representations of $\mathrm{GL}_n(K)$ of Swan conductor 1 are equivalent to the simple supercuspidal representations in the sense of Adrian and Liu [2016] (see [Gross and Reeder 2010; Reeder and Yu 2014]). Such representations are called “epipelagic” in [Bushnell and Henniart 2014].

Let p be the characteristic of the residue field k of K . If n is prime to p , the simple supercuspidal representations of $\mathrm{GL}_n(K)$ are essentially tame. Hence, this case is covered by the work of Bushnell and Henniart. See also [Adrian and Liu 2016]. It is discussed in [Kaletha 2015] to generalize the construction of the local Langlands correspondence for essentially tame epipelagic representations to other reductive groups.

MSC2020: primary 11F80; secondary 11F70.

Keywords: Galois representation, simple supercuspidal representation, local Langlands correspondence.

In this paper, we consider the case where p divides n . In this case, the simple supercuspidal representations of $\mathrm{GL}_n(K)$ are not essentially tame. Moreover, if n is a power of p , the irreducible representations of the Weil group W_K of Swan conductor 1, which correspond to the simple supercuspidal representations via the local Langlands correspondence, cannot be induced from any proper subgroup. Such representations are called primitive (see [Koch 1977]). For simple supercuspidal representations, we have a straightforward characterization of the local Langlands correspondence given in [Bushnell and Henniart 2014]. Further, Bushnell and Henniart study the restriction to the wild inertia subgroup of the Langlands parameters for the simple supercuspidal representations explicitly. Actually, the restriction to the wild inertia subgroup already determines the original Langlands parameters up to character twists, but we need additional data, which appear in Bushnell and Henniart's characterization, to pin down the correct Langlands parameters. On the other hand, the construction of the irreducible representations of W_K of Swan conductor 1 is a nontrivial problem. What we will do in this paper is

- to construct the irreducible representations of W_K of Swan conductor 1 without appealing to the existence of the local Langlands correspondence, and
- to give a description of the Langlands parameters themselves for the simple supercuspidal representations.

Let ℓ be a prime number different from p . For the construction of the irreducible representations of W_K of Swan conductor 1, we use étale cohomology of an Artin–Schreier ℓ -adic sheaf on $\mathbb{A}_{k^{\mathrm{ac}}}^1$, where k^{ac} is an algebraic closure of k . It will be possible to avoid usage of geometry in the construction of the irreducible representations of W_K of Swan conductor 1. However, we prefer this approach, because

- we can use geometric tools such as the Lefschetz trace formula and the product formula of Deligne–Laumon to study the constructed representations, and
- the construction works also for ℓ -adic integral coefficients and mod ℓ coefficients.

A description of the local Langlands correspondence for the simple supercuspidal representations is discussed in [Imai and Tsushima 2022] in the special case where $n = p = 2$. Even in the special case, our method in this paper is totally different from that in [Imai and Tsushima 2022].

We explain the main result. We write $n = p^e n'$, where n' is prime to p . We fix a uniformizer ϖ of K and an isomorphism $\iota : \overline{\mathbb{Q}}_\ell \simeq \mathbb{C}$.

Let \mathcal{L}_ψ be the Artin–Schreier $\overline{\mathbb{Q}}_\ell$ -sheaf on $\mathbb{A}_{k^{\mathrm{ac}}}^1$ associated to a nontrivial character ψ of \mathbb{F}_p . Let $\pi : \mathbb{A}_{k^{\mathrm{ac}}}^1 \rightarrow \mathbb{A}_{k^{\mathrm{ac}}}^1$ be the morphism defined by $\pi(y) = y^{p^e+1}$. Let $\zeta \in \mu_{q-1}(K)$, where $q = |k|$. We put $E_\zeta = K[X]/(X^{n'} - \zeta\varpi)$. Then we can define

a natural action of W_{E_ζ} on $H_C^1(\mathbb{A}_{k^{\text{ac}}}^1, \pi^* \mathcal{L}_\psi)$. Using this action, we can associate a primitive representation $\tau_{n,\zeta,\chi,c}$ of W_{E_ζ} to $\zeta \in \mu_{q-1}(K)$, a character χ of k^\times and $c \in \mathbb{C}^\times$. We construct an irreducible representation $\tau_{\zeta,\chi,c}$ of Swan conductor 1 as the induction of $\tau_{n,\zeta,\chi,c}$ to W_K .

We can associate a simple supercuspidal representation $\pi_{\zeta,\chi,c}$ of $\text{GL}_n(K)$ to the same triple (ζ, χ, c) by type theory. Any simple supercuspidal representation can be written in this form uniquely (see [Imai and Tsushima 2018, Proposition 1.3]).

Theorem. *The representations $\tau_{\zeta,\chi,c}$ and $\pi_{\zeta,\chi,c}$ correspond via the local Langlands correspondence.*

In Section 1, we recall a general fact on representations of a semidirect product of a Heisenberg group with a cyclic group. In Section 2, we give a construction of the irreducible representations of W_K of Swan conductor 1. To construct a representation of W_K which naturally fits a description of the local Langlands correspondence, we need a subtle character twist. Such a twist appears also in the essentially tame case in [Bushnell and Henniart 2010], where it is called a rectifier. Our twist can be considered as an analogue of the rectifier. We construct the representations of W_K using geometry, but we give also a representation theoretic characterization of the constructed representations. In Section 3, we give a construction of the simple supercuspidal representations of $\text{GL}_n(K)$ using the type theory.

In Section 4, we state the main theorem and recall a characterization of the local Langlands correspondence for simple supercuspidal representations given in [Bushnell and Henniart 2014]. The characterization consists of the three equalities of (i) the determinant and the central character, (ii) the refined Swan conductors, and (iii) the epsilon factors.

In Section 5, we recall some general facts on epsilon factors. In Section 6, we recall facts on Stiefel–Whitney classes, multiplicative discriminants and additive discriminants. We use these facts to calculate Langlands constants of wildly ramified extensions. In Section 7, we recall the product formula of Deligne–Laumon. In Section 8, we show the equality of the determinant and the central character using the product formula of Deligne–Laumon.

In Section 9, we construct a field extension T_ζ^u of E_ζ such that the restriction of $\tau_{n,\zeta,\chi,c}$ to $W_{T_\zeta^u}$ is an induction of a character and $p \nmid [T_\zeta^u : E_\zeta]$, which we call an imprimitive field. In Section 10, we show the equality of the refined Swan conductors. We see also that the constructed representations of W_K are actually of Swan conductor 1.

In Section 11, we show the equality of the epsilon factors. It is difficult to calculate the epsilon factors of irreducible representations of W_K of Swan conductor 1 directly, because primitive representations are involved. However, we know the equality of the epsilon factors up to p^e -th roots of unity if $n = p^e$, since we have already

checked the conditions (i) and (ii) in the characterization. Using this fact and $p \nmid [T_\zeta^u : E_\zeta]$, the problem is reduced to study an epsilon factor of a character. Next we reduce the problem to the case where the characteristic of K is p and $k = \mathbb{F}_p$. At this stage, it is possible to calculate the epsilon factor if $p \neq 2$. However, it is still difficult if $p = 2$, because the direct calculation of the epsilon factor involves an explicit study of the Artin reciprocity map for a wildly ramified extension with a nontrivial ramification filtration. This is a special phenomenon in the case where $p = 2$. We will avoid this difficulty by reducing the problem to the case where $e = 1$. In this case, we have already known the equality up to sign. Hence, it suffices to show the equality of nonzero real parts. This is easy, because the difficult study of the Artin reciprocity map involves only the imaginary part of the equality.

In [Appendix](#), we discuss a construction of irreducible representations of W_K of Swan conductor 1 in the cohomology of Artin–Schreier varieties. This geometric construction incorporates a twist by a “rectifier”. We see that the “rectifier” parts come from the cohomology of Artin–Schreier varieties associated to quadratic forms. The Artin–Schreier varieties which we use have origins in studies of Lubin–Tate spaces in [[Imai and Tsushima 2017; 2021](#)].

Notation. Let A^\vee denote the character group $\text{Hom}_{\mathbb{Z}}(A, \mathbb{C}^\times)$ for a finite abelian group A . For a nonarchimedean local field K , let

- \mathcal{O}_K denote the ring of integers of K ,
- \mathfrak{p}_K denote the maximal ideal of \mathcal{O}_K ,
- v_K denote the normalized valuation of K which sends a uniformizer of K to 1,
- $\text{ch } K$ denote the characteristic of K ,
- G_K denote the absolute Galois group of K ,
- W_K denote the Weil group of K ,
- I_K denote the inertia subgroup of W_K ,
- P_K denote the wild inertia subgroup of W_K ,

and we put $U_K^m = 1 + \mathfrak{p}_K^m$ for any positive integer m .

1. Representations of finite groups

First, we recall a fact on representations of Heisenberg groups. Let G be a finite group with center Z . We assume:

- (i) The group G/Z is an elementary abelian p -group.
- (ii) For any $g \in G \setminus Z$, the map $c_g : G \rightarrow Z$, $g' \mapsto [g, g']$ is surjective.

Remark 1.1. The map c_g in (ii) is a group homomorphism. Hence, Z is automatically an elementary abelian p -group.

Let $\psi \in Z^\vee$ be a nontrivial character.

Proposition 1.2. *There is a unique irreducible representation ρ_ψ of G such that $\rho_\psi|_Z$ contains ψ . Moreover, we have $(\dim \rho_\psi)^2 = [G : Z]$ and we can construct ρ_ψ as follows: Take an abelian subgroup G_1 of G such that $Z \subset G_1$ and $2 \dim_{\mathbb{F}_p}(G_1/Z) = \dim_{\mathbb{F}_p}(G/Z)$. Extend ψ to a character ψ_1 of G_1 . Then $\rho_\psi = \text{Ind}_{G_1}^G \psi_1$.*

Proof. The claims other than the construction of ρ_ψ is Proposition 8.3.3 in [Bushnell and Fröhlich 1983]. Note that if an abelian subgroup G_1 of G satisfies the conditions in the claim, then G_1/Z is a maximal totally isotropic subspace of G/Z under the pairing

$$(G/Z) \times (G/Z) \rightarrow \mathbb{C}^\times, \quad (gZ, g'Z) \mapsto \psi([g, g']).$$

Hence the construction follows from the proof of [Bushnell and Fröhlich 1983, Proposition 8.3.3]. \square

Next, we consider representations of a semidirect product of a Heisenberg group with a cyclic group. Let $A \subset \text{Aut}(G)$ be a cyclic subgroup of order $p^e + 1$ where $e = \frac{1}{2}(\log_p [G : Z])$. We assume:

- (3) The group A acts on Z trivially.
- (4) For any nontrivial element $a \in A$, the action of a on G/Z fixes only the unit element.

We consider the semidirect product $A \ltimes G$ by the action of A on G .

Lemma 1.3. *There is a unique irreducible representation ρ'_ψ of $A \ltimes G$ such that $\rho'_\psi|_G \simeq \rho_\psi$ and $\text{tr } \rho'_\psi(a) = -1$ for every nontrivial element $a \in A$.*

Proof. The claim is proved in the proof of Lemma 22.2 in [Bushnell and Henniart 2006] if Z is cyclic and ψ is a faithful character. In fact, the same proof works also in our case. \square

Corollary 1.4. *There exists a unique representation ρ'_ψ of $A \ltimes G$ such that*

$$\rho'_\psi|_Z \simeq \psi^{\oplus p^e} \quad \text{and} \quad \text{tr } \rho'_\psi(a) = -1$$

for every nontrivial element $a \in A$. Further, the representation $\rho'_\psi|_G$ is irreducible.

Proof. First we show the existence. We take the representation ρ'_ψ in Lemma 1.3. Then ρ'_ψ has a central character equal to ψ by Proposition 1.2. This shows the existence.

We show the uniqueness and the irreducibility of $\rho'_\psi|_G$. Assume that ρ'_ψ satisfies the condition in the claim. Take an irreducible subrepresentation ρ_ψ of $\rho'_\psi|_G$. Then ρ_ψ satisfies the condition of Proposition 1.2. Hence, $\dim \rho_\psi = p^e$. Then we have $\rho_\psi = \rho'_\psi|_G$ and $\rho'_\psi|_G$ is irreducible. Such ρ_ψ is unique by Lemma 1.3. \square

2. Galois representations

2A. Swan conductor. Let K be a nonarchimedean local field with residue field k . Let p be the characteristic of k . Let f be the extension degree of k over \mathbb{F}_p . We put $q = p^f$.

Let

$$\text{Art}_K : K^\times \xrightarrow{\sim} W_K^{\text{ab}}$$

be the Artin reciprocity map, which sends a uniformizer to a lift of the geometric Frobenius element.

Let τ be a finite dimensional irreducible continuous representation of W_K over \mathbb{C} . Let $\Psi : K \rightarrow \mathbb{C}^\times$ be a nontrivial additive character. Let $\varepsilon(\tau, s, \Psi)$ denote the Deligne–Langlands local constant of τ with respect to Ψ . We simply write $\varepsilon(\tau, \Psi)$ for $\varepsilon(\tau, \frac{1}{2}, \Psi)$.

We define an unramified character $\omega_s : K^\times \rightarrow \mathbb{C}^\times$ by $\omega_s(\varpi) = q^{-s}$ for $s \in \mathbb{R}$, where ϖ is a uniformizer of K . We recall that

$$(2-1) \quad \varepsilon(\tau, s, \Psi) = \varepsilon(\tau \otimes \omega_s, 0, \Psi)$$

(see [Tate 1979, (3.6.4)]).

Let $\psi_0 \in \mathbb{F}_p^\vee$ by $\psi_0(1) = e^{2\pi\sqrt{-1}/p}$. We take an additive character $\psi_K : K \rightarrow \mathbb{C}^\times$ such that $\psi_K(x) = \psi_0(\text{Tr}_{k/\mathbb{F}_p}(\bar{x}))$ for $x \in \mathcal{O}_K$. By [Bushnell and Henniart 2006, Proposition 29.4], there exists an integer $\text{sw}(\tau)$ such that

$$\varepsilon(\tau, s, \psi_K) = q^{-\text{sw}(\tau)s} \varepsilon(\tau, 0, \psi_K).$$

We put $\text{Sw}(\tau) = \max\{\text{sw}(\tau), 0\}$, which we call the Swan conductor of τ .

2B. Construction. We construct a group Q which acts on a curve C over an algebraic closure of k . By using this action of Q and Frobenius action, we construct a representation of a semidirect product $Q \rtimes \mathbb{Z}$ in étale cohomology of C . Then we use the representation of $Q \rtimes \mathbb{Z}$ to construct a representation of a Weil group.

We fix an algebraic closure K^{ac} of K . Let k^{ac} be the residue field of K^{ac} . Let n be a positive integer. We write $n = p^e n'$ with $(p, n') = 1$. Throughout this paper, we assume that $e \geq 1$. Let

$$Q = \{(a, b, c) \mid a \in \mu_{p^e+1}(k^{\text{ac}}), b, c \in k^{\text{ac}}, b^{p^{2e}} + b = 0, c^p - c + b^{p^e+1} = 0\}$$

be the group whose multiplication is given by

$$(a_1, b_1, c_1) \cdot (a_2, b_2, c_2) = \left(a_1 a_2, b_1 + a_1 b_2, c_1 + c_2 + \sum_{i=0}^{e-1} (a_1 b_1^{p^e} b_2)^{p^i} \right).$$

Remark 2.1. The construction of the group Q has its origin in a study of the automorphism of a curve C defined below. We can check that the above multiplication gives a group structure on Q directly, but it's also possible to show this by

checking that the inclusion from Q to the automorphism group of C defined below is compatible with the multiplications.

Note that $|Q| = p^{2e+1}(p^e + 1)$. Let $Q \rtimes \mathbb{Z}$ be a semidirect product, where $m \in \mathbb{Z}$ acts on Q by $(a, b, c) \mapsto (a^{p^{-m}}, b^{p^{-m}}, c^{p^{-m}})$. We put

$$(2-2) \quad \text{Fr}(m) = ((1, 0, 0), m) \in Q \rtimes \mathbb{Z} \quad \text{for } m \in \mathbb{Z}.$$

Let C be the smooth affine curve over k^{ac} defined by

$$x^p - x = y^{p^e+1} \quad \text{in } \mathbb{A}_{k^{\text{ac}}}^2.$$

We define a right action of $Q \rtimes \mathbb{Z}$ on C by

$$(x, y)((a, b, c), 0) = \left(x + \sum_{i=0}^{e-1} (by)^{p^i} + c, a(y + b^{p^e}) \right), \quad (x, y) \text{Fr}(1) = (x^p, y^p).$$

We consider the morphisms

$$h : \mathbb{A}_{k^{\text{ac}}}^1 \rightarrow \mathbb{A}_{k^{\text{ac}}}^1, \quad x \mapsto x^p - x, \quad \pi : \mathbb{A}_{k^{\text{ac}}}^1 \rightarrow \mathbb{A}_{k^{\text{ac}}}^1, \quad y \mapsto y^{p^e+1}.$$

Then we have the fiber product

$$\begin{array}{ccc} C & \xrightarrow{h'} & \mathbb{A}_{k^{\text{ac}}}^1 \\ \pi' \downarrow & \square & \downarrow \pi \\ \mathbb{A}_{k^{\text{ac}}}^1 & \xrightarrow{h} & \mathbb{A}_{k^{\text{ac}}}^1 \end{array}$$

where π' and h' are the natural projections to the first and second coordinates respectively. Let $g = ((a, b, c), m) \in Q \rtimes \mathbb{Z}$. We consider the morphism

$$g_0 : \mathbb{A}_{k^{\text{ac}}}^1 \rightarrow \mathbb{A}_{k^{\text{ac}}}^1, \quad y \mapsto (a(y + b^{p^e}))^{p^m}.$$

Let ℓ be a prime number different from p . Then we have a natural isomorphism

$$c_g : g_0^* h'_* \bar{\mathbb{Q}}_\ell \xrightarrow{\sim} h'_* g^* \bar{\mathbb{Q}}_\ell \xrightarrow{\sim} h'_* \bar{\mathbb{Q}}_\ell.$$

We take an isomorphism $\iota : \bar{\mathbb{Q}}_\ell \simeq \mathbb{C}$. We sometimes view a character over \mathbb{C} as a character over $\bar{\mathbb{Q}}_\ell$ by ι . Let $\psi \in \mathbb{F}_p^\vee$. We write \mathcal{L}_ψ for the Artin–Schreier $\bar{\mathbb{Q}}_\ell$ -sheaf on $\mathbb{A}_{k^{\text{ac}}}^1$ associated to ψ , which is equal to $\mathfrak{F}(\psi)$ in the notation of [Deligne 1977, Sommes trig. 1.8(i)]. Then we have a decomposition $h_* \bar{\mathbb{Q}}_\ell = \bigoplus_{\psi \in \mathbb{F}_p^\vee} \mathcal{L}_\psi$. This decomposition gives canonical isomorphisms

$$(2-3) \quad h'_* \bar{\mathbb{Q}}_\ell \cong \pi^* h_* \bar{\mathbb{Q}}_\ell \cong \bigoplus_{\psi \in \mathbb{F}_p^\vee} \pi^* \mathcal{L}_\psi.$$

The isomorphisms c_g and (2-3) induce $c_{g,\psi} : g_0^* \pi^* \mathcal{L}_\psi \rightarrow \pi^* \mathcal{L}_\psi$. We define a left action of $Q \times \mathbb{Z}$ on $H_c^1(\mathbb{A}_{k^{\text{ac}}}^1, \pi^* \mathcal{L}_\psi)$ by

$$H_c^1(\mathbb{A}_{k^{\text{ac}}}^1, \pi^* \mathcal{L}_\psi) \xrightarrow{g_0^*} H_c^1(\mathbb{A}_{k^{\text{ac}}}^1, g_0^* \pi^* \mathcal{L}_\psi) \xrightarrow{c_{g,\psi}} H_c^1(\mathbb{A}_{k^{\text{ac}}}^1, \pi^* \mathcal{L}_\psi) \quad \text{for } g \in Q \times \mathbb{Z}.$$

Let τ_ψ be the representation of $Q \times \mathbb{Z}$ over \mathbb{C} defined by $H_c^1(\mathbb{A}_{k^{\text{ac}}}^1, \pi^* \mathcal{L}_\psi)$ and ι . For $\theta \in \mu_{p^e+1}(k^{\text{ac}})^\vee$, let \mathcal{K}_θ be the smooth Kummer $\overline{\mathbb{Q}}_\ell$ -sheaf on $\mathbb{G}_{m,k^{\text{ac}}}$ associated to θ . We view $\mu_{p^e+1}(k^{\text{ac}}) \times \mathbb{F}_p$ as a subgroup of Q by $(a, c) \mapsto (a, 0, c)$.

Lemma 2.2. *We have a natural isomorphism*

$$H_c^1(\mathbb{A}_{k^{\text{ac}}}^1, \pi^* \mathcal{L}_\psi) \simeq \bigoplus_{\theta \in \mu_{p^e+1}(k^{\text{ac}})^\vee \setminus \{1\}} H_c^1(\mathbb{G}_{m,k^{\text{ac}}}, \mathcal{L}_\psi \otimes \mathcal{K}_\theta),$$

which is compatible with the actions of $\mu_{p^e+1}(k^{\text{ac}}) \times \mathbb{F}_p$ where

$$(a, c) \in \mu_{p^e+1}(k^{\text{ac}}) \times \mathbb{F}_p$$

acts on $H_c^1(\mathbb{G}_{m,k^{\text{ac}}}, \mathcal{L}_\psi \otimes \mathcal{K}_\theta)$ by $\theta(a) \psi(c)$. Further, we have

$$\dim H_c^1(\mathbb{G}_{m,k^{\text{ac}}}, \mathcal{L}_\psi \otimes \mathcal{K}_\theta) = 1$$

for any $\theta \in \mu_{p^e+1}(k^{\text{ac}})^\vee \setminus \{1\}$.

Proof. By the projection formula, we have natural isomorphisms

$$\pi_* \pi^* \mathcal{L}_\psi \simeq \pi_*(\pi^* \mathcal{L}_\psi \otimes \overline{\mathbb{Q}}_\ell) \simeq \mathcal{L}_\psi \otimes \pi_* \overline{\mathbb{Q}}_\ell \quad \text{on } \mathbb{A}_{k^{\text{ac}}}^1.$$

Further, we have

$$\pi_* \overline{\mathbb{Q}}_\ell \simeq \bigoplus_{\theta \in \mu_{p^e+1}(k^{\text{ac}})^\vee} \mathcal{K}_\theta \quad \text{on } \mathbb{G}_{m,k^{\text{ac}}},$$

since π is a finite étale $\mu_{p^e+1}(k^{\text{ac}})$ -covering over $\mathbb{G}_{m,k^{\text{ac}}}$. Therefore, we have

$$(2-4) \quad \pi_* \pi^* \mathcal{L}_\psi \simeq \mathcal{L}_\psi \otimes \pi_* \overline{\mathbb{Q}}_\ell \simeq \bigoplus_{\theta \in \mu_{p^e+1}(k^{\text{ac}})^\vee} \mathcal{L}_\psi \otimes \mathcal{K}_\theta$$

on $\mathbb{G}_{m,k^{\text{ac}}}$. Let $\{0\}$ denote the origin of $\mathbb{A}_{k^{\text{ac}}}^1$. Let $i : \{0\} \rightarrow \mathbb{A}_{k^{\text{ac}}}^1$ and $j : \mathbb{G}_{m,k^{\text{ac}}} \rightarrow \mathbb{A}_{k^{\text{ac}}}^1$ be the natural immersions. From the exact sequence

$$0 \rightarrow j_* j^* \pi^* \mathcal{L}_\psi \rightarrow \pi^* \mathcal{L}_\psi \rightarrow i_* i^* \pi^* \mathcal{L}_\psi \rightarrow 0,$$

we have the exact sequence

$$(2-5) \quad 0 \rightarrow H^0(\{0\}, i^* \pi^* \mathcal{L}_\psi) \rightarrow H_c^1(\mathbb{G}_{m,k^{\text{ac}}}, \pi^* \mathcal{L}_\psi) \rightarrow H_c^1(\mathbb{A}_{k^{\text{ac}}}^1, \pi^* \mathcal{L}_\psi) \rightarrow 0,$$

since

$$H_c^0(\mathbb{A}_{k^{\text{ac}}}^1, \pi^* \mathcal{L}_\psi) = 0 \quad \text{and} \quad H^1(\{0\}, i^* \pi^* \mathcal{L}_\psi) = 0.$$

Note that $H^0(\{0\}, i^* \pi^* \mathcal{L}_\psi) \simeq \psi$. By (2-4), we have isomorphisms

$$(2-6) \quad \begin{aligned} H_c^1(\mathbb{G}_{m,k^{\text{ac}}}, \pi^* \mathcal{L}_\psi) &\simeq H_c^1(\mathbb{G}_{m,k^{\text{ac}}}, \pi_* \pi^* \mathcal{L}_\psi) \\ &\simeq \bigoplus_{\theta \in \mu_{p^e+1}(k^{\text{ac}})^\vee} H_c^1(\mathbb{G}_{m,k^{\text{ac}}}, \mathcal{L}_\psi \otimes \mathcal{K}_\theta). \end{aligned}$$

We know that

$$(2-7) \quad \dim H_c^1(\mathbb{G}_{m,k^{\text{ac}}}, \mathcal{L}_\psi \otimes \mathcal{K}_\theta) = 1$$

for any $\theta \in \mu_{p^e+1}(k^{\text{ac}})^\vee$ by the proof of [Imai and Tsushima 2017, Lemma 7.1] (see [Imai and Tsushima 2023, (2.3)]). Since the composition of

$$H^0(\{0\}, i^* \pi^* \mathcal{L}_\psi) \rightarrow H_c^1(\mathbb{G}_{m,k^{\text{ac}}}, \pi^* \mathcal{L}_\psi)$$

and (2-6) is compatible with the actions of $\mu_{p^e+1}(k^{\text{ac}}) \times \mathbb{F}_p$, it factors through an isomorphism $H^0(\{0\}, i^* \pi^* \mathcal{L}_\psi) \simeq H_c^1(\mathbb{G}_{m,k^{\text{ac}}}, \mathcal{L}_\psi)$ by (2-7). Then the claim follows from (2-5), (2-6) and (2-7). \square

Let $\varrho : \mu_2(k) \hookrightarrow \mathbb{C}^\times$ be the nontrivial group homomorphism if $p \neq 2$. We define a character $\theta_0 \in \mu_{p^e+1}(k^{\text{ac}})^\vee$ by

$$(2-8) \quad \theta_0(a) = \begin{cases} \varrho(a^{(p^e+1)/2}) & \text{if } p \neq 2, \\ 1 & \text{if } p = 2 \end{cases}$$

for $a \in \mu_{p^e+1}(k^{\text{ac}})$. For an integer m and a positive odd integer m' , let $\left(\frac{m}{m'}\right)$ denote the Jacobi symbol. For an odd prime p , we set

$$\epsilon(p) = \begin{cases} 1 & \text{if } p \equiv 1 \pmod{4}, \\ \sqrt{-1} & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

We have $\epsilon(p)^2 = \left(\frac{-1}{p}\right)$. We define a representation τ_n of $Q \rtimes \mathbb{Z}$ as the twist of τ_{ψ_0} by the character

$$(2-9) \quad \begin{aligned} Q \rtimes \mathbb{Z} &\rightarrow \mathbb{C}^\times, \\ ((a, b, c), m) &\mapsto \begin{cases} \theta_0(a)^n \left((-\epsilon(p) \left(\frac{-2n'}{p}\right))^n p^{-\frac{1}{2}} \right)^m & \text{if } p \neq 2, \\ ((-1)^{\frac{n(n-2)}{8}} p^{-\frac{1}{2}})^m & \text{if } p = 2. \end{cases} \end{aligned}$$

The value of this character is related to a quadratic Gauss sum. A geometric origin of this character is given in (A-3). Let $(\zeta, \chi, c) \in \mu_{q-1}(K) \times (k^\times)^\vee \times \mathbb{C}^\times$. We take a uniformizer ϖ of K . We choose an element $\varphi'_\zeta \in K^{\text{ac}}$ such that $\varphi_\zeta^{m'} = \zeta \varpi$ and set $E_\zeta = K(\varphi'_\zeta)$. We choose elements $\alpha_\zeta, \beta_\zeta, \gamma_\zeta \in K^{\text{ac}}$ such that

$$(2-10) \quad \alpha_\zeta^{p^e+1} = -\varphi'_\zeta, \quad \beta_\zeta^{p^2e} + \beta_\zeta = -\alpha_\zeta^{-1}, \quad \gamma_\zeta^p - \gamma_\zeta = \beta_\zeta^{p^e+1}.$$

For $\sigma \in W_{E_\zeta}$, we set

$$(2-11) \quad \begin{aligned} a_\sigma &= \sigma(\alpha_\zeta)/(\alpha_\zeta), & b_\sigma &= a_\sigma \sigma(\beta_\zeta) - \beta_\zeta, \\ c_\sigma &= \sigma(\gamma_\zeta) - \gamma_\zeta + \sum_{i=0}^{e-1} (b_\sigma^{p^e} (\beta_\zeta + b_\sigma))^{p^i}. \end{aligned}$$

Then we have $a_\sigma, b_\sigma, c_\sigma \in \mathcal{O}_{K^{\text{ac}}}$. For $\sigma \in W_{E_\zeta}$, we put $n_\sigma = v_{E_\zeta}(\text{Art}_{E_\zeta}^{-1}(\sigma))$. We have the homomorphism

$$(2-12) \quad \Theta_\zeta : W_{E_\zeta} \rightarrow Q \times \mathbb{Z}, \quad \sigma \mapsto ((\bar{a}_\sigma, \bar{b}_\sigma, \bar{c}_\sigma), f n_\sigma).$$

Lemma 2.3. *The image of the homomorphism Θ_ζ is $Q \times (f\mathbb{Z})$.*

Proof. It suffices to show that the image of $I_{E_\zeta} \subset W_{E_\zeta}$ under Θ_ζ is equal to $Q \subset Q \times \mathbb{Z}$, since the homomorphism $W_{E_\zeta} \rightarrow f\mathbb{Z}$, $\sigma \mapsto f n_\sigma$ is surjective. We put $N_\zeta = E_\zeta(\alpha_\zeta, \beta_\zeta, \gamma_\zeta)$. Then the kernel of Θ_ζ is equal to I_{N_ζ} by the definition. Hence we have an injection $I_{E_\zeta}/I_{N_\zeta} \hookrightarrow Q$. This injection is actually a bijection, since N_ζ is a totally ramified extension over E_ζ of degree $p^{2e+1}(p^e + 1)$, which equals to $|Q|$. Therefore, we obtain the claim. \square

We write $\tau_{n,\zeta}$ for the representation of W_{E_ζ} given by Θ_ζ and τ_n . Recall that c is an element of \mathbb{C}^\times . Let $\phi_c : W_{E_\zeta} \rightarrow \mathbb{C}^\times$ be the character defined by $\phi_c(\sigma) = c^{n_\sigma}$. We have the isomorphism $\varphi_\zeta^{\mathbb{Z}} \times \mathcal{O}_{E_\zeta}^\times \simeq E_\zeta^\times$ given by the multiplication. Let $\text{Frob}_p : k^\times \rightarrow k^\times$ be the inverse of the p -th power map. We consider the following composition:

$$\lambda_\zeta : W_{E_\zeta}^{\text{ab}} \simeq E_\zeta^\times \simeq \varphi_\zeta^{\mathbb{Z}} \times \mathcal{O}_{E_\zeta}^\times \xrightarrow{\text{pr}_2} \mathcal{O}_{E_\zeta}^\times \xrightarrow{\text{can.}} k^\times \xrightarrow{\text{Frob}_p^e} k^\times.$$

We put

$$(2-13) \quad \tau_{n,\zeta,\chi,c} = \tau_{n,\zeta} \otimes (\chi \circ \lambda_\zeta) \otimes \phi_c \quad \text{and} \quad \tau_{\zeta,\chi,c} = \text{Ind}_{E_\zeta/K} \tau_{n,\zeta,\chi,c}.$$

We will see that $\tau_{\zeta,\chi,c}$ is an irreducible representation of Swan conductor 1 in [Proposition 10.8](#). This Galois representation $\tau_{\zeta,\chi,c}$ is our main object in this paper. We will study several invariants associated to this, for example, its determinant and epsilon factor.

2C. Characterization. We put

$$Q_0 = \{(1, b, c) \in Q\}, \quad F = \{(1, 0, c) \in Q \mid c \in \mathbb{F}_p\}.$$

We identify \mathbb{F}_p with F by $c \mapsto (1, 0, c)$.

Lemma 2.4. *For any $g = (1, b, c) \in Q_0$ with $b \neq 0$, the map $Q_0 \rightarrow F$, $g' \mapsto [g, g']$ is surjective.*

Proof. For $(1, b_1, c_1), (1, b_2, c_2) \in Q_0$, we have

$$[(1, b_1, c_1), (1, b_2, c_2)] = \left(1, 0, \sum_{i=0}^{e-1} (b_1^{p^e} b_2 - b_1 b_2^{p^e})^{p^i}\right).$$

If $b_1 \neq 0$, then

$$\{b \in k^{\text{ac}} \mid b^{p^{2e}} + b = 0\} \rightarrow \mathbb{F}_{p^e}, \quad b_2 \rightarrow b_1^{p^e} b_2 - b_1 b_2^{p^e}$$

is surjective. The claim follows from the surjectivity of $\text{Tr}_{\mathbb{F}_{p^e}/\mathbb{F}_p}$. \square

By this lemma, we can apply the results from [Section 1](#) to our situation with $G = Q_0$, $Z = F$ and $A = \mu_{p^e+1}(k^{\text{ac}})$, where the action of $\mu_{p^e+1}(k^{\text{ac}})$ on Q_0 is given by the embedding

$$\mu_{p^e+1}(k^{\text{ac}}) \rightarrow Q, \quad a \mapsto (a, 0, 0)$$

and the conjugation. Let τ^0 denote the unique representation of Q characterized by

$$(2-14) \quad \tau^0|_F \simeq \psi_0^{\oplus p^e}, \quad \text{Tr } \tau^0((a, 0, 0)) = -1$$

for $a \in \mu_{p^e+1}(k^{\text{ac}}) \setminus \{1\}$ (see [Corollary 1.4](#)).

We have a decomposition

$$(2-15) \quad \tau^0 = \bigoplus_{\theta \in \mu_{p^e+1}(k^{\text{ac}})^\vee \setminus \{1\}} L_\theta$$

such that $a \in \mu_{p^e+1}(k^{\text{ac}})$ acts on L_θ by $\theta(a)$, since the both sides of (2-15) have the same character as representations of $\mu_{p^e+1}(k^{\text{ac}})$. For a positive integer m dividing p^e+1 , we consider $\mu_m(k^{\text{ac}})^\vee$ as a subset of $\mu_{p^e+1}(k^{\text{ac}})^\vee$ by the dual of the surjection

$$\mu_{p^e+1}(k^{\text{ac}}) \rightarrow \mu_m(k^{\text{ac}}), \quad x \rightarrow x^{(p^e+1)/m}.$$

We simply write Q for the subgroup $Q \times \{0\} \subset Q \rtimes \mathbb{Z}$.

Lemma 2.5. *We have $\tau_{\psi_0}|_Q \simeq \tau^0$.*

Proof. The representation $\tau_{\psi_0}|_Q$ satisfies the characterization (2-14) by [Lemma 2.2](#). Hence $\tau_{\psi_0}|_Q$ is isomorphic to τ^0 . \square

Corollary 2.6. *The representation $\tau_{\psi_0}|_{Q_0}$ is irreducible.*

Proof. This follows from [Corollary 1.4](#), equation (2-14) and [Lemma 2.5](#). \square

For any odd prime p , we have

$$(2-16) \quad \sum_{x \in \mathbb{F}_p^\times} \psi_0(x^2) = \sum_{x \in \mathbb{F}_p^\times} \left(\frac{x}{p}\right) \psi_0(x) = \epsilon(p) \sqrt{p}$$

by Gauss.

Lemma 2.7. *We have*

$$\mathrm{Tr} \tau_{\psi_0}(\mathrm{Fr}(1)) = \begin{cases} -\epsilon(p)\sqrt{p} & \text{if } p \neq 2, \\ 0 & \text{if } p = 2. \end{cases}$$

Proof. By the Lefschetz trace formula, we have

$$\sum_{x \in \mathbb{A}^1(\mathbb{F}_p)} \mathrm{Tr}(\mathrm{Fr}_p, (\pi^* \mathcal{L}_{\psi_0})_x) = \sum_{i=0}^2 (-1)^i \mathrm{Tr}(\mathrm{Fr}_p, H_c^i(\mathbb{A}_{k^{\mathrm{ac}}}^1, \pi^* \mathcal{L}_{\psi})),$$

where Fr_p is the geometric p -th power Frobenius morphism. Since $H_c^i(\mathbb{A}_{k^{\mathrm{ac}}}^1, \pi^* \mathcal{L}_{\psi})$ vanishes for $i = 0, 2$, we have

$$\begin{aligned} \mathrm{Tr} \tau_{\psi_0}(\mathrm{Fr}(1)) &= - \sum_{x \in \mathbb{A}^1(\mathbb{F}_p)} \mathrm{Tr}(\mathrm{Fr}_p, (\pi^* \mathcal{L}_{\psi_0})_x) \\ &= - \sum_{x \in \mathbb{F}_p} \psi_0(x^{p^e+1}) = - \sum_{x \in \mathbb{F}_p} \psi_0(x^2) = \begin{cases} -\epsilon(p)\sqrt{p} & \text{if } p \neq 2, \\ 0 & \text{if } p = 2, \end{cases} \end{aligned}$$

where we use (2-16) in the last equality. \square

We assume $p = 2$ in this paragraph. We take $b_0 \in \mathbb{F}_{2^{2e}}$ such that $\mathrm{Tr}_{\mathbb{F}_{2^{2e}}/\mathbb{F}_2}(b_0) = 1$. Further, we put

$$(2-17) \quad c_0 = b_0^{2^e} + \sum_{0 \leq i < j \leq e-1} b_0^{2^{e+i}+2^j}.$$

Then we have

$$\begin{aligned} (2-18) \quad c_0^2 - c_0 &= b_0^{2^{e+1}} + b_0^{2^e} + \sum_{0 \leq i < j \leq e-1} b_0^{2^{e+i+1}+2^{j+1}} + \sum_{0 \leq i < j \leq e-1} b_0^{2^{e+i}+2^{j+1}} \\ &= b_0^{2^{e+1}} + b_0^{2^e} + \sum_{i=0}^{e-2} b_0^{2^{e+i+1}+2^e} + \sum_{j=1}^{e-1} b_0^{2^e+2^j} \\ &= b_0^{2^{e+1}} + b_0^{2^e} + b_0^{2^e}(1 + b_0 + b_0^{2^e}) = b_0^{2^e+1}, \end{aligned}$$

where we use $\mathrm{Tr}_{\mathbb{F}_{2^{2e}}/\mathbb{F}_2}(b_0) = 1$ at the third equality. We put

$$\mathbf{g} = ((1, b_0, c_0), -1) \in Q \times \mathbb{Z}.$$

Lemma 2.8. *We assume that $p = 2$. Then we have $\mathrm{Tr} \tau_{\psi_0}(\mathbf{g}^{-1}) = -2$.*

Proof. We note that

$$(2-19) \quad \mathbf{g}^{-1} = \mathrm{Fr}(1) \left(\left(1, b_0, c_0 + \sum_{i=0}^{e-1} (b_0^{2^e+1})^{2^i} \right), 0 \right).$$

For $y \in k^{\text{ac}}$ satisfying $y^2 + b_0^{2^e} = y$, we take $x_y \in k^{\text{ac}}$ such that $x_y^2 - x_y = y^{2^e+1}$. We take $y_0 \in k^{\text{ac}}$ such that $y_0^2 + b_0^{2^e} = y_0$. Then, by the Lefschetz trace formula and (2-19), we have

$$\begin{aligned} \text{Tr } \tau_{\psi_0}(\mathbf{g}^{-1}) &= - \sum_{y^2 + b_0^{2^e} = y} \text{Tr}(\mathbf{g}^{-1}, (\pi^* \mathcal{L}_{\psi_0})_y) \\ &= - \sum_{y^2 + b_0^{2^e} = y} \psi_0 \left(x_y^2 - x_y + \sum_{i=0}^{e-1} (b_0 y^2)^{2^i} + c_0 + \sum_{i=0}^{e-1} (b_0^{2^e+1})^{2^i} \right) \\ &= - \sum_{z \in \mathbb{F}_2} \psi_0 \left((y_0 + z)^{2^e+1} + \sum_{i=0}^{e-1} (b_0 (y_0 + z))^{2^i} + c_0 \right) = -2, \end{aligned}$$

where we change a variable by $y = y_0 + z$ at the second equality, and use

$$\begin{aligned} y_0^{2^e+1} + \sum_{i=0}^{e-1} (b_0 y_0)^{2^i} &= y_0 \left(y_0 + \sum_{i=0}^{e-1} b_0^{2^e+i} \right) + \sum_{i=0}^{e-1} b_0^{2^i} \left(y_0 + \sum_{j=0}^{i-1} b_0^{2^e+j} \right) = c_0, \\ y_0^{2^e} + y_0 + \sum_{i=0}^{e-1} b_0^{2^i} &= \sum_{i=0}^{e-1} (y_0^2 + y_0)^{2^i} + \sum_{i=0}^{e-1} b_0^{2^i} = \text{Tr}_{\mathbb{F}_{2^{2^e}}/\mathbb{F}_2}(b_0) = 1 \end{aligned}$$

at the last equality. □

Proposition 2.9. *The representation τ_{ψ_0} is characterized by $\tau_{\psi_0}|_Q \simeq \tau^0$ and*

$$\begin{cases} \text{Tr } \tau_{\psi_0}(\text{Fr}(1)) = -\epsilon(p)\sqrt{p} & \text{if } p \neq 2, \\ \text{Tr } \tau_{\psi_0}(\mathbf{g}^{-1}) = -2 & \text{if } p = 2. \end{cases}$$

In particular, τ_{ψ_0} does not depend on the choice of ℓ and ι .

Proof. This follows from Lemmas 2.5, 2.7 and 2.8. □

3. Representations of general linear algebraic groups

3A. Simple supercuspidal representation. Let π be an irreducible supercuspidal representation of $\text{GL}_n(K)$ over \mathbb{C} . Let $\varepsilon(\pi, s, \Psi)$ denote the Godement–Jacquet local constant of π with respect to the nontrivial character $\Psi : K \rightarrow \mathbb{C}^\times$. We simply write $\varepsilon(\pi, \Psi)$ for $\varepsilon(\pi, \frac{1}{2}, \Psi)$. By [Godement and Jacquet 1972, Theorem 3.3(4)], there exists an integer $\text{sw}(\pi)$ such that

$$\varepsilon(\pi, s, \psi_K) = q^{-\text{sw}(\pi)s} \varepsilon(\pi, 0, \psi_K).$$

We put $\text{Sw}(\pi) = \max\{\text{sw}(\pi), 0\}$, which we call the Swan conductor of π .

Definition 3.1. An irreducible supercuspidal representation π of $\text{GL}_n(K)$ over \mathbb{C} is called simple supercuspidal if $\text{Sw}(\pi) = 1$.

This definition is equivalent to [Imai and Tsushima 2018, Definition 1.1] by [Imai and Tsushima 2018, Proposition 1.3].

3B. Construction. In the following, we construct a smooth representation $\pi_{\zeta, \chi, c}$ of $\mathrm{GL}_n(K)$ for each triple $(\zeta, \chi, c) \in \mu_{q-1}(K) \times (k^\times)^\vee \times \mathbb{C}^\times$.

Let $B \subset M_n(k)$ be the subring consisting of upper triangular matrices. Let $\mathfrak{J} \subset M_n(\mathcal{O}_K)$ be the inverse image of B under the reduction map $M_n(\mathcal{O}_K) \rightarrow M_n(k)$. Then \mathfrak{J} is a hereditary \mathcal{O}_K -order (see [Bushnell and Kutzko 1993, (1.1)]). Let \mathfrak{P} denote the Jacobson radical of the order \mathfrak{J} . We put $U_{\mathfrak{J}}^1 = 1 + \mathfrak{P} \subset \mathrm{GL}_n(\mathcal{O}_K)$. We set

$$\varphi_\zeta = \begin{pmatrix} \mathbf{0} & I_{n-1} \\ \zeta \varpi & \mathbf{0} \end{pmatrix} \in M_n(K) \quad \text{and} \quad L_\zeta = K(\varphi_\zeta).$$

Then, L_ζ is a totally ramified extension of K of degree n .

We put $\varphi_{\zeta, n} = n' \varphi_\zeta$ and

$$\epsilon_0 = \begin{cases} \frac{1}{2}(n' + 1) & \text{if } p^e = 2, \\ 0 & \text{if } p^e \neq 2. \end{cases}$$

We define a character $\Lambda_{\zeta, \chi, c} : L_\zeta^\times U_{\mathfrak{J}}^1 \rightarrow \mathbb{C}^\times$ by

$$\begin{aligned} \Lambda_{\zeta, \chi, c}(\varphi_\zeta) &= (-1)^{n-1+\epsilon_0 f} c, & \Lambda_{\zeta, \chi, c}(x) &= \chi(\bar{x}) \quad \text{for } x \in \mathcal{O}_K^\times, \\ \Lambda_{\zeta, \chi, c}(x) &= (\psi_K \circ \mathrm{tr})(\varphi_{\zeta, n}^{-1}(x-1)) \quad \text{for } x \in U_{\mathfrak{J}}^1, \end{aligned}$$

where tr means the trace as an element of $M_n(K)$. We put

$$\pi_{\zeta, \chi, c} = \mathrm{c}\text{-Ind}_{L_\zeta^\times U_{\mathfrak{J}}^1}^{\mathrm{GL}_n(K)} \Lambda_{\zeta, \chi, c}.$$

Then, $\pi_{\zeta, \chi, c}$ is a simple supercuspidal representation of $\mathrm{GL}_n(K)$, and every simple supercuspidal representation is isomorphic to $\pi_{\zeta, \chi, c}$ for a uniquely determined $(\zeta, \chi, c) \in \mu_{q-1}(K) \times (k^\times)^\vee \times \mathbb{C}^\times$ by [Imai and Tsushima 2018, Proposition 1.3]. The representation $\pi_{\zeta, \chi, c}$ contains the m -simple stratum $[\mathfrak{J}, 1, 0, \varphi_{\zeta, n}^{-1}]$ in the sense of [Bushnell and Henniart 2014, Section 2.1].

Proposition 3.2. $\varepsilon(\pi_{\zeta, \chi, c}, \psi_K) = (-1)^{n-1+\epsilon_0 f} \chi(n') c.$

Proof. This follows from [Bushnell and Henniart 1999, Section 6.1, Lemma 2 and Section 6.3, Proposition 1]. \square

4. Local Langlands correspondence

Our main theorem is the following.

Theorem 4.1. *The representations $\pi_{\zeta, \chi, c}$ and $\tau_{\zeta, \chi, c}$ correspond via the local Langlands correspondence.*

To prove this theorem, we recall a characterization of the local Langlands correspondence for epipelagic representations due to Bushnell–Henniart. Recall that $\Psi : K \rightarrow \mathbb{C}^\times$ is a nontrivial character. The following lemma is a special case of [Deligne and Henniart 1981, Proposition 4.13].

Lemma 4.2 [Bushnell and Henniart 2014, Lemma 2.3]. *Let τ be an irreducible smooth representation of W_K such that $\text{sw}(\tau) \geq 1$. Then, there exists $\gamma_{\tau, \Psi} \in K^\times$ such that*

$$\varepsilon(\chi \otimes \tau, s, \Psi) = \chi(\gamma_{\tau, \Psi})^{-1} \varepsilon(\tau, s, \Psi)$$

for any tamely ramified character χ of W_K . This property determines the coset $\gamma_{\tau, \Psi} U_K^1$ uniquely.

Definition 4.3. Let τ be an irreducible smooth representation of W_K such that $\text{sw}(\tau) \geq 1$. We take $\gamma_{\tau, \Psi}$ as in Lemma 4.2. We put

$$\text{rsw}(\tau, \Psi) = \gamma_{\tau, \Psi}^{-1} \in K^\times / U_K^1,$$

which we call the refined Swan conductor of τ with respect to Ψ .

Remark 4.4. By (2-1), we have $v_K(\text{rsw}(\tau, \psi_K)) = \text{Sw}(\tau)$ in Definition 4.3.

Lemma 4.5. *Let π be an irreducible supercuspidal representation of $\text{GL}_n(K)$ such that $\text{sw}(\pi) \geq 1$.*

(1) *There exists $\gamma_{\pi, \Psi} \in K^\times$ such that*

$$\varepsilon(\chi \otimes \pi, s, \Psi) = \chi(\gamma_{\pi, \Psi})^{-1} \varepsilon(\pi, s, \Psi)$$

for any tamely ramified character χ of K^\times . This property determines the coset $\gamma_{\pi, \Psi} U_K^1$ uniquely.

(2) *Let $[\mathfrak{A}, m, 0, \alpha]$ be a simple stratum contained in π . Then we have $\gamma_{\pi, \Psi} \equiv \det \alpha \pmod{U_K^1}$.*

Proof. The first statement is [Bushnell and Henniart 1999, Theorem 1.4(i)]. The second statement follows from [Bushnell and Henniart 1999, Remark 1.4]. \square

Definition 4.6. Let π be an irreducible supercuspidal representation of $\text{GL}_n(K)$ such that $\text{sw}(\pi) \geq 1$. We take $\gamma_{\pi, \Psi}$ as in Lemma 4.5. Then we put

$$\text{rsw}(\pi, \Psi) = \gamma_{\pi, \Psi}^{-1} \in K^\times / U_K^1,$$

which we call the refined Swan conductor of π with respect to Ψ .

Remark 4.7. We have $v_K(\text{rsw}(\pi, \psi_K)) = \text{Sw}(\pi)$ in Definition 4.6.

For an irreducible supercuspidal representation π of $\text{GL}_n(K)$, let ω_π denote the central character of π .

Proposition 4.8 [Bushnell and Henniart 2014, Proposition 2.3]. *Let π be a simple supercuspidal representation of $\text{GL}_n(K)$. The Langlands parameter for π is characterized as the n -dimensional irreducible smooth representation τ of W_K satisfying*

$$\det \tau = \omega_\pi, \quad \text{rsw}(\tau, \psi_K) = \text{rsw}(\pi, \psi_K), \quad \varepsilon(\tau, \psi_K) = \varepsilon(\pi, \psi_K).$$

We will show that $\tau_{\xi, \chi, c}$ and $\pi_{\xi, \chi, c}$ satisfy the conditions of [Proposition 4.8](#) in [Propositions 8.6, 10.5, Lemma 10.7](#) and [Proposition 11.6](#).

5. General facts on epsilon factors

In this section, we recall some general facts on epsilon factors.

For a finite separable extension L over K , we put $\Psi_L = \Psi \circ \text{Tr}_{L/K}$ and let

$$\lambda(L/K, \Psi) = \frac{\varepsilon(\text{Ind}_{L/K} 1, s, \Psi)}{\varepsilon(1, s, \Psi_L)}$$

denote the Langlands constant which is independent of s , where 1 is the trivial representation of W_L (see [\[Bushnell and Henniart 2006, Section 30.4\]](#)).

Proposition 5.1. *Let τ be a finite dimensional smooth representation of W_K such that $\tau|_{P_K}$ is irreducible and nontrivial. Let L be a tamely ramified finite extension of K . Then we have*

$$\varepsilon(\tau|_{W_L}, \Psi_L) = \lambda(L/K, \Psi)^{-\dim \tau} \delta_{L/K}(\text{rsw}(\tau, \Psi)) \varepsilon(\tau, \Psi)^{[L:K]}.$$

Proof. This is proved by the same arguments as in [\[Bushnell and Henniart 2006, Proposition 48.3\]](#). \square

Proposition 5.2. *Let τ be a finite dimensional smooth representation of W_K such that $\tau|_{P_K}$ does not contain the trivial character.*

- (1) *If ϕ is a tamely ramified character of W_K , then $\text{rsw}(\tau \otimes \phi, \Psi) = \text{rsw}(\tau, \Psi)$.*
- (2) *Let L be a tamely ramified finite extension of K . Then we have*

$$\text{rsw}(\tau|_{W_L}, \Psi_L) = \text{rsw}(\tau, \Psi) \pmod{U_L^1}.$$

Proof. This is [\[Bushnell and Henniart 2006, Theorem 48.1\(2\), \(3\)\]](#). \square

For a nontrivial character ξ of K^\times , the level of ξ means the least integer $m \geq 0$ such that ξ is trivial on U_K^{m+1} .

Proposition 5.3. *Let ξ be a character of K^\times of level $m \geq 1$. Assume that $\gamma \in K^\times$ satisfies*

$$\xi(1+x) = \Psi(\gamma x) \quad \text{for } x \in \mathfrak{p}_K^{[m/2]+1}.$$

- (1) *We have $\text{rsw}(\xi, \Psi) = \gamma^{-1}$.*
- (2) *We have*

$$\varepsilon(\xi, \Psi) = q^{[(m+1)/2] - (m+1)/2} \sum_{y \in U_K^{[(m+1)/2]} / U_K^{[m/2]+1}} \xi(\gamma y)^{-1} \Psi(\gamma y).$$

Proof. Claim (1) follows from [\[Bushnell and Henniart 2006, Stability theorem 23.8\]](#). Claim (2) follows from [\[Bushnell and Henniart 2006, Section 23.5, Lemma 1, \(23.6.2\) and Proposition 23.6\]](#). \square

For a finite Galois extension L of K , let $\psi_{L/K}$ denote the Herbrand function of L/K and $\text{Gal}(L/K)_i$ denote the lower numbering i -th ramification subgroup of $\text{Gal}(L/K)$ for $i \geq 0$ (see [Serre 1968, Chapter IV]). We use the following lemmas to calculate the refined Swan conductor of a character of a Weil group.

Lemma 5.4. *Let m be a positive integer dividing f . Let h be a positive integer that is prime to p and less than $p^m v_K(p)/(p^m - 1)$. Let L be a Galois extension of K defined by $x^{p^m} - x = 1/\varpi^h$. Then we have*

$$\text{Gal}(L/K)_i = \begin{cases} \text{Gal}(L/K) & \text{if } i \leq h, \\ \{1\} & \text{if } i > h \end{cases}$$

and

$$\psi_{L/K}(v) = \begin{cases} v & \text{if } v \leq h, \\ p^m(v - h) + h & \text{if } v > h. \end{cases}$$

Proof. Take an integer l such that $lh \equiv 1 \pmod{p^m}$. Then we have

$$v_L\left(\frac{1}{x^l \varpi^{(lh-1)/p^m}}\right) = 1.$$

Hence, for $\sigma \in \text{Gal}(L/K)$ and $i \geq 0$, we have $\sigma \in \text{Gal}(L/K)_i$ if and only if

$$(5-1) \quad i + 1 \leq v_L\left(\sigma\left(\frac{1}{x^l \varpi^{(lh-1)/p^m}}\right) - \frac{1}{x^l \varpi^{(lh-1)/p^m}}\right) = v_L(\sigma(x)^l - x^l) + hl + 1.$$

The right-hand side of (5-1) is $h + 1$ if $\sigma \neq 1$. Hence the first claim follows. The second claim follows from the first claim. \square

Lemma 5.5. *Let L be a totally ramified finite abelian extension of K . Let $m \geq 1$.*

(1) *We have*

$$\begin{aligned} \text{Nr}_{L/K}(U_L^{\psi_{L/K}(m)}) &\subset U_K^m, & \text{Nr}_{L/K}(U_L^{\psi_{L/K}(m)+1}) &\subset U_K^{m+1}, \\ \text{Art}_K(U_K^m) &\subset \text{Gal}(L/K)_{\psi_{L/K}(m)}. \end{aligned}$$

(2) *We take $\alpha \in K$ and $\beta \in L$ such that $v_K(\alpha) = m$ and $v_L(\beta) = \psi_{L/K}(m)$. We put $P(z) = z^p - z$ for $z \in k$. Assume that*

$$\begin{array}{ccc} U_L^{\psi_{L/K}(m)} & \xrightarrow{\text{Nr}_{L/K}} & U_K^m \\ \downarrow P_{L,\beta} & & \downarrow P_{K,\alpha} \\ k & \xrightarrow{P} & k \end{array}$$

is commutative, where

$$\begin{aligned} P_{K,\alpha} : U_K^m &\rightarrow k, & 1 + \alpha x &\mapsto \bar{x}, \\ P_{L,\beta} : U_L^{\psi_{L/K}(m)} &\rightarrow k, & 1 + \beta x &\mapsto \bar{x}. \end{aligned}$$

Let ϖ_L be a uniformizer of L . Then we have

$$p_{L,\beta} \left(\frac{\text{Art}_K(1 + \alpha x)(\varpi_L)}{\varpi_L} \right) = \text{Tr}_{k/\mathbb{F}_p}(\bar{x})$$

for $x \in \mathcal{O}_K$.

Proof. The first claim follows from [Serre 1968, Chapter V, Section 3, Proposition 4 and Chapter XV, Section 2, Corollaire 3 of Théorème 1]. We note that our normalization of the Artin reciprocity map is inverse to that in [Serre 1968, Chapter XIII, Section 4]. Let $x \in \mathcal{O}_K$. By [Serre 1968, Chapter XV, Section 3, Proposition 4] and the construction of the isomorphism of [Serre 1968, Chapter XV, Section 2, Proposition 3], we have

$$p_{L,\beta} \left(\frac{\text{Art}_K(1 + \alpha x)(\varpi_L)}{\varpi_L} \right) = z_x^q - z_x,$$

where we take $z_x \in k^{\text{ac}}$ such that $z_x^p - z_x = \bar{x}$. Then we have the second claim, since

$$z_x^q - z_x = \text{Tr}_{k/\mathbb{F}_p}(z_x^p - z_x) = \text{Tr}_{k/\mathbb{F}_p}(\bar{x})$$

for such z_x . □

6. Stiefel–Whitney class and discriminant

6A. Stiefel–Whitney class. Let $R(W_K, \mathbb{R})$ be the Grothendieck group of finite-dimensional representations of W_K over \mathbb{R} with finite images. For $V \in R(W_K, \mathbb{R})$, we put $V_{\mathbb{C}} = V \otimes_{\mathbb{R}} \mathbb{C}$ and define $\varepsilon(V_{\mathbb{C}}, \Psi)$ by the additivity using the epsilon factors in Section 2A. For $V \in R(W_K, \mathbb{R})$, we define the i -th Stiefel–Whitney class $w_i(V) \in H^i(G_K, \mathbb{Z}/2\mathbb{Z})$ for $i \geq 0$ as in [Deligne 1976, (1.3)]. Let

$$\text{cl} : H^2(G_K, \mathbb{Z}/2\mathbb{Z}) \rightarrow H^2(G_K, K^{\text{ac}, \times}) \xrightarrow{\sim} \mathbb{Q}/\mathbb{Z},$$

where the first map is induced by $\mathbb{Z}/2\mathbb{Z} \rightarrow K^{\text{ac}, \times}$, $m \mapsto (-1)^m$ and the second isomorphism is the invariant map.

Theorem 6.1 [Deligne 1976, Théorème 1.5]. *Assume that $V \in R(W_K, \mathbb{R})$ has dimension 0 and determinant 1. Then we have*

$$\varepsilon(V_{\mathbb{C}}, \Psi) = \exp(2\pi\sqrt{-1} \text{cl}(w_2(V))).$$

In particular, we have $\varepsilon(V_{\mathbb{C}}, \Psi) = 1$ if $\text{ch } K = 2$.

6B. Discriminant. Let L be a finite separable extension of K . We put

$$\delta_{L/K} = \det(\text{Ind}_{L/K} 1).$$

6B1. Multiplicative discriminant. Assume $\text{ch } K \neq 2$. We define $d_{L/K} \in K^\times / (K^\times)^2$ as the discriminant of the quadratic form $\text{Tr}_{L/K}(x^2)$ on L . For $a \in K^\times / (K^\times)^2$, let $\{a\} \in H^1(G_K, \mathbb{Z}/2\mathbb{Z})$ and $\kappa_a \in \text{Hom}(W_K, \{\pm 1\})$ be the elements corresponding to a under the natural isomorphisms

$$K^\times / (K^\times)^2 \simeq H^1(G_K, \mathbb{Z}/2\mathbb{Z}) \simeq \text{Hom}(W_K, \{\pm 1\}).$$

We have

$$(6-1) \quad \delta_{L/K} = \kappa_{d_{L/K}}$$

by [Bourbaki 1981, Chapter V, Section 10, Example 2(6)] (see [Serre 1984, Section 1.4]). For $a, b \in K^\times / (K^\times)^2$, we put

$$\{a, b\} = \{a\} \cup \{b\} \in H^2(G_K, \mathbb{Z}/2\mathbb{Z}).$$

Proposition 6.2 [Abbes and Saito 2010, Proposition 6.5]. *Let m be the extension degree of L over K . We take a generator a of L over K . Let $f(x) \in K[x]$ be the minimal polynomial of a . We put $D = f'(a) \in L$. Then we have*

$$d_{L/K} = (-1)^{\binom{m}{2}} \text{Nr}_{L/K}(D) \in K^\times / (K^\times)^2,$$

$$w_2(\text{Ind}_{L/K} \kappa_D) = \binom{m}{4} \{-1, -1\} + \{d_{L/K}, 2\} \in H^2(G_K, \mathbb{Z}/2\mathbb{Z}).$$

6B2. Additive discriminant. We put $P_m(x) = x^m - x$ for any positive integer m . We assume that $\text{ch } K = 2$.

Definition 6.3 [Bergé and Martinet 1985, Définition 2.7]. Let m be the extension degree of L over K . Let $f(x) \in K[x]$ be the minimal polynomial of a generator of L over K . We have a decomposition $f(x) = \prod_{1 \leq i \leq m} (x - a_i)$ over the Galois closure of L over K . We put

$$d_{L/K}^+ = \sum_{1 \leq i < j \leq m} \frac{a_i a_j}{(a_i + a_j)^2} \in K/P_2(K),$$

which we call the additive discriminant of L over K .

Theorem 6.4 [Bergé and Martinet 1985, Théorème 2.7]. *Let L' be the subextension of K^{ac} over K corresponding to $\text{Ker } \delta_{L/K}$. Then the extension L' over K corresponds to $d_{L/K}^+ \in K/P_2(K)$ by the Artin–Schreier theory.*

7. Product formula of Deligne–Laumon

We recall a statement of the product formula of Deligne–Laumon. In this paper, we need only the rank one case, which is proved in [Deligne 1973, Proposition 10.12.1], but we follow the notation from [Laumon 1987].

7A. Local factor. We consider a triple (T, \mathcal{F}, ω) which consists of the following.

- The affine scheme $T = \text{Spec } \mathcal{O}_{K_T}$ where \mathcal{O}_{K_T} is the ring of integers in a local field K_T of characteristic p whose residue field contains k .
- A constructible $\overline{\mathbb{Q}}_\ell$ -sheaf \mathcal{F} on T .
- A nonzero meromorphic 1-form ω on T .

Then we can associate $\varepsilon_{\psi_0}(T, \mathcal{F}, \omega) \in \mathbb{C}^\times$ to the triple (T, \mathcal{F}, ω) as in [Laumon 1987, Théorème 3.1.5.4] using ι .

Assume that $K_T = k((t))$. Let $\eta = \text{Spec } k((t))$ be the generic point of T with the natural inclusion $j : \eta \rightarrow T$. We define a character $\Psi_\omega : k((t)) \rightarrow \mathbb{C}^\times$ by

$$\Psi_\omega(a) = (\psi_0 \circ \text{Tr}_{k/\mathbb{F}_p})(\text{Res}(a\omega)) \quad \text{for } a \in k((t)).$$

Let $l(\Psi_\omega)$ be the level of Ψ_ω in the sense of [Bushnell and Henniart 2006, Definition 1.7]. We fix an algebraic closure $k((t))^{\text{ac}}$ of $k((t))$. For a rank 1 smooth $\overline{\mathbb{Q}}_\ell$ -sheaf V on η corresponding to a character $\chi : G_{k((t))} \rightarrow \mathbb{C}^\times$ via ι , we have

$$(7-1) \quad \varepsilon_{\psi_0}(T, j_* V, \omega) = q^{-l(\Psi_\omega)/2} \varepsilon(\chi \omega_{-1/2}, \Psi_\omega)$$

by [Laumon 1987, Théorème 3.1.5.4(v); Tate 1979, (3.6.2)] and [Bushnell and Henniart 2006, Proposition 23.1(3)].

7B. Product formula. Let X be a geometrically connected proper smooth curve over k of genus g . Let \mathcal{F} be a constructible $\overline{\mathbb{Q}}_\ell$ -sheaf on X . Let $\text{Frob}_q \in G_k$ be the geometric Frobenius element. We put

$$\varepsilon(X, \mathcal{F}) = \iota \left(\prod_{i=0}^2 \det(-\text{Frob}_q; H^i(X \otimes_k k^{\text{ac}}, \mathcal{F}))^{(-1)^{i-1}} \right).$$

Let $\text{rk}(\mathcal{F})$ be the generic rank of \mathcal{F} .

Theorem 7.1 [Laumon 1987, Théorème 3.2.1.1]. *Let ω be a nonzero meromorphic 1-form on X . Then we have*

$$\varepsilon(X, \mathcal{F}) = q^{\text{rk}(\mathcal{F})(1-g)} \prod_{x \in |X|} \varepsilon_{\psi_0}(X(x), \mathcal{F}|_{X(x)}, \omega|_{X(x)}),$$

where $|X|$ is the set of closed points of X , and $X(x)$ is the completion of X at x .

8. Determinant

In this section, we study $\det \tau_{\psi_0}$ to show the equality $\omega_{\pi_{\zeta, \chi, c}} = \det \tau_{\zeta, \chi, c}$ of the central character and the determinant. We use the product formula of Deligne–Laumon to study $\det \tau_{\psi_0}(\text{Fr}(1))$, where $\text{Fr}(1)$ is defined in (2-2).

Lemma 8.1. *We have $Q^{\text{ab}} = Q/Q_0$.*

Proof. By Lemma 2.4, we have $Q^{\text{ab}} = (Q/F)^{\text{ab}}$. For $(a, b, c) \in Q$, let (a, b) be the image of (a, b, c) in Q/F . Then we have

$$(a, 0)(a, b)(a, 0)^{-1}(a, b)^{-1} = (1, (a-1)b).$$

Hence, we obtain the claim. \square

We view θ_0 defined in (2-8) as a character of Q by $(a, b, c) \mapsto \theta_0(a)$. Recall that τ^0 is the representation of Q defined in (2-14).

Lemma 8.2. *We have $\det \tau^0 = \theta_0$.*

Proof. By Lemma 8.1, it suffices to show $\det \tau^0 = \theta_0$ on $\mu_{p^e+1}(k^{\text{ac}})$. By Lemma 2.2 and Lemma 2.5, we have

$$\det \tau^0(a) = \prod_{\chi \in \mu_{p^e+1}(k^{\text{ac}})^\vee \setminus \{1\}} \chi(a)$$

for $a \in \mu_{p^e+1}(k^{\text{ac}})$. Hence, the claim follows. \square

For $a \in k^\times$, let $\left(\frac{a}{k}\right)$ denote the quadratic residue symbol of k defined by

$$\left(\frac{a}{k}\right) = \begin{cases} 1 & \text{if } a \text{ is square in } k, \\ -1 & \text{if } a \text{ is not square in } k. \end{cases}$$

Lemma 8.3. *Let m be a positive integer that is prime to p . We take an m -th root $\varpi^{1/m}$ of ϖ , and put $L = K(\varpi^{1/m})$.*

(1) *If m is odd, then $\delta_{L/K}$ is the unramified character satisfying $\delta_{L/K}(\varpi) = \left(\frac{q}{m}\right)$.*

(2) *If m is even, we have $\delta_{L/K}(\varpi) = \left(\frac{-1}{q}\right)^{m/2}$ and $\delta_{L/K}(x) = \left(\frac{x}{k}\right)$ for $x \in \mathcal{O}_K^\times$.*

Proof. These are proved in [Bushnell and Fröhlich 1983, (10.1.6)] if $\text{ch } K = 0$. Actually, the same proof works also in the positive characteristic case. \square

Lemma 8.4. *Let m, m' be positive integers that are prime to p . We take an m -th root $\varpi^{1/m}$ of ϖ , and put $L = K(\varpi^{1/m})$. Let $\psi'_K : K \rightarrow \mathbb{C}^\times$ be a character such that $\psi'_K(x) = \psi_0(\text{Tr}_{k/\mathbb{F}_p}(m'\bar{x}))$ for $x \in \mathcal{O}_K$. Then we have*

$$\lambda(L/K, \psi'_K) = \begin{cases} \left(\frac{q}{m}\right) & \text{if } m \text{ is odd,} \\ -(-\epsilon(p))\left(\frac{2mm'}{p}\right)\left(\frac{-1}{p}\right)^{(m/2)-1} & \text{if } m \text{ is even.} \end{cases}$$

Proof. If m is odd, we have

$$\lambda(L/K, \psi'_K) = \varepsilon(\delta_{L/K}, \psi'_K) = \left(\frac{q}{m}\right)$$

by [Henniart 1984, Proposition 2] and Lemma 8.3(1).

Assume that m is even. Note that $p \neq 2$ in this case. Then we have

$$(8-1) \quad d_{L/K} = (-1)^{m/2} \text{Nr}_{L/K}(m(\varpi^{1/m})^{m-1}) = -(-1)^{m/2} \varpi \in K^\times / (K^\times)^2$$

by [Proposition 6.2](#). For $\chi \in (\mathbb{F}_q^\times)^\vee$ and $\psi \in \mathbb{F}_q^\vee \setminus \{1\}$, we set

$$\tau(\chi, \psi) = - \sum_{x \in \mathbb{F}_q^\times} \chi^{-1}(x) \psi(x)$$

and have the Hasse–Davenport formula

$$(8-2) \quad \tau(\chi \circ \mathrm{Nr}_{\mathbb{F}_{q^n}/\mathbb{F}_q}, \psi \circ \mathrm{Tr}_{\mathbb{F}_{q^n}/\mathbb{F}_q}) = \tau(\chi, \psi)^n.$$

Let

$$(\cdot, \cdot)_K : K^\times / (K^\times)^2 \times K^\times / (K^\times)^2 \rightarrow \{\pm 1\}$$

denote the Hilbert symbol. By [\(6-1\)](#) and [\(8-1\)](#), we have

$$\delta_{L/K}(x) = \kappa_{d_{L/K}}(x) = (x, d_{L/K})_K = (x, \varpi)_K = \left(\frac{\bar{x}}{k}\right)$$

for $x \in \mathcal{O}_K^\times$. By [\[Bushnell and Henniart 2006, Theorem 23.5\]](#), we have

$$\varepsilon(\delta_{L/K}, \psi'_K) = q^{-1/2} \sum_{x \in \mathcal{O}_K^\times / U_K^1} \delta_{L/K}(x) \psi'_K(x) = q^{-1/2} \sum_{x \in k^\times} \left(\frac{x}{k}\right) \psi_0(\mathrm{Tr}_{k/\mathbb{F}_p}(m'x)).$$

By applying [\(8-2\)](#) to the extension k over \mathbb{F}_p and using [\(2-16\)](#), we have

$$q^{-1/2} \sum_{x \in k^\times} \left(\frac{x}{k}\right) \psi_0(\mathrm{Tr}_{k/\mathbb{F}_p}(m'x)) = - \left(-\varepsilon(p) \left(\frac{m'}{p}\right)\right)^f.$$

Hence, we have

$$\begin{aligned} \lambda(L/K, \psi'_K) &= \varepsilon(\delta_{L/K}, \psi'_K) \left(\frac{m}{q}\right) \left(\frac{-1}{q}\right)^{(m/2)-1} (d_{L/K}, 2)_K \\ &= - \left(-\varepsilon(p) \left(\frac{2mm'}{p}\right) \left(\frac{-1}{p}\right)^{(m/2)-1}\right)^f \end{aligned}$$

by [\[Saito 1995, Theorem II.2B\]](#) and [\[Tate 1979, \(3.6.1\)\]](#). □

Lemma 8.5. *We have*

$$\det \tau_{\psi_0}(\mathrm{Fr}(1)) = \begin{cases} \left(-\varepsilon(p) \left(\frac{2}{p}\right)\right)^f q^{p^e/2} & \text{if } p \neq 2, \\ q^{2^e-1} & \text{if } p = 2. \end{cases}$$

Proof. Let x be the standard coordinate of \mathbb{A}_k^1 . Let j be the open immersion $\mathbb{A}_k^1 \hookrightarrow \mathbb{P}_k^1$. We put $t = 1/x$. As in [Section 7A](#), we put $T = \mathrm{Spec} k[[t]]$ and $\eta = \mathrm{Spec} k((t))$ with the open immersion $j : \eta \rightarrow T$.

We consider $k((s))$ as a subfield of $k((t))$ by $s = t^{p^e+1}$. Let $\tilde{\xi} : G_{k((s))} \rightarrow \mathbb{C}^\times$ be the Artin–Schreier character associated to $y^p - y = 1/s$ and ψ_0 , which means the composite of

$$G_{k((s))} \rightarrow \mathbb{F}_p, \quad \sigma \mapsto \sigma(y) - y$$

and ψ_0^{-1} where y is an element of $k((t))^{\mathrm{ac}}$ such that $y^p - y = 1/s$.

We use the notation in [Lemma 5.5](#). Note that $\psi_{k((s))(y)/k((s))}(1) = 1$ by [Lemma 5.4](#). We can check that

$$\mathrm{Nr}_{k((s))(y)/k((s))}(1 + y^{-1}x) = 1 + s(x^p - x)$$

for $x \in k$. For $x \in \mathcal{O}_{k((s))}$, we have

$$\begin{aligned} \tilde{\xi}(1 + sx) &= \psi_0^{-1}(\mathrm{Art}_{k((s))}(1 + sx)(y) - y) \\ &= \psi_0^{-1}\left(-p_{k((s))(y), y^{-1}}\left(\frac{\mathrm{Art}_{k((s))}(1 + sx)(y^{-1})}{y^{-1}}\right)\right) = \psi_0(\mathrm{Tr}_{k/\mathbb{F}_p}(\bar{x})), \end{aligned}$$

where we use [Lemma 5.5](#) with $\alpha = s$, $\beta = \varpi_{k((s))} = y^{-1}$. Hence, we have $\mathrm{rsw}(\tilde{\xi}, \Psi_{s^{-1}ds}) = s$ by [Proposition 5.3\(1\)](#).

Let $\xi : G_{k((t))} \rightarrow \mathbb{C}^\times$ be the restriction of $\tilde{\xi}$ to $G_{k((t))}$. Then ξ is the Artin–Schreier character associated to $y^p - y = 1/t^{p^e+1}$ and ψ_0 .

Let V_ξ be the smooth $\overline{\mathbb{Q}}_\ell$ -sheaf on η corresponding to ξ via ι . Then we have $V_\xi \simeq \mathcal{L}_{\psi_0}|_\eta$ by [\[Deligne 1977, Définition 1.7 in Sommes trig.\]](#). Let the notation be as in [Lemma 2.2](#). We write ω for the meromorphic 1-form dx on \mathbb{P}_k^1 . By [\[Laumon 1987, Théorème 3.1.5.4\(v\)\]](#), we have

$$\varepsilon_{\psi_0}(X_{(x)}, (j! \pi^* \mathcal{L}_{\psi_0})|_{X_{(x)}}, \omega|_{X_{(x)}}) = 1$$

for any $x \in |\mathbb{A}_k^1|$ with $X = \mathbb{P}_k^1$ in the notation of [Theorem 7.1](#). We simply write ω for $\omega|_T$. Then we have

$$\det \tau_{\psi_0}(\mathrm{Fr}(1)) = (-1)^{p^e} \varepsilon(\mathbb{P}_k^1, j! \pi^* \mathcal{L}_{\psi_0}) = (-1)^p q \varepsilon_{\psi_0}(T, j! V_\xi, \omega)$$

by [Theorem 7.1](#). Since ξ is a ramified character, we have $j! V_\xi \simeq j_* V_\xi$. Hence,

$$\varepsilon_{\psi_0}(T, j! V_\xi, \omega) = \varepsilon_{\psi_0}(T, j_* V_\xi, \omega) = q^{-1} \varepsilon(\xi \omega_{-1/2}, \Psi_\omega)$$

by (7-1). Since $\omega = -t^{-2}dt$ on T , we have

$$\varepsilon(\xi \omega_{-1/2}, \Psi_\omega) = (\xi \omega_{-1/2})(-t^{-1}) \varepsilon(\xi \omega_{-1/2}, \Psi_{t^{-1}dt})$$

by [\[Bushnell and Henniart 2006, 23.5 Lemma 1\]](#). We have

$$\xi(-t^{-1}) = \xi(-t^{p^e}) = \xi(-t)^{p^e} = 1,$$

since $\mathrm{Nr}_{k((t))(y)/k((t))}(y) = 1/t^{p^e+1}$. Hence we obtain

$$(\xi \omega_{-1/2})(-t^{-1}) \varepsilon(\xi \omega_{-1/2}, \Psi_{t^{-1}dt}) = q^{p^e/2} \varepsilon(\xi, \Psi_{t^{-1}dt})$$

by [Lemma 4.2](#), since $\mathrm{rsw}(\xi, \Psi_{t^{-1}dt}) = s$ by $\mathrm{rsw}(\tilde{\xi}, \Psi_{s^{-1}ds}) = s$ and [Proposition 5.2\(2\)](#). By [Proposition 5.3\(2\)](#), we have $\varepsilon(\tilde{\xi}, \Psi_{s^{-1}ds}) = \tilde{\xi}(s) = 1$, since the level of $\tilde{\xi}$ is 1 and $\mathrm{Nr}_{k((s))(y)/k((s))}(y^{-1}) = s$. Hence, we obtain

$$\varepsilon(\xi, \Psi_{t^{-1}dt}) = \lambda(k((t))/k((s)), \Psi_{s^{-1}ds})^{-1} \delta_{k((t))/k((s))}(\mathrm{rsw}(\tilde{\xi}, \Psi_{s^{-1}ds}))$$

by [Proposition 5.1](#). By [Lemmas 8.4](#) and [8.3](#), we respectively have

$$\lambda(k((t))/k((s)), \Psi_{s^{-1}ds}) = \begin{cases} -(-\epsilon(p)\left(\frac{2}{p}\right)\left(\frac{-1}{p}\right)^{(p^e-1)/2})^f & \text{if } p \neq 2, \\ \left(\frac{q}{p^e+1}\right) & \text{if } p = 2 \end{cases}$$

and

$$\delta_{k((t))/k((s))}(\text{rsw}(\tilde{\xi}, \Psi_{s^{-1}ds})) = \begin{cases} \left(\frac{-1}{q}\right)^{(p^e+1)/2} & \text{if } p \neq 2, \\ \left(\frac{q}{p^e+1}\right) & \text{if } p = 2. \end{cases}$$

The claim follows from the above equalities. \square

We simply write τ_ζ for $\tau_{\zeta,1,1}$.

Proposition 8.6. *We have $\omega_{\pi_{\zeta,\chi,c}} = \det \tau_{\zeta,\chi,c}$.*

Proof. By [\(2-13\)](#) and [\[Gallagher 1965, \(1\)\]](#), we have

$$(8-3) \quad \det \tau_{\zeta,\chi,c} = \delta_{E_\zeta/K}^{p^e} (\det \tau_{n,\zeta,\chi,c})|_{K^\times},$$

since $\delta_{E_\zeta/K} = \det(\text{Ind}_{E_\zeta/K} 1)$ and the transfer homomorphism $W_K^{\text{ab}} \rightarrow W_{E_\zeta}^{\text{ab}}$ is compatible with the natural inclusion $K^\times \rightarrow E_\zeta^\times$ under the Artin reciprocity maps. Hence, we may assume $\chi = 1$ and $c = 1$ by twist (see [\(2-13\)](#)). Then it suffices to show $\det \tau_\zeta = 1$. We see that $\det \tau_\zeta$ is unramified by [\(2-9\)](#), [Lemmas 2.5, 8.2, 8.3](#) and equation [\(8-3\)](#).

If p and n' are odd, then we have

$$\begin{aligned} \det \tau_\zeta(\varpi) &= \left(\frac{q}{n'}\right)^{p^e} \left(-\epsilon(p)\left(\frac{2}{p}\right)p^{p^e/2}\right)^{fn'} \left(\left(-\epsilon(p)\left(\frac{-2n'}{p}\right)\right)^n p^{-\frac{1}{2}}\right)^{fn'p^e} \\ &= \left(\left(\frac{p}{n'}\right)\left(\frac{n'}{p}\right)\epsilon(p)^{p^en-1}\right)^{fn'} = \left(\left(\frac{p}{n'}\right)\left(\frac{n'}{p}\right)(-1)^{\frac{1}{2}(p-1)\frac{1}{2}(n'-1)}\right)^{fn'} = 1 \end{aligned}$$

by [\(8-3\)](#), [Lemmas 8.3\(1\)](#) and [8.5](#). We see that $\det \tau_\zeta(\varpi) = 1$ similarly also in the other case using [\(8-3\)](#), [Lemmas 8.3](#) and [8.5](#). \square

9. Imprimitve field

In this section, we construct a field extension T_ζ^u of E_ζ such that $\tau_{n,\zeta}|_{W_{T_\zeta^u}}$ is an induction of a character. We call T_ζ^u an imprimitive field of $\tau_{n,\zeta}$, since $\tau_{n,\zeta}|_{W_{T_\zeta^u}}$ is not primitive.

9A. Construction of character. Here we construct subgroups $R \subset Q' \subset Q \rtimes \mathbb{Z}$ and a character ϕ_n of R . Later (see [Section 9B](#)) we will see that $\tau_n|_{Q'} \simeq \text{Ind}_R^{Q'} \phi_n$. Our imprimitive field T_ζ^u will correspond to the subgroup $Q' \subset Q \rtimes \mathbb{Z}$.

Let e_0 be the positive integer such that $e_0 \in 2^{\mathbb{N}}$ and e/e_0 is odd.

Lemma 9.1. *Assume $p \neq 2$. Then we have $\text{Tr} \tau_{\psi_0}(\text{Fr}(2e_0)) = p^{e_0}$.*

Proof. For $a \in k^{\text{ac}}$ and $b \in \mathbb{F}_{p^{2e_0}}$ such that $a^p - a = b^{p^e+1}$, we have that

$$(9-1) \quad a^{p^{2e_0}} - a = \text{Tr}_{\mathbb{F}_{p^{2e_0}}/\mathbb{F}_p}(b^{p^e+1}).$$

By (9-1) and the Lefschetz trace formula, we see that

$$\begin{aligned} \text{Tr } \tau_{\psi_0}(\text{Fr}(2e_0)) &= - \sum_{b \in \mathbb{F}_{p^{2e_0}}} (\psi_0 \circ \text{Tr}_{\mathbb{F}_{p^{e_0}}/\mathbb{F}_p})(\text{Tr}_{\mathbb{F}_{p^{2e_0}}/\mathbb{F}_{p^{e_0}}}(b^{p^e+1})) \\ &= -1 - (p^{e_0} + 1) \sum_{x \in \mathbb{F}_{p^{e_0}}^\times} (\psi_0 \circ \text{Tr}_{\mathbb{F}_{p^{e_0}}/\mathbb{F}_p})(x) = p^{e_0} \end{aligned}$$

using $(p^e + 1, p^{2e_0} - 1) = p^{e_0} + 1$. \square

Corollary 9.2. *Assume $p \neq 2$. Then we have $\text{Tr } \tau_n(\text{Fr}(2e_0)) = (-1)^{ne_0(p-1)/2}$.*

Proof. This follows from (2-9) and Lemma 9.1. \square

Let n_0 be the biggest integer such that 2^{n_0} divides $p^{e_0} + 1$. We take $r \in k^{\text{ac}}$ such that $r^{2^{n_0}} = -1$. We define a subgroup R_0 of Q_0 by

$$R_0 = \{(1, b, c) \in Q_0 \mid b^{p^e} - rb = 0\}.$$

Lemma 9.3. (1) *If $p \neq 2$, then the action of $2e_0\mathbb{Z} \subset \mathbb{Z}$ on Q stabilizes R_0 .*

(2) *If $p = 2$, then the action of \mathfrak{g} on $Q \rtimes \mathbb{Z}$ by conjugation stabilizes R_0 .*

Proof. The first claim follows from $r^{p^{2e_0}-1} = 1$. We can see the second claim easily using (2-19). \square

We put

$$Q' = \begin{cases} Q_0 \rtimes (2e_0\mathbb{Z}) & \text{if } p \neq 2, \\ Q_0 \rtimes \mathbb{Z} & \text{if } p = 2, \end{cases} \quad R = \begin{cases} R_0 \rtimes (2e_0\mathbb{Z}) & \text{if } p \neq 2, \\ R_0 \cdot \langle \mathfrak{g} \rangle & \text{if } p = 2 \end{cases}$$

as subgroups of $Q \rtimes \mathbb{Z}$, which are well-defined by Lemma 9.3. We are going to construct a character ϕ_n of R in this subsection. Then, we will show that $\tau_n|_{Q'} \simeq \text{Ind}_R^{Q'} \phi_n$ in the next subsection.

First, we consider the case where p is odd. We define a homomorphism $\phi_n : R \rightarrow \mathbb{C}^\times$ by

$$(9-2) \quad \begin{aligned} \phi_n((1, b, c), 0) &= \psi_0\left(c - \frac{1}{2} \sum_{i=0}^{e-1} (rb^2)^{p^i}\right) \quad \text{for } (1, b, c) \in R_0, \\ \phi_n(\text{Fr}(2e_0)) &= (-1)^{ne_0((p-1)/2)}. \end{aligned}$$

Then ϕ_n extends the character ψ_0 of F .

Next, we consider the case where $p = 2$. We define an abelian group R'_0 as

$$R'_0 = \{(b, c) \mid b \in \mathbb{F}_2, c \in \mathbb{F}_{2^{2e}}, c^{2^e} - c = b\},$$

with the multiplication given by

$$(b_1, c_1) \cdot (b_2, c_2) = (b_1 + b_2, c_1 + c_2 + b_1 b_2).$$

We define $\phi : R_0 \rightarrow R'_0$ by

$$\phi((1, b, c)) = \left(\text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}(b), c + \sum_{0 \leq i < j \leq e-1} b^{2^i+2^j} \right) \quad \text{for } (1, b, c) \in R_0,$$

which is a homomorphism by

$$(9-3) \quad \text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}(b) \text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}(b') = \text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}(bb') + \sum_{0 \leq i < j \leq e-1} (b^{2^i} b'^{2^j} + b'^{2^i} b^{2^j})$$

for $b, b' \in \mathbb{F}_{2^e}$. Let $b_0 \in \mathbb{F}_{2^{2e}}$ be as before [Lemma 2.8](#). Let F' be the kernel of the homomorphism

$$\mathbb{F}_{2^e} \rightarrow \mathbb{F}_2, \quad c \mapsto \text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}((b_0 + b_0^{2^e})c).$$

We put $R''_0 = R'_0/F'$, where we consider F' as a subgroup of R'_0 by $c \mapsto (0, c)$. Then R''_0 is a cyclic group of order 4. We write $\bar{g}(b, c)$ for the image of $(b, c) \in R'_0$ under the projection $R'_0 \rightarrow R''_0$. Let $\phi' : R_0 \rightarrow R''_0$ be the composite of ϕ and the projection $R'_0 \rightarrow R''_0$. We put

$$(9-4) \quad s = \sum_{i=0}^{e-1} b_0^{2^i}, \quad t = \text{Tr}_{\mathbb{F}_{2^{2e}}/\mathbb{F}_{2^e}}(b_0).$$

We have $s^2 + s = t$ and $\text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}(t) = \text{Tr}_{\mathbb{F}_{2^{2e}}/\mathbb{F}_2}(b_0) = 1$. We have

$$\left(1, s^2 + \sum_{0 \leq i < j \leq e-1} t^{2^i+2^j} \right) \in R'_0,$$

which is of order 4. The element $\bar{g}(1, s^2 + \sum_{0 \leq i < j \leq e-1} t^{2^i+2^j})$ is a generator of R''_0 , because

$$2\bar{g}\left(1, s^2 + \sum_{0 \leq i < j \leq e-1} t^{2^i+2^j}\right) = \bar{g}(0, 1) \neq 0.$$

Let $\tilde{\psi}_0 : R''_0 \rightarrow \mathbb{C}^\times$ be the faithful character satisfying

$$\tilde{\psi}_0\left(\bar{g}\left(1, s^2 + \sum_{0 \leq i < j \leq e-1} t^{2^i+2^j}\right)\right) = -\sqrt{-1}.$$

We define a homomorphism $\phi_n : R \rightarrow \mathbb{C}^\times$ by

$$(9-5) \quad \begin{aligned} \phi_n(((1, b, c), 0)) &= (\tilde{\psi}_0 \circ \phi')((1, b, c)) \quad \text{for } (1, b, c) \in R_0, \\ \phi_n(\mathbf{g}) &= (-1)^{\frac{1}{8}n(n-2)} \frac{-1+\sqrt{-1}}{\sqrt{2}}, \end{aligned}$$

which is a character of order 8. Then ϕ_n extends the character ψ_0 of F .

9B. Induction of character.

Lemma 9.4. *We have $\tau_n|_{Q'} \simeq \text{Ind}_R^{Q'} \phi_n$.*

Proof. We write $\tilde{\psi}_n$ for $\phi_n|_{R_0}$. We know that $\tau_n|_{Q_0} \cong \text{Ind}_{R_0}^{Q_0} \tilde{\psi}_n$ by [Proposition 1.2](#), since R_0 is an abelian group such that $2 \dim_{\mathbb{F}_p}(R_0/F) = \dim_{\mathbb{F}_p}(Q_0/F)$.

First, we consider the case where p is odd. The claim for general f follows from the claim for $f = 1$ by the restriction. Hence, we may assume that $f = 1$.

If $\tilde{\psi} \in R_0^\vee$ satisfies $\tilde{\psi}|_F = \psi_0$, then we have $\tau_n|_{Q_0} \cong \text{Ind}_{R_0}^{Q_0} \tilde{\psi}$ by [Proposition 1.2](#), and obtain an injective homomorphism $\tilde{\psi} \hookrightarrow \tau_n|_{R_0}$ as representations of R_0 by Frobenius reciprocity. Hence we have a decomposition

$$(9-6) \quad \tau_n|_{R_0} = \bigoplus_{\tilde{\psi} \in R_0^\vee, \tilde{\psi}|_F = \psi_0} \tilde{\psi},$$

since the number of $\tilde{\psi} \in R_0^\vee$ such that $\tilde{\psi}|_F = \psi_0$ is p^e .

We put

$$\bar{R}_0 = \{b \in k^{\text{ac}} \mid b^{p^e} - rb = 0\}.$$

The $\tilde{\psi}_n$ -component in (9-6) is the unique component that is stable by the action of $((1, 0, 0), 2e_0)$, since the homomorphism

$$\bar{R}_0 \rightarrow \bar{R}_0, \quad b \mapsto b^{p^{2e_0}} - b$$

is an isomorphism. Hence, we have a nontrivial homomorphism $\phi_n \rightarrow \tau_n|_R$ by [Corollary 9.2](#). Then we have a nontrivial homomorphism $\text{Ind}_R^{Q'} \phi_n \rightarrow \tau_n|_{Q'}$ by Frobenius reciprocity. The representation $\tau_n|_{Q'}$ is irreducible by [Corollary 2.6](#). Then we obtain the claim, since $[Q' : R] = p^e$.

Next we consider the case where $p = 2$. Then it suffices to show that

$$\text{Tr}(\text{Ind}_R^{Q'} \phi_n)(g^{-1}) = -(-1)^{\frac{1}{8}n(n-2)} \sqrt{2}$$

by (2-9) and [Proposition 2.9](#). We have a decomposition

$$(9-7) \quad (\text{Ind}_R^{Q'} \phi_n)|_{R_0} = \bigoplus_{\phi \in R_0^\vee, \phi|_F = \psi_0} \phi.$$

Let $\tilde{\psi}'_n$ be the twist of $\tilde{\psi}_n$ by the character

$$R_0 \rightarrow \bar{\mathbb{Q}}_\ell^\times, \quad (1, b, c) \mapsto \psi_0(\text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}(b)).$$

Then only the $\tilde{\psi}_n$ -component and the $\tilde{\psi}'_n$ -component in (9-6) are stable by the action of $((1, b_0, c_0), 1)$, since the image of the homomorphism

$$\mathbb{F}_{2^e} \rightarrow \mathbb{F}_{2^e}, \quad b \mapsto b^2 - b$$

is equal to $\text{Ker Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}$. The action of $\text{Fr}(e)$ permutes the $\tilde{\psi}_n$ -component and the $\tilde{\psi}'_n$ -component. Hence, \mathbf{g} acts on the $\tilde{\psi}'_n$ -component by $\phi_n(\mathbf{g})$ times

$$\phi_n(\text{Fr}(e)^{-1} \mathbf{g} \text{Fr}(e) \mathbf{g}^{-1}) = \phi_n\left(\left(\left(1, t, c_0 + c_0^{2^e} + \sum_{i=0}^{e-1} (b_0^{2^e+1} + b_0^{2^e+1})^{2^i}\right), 0\right)\right) = \sqrt{-1}.$$

Hence we have

$$\text{Tr}(\text{Ind}_R^Q \phi_n)(\mathbf{g}^{-1}) = (1 - \sqrt{-1}) \phi_n(\mathbf{g}^{-1}) = -(-1)^{\frac{1}{8}n(n-2)} \sqrt{2}. \quad \square$$

We use the notations from equation (2-10). We set $T_\zeta = E_\zeta(\alpha_\zeta)$, $M_\zeta = T_\zeta(\beta_\zeta)$ and $N_\zeta = M_\zeta(\gamma_\zeta)$. Let f_0 be the positive integer such that $f_0 \in 2^{\mathbb{N}}$ and f/f_0 is odd. We put

$$N = \begin{cases} 2e_0/f_0 & \text{if } p \neq 2 \text{ and } f_0 | 2e_0, \\ 1, & \text{otherwise.} \end{cases}$$

Let K^{ur} be the maximal unramified extension of K in K^{ac} . Let $K^{\text{u}} \subset K^{\text{ur}}$ be the unramified extension of degree N over K . Let k_N be the residue field of K^{u} . For a finite field extension L of K in K^{ac} , we write L^{u} for the composite field of L and K^{u} in K^{ac} . For $a \in k^{\text{ac}}$, we write $\hat{a} \in \mathcal{O}_{K^{\text{ur}}}$ for the Teichmüller lift of a . We put

$$(9-8) \quad \delta'_\zeta = \begin{cases} \beta_\zeta^{p^e} - \hat{r} \beta_\zeta & \text{if } p \neq 2, \\ \beta_\zeta^{2^e} - \beta_\zeta + \sum_{i=0}^{e-1} \hat{b}^{2^i} & \text{if } p = 2, \end{cases} \quad \epsilon_1 = \begin{cases} 0 & \text{if } p \neq 2, \\ 1 & \text{if } p = 2. \end{cases}$$

Then we have

$$\delta_\zeta'^{p^e} - \hat{r}^{-1} \delta'_\zeta \equiv -\alpha_\zeta^{-1} + \epsilon_1 \pmod{\mathfrak{p}_{T_\zeta^{\text{u}}}(\delta'_\zeta)}.$$

We take $\delta_\zeta \in T_\zeta^{\text{u}}(\delta'_\zeta)$ such that

$$(9-9) \quad \delta_\zeta^{p^e} - \hat{r}^{-1} \delta_\zeta = -\alpha_\zeta^{-1} + \epsilon_1, \quad \delta_\zeta \equiv \delta'_\zeta \pmod{\mathfrak{p}_{T_\zeta^{\text{u}}}(\delta'_\zeta)}.$$

We put $M_\zeta^{\text{u}} = T_\zeta^{\text{u}}(\delta_\zeta)$. The image of $\Theta_\zeta|_{W_{M_\zeta^{\text{u}}}}$ is contained in R . Let $\xi_{n,\zeta} : W_{M_\zeta^{\text{u}}} \rightarrow \mathbb{C}^\times$ be the composite of the restrictions $\Theta_\zeta|_{W_{M_\zeta^{\text{u}}}}$ and $\phi_n|_R$. By the local class field theory, we regard $\xi_{n,\zeta}$ as a character of $M_\zeta^{\text{u}\times}$.

Proposition 9.5. *We have $\tau_{n,\zeta}|_{W_{T_\zeta^{\text{u}}}} \simeq \text{Ind}_{M_\zeta^{\text{u}}/T_\zeta^{\text{u}}} \xi_{n,\zeta}$.*

Proof. This follows from Lemma 9.4. □

Remark 9.6. Our imprimitive field is different from that in [Bushnell and Henniart 2014, Section 5.1]. In our case, T_ζ^{u} need not be normal over K . This choice is technically important in our proof of the main result.

9C. Study of character. Here we study the character $\xi_{n,\zeta}$ in detail.

Assume that $\text{ch } K = p$ and $f = 1$ in this subsection. We will use results in this subsection to compute the epsilon factor of $\xi_{n,\zeta}$ later after a reduction to the case where $\text{ch } K = p$ and $f = 1$. By (2-10), (9-8), (9-9) and $\text{ch } K = p$, we have that $\delta_\zeta = \delta'_\zeta$.

9C1. Odd case. Assume $p \neq 2$. We put

$$(9-10) \quad \theta_\zeta = \gamma_\zeta + \frac{1}{2} \sum_{i=0}^{e-1} (r\beta_\zeta^2)^{p^i}.$$

Since $r^{p^{e_0}+1} = -1$ and $(p^e + 1)/(p^{e_0} + 1)$ is an odd integer, we have $r^{p^e+1} = -1$. Then we have

$$(9-11) \quad \begin{aligned} \theta_\zeta^p - \theta_\zeta &= \beta_\zeta^{p^e+1} - \frac{1}{2r} (\beta_\zeta^{2p^e} + r^2 \beta_\zeta^2) \\ &= -\frac{1}{2r} (\beta_\zeta^{2p^e} - 2r\beta_\zeta^{p^e+1} + r^2 \beta_\zeta^2) = -\frac{1}{2r} \delta_\zeta^2. \end{aligned}$$

We put $N_\zeta^u = M_\zeta^u(\theta_\zeta)$. Let $\xi'_{n,\zeta}$ be the twist of $\xi_{n,\zeta}$ by the unramified character

$$W_{M_\zeta^u} \rightarrow \mathbb{C}^\times, \quad \sigma \mapsto \sqrt{-1}^{n n_\sigma (p-1)/2},$$

where n_σ is as before (2-12).

Lemma 9.7. *If $p \neq 2$, then $\xi'_{n,\zeta}$ factors through $\text{Gal}(N_\zeta^u/M_\zeta^u)$.*

Proof. Let $\sigma \in \text{Ker } \xi'_{n,\zeta}$. Recall that $a_\sigma, b_\sigma, c_\sigma$ are defined in (2-11). Then we have $(\bar{a}_\sigma, \bar{b}_\sigma, \bar{c}_\sigma) \in R_0$ and

$$\bar{c}_\sigma - \frac{1}{2} \sum_{i=0}^{e-1} (r\bar{b}_\sigma^2)^{p^i} = 0$$

by (9-2). Hence, we see that

$$\begin{aligned} \sigma(\theta_\zeta) - \theta_\zeta &= c_\sigma - \sum_{i=0}^{e-1} (r b_\sigma (\beta_\zeta + b_\sigma))^{p^i} + \frac{1}{2} \sum_{i=0}^{e-1} (r((\beta_\zeta + b_\sigma)^2 - \beta_\zeta^2))^{p^i} \\ &= c_\sigma - \frac{1}{2} \sum_{i=0}^{e-1} (r b_\sigma^2)^{p^i} \equiv 0 \pmod{\mathfrak{p}_{N_\zeta^u}} \end{aligned}$$

by (2-11). Therefore, we obtain the claim by $\sigma(\delta_\zeta) = \delta_\zeta$ and (9-11). \square

9C2. Even case. Assume $p = 2$. Let $\xi'_{n,\zeta}$ be the twist of $\xi_{n,\zeta}$ by the character

$$(9-12) \quad W_{M_\zeta^u} \rightarrow \mathbb{C}^\times, \quad \sigma \mapsto \left((-1)^{\frac{1}{8}n(n-2) - \frac{1+\sqrt{-1}}{\sqrt{2}}} \right)^{n_\sigma}.$$

We take $b_1, b_2 \in k^{\text{ac}}$ such that

$$(9-13) \quad b_1^2 - b_1 = s, \quad b_2^2 - b_2 = t \left(b_1^2 + \sum_{i=0}^{e-1} (b_1 s)^{2^i} \right).$$

We put

$$(9-14) \quad \eta_\zeta = \sum_{i=0}^{e-1} \beta_\zeta^{2^i} + b_1, \quad \gamma'_\zeta = \gamma_\zeta + \sum_{0 \leq i < j \leq e-1} \beta_\zeta^{2^i+2^j},$$

and

$$(9-15) \quad \theta'_\zeta = \sum_{i=0}^{e-1} (t\gamma'_\zeta)^{2^i} + \sum_{0 \leq i \leq j \leq e-2} t^{2^i} (\delta_\zeta \eta_\zeta)^{2^j} + \sum_{0 \leq j < i \leq e-1} t^{2^i} (b_1 \delta_\zeta + s \eta_\zeta)^{2^j} + b_1^2 \eta_\zeta + b_2.$$

Lemma 9.8. *We have $\eta_\zeta^2 - \eta_\zeta = \delta_\zeta$ and $\theta_\zeta'^2 - \theta'_\zeta = (\delta_\zeta \eta_\zeta)^{2^{e-1}}$.*

Proof. We can check the first claim easily. We show the second claim. We use P_m in [Section 6B2](#). We have

$$(9-16) \quad P_2(\gamma'_\zeta) = (\beta_\zeta^{2^e} - \beta_\zeta) \sum_{i=0}^{e-1} \beta_\zeta^{2^i} + \beta_\zeta^2 = (\delta_\zeta - s)(\eta_\zeta - b_1) + \beta_\zeta^2.$$

Hence, we have

$$(9-17) \quad P_{2^e}(\gamma'_\zeta) = \sum_{i=0}^{e-1} ((\delta_\zeta - s)(\eta_\zeta - b_1))^{2^i} + (\eta_\zeta - b_1)^2.$$

By $b_1^4 + b_1 = s^2 + s = t$ and $\eta_\zeta^2 - \eta_\zeta = \delta_\zeta$, we have

$$(b_1^2 \eta_\zeta)^2 + b_1^2 \eta_\zeta = t \eta_\zeta^2 + b_1 \eta_\zeta^2 + b_1^2 \eta_\zeta = t \eta_\zeta^2 + b_1(\eta_\zeta^2 + \eta_\zeta) + s \eta_\zeta = t \eta_\zeta^2 + b_1 \delta_\zeta + s \eta_\zeta.$$

Hence, by using $\sum_{i=1}^{e-1} t^{2^i} = 1 - t$ and $t \in \mathbb{F}_{2^e}$, we have

$$\begin{aligned} \theta_\zeta'^2 - \theta'_\zeta &= t P_{2^e}(\gamma'_\zeta) + t \sum_{i=0}^{e-1} (\delta_\zeta \eta_\zeta + b_1 \delta_\zeta + s \eta_\zeta)^{2^i} + (\delta_\zeta \eta_\zeta)^{2^{e-1}} + t \eta_\zeta^2 + b_2^2 - b_2 \\ &= t \left(\sum_{i=0}^{e-1} (b_1 s)^{2^i} + \eta_\zeta^2 + b_1^2 \right) + (\delta_\zeta \eta_\zeta)^{2^{e-1}} + t \eta_\zeta^2 + b_2^2 - b_2 = (\delta_\zeta \eta_\zeta)^{2^{e-1}}, \end{aligned}$$

where we use [\(9-17\)](#) at the second equality and [\(9-13\)](#) at the third one. \square

We take $\theta_\zeta \in K^{\text{ac}}$ such that $\theta'_\zeta = \theta_\zeta^{2^{e-1}}$. Then we have $\theta_\zeta^2 - \theta_\zeta = \delta_\zeta \eta_\zeta$. We put $N_\zeta^{\text{u}} = M_\zeta^{\text{u}}(\eta_\zeta, \theta_\zeta)$, which is a cyclic extension of M_ζ^{u} of order 4 by [Lemma 9.8](#).

Lemma 9.9. *The character $\xi'_{n,\zeta}$ factors through $\text{Gal}(N_\zeta^{\text{u}}/M_\zeta^{\text{u}})$.*

Proof. Let $\sigma \in \text{Ker } \xi'_{n,\zeta}$. We take $\sigma_1, \sigma_2 \in \text{Ker } \xi'_{n,\zeta}$ such that $\sigma = \sigma_1 \sigma_2^{-n\sigma}$, $\sigma_1 \in I_{M_\zeta^{\text{u}}}$ and $\Theta_\zeta(\sigma_2) = ((1, b_0, c_0), -1)$. Then we have $(\bar{a}_{\sigma_1}, \bar{b}_{\sigma_1}, \bar{c}_{\sigma_1}) \in R_0$, $\text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}(\bar{b}_{\sigma_1}) = 0$ and

$$(9-18) \quad \text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2} \left(t \left(\bar{c}_{\sigma_1} + \sum_{0 \leq i < j \leq e-1} \bar{b}_{\sigma_1}^{2^i + 2^j} \right) \right) = 0$$

by [\(9-5\)](#). It suffices to show that $\sigma_i(\eta_\zeta) = \eta_\zeta$ and $\sigma_i(\theta'_\zeta) = \theta'_\zeta$ for $i = 1, 2$.

We have

$$\begin{aligned}\sigma_1(\eta_\zeta) - \eta_\zeta &\equiv \sum_{i=0}^{e-1} b_{\sigma_1}^{2^i} \equiv 0 \pmod{\mathfrak{p}_{N_\zeta^u}}, \\ \sigma_2(\eta_\zeta) - \eta_\zeta &\equiv \sum_{i=0}^{e-1} b_0^{2^i} + b_1^2 - b_1 \equiv 0 \pmod{\mathfrak{p}_{N_\zeta^u}}\end{aligned}$$

by $\text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}(\bar{b}_{\sigma_1}) = 0$ and $b_1^2 - b_1 = s$. By [Lemma 9.8](#), we have

$$\sigma_i(\eta_\zeta) - \eta_\zeta \in \mathbb{F}_2 \quad \text{for } i = 1, 2.$$

Hence, we have $\sigma_i(\eta_\zeta) = \eta_\zeta$ for $i = 1, 2$. We have

$$\sigma_1(\theta'_\zeta) - \theta'_\zeta = \sum_{i=0}^{e-1} \left(t(\sigma_1(\gamma'_\zeta) - \gamma'_\zeta) \right)^{2^i}.$$

Further, we have

$$\begin{aligned}\sigma_1(\gamma'_\zeta) - \gamma'_\zeta &\equiv c_{\sigma_1} + \sum_{i=0}^{e-1} (b_{\sigma_1})^{2^{i+1}} + \sum_{i=0}^{e-1} b_{\sigma_1}^{2^i} \sum_{i=0}^{e-1} \beta_\zeta^{2^i} + \sum_{0 \leq i < j \leq e-1} b_{\sigma_1}^{2^i+2^j} \\ &\equiv c_{\sigma_1} + \sum_{0 \leq i < j \leq e-1} b_{\sigma_1}^{2^i+2^j} \pmod{\mathfrak{p}_{N_\zeta^u}},\end{aligned}$$

where we use [\(2-11\)](#) and $\bar{b}_{\sigma_1} \in \mathbb{F}_{2^e}$ at the first equality, and use $\text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}(\bar{b}_{\sigma_1}) = 0$ at the second one. This implies $\sigma_1(\theta'_\zeta) \equiv \theta'_\zeta \pmod{\mathfrak{p}_{N_\zeta^u}}$ by [\(9-18\)](#). By a similar argument as above using [Lemma 9.8](#), we obtain $\sigma_1(\theta'_\zeta) = \theta'_\zeta$.

It remains to show $\sigma_2(\theta'_\zeta) = \theta'_\zeta$. Using [\(9-16\)](#) and $\text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}(t) = 1$, we see that

$$(9-19) \quad \sum_{i=0}^{e-1} (t\gamma'_\zeta)^{2^i} = \gamma'_\zeta + \sum_{1 \leq i \leq j \leq e-1} t^{2^j} \beta_\zeta^{2^i} + \sum_{0 \leq i < j \leq e-1} t^{2^j} ((\delta_\zeta - s)(\eta_\zeta - b_1))^{2^i}.$$

We put

$$\gamma''_\zeta = \gamma'_\zeta + \sum_{1 \leq i \leq j \leq e-1} t^{2^j} \beta_\zeta^{2^i}.$$

By $c_0^2 + c_0 = b_0^{2^e+1}$ and $t = b_0 + b_0^{2^e}$ (see [\(2-18\)](#), [\(9-4\)](#)), we have

$$\sigma_2(\gamma_\zeta) - \gamma_\zeta \equiv c_0 + \sum_{i=0}^{e-1} (b_0^{2^e}(\beta_\zeta + b_0))^{2^i} \equiv c_0^{2^e} + \sum_{i=0}^{e-1} ((b_0 + t) \beta_\zeta)^{2^i} \pmod{\mathfrak{p}_{N_\zeta^u}}.$$

Then we have

$$\sigma_2(\gamma'_\zeta) - \gamma'_\zeta \equiv c_0^{2^e} + s(\eta_\zeta - b_1) + \sum_{i=0}^{e-1} (t\beta_\zeta)^{2^i} + \sum_{0 \leq i < j \leq e-1} b_0^{2^i+2^j} \pmod{\mathfrak{p}_{N_\zeta^u}}$$

by (9-4) and (9-14). Hence, we have

$$\begin{aligned}
\sigma_2(\gamma'_\zeta) - \gamma'_\zeta &\equiv \sigma_2(\gamma'_\zeta) - \gamma'_\zeta + \sum_{1 \leq i \leq j \leq e-1} t^{2^{j+1}} (\beta_\zeta + b_0)^{2^i} - \sum_{1 \leq i \leq j \leq e-1} t^{2^j} \beta_\zeta^{2^i} \\
&\equiv \sigma_2(\gamma'_\zeta) - \gamma'_\zeta + t(\eta_\zeta - b_1) + \sum_{i=0}^{e-1} (t\beta_\zeta)^{2^i} + \sum_{1 \leq i < j \leq e} b_0^{2^i} t^{2^j} \\
&\equiv c_0^{2^e} + s^2(\eta_\zeta - b_1) + \sum_{0 \leq i < j \leq e-1} b_0^{2^i+2^j} + \sum_{1 \leq i < j \leq e} b_0^{2^i} t^{2^j} \pmod{\mathfrak{p}_{N_\zeta^u}},
\end{aligned}$$

where we use (9-14) and $t \in \mathbb{F}_{2^e}$ at the second equality and $s^2 + s = t$ at the last equality. We can check that

$$c_0^{2^e} + \sum_{0 \leq i < j \leq e-1} b_0^{2^i+2^j} + \sum_{1 \leq i < j \leq e} b_0^{2^i} t^{2^j} = st$$

by (2-17), (9-4) and $\text{Tr}_{\mathbb{F}_{2^{2e}}/\mathbb{F}_2}(b_0) = 1$. As a result, we obtain

$$\sigma_2(\gamma''_\zeta) - \gamma''_\zeta \equiv s^2 \eta_\zeta + b_1 s^2 + st \pmod{\mathfrak{p}_{N_\zeta^u}}.$$

Hence, by (9-15) and (9-19), we have

$$\sigma_2(\theta'_\zeta) - \theta'_\zeta \equiv \sum_{i=0}^{2^{e-1}+2^{e-2}} d_i \eta_\zeta^i \pmod{\mathfrak{p}_{N_\zeta^u}}$$

for some $d_i \in k^{\text{ac}}$. We have

$$d_0 = b_1 s^2 + st + t \sum_{j=1}^{e-1} (b_1 s)^{2^j} + b_1 s \sum_{l=1}^{e-1} t^{2^l} + b_2^2 - b_2 = 0.$$

This implies $\sigma_2(\theta'_\zeta) = \theta'_\zeta$, since we know that $\sigma_2(\theta'_\zeta) - \theta'_\zeta \in \mathbb{F}_2$ by Lemma 9.8. \square

10. Refined Swan conductor

Let $\tilde{K} \subset K^{\text{ur}}$ be the unramified extension of K^{u} generated by $\mu_{p^{4pe-1}}(K^{\text{ur}})$. For a finite field extension L of K in K^{ac} , we write \tilde{L} for the composite field of L and \tilde{K} in K^{ac} . We write \tilde{M}'_ζ for $\tilde{M}_\zeta^{\text{u}}$. Then \tilde{N}_ζ is a Galois extension of \tilde{M}'_ζ . By equations (9-8) and (9-9), we can take $\beta'_\zeta \in \tilde{M}'_\zeta$ such that

$$(10-1) \quad \beta'^{p^e}_\zeta - \hat{r} \beta'_\zeta = \delta_\zeta, \quad \beta'_\zeta \equiv \beta_\zeta \pmod{\mathfrak{p}_{\tilde{M}'_\zeta}},$$

since there is $x \in \mathbb{F}_{2^{4e}}$ such that $x^{2^e} - x = \sum_{i=0}^{e-1} b_0^{2^i}$ if $p = 2$. Then we have $\tilde{M}_\zeta = \tilde{M}'_\zeta(\beta'_\zeta)$ by Krasner's lemma.

Lemma 10.1. (1) *We have*

$$(10-2) \quad \psi_{\tilde{N}_\zeta/\tilde{M}'_\zeta}(v) = \begin{cases} v & \text{if } v \leq 1, \\ p^e(v-1) + 1 & \text{if } 1 < v \leq 2, \\ p^{e+1}(v-2) + p^e + 1 & \text{if } 2 < v. \end{cases}$$

(2) We have

$$\mathrm{Gal}(\tilde{N}_\zeta/\tilde{M}'_\zeta)_i = \begin{cases} \mathrm{Gal}(\tilde{N}_\zeta/\tilde{M}'_\zeta) & \text{if } i \leq 1, \\ \mathrm{Gal}(\tilde{N}_\zeta/\tilde{M}_\zeta) & \text{if } 2 \leq i \leq p^e + 1, \\ \{1\} & \text{if } p^e + 2 \leq i. \end{cases}$$

Proof. We have

$$\psi_{\tilde{M}'_\zeta/\tilde{M}'_\zeta}(v) = \begin{cases} v & \text{if } v \leq 1, \\ p^e(v-1)+1 & \text{if } v > 1, \end{cases}$$

$$\psi_{\tilde{N}_\zeta/\tilde{M}_\zeta}(v) = \begin{cases} v & \text{if } v \leq p^e + 1, \\ p(v-p^e-1)+p^e+1 & \text{if } v > p^e + 1 \end{cases}$$

by (2-10), (10-1) and Lemma 5.4 noting that \hat{r} has a $(p^e - 1)$ -st root in \tilde{M}'_ζ . Hence, claim (1) follows from $\psi_{\tilde{N}_\zeta/\tilde{M}'_\zeta} = \psi_{\tilde{N}_\zeta/\tilde{M}_\zeta} \circ \psi_{\tilde{M}_\zeta/\tilde{M}'_\zeta}$. Claim (2) follows from claim (1) and

$$\mathrm{Gal}(\tilde{N}_\zeta/\tilde{M}'_\zeta)_{p^e+1} \supset \mathrm{Gal}(\tilde{N}_\zeta/\tilde{M}_\zeta)_{p^e+1} = \mathrm{Gal}(\tilde{N}_\zeta/\tilde{M}_\zeta). \quad \square$$

We set

$$\varpi_{\tilde{M}'_\zeta} = \delta_\zeta^{-1}, \quad \varpi_{\tilde{M}_\zeta} = \beta_\zeta^{-1} \quad \text{and} \quad \varpi_{\tilde{N}_\zeta} = (\gamma_\zeta \varpi_{\tilde{M}_\zeta}^{p^e-1})^{-1}.$$

Then the elements $\varpi_{\tilde{M}'_\zeta}$, $\varpi_{\tilde{M}_\zeta}$ and $\varpi_{\tilde{N}_\zeta}$ are uniformizers of \tilde{M}'_ζ , \tilde{M}_ζ and \tilde{N}_ζ respectively. Let \tilde{k} be the residue field of \tilde{K} .

Lemma 10.2. *We have a commutative diagram*

$$\begin{array}{ccc} U_{\tilde{N}_\zeta}^{p^e+1} & \xrightarrow{\mathrm{Nr}_{\tilde{N}_\zeta/\tilde{M}'_\zeta}} & U_{\tilde{M}'_\zeta}^2 \\ \downarrow & & \downarrow \\ \tilde{k} & \xrightarrow{P} & \tilde{k} \end{array}$$

where the map P is given by $x \mapsto x^p - x$ and the vertical maps are given by

$$P_{\tilde{N}_\zeta, -\gamma_\zeta^{-1}} : U_{\tilde{N}_\zeta}^{p^e+1} \rightarrow \tilde{k}, \quad 1 - x\gamma_\zeta^{-1} \mapsto \bar{x},$$

$$P_{\tilde{M}'_\zeta, \hat{r}\varpi_{\tilde{M}'_\zeta}^2} : U_{\tilde{M}'_\zeta}^2 \rightarrow \tilde{k}, \quad 1 + x\hat{r}\varpi_{\tilde{M}'_\zeta}^2 \mapsto \bar{x}.$$

Proof. The norm maps $\mathrm{Nr}_{\tilde{N}_\zeta/\tilde{M}_\zeta}$ and $\mathrm{Nr}_{\tilde{M}_\zeta/\tilde{M}'_\zeta}$ induce

$$U_{\tilde{N}_\zeta}^{p^e+1}/U_{\tilde{N}_\zeta}^{p^e+2} \rightarrow U_{\tilde{M}_\zeta}^{p^e+1}/U_{\tilde{M}_\zeta}^{p^e+2}, \quad 1 - u\gamma_\zeta^{-1} \mapsto 1 - (u^p - u)\varpi_{\tilde{M}_\zeta}^{p^e+1},$$

$$U_{\tilde{M}_\zeta}^{p^e+1}/U_{\tilde{M}_\zeta}^{p^e+2} \rightarrow U_{\tilde{M}'_\zeta}^2/U_{\tilde{M}'_\zeta}^3, \quad 1 - u\varpi_{\tilde{M}_\zeta}^{p^e+1} = 1 - u\beta_\zeta^{-1}\varpi_{\tilde{M}'_\zeta} \mapsto 1 + u\hat{r}\varpi_{\tilde{M}'_\zeta}^2$$

respectively by Lemma 5.5(1) and calculations of the norms. Hence, the claim follows. \square

For any finite extension M of K , we write ψ_M for the composite $\psi_K \circ \text{Tr}_{M/K}$.

Lemma 10.3. *We have $\text{rsw}(\xi_{n,\zeta}|_{W_{M_\zeta^u}}, \psi_{M_\zeta^u}) = -n'\delta_\zeta^{-(p^e+1)} \pmod{U_{M_\zeta^u}^1}$.*

Proof. We put $\tilde{\xi}_{n,\zeta} = \xi_{n,\zeta}|_{W_{\tilde{M}'_\zeta}}$, and regard it as a character of $\tilde{M}'_\zeta{}^\times$. By (2-12), Lemmas 5.5(1) and Lemma 10.1, the restriction of $\tilde{\xi}_{n,\zeta}$ to $U_{\tilde{M}'_\zeta}^2$ is given by the composition

$$U_{\tilde{M}'_\zeta}^2 \xrightarrow{\text{Art}_{\tilde{M}'_\zeta}} \text{Gal}(\tilde{N}_\zeta/\tilde{M}'_\zeta) \simeq \mathbb{F}_p \xrightarrow{\psi_0} \overline{\mathbb{Q}}_\ell^\times,$$

where the isomorphism $\text{Gal}(\tilde{N}_\zeta/\tilde{M}'_\zeta) \simeq \mathbb{F}_p$ is given by $\sigma \mapsto \overline{\sigma(\gamma_\zeta) - \gamma_\zeta}$. We define $p_{\tilde{N}_\zeta, -\gamma_\zeta^{-1}}$ as in Lemma 10.2. For $u \in \mathcal{O}_{\tilde{M}'_\zeta}$, we put $\sigma_u = \text{Art}_{\tilde{M}'_\zeta}(1 + u\hat{r}\overline{\omega}_{\tilde{M}'_\zeta}^2)$ and then have

$$\begin{aligned} (10-3) \quad \tilde{\xi}_{n,\zeta}(1 + u\hat{r}\overline{\omega}_{\tilde{M}'_\zeta}^2) &= \psi_0(\overline{\sigma_u(\gamma_\zeta) - \gamma_\zeta}) \\ &= \psi_0\left(p_{\tilde{N}_\zeta, -\gamma_\zeta^{-1}}\left(\frac{\gamma_\zeta}{\sigma_u(\gamma_\zeta)}\right)\right) \\ &= \psi_0\left(p_{\tilde{N}_\zeta, -\gamma_\zeta^{-1}}\left(\frac{\sigma_u(\overline{\omega}_{\tilde{N}_\zeta})}{\overline{\omega}_{\tilde{N}_\zeta}}\right)\right) = \psi_0 \circ \text{Tr}_{\tilde{k}/\mathbb{F}_p}(\bar{u}), \end{aligned}$$

where we use Lemmas 5.5(2) and 10.2 at the last equality. Since we have

$$\overline{\text{Tr}_{\tilde{M}'_\zeta/\tilde{T}_\zeta}(\delta_\zeta^{p^e-1}u)} = -r^{-1}\bar{u}$$

for $u \in \mathcal{O}_{\tilde{M}'_\zeta}$, we obtain

$$\tilde{\xi}_{n,\zeta}(1+x) = \psi_{\tilde{M}'_\zeta}(-n'^{-1}\delta_\zeta^{p^e+1}x)$$

for $x \in \mathfrak{p}_{\tilde{M}'_\zeta}^2$ by (10-3). This implies

$$(10-4) \quad \xi_{n,\zeta}(1+x) = \psi_{M_\zeta^u}(-n'^{-1}\delta_\zeta^{p^e+1}x)$$

for $x \in \mathfrak{p}_{M_\zeta^u}^2$, because $\text{Tr}_{\tilde{k}/k_N} : \tilde{k} \rightarrow k_N$ is surjective. The claim follows from (10-4) and Proposition 5.3(1). \square

Lemma 10.4. *We have $\text{rsw}(\tau_{n,\zeta,\chi,c}, \psi_{E_\zeta}) = n'\varphi'_\zeta \pmod{U_{E_\zeta}^1}$.*

Proof. By Proposition 5.2(1), we may assume that $\chi = 1$, $c = 1$. By Proposition 9.5 and Lemma 10.3, we have

$$(10-5) \quad \text{rsw}(\tau_{n,\zeta}|_{W_{T_\zeta^u}}, \psi_{T_\zeta^u}) = \text{Nr}_{M_\zeta^u/T_\zeta^u}(\text{rsw}(\xi_{n,\zeta}, \psi_{M_\zeta^u})) = n'\varphi'_\zeta \pmod{U_{T_\zeta^u}^1}.$$

Since T_ζ^u is a tamely ramified extension of E_ζ , we have

$$(10-6) \quad \text{rsw}(\tau_{n,\zeta}, \psi_{E_\zeta}) = \text{rsw}(\tau_{n,\zeta}|_{W_{T_\zeta^u}}, \psi_{T_\zeta^u}) \pmod{U_{T_\zeta^u}^1}$$

by Proposition 5.2(2). The claim follows from (10-5) and (10-6). \square

Proposition 10.5. *We have $\text{rsw}(\tau_{\zeta, \chi, c}, \psi_K) = \text{rsw}(\pi_{\zeta, \chi, c}, \psi_K)$.*

Proof. By $\tau_{\zeta, \chi, c} = \text{Ind}_{E_{\zeta}/K} \tau_{n, \zeta, \chi, c}$, we have

$$(10-7) \quad \text{rsw}(\tau_{\zeta, \chi, c}, \psi_K) = \text{Nr}_{E_{\zeta}/K}(\text{rsw}(\tau_{n, \zeta, \chi, c}, \psi_{E_{\zeta}})).$$

Hence, the claim follows from Lemmas 4.5 and 10.4. \square

Lemma 10.6. *We have $\text{Sw}(\tau_{\zeta, \chi, c}) = 1$.*

Proof. This follows from Lemma 10.4 and (10-7). \square

Lemma 10.7. *The representation $\tau_{\zeta, \chi, c}$ is irreducible.*

Proof. We know that the restriction of $\tau_{n, \zeta, \chi, c}$ to the wild inertia subgroup of $W_{E_{\zeta}}$ is irreducible by Corollary 2.6. Assume that $\tau_{\zeta, \chi, c}$ is not irreducible. Then we have an irreducible factor τ' of $\tau_{\zeta, \chi, c}$ such that $\text{Sw}(\tau') = 0$, by Lemma 10.6 and the additivity of Sw . Then, the restriction of τ' to the wild inertia subgroup of W_K is trivial by $\text{Sw}(\tau') = 0$. On the other hand, we have an injective homomorphism $\tau_{n, \zeta, \chi, c} \rightarrow \tau'|_{W_{E_{\zeta}}}$ by Frobenius reciprocity. This is a contradiction. \square

Proposition 10.8. *The representation $\tau_{\zeta, \chi, c}$ is irreducible of Swan conductor 1.*

Proof. This follows from Lemmas 10.6 and 10.7. \square

11. Epsilon factor

11A. Reduction to special cases. In this subsection, we show the equality

$$\varepsilon(\tau_{\zeta, \chi, c}, \psi_K) = \varepsilon(\pi_{\zeta, \chi, c}, \psi_K)$$

of epsilon factors assuming some results in the special case where $n = p^e$, $\text{ch } K = p$ and $f = 1$. The results in the special case will be proved in the next subsection.

Lemma 11.1. *We have*

$$\lambda(E_{\zeta}/K, \psi_K) = \begin{cases} \left(\frac{q}{n'}\right) & \text{if } n' \text{ is odd,} \\ -(-\epsilon(p))\left(\frac{2n'}{p}\right)\left(\frac{-1}{p}\right)^{(n'/2)-1} & \text{if } n' \text{ is even,} \end{cases}$$

$$\lambda(T_{\zeta}^u/E_{\zeta}, \psi_{E_{\zeta}}) = \begin{cases} -(-1)^{\frac{1}{4}(p-1)fN} & \text{if } p \neq 2, \\ \left(\frac{q}{p^e+1}\right) & \text{if } p = 2. \end{cases}$$

Proof. We have

$$\lambda(T_{\zeta}^u/E_{\zeta}, \psi_{E_{\zeta}}) = \lambda(T_{\zeta}^u/E_{\zeta}^u, \psi_{E_{\zeta}^u}) \lambda(E_{\zeta}^u/E_{\zeta}, \psi_{E_{\zeta}})^{p^e+1} = \lambda(T_{\zeta}^u/E_{\zeta}^u, \psi_{E_{\zeta}^u}).$$

If $p \neq 2$, then we have

$$\lambda(T_{\zeta}^u/E_{\zeta}^u, \psi_{E_{\zeta}^u}) = -\left(-\epsilon(p)\left(\frac{2n'}{p}\right)\left(\frac{-1}{p}\right)^{(p^e-1)/2}\right)^{fN} = -(-1)^{\frac{1}{4}(p-1)fN}$$

by Lemma 8.4, since fN is even. The other assertions immediately follow from Lemma 8.4. \square

Lemma 11.2. *We have*

$$\lambda(M_\zeta^u/T_\zeta^u, \psi_{T_\zeta^u}) = \begin{cases} (-1)^f & \text{if } p = 2 \text{ and } e \leq 2, \\ \left(\frac{r}{k_N}\right), & \text{otherwise.} \end{cases}$$

Proof. Let $K_{(0)}$ and $K_{(p)}$ be nonarchimedean local fields of characteristic 0 and p respectively. Assume that the residue fields of $K_{(0)}$ and $K_{(p)}$ are isomorphic to k . We take uniformizers $\varpi_{(0)}$ and $\varpi_{(p)}$ of $K_{(0)}$ and $K_{(p)}$ respectively. We define $T_{\zeta,(0)}^u$ similarly as T_ζ^u starting from $K_{(0)}$. We use similar notations also for other objects in the characteristic zero side and the positive characteristic side. We have the isomorphism

$$\mathcal{O}_{T_{\zeta,(p)}^u}/\mathfrak{p}_{T_{\zeta,(p)}^u}^2 \xrightarrow{\sim} \mathcal{O}_{T_{\zeta,(0)}^u}/\mathfrak{p}_{T_{\zeta,(0)}^u}^2, \quad \xi_0 + \xi_1 \varpi_{T_{\zeta,(p)}^u} \mapsto \hat{\xi}_0 + \hat{\xi}_1 \varpi_{T_{\zeta,(0)}^u}$$

of algebras, where $\xi_0, \xi_1 \in k$. Hence, it suffices to show the claim in one of the characteristic zero side and the positive characteristic side by [Deligne 1984, Proposition 3.7.1], since $\text{Gal}(M_{\zeta,(p)}^u/T_{\zeta,(p)}^u)^2 = 1$ and $\text{Gal}(M_{\zeta,(0)}^u/T_{\zeta,(0)}^u)^2 = 1$, where we use upper numbering filtration of Galois groups.

First, we consider the case where $p \neq 2$ and $\text{ch } K = p$. Then, we have $d_{M_\zeta^u/T_\zeta^u} = \hat{r}$ by Proposition 6.2 and the fact that fN is even. Hence, $\delta_{M_\zeta^u/T_\zeta^u}$ is unramified by (6-1). Hence, we have

$$\lambda(M_\zeta^u/T_\zeta^u, \psi_{T_\zeta^u}) = \varepsilon(\delta_{M_\zeta^u/T_\zeta^u}, \psi_{T_\zeta^u})^{p^e} = \left(\frac{r}{k_N}\right)$$

by [Henniart 1984, Proposition 2; Bushnell and Henniart 2006, Proposition 23.5] and (6-1).

We consider the case where $p = 2$. Assume that $e \geq 3$ and $\text{ch } K = 0$. We have $D = 2^e \delta_\zeta^{2^e - 1} + 1$ in the notation of Proposition 6.2 with $(L, K, a) = (M_\zeta^u, T_\zeta^u, \delta_\zeta)$. Then, we have $D \in (M_\zeta^u)^\times$. Hence, we have $\kappa_D = 1$, $d_{M_\zeta^u/T_\zeta^u} = 1$ and

$$w_2(\text{Ind}_{M_\zeta^u/T_\zeta^u} 1) = 1$$

by Proposition 6.2 and $\binom{p^e}{4} \equiv 0 \pmod{2}$. Therefore we have

$$\lambda(M_\zeta^u/T_\zeta^u, \psi_{T_\zeta^u}) = \varepsilon(\text{Ind}_{M_\zeta^u/T_\zeta^u} 1, \psi_{T_\zeta^u}) = \varepsilon(1^{\oplus p^e}, \psi_{T_\zeta^u}) = 1$$

by Theorem 6.1.

Assume that $e = 2$ and $\text{ch } K = 2$. Then we see that $d_{M_\zeta^u/T_\zeta^u}^+ = 1$ by Definition 6.3. Hence, $\delta_{M_\zeta^u/T_\zeta^u}$ is the unramified character satisfying

$$\delta_{M_\zeta^u/T_\zeta^u}(\varpi_{T_\zeta^u}) = (-1)^f$$

by Theorem 6.4. Then we see that

$$\lambda(M_\zeta^u/T_\zeta^u, \psi_{T_\zeta^u}) = \varepsilon(\text{Ind}_{M_\zeta^u/T_\zeta^u} 1, \psi_{T_\zeta^u}) = \varepsilon(\delta_{M_\zeta^u/T_\zeta^u} \oplus 1^{\oplus 3}, \psi_{T_\zeta^u}) = (-1)^f,$$

where we use Theorem 6.1 at the second equality.

Assume that $e = 1$ and $\text{ch } K = 2$. Let $\kappa_{M_\zeta^u/T_\zeta^u}$ be the quadratic character associated to the extension M_ζ^u over T_ζ^u . Then we have

$$\lambda(M_\zeta^u/T_\zeta^u, \psi_{T_\zeta^u}) = \varepsilon(\kappa_{M_\zeta^u/T_\zeta^u}, \psi_{T_\zeta^u})$$

by [Theorem 6.1](#) similarly as above. We can check that the norm map $\text{Nr}_{M_\zeta^u/T_\zeta^u}$ induces

$$U_{M_\zeta^u}^1/U_{M_\zeta^u}^2 \rightarrow U_{T_\zeta^u}^1/U_{T_\zeta^u}^2, \quad 1 + u\delta_\zeta^{-1} \mapsto 1 + (u^2 - u)\alpha_\zeta.$$

Then, by [Lemma 5.5](#), we have

$$\begin{aligned} (11-1) \quad \kappa_{M_\zeta^u/T_\zeta^u}(1 + \alpha_\zeta x) &= \psi_0(\text{Art}_{T_\zeta^u}(1 + \alpha_\zeta x)(\delta_\zeta) - \delta_\zeta) \\ &= \psi_0\left(p_{M_\zeta^u, \delta_\zeta^{-1}}\left(\frac{\text{Art}_{T_\zeta^u}(1 + \alpha_\zeta x)(\delta_\zeta^{-1})}{\delta_\zeta^{-1}}\right)\right) \\ &= \psi_0(\text{Tr}_{k/\mathbb{F}_p}(\bar{x})) \end{aligned}$$

for $x \in \mathcal{O}_{T_\zeta^u}$ noting that $k_N = k$. Hence, we have $\text{rsw}(\kappa_{M_\zeta^u/T_\zeta^u}, \psi_{T_\zeta^u}) = \alpha_\zeta$ by [Proposition 5.3\(1\)](#). By [Proposition 5.3\(2\)](#), we have

$$\varepsilon(\kappa_{M_\zeta^u/T_\zeta^u}, \psi_{T_\zeta^u}) = \kappa_{M_\zeta^u/T_\zeta^u}(\alpha_\zeta) = \kappa_{M_\zeta^u/T_\zeta^u}(1 + \alpha_\zeta) = (-1)^f,$$

where we use $\text{Nr}_{M_\zeta^u/T_\zeta^u}(\delta_\zeta) = \alpha_\zeta^{-1} + 1$ and (11-1) at the last equality. \square

Lemma 11.3. *We have*

$$\text{Tr}_{M_\zeta^u/T_\zeta^u}(\delta_\zeta^i) = \begin{cases} 0 & \text{if } 1 \leq i \leq p^e - 2, \\ \hat{r}^{-1}(p^e - 1) & \text{if } i = p^e - 1. \end{cases}$$

Proof. Vanishing for $1 \leq i \leq p^e - 2$ follows from [\(9-9\)](#). We have also

$$\text{Tr}_{M_\zeta^u/T_\zeta^u}(\delta_\zeta^{p^e-1}) = \text{Tr}_{M_\zeta^u/T_\zeta^u}(\hat{r}^{-1} + \delta_\zeta^{-1}(-\alpha_\zeta^{-1} + \epsilon_1)) = \hat{r}^{-1}(p^e - 1)$$

by [\(9-9\)](#). \square

Lemma 11.4. *We have*

$$\delta_{T_\zeta^u/E_\zeta}(\text{rsw}(\tau_{n,\zeta}, \psi_{E_\zeta})) = \begin{cases} 1 & \text{if } p \neq 2, \\ \left(\frac{q}{p^e+1}\right) & \text{if } p = 2. \end{cases}$$

Proof. If $p = 2$, the claim follows from [Lemmas 8.3\(1\) and 10.4](#), since T_ζ^u is totally ramified over E_ζ .

Assume that $p \neq 2$. Then we have $d_{T_\zeta^u/E_\zeta^u} = (-1)^{(p^e+1)/2}\varphi'_\zeta$ by [Proposition 6.2](#). Hence, we have $\delta_{T_\zeta^u/E_\zeta^u}((-1)^{(p^e-1)/2}\varphi'_\zeta) = 1$ by [Lemma 8.3\(2\)](#). Therefore, we have

$$\begin{aligned} \delta_{T_\zeta^u/E_\zeta}(\text{rsw}(\tau_{n,\zeta}, \psi_{E_\zeta})) &= \delta_{T_\zeta^u/E_\zeta^u}(n'\varphi'_\zeta) \\ &= \delta_{T_\zeta^u/E_\zeta^u}(n'(-1)^{(p^e-1)/2}) = \left(\frac{n'(-1)^{(p^e-1)/2}}{q^N}\right) = 1 \end{aligned}$$

by [\[Gallagher 1965, \(1\)\]](#), [Lemmas 8.3\(2\), 10.4](#) and the fact that fN is even. \square

Lemma 11.5. *Assume that $n = p^e$. Then we have $\varepsilon(\tau_{\zeta, \chi, c}, \psi_K) \equiv \varepsilon(\pi_{\zeta, \chi, c}, \psi_K) \pmod{\mu_{p^e}(\mathbb{C})}$.*

Proof. Let π be the representation of $\mathrm{GL}_n(K)$ corresponding to $\tau_{\zeta, \chi, c}$ by the local Langlands correspondence. By the proof of [Bushnell and Henniart 2014, Proposition 2.2], Propositions 8.6 and 10.5, we have

$$\pi \simeq c\text{-Ind}_{L_{\zeta}^{\times} U_{\mathfrak{J}}^1}^{\mathrm{GL}_n(K)} \Lambda$$

for a character $\Lambda : L_{\zeta}^{\times} U_{\mathfrak{J}}^1 \rightarrow \mathbb{C}^{\times}$ which coincides with $\Lambda_{\zeta, \chi, c}$ on $K^{\times} U_{\mathfrak{J}}^1$. Then, the claim follows from [Bushnell and Henniart 2014, Lemma 2.2(1)], because $L_{\zeta}^{\times} U_{\mathfrak{J}}^1 / (K^{\times} U_{\mathfrak{J}}^1)$ is the cyclic group of order p^e . \square

Proposition 11.6. *We have $\varepsilon(\tau_{\zeta, \chi, c}, \psi_K) = \varepsilon(\pi_{\zeta, \chi, c}, \psi_K)$.*

Proof. By Proposition 3.2 and $\tau_{\zeta, \chi, c} \simeq \mathrm{Ind}_{E_{\zeta}/K} \tau_{n, \zeta, \chi, c}$, it suffices to show that

$$\lambda(E_{\zeta}/K, \psi_K)^{p^e} \varepsilon(\tau_{n, \zeta, \chi, c}, \psi_{E_{\zeta}}) = (-1)^{n-1+\epsilon_0 f} \chi(n') c.$$

By Lemma 10.4, we may assume $\chi = 1$ and $c = 1$. Hence, it suffices to show

$$(11-2) \quad \lambda(E_{\zeta}/K, \psi_K)^{p^e} \varepsilon(\tau_{n, \zeta}, \psi_{E_{\zeta}}) = (-1)^{n-1+\epsilon_0 f}.$$

Assuming that (11-2) is proved for $n = p^e$, we show (11-2) for general n . Let $\tau'_{n, \zeta}$ denote the representation of $W_{E_{\zeta}}$ given by Θ_{ζ} in (2-12) and τ_{p^e} . We put $\psi'_{E_{\zeta}} = n'^{-1} \psi_{E_{\zeta}}$. Applying the result for $n = p^e$ to $E_{\zeta}, \varphi'_{\zeta}$ in place of K, ϖ , we have

$$\varepsilon(\tau'_{n, \zeta}, \psi'_{E_{\zeta}}) = (-1)^{p^e-1+\epsilon'_0 f},$$

where ϵ'_0 denotes ϵ_0 for $n = p^e$. Since $\det \tau'_{n, \zeta}$ is unramified as in the proof of Proposition 8.6, we have

$$(11-3) \quad \varepsilon(\tau'_{n, \zeta}, \psi_{E_{\zeta}}) = \det \tau'_{n, \zeta}(n') \varepsilon(\tau'_{n, \zeta}, \psi'_{E_{\zeta}}) = (-1)^{p^e-1+\epsilon'_0 f}.$$

We note that the inflation of the character in (2-9) by Θ_{ζ} factors through

$$W_{E_{\zeta}} \rightarrow \{\pm 1\} \times \mathbb{Z}, \quad \sigma \mapsto (a_{\sigma}^{(p^e+1)/2}, f n_{\sigma}).$$

If $p \neq 2$, then we have $(n' \varphi'_{\zeta}, -\varphi'_{\zeta})_{E_{\zeta}} = \left(\frac{n'}{q}\right)$, where

$$(\cdot, \cdot)_{E_{\zeta}} : E_{\zeta}^{\times} / (E_{\zeta}^{\times})^2 \times E_{\zeta}^{\times} / (E_{\zeta}^{\times})^2 \rightarrow \{\pm 1\}$$

denotes the Hilbert symbol. Hence, we have

$$(11-4) \quad \frac{\varepsilon(\tau_{n, \zeta}, \psi_{E_{\zeta}})}{\varepsilon(\tau'_{n, \zeta}, \psi_{E_{\zeta}})} = \begin{cases} \left(\frac{n'}{q}\right)^{n-p^e} \left(\left(\frac{n'}{p}\right)^n (-\epsilon(p) \left(\frac{-2}{p}\right))^{n-p^e}\right)^f & \text{if } p \neq 2, \\ (-1)^{\left(\frac{1}{8}n(n-2) - \frac{1}{8}2^e(2^e-2)\right)f} & \text{if } p = 2 \end{cases}$$

by (2-9), Lemmas 4.2 and 10.4. Then we have (11-2) by Lemma 11.1, equations (11-3) and (11-4).

Therefore, we may assume that $n = p^e$. By Lemmas 11.1 and 11.5, it suffices to show that

$$\varepsilon(\tau_{n,\zeta}, \psi_{E_\zeta})^{N(p^e+1)} = \begin{cases} 1 & \text{if } p \neq 2, \\ (-1)^{1+\epsilon_0 f} & \text{if } p = 2. \end{cases}$$

By Proposition 5.1, we have

$$\varepsilon(\tau_{n,\zeta}, \psi_{E_\zeta})^{N(p^e+1)} = \delta_{T_\zeta^u/E_\zeta}(\text{rsw}(\tau_{n,\zeta}, \psi_{E_\zeta}))^{-1} \lambda(T_\zeta^u/E_\zeta, \psi_{E_\zeta})^{p^e} \varepsilon(\tau_{n,\zeta}|_{W_{T_\zeta^u}}, \psi_{T_\zeta^u}).$$

By this, Lemmas 11.1 and 11.4, it suffices to show that

$$\varepsilon(\tau_{n,\zeta}|_{W_{T_\zeta^u}}, \psi_{T_\zeta^u}) = \begin{cases} -(-1)^{\frac{1}{4}(p-1)fN} & \text{if } p \neq 2, \\ (-1)^{1+\epsilon_0 f} \left(\frac{q}{p^e+1}\right) & \text{if } p = 2. \end{cases}$$

This follows from Lemma 11.2 and Proposition 11.7. \square

We set $\varpi_{M_\zeta^u} = \delta_\zeta^{-1}$.

Proposition 11.7. *Assume that $n = p^e$. Then we have*

$$\varepsilon(\xi_{n,\zeta}, \psi_{M_\zeta^u}) = \begin{cases} -(-1)^{\frac{1}{4}(p-1)fN} \left(\frac{r}{kN}\right) & \text{if } p \neq 2, \\ (-1)^{1+\epsilon_0 f} & \text{if } p = 2. \end{cases}$$

Proof. First, we reduce the problem to the positive characteristic case. Assume that $\text{ch } K = 0$. Take a positive characteristic local field $K_{(p)}$ whose residue field is isomorphic to k . We define $M_{\zeta,(p)}^u$ similarly as M_ζ^u starting from $K_{(p)}$. We use similar notations also for other objects in the positive characteristic side. Then we have the isomorphism

$$\mathcal{O}_{M_{\zeta,(p)}^u}/\mathfrak{p}_{M_{\zeta,(p)}^u}^3 \xrightarrow{\sim} \mathcal{O}_{M_\zeta^u}/\mathfrak{p}_{M_\zeta^u}^3, \quad \xi_0 + \xi_1 \varpi_{M_{\zeta,(p)}^u} + \xi_2 \varpi_{M_{\zeta,(p)}^u}^2 \mapsto \hat{\xi}_0 + \hat{\xi}_1 \varpi_{M_\zeta^u} + \hat{\xi}_2 \varpi_{M_\zeta^u}^2$$

of algebras, where $\xi_1, \xi_2, \xi_3 \in k$. Hence, the problem is reduced to the positive characteristic case by [Deligne 1984, Proposition 3.7.1].

We may assume $K = \mathbb{F}_q((t))$. We put $K_{(1)} = \mathbb{F}_p((t))$. We define $M_{\zeta,(1)}^u$ similarly as M_ζ^u starting from $K_{(1)}$. We use similar notations also for other objects in the $K_{(1)}$ -case. We put $f' = [M_\zeta^u : M_{\zeta,(1)}^u]$. We have

$$\delta_{M_\zeta^u/M_{\zeta,(1)}^u}(\text{rsw}(\xi_{n,\zeta,(1)}, \psi_{M_{\zeta,(1)}^u})) = (-1)^{f'-1}$$

by Lemma 10.3. We have $\lambda(M_\zeta^u/M_{\zeta,(1)}^u, \psi_{M_{\zeta,(1)}^u}) = 1$, since the level of $\psi_{M_{\zeta,(1)}^u}$ is $2 - p^e$ by Lemma 11.3. Then, we obtain

$$(11-5) \quad \varepsilon(\xi_{n,\zeta}, \psi_{M_\zeta^u}) = (-1)^{f'-1} \varepsilon(\xi_{n,\zeta,(1)}, \psi_{M_{\zeta,(1)}^u})^{f'}$$

by Proposition 5.1. By (11-5), the problem is reduced to the case where $f = 1$. In this case, the claim follows from Lemmas 11.11 and 11.16. \square

11B. Special cases. We assume that $n = p^e$, $\text{ch } K = p$ and $f = 1$ in this subsection.

11B1. *Odd case.* Assume that $p \neq 2$.

Lemma 11.8. *We have $\psi_{M_\zeta^u}(-\delta_\zeta^{p^e+1}(1+x\varpi_{M_\zeta^u})) = 1$ for $x \in k_N$.*

Proof. For $x \in k_N$, we have

$$\psi_{M_\zeta^u}(-\delta_\zeta^{p^e+1}(1+x\varpi_{M_\zeta^u})) = \psi_{M_\zeta^u}(-(r^{-1}\delta_\zeta - \alpha_\zeta^{-1})(\delta_\zeta + x)) = \psi_{M_\zeta^u}(-r^{-1}\delta_\zeta^2),$$

because $\text{Tr}_{M_\zeta^u/T_\zeta^u}(\delta_\zeta) = 0$ and $[M_\zeta^u : T_\zeta^u] = p^e$. If $p^e \neq 3$, then we have the claim, because $\text{Tr}_{M_\zeta^u/T_\zeta^u}(\delta_\zeta^2) = 0$.

We assume that $p^e = 3$. Then we have

$$\psi_{M_\zeta^u}(-r^{-1}\delta_\zeta^2) = \psi_{T_\zeta^u}(-2r^{-2}) = \psi_0(\text{Tr}_{k_N/\mathbb{F}_p}(-2r^{-2})) = \psi_0(-N \text{Tr}_{\mathbb{F}_{p^2}/\mathbb{F}_p}(r^{-2})) = 1$$

by $\text{Tr}_{M_\zeta^u/T_\zeta^u}(\delta_\zeta^2) = 2r^{-1}$ and $r^4 = -1$. \square

Let θ_ζ be as in (9-10).

Lemma 11.9. *We have*

$$\text{Nr}_{N_\zeta^u/M_\zeta^u}(1+x\theta_\zeta^{(p-1)/2}\varpi_{M_\zeta^u}) \equiv 1 + (-2r)^{(1-p)/2}x^p\varpi_{M_\zeta^u} + \frac{x^2}{2}\varpi_{M_\zeta^u}^2 \pmod{\mathfrak{p}_{M_\zeta^u}^3}$$

for $x \in k_N$.

Proof. We put $T = 1 + x\theta_\zeta^{(p-1)/2}\varpi_{M_\zeta^u}$. By $\theta_\zeta^p - \theta_\zeta = (-2r)^{-1}\delta_\zeta^2$ in (9-11), we have

$$\theta_\zeta = -\frac{1}{2r}\delta_\zeta^2((x^{-1}(T-1)\delta_\zeta)^2 - 1)^{-1}.$$

Substituting this to $x^{-1}(T-1)\delta_\zeta = \theta_\zeta^{(p-1)/2}$, we have

$$(T^2 - 2T + 1 - x^2\varpi_{M_\zeta^u}^2)^{(p-1)/2}(T-1) - (-2r)^{(1-p)/2}x^p\varpi_{M_\zeta^u} = 0.$$

The claim follows from this. \square

Lemma 11.10. *We have*

$$\sum_{x \in k_N} \xi_{n,\zeta}(1+x\varpi_{M_\zeta^u})^{-1} = -((-1)^{(p-1)/2}p)^{e_0} \left(\frac{r}{k_N}\right).$$

Proof. Let $\xi'_{n,\zeta}$ be as in Section 9C. We note that the left-hand side of the claim does not change even if we replace $\xi_{n,\zeta}$ by $\xi'_{n,\zeta}$. We have

$$\begin{aligned} (11-6) \quad \sum_{x \in k_N} \xi'_{n,\zeta}(1+x\varpi_{M_\zeta^u})^{-1} &= \sum_{x \in k_N} \xi'_{n,\zeta}(1+(-2r)^{(1-p)/2}x^p\varpi_{M_\zeta^u})^{-1} \\ &= \sum_{x \in k_N} \xi'_{n,\zeta} \left(1 - \frac{x^2}{2}\varpi_{M_\zeta^u}^2\right)^{-1} \\ &= \sum_{x \in k_N} \psi_{M_\zeta^u} \left(-\frac{x^2}{2}\delta_\zeta^{p^e-1}\right), \end{aligned}$$

where we use Lemmas 9.7 and 11.9 at the second equality and (10-4) at the last equality. The last expression in (11-6) is equal to

$$\sum_{x \in k_N} \psi_{T_\zeta^u}(- (2r)^{-1} (p^e - 1) x^2) = \sum_{x \in k_N} \psi_0(\text{Tr}_{k_N/\mathbb{F}_p}(rx^2)) = -((-1)^{(p-1)/2} p)^{e_0} \left(\frac{r}{k_N} \right)$$

by (2-16), (8-2), Lemma 11.3 and $N = 2e_0$. \square

Lemma 11.11. *We have $\varepsilon(\xi_{n,\zeta}, \psi_{M_\zeta^u}) = -(-1)^{\frac{1}{2}((p-1)e_0)} \left(\frac{r}{k_N} \right)$.*

Proof. We have

$$\begin{aligned} \varepsilon(\xi_{n,\zeta}, \psi_{M_\zeta^u}) &= p^{-e_0} \sum_{x \in k_N} \xi_{n,\zeta}(-\delta_\zeta^{p^e+1} (1 + x \varpi_{M_\zeta^u}))^{-1} \psi_{M_\zeta^u}(-\delta_\zeta^{p^e+1} (1 + x \varpi_{M_\zeta^u})) \\ &= -(-1)^{\frac{1}{2}((p-1)e_0)} \left(\frac{r}{k_N} \right) \xi_{n,\zeta}(-\delta_\zeta^{p^e+1})^{-1} \end{aligned}$$

by Proposition 5.3(2), Lemmas 11.8 and 11.10. We have

$$\begin{aligned} \xi_{n,\zeta}(-\delta_\zeta^{p^e+1}) &= \xi'_{n,\zeta}(-\delta_\zeta^{p^e+1}) (-1)^{\frac{1}{2}(p-1)\frac{1}{2}(p^e+1)N} \\ &= \xi'_{n,\zeta}(-\delta_\zeta^{p^e+1}) = \xi'_{n,\zeta}(-(-2r)^{(p^e+1)\frac{1}{2}(1-p)}) = 1, \end{aligned}$$

where we use

$$\text{Nr}_{N_\zeta^u/M_\zeta^u}(\theta_\zeta^{(p-1)/2} \varpi_{M_\zeta^u}) = (-2r)^{(1-p)/2} \varpi_{M_\zeta^u}$$

at the third equality and $k_N^\times \subset \text{Nr}_{N_\zeta^u/M_\zeta^u}((N_\zeta^u)^\times)$ at the last equality. Thus, we have the claim. \square

11B2. Even case. Assume that $p = 2$.

Lemma 11.12. *We have $\text{Tr}_{M_\zeta^u/K}(\delta_\zeta^{2^e+1}) = 0$ and*

$$\text{Tr}_{M_\zeta^u/K}(\delta_\zeta^{2^e}) = \begin{cases} 1 & \text{if } e = 1, \\ 0 & \text{if } e \geq 2. \end{cases}$$

Proof. These follow from $\delta_\zeta^{2^e} - \delta_\zeta = \alpha_\zeta^{-1} + 1$. \square

Lemma 11.13. *We have $\text{Nr}_{N_\zeta^u/M_\zeta^u}(\theta_\zeta \delta_\zeta^{-1}) = \delta_\zeta^{-1}$.*

Proof. We have $\text{Nr}_{N_\zeta^u/M_\zeta^u}(\theta_\zeta) = \delta_\zeta^3$ by $\theta_\zeta^2 - \theta_\zeta = \delta_\zeta \eta_\zeta$ and $\eta_\zeta^2 - \eta_\zeta = \delta_\zeta$. The claim follows from this. \square

Let $\sigma_0 \in \text{Gal}(N_\zeta^u/M_\zeta^u)$ be a generator of $\text{Gal}(N_\zeta^u/M_\zeta^u)$ determined by

$$\sigma_0(\eta_\zeta) - \eta_\zeta = 1 \quad \text{and} \quad \sigma_0(\theta_\zeta) - \theta_\zeta = \eta_\zeta.$$

Lemma 11.14. *Let $\iota_{n,\zeta} : \text{Gal}(N_\zeta^u/M_\zeta^u) \rightarrow \mathbb{C}^\times$ be the homomorphism induced by $\xi'_{n,\zeta}$ (see Lemma 9.9). Then we have $\iota_{n,\zeta}(\sigma_0) = -\sqrt{-1}$.*

Proof. Let s, t be as in (9-4). We take $\sigma \in I_{M_\zeta^u}$ such that $\Theta_\zeta(\sigma) = ((1, t, s^2), 0)$. Recall that

$$\phi'((1, t, s^2)) = \bar{g} \left(1, s^2 + \sum_{0 \leq i < j \leq e-1} t^{2^i+2^j} \right) \in R_0''$$

is a generator. Then it suffices to show that $\sigma(\eta_\zeta) - \eta_\zeta = 1$ and $\sigma(\theta_\zeta) - \theta_\zeta = \eta_\zeta$. We can check the first equality easily. To show the second equality, it suffices to show that $\sigma(\theta'_\zeta) - \theta'_\zeta = \eta_\zeta^{2^{e-1}}$. By (2-11), we have

$$\sigma(\gamma_\zeta) - \gamma_\zeta \equiv s^2 + \sum_{i=0}^{e-1} (t\beta_\zeta + t^2)^{2^i} \pmod{\mathfrak{p}_{N_\zeta^u}}.$$

By $t = \sigma(\beta_\zeta) - \beta_\zeta$, $\text{Tr}_{\mathbb{F}_{2^e}/\mathbb{F}_2}(t) = 1$ and (9-14), we have

$$\sigma(\gamma'_\zeta) - \gamma'_\zeta = \eta_\zeta - b_1 + s^2 + \sum_{0 \leq i \leq j \leq e-1} t^{2^i+2^j} \pmod{\mathfrak{p}_{N_\zeta^u}}.$$

Hence, by (9-15) and (9-19), we have

$$\sigma(\theta'_\zeta) - \theta'_\zeta \equiv \sum_{i=0}^{2^e-1} d_i \eta_\zeta^i \pmod{\mathfrak{p}_{N_\zeta^u}}$$

with some $d_i \in k^{\text{ac}}$. By (9-19), we have

$$\sum_{i=0}^{e-1} (t(\sigma(\gamma'_\zeta) - \gamma'_\zeta))^{2^i} = \sigma(\gamma'_\zeta) - \gamma'_\zeta + \sum_{1 \leq i \leq j \leq e-1} t^{2^i+2^j} + \sum_{0 \leq i < j \leq e-1} t^{2^j} (\delta_\zeta - s)^{2^i}.$$

Therefore, again by (9-15) and (9-19), we have

$$d_0 = b_1 + s^2 + \sum_{0 \leq i \leq j \leq e-1} t^{2^i+2^j} + \sum_{1 \leq i \leq j \leq e-1} t^{2^i+2^j} + b_1^2 = s + s^2 + t = 0.$$

This implies $\sigma(\theta'_\zeta) - \theta'_\zeta = \eta_\zeta^{2^{e-1}}$, since we know that $\sigma(\theta'_\zeta) - \theta'_\zeta - \eta_\zeta^{2^{e-1}} \in \mathbb{F}_2$ by Lemma 9.8 and $\sigma(\eta_\zeta) - \eta_\zeta = 1$. \square

Lemma 11.15. *We have*

$$\varepsilon(\xi'_{n,\zeta}, \psi_{M_\zeta^u}) = \begin{cases} \frac{1+\sqrt{-1}}{\sqrt{2}} & \text{if } e = 1, \\ \frac{1-\sqrt{-1}}{\sqrt{2}} & \text{if } e \geq 2. \end{cases}$$

Proof. By Proposition 5.3, equation (10-4), Lemmas 11.3, 11.12 and 11.13, we have

$$(11-7) \quad \varepsilon(\xi'_{n,\zeta}, \psi_{M_\zeta^u}) = 2^{-1/2} \sum_{x \in \mathbb{F}_2} \xi'_{n,\zeta}(\delta_\zeta^{2^e+1}(1+x\delta_\zeta^{-1}))^{-1} \psi_{M_\zeta^u}(\delta_\zeta^{2^e+1}(1+x\delta_\zeta^{-1})) \\ = \begin{cases} 2^{-1/2}(1-\xi'_{n,\zeta}(1+\delta_\zeta^{-1})^{-1}) & \text{if } e = 1, \\ 2^{-1/2}(1+\xi'_{n,\zeta}(1+\delta_\zeta^{-1})^{-1}) & \text{if } e \geq 2. \end{cases}$$

First assume that $e = 1$. Then we know the equality in the claim modulo $\mu_2(\mathbb{C})$ by [Lemma 11.5](#). Hence it suffices to show the equality of the real parts. This follows from (11-7). In particular, we have $\xi'_{2,\zeta}(1 + \delta_\zeta^{-1}) = \sqrt{-1}$.

Next, we consider the general case. We put $\alpha'_1 = 1/(\delta_\zeta^2 - \delta_\zeta + 1)$ and $\varpi' = \alpha_1'^3$. Let $\xi'_{2,1,\zeta}$ denote $\xi'_{2,1}$ in the case where K and ϖ are replaced by $\mathbb{F}_2((\varpi'))$ and ϖ' . By applying [Lemma 11.14](#) to $\xi'_{n,\zeta}$ and $\xi'_{2,1,\zeta}$, we have $\xi'_{n,\zeta} = \xi'_{2,1,\zeta}$. We know that $\xi'_{2,1,\zeta}(1 + \delta_\zeta^{-1}) = \sqrt{-1}$ by the result in the case $e = 1$. Hence, we have $\xi'_{n,\zeta}(1 + \delta_\zeta^{-1}) = \sqrt{-1}$, which shows the claim. \square

Lemma 11.16. *We have*

$$\varepsilon(\xi_{n,\zeta}, \psi_{M_\zeta^n}) = (-1)^{1+\epsilon_0}.$$

Proof. The epsilon factor $\varepsilon(\xi_{n,\zeta}, \psi_{M_\zeta^n})$ equals $\varepsilon(\xi'_{n,\zeta}, \psi_{M_\zeta^n})$ times

$$\begin{cases} \left(\frac{1+\sqrt{-1}}{\sqrt{2}}\right)^{-3(2^e+1)} & \text{if } e \neq 2, \\ -\left(\frac{1+\sqrt{-1}}{\sqrt{2}}\right)^{-3(2^e+1)} & \text{if } e = 2 \end{cases}$$

by [Lemma 4.2](#), equation (9-12) and [Lemma 10.3](#). Hence, the claim follows from [Lemma 11.15](#). \square

Appendix: Realization in cohomology of Artin–Schreier variety

We realize τ_n in the cohomology of an Artin–Schreier variety. Let v_{n-2} be the quadratic form on $\mathbb{A}_{k^{\text{ac}}}^{n-2}$ defined by

$$v_{n-2}((y_i)_{1 \leq i \leq n-2}) = -\frac{1}{n'} \sum_{1 \leq i < j \leq n-2} y_i y_j.$$

Let X be the smooth affine variety over k^{ac} defined by

$$x^p - x = y^{p^e+1} + v_{n-2}((y_i)_{1 \leq i \leq n-2}) \quad \text{in } \mathbb{A}_{k^{\text{ac}}}^n.$$

We define a right action of $Q \rtimes \mathbb{Z}$ on X by

$$\begin{aligned} (x, y, (y_i)_{1 \leq i \leq n-2})((a, b, c), 0) \\ = \left(x + \sum_{i=0}^{e-1} (by)^{p^i} + c, a(y + b^{p^e}), (a^{(p^e+1)/2} y_i)_{1 \leq i \leq n-2} \right), \\ (x, y, (y_i)_{1 \leq i \leq n-2}) \text{Fr}(1) = (x^p, y^p, (y_i^p)_{1 \leq i \leq n-2}). \end{aligned}$$

We consider the morphism

$$\pi_{n-2} : \mathbb{A}_{k^{\text{ac}}}^{n-1} \rightarrow \mathbb{A}_{k^{\text{ac}}}^1, \quad (y, (y_i)_{1 \leq i \leq n-2}) \mapsto y^{p^e+1} + v_{n-2}((y_i)_{1 \leq i \leq n-2}).$$

Then we have a decomposition

$$(A-1) \quad H_c^{n-1}(X, \overline{\mathbb{Q}}_\ell) \cong \bigoplus_{\psi \in \mathbb{F}_p^\times \setminus \{1\}} H_c^{n-1}(\mathbb{A}_{k^{\text{ac}}}^{n-1}, \pi_{n-2}^* \mathcal{L}_\psi)$$

as $Q \rtimes \mathbb{Z}$ representations. Let ρ_n be the representation over \mathbb{C} of $Q \rtimes \mathbb{Z}$ defined by

$$H_c^{n-1}(\mathbb{A}_{k^{\text{ac}}}^{n-1}, \pi_{n-2}^* \mathcal{L}_{\psi_0}) \left(\frac{n-1}{2} \right)$$

and ι , where $\left(\frac{n-1}{2} \right)$ means the twist by the character $((a, b, c), m) \mapsto p^{m(n-1)/2}$.

Lemma A.1. *If $p \neq 2$, then we have $\det v_{n-2} = -(-2n')^n \in \mathbb{F}_p^\times / (\mathbb{F}_p^\times)^2$.*

Proof. This is an easy calculation. □

Proposition A.2. *We have $\tau_n \simeq \rho_n$.*

Proof. Let Y be the smooth affine variety over k^{ac} defined by

$$x^p - x = v_{n-2}((y_i)_{1 \leq i \leq n-2}) \quad \text{in } \mathbb{A}_{k^{\text{ac}}}^{n-1}.$$

We define a right action of $Q \rtimes \mathbb{Z}$ on Y by

$$\begin{aligned} (x, (y_i)_{1 \leq i \leq n-2})((a, b, c), 0) &= (x, (a^{(p^e+1)/2} y_i)_{1 \leq i \leq n-2}), \\ (x, (y_i)_{1 \leq i \leq n-2}) \text{Fr}(1) &= (x^p, (y_i^p)_{1 \leq i \leq n-2}). \end{aligned}$$

Using the action of $Q \rtimes \mathbb{Z}$ on Y , we can define an action of $Q \rtimes \mathbb{Z}$ on

$$H_c^{n-2}(\mathbb{A}_{k^{\text{ac}}}^{n-2}, v_{n-2}^* \mathcal{L}_{\psi_0}).$$

Then we have

$$(A-2) \quad H_c^{n-1}(\mathbb{A}_{k^{\text{ac}}}^{n-1}, \pi_{n-2}^* \mathcal{L}_{\psi_0}) \cong H_c^1(\mathbb{A}_{k^{\text{ac}}}^1, \pi^* \mathcal{L}_{\psi_0}) \otimes H_c^{n-2}(\mathbb{A}_{k^{\text{ac}}}^{n-2}, v_{n-2}^* \mathcal{L}_{\psi_0})$$

by the Künneth formula, where the isomorphism is compatible with the actions of $Q \rtimes \mathbb{Z}$. By (A-2), it suffices to show the action of $Q \rtimes \mathbb{Z}$ on

$$(A-3) \quad H_c^{n-2}(\mathbb{A}_{k^{\text{ac}}}^{n-2}, v_{n-2}^* \mathcal{L}_{\psi_0}) \left(\frac{n-1}{2} \right)$$

is equal to the character (2-9) via ι .

First, consider the case where $p \neq 2$. The equality of the actions of Q follows from [Deneff and Loeser 1998, Lemma 2.2.3]. We have

$$(A-4) \quad \begin{aligned} (-1)^{n-2} \sum_{\mathbf{y} \in \mathbb{F}_p^{n-2}} \psi_0(v_{n-2}(\mathbf{y})) &= \left(\frac{-1}{p} \right) \left(- \left(\frac{-2n'}{p} \right) \right)^n (\epsilon(p) \sqrt{p})^{n-2} \\ &= \left(-\epsilon(p) \left(\frac{-2n'}{p} \right) \right)^n \sqrt{p}^{n-2} \end{aligned}$$

by Lemma A.1. The equality of the actions of $\text{Fr}(1) \in Q \rtimes \mathbb{Z}$ follows from [Deligne 1977, Sommes trig. Scholie 1.9] and (A-4).

If $p = 2$, the equality follows from [Imai and Tsushima 2020, Proposition 4.5] and $\left(\frac{2}{n-1} \right) = (-1)^{\frac{1}{8}n(n-2)}$. □

Acknowledgements

The authors would like to thank the handling editor Don Blasius and referees for helpful comments and suggestions. This work was supported by Japan Society for the Promotion of Science (JSPS) KAKENHI grants 26707003, 20K03529, 21H00973.

References

- [Abbes and Saito 2010] A. Abbes and T. Saito, “Local Fourier transform and epsilon factors”, *Compos. Math.* **146**:6 (2010), 1507–1551. [MR](#) [Zbl](#)
- [Adrian and Liu 2016] M. Adrian and B. Liu, “Some results on simple supercuspidal representations of $GL_n(F)$ ”, *J. Number Theory* **160** (2016), 117–147. [MR](#) [Zbl](#)
- [Bergé and Martinet 1985] A.-M. Bergé and J. Martinet, “Formes quadratiques et extensions en caractéristique 2”, *Ann. Inst. Fourier (Grenoble)* **35**:2 (1985), 57–77. [MR](#)
- [Bourbaki 1981] N. Bourbaki, *Éléments de mathématique*, Masson, Paris, 1981. [MR](#) [Zbl](#)
- [Bushnell and Fröhlich 1983] C. J. Bushnell and A. Fröhlich, *Gauss sums and p -adic division algebras*, Lecture Notes in Mathematics **987**, Springer, Berlin, 1983. [MR](#) [Zbl](#)
- [Bushnell and Henniart 1999] C. J. Bushnell and G. Henniart, *Local tame lifting for $GL(n)$, II: wildly ramified supercuspidals*, Astérisque **254**, Société Mathématique de France, Paris, 1999. [MR](#) [Zbl](#)
- [Bushnell and Henniart 2005a] C. J. Bushnell and G. Henniart, “The essentially tame local Langlands correspondence, I”, *J. Amer. Math. Soc.* **18**:3 (2005), 685–710. [MR](#) [Zbl](#)
- [Bushnell and Henniart 2005b] C. J. Bushnell and G. Henniart, “The essentially tame local Langlands correspondence, II: totally ramified representations”, *Compos. Math.* **141**:4 (2005), 979–1011. [MR](#) [Zbl](#)
- [Bushnell and Henniart 2006] C. J. Bushnell and G. Henniart, *The local Langlands conjecture for $GL(2)$* , Grundlehren der Mathematischen Wissenschaften **335**, Springer, Berlin, 2006. [MR](#) [Zbl](#)
- [Bushnell and Henniart 2010] C. J. Bushnell and G. Henniart, “The essentially tame local Langlands correspondence, III: the general case”, *Proc. Lond. Math. Soc.* (3) **101**:2 (2010), 497–553. [MR](#) [Zbl](#)
- [Bushnell and Henniart 2014] C. J. Bushnell and G. Henniart, “Langlands parameters for epipelagic representations of GL_n ”, *Math. Ann.* **358**:1-2 (2014), 433–463. [MR](#) [Zbl](#)
- [Bushnell and Kutzko 1993] C. J. Bushnell and P. C. Kutzko, *The admissible dual of $GL(N)$ via compact open subgroups*, Annals of Mathematics Studies **129**, Princeton University Press, Princeton, NJ, 1993. [MR](#) [Zbl](#)
- [Deligne 1973] P. Deligne, “Les constantes des équations fonctionnelles des fonctions L ”, pp. 501–597 in *Modular functions of one variable, II: Proc. Internat. Summer School* (Univ. Antwerp, Antwerp, 1972), Lecture Notes in Math. **349**, Springer, Berlin, 1973. [MR](#) [Zbl](#)
- [Deligne 1976] P. Deligne, “Les constantes locales de l’équation fonctionnelle de la fonction L d’Artin d’une représentation orthogonale”, *Invent. Math.* **35** (1976), 299–316. [MR](#) [Zbl](#)
- [Deligne 1977] P. Deligne, *Cohomologie étale*, Lecture Notes in Mathematics **569**, Springer, Berlin, 1977. [MR](#) [Zbl](#)
- [Deligne 1984] P. Deligne, “Les corps locaux de caractéristique p , limites de corps locaux de caractéristique 0”, pp. 119–157 in *Representations of reductive groups over a local field*, Hermann, Paris, 1984. [MR](#) [Zbl](#)

- [Deligne and Henniart 1981] P. Deligne and G. Henniart, “Sur la variation, par torsion, des constantes locales d’équations fonctionnelles de fonctions L ”, *Invent. Math.* **64**:1 (1981), 89–118. [MR](#) [Zbl](#)
- [Denef and Loeser 1998] J. Denef and F. Loeser, “Character sums associated to finite Coxeter groups”, *Trans. Amer. Math. Soc.* **350**:12 (1998), 5047–5066. [MR](#)
- [Gallagher 1965] P. X. Gallagher, “Determinants of representations of finite groups”, *Abh. Math. Sem. Univ. Hamburg* **28** (1965), 162–167. [MR](#) [Zbl](#)
- [Godement and Jacquet 1972] R. Godement and H. Jacquet, *Zeta functions of simple algebras*, Lecture Notes in Mathematics **260**, Springer, Berlin, 1972. [MR](#) [Zbl](#)
- [Gross and Reeder 2010] B. H. Gross and M. Reeder, “Arithmetic invariants of discrete Langlands parameters”, *Duke Math. J.* **154**:3 (2010), 431–508. [MR](#) [Zbl](#)
- [Harris and Taylor 2001] M. Harris and R. Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies **151**, Princeton University Press, Princeton, NJ, 2001. [MR](#) [Zbl](#)
- [Henniart 1984] G. Henniart, “Galois ε -factors modulo roots of unity”, *Invent. Math.* **78**:1 (1984), 117–126. [MR](#) [Zbl](#)
- [Imai and Tsushima 2017] N. Imai and T. Tsushima, “Stable models of Lubin–Tate curves with level three”, *Nagoya Math. J.* **225** (2017), 100–151. [MR](#) [Zbl](#)
- [Imai and Tsushima 2018] N. Imai and T. Tsushima, “Local Jacquet–Langlands correspondences for simple supercuspidal representations”, *Kyoto J. Math.* **58**:3 (2018), 623–638. [MR](#) [Zbl](#)
- [Imai and Tsushima 2020] N. Imai and T. Tsushima, “Affinoids in the Lubin–Tate perfectoid space and simple supercuspidal representations, I: tame case”, *Int. Math. Res. Not.* **2020**:22 (2020), 8251–8291. [MR](#) [Zbl](#)
- [Imai and Tsushima 2021] N. Imai and T. Tsushima, “Affinoids in the Lubin–Tate perfectoid space and simple supercuspidal representations II: wild case”, *Math. Ann.* **380**:1-2 (2021), 751–788. [MR](#) [Zbl](#)
- [Imai and Tsushima 2022] N. Imai and T. Tsushima, “Geometric realization of the local Langlands correspondence for representations of conductor three”, *Publ. Res. Inst. Math. Sci.* **58**:1 (2022), 49–77. [MR](#) [Zbl](#)
- [Imai and Tsushima 2023] N. Imai and T. Tsushima, “Geometric construction of Heisenberg–Weil representations for finite unitary groups and Howe correspondences”, *Eur. J. Math.* **9**:2 (2023), art. id. 31. [MR](#) [Zbl](#)
- [Kaletha 2015] T. Kaletha, “Epipelagic L -packets and rectifying characters”, *Invent. Math.* **202**:1 (2015), 1–89. [MR](#) [Zbl](#)
- [Koch 1977] H. Koch, “Classification of the primitive representations of the Galois group of local fields”, *Invent. Math.* **40**:2 (1977), 195–216. [MR](#) [Zbl](#)
- [Laumon 1987] G. Laumon, “Transformation de Fourier, constantes d’équations fonctionnelles et conjecture de Weil”, *Inst. Hautes Études Sci. Publ. Math.* **65** (1987), 131–210. [MR](#) [Zbl](#)
- [Laumon et al. 1993] G. Laumon, M. Rapoport, and U. Stuhler, “ \mathcal{D} -elliptic sheaves and the Langlands correspondence”, *Invent. Math.* **113**:2 (1993), 217–338. [MR](#) [Zbl](#)
- [Reeder and Yu 2014] M. Reeder and J.-K. Yu, “Epipelagic representations and invariant theory”, *J. Amer. Math. Soc.* **27**:2 (2014), 437–477. [MR](#) [Zbl](#)
- [Saito 1995] T. Saito, “Local constant of $\text{Ind}_K^L 1$ ”, *Comment. Math. Helv.* **70**:4 (1995), 507–515. [MR](#) [Zbl](#)
- [Serre 1968] J.-P. Serre, *Corps locaux*, 2nd ed., Publications de l’Université de Nancago **8**, Hermann, Paris, 1968. [MR](#) [Zbl](#)

[Serre 1984] J.-P. Serre, “L’invariant de Witt de la forme $\text{Tr}(x^2)$ ”, *Comment. Math. Helv.* **59**:4 (1984), 651–676. [MR](#) [Zbl](#)

[Tate 1979] J. Tate, “Number theoretic background”, pp. 3–26 in *Automorphic forms, representations and L-functions* (Oregon State Univ., Corvallis, Ore., 1977), Proc. Sympos. Pure Math., II **33**, Amer. Math. Soc., Providence, RI, 1979. [MR](#) [Zbl](#)

Received November 14, 2023.

NAOKI IMAI
GRADUATE SCHOOL OF MATHEMATICAL SCIENCES
THE UNIVERSITY OF TOKYO
TOKYO
JAPAN
naoki@ms.u-tokyo.ac.jp

TAKAHIRO TSUSHIMA
DEPARTMENT OF MATHEMATICS AND INFORMATICS
FACULTY OF SCIENCE
CHIBA UNIVERSITY
CHIBA
JAPAN
tsushima@math.s.chiba-u.ac.jp

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2023 is US \$605/year for the electronic version, and \$820/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 326 No. 1 September 2023

Spin Lefschetz fibrations are abundant	1
MIHAIL ARABADJI and R. İNANÇ BAYKUR	
Some arithmetical properties of convergents to algebraic numbers	17
YANN BUGEAUD and KHOA D. NGUYEN	
Local Galois representations of Swan conductor one	37
NAOKI IMAI and TAKAHIRO TSUSHIMA	
Divisors of Fourier coefficients of two newforms	85
ARVIND KUMAR and MONI KUMARI	
Desingularizations of quiver Grassmannians for the equioriented cycle quiver	109
ALEXANDER PÜTZ and MARKUS REINEKE	
Varieties of chord diagrams, braid group cohomology and degeneration of equality conditions	135
VICTOR A. VASSILIEV	
Positively curved Finsler metrics on vector bundles, II	161
KUANG-RU WU	