

*Pacific
Journal of
Mathematics*

**MORE PROPERTIES OF
OPTIMAL POLYNOMIAL APPROXIMANTS
IN HARDY SPACES**

RAYMOND CHENG AND CHRISTOPHER FELDER

Volume 327 No. 2

December 2023

MORE PROPERTIES OF OPTIMAL POLYNOMIAL APPROXIMANTS IN HARDY SPACES

RAYMOND CHENG AND CHRISTOPHER FELDER

We study optimal polynomial approximants (OPAs) in the classical Hardy spaces on the unit disk, H^p ($1 < p < \infty$). For fixed $f \in H^p$ and $n \in \mathbb{N}$, the OPA of degree n associated to f is the polynomial which minimizes the quantity $\|qf - 1\|_p$ over all complex polynomials q of degree less than or equal to n . We begin with some examples which illustrate, when $p \neq 2$, how the Banach space geometry makes the above minimization problem interesting. We then weave through various results concerning limits and roots of these polynomials, including results which show that OPAs can be witnessed as solutions of certain fixed-point problems. Finally, using duality arguments, we provide several bounds concerning the error incurred in the OPA approximation.

1. Introduction	267
2. Preliminaries and geometric oddities	269
3. Limits and continuity	274
4. Error bounds and duality arguments	285
References	294

1. Introduction

This paper concerns a minimization problem in classical Hardy spaces on the unit disk \mathbb{D} ,

$$H^p := \left\{ f \in \text{Hol}(\mathbb{D}) : \sup_{0 \leq r < 1} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta < \infty \right\},$$

where $\text{Hol}(\mathbb{D})$ denotes the collection of holomorphic functions on \mathbb{D} . As is standard, for $1 \leq p < \infty$, we define the norm of $f \in H^p$ as

$$\|f\|_p := \left(\sup_{0 \leq r < 1} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta \right)^{1/p}.$$

MSC2020: primary 30E10; secondary 46E30.

Keywords: optimal polynomial approximant, Pythagorean inequality, duality, fixed point.

When $p = \infty$, we have the set of bounded analytic functions

$$H^\infty := \{f \in \text{Hol}(\mathbb{D}) : \sup_{z \in \mathbb{D}} |f(z)| < \infty\},$$

with corresponding norm

$$\|f\|_\infty := \sup_{z \in \mathbb{D}} |f(z)|.$$

We will frequently view these spaces as subspaces of the Lebesgue spaces $L^p := L^p(\mathbb{T}, dm)$, where dm is normalized Lebesgue measure on the unit circle \mathbb{T} .

Our main objects of study are *optimal polynomial approximants* (OPAs) in Hardy spaces; these are solutions to the minimization problem

$$\inf_{q \in \mathcal{P}_n} \|qf - 1\|_p,$$

where $f \in H^p$ and \mathcal{P}_n is the set of complex polynomials of degree less than or equal to n . We point out that the infimum above is actually a minimum. In our context, this means the problem of finding a degree n OPA can be restated as finding the solution to

$$\inf_{h \in \mathcal{P}_n} \|h - 1\|_p,$$

which is given by the *metric projection* of 1 on the subspace \mathcal{P}_n . A priori, the minimizing argument may not be unique. However, when $1 < p < \infty$, it is well known that there is, in fact, a unique minimizing polynomial due to the uniform convexity of the space; for in a such a space, any closed subspace enjoys a unique nearest-point property. When $p \neq 2$, the projection is *nonlinear*, which starkly contrasts with the Hilbert space setting.

For $p \neq 2$, this problem was originally studied by Centner [10], and considered again in an additional paper by Centner and the authors [11]. We will give some background now, but point the reader to [10; 11] for more thorough exposition, and to [5; 6; 7; 8; 9; 10; 15; 17; 18] for relevant work on OPAs in various Hilbert spaces.

When $p = 2$, this problem was first studied by engineers in work related to digital filter design. The problem reemerged later as a potential way to study cyclic vectors for the forward shift (see [4] for historical discussion). This renewed interest is evidenced by many papers over the last decade (again, see [4], as well as [1; 3] for recent results in the weighted and noncommutative settings, respectively). Other than the work in [14], these results concern only Hilbert spaces, where the geometry makes computation of OPAs an explicit (but nontrivial!) linear algebra exercise. For example, in H^2 , the coefficients (say, a_0, \dots, a_n) of the OPA of degree n associated to a function $f \in H^2$ can be found via the linear system

$$(1.0.1) \quad \langle (S^j f, S^k f)_{H^2} \rangle_{0 \leq j, k \leq n} (a_0, \dots, a_n)^T = (\overline{f(0)}, 0, \dots, 0)^T,$$

where S is the forward shift operator, given by $f(z) \mapsto zf(z)$ (see, e.g., [15, Theorem 2.1]).

In the Banach space setting (for example, H^p , $p \neq 2$), there is not a direct analogue of this exercise, and the nonlinearity of the metric projection makes explicit calculation of OPAs a highly nontrivial task. In the next section, we will state precisely the definition of optimal polynomial approximant. Before moving there, let us give an outline of the paper:

- Section 2 will formally introduce the OPA problem, give some background information concerning the geometry of Banach spaces, and provide some examples illustrating how this geometry differs from that of Hilbert spaces.
- The results of Section 3 are broken into three parts:
 - convergence of OPAs under variance of the parameters of the OPA problem (e.g., n , p , and f),
 - the location of roots of OPAs,
 - constant and linear OPAs as solutions to a fixed-point problem.
- Using duality, Section 4 establishes various bounds for the error $\|qf - 1\|_p$.

2. Preliminaries and geometric oddities

We begin here by providing some background material concerning the geometry of Banach spaces, followed by several examples in H^p which illustrate some oddities that arise when $p \neq 2$.

Let \mathbf{x} and \mathbf{y} be vectors belonging to a normed linear space \mathcal{X} . We say that \mathbf{x} is *orthogonal* to \mathbf{y} in the Birkhoff–James sense [2; 16] if

$$(2.0.1) \quad \|\mathbf{x} + \beta\mathbf{y}\|_{\mathcal{X}} \geq \|\mathbf{x}\|_{\mathcal{X}}$$

for all scalars β . In this situation we write $\mathbf{x} \perp_{\mathcal{X}} \mathbf{y}$. In the case $\mathcal{X} = L^p$, let us write \perp_p instead of \perp_{L^p} , and similarly for $\mathcal{X} = H^p$.

For $1 < p < \infty$, there is also a function-theoretic test for p -orthogonality, which we note now.

Theorem 2.0.2 (James [16]). *Suppose $1 < p < \infty$. Then for f and g belonging to L^p , we have*

$$(2.0.3) \quad f \perp_p g \iff \int_{\mathbb{T}} |f|^{p-2} \bar{f} g \, dm = 0,$$

where any occurrence of “ $|0|^{p-2}0$ ” in the integrand is interpreted as zero.

In light of (2.0.3) we define, for a measurable function f and any $s > 0$,

$$(2.0.4) \quad f^{(s)} := |f|^{s-1} \bar{f}.$$

If $f \in L^p$, then $f^{(p-1)} \in L^q$, where q is the classical Hölder conjugate to p , satisfying $\frac{1}{p} + \frac{1}{q} = 1$. For $g \in L^p$ and $f \in L^q$, we use the standard notation for the dual pairing

$$\langle f, g \rangle = \int_{\mathbb{T}} f \bar{g} \, dm,$$

and from (2.0.3), we have

$$(2.0.5) \quad f \perp_p g \iff \langle g, f^{(p-1)} \rangle = 0.$$

Consequently, the relation \perp_p is linear in its second argument when $1 < p < \infty$, and it then makes sense to speak of a vector being orthogonal to a subspace. We use this now to formally define OPAs.

Definition 2.0.6 (OPA). Let $1 < p < \infty$ and let $f \in H^p \setminus \{0\}$. Given a nonnegative integer n , the n -th optimal polynomial approximant to $1/f$ in H^p is the polynomial solving the minimization problem

$$\min_{q \in \mathcal{P}_n} \|qf - 1\|_p,$$

where \mathcal{P}_n is the set of complex polynomials of degree less than or equal to n . This polynomial exists, is unique, and will be denoted by

$$q_{n,p}[f].$$

Given f, n and p , we refer to the problem of finding the corresponding OPA as the *OPA problem*.

Considering previous discussion on the metric projection, it is immediate that

$$1 - q_{n,p}[f]f \perp_p \bigvee \{f, zf, z^2f, \dots, z^n f\}.$$

We will use the notation $z^k f$ and $S^k f$ interchangeably if there is no risk of confusion. We will also use the notation $[f]_p$ to denote the closure of $\bigvee \{f, zf, z^2f, z^3f, \dots\}$ in H^p , i.e.,

$$[f]_p := \overline{\bigvee \{f, zf, z^2f, z^3f, \dots\}}^{H^p}.$$

In order to avoid trivialities, we will also often ask that $f(0) \neq 0$; in the case that $f(0) = 0$, this is equivalent to $1 \perp_p f$, and so the metric projection of 1 onto $f \mathcal{P}_n$ is identically zero.

In connection with Birkhoff–James orthogonality, there is a version of the Pythagorean theorem for L^p . This theorem takes the form of a family of inequalities relating the lengths of orthogonal vectors with that of their sum [12, Corollary 3.4].

Theorem 2.0.7. *Suppose that $x \perp_p y$ in L^p . If $p \in (1, 2]$, then*

$$\|x + y\|_p^p \leq \|x\|_p^p + \frac{1}{2^{p-1} - 1} \|y\|_p^p,$$

$$\|x + y\|_p^2 \geq \|x\|_p^2 + (p - 1)\|y\|_p^2.$$

If $p \in [2, \infty)$, then

$$\|x + y\|_p^p \geq \|x\|_p^p + \frac{1}{2^{p-1} - 1} \|y\|_p^p,$$

$$\|x + y\|_p^2 \leq \|x\|_p^2 + (p - 1)\|y\|_p^2.$$

These Pythagorean inequalities enable us to obtain bounds and estimates when $p \neq 2$, in lieu of exact calculations possible in the Hilbert space case.

The following examples illustrate some of the ways the geometry of H^p ($p \neq 2$) can run counterintuitive to experience in Hilbert space. Although these examples may not be immediately surprising to the Banach space enthusiast, we relay them for the general functional analyst, especially working in linear approximation problems, as interesting observations related to natural geometric questions.

Example 2.0.8. In a Hilbert space, an orthogonal projection is always a contraction. However, when $p \neq 2$, the norm of the metric projection of a vector can exceed the length of the vector itself.

Consider the linear OPA for $f(z) = 1 + 0.5z$ in H^4 . Numerically, we find that $Q(z) := q_{1,4}[f] \approx 0.9771018 - 0.4339644z$, and thus

$$\|Qf\|_4^4 \approx 1.10294 > 1.$$

Example 2.0.9. In H^2 , it is simple to verify that if $F(0) = 0$, then, for $c \in \mathbb{C}$, the quantity

$$\|c + F\|_2$$

is minimized when $c = 0$. However, this is not the case when $p \neq 2$.

For example, let $p = 4$ and $F(z) = z + 2z^2$. Note that since F has real coefficients, the value of c which minimizes $\|c + F\|_p$ must also be real, or else uniqueness of nearest points is violated. In turn, we have

$$\|c + F\|_4^4 = 33 + 8c + 20c^2 + c^4.$$

Numerically, this is minimized when $c \approx -0.199209$. In particular, the value of the minimizing argument can be nonzero when $p \neq 2$.

Example 2.0.10. Notice for $f \in H^2$ and any $n > 0$, we have

$$1 - q_{n,2}[f]f \perp_2 zf \quad \text{and} \quad 1 \perp_2 zf.$$

Using linearity, we have

$$q_{n,2}[f]f = 1 - (1 - q_{n,2}[f]f) \perp_2 zf.$$

It is natural to ask if this is true when $p \neq 2$, that is, is it true in general that

$$q_{n,p}[f]f \perp_p zf?$$

Let us take $p = 4, n = 1$, and $f(z) = 1 + 2z + z^8$. Numerically, one can find that

$$\int_{\mathbb{T}} (q_{1,p}[f]f)^{(p-1)}zf \, dm \approx 0.00355837,$$

which is nonzero, and so orthogonality fails. This illustrates that for $p \neq 2$, the relation \perp_p fails to be linear in its first argument, and so $q_{n,p}[f]f$ is not necessarily orthogonal to zf .

Example 2.0.11. For $f, g \in H^2$, an exercise shows that if $\|f\|_2 \leq \|g\|_2$, then $\|1 + zf\|_2 \leq \|1 + zg\|_2$. Might a similar statement hold for $p \neq 2$? The following example shows that the answer is *no*.

Let $p = 4$ and choose

$$\begin{aligned} f(z) &= 0.9(1 + z + z^2), \\ g(z) &= -1 - z - z^2. \end{aligned}$$

It is immediate that $\|f\|_4 < \|g\|_4$. However, numerically, we find

$$\begin{aligned} \|1 + zf\|_4^4 &\approx 31.9339, \\ \|1 + zg\|_4^4 &\approx 20.0000. \end{aligned}$$

With these examples in hand, it may now be reasonable to suspect that OPAs have a dependence on p which is highly nonlinear. In general, this is true. Let us demonstrate this with what we describe as the OPA “error”—the quantity $\|q_{n,p}[f]f - 1\|_p$. We use this as motivation in Section 3, where we study the p -dependence of OPAs.

Example 2.0.12. For m a positive integer, consider $f(z) = 1 + 2z^m$. Let us show

$$\|q_{0,2}[f]f - 1\|_2 \neq \|q_{0,4}[f]f - 1\|_4.$$

For $p = 2$ and any scalar $a \in \mathbb{R}$, we have

$$\begin{aligned} \|af - 1\|_2^2 &= \int_0^{2\pi} ([a - 1]^2 + 2ae^{im\theta})([a - 1]^2 + 2ae^{-im\theta}) \frac{d\theta}{2\pi} \\ &= (a - 1)^2 + 4a^2. \end{aligned}$$

Notice that the result of the integration is the extraction of the constant OPA (again, this constant must be real, as the coefficients of f are real). Minimizing this expression (by differentiating with respect to a), we find

$$a = \frac{1}{5}.$$

This yields

$$\|af - 1\|_2 = \sqrt{\left(\frac{1}{5} - 1\right)^2 + 4\left(\frac{1}{5}\right)^2} = \sqrt{\frac{4}{5}} \approx 0.894427.$$

For $p = 4$, we have

$$\|af - 1\|_4^4 = \int_0^{2\pi} (a + 2ae^{im\theta} - 1)^2 (a + 2ae^{-im\theta} - 1)^2 \frac{d\theta}{2\pi},$$

and one may extract the constant term as

$$(a - 1)^4 + 16a^2(a - 1)^2 + 16a^4.$$

Next, one may numerically find that $a \approx 0.121991$ minimizes the above expression.

Finally, this yields

$$\|q_{0,4}[f]f - 1\|_4 \approx \sqrt[4]{0.781388} \approx 0.940192.$$

In addition to the error, one may also notice that the OPAs themselves vary with p . For example, letting $f(z) = 1 + 2z + z^8$, one may numerically find that

$$\begin{aligned} q_{0,4}[f] &\approx 0.0970262, \\ q_{0,6}[f] &\approx 0.0674066. \end{aligned}$$

However, this is not always the case(!). The following example is a generalization of [10, Example 6.1], which showed that the constant OPAs for $f(z) = 1 - z$ do not vary with p .

Example 2.0.13. Let $1 < p < \infty$ and let $f \in H^p$. Let $\lambda \in \mathbb{C}$ and let $h = 1 + f$. Suppose that $|f(e^{it})| = 1$ a.e. on \mathbb{T} and $\overline{f(e^{-it})} = f(e^{it})$ (i.e., the Fourier coefficients of f are real).

Putting $a = q_{0,p}[h]$, we observe

$$\begin{aligned} \|a(1 + f(e^{it})) - 1\|_p &= \|af(e^{it}) + a - 1\|_p \\ &= \|a + \overline{f(e^{it})}(a - 1)\|_p && \text{(multiply by } \bar{f}, \text{ inner)} \\ &= \|a + f(e^{-it})(a - 1)\|_p && \text{(real coefficients)} \\ &= \|a + f(e^{it})(a - 1)\|_p && (t \mapsto -t) \\ &= \|-a + 1 - f(e^{it})(a - 1)\|_p && \text{(multiply by } -1 \text{ and add } 0) \\ &= \|(1 - a)(1 + f(e^{it})) - 1\|_p. \end{aligned}$$

This tells us that $a = \frac{1}{2}$, which is independent of p . Note that if f is any Blaschke product with real zeros, the hypotheses above are satisfied.

3. Limits and continuity

In this section, we provide results which relate to varying the parameters in the OPA problem (i.e., the degree n , the value of p , and the function f). We first deal with this directly. Then, as a corollary, the first subsection below discusses the possible set of roots for OPAs. In the final subsection, we show that OPAs (in certain cases) are solutions to a fixed-point problem. All of these results enable us to make estimates concerning OPAs, knowing that exact computation is difficult when $p \neq 2$.

We begin by recording, without proof, a known result about metric projections (see, e.g., [13, Proposition 4.8.3]).

Proposition 3.0.1. *Let $1 < p < \infty$. Let $f \in H^p$ with $f(0) \neq 0$ and let h be the metric projection of 1 onto $[f]_p$. Then, in norm,*

$$q_{n,p}[f]f \rightarrow h \quad \text{as } n \rightarrow \infty.$$

In the following proposition, for $1 < p < \infty$ and $f \in H^p$, we write $Q_\infty f$ for the metric projection of 1 onto $[f]_p$, understanding that Q_∞ need not be a bona fide H^p function. The next result tells us something about the error incurred by approximating $q_{n,p}[f]$ using the Taylor polynomials of Q_∞ , when the (rather strict) assumption of norm convergence holds.

Proposition 3.0.2. *Let $1 < p < \infty$, and $f \in H^p$. Suppose that the representation*

$$Q_\infty(z)f(z) = \sum_{k=0}^{\infty} \alpha_k z^k f(z)$$

converges in norm. Then there exist a positive constant C and an index N such that

$$\|q_{n,p}[f]f - Q_{(n)}f\|_p^r \leq C \|Q_\infty f - Q_{(n)}f\|_p$$

for all $n \geq N$, where $Q_{(n)}(z) = \sum_{k=0}^n \alpha_k z^k$, and r and K are the applicable Pythagorean parameters.

Proof. From the orthogonality relation

$$1 - q_{n,p}[f]f \perp_p q_{n,p}[f]f - Q_{(n)}f,$$

the Pythagorean inequality gives

$$\|1 - q_{n,p}[f]f\|_p^r + K \|q_{n,p}[f]f - Q_{(n)}f\|_p^r \leq \|1 - Q_{(n)}f\|_p^r.$$

Rearrange and estimate to get

$$\begin{aligned}
 K \|q_{n,p}[f]f - Q_{(n)}f\|_p^r &\leq \|1 - Q_{(n)}f\|_p^r - \|1 - q_{n,p}[f]f\|_p^r \\
 &\leq \|1 - Q_{(n)}f\|_p^r - \|1 - Q_\infty f\|_p^r \\
 &\leq r \|1 - Q_{(n)}f\|_p^{r-1} (\|1 - Q_{(n)}f\|_p - \|1 - Q_\infty f\|_p) \\
 &\leq r \|1 - Q_{(n)}f\|_p^{r-1} \|Q_\infty f - Q_{(n)}f\|_p \\
 &\leq 2r \|1 - Q_\infty f\|_p^{r-1} \|Q_\infty f - Q_{(n)}f\|_p,
 \end{aligned}$$

for n sufficiently large. In the third step we applied the elementary inequality

$$a^r - b^r \leq r a^{r-1} (a - b),$$

for $0 < b < a$ and $r > 1$.

This verifies the claim, with $C = 2r \|1 - Q_\infty f\|_p^{r-1} / K$. □

The previous proposition can be applied when f is any polynomial; we record that result now.

Proposition 3.0.3. *Suppose that $1 < p < \infty$. Let z_1, z_2, \dots, z_N be a sequence of nonzero points of \mathbb{D} , and define*

$$f(z) := \left(1 - \frac{z}{z_1}\right) \left(1 - \frac{z}{z_2}\right) \cdots \left(1 - \frac{z}{z_N}\right).$$

Set $r = 2$ if $1 < p \leq 2$, and set $r = p$ if $2 < p < \infty$. Then the metric projection h of the unit constant function 1 onto the subspace $[f]_p$ of H^p has a norm convergent representation

$$h(z) = \sum_{k=0}^{\infty} b_k z^k f(z),$$

and there exists a positive constant C such that

$$(3.0.4) \quad \left\| q_{n,p}[f]f - \sum_{k=0}^n b_k z^k f \right\|_p^r \leq C \left\| \sum_{k=n+1}^{\infty} b_k z^k f \right\|_p^r$$

for all positive integers n .

We omit the proof here, but note that the metric projection h must vanish at the zeros of f ; in turn, boundedness of the difference-quotient operator, given by

$$(B_w f)(z) = \frac{f(z) - f(w)}{z - w}, \quad z, w \in \mathbb{D},$$

(applied where $f(w) = 0$) then ensures the norm convergent representation.

3.1. Continuity. As discussed earlier, OPAs generally vary with p . We discuss this variance here, first showing that when f is a bounded function, $q_{n,p}[f]$ varies continuously with respect to p .

Lemma 3.1.1. *Let $f \in H^\infty$ with $f(0) \neq 0$ and let $d \in \mathbb{N}$. If $(p_k)_k \subseteq (1, \infty)$ with $p_k \rightarrow p \in (1, \infty)$, then $q_{d,p_k}[f]$ converges to $q_{d,p}[f]$ uniformly as $k \rightarrow \infty$.*

Proof. Let us write

$$f(z) = \sum_{j=0}^\infty f_j z^j,$$

$$q_{d,p}[f](z) = a_0 + a_1 z + a_2 z^2 + \cdots + a_d z^d,$$

$$q_{d,p_k}[f](z) = a_0^{(k)} + a_1^{(k)} z + a_2^{(k)} z^2 + \cdots + a_d^{(k)} z^d.$$

Since $f \in H^\infty$, Hölder yields

$$\left| \int_{\mathbb{T}} f z^{-j} dm \right| \leq \|f\|_\infty$$

for all $j \geq 0$. Hence, all of the coefficients f_k are bounded by $\|f\|_\infty$.

Letting p'_k be the dual exponent of p_k , we observe

$$|a_0^{(k)} f_0 - 1| = \left| \int_{\mathbb{T}} (q_{d,p_k}[f]f - 1) dm \right| \leq \|q_{d,p_k}[f]f - 1\|_{p_k} \|1\|_{p'_k} \leq 1,$$

and so the sequence $\{a_0^{(k)}\}$ is bounded.

Further, since

$$|a_0^{(k)} f_j + a_1^{(k)} f_{j-1} + \cdots + a_j^{(k)} f_0| \leq \left| \int_{\mathbb{T}} (q_{d,p_k}[f]f - 1) z^{-j} dm \right| \leq 1,$$

it follows

$$|a_j^{(k)}| \leq \frac{|a_0^{(k)} f_j + a_1^{(k)} f_{j-1} + \cdots + a_{j-1}^{(k)} f_1| + 1}{|f_0|},$$

for all $k \in \mathbb{N}$ and $1 \leq j \leq d$. That is, $\{a_j^{(k)}\}_{k=1}^\infty$ is also a bounded sequence for $1 \leq j \leq d$. By passing to a subsequence and relabeling, we can assume that $\{q_{n,p_k}[f]\}_{k=1}^\infty$ is a uniformly convergent sequence of polynomials, which converges to some polynomial, say, $A(z) = a_0 + a_1 z + \cdots + a_d z^d \in \mathcal{P}_d$.

Now, for $0 \leq j \leq d$, recall the orthogonality equations

$$\int_{\mathbb{T}} [q_{d,p_k}[f]f - 1]^{(p_k-1)} z^j f dm = 0.$$

Taking $k \rightarrow \infty$ and invoking uniform convergence, we find that

$$\int_{\mathbb{T}} [Af - 1]^{(p_k-1)} z^j f \, dm = 0,$$

for $0 \leq j \leq d$ (the taking of $(p_k - 1)$ powers also being well behaved). By uniqueness of the optimal polynomial, it must be that

$$A(z) = q_{d,p}[f](z).$$

Since every subsequence of the originally given sequence $\{q_{d,p_k}[f]\}_{k=1}^\infty$ has a further subsequence that converges to the same limit $q_{d,p}[f]$, it must be that

$$q_{d,p_k}[f] \rightarrow q_{d,p}[f]$$

uniformly. □

We now present another continuity result—continuity in f . In particular, if $f_k \rightarrow f$ in H^p , then $q_{n,p}[f_k] \rightarrow q_{n,p}[f]$. Before establishing this result, we need a couple of lemmas.

Lemma 3.1.2. *Let $1 < p < \infty$ and $\frac{1}{p} + \frac{1}{q} = 1$. If $\varphi_k \rightarrow \varphi$ in L^p , then $\varphi_k^{(p-1)} \rightarrow \varphi^{(p-1)}$ in L^q .*

Proof. First, we check that

$$\int_{\mathbb{T}} |\varphi^{(p-1)}|^q \, dm = \int_{\mathbb{T}} |\varphi|^{(p-1)q} \, dm = \int_{\mathbb{T}} |\varphi|^p \, dm,$$

and so $\varphi^{(p-1)} \in L^q$; similarly $\varphi_k^{(p-1)} \in L^q$.

Next, we apply the generalized dominated convergence theorem, using the sequential bound

$$|\varphi_k^{(p-1)} - \varphi^{(p-1)}|^q \leq 2^{q-1} (|\varphi_k|^p + |\varphi|^p) \quad \text{a.e.-}dm,$$

with the Carleson–Hunt theorem supplying pointwise convergence almost everywhere. The conclusion is

$$\int_{\mathbb{T}} |\varphi_k^{(p-1)} - \varphi^{(p-1)}|^q \, dm \rightarrow 0,$$

as claimed. □

Below, we use the standard notation $\hat{f}(n)$ to denote the n -th Fourier coefficient of a function $f \in L^p$.

Lemma 3.1.3. *Let $1 < p < \infty$. If $\varphi \in H^p$, then*

$$\|\varphi\|_p^r \geq |\hat{\varphi}(0)|^r + K |\hat{\varphi}(1)|^r + K^2 |\hat{\varphi}(2)|^r + \dots,$$

where r and K are the lower Pythagorean parameters.

Proof. This follows immediately from the orthogonality relations

$$z^k \perp_p z^m H^p \quad \text{for all } m > k \geq 0,$$

and repeated application of the lower Pythagorean inequality. □

We are now prepared to prove the aforementioned result.

Theorem 3.1.4. *Suppose that $1 < p < \infty$ and $n \in \mathbb{N}$. Let $f_k \in H^p$ and let $Q_k := q_{n,p}[f_k]$ for each $k \in \mathbb{N}$. If $f_k \rightarrow f$ in H^p , and $f(0) \neq 0$, then $Q_k \rightarrow Q := q_{n,p}[f]$.*

Proof. Let us first handle the case $n = 1$, and write $Q_k(z) = a_k + b_k z$ for $q_{1,p}[f]$.

Since $f_k(0) \rightarrow f(0)$, and $f(0) \neq 0$, there is no harm in assuming that there exists $c > 0$ such that $|f_k(0)| \geq c$ for all k .

From the relation

$$1 \perp_p z H^p$$

we see that

$$1 \geq \|1 - Q_k f_k\|_p^r \geq |1 - a_k f_k(0)|^r + K \|Q_k f_k - a_k f_k(0)\|_p^r,$$

where r and K are the lower Pythagorean parameters. It follows that

$$1 \geq |1 - a_k f_k(0)|,$$

implying that

$$|a_k| \leq \frac{2}{c}$$

for all k . Thus $\{a_k\}$ is a bounded complex sequence, from which we can extract a convergent subsequence, which for now we relabel as the original sequence.

Next, subharmonicity and the triangle inequality yield

$$c |b_k| \leq |b_k f_k(0)| \leq \|b_k z f_k\|_p \leq \|1 - (a_k + b_k z) f_k\|_p + \|1 - a_k f_k\|_p.$$

The last expression on the right side is uniformly bounded as k varies through \mathbb{N} , and hence $\{b_k\}$ is a bounded sequence. Once again we may draw a convergent subsequence, and relabel it so that

$$Q_k = a_k + b_k z$$

converges uniformly to some $R(z) = a + bz$.

It needs to be shown that $R = Q := q_{1,p}[f]$. For this we rely on the elementary result that if $v_k \rightarrow v$ in a Banach space and $\lambda_k \rightarrow \lambda$ in its dual space, then $\lambda_k(v_k) \rightarrow \lambda(v)$.

We apply this, identifying

$$\begin{aligned} v_k &= f_k, \\ v &= f, \\ \lambda_k(\cdot) &= \int_{\mathbb{T}} (1 - [a_k + b_k z] f_k)^{(p-1)}(\cdot) dm, \\ \lambda(\cdot) &= \int_{\mathbb{T}} (1 - [a + bz] f)^{(p-1)}(\cdot) dm. \end{aligned}$$

Then Lemma 3.1.2 ensures that $\lambda_k \rightarrow \lambda$, as needed.

The conclusion is that $\lambda(v) = \lim_{k \rightarrow \infty} \lambda_k(v_k) = \lim_{k \rightarrow \infty} 0 = 0$, or

$$1 - (a + bz) f \perp_p f.$$

Repeat this argument with the choices

$$v_k = z f_k \quad \text{and} \quad v = z f$$

to see that

$$1 - (a + bz) f \perp_p z f$$

as well. This forces $R(z) = Q(z) = a + bz = q_{1,p}[f](z)$, as claimed.

So far, we only know that there is a subsequence that satisfies the claim. However, we see that every subsequence of the original sequence $\{f_k\}$ has a further subsequence for which the linear OPAs tend to the same limit $a + bz$, the linear OPA from f being unique. This proves that in fact the full sequence $\{a_k + b_k z\}$ converges to $a + bz$.

This verifies the claim when $n = 1$.

More generally, for arbitrary $n \in \mathbb{N}$, let

$$Q_k(z) = q_{n,p}[f](z) = a_0^{(k)} + a_1^{(k)} z + \dots + a_n^{(k)} z^n.$$

From Lemma 3.1.3, we get

$$\begin{aligned} 1 &\geq \|1 - Q_k f_k\|_p^r \\ &\geq |1 - a_0^{(k)} f_0|^r + \sum_{m=1}^{\infty} K^m |a_0^{(k)} f_m + a_1^{(k)} f_{m-1} + \dots + a_m^{(k)} f_0|^r, \end{aligned}$$

which implies

$$\frac{1}{K^{m/r}} \geq |a_0^{(k)} f_m + a_1^{(k)} f_{m-1} + \dots + a_m^{(k)} f_0|$$

for all m .

We know that $a_0^{(k)}$ is bounded in k . It is also easy to see that $|f_j| \leq \|f\|_p$ for all j . If $a_0^{(k)}, a_1^{(k)}, \dots, a_j^{(k)}$ are also bounded in k , then the relation

$$|a_{j+1}^{(k)}| \leq \frac{1}{K^{m/r}|f_0|} + \left| \frac{a_0^{(k)} f_{j+1}}{f_0} + \frac{a_1^{(k)} f_j}{f_0} + \dots + \frac{a_j^{(k)} f_1}{f_0} \right|$$

ensures that $a_{j+1}^{(k)}$ is bounded as well. This proves that all of the coefficients of Q_k are uniformly bounded in k .

Arguing as before, we may find a subsequence from $\{Q_k\}$ that converges uniformly, and the limit must be $q_{n,p}[f]$. In fact, this must be the limit of the original sequence. \square

3.2. Roots of OPAs. As a corollary to the last continuity theorem, we begin this subsection with two results concerning the set of possible OPA roots. Let us first establish some notation.

Definition 3.2.1. For $1 < p < \infty$ and $n \geq 0$, we define the set of possible roots of OPAs of degree n in H^p as

$\Omega_{n,p} := \{w \in \mathbb{C} : \text{there exists } f \in H^p \text{ such that } f(0) \neq 0 \text{ with } q_{n,p}[f](w) = 0\}$, and let

$$\Omega_p := \bigcup_{n \geq 0} \Omega_{n,p}.$$

Note that $\Omega_{0,p} = \emptyset$ for all $p \in (1, \infty)$. We have an immediate proposition concerning these sets.

Proposition 3.2.2. For $1 < p < \infty$ and each $n \geq 1$, we have $\Omega_{n,p} \subseteq \Omega_{1,p}$, and therefore

$$\Omega_p = \Omega_{1,p}.$$

Proof. Suppose $w \in \Omega_{n,p}$ with $q_{n,p}[f](w) = 0$. Put $q_{n,p}[f] = (z - w)\tilde{q}$. Then, by optimality, we have

$$\begin{aligned} \|q_{1,p}[\tilde{q}f]\tilde{q}f - 1\|_p &\leq \|(z - w)\tilde{q}f - 1\|_p \\ &= \|q_{n,p}[f]f - 1\|_p \\ &\leq \|q_{1,p}[\tilde{q}f]\tilde{q}f - 1\|_p, \end{aligned}$$

and we deduce that $q_{1,p}[\tilde{q}f] = q_{n,p}[f]/\tilde{q} = z - w$, which implies that $w \in \Omega_{1,p}$. \square

Presently, we see that the set of OPA roots must contain the set $\mathbb{C} \setminus \bar{\mathbb{D}}$.

Proposition 3.2.3. Let $1 < p < \infty$. If $w \in \mathbb{C} \setminus \bar{\mathbb{D}}$, then there exists $f \in H^p$ such that $q_{1,p}[f]$ has the root w , and so

$$\mathbb{C} \setminus \bar{\mathbb{D}} \subseteq \Omega_p.$$

Proof. Let $w \in \mathbb{C} \setminus \overline{\mathbb{D}}$ and let

$$f(z) := \frac{1}{z - w},$$

which belongs to H^p for all $p \in (1, \infty)$. Further,

$$\|1 - Qf\|_p = 0$$

when

$$Q(z) = z - w.$$

Therefore, it must be that $q_{n,p}[f](z) = z - w$ for all $n \geq 1$. Hence, $w \in \Omega_p$. \square

We now show that this set is connected and symmetric under rotation.

Proposition 3.2.4. *For $1 < p < \infty$, the set Ω_p is rotationally symmetric and connected.*

Proof. We begin by establishing rotational symmetry. Let $f \in H^p$, and suppose $q_{1,p}[f] = a(z - w)$ (by Proposition 3.2.2, it suffices to take the linear OPA). Then, for any γ with $|\gamma| = 1$,

$$\begin{aligned} \|1 - a(z - w)\|_p^p &= \int_0^{2\pi} |1 - a(z - w)f(z)|^p dm(z) \\ &= \int_0^{2\pi} |1 - a(\gamma\xi - w)f(\gamma\xi)|^p |\gamma|^p dm(\xi) \\ &= \int_0^{2\pi} |1 - a(\gamma\xi - w)f(\gamma\xi)|^p dm(\xi) \\ &= \int_0^{2\pi} |1 - (a\gamma)(\xi - \bar{\gamma}w)f(\gamma\xi)|^p dm(\xi). \end{aligned}$$

It must be that $(a\gamma)(z - \bar{\gamma}w)$ is the linear OPA for $f(\gamma z)$, for otherwise, by reversing these steps from

$$\|1 - q_{1,p}[f(\gamma z)]f(\gamma z)\|_p^p$$

we obtain a contradiction.

This shows that if w is an OPA root, then so is $\bar{\gamma}w$ for all γ , $|\gamma| = 1$. That is, the set Ω_p is rotationally symmetric.

Next, suppose that f and g belong to H^p , with real coefficients, and with $f(0) > 0$ and $g(0) > 0$. Let their linear OPA roots be r and R , respectively, where $0 < r < R$. By the continuity of the map $F \mapsto q_{1,p}[F]$, we see that the set of linear OPA roots of the collection of functions $tf + (1 - t)g$, $0 \leq t \leq 1$, must be an interval containing $[r, R]$; this is because the collection of functions is connected, and continuous maps preserve connectivity. Note that $tf(0) + (1 - t)g(0) > 0$ for all t , as required for the linear OPA to be nontrivial. Consequently, Ω_p is path connected, and hence connected. \square

3.3. Fixed-point approach. Again, we mention that computing OPAs when $p \neq 2$ is a challenging task. Here, we explore the idea of OPAs being fixed points of an iterative process. We begin with the degree-zero case and then move to the degree-one case.

Theorem 3.3.1. *Let $2 < p < \infty$, and let $f \in H^p$ be a nonconstant function. Then the degree-zero OPA $q_{0,p}[f]$ is the unique solution to the fixed-point equation*

$$\zeta = \Phi(\zeta),$$

where $\Phi : \mathbb{C} \mapsto \mathbb{C}$ is given by

$$\Phi(\zeta) := \left(\int_{\mathbb{T}} |1 - \zeta f|^{p-2} \bar{f} \, dm \right) \left(\int_{\mathbb{T}} |1 - \zeta f|^{p-2} |f|^2 \, dm \right)^{-1}.$$

For any $\lambda_1 \in \mathbb{C}$, the sequence $\{\lambda_k\}$ given by $\lambda_{k+1} = \Phi(\lambda_k)$ converges to $q_{0,p}[f]$.

Proof. Since $p > 2$, and

$$1 = \frac{2}{p} + \frac{p-2}{p},$$

the parameters $\frac{p}{2}$ and $\frac{p}{p-2}$ are Hölder conjugates of each other. Hence Hölder's inequality gives

$$\begin{aligned} \int_{\mathbb{T}} |1 - \zeta f|^{p-2} |f|^2 \, dm &\leq \left(\int_{\mathbb{T}} |1 - \zeta f|^{(p-2)p/(p-2)} \, dm \right)^{(p-2)/p} \left(\int_{\mathbb{T}} |f|^{2(p/2)} \, dm \right)^{2/p} < \infty. \end{aligned}$$

Furthermore, since f is nonconstant, the integral in the denominator of Φ is nonzero for any value of ζ .

When $\zeta \neq 0$, we have

$$\begin{aligned} |\zeta| \int_{\mathbb{T}} |1 - \zeta f|^{p-2} |f| \, dm &\leq \int_{\mathbb{T}} |1 - \zeta f|^{p-2} (|1 - \zeta f| + 1) \, dm \\ &= \int_{\mathbb{T}} |1 - \zeta f|^{p-1} \, dm + \int_{\mathbb{T}} |1 - \zeta f|^{p-2} \, dm \\ &< \infty; \end{aligned}$$

and, when $\zeta = 0$,

$$\int_{\mathbb{T}} |1 - \zeta f|^{p-2} |f| \, dm = \int_{\mathbb{T}} |1 - 0 \cdot f|^{p-2} |f| \, dm = \int_{\mathbb{T}} |f| \, dm < \infty.$$

This verifies that Φ is well defined for all $\zeta \in \mathbb{C}$.

In fact, Φ is continuous and bounded. Continuity of the numerator and denominator of Φ at any point ζ_0 can be established by a dominated convergence argument,

with respective dominating functions

$$2^{p-2}(1 + C|f|^{p-2})|f| \quad \text{and} \quad 2^{p-2}(1 + C|f|^{p-2})|f|^2,$$

where $C > |\zeta_0|^{p-2}$. Continuity at infinity is established by

$$\begin{aligned} \lim_{\zeta \rightarrow \infty} \frac{\int_{\mathbb{T}} |1 - \zeta f|^{p-2} \bar{f} \, dm}{\int_{\mathbb{T}} |1 - \zeta f|^{p-2} |f|^2 \, dm} &= \lim_{\zeta \rightarrow \infty} \frac{\int_{\mathbb{T}} |1/\zeta - f|^{p-2} \bar{f} \, dm}{\int_{\mathbb{T}} |1/\zeta - f|^{p-2} |f|^2 \, dm} \\ &= \lim_{\zeta \rightarrow \infty} \frac{\int_{\mathbb{T}} |f|^{p-2} \bar{f} \, dm}{\int_{\mathbb{T}} |f|^p \, dm}. \end{aligned}$$

Consequently, Φ is a bounded function. For any choice of $\lambda_1 \in \mathbb{C}$, define $\lambda_{k+1} = \Phi(\lambda_k)$ for all $k = 1, 2, 3, \dots$. The resulting sequence $\{\lambda_k\}$ is a bounded sequence, and must contain a convergent subsequence, $\{\lambda_{n_k}\}$, with $\lambda_k \rightarrow \lambda \in \mathbb{C}$. Continuity ensures that

$$\lambda = \Phi(\lambda),$$

which is to say that

$$\begin{aligned} \lambda \int_{\mathbb{T}} |1 - \lambda f|^{p-2} f \bar{f} \, dm &= \int_{\mathbb{T}} |1 - \lambda f|^{p-2} \bar{f} \, dm \\ 0 &= \int_{\mathbb{T}} |1 - \lambda f|^{p-2} (1 - \lambda f) \bar{f} \, dm, \end{aligned}$$

or $1 - \lambda f \perp_p f$. This shows that $\lambda = q_{0,p}[f]$.

But any subsequence of $\{\lambda_k\}$ must have a further subsequence that converges to the same limit. Thus the sequence $\{\lambda_k\}$ itself must converge to $\lambda = q_{0,p}[f]$. \square

We now discuss the linear case, first recording some notation.

Let a linear polynomial $Q_1(z) = a_1 + b_1 z$ be given and, for $k \geq 1$, let

$$\begin{aligned} (3.3.2) \quad \begin{bmatrix} a_{k+1} \\ b_{k+1} \end{bmatrix} &= \begin{bmatrix} C_k & \overline{D_k} \\ D_k & C_k \end{bmatrix}^{-1} \begin{bmatrix} A_k \\ B_k \end{bmatrix} \\ &= \frac{1}{|C_k|^2 + |D_k|^2} \begin{bmatrix} C_k & -\overline{D_k} \\ -D_k & C_k \end{bmatrix} \begin{bmatrix} A_k \\ B_k \end{bmatrix} \\ &= \frac{1}{|C_k|^2 + |D_k|^2} \begin{bmatrix} c A_k C_k - B_k \overline{D_k} \\ B_k C_k - A_k D_k \end{bmatrix}, \end{aligned}$$

where

$$(3.3.3) \quad \begin{aligned} A_k &= \int_{\mathbb{T}} |1 - Q_k f|^{p-2} \bar{f} \, dm, & B_k &= \int_{\mathbb{T}} |1 - Q_k f|^{p-2} \bar{z} f \, dm, \\ C_k &= \int_{\mathbb{T}} |1 - Q_k f|^{p-2} |f|^2 \, dm, & D_k &= \int_{\mathbb{T}} |1 - Q_k f|^{p-2} \bar{z} |f|^2 \, dm, \end{aligned}$$

and $Q_k(z) = a_k + b_k z$. This determines a sequence of linear polynomials.

Theorem 3.3.4. *Let $2 < p < \infty$, and suppose that $f \in H^p$ is a nonconstant polynomial with $f(0) \neq 0$. If $Q_k(z) = a_k + b_k z$, $k \geq 0$, is the sequence of linear polynomials arising from (3.3.2), then Q_k converges to $q_{1,p}[f]$.*

Proof. If Q_1 is identically zero, then by inspection we see that Q_2 is not the zero polynomial. Thus, by relabeling if necessary, let us assume Q_1 is not identically zero.

By the hypotheses on f , the expression $w - Qf$ is a nonconstant polynomial for any complex number w and linear polynomial Q ; hence $|w - Qf|^{p-2}$ will be integrable on the unit circle.

Consider the expression, with integrals being taken over the circle,

$$\Phi(Q) := \left| \frac{\int |Qf|^{p-2} \bar{f} \, dm \int |Qf|^{p-2} |f|^2 \, dm - \int |Qf|^{p-2} \bar{z} \bar{f} \, dm \int |Qf|^{p-2} \bar{z} |f|^2 \, dm}{\left| \int |Qf|^{p-2} |f|^2 \, dm \right|^2 + \left| \int |Qf|^{p-2} \bar{z} |f|^2 \, dm \right|^2} \right|,$$

as Q varies over set

$$\mathcal{Q} := \{a + bz \in \mathcal{P}_1 : \max\{|a|, |b|\} = 1\}.$$

Under the assumptions on f , the denominator is bounded away from zero. Thus $\Phi(Q)$ is a continuous function on a compact set, and achieves its maximum. In fact, the value of $\Phi(Q)$ is indifferent to rescaling Q , except for multiplying it by zero.

From this we can further deduce that the values of

$$\Psi(Q, w) := \left| \frac{\int |w - Qf|^{p-2} \bar{f} \, dm \int |w - Qf|^{p-2} |f|^2 \, dm - \int |w - Qf|^{p-2} \bar{z} \bar{f} \, dm \int |w - Qf|^{p-2} \bar{z} |f|^2 \, dm}{\left| \int |w - Qf|^{p-2} |f|^2 \, dm \right|^2 + \left| \int |w - Qf|^{p-2} \bar{z} |f|^2 \, dm \right|^2} \right|$$

are uniformly bounded for $Q \in \mathcal{Q}$ and $|w| \leq 1$.

Next, notice that for any nonzero linear polynomial $Q(z) = a + bz$ we have

$$\begin{aligned} \int_{\mathbb{T}} |1 - Qf|^{p-2} \bar{f} \, dm &= \int_{\mathbb{T}} |1 - (a + bz)f|^{p-2} \bar{f} \, dm \\ &= |c|^{p-2} \int_{\mathbb{T}} \left| \frac{1}{c} - \left(\frac{a}{c} + \left[\frac{b}{c} \right] z \right) f \right|^{p-2} \bar{f} \, dm, \end{aligned}$$

where $c := \max\{|a|, |b|\}$. This is to say that the value of A_k in (3.3.3) scales in a simple way with c , with the result that f is multiplied by a member of \mathcal{Q} , and the 1 inside the integrand is replaced by $\frac{1}{c}$. Similar remarks apply to the formulas for B_k , C_k , and D_k .

Consequently, when A_k , B_k , C_k and D_k are assembled together to yield a_{k+1} and b_{k+1} , the scaling factors $|c|^{p-2}$ attached to each integral cancel.

Let us write $c_k := \max\{|a_k|, |b_k|\}$. The above observations establish that $|c_{k+1}|$ is uniformly bounded as k varies over such indices that $|c_k| \geq 1$.

For the other values of k , for which $|c_k| < 1$, the corresponding expressions for $|1 - Q_k f|^{p-2}$ are again uniformly bounded in the obvious way, implying that the resulting c_{k+1} are also uniformly bounded.

This shows that $\{Q_k\}$ is a bounded sequence of linear polynomials, which must therefore have a convergent subsequence. The limit is a linear polynomial Q_∞ , which satisfies the orthogonality conditions for $q_{1,p}[f]$, and hence must be the OPA. Uniqueness of the OPA ensures that, in fact, every subsequence of $\{Q_k\}$ has a further subsequence that converges to $q_{1,p}[f]$. In conclusion, we have

$$\lim_{k \rightarrow \infty} Q_k = q_{1,p}[f]. \quad \square$$

We end this section by noting that Theorems 3.3.1 and 3.3.4 are only established for $2 < p < \infty$, and, in the degree-one case, only for polynomials. It is currently unclear if these results extend to $1 < p < 2$, or if analogous results hold for higher-degree OPAs.

4. Error bounds and duality arguments

The present section is concerned with estimating (both above and below) the quantity $\|q_{n,p}[f]f - 1\|_p$, i.e., the “error” in the optimal polynomial approximation algorithm. We begin by employing some duality methods, first recalling a fundamental result from classical functional analysis, tailored to our setting.

Lemma 4.0.1. *Let $1 < p < \infty$ and $f \in H^p$. For any $n \in \mathbb{N}$, we have*

$$\|q_{n,p}[f]f - 1\|_p = \left[\inf_{\psi \in L^q} \{ \|\psi\|_{L^q} : \psi_0 = 1, \langle z^k f, \psi \rangle = 0 \text{ for all } 0 \leq k \leq n \} \right]^{-1}.$$

Proof. By an elementary duality theorem of functional analysis, with respect to the pairing

$$\langle f, g \rangle = \int_0^{2\pi} f(e^{i\theta}) \overline{g(e^{i\theta})} \frac{d\theta}{2\pi},$$

we have

$$\begin{aligned} & \|q_{n,p}[f]f - 1\|_p \\ &= \inf\{\|Qf - 1\|_p : Q \in \mathcal{P}_n\} \\ &= \text{dist}_{H^p}(1, \mathcal{P}_n f) \\ &= \text{dist}_{L^p}(1, \mathcal{P}_n f) \\ &= \|1\|_{[(\mathcal{P}_n f)^\perp]^*} \\ &= \sup \left\{ \frac{|\langle \psi, 1 \rangle|}{\|\psi\|_{L^q}} : \psi \in (\mathcal{P}_n f)^\perp \setminus \{0\} \right\} \end{aligned}$$

$$\begin{aligned}
 &= \sup \left\{ \frac{|\psi_0|}{\|\psi\|_{L^q}} : \psi \in (\mathcal{P}_n f)^\perp \setminus \{0\} \right\} \\
 &= [\inf\{\|\psi/\psi_0\|_{L^q} : \psi \in (\mathcal{P}_n f)^\perp \setminus \{0\}\}]^{-1} \\
 &= [\inf\{\|\psi\|_{L^q} : \psi \in (\mathcal{P}_n f)^\perp, \psi_0 = 1\}]^{-1} \\
 &= [\inf\{\|\psi\|_{L^q} : \psi \in L^q, \psi_0 = 1, \langle z^k f, \psi \rangle = 0 \text{ for all } 0 \leq k \leq n\}]^{-1}. \quad \square
 \end{aligned}$$

Remark 4.0.2. The reason to move to L^p in the third equality is that the dual of L^p is L^q . If we stick with the norm in H^p , then (caution!) the relevant dual space is the quotient space L^q/H^q , rather than H^q . These spaces are isometrically isomorphic only when $p = 2$.

In the fourth equality, we mean “the norm of the unit constant function, viewed as a bounded linear functional on the annihilator of the subspace spanned by $f \mathcal{P}_n$ ”.

We first apply this duality to provide a lower bound for the OPA error in the case that we are approximating a polynomial with zeros in the disk.

Proposition 4.0.3. *Suppose f is a polynomial*

$$(4.0.4) \quad f(z) = (z - w_1)(z - w_2) \cdots (z - w_d),$$

with the roots being distinct, nonzero, and contained inside \mathbb{D} . Then, for $1 < p < \infty$ and $\lambda := q_{0,p}[f]$, we have

$$(4.0.5) \quad \|1 - \lambda f\|_p \geq 1 - |w_1 w_2 \cdots w_d|.$$

Proof. The space of functions in L^q which annihilate f contains functions of the form

$$c_1 \Lambda_{w_1} + c_2 \Lambda_{w_2} + \cdots + c_d \Lambda_{w_d},$$

where Λ_{w_j} denotes the point evaluation functional (or Szegő kernel) at the point $w_j \in \mathbb{D}$. Thus, by the Lemma 4.0.1, we have

$$\|1 - \lambda f\|_p = [\inf \|\psi\|_q]^{-1},$$

where the infimum is over $\psi \in L^q$ satisfying $\langle f, \psi \rangle = 0$ and $\psi_0 = 1$.

Let

$$B(z) = a \prod_{k=1}^d \frac{w_k - z}{1 - \bar{w}_k z},$$

a constant multiple of the Blaschke product with the same zeros as f . Its numerator has leading term $\pm a z^d$, while the denominator has leading term $\pm \bar{w}_1 w_2 \cdots w_d z^d$ (with matching signs). Thus long division followed by partial fractions expansion

results in an expression of the form

$$B(z) = \frac{a}{w_1 w_2 \cdots w_d} + c_1 \Lambda_{w_1}(z) + c_2 \Lambda_{w_2}(z) + \cdots + c_d \Lambda_{w_d}(z).$$

Evaluating this equation at $z = 0$ tells us that

$$a w_1 w_2 \cdots w_d = \frac{a}{w_1 w_2 \cdots w_d} + c_1 + c_2 + \cdots + c_d.$$

This suggests making the specific choice of

$$\psi(z) = c_1 \Lambda_{w_1}(z) + c_2 \Lambda_{w_2}(z) + \cdots + c_d \Lambda_{w_d}(z)$$

with the coefficients determined above. The requirement of $\psi(0) = 1$ therefore gives

$$a w_1 w_2 \cdots w_d = \frac{a}{w_1 w_2 \cdots w_d} + 1,$$

which will furnish the value of a , namely,

$$a = \left[w_1 w_2 \cdots w_d - \frac{1}{w_1 w_2 \cdots w_d} \right]^{-1}.$$

Finally, an application of the triangle inequality yields

$$\begin{aligned} \|1 - \lambda f\|_p &\geq \|\psi\|_q^{-1} \geq \left\| B - \frac{a}{w_1 w_2 \cdots w_d} \right\|_q^{-1} \\ &\geq \left[\|B\|_q + \left\| \frac{a}{w_1 w_2 \cdots w_d} \right\|_q \right]^{-1} = \left[|a| + \left| \frac{a}{w_1 w_2 \cdots w_d} \right| \right]^{-1} \\ &= \frac{1 - |f(0)|^2}{1 + |f(0)|} = 1 - |f(0)| = 1 - |w_1 w_2 \cdots w_d|. \quad \square \end{aligned}$$

Remark 4.0.6. The above result holds for any function f vanishing at the points w_1, \dots, w_d . Further, by Theorem 3.1.4, the result also extends to any infinite Blaschke product.

Let us now use duality to further investigate OPA errors for more general functions.

Proposition 4.0.7. *Let $1 < p < \infty$, $\frac{1}{p} + \frac{1}{q} = 1$, $n \in \mathbb{N}$, and $f \in H^p$ with $f(0) = 1$. Then*

$$\|q_{n-1,p}[f]f - 1\|_p \geq \frac{1}{\|1 + \psi_1 z + \psi_2 z^2 + \cdots + \psi_n z^n\|_q},$$

where the coefficients $\psi_k, 1 \leq k \leq n$, satisfy the matrix equation

$$\begin{bmatrix} f_1 & f_2 & f_3 & \cdots & f_n \\ f_0 & f_1 & f_2 & \cdots & f_{n-1} \\ 0 & f_0 & f_1 & \cdots & f_{n-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & f_1 \end{bmatrix} \begin{bmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \vdots \\ \psi_n \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

Proof. It suffices to check that the function

$$\psi(z) = 1 + \psi_1 z + \psi_2 z^2 + \cdots + \psi_n z^n$$

satisfies the hypotheses of Lemma 4.0.1. That is, for $0 \leq k \leq n$, that

$$\langle z^k f, \psi \rangle = 0.$$

This is ensured precisely by the linear system in the statement of the proposition. \square

With further calculation, the approach in the previous proposition can be used to show the following:

Proposition 4.0.8. *Let $1 < p < \infty, \frac{1}{p} + \frac{1}{q} = 1, n \in \mathbb{N}$, and $f \in H^p$ with $f(0) = 1$. Let*

$$\frac{1}{f(z)} = 1 + g_1 z + g_2 z^2 + \cdots$$

be the power series of $\frac{1}{f}$ about the origin. Then

$$\|q_{n-1,p}[f]f - 1\|_p \geq \frac{|g_n|}{\|1 + g_1 z + g_2 z^2 + \cdots + g_n z^n\|_q}.$$

Proof. Let us begin with, from Proposition 4.0.7, the matrix equation

$$\begin{bmatrix} f_1 & f_2 & f_3 & \cdots & f_n \\ f_0 & f_1 & f_2 & \cdots & f_{n-1} \\ 0 & f_0 & f_1 & \cdots & f_{n-2} \\ \vdots & \vdots & \cdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & f_1 \end{bmatrix} \begin{bmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \vdots \\ \psi_n \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

There is no harm in multiplying both sides of the equation on the left by the elementary permutation matrix

$$\begin{bmatrix} 0 & 0 & 0 & \cdots & 1 \\ 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 \end{bmatrix}$$

(the next to last entry of the bottom row is 1), which has the effect of changing the equation to

$$\begin{bmatrix} f_0 & f_1 & f_2 & \cdots & f_{n-1} \\ 0 & f_0 & f_1 & \cdots & f_{n-2} \\ 0 & 0 & f_0 & \cdots & f_{n-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & f_1 \\ f_1 & f_2 & f_3 & \cdots & f_n \end{bmatrix} \begin{bmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \psi_4 \\ \vdots \\ \psi_n \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ \vdots \\ -1 \end{bmatrix}.$$

By successively subtracting multiples of the other rows, the bottom row can be placed in the form

$$[0 \ 0 \ 0 \ \cdots \ C]$$

for some constant C , which could be zero. In fact, recalling that $f_0 = 1$, we see that C must be given by

$$C = \det \begin{bmatrix} f_1 & f_2 & f_3 & \cdots & f_n \\ f_0 & f_1 & f_2 & \cdots & f_{n-1} \\ 0 & f_0 & f_1 & \cdots & f_{n-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & f_1 \end{bmatrix}.$$

Furthermore, the sequence of row operations to diagonalize the matrix leaves the right side unchanged as

$$[0 \ 0 \ 0 \ \cdots \ -1]^T.$$

Assuming that $C \neq 0$, and again recalling that $f_0 = 1$, our matrix equation can be written as

$$\begin{bmatrix} f_0 & f_1 & f_2 & \cdots & f_{n-1} \\ 0 & f_0 & f_1 & \cdots & f_{n-2} \\ 0 & 0 & f_0 & \cdots & f_{n-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & f_0 \end{bmatrix} \begin{bmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \vdots \\ \psi_n \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ -1/C \end{bmatrix}.$$

The inverse of the transposed (Toeplitz) matrix on the left is simply

$$\begin{bmatrix} g_0 & g_1 & g_2 & \cdots & g_{n-1} \\ 0 & g_0 & g_1 & \cdots & g_{n-2} \\ 0 & 0 & g_0 & \cdots & g_{n-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & g_0 \end{bmatrix},$$

where $g(0) = 1$ and $g(z) = g_0 + g_1z + g_2z^2 + \cdots$ is the Taylor expansion of $1/f(z)$, valid for some disk centered at the origin. The conclusion is that

$$\psi_k = -\frac{g_{n-k}}{C} \quad \text{for all } 0 \leq k \leq n-1.$$

Our next challenge is to find an analytical meaning for the constant C . But notice that the row operations needed to clear entries from the bottom row of

$$(4.0.9) \quad \begin{bmatrix} f_0 & f_1 & f_2 & \cdots & f_{n-1} \\ 0 & f_0 & f_1 & \cdots & f_{n-2} \\ 0 & 0 & f_0 & \cdots & f_{n-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & f_1 \\ f_1 & f_2 & f_3 & \cdots & f_n \end{bmatrix}$$

would (suitably modified) similarly clear the second through the last entries from the top row. Performing all of these (suitably modified) row operations on the identity matrix would have to result in

$$\begin{bmatrix} 1 & g_1 & g_2 & \cdots & g_{n-1} \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{bmatrix}$$

Then following carefully what operations are correspondingly performed on the last column in (4.0.9), we conclude that

$$C = f_n + g_1f_{n-1} + g_2f_{n-2} + \cdots + g_{n-1}f_1 = -g_n.$$

Finally, note that

$$\begin{aligned} \left\| 1 + \frac{g_{n-1}}{g_n}z + \frac{g_{n-2}}{g_n}z^2 + \cdots + \frac{1}{g_n}z^n \right\|_q &= \frac{1}{|g_n|} \|g_n + g_{n-1}z + g_{n-2}z^2 + \cdots + z^n\|_q \\ &= \frac{1}{|g_n|} \|z^{-n}(g_n + g_{n-1}z + g_{n-2}z^2 + \cdots + z^n)\|_q \\ &= \frac{1}{|g_n|} \|1 + g_1z^{-1} + g_2z^{-2} + \cdots + g_nz^{-n}\|_q \\ &= \frac{1}{|g_n|} \|1 + g_1z + g_2z^2 + \cdots + g_nz^n\|_q, \end{aligned}$$

where in the last step, the change of variable $\theta \mapsto -\theta$, for $z = e^{i\theta}$, leaves the norm integral unchanged. \square

We will provide an improvement to the above proposition, but we must first consider the problem of finding $G \in zH^p$ such that

$$\|\overline{G} + F\|_p$$

is minimized. Duality tells us that (as we continue to mark extremal functions with *)

$$\begin{aligned} \|\overline{G^*} + F\|_p &= \sup \left\{ \frac{|\langle F, \psi \rangle|}{\|\psi\|_q} : \psi \in H^q \right\} \\ &= \frac{|\langle F, \psi^* \rangle|}{\|\psi^*\|_q} \\ &= \frac{|\langle F, \psi^* \rangle|}{\inf\{\|\psi^* + K\|_q : K \in \overline{zH^q}\}}. \end{aligned}$$

Once again, we are up against the dual of H^p being isomorphic to H^q , but not isometrically.

Nonetheless, we must consider the metric projection of F onto the subspace $\overline{zH^p}$. Notice that $(\overline{G^*} + F)^{\langle p-1 \rangle}$ annihilates any negative frequencies. Therefore, there exists $h \in H^q$ such that

$$(\overline{G^*} + F)^{\langle p-1 \rangle} = h,$$

and thus, taking a $\langle q - 1 \rangle$ power, we have

$$|h|^{\langle q-2 \rangle} h = \overline{G^*} + F.$$

In turn, we see that finding G^* amounts to solving the above highly unpleasant functional equation.

Let us record this in the following result, where we write P_+ for the Riesz projection, given by

$$\sum_{k=-\infty}^{\infty} c_k z^k \mapsto \sum_{k=0}^{\infty} c_k z^k,$$

which is bounded from $L^p \rightarrow H^p$.

Proposition 4.0.10. *Let $1 < p < \infty$ and $\frac{1}{p} + \frac{1}{q} = 1$. Suppose $h \in H^q$, and define $F := P_+ h^{\langle q-1 \rangle}$. Then*

$$\inf\{\|F + \overline{G}\|_p : G \in zH^p\}$$

is attained by taking $F + \overline{G} = \overline{h^{\langle q-1 \rangle}}$. In this case, the value of the infimum is given by

$$\inf\{\|F + \overline{G}\|_p : G \in zH^p\} = \|h\|_q^{q-1}.$$

This warrants the following observation:

Proposition 4.0.11. *For $1 < p < \infty$, the set of images $P_+ \overline{(H^q)^{\langle q-1 \rangle}}$ is dense in H^p .*

Proof. Suppose $g \in H^q$ has the property that

$$\langle P_+ \overline{h^{(q-1)}}, g \rangle = 0$$

for all $h \in H^q$. Then

$$0 = \langle P_+ \overline{h^{(q-1)}}, g \rangle = \int_{\mathbb{T}} P_+ \overline{h^{(q-1)}} \bar{g} \, dm = \int_{\mathbb{T}} \overline{h^{(q-1)}} \bar{g} \, dm.$$

We are able to drop the projection in the last line since integration against \bar{g} will annihilate the negative frequencies of $\overline{h^{(q-1)}}$. In particular, this must hold for $h = g$, and hence $0 = \|g\|_q^q$. This forces g to be identically zero. \square

In turn, we can make the following improvement to Proposition 4.0.8.

Proposition 4.0.12. *Let $1 < p < \infty$, $\frac{1}{p} + \frac{1}{q} = 1$, $n \in \mathbb{N}$, and $f \in H^p$ with $f(0) = 1$. Let*

$$\frac{1}{f(z)} = 1 + g_1 z + g_2 z^2 + \dots$$

be the power series of $\frac{1}{f}$ about the origin. If $g_n \neq 0$, then

$$\|q_{n-1,p}[f]f - 1\|_p \geq \frac{1}{\|h^{(p-1)}\|_q},$$

where $h \in H^p$ satisfies

$$P_+ \overline{h^{(p-1)}} = 1 + \frac{g_{n-1}}{g_n} z + \frac{g_{n-2}}{g_n} z^2 + \dots + \frac{g_0}{g_n} z^n.$$

Let us now consider the case where $n \rightarrow \infty$. Then, by writing $f = JG$ for J inner and G outer, we see, for all $k \geq 0$, that $\psi \in L^q$ satisfies

$$0 = \langle z^k f, \psi \rangle = \langle z^k JG, \psi \rangle = \langle z^k G, \bar{J}\psi \rangle.$$

As $\{z^k G : k \geq 0\}$ is dense in H^p , we have, for any $n \geq 0$,

$$0 = \langle z^n, \bar{J}\psi \rangle = \int_0^{2\pi} J(e^{i\theta}) \overline{\psi(e^{i\theta})} e^{in\theta} \frac{d\theta}{2\pi}.$$

From this, it follows that $K(z) := J(z) \overline{\psi(z)}/z$ is an element of H^q . We further divine that ψ must be determined by

$$\psi(z) = J(z) \overline{zK(z)}, \quad z \in \mathbb{T},$$

for some $K \in H^q$. The condition $\psi(0) = 1$ takes the form

$$1 = J_1 \bar{K}_0 + J_2 \bar{K}_1 + J_3 \bar{K}_2 + \dots.$$

We now must minimize $\|J\bar{z}\bar{K}\|_q$ subject to $K \in H^q$ satisfying the above constraint. It is tempting to try $K = \bar{J}$, but this will not work, since

$$0 = J_1\bar{J}_0 + J_2\bar{J}_1 + J_3\bar{J}_2 + \dots$$

Instead, take $K(z) = c[J(z) - J(0)]/z$, where $c^{-1} = |J_1|^2 + |J_2|^2 + |J_3|^2 + \dots$. Then

$$J_1\bar{K}_0 + J_2\bar{K}_1 + J_3\bar{K}_2 + \dots = c(|J_1|^2 + |J_2|^2 + |J_3|^2 + \dots) = 1,$$

as needed. Using this choice of K to compute ψ , we obtain

$$\psi(z) = J(z)\overline{zK(z)} = cJ(z)\bar{z} \frac{\overline{J(z)} - \overline{J(0)}}{\bar{z}} = c(1 - J(z)\overline{J(0)}).$$

Since $\|q_{n,p}[f]f - 1\|_p \geq 1/\|\psi\|_q$, this furnishes the following bound.

Proposition 4.0.13. *Let $1 < p < \infty$, $\frac{1}{p} + \frac{1}{q} = 1$, $n \in \mathbb{N}$, and $f \in H^p$ with $f(0) = 1$. Then*

$$\|q_{n,p}[f]f - 1\|_p \geq \frac{|J_1|^2 + |J_2|^2 + |J_3|^2 + \dots}{\|1 - \overline{J(0)}J(z)\|_q},$$

where J is the inner part of f .

Incidentally, $c^{-1} = (|J_1|^2 + |J_2|^2 + |J_3|^2 + \dots) = \|J\|_2^2 - |J(0)|^2 = 1 - |J(0)|^2$, so the lower bound above could be written equivalently as

$$\frac{1 - |J(0)|^2}{\|(1 - |J(0)|^2) - \overline{J(0)}(J_1z + J_2z^2 + \dots)\|_q},$$

which is obviously no greater than 1, as needed.

We now step away from duality. Our final results concern OPA errors, but are proven with H^2 methods. The following proposition should be compared with Proposition 4.0.3; although the result below provides a better bound, it holds only for $p > 2$.

Proposition 4.0.14. *Let $2 < p < \infty$, and suppose $f \in H^p$ has a factorization $f = JG$, where J is inner and G is outer. If $f(0) \neq 0$, then, for any $n \in \mathbb{N}$,*

$$\|q_{n,p}[f]f - 1\|_p \geq \sqrt{1 - |J(0)|^2}.$$

Proof. Let \mathcal{P} be the collection of all polynomials. Then

$$\begin{aligned} \|q_{n,p}[f]f - 1\|_p &\geq \inf_{Q \in \mathcal{P}} \|Qf - 1\|_p \geq \inf_{Q \in \mathcal{P}} \|Qf - 1\|_2 \\ &= \inf_{Q \in \mathcal{P}} \|QJG - 1\|_2 = \inf_{Q \in \mathcal{P}} \|QG - \bar{J}\|_2 = \|\overline{J(0)} - \bar{J}\|_2, \end{aligned}$$

with the last equality following from the fact that since G is outer, $\{QG : Q \in \mathcal{P}\}$ is dense in H^2 . Now use

$$1 = \|J\|_2^2 = |J(0)|^2 + \|J - J(0)\|_2^2. \quad \square$$

Note further that if $J = B$ is a Blaschke product, then this implies that

$$\|q_{n,p}[f]f - 1\|_p \geq \|J(0) - J\|_2 = \sqrt{1 - |B(0)|^2} = \sqrt{1 - |w_1 w_2 w_3 \cdots|^2},$$

where w_1, w_2, w_3, \dots are the zeros of B .

We end by providing a related result when $1 < p < 2$.

Proposition 4.0.15. *Let $1 < p < 2$ and suppose $f(0) \neq 0$. Then, for any $n \in \mathbb{N}$,*

$$\|q_{n,p}[f]f - 1\|_p \leq \sqrt{1 - (q_{n,2}[f]f)(0)}.$$

Proof. Routine bounds yield

$$\|q_{n,p}[f]f - 1\|_p \leq \|q_{n,2}[f]f - 1\|_p \leq \|q_{n,2}[f]f - 1\|_2 = \sqrt{1 - (q_{n,2}[f]f)(0)},$$

where the last equality is a consequence of the linear system described in (1.0.1). \square

Noting that $\|q_{n,2}[f]f - 1\|_2 \leq \|q_{0,2}[f]f - 1\|_2$, we can also establish the simple bound

$$\|q_{n,p}[f]f - 1\|_p \leq \left(1 - \frac{|f(0)|^2}{\|f\|_2^2}\right)^{1/2}.$$

References

- [1] A. Acuaviva and D. Seco, “Equidistribution of zeros of some polynomials related to cyclic functions”, *Anal. Math. Phys.* **12**:2 (2022), art. id. 53. MR Zbl
- [2] J. Alonso, H. Martini, and S. Wu, “On Birkhoff orthogonality and isosceles orthogonality in normed linear spaces”, *Aequationes Math.* **83**:1-2 (2012), 153–189. MR Zbl
- [3] P. Arora, M. Augat, M. T. Jury, and M. Sargent, “An optimal approximation problem for free polynomials”, *Proc. Amer. Math. Soc.* **152**:2 (2024), 455–470. MR Zbl
- [4] C. Bénéteau and R. Centner, “A survey of optimal polynomial approximants, applications to digital filter design, and related open problems”, *Complex Anal. Synerg.* **7**:2 (2021), art. id. 16. MR Zbl
- [5] C. Bénéteau, A. A. Condori, C. Liaw, D. Seco, and A. A. Sola, “Cyclicity in Dirichlet-type spaces and extremal polynomials”, *J. Anal. Math.* **126** (2015), 259–286. MR Zbl
- [6] C. Bénéteau, A. A. Condori, C. Liaw, D. Seco, and A. A. Sola, “Cyclicity in Dirichlet-type spaces and extremal polynomials, II: Functions on the bidisk”, *Pacific J. Math.* **276**:1 (2015), 35–58. MR Zbl
- [7] C. Bénéteau, G. Knese, L. u. Kosiński, C. Liaw, D. Seco, and A. Sola, “Cyclic polynomials in two variables”, *Trans. Amer. Math. Soc.* **368**:12 (2016), 8737–8754. MR Zbl
- [8] C. Bénéteau, M. C. Fleeman, D. S. Khavinson, D. Seco, and A. A. Sola, “Remarks on inner functions and optimal approximants”, *Canad. Math. Bull.* **61**:4 (2018), 704–716. MR Zbl

- [9] C. Bénéteau, D. Khavinson, C. Liaw, D. Seco, and B. Simanek, “Zeros of optimal polynomial approximants: Jacobi matrices and Jentzsch-type theorems”, *Rev. Mat. Iberoam.* **35**:2 (2019), 607–642. MR Zbl
- [10] R. Centner, “Optimal polynomial approximants in L^p ”, *Concr. Oper.* **9**:1 (2022), 96–113. MR Zbl
- [11] R. Centner, R. Cheng, and C. Felder, “Optimal polynomial approximants in H^p ”, preprint, 2023. arXiv 2305.16068
- [12] R. Cheng and W. T. Ross, “Weak parallelogram laws on Banach spaces and applications to prediction”, *Period. Math. Hungar.* **71**:1 (2015), 45–58. MR Zbl
- [13] R. Cheng, J. Mashreghi, and W. T. Ross, *Function theory and ℓ^p spaces*, University Lecture Series **75**, American Mathematical Society, Providence, RI, 2020. MR Zbl
- [14] R. Cheng, W. T. Ross, and D. Seco, “Zeros of optimal polynomial approximants in ℓ_A^p ”, *Adv. Math.* **404**:part A (2022), art. id. 108396. MR Zbl
- [15] E. Fricain, J. Mashreghi, and D. Seco, “Cyclicity in reproducing kernel Hilbert spaces of analytic functions”, *Comput. Methods Funct. Theory* **14**:4 (2014), 665–680. MR Zbl
- [16] R. C. James, “Orthogonality and linear functionals in normed linear spaces”, *Trans. Amer. Math. Soc.* **61** (1947), 265–292. MR Zbl
- [17] M. Sargent and A. A. Sola, “Optimal approximants and orthogonal polynomials in several variables, II: Families of polynomials in the unit ball”, *Proc. Amer. Math. Soc.* **149**:12 (2021), 5321–5330. MR Zbl
- [18] M. Sargent and A. A. Sola, “Optimal approximants and orthogonal polynomials in several variables”, *Canad. J. Math.* **74**:2 (2022), 428–456. MR Zbl

Received November 25, 2023. Revised February 4, 2024.

RAYMOND CHENG
 DEPARTMENT OF MATHEMATICS AND STATISTICS
 OLD DOMINION UNIVERSITY
 NORFOLK, VA
 UNITED STATES
 rcheng@odu.edu

CHRISTOPHER FELDER
 DEPARTMENT OF MATHEMATICS
 INDIANA UNIVERSITY BLOOMINGTON
 BLOOMINGTON, IN
 UNITED STATES
 cfelder@iu.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2023 is US \$605/year for the electronic version, and \$820/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 327 No. 2 December 2023

Co-Hopfian and boundedly endo-rigid mixed abelian groups	183
MOHSEN ASGHARZADEH, MOHAMMAD GOLSHANI and SAHARON SHELAH	
The fundamental solution to \square_b on quadric manifolds with nonzero eigenvalues and null variables	233
ALBERT BOGGESS and ANDREW RAICH	
More properties of optimal polynomial approximants in Hardy spaces	267
RAYMOND CHENG and CHRISTOPHER FELDER	
The Griffiths double cone group is isomorphic to the triple	297
SAMUEL M. CORSON	
Arithmetic modular links	337
TALI PINSKY, JESSICA S. PURCELL and JOSÉ ANDRÉS RODRÍGUEZ-MIGUELES	
Tensor triangulated category structures in the derived category of a variety with big (anti)canonical bundle	359
ANGEL TOLEDO	