



Complements of tori and Klein bottles in the 4-sphere that have hyperbolic structure

DUBRAVKO IVANŠIĆ
 JOHN G. RATCLIFFE
 STEVEN T. TSCHANTZ

Abstract Many noncompact hyperbolic 3-manifolds are topologically complements of links in the 3-sphere. Generalizing to dimension 4, we construct a dozen examples of noncompact hyperbolic 4-manifolds, all of which are topologically complements of varying numbers of tori and Klein bottles in the 4-sphere. Finite covers of some of those manifolds are then shown to be complements of tori and Klein bottles in other simply-connected closed 4-manifolds. All the examples are based on a construction of Ratcliffe and Tschantz, who produced 1171 noncompact hyperbolic 4-manifolds of minimal volume. Our examples are finite covers of some of those manifolds.

AMS Classification 57M50, 57Q45

Keywords Hyperbolic 4-manifolds, links in the 4-sphere, links in simply-connected closed 4-manifolds

1 Introduction

Let \mathbf{H}^n be the n -dimensional hyperbolic space and let G be a discrete subgroup of $\text{Isom } \mathbf{H}^n$, the isometries of \mathbf{H}^n . If G is torsion-free, then $M = \mathbf{H}^n/G$ is a hyperbolic manifold of dimension n . A hyperbolic manifold of interest in this paper is also complete, noncompact and has finite volume and the term “hyperbolic manifold” will be understood to include those additional properties. Such a manifold M is the interior of a compact manifold with boundary \overline{M} . Every boundary component E of \overline{M} is a compact flat (Euclidean) manifold, i.e. a manifold of the form \mathbf{R}^{n-1}/K , where K is a discrete subgroup of $\text{Isom } \mathbf{R}^{n-1}$, the isometries of \mathbf{R}^{n-1} . For simplicity, we sometimes inaccurately call E a boundary component of M (rather than \overline{M} , whose boundary component it really is).

When $n = 3$, M is the interior of a compact 3-manifold \overline{M} whose boundary consists of tori and Klein bottles. When all the boundary components are tori,

M may be diffeomorphic to S^3 with several closed solid tori removed, which in turn is diffeomorphic to $S^3 - A$, where A is the link consisting of the core circles of the solid tori. Indeed, it is a well known fact that many hyperbolic 3-manifolds are link complements in S^3 .

The first example of a generalization of this situation to dimension 4 was given by Ivanšić [6], where it was shown that \widetilde{M}_{1011} , the orientable double cover of one of the 1049 nonorientable hyperbolic 4-manifolds that Ratcliffe and Tschantz constructed [9], is a complement of 5 tori in S^4 . Finite covers of \widetilde{M}_{1011} were also shown to be complements of a collection of tori in a simply-connected 4-manifold with even Euler characteristic.

In this paper, we find 11 additional examples of hyperbolic 4-manifolds that are complements in S^4 of a set of tori or a combination of tori and Klein bottles. The list of examples is in Theorem 4.4. Like \widetilde{M}_{1011} , all of them are orientable double covers of some of Ratcliffe and Tschantz's nonorientable manifolds. Proving that those manifolds are complements in S^4 is typically trickier than in [6], since the many symmetries of M_{1011} enabled an easier computation.

By taking finite covers of some of the link complements we also find additional examples of hyperbolic 4-manifolds that are complements of a collection of tori and Klein bottles in a simply-connected 4-manifold with even Euler characteristic. Example 4.10 displays 8 different families of examples, illustrating the richness of such objects, many more of which likely exist.

Manifolds \widetilde{M}_{1091} and \widetilde{M}_{1011} were used by Ratcliffe and Tschantz [10] to construct the first examples (infinitely many) of aspherical homology 4-spheres. Three other examples in this paper can be used for the same purpose, see Example 4.11.

When looking for examples of link complements in S^4 among the orientable double covers of the 1049 nonorientable manifolds, the main difficulty is to reduce the number of potential examples with which to experiment. We use a homological and a group-theoretical criterion (see §4) to rule out all except 49 manifolds from having the desired property. Then we show that some of the remaining 49 manifolds have orientable double covers that are complements of tori and Klein bottles in the 4-sphere. Essentially, we use group presentations to show that the manifold N resulting from closing up the boundary components of the double covers is simply-connected. This, along with the fact that the Euler characteristic $\chi(N)$ is 2, guarantees that N is homeomorphic to S^4 .

Ratcliffe and Tschantz constructed their manifolds by side-pairings of a hyperbolic noncompact right-angled 24-sided polyhedron, making computations with

them laborious, especially if they have to be done repeatedly. The two criteria were checked using a computer, but we emphasize that the results of this paper do not depend on a computer calculation, since it is verified by hand that some of the “promising” examples indeed have a double cover that is a complement in the 4-sphere.

2 Some preliminary facts

Let M be hyperbolic n -manifold. We say that M is a (codimension-2) complement in N if $M = N - A$, where N is a closed n -manifold and A is a closed $(n - 2)$ -submanifold of N that has a tubular neighborhood in N and has as many components as $\partial\overline{M}$. Then every component E of $\partial\overline{M}$, being the boundary of a tubular neighborhood, must be an S^1 -bundle over a component of A . The $(n - 1)$ -manifold E is flat. The following theorem (see [5] or [1] Theorem 6.40) summarizes when a flat manifold is an S^1 -bundle and, when it is, what its S^1 -fibers are.

Theorem 2.1 *Let $E = \mathbf{R}^{n-1}/K$ be a compact flat $(n - 1)$ -manifold, where K is a discrete subgroup of $\text{Isom } \mathbf{R}^{n-1}$. Then E is an S^1 -bundle over a manifold B if and only if there exists a translation $t \in K$ such that $\langle t \rangle$ is a normal subgroup of K and t is not a power of any element of K other than $t^{\pm 1}$. If the translation is given by $x \mapsto x + v$ and any element of K by $x \mapsto Ax + a$, where $A \in O(n - 1)$, $a \in \mathbf{R}^{n-1}$, the normality of $\langle t \rangle$ can be expressed as $Av = \pm v$ for every A such that $Ax + a$ is an element of K . Furthermore, if $n - 1 \neq 4, 5$ (we are interested in $n - 1 = 3$ in this paper) the manifold B above is a flat manifold. \square*

We say that a translation t is *normal in E* if $\langle t \rangle$ is a normal subgroup, we say it is *primitive in E* if it is not a power of any other element of K .

By Theorem 2.1, if M is a hyperbolic 4-manifold that is a complement in N , it must be a complement of flat 2-manifolds, that is, tori and Klein bottles. We are interested in the case when $N = S^4$. When $M = N - A$, an immediate consequence of the fact that components of A are flat is that $\chi(M) = \chi(N)$, so examples of complements in S^4 must be searched for among manifolds with $\chi(M) = 2$. Nonorientable examples of Ratcliffe and Tschantz are very suitable for this purpose, since their Euler characteristic is 1: passing to the orientable double cover will give us the required Euler characteristic.

Once we have found a hyperbolic manifold M all of whose boundary components are S^1 -bundles, $M = N - A$. To show that N is a topological S^4 , it suffices to show that $\pi_1 N = 1$ and $\chi(N) = 2$ (see [3] or [2]). The following application of van Kampen's theorem shows how to compute $\pi_1 N$ from $\pi_1 M$; details are in [6].

Proposition 2.2 *Let M be a complement in N and let $\partial\overline{M} = E_1 \cup \dots \cup E_m$. If $t_1, \dots, t_m \in \pi_1 M$ represent fibers of the S^1 -bundles E_1, \dots, E_m then $\pi_1 N = \pi_1 M / \langle\langle t_1, \dots, t_m \rangle\rangle$, where $\langle\langle A \rangle\rangle$ denotes the normal closure of a subset $A \subset \pi_1 M$. In other words, if we have a presentation for M , the presentation for N is obtained by adding relations $t_1 = 1, \dots, t_m = 1$. \square*

Hyperbolic manifolds M of interest in this paper are given by side-pairings of a hyperbolic finite-volume noncompact polyhedron Q . Side-pairings determine sets of points on the polyhedron, called *cycles*, that are identified in the manifold. Boundary components of \overline{M} correspond to cycles of ideal vertices of Q . If $[v_i]$ is the cycle of an ideal vertex corresponding to E_i , then $\text{stab } v_i$ is isomorphic to $\pi_1 E_i$, making the inclusion $\pi_1 E_i \rightarrow \pi_1 M$ injective. Elements of $\pi_1 E_i$ are parabolic isometries, thus, they preserve horospheres centered at v_i and act like Euclidean isometries on them.

Let Γ be a discrete subgroup of $\text{Isom } \mathbf{H}^n$ with fundamental polyhedron P . Then Γ is generated by side-pairings of P , isometries that send a side of P to another side of P . If s is an isometry sending side S of P to S' , then, if we start moving from P and pass through S , we wind up in $s^{-1}P$. Similarly, if we take a path P to γP that passes through interiors of translates of sides S_1, \dots, S_m whose side-pairings are s_1, \dots, s_m , then $\gamma = s_1^{-1} \dots s_m^{-1}$. Furthermore, if G is a finite-index subgroup of Γ with transversal X (that is, a set of right-coset representatives, so $\Gamma = \cup_{x \in X} Gx$), then the fundamental polyhedron of G is $Q = \cup_{x \in X} xP$, where elements of X can be chosen so that Q is connected.

The fundamental groups G of Ratcliffe and Tschantz's manifolds are all minimal index torsion-free subgroups of a certain reflection group Γ , which is generated by reflections in a 10-sided right-angled polyhedron P . The side-pairings generating Γ are, of course, reflections in the sides of P . Every subgroup G corresponding to a Ratcliffe-Tschantz manifold has the same transversal in Γ , a finite group K isomorphic to \mathbf{Z}_2^4 and generated by reflections in the four coordinate planes in the ball model of \mathbf{H}^4 . The fundamental polyhedron of G is $Q = KP$, a 24-sided regular right-angled hyperbolic polyhedron that we call the *24-cell*. Let f be a side-pairing of Q that sends side R to side R' . As was shown in [9] and [6], f has to have form $f = kr = r'k$, where $k \in K$ is an

element such that $k(R) = R'$, and r, r' are reflections in R, R' , respectively. We will call k the K -part of f . Thus, it is enough to specify elements $k \in K$ in order to define a side-pairing of Q .

As is clear from Theorem 2.1 and Proposition 2.2, we will need the ability to recognize when a composite of side-pairings is a translation. The following proposition, proved in [6], will be used for that purpose:

Proposition 2.3 *Let Γ be generated by reflections in the sides of a polyhedron P and let G be a finite-index subgroup of Γ so that its transversal is a finite group K , making $Q = KP$ the fundamental polyhedron of G . If a path from Q to gQ passes through translates of sides R_1, \dots, R_m of Q , then g can be written as $g = (r_m \dots r_1)(k_1^{-1} \dots k_m^{-1})$, where r_i is a reflection in a translate of R_i and $k_i \in K$ is the K -part of the side-pairing of R_i . \square*

In this paper we work with presentations of groups and their subgroups. If a group G is given by presentation $\langle Y | R \rangle$, and H is a finite-index subgroup of G with Schreier transversal X (i.e. $G = \cup_{x \in X} Hx$ and every initial segment of an element $x \in X$ is also an element of X), then the presentation of H can be derived from the presentation of G . Let $g \mapsto \bar{g}$ be the map $G \rightarrow X$ sending g to its coset representative, that is $Hg = H\bar{g}$, and let $\gamma : G \rightarrow H$ be the map $\gamma(g) = g\bar{g}^{-1}$.

Proposition 2.4 (The Reidemeister-Schreier method, [7] Proposition 4.1) *The presentation of H is given by $\langle Z | S \rangle$, where $Z = \{\gamma(xy) \neq 1 \mid x \in X, y \in Y\}$ and S is the set of all words rxr^{-1} written in terms of elements of Z , for all $x \in X$ and $r \in R$. \square*

3 The 24-cell, its side-pairings and cycle relations

Working in the ball model of \mathbf{H}^4 , let S_{****} be one of the 24 spheres of radius 1 centered at a point in \mathbf{R}^4 whose coordinates have two zeroes and two ± 1 's. The string $****$ consists of symbols $+, -, 0$ and identifies the sphere by its center. For example, S_{+0-0} is the sphere centered at $(1, 0, -1, 0)$. These spheres are orthogonal to $\partial\mathbf{H}^4$, hence they determine hyperplanes in \mathbf{H}^4 . The 24-sided polyhedron Q is defined to be the intersection of the half-spaces corresponding to those hyperplanes that contain the origin. The side of Q lying on the sphere S_{****} is also denoted by S_{****} . All the dihedral angles of Q are $\pi/2$ and its 24 ideal vertices are $v_{\pm 000} = (\pm 1, 0, 0, 0)$, $v_{0\pm 00} = (0, \pm 1, 0, 0)$, $v_{00\pm 0} =$

$(0, 0, \pm 1, 0)$, $v_{000\pm} = (0, 0, 0, \pm 1)$ and $v_{\pm\pm\pm\pm} = (\pm 1/2, \pm 1/2, \pm 1/2, \pm 1/2)$. The polyhedron Q has no real vertices.

Note that two sides intersect if and only if their identifying strings have equal nonzero entries in one position and the positions of the remaining nonzero entries are different. Two sides touch at $\partial\mathbf{H}^4$ when they have equal nonzero entries in one position and opposite nonzero entries in another position, or if the positions where they have nonzero entries are complementary. Furthermore, an ideal vertex is on a side if its Euclidean distance from the center of the sphere defining the side is 1. In the case of vertices whose label has one nonzero entry, a vertex is on a side if and only if the side has an equal nonzero entry in the same position as the vertex; in the case of vertices $v_{\pm\pm\pm\pm}$, a vertex is on a side if and only if the nonzero entries in the label of the side coincide with the entries in the label of the vertex in the same position. For example, S_{0-+0} and S_{+-00} intersect, S_{0-+0} and S_{0+0+} are disjoint, the side S_{0+0-} has six ideal vertices v_{0+00} , v_{000-} , $v_{\pm\pm\pm-}$; S_{0-+0} and S_{0--0} touch at v_{0-00} , and S_{0-+0} and S_{-00-} touch at v_{--+-} .

As mentioned in §2, every Ratcliffe-Tschantz example has side-pairing isometries of form rk , where $k \in \mathbf{Z}_2^4$ is a composite of reflections in the coordinate planes $x_1 = 0, \dots, x_4 = 0$. The label $\pm\pm\pm\pm$ identifies an element of K , for example, k_{+--+} is the composite of reflections in x_2 and x_3 . Ratcliffe and Tschantz have found in [9] that all sides whose labels have nonzero entries in the same two positions must have the same K -part for their side-pairing.

Side-pairing labeling convention Group letters a, \dots, l and the sides of Q as follows:

$$\{a, b, S_{\pm\pm 00}\}, \{c, d, S_{\pm 0 \pm 0}\}, \{e, f, S_{0 \pm \pm 0}\}, \\ \{g, h, S_{\pm 00 \pm}\}, \{i, j, S_{0 \pm 0 \pm}\}, \{k, l, S_{00 \pm \pm}\}.$$

The two letters in each group denote the side-pairings of the four sides from the group. The first letter always pairs the side labeled $++$, and the second letter pairs the side whose label is the next unused one, where the symbols $\pm\pm$ have been ordered in the dictionary order: $++$, $+-$, $-+$, $--$.

The encoding of the side-pairing transformation is done like in [9]. A string of six characters from $\{1, \dots, 9, A, \dots, F\}$, one for each group above, encodes the K -part of each side-pairing. Each character stands for a hexadecimal number, which is turned into binary form and the *order of digits is reversed*. Converting every 0 into a plus and every 1 into a minus obtains the label of k . For example, 1 stands for k_{-+++} ; $C = 12$ stands for k_{++--} . (See Example 4.6.)

The presentation of a group G generated by side-pairings of a polyhedron is obtained by applying Poincaré’s polyhedron theorem ([8] Theorem 11.2.2). Finding presentations of the groups and identifying usable translations in parabolic groups is crucial to our proof of Theorem 4.4. An efficient method for these two tasks is essential if one is to deal with a dozen examples, so we elaborate on it.

Let C be a small enough horosphere centered at v_{++++} that intersects only the sides of Q that contain v_{++++} . Then $Q_C = C \cap Q$ is a cube with opposing sides S_{++00} and S_{00++} , S_{+0+0} and S_{0+0+} , S_{0++0} and S_{+00+} (see Figure 1). The tiling of \mathbf{H}^4 by translates of Q intersected with C gives a tiling of C by cubes, one of which is Q_C .




Figure 1: The cube Q_C when C is centered at v_{++++}

We use the term *ridge* to denote a codimension-2 face of a polyhedron, while an *edge* is a 1-face. Every edge of the cube Q_C represents a ridge of Q . Circling around the edge once gives a cycle relation for the presentation in Poincaré’s theorem. To easily determine which sides we pass through, use Figure 2. The left column depicts the intersections of the tiling of C with the three planes that pass through the center of Q_C and are parallel to its sides. The side-pairings in that figure are the ones for manifold M_{56} from Example 4.6.

Let s be the side-pairing sending S to S' , r_S the reflection in S and let k_S be the K -part of s . Consider a ridge $U \cap S$, where U is another side of Q . The ridge $U \cap S$ is sent via s to a ridge $V \cap S'$, where V is another side of Q . Now $V \cap S' = k_S r_S(U \cap S) = k_S(U \cap S) = k_S(U) \cap S'$ so we conclude $V = k_S(U)$. Therefore, adjacent to Q on the other side of S is $s^{-1}Q$, whose sides $s^{-1}V$ and $s^{-1}S'$ meet at ridge $U \cap S = s^{-1}(V \cap S')$.

The observation $V = k_S(U)$ allows us to easily find the labelings on all the struts in the diagrams in the left column of Figure 2 (labelings inside the square come from Q_C). For example, consider the sides around the vertex labeled 1 in the topmost diagram. The bottom left strut is the translate of the side $k_{S_{+00+}}(S_{0+0+}) = k_{-++-}(S_{0+0+}) = S_{0+0-}$, whose side-pairing is j . The left bottom strut is the translate of $k_{S_{0+0+}}(S_{+00+}) = k_{--++}(S_{+00+}) = S_{-00+}$, whose side-pairing is h^{-1} .




Figure 2: Diagrams that produce cycle relations for M_{56}

Cycle relations are words obtained circling an edge of Q_C and successively adding the side-pairings at the end of the word. Care must be taken to add s^{-1} at the end of the word whenever we *exit* through a side labeled s and to add s whenever we *enter* into a side labeled s . Going in the direction of the arrow around vertex 1 we exit through g and j and then enter into h^{-1} and i , so the cycle relation is $g^{-1}j^{-1}h^{-1}i = 1$.

We thus obtain 12 relations for G from the first column of Figure 2. Because Q has 96 ridges that fall into 24 ridge cycles, each corresponding to a relation, 12 further relations are needed. They are found by going around other edges

in the tiling of C . If we reflect the three planes we used for sections of Q_C in the sides of Q_C parallel to them, we get new sections of the tiling represented by diagrams in the middle and right columns of Figure 2. The fact $V = k_S(U)$ from above is now used to see that the labelings on the new diagrams are obtained by applying k_S to every label in the original diagram, where S is the side used to reflect the section plane.

The middle and right columns do not include struts because cycle relations follow from only the labels on the square part and the cycle relations for the left column. If $A = \{S_1, kS_1, S_2, kS_2\}$ is the set of four sides whose labels all have the same nonzero positions, and are hence paired by the same element $k \in K$, we note that any element $k' \in K$ acts on A and, due to commutativity of K , preserves or interchanges the subsets of A containing paired sides. Along with the fact that $k'^2 = 1$ for every $k' \in K$ this means that action of k' on A can be inferred just from knowing $k'S_1$. For example, in the third row in Fig. 1, we see that by applying $k_{S_{0++0}}$ to the sides labeled i and a we get sides labeled j and a^{-1} . We infer that applying $k_{S_{0++0}}$ to sides labeled j and b gives sides labeled i and b^{-1} . Thus, the cycle relation for vertex 2 in the new diagram is the cycle relation $i^{-1}b^{-1}ia$ for vertex 2 in the old diagram with letters replaced in the above pattern, obtaining $j^{-1}bja^{-1}$. Now, if a cycle relation coming from the new diagram is the inverse, a cyclic permutation or a combination of the two of a corresponding cycle relation in the old diagram, we have found the same ridge cycle, and have to reflect the section plane in the other side of the cube Q_C .

We argue that the nine diagrams thus obtained will contain all the cycle relations (multiple times, in fact). Note that in Q every ridge is an ideal triangle, two of whose ideal vertices are of type $v_{\pm\pm\pm\pm}$. For example, the ridge $R = S_{0+0+} \cap S_{0++0}$ is an ideal triangle with vertices v_{0+00} , v_{++++} and v_{-+++} and edges $E_1 = S_{0+0+} \cap S_{0++0} \cap S_{00++}$, $E_2 = S_{0+0+} \cap S_{0++0} \cap S_{++00}$, and $E_3 = S_{0+0+} \cap S_{0++0} \cap S_{-+00}$. The intersection of translates of edges of Q with C are the vertices in the tiling of C . An edge of the cube Q_C is an arc in the ideal triangle going from one edge of the triangle to another. Because elements of K do not change the nonzero positions in the labeling of sides, only E_2 and E_3 might be in the same cycle of edges. Thus, in the manifold \mathbf{H}^4/G the ideal triangle might only have its edges E_2 and E_3 identified, since no points on E_1 are identified. If ridge R and the ridge represented by reflection of $R \cap C$ in the side S_{00++} of Q_C were in the same cycle, the segment consisting of $R \cap C$ and its reflection would project to a smooth curve in the ideal triangle that crosses E_1 twice, which is impossible. Since in Q there are only 8 ridges of type $S_{0\pm0\pm} \cap S_{0\pm\pm0}$ we conclude that every one of them is either in the cycle

of R , or the cycle of the ridge represented by the reflection of $C \cap R$ in S_{00++} . Reasoning similarly for all the other ridge cycles we see that the opportunity for cycle relation repetitions occurs only for ridges represented by vertices labeled 2 and 3 in the middle column. Often there are no repetitions: in Example 4.6 we needed three section planes only in the first row.

Now we describe how to identify and choose suitable translations in parabolic subgroups. Let G_v be the parabolic group that is the stabilizer of an ideal vertex v of Q , and let C be a horosphere centered at v . The intersection $Q_C = Q \cap C$ is always a Euclidean cube. Position a coordinate system with origin in the center of Q_C and axes perpendicular to its sides, and consider the tiling of C by translates of Q_C . To simplify language, we identify a cube gQ_C and the element g . According to Proposition 2.3, by considering a path from Q_C to gQ_C that passes only through sides of cubes (e.g. a path that is piecewise parallel to the coordinate axes), we may write $g = (r_m \dots r_1)(k_1^{-1} \dots k_m^{-1})$. Since $r_m \dots r_1$ preserves v , $g \in G_v$ if and only if $k_1^{-1} \dots k_m^{-1}v = v$.

Let $v = v_{+000}$, v_{0+00} , v_{00+0} or v_{000+} . The coordinate axes perpendicular to sides of Q_C may be identified with coordinate axes x_1 , x_2 , x_3 and x_4 , in the arrangement suggested by Figure 3. The 4-symbol labels on the axes indicate which side we pass through if we move from the center of Q_C in the direction of the arrow. Parallel sides of each Q_C have same nonzero positions but differ in sign in positions other than the nonzero position of the vertex label (a side and a vertex have the same symbol in the nonzero position of the vertex label when the vertex is on the side). Moving two steps in the directions of the axes produces translations since $k_1 = k_2$ in every case. Let T_v be the subgroup of G_v generated by those three translations and let TG_v be the translation subgroup of G_v . For our purposes we look for translations only in $8Q_C$, the $2 \times 2 \times 2$ block of cubes that is the fundamental polyhedron for T_v . A Euclidean transformation $Ax + a$ is a translation if and only if its rotational part A is the identity. We get the rotational part of $r_m \dots r_1 k_1^{-1} \dots k_m^{-1}$ by replacing r_i by an element k_{r_i} that represents the reflection in the coordinate plane parallel to the reflection plane of r_i and computing the product. Note that it is enough to find the rotational parts of the three cubes adjacent to Q_C in the direction of the three axes. Any other rotational part is obtained by multiplying those three: every time we move by one step in the direction of an axis, we multiply by the rotational part of one of the three cubes that corresponds to that axis. The dots in Figure 3 represent centers of cubes comprising $8Q_C$; the dot is filled if it corresponds to an element preserving v . Labels on the dots give the rotational part of the group element corresponding to the dot. The three zero positions in the vertex label give the rotational part of the transformation,




Figure 3: Translations in $8Q_C$ for manifold M_{56}

while the nonzero position tells whether the transformation preserves v . For example, in the topmost diagram in Figure 3 the label $++-+$ on a vertex reveals an element that preserves v whose rotational part is a reflection in the x_3 -plane. The label $--++$ reveals that the corresponding element does not preserve v . Arrows in the middle column indicate translations found in $8Q_C$, they correspond to dots labeled $++++$. The right column of the diagram computes the group elements corresponding to the dots of interest. This is done by considering which translates of sides we pass through as we go from Q_C to the dot.




Figure 4: Translations in $64Q_C$ for manifold M_{56}

Let $v = v_{++++}$, then Q_C is the cube from Figure 1. Here the condition $k_1^{-1} \dots k_m^{-1} v = v$ becomes $k_1^{-1} \dots k_m^{-1} = 1$, since K acts freely on the set of vertices $v_{\pm\pm\pm\pm}$. Moving four steps in the directions of the three axes produces translations because $k_1 = k_3$, $k_2 = k_4$ and $r_1 \dots r_4$ is a composite of reflections in four parallel planes. Let T_v be the subgroup of G_v generated by those three translations and TG_v be the translation subgroup of G_v . For our purposes, we look for translations in $64Q_C$, the $4 \times 4 \times 4$ block of cubes that is the fundamental polyhedron for T_v . It is clear that we get a translation if and only if we move an even number of steps in the direction of each axis, and have $k_1^{-1} \dots k_m^{-1} = 1$. Therefore, we have to check $k_1^{-1} \dots k_m^{-1} = 1$ for the 7 cubes obtained by moving Q_C by a combination of two units in every direction. The dots in Figure 4 depict the centers of some cubes in $64Q_C$, and the ones at the vertices of the cube are the interesting ones. Labels indicate the products of the K -parts, so a $++++$ label means the dot corresponds to a translation. At right we compute the group elements corresponding to the dots of interest. One of the translations is omitted, since it is the sum of the other two in the diagram.

4 Examples of “link” complements in the 4-sphere

In this section we show that 12 nonorientable Ratcliffe-Tschantz manifolds have orientable double covers that are complements of tori and Klein bottles in S^4 . First of all, we need criteria that rule out a manifold from having this property. The first criterion is:

Proposition 4.1 *Let M be a complement of r tori and s Klein bottles in S^4 . Then (coefficients are in \mathbf{Z}):*

$$H_1M = \mathbf{Z}^r \oplus \mathbf{Z}_2^s, \quad H_2M = \mathbf{Z}^{2r} \oplus \mathbf{Z}^s, \quad H_3M = \mathbf{Z}^{r+s-1}.$$

Proof The proof is a simple application of Alexander’s duality: if A is a compact submanifold of S^n , then $\tilde{H}_i(S^4 - A) \cong \tilde{H}^{3-i}(A)$. \square

The other criterion was developed in [6] and exploits a homomorphism that exists for every group G of a Ratcliffe-Tschantz manifold. It is defined by $\phi : G \rightarrow \mathbf{Z}_2^6$, $a, b \mapsto e_1$, $c, d \mapsto e_2, \dots, k, l \mapsto e_6$, where e_i is a canonical generator of \mathbf{Z}_2^6 .

Proposition 4.2 *Let T be the set of all translations in G . If $\dim_{\mathbf{Z}_2} \langle \phi(T) \rangle < 5$ then the orientable double cover of M is not a complement in the 4-sphere. \square*

Using a computer to test the 1149 nonorientable Ratcliffe-Tschantz manifolds for both criteria, we found that only 49 remained eligible to have double-cover complements in S^4 , namely, manifolds numbered

23	25	28	29	32	33	35	36	40	46	47	51	56
60	71	72	73	74	80	92	93	94	96	109	112	116
121	131	132	139	170	171	231	235	236	251	292	296	297
299	426	427	434	435	1011	1091	1092	1094	1095			

If the double cover of any of these manifolds is a complement in S^4 , the potential link structure is visible from its homology by Proposition 4.1. The computer calculation showed that only twelve different link structures are possible. We then proceeded to prove (manually) that for each of those link structures there is at least one manifold from the above list whose orientable double cover is the complement of such a link in S^4 . Those manifolds are listed in Theorem 4.4. Most likely, other manifolds from the list above have the same property, but it is not easy to ascertain that we will be getting examples different from the ones

in Theorem 4.4, since orientable double covers of nonisometric manifolds may be isometric.

Let G be any group, H a normal subgroup of G and let $\langle\langle A \rangle\rangle_H$ be the normal closure of a set $A \subset H$ in H . If G is the biggest group under consideration, $\langle\langle A \rangle\rangle$ or $\langle\langle A \rangle\rangle_G$ will denote the normal closure of A in G . Clearly, $\langle\langle A \rangle\rangle_H$ is the set of all elements of the form $\prod h_i a_i h_i^{-1}$, where $h_i \in H$, $a_i \in A \cup A^{-1}$. If X is a transversal of H in G it is easy to see that for a set $A \subset H$, $\langle\langle A \rangle\rangle_G = \langle\langle xAx^{-1}, x \in X \rangle\rangle_H$. Note that $\langle\langle A \rangle\rangle_G$ is a subgroup of H , since H is normal in G .

While $\langle\langle A \rangle\rangle_H \subseteq \langle\langle A \rangle\rangle_G$, in general, $\langle\langle A \rangle\rangle_H \neq \langle\langle A \rangle\rangle_G$. For example, if $H = \langle a \rangle * \langle b \rangle$, $G = H \rtimes \mathbf{Z}_2$, where \mathbf{Z}_2 acts on H by swapping the generators a and b , then clearly $\langle\langle a \rangle\rangle_G = H$, while $\langle\langle a \rangle\rangle_H$ is a subgroup of H with infinite index.

The following proposition will prove useful.

Proposition 4.3 *Let H be a normal subgroup of G with transversal X . Suppose a set A has the property that $\langle\langle A \rangle\rangle_H = \langle\langle A \rangle\rangle_G$. Let $B \subset H$ have the property: for every $b \in B$ and every $x \in X$, $[xbx^{-1}] = [b^{\pm 1}]$, where $[]$ denotes classes in $G/\langle\langle A \rangle\rangle_G$.*

Then $\langle\langle A \cup B \rangle\rangle_H = \langle\langle A \cup B \rangle\rangle_G$.

Proof Elements of type $hxa(hx)^{-1}$ and $hxb(hx)^{-1}$ generate the subgroup $\langle\langle A \cup B \rangle\rangle_G$, where $a \in A \cup A^{-1}$, $b \in B \cup B^{-1}$, $x \in X$, $h \in H$. Now $\langle\langle xAx^{-1}, x \in X \rangle\rangle_H = \langle\langle A \rangle\rangle_G = \langle\langle A \rangle\rangle_H$ implies that every element xax^{-1} is a product of elements of type hah^{-1} , so every $hxa(hx)^{-1} = h(xax^{-1})h^{-1}$ has the same property. Since $[xbx^{-1}] = [b^{\pm 1}]$ in $G/\langle\langle A \rangle\rangle_G$ means that $xbx^{-1} = b^{\pm 1}u$, for some $u \in \langle\langle A \rangle\rangle_G = \langle\langle A \rangle\rangle_H$, xbx^{-1} is a product of $b^{\pm 1}$ and elements of type hah^{-1} , making $hxb(hx)^{-1}$ a product of elements of type hbh^{-1} and hah^{-1} . Therefore, $\langle\langle A \cup B \rangle\rangle_G$ is generated by elements of type hah^{-1} and hbh^{-1} , so $\langle\langle A \cup B \rangle\rangle_G = \langle\langle A \cup B \rangle\rangle_H$. \square

Theorem 4.4 *The orientable double covers of the nonorientable Ratcliffe-Tschantz manifolds listed in Table 1 are complements of the indicated combination of tori and Klein bottles in a manifold that is homeomorphic to the 4-sphere.*

The orientable double cover of manifold	is a complement in S^4 of:
no. 1011	5 tori
no. 71	6 tori
no. 23	7 tori
no. 1092	8 tori
no. 1091	9 tori
no. 231	3 tori, 4 Klein bottles
no. 112	4 tori, 2 Klein bottles
no. 56	4 tori, 3 Klein bottles
no. 92	5 tori, 1 Klein bottle
no. 51	5 tori, 2 Klein bottles
no. 40	6 tori, 1 Klein bottle
no. 36	6 tori, 2 Klein bottles

Table 1

Remark 4.5 Note that Theorem 4.4 claims that double covers of various Ratcliffe-Tschantz manifolds are complements in a manifold that is *homeomorphic* to the 4-sphere, rather than diffeomorphic. This is because we used Freedman's theory, applicable only to the topological category, to recognize a 4-sphere. It is still unknown whether manifolds homeomorphic to the 4-sphere are diffeomorphic to the S^4 with the standard differentiable structure. Using Kirby diagrams of a 4-manifold, Ivanišić has shown that the topological S^4 inside which \widetilde{M}_{1011} is a complement of 5 tori is indeed diffeomorphic to the standard differentiable S^4 . We anticipate that the same is true for the examples treated here.

Proof Let M be a nonorientable Ratcliffe-Tschantz manifold, \widetilde{M} its orientable double cover, $G = \pi_1 M$, $H = \pi_1 \widetilde{M}$. Clearly, H is the index-2 subgroup of G containing all the orientation preserving isometries. If we can find primitive normal translations t_1, \dots, t_m , one in each boundary component of \widetilde{M} , then \widetilde{M} is a complement in a closed manifold N . As was shown in [5], the Euler characteristics of \widetilde{M} and N are equal. By Proposition 2.2 $\pi_1 N = H / \langle\langle t_1, \dots, t_m \rangle\rangle_H$. The classification of 4-dimensional simply-connected manifolds (see [2] or [3]) gives that if $\pi_1 N = 1$ and $\chi(N) = 2$, then N is homeomorphic to S^4 . Since $\chi(\widetilde{M}) = 2$, it will be enough to show that $H / \langle\langle t_1, \dots, t_m \rangle\rangle_H = 1$.

The presentation of G has 12 generators and 24 relations so the presentation of the index-2 subgroup H obtained by the Reidemeister-Schreier method will have 23 generators and 48 relations, very tedious to compute with. Thus, the challenge is to show that $H / \langle\langle t_1, \dots, t_m \rangle\rangle_H = 1$ while working with the

presentation of G as much as possible.

The ten homeomorphism types of flat 3-manifolds are identified by the letters A–J, where the manifolds are ordered in the same way as in [4] or [11]. Every nonorientable Ratcliffe-Tschantz manifold that passes both criteria from above has boundary components of type A, B, G, H, I or J. Type A is the 3-torus and type B is an (orientable) S^1 -bundle over a Klein bottle. The nonorientable types G and H have type A as their orientable double cover and the nonorientable types I and J have type B as their orientable double cover. Holonomy groups of boundary components of M are computed in the course of searching for suitable translations. They help determine whether chosen translations are normal in a boundary component and they distinguish types A, B, G or H, and I or J. Although not essential for our proof, types G and H can further be distinguished by the fact that at least one generator of the translation subgroup of H is not a normal translation. There is a criterion that distinguishes I and J as well.

Note that orientable boundary components of M lift to two homeomorphic boundary components in \widetilde{M} and nonorientable ones lift to their orientable double covers. Thus, knowing the boundary components of M immediately gives the link structure that \widetilde{M} is a complement of (a torus or a Klein bottle for every component of type A or B, respectively).

Translations in boundary components of \widetilde{M} have to be carefully chosen in order to be S^1 -fibers, to produce $H/\langle\langle t_1, \dots, t_m \rangle\rangle_H = 1$ and to allow us to work in G . It was shown in [6] that the homomorphism ϕ from Proposition 4.2 sends to 0 every subgroup T_v in §3, and that $H/\langle\langle t_1, \dots, t_m \rangle\rangle_H = 1$ implies $\dim_{\mathbf{Z}_2} \langle\phi(t_1), \dots, \phi(t_m)\rangle \geq 5$. Therefore, one should choose translations in $8Q_C$ or $64Q_C$ whenever possible, since one risks otherwise that $\dim_{\mathbf{Z}_2} \langle\phi(t_1), \dots, \phi(t_m)\rangle$ comes short of 5. A translation of type $t = t' + t''$ where $t' \in T_v$ and $t'' \in 8Q_C$ or $64Q_C$, is a possible choice, but we avoid them, because its length in terms of generators a – l is big, possibly complicating the computation.

Let E be a boundary component of M and \widetilde{E} a component of its lift in \widetilde{M} . In order for \widetilde{M} to be a complement, we must choose a translation t that is normal and primitive in \widetilde{E} . Note that t need not be normal in E , but it is very useful if it is when E is nonorientable. For in that case, there is an orientation-reversing $x \in G$ so that $xtx^{-1} = t^{\pm 1}$, which by Proposition 4.3 implies $\langle\langle A, t \rangle\rangle_H = \langle\langle A, t \rangle\rangle_G$ for any subset A of H satisfying $\langle\langle A \rangle\rangle_H = \langle\langle A \rangle\rangle_G$. This allows us to continue computing in G . We introduce a term for this

property: we say that t is *self-conjugate* if there exists an orientation-reversing $x \in G$ so that $xtx^{-1} = t^{\pm 1}$.

Checking that t is primitive in \tilde{E} is easy because \tilde{E} is either of type A or B. In the first case, t must be the shortest translation in its direction, in the second, it must not point in the same direction as the axis of rotation by π that generates the holonomy group.

If E is orientable, its lift has two components \tilde{E} and \tilde{E}' , whose fundamental groups are conjugate by an orientation reversing $x \in G$. Choosing a primitive normal translation t in \tilde{E} gives the obvious choice xtx^{-1} for \tilde{E}' . This choice works most of the time, but on occasion, like in Example 4.8, one must choose a different translation in \tilde{E}' in order to satisfy the condition $\dim_{\mathbf{Z}_2} \langle \phi(t_1), \dots, \phi(t_m) \rangle \geq 5$.

To prove the theorem, each example needs to be handled separately. We show three of the most difficult examples in decreasing detail and indicate how to proceed on others.

Example 4.6 The side-pairing of manifold M_{56} is encoded by 13D935. According to our side-pairing convention, the side-pairings are (the K -parts are under the arrow):

$$\begin{array}{ll}
 S_{++00} \xrightarrow[-++++]{a} S_{-+00} & S_{+-00} \xrightarrow[-++++]{b} S_{--00} \\
 S_{+0+0} \xrightarrow[-+++]{c} S_{-0+0} & S_{+0-0} \xrightarrow[-+++]{d} S_{-0-0} \\
 S_{0++0} \xrightarrow[-+---]{e} S_{0+-0} & S_{0-+0} \xrightarrow[-+---]{f} S_{0--0} \\
 S_{+00+} \xrightarrow[-++-]{g} S_{-00-} & S_{+00-} \xrightarrow[-++-]{h} S_{-00+} \\
 S_{0+0+} \xrightarrow[-+++]{i} S_{0-0+} & S_{0+0-} \xrightarrow[-+++]{j} S_{0-0-} \\
 S_{00++} \xrightarrow[-+++]{k} S_{00-+} & S_{00+-} \xrightarrow[-+++]{l} S_{00--}
 \end{array}$$

The orientation-reversing generators are c , d and $g-l$ (those whose K -part has an even number of minuses in its label).

We use diagrams in Figure 2 to find the cycle relations as described in §3. The arrangement of groups of relators below corresponds to the arrangement of diagrams used to obtain them and the numbers 1–4 are labels of the vertices in the diagrams corresponding to each cycle relation. The cycle relations marked

by • are repetitions of those from the left column, so we reflect the section plane in S_{00++} and obtain the two new relations in the right column.

$$\begin{array}{lll}
 1) & g^{-1}j^{-1}h^{-1}i = 1 & hj^{-1}gi = 1 \\
 2) & g^{-1}ch^{-1}c = 1 & \bullet hc^{-1}gc^{-1} = 1 \quad hd^{-1}gd^{-1} = 1 \\
 3) & e^{-1}j^{-1}fi = 1 & \bullet e^{-1}j^{-1}fi = 1 \quad ej^{-1}f^{-1}i = 1 \\
 4) & e^{-1}dfc = 1 & e^{-1}d^{-1}fc^{-1} = 1 \\
 \\
 1) & g^{-1}l^{-1}h^{-1}k = 1 & hl^{-1}gk = 1 \\
 2) & g^{-1}ah^{-1}a = 1 & hb^{-1}gb^{-1} = 1 \\
 3) & e^{-1}le^{-1}k = 1 & f^{-1}lf^{-1}k = 1 \\
 4) & e^{-1}aea = 1 & f^{-1}b^{-1}fb^{-1} = 1 \\
 \\
 1) & i^{-1}k^{-1}ik = 1 & j^{-1}ljl^{-1} = 1 \\
 2) & i^{-1}bia = 1 & j^{-1}b^{-1}ja^{-1} = 1 \\
 3) & c^{-1}k^{-1}d^{-1}k = 1 & dlcl^{-1} = 1 \\
 4) & c^{-1}bc^{-1}a = 1 & db^{-1}da^{-1} = 1
 \end{array}$$

The diagrams in Figures 3 and 4 find the translations of interest for M_{56} . The following table identifies boundary component types and the translations. \widetilde{M}_{56} has boundary components of type BBBAAAA, making it a complement of 3 Klein bottles and 4 tori. Boundary components E_1 – E_4 are the ones corresponding to ideal vertices $v_{+000}-v_{000+}$, while E_5 corresponds to vertex v_{++++} . Elements of the holonomy group are identified by their planes of reflection, for example, x_2x_3 is the composite of reflections in planes x_2 and x_3 .

comp.	type	translations in $8Q_C$ or $64Q_C$	holonomy	self-conj?
E_1	I	none	x_2, x_3, x_2x_3	n/a
E_2	B	a	x_1x_4	no
E_3	H	$e^{-1}dl$	x_2	no
E_4	G	$i^{-1}k^{-1}$	x_1	yes
E_5	A	$c^{-1}i, a^{-1}k^{-1}eg$	trivial	no

Figure 3 shows why $e^{-1}dl$ is not normal in E_3 : reflecting the vector in the third cube in the plane $x_2 = 0$ produces a nonparallel vector. Similarly, $i^{-1}k^{-1}$ is normal in E_4 because it is invariant under reflection in the plane $x_1 = 0$. Note that there are no translations in E_1 that are in the cube $8Q_C$, so we use one of the translations $a^{-1}b, c^{-1}d^{-1}$, or $g^{-1}h$ that generate $T_{v_{+000}}$. The nontrivial element of holonomy of \widetilde{E}_1 is a rotation in the x_2x_3 -plane. All three

translations are thus normal in \tilde{E}_1 , but the third one is not primitive, since it is a power of a slide-rotation. Make the following choices for translations:

$$t_1 = c^{-1}d, \quad t_2 = a, \quad t_3 = e^{-1}dl, \quad t_4 = i^{-1}k^{-1}, \quad t_5 = c^{-1}i, \\ t'_2 = ct_2c^{-1}, \quad t'_5 = c(a^{-1}k^{-1}eg)c^{-1}.$$

Because t_1 and t_4 are self-conjugate, we have $\langle\langle t_1, t_2, t_4 \rangle\rangle_G = \langle\langle t_1, t_2, t'_2, t_4 \rangle\rangle_H$. Adding $t_1 = t_2 = t_4 = 1$ to the relators we get $d = c$, $k = i^{-1}$ and $a = 1$, which immediately yields $b = 1$ and $h = g^{-1}$, so the presentation for $G/\langle\langle t_1, t_2, t_4 \rangle\rangle$ has generators c, f, g, i, j, l and relations (repeating or trivial relations omitted):

$$\left\{ \begin{array}{l} g^{-1}j^{-1}gi = 1 \\ g^{-1}cgc = 1 \\ e^{-1}j^{-1}fi = 1 \\ e^{-1}cfc = 1 \\ ej^{-1}f^{-1}i = 1 \\ e^{-1}c^{-1}fc^{-1} = 1 \end{array} \right\} \left\{ \begin{array}{l} g^{-1}l^{-1}gi^{-1} = 1 \\ e^{-1}le^{-1}i^{-1} = 1 \\ f^{-1}lf^{-1}i^{-1} = 1 \end{array} \right\} \left\{ \begin{array}{l} c^{-1}ic^{-1}i^{-1} = 1 \\ c^{-2} = 1 \\ j^{-1}ljl^{-1} = 1 \\ clcl^{-1} = 1 \end{array} \right.$$

Taking quotients is equivalent to adding relators so, as is customary in working with group presentations, we usually omit class notation. Taking into account $c^2 = 1$, note that one relation in the group at right says that i and c commute in $G/\langle\langle t_1, t_2, t_4 \rangle\rangle$, and one relation in the left group says g and c commute ($c = c^{-1}$). But then $ct_5c^{-1} = cc^{-1}ic^{-1} = t_5$, fulfilling conditions of Proposition 4.3, so we conclude $\langle\langle t_1, t_2, t_4, t_5 \rangle\rangle_G = \langle\langle t_1, t_2, t'_2, t_4, t_5 \rangle\rangle_H$.

In the group $G/\langle\langle t_1, t_2, t_4, t_5 \rangle\rangle$, we now have $i = c$, so $j = gig^{-1} = gcg^{-1} = c$ and $l = gi^{-1}g^{-1} = gc^{-1}g^{-1} = c$, which in conjunction with $e^{-1}le^{-1}i^{-1} = 1$ implies $e^{-1}ce^{-1}c^{-1} = 1$, in other words, $cec^{-1} = e^{-1}$. Since now $t_3 = e^{-1}$, we have $ct_3c^{-1} = ce^{-1}c^{-1} = e = t_3^{-1}$, so Proposition 4.3 yields $\langle\langle t_1, t_2, t_3, t_4, t_5 \rangle\rangle_G = \langle\langle t_1, t_2, t'_2, t_3, t_4, t_5 \rangle\rangle_H$.

The presentation of the group $G/\langle\langle t_1, t_2, t_3, t_4, t_5 \rangle\rangle$ is obtained from the above relations combined with $i = c$ and $e = 1$ (following from $t_5 = t_3 = 1$). Then $f = 1$ follows. Using $j = l = c$, the presentation simplifies to

$$\langle c, g \mid c^2 = 1, cgc^{-1} = g \rangle = \mathbf{Z} \oplus \mathbf{Z}_2.$$

In this group, $t'_5 = ca^{-1}k^{-1}egc^{-1} = c1c1gc^{-1} = gc^{-1} = gc$, so $ct'_5c^{-1} = t'_5$. Proposition 4.3 says $\langle\langle t_1, t_2, t_3, t_4, t_5, t'_5 \rangle\rangle_G = \langle\langle t_1, t_2, t'_2, t_3, t_4, t_5, t'_5 \rangle\rangle_H$. Now $G/\langle\langle t_1, t_2, t_3, t_4, t_5, t'_5 \rangle\rangle = \langle c \mid c^2 = 1 \rangle = \mathbf{Z}_2$, so $H/\langle\langle t_1, t_2, t'_2, t_3, t_4, t_5, t'_5 \rangle\rangle_H = H/\langle\langle t_1, t_2, t_3, t_4, t_5, t'_5 \rangle\rangle_G = 1$. Therefore, \tilde{M}_{56} is a complement in the 4-sphere.

Example 4.7 The side-pairing of manifold M_{1091} is encoded by 53RR35. All the generators are orientation reversing. The presentation of G is given by the following relations, which are grouped by row of diagrams like in Figure 2.

$$\left\{ \begin{array}{l} g^{-1}jh^{-1}i = 1 \\ g^{-1}dh^{-1}c = 1 \\ e^{-1}jf^{-1}i = 1 \\ e^{-1}df^{-1}c = 1 \\ hjgi = 1 \\ hc^{-1}gd^{-1} = 1 \\ f^{-1}je^{-1}i = 1 \\ f^{-1}c^{-1}e^{-1}d^{-1} = 1 \end{array} \right. \left\{ \begin{array}{l} g^{-1}lh^{-1}k = 1 \\ g^{-1}bh^{-1}a = 1 \\ e^{-1}lfk = 1 \\ e^{-1}bfa = 1 \\ hlgk = 1 \\ ha^{-1}gb^{-1} = 1 \\ fle^{-1}k = 1 \\ fa^{-1}e^{-1}b^{-1} = 1 \end{array} \right. \left\{ \begin{array}{l} i^{-1}k^{-1}ik = 1 \\ i^{-1}bia = 1 \\ c^{-1}k^{-1}d^{-1}k = 1 \\ c^{-1}bd^{-1}a = 1 \\ jlj^{-1}l^{-1} = 1 \\ ja^{-1}j^{-1}b^{-1} = 1 \\ dlcl^{-1} = 1 \\ da^{-1}cb^{-1} = 1 \end{array} \right.$$

The table below lists the translations of interest to us in the boundary components. The boundary components of \widetilde{M}_{1091} are AAAAAAAAAA, making it a complement of 9 tori. The boundary component E_6 is the one corresponding to vertex v_{+++-} , for which the search for translations is the same as for v_{++++} .

comp.	type	translations in $8Q_C$ or $64Q_C$	holonomy	self-conj?
E_1	A	$c^{-1}h, a^{-1}h, c^{-1}b$	trivial	no
E_2	H	$e^{-1}j$	x_3	no
E_3	H	$e^{-1}l$	x_2	no
E_4	G	$i^{-1}k^{-1}$	x_1	yes
E_5	A	$a^{-1}k, c^{-1}i, e^{-1}g$	trivial	no
E_6	A	$a^{-1}l, c^{-1}j, e^{-1}h$	trivial	no

Choose the translations:

$$t_1 = c^{-1}b, \quad t_2 = e^{-1}j, \quad t_3 = e^{-1}l, \quad t_4 = i^{-1}k^{-1}, \quad t_5 = c^{-1}i, \quad t_6 = e^{-1}h, \\ t'_1 = ct_1c^{-1}, \quad t'_5 = ct_5c^{-1}, \quad t'_6 = ct_6c^{-1}.$$

Due to self-conjugacy of t_4 , $\langle\langle t_1, t_4, t_5, t_6 \rangle\rangle_G = \langle\langle t_1, t'_1, t_4, t_5, t'_5, t_6, t'_6 \rangle\rangle_H$. In what follows, we refer to the relations above as entries in an 8×3 matrix. In the group $G/\langle\langle t_1, t_4, t_5, t_6 \rangle\rangle$ we have $k^{-1} = i = c = b$ and $h = e$, so equation 43 gives $d = a$ and equation 23 gives $b = a^{-1}$. Equations 32 and 42 convert to $l = ea^{-1}f^{-1}$ and $a^{-1} = ea^{-1}f^{-1}$, so $l = a^{-1}$. Similarly, comparing 11 and 21 gives $j = a$. The presentation of $G/\langle\langle t_1, t_4, t_5, t_6 \rangle\rangle$ then reduces to generators a, e, g and f and the relations, grouped as above (third column reduces to trivial relations):

$$\left\{ \begin{array}{l} g^{-1}ae^{-1}a^{-1} = 1 \\ e^{-1}af^{-1}a^{-1} = 1 \\ eaga^{-1} = 1 \\ f^{-1}ae^{-1}a^{-1} = 1 \end{array} \right. \left\{ \begin{array}{l} g^{-1}a^{-1}e^{-1}a = 1 \\ e^{-1}a^{-1}fa = 1 \\ ea^{-1}ga = 1 \\ fa^{-1}e^{-1}a = 1 \end{array} \right.$$

In this array equations 11 and 41 give $f = g$, equations 22 and 32 imply $e = e^{-1}$, and equations 12 and 42 give $g = f^{-1}$, further simplifying the presentation to

$$\langle a, e, f \mid e^2 = f^2 = 1, aea^{-1} = f, afa^{-1} = e \rangle \cong (\mathbf{Z}_2 * \mathbf{Z}_2) \rtimes \mathbf{Z},$$

where \mathbf{Z} acts on $\mathbf{Z}_2 * \mathbf{Z}_2$ by swapping the generators. One can now show (next paragraph) that $H / \langle \langle t_1, t_4, t_5, t_6 \rangle \rangle \cong \mathbf{Z} \oplus \mathbf{Z}$, where the generators are $[t_2] = e^{-1}a$ and $[t_3] = e^{-1}a^{-1}$. Let $H_1 = H / \langle \langle t_1, t'_1, t_4, t_5, t'_5, t_6, t'_6 \rangle \rangle_H$. We now have

$$\begin{aligned} H / \langle \langle t_1, t'_1, t_2, t_3, t_4, t_5, t'_5, t_6, t'_6 \rangle \rangle_H &= \\ (H / \langle \langle t_1, t'_1, t_4, t_5, t'_5, t_6, t'_6 \rangle \rangle_H) / \langle \langle [t_2], [t_3] \rangle \rangle_{H_1} &= \\ (H / \langle \langle t_1, t_4, t_5, t_6 \rangle \rangle) / \langle \langle [t_2], [t_3] \rangle \rangle_{H_1} &= \\ \langle \langle [t_2] \rangle \oplus \langle [t_3] \rangle \rangle / \langle \langle [t_2], [t_3] \rangle \rangle_{H_1} &= 1. \end{aligned}$$

To show that $H / \langle \langle t_1, t_4, t_5, t_6 \rangle \rangle \cong \mathbf{Z} \oplus \mathbf{Z}$ one can use either the explicit isomorphism of $G / \langle \langle t_1, t_4, t_5, t_6 \rangle \rangle$ with $(\mathbf{Z}_2 * \mathbf{Z}_2) \rtimes \mathbf{Z}$, or apply the Reidemeister-Schreier method to the presentation above using $\{1, e\}$ as the transversal. In the latter case, we get the presentation

$$\langle a_0, f_0, a_1, f_1 \mid f_0 f_1 = a_0 a_1^{-1} f_0^{-1} = a_0 f_1 a_1^{-1} = a_1 a_0^{-1} f_1^{-1} = a_1 f_0 a_0^{-1} = 1 \rangle,$$

where $a_0 = ae^{-1}$, $f_0 = fe^{-1}$, $a_1 = ea$, and $f_1 = ef$. This easily reduces to $\langle a_0, a_1 \mid a_0 a_1 = a_1 a_0 \rangle$. We also note $[t_2] = e^{-1}a = ea = a_1$ and $[t_3] = e^{-1}a^{-1} = (ae^{-1})^{-1} = a_0^{-1}$.

Example 4.8 The side-pairing of manifold M_{36} is encoded by 1468AF. The orientation-reversing generators are e, f, i, j, k, l . The presentation of G is given by the following relations.

$$\left\{ \begin{array}{l} g^{-1}j^{-1}g^{-1}i = 1 \\ g^{-1}c^{-1}gc = 1 \\ e^{-1}j f^{-1}i = 1 \\ e^{-1}c f c = 1 \\ h^{-1}j^{-1}h^{-1}i = 1 \\ h^{-1}d^{-1}hd = 1 \\ e j^{-1} f i^{-1} = 1 \\ e^{-1}d f d = 1 \end{array} \right\} \left\{ \begin{array}{l} g^{-1}l^{-1}h^{-1}k = 1 \\ g^{-1}a^{-1}ha = 1 \\ e^{-1}l e^{-1}k = 1 \\ e^{-1}b^{-1}ea = 1 \\ g^{-1}k h^{-1}l^{-1} = 1 \\ gb^{-1}h^{-1}b = 1 \\ f^{-1}k^{-1}f^{-1}l^{-1} = 1 \\ f^{-1}b^{-1}fa = 1 \end{array} \right\} \left\{ \begin{array}{l} i^{-1}l^{-1}i^{-1}k = 1 \\ i^{-1}b^{-1}ia = 1 \\ c^{-1}l d^{-1}k = 1 \\ c^{-1}a^{-1}da = 1 \\ j k j l^{-1} = 1 \\ j a^{-1} j^{-1} b = 1 \\ c^{-1}k d^{-1}l = 1 \\ c b^{-1} d^{-1} b = 1 \end{array} \right.$$

The following table identifies boundary component types and translations of interest. The boundary components of \bar{M}_{36} are AAAABBAA making it a complement of 6 tori and 2 Klein bottles.

comp.	type	translations in $8Q_C$ or $64Q_C$	holonomy	self-conj?
E_1	A	c^{-1}, g^{-1}	trivial	no
E_2	A	$a^{-1}, e^{-1}j$	trivial	no
E_3	J	none	x_1, x_2, x_1x_2	n/a
E_4	J	none	x_1, x_2, x_1x_2	n/a
E_5	A	$c^{-1}i^{-1}eg, a^{-1}k^{-1}ci, a^{-1}k^{-1}eg$	trivial	no

Choose the translations:

$$t_1 = a^{-1}b, \quad t_2 = e^{-1}j, \quad t_3 = e^{-1}f^{-1}, \quad t_4 = i^{-1}j^{-1}, \quad t_5 = a^{-1}k^{-1}ci,$$

$$t'_1 = kck^{-1}, \quad t'_2 = eae^{-1}, \quad t'_5 = k(c^{-1}i^{-1}eg)k^{-1}.$$

Note that for boundary component E_1 , although there are two translations in $8Q_C$, we choose $a^{-1}b \in \ker \phi$ because neither c nor g is self-conjugate. Note also that in boundary components E_3 and E_4 there are no translations in $8Q_C$ so we are forced to choose translations in $\ker \phi$. Lifts of E_3 and E_4 are of type B , so care must be taken that the translations are primitive in the lifts. In other words, the translations must be in the planes of rotation (in this case, the plane x_1x_2).

Since t_3 and t_4 are normal in nonorientable E_3 and E_4 , they are self-conjugate, so $\langle\langle t_3, t_4 \rangle\rangle_H = \langle\langle t_3, t_4 \rangle\rangle_G$. In $G/\langle\langle t_3, t_4 \rangle\rangle$ we have $f = e^{-1}$ and $j = i^{-1}$. Equations 42 and 82 then give $eae^{-1} = b$, $ebe^{-1} = a$, so $e(a^{-1}b)e^{-1} = b^{-1}a = (a^{-1}b)^{-1}$, that is, $et_1e^{-1} = t_1^{-1}$. Equation 31 reads as $e^{-1}i^{-1}ei = 1$, so $et_2e^{-1} = t_2$ because e and $j = i^{-1}$ commute. Then $\langle\langle t_1, t_2, t_3, t_4 \rangle\rangle_H = \langle\langle t_1, t_2, t_3, t_4 \rangle\rangle_G$.

In $G/\langle\langle t_1, t_2, t_3, t_4 \rangle\rangle$ we have $a = b$, which gives $eae^{-1} = a$, implying

$$\langle\langle t_1, t_2, t'_2, t_3, t_4 \rangle\rangle_H = \langle\langle t_1, t_2, t'_2, t_3, t_4 \rangle\rangle_G.$$

In $G/\langle\langle t_1, t_2, t'_2, t_3, t_4 \rangle\rangle$ we then have $a = 1$ which implies $b = 1$, $h = g$ and $d = c$. Equation 32 gives $l = ek^{-1}e$. Simplifying the presentation, we get

$$G/\langle\langle t_1, t_2, t'_2, t_3, t_4 \rangle\rangle = \langle c, e, g, k \mid egeg = g^{-1}c^{-1}gc = e^{-1}ce^{-1}c = kgekeg = k^2 = c^{-1}ekc^{-1}k = 1 \rangle.$$

Unfortunately, the remaining translations $t'_1 = kck^{-1}$, $t_5 = kce^{-1}$ and $t'_5 = kc^{-1}e^2gk^{-1}$ cannot easily be shown to be self-conjugate in this group, so we resort to the Reidemeister-Schreier method to get $H/\langle\langle t_1, t_2, t'_2, t_3, t_4 \rangle\rangle$ before we continue. Using $\{1, k\}$ as the transversal, it is not hard to see that the presentation for $H/\langle\langle t_1, t_2, t'_2, t_3, t_4 \rangle\rangle$ has generators $c_0 = c$, $g_0 = g$, $e_0 = ek^{-1}$, $c_1 = kck^{-1}$, $g_1 = k g k^{-1}$, $e_1 = ke$ and relations

$$\begin{aligned} e_0g_1e_1g_0 = 1 & \quad g_0^{-1}c_0^{-1}g_0c_0 = 1 & \quad e_1^{-1}c_1e_0^{-1}c_0 = 1 & \quad g_1e_1^2g_0 = 1 & \quad c_0^{-1}e_0^2c_1^{-1} = 1 \\ e_1g_0e_0g_1 = 1 & \quad g_1^{-1}c_1^{-1}g_1c_1 = 1 & \quad e_0^{-1}c_0e_1^{-1}c_1 = 1 & \quad g_0e_0^2g_1 = 1 & \quad c_1^{-1}e_1^2c_0^{-1} = 1 \end{aligned}$$

The translations take the form $t'_1 = c_1$, $t_5 = c_1 e_0^{-1}$, $t'_5 = c_1^{-1} e_1 e_0 g_1$. Adding $t'_1 = t_5 = t'_5 = 1$ to the above relations immediately gives $c_1 = e_0 = 1$, $g_1 = e_1^{-1}$. Relation 15 then reduces to $c_0 = 1$ so 13 now says $e_1 = 1$, hence $g_1 = 1$. Then $g_0 = 1$ follows from relation 21, therefore, $H / \langle \langle t_1, t'_1, t_2, t'_2, t_3, t_4, t_5, t'_5 \rangle \rangle_H = 1$.

In this example, infinitely many other choices for t_5 and t'_5 also give the 4-sphere. If we only add $c_1 = 1$ to the above equations, we get $c_0 = e_0^2$, $e_1 = e_0$ and $g_1 = e_0^{-2} g_0^{-1}$, and the presentation simplifies to $\langle e_0, g_0 \mid e_0 g_0 e_0^{-1} g_0^{-1} = 1 \rangle$. Thus, $H / \langle \langle t_1, t'_1, t_2, t'_2, t_3, t_4 \rangle \rangle_H = \langle e_0 \rangle \oplus \langle g_0 \rangle$. The basis elements of $\pi_1 \tilde{E}_5 = \mathbf{Z}^3$ are $v_1 = c^{-1} i^{-1} e g$, $v_2 = a^{-1} k^{-1} c i$, and $v_3 = a^{-1} k^{-1} e g$; they are sent to g_0 , e_0^{-1} and $e_0 g_0$. The conjugates $v'_i = k v_i k^{-1}$ generate $\pi_1 \tilde{E}'_5$; they are sent to g_0^{-1} , e_0 and $e_0^{-1} g_0^{-1}$. Any primitive translations in \tilde{E}_5 , \tilde{E}'_5 then have form $t_5 = p v_1 - q v_2 + r v_3$ or $t'_5 = -p' v'_1 + q' v'_2 - r' v'_3$ where $\gcd(p, q, r) = \gcd(p', q', r') = 1$. If $\psi : H \rightarrow H / \langle \langle t_1, t'_1, t_2, t'_2, t_3, t_4 \rangle \rangle_H$ is the quotient map, then $\psi(t_5) = (q + r, p + r)$ and $\psi(t'_5) = (q' + r', p' + r')$, as written in terms of the basis $\{e_0, g_0\}$. Then

$$H / \langle \langle t_1, t'_1, t_2, t'_2, t_3, t_4, t_5, t'_5 \rangle \rangle_H = 1 \iff \begin{vmatrix} q + r & q' + r' \\ p + r & p' + r' \end{vmatrix} = 1.$$

Infinitely many choices of p, q, r, p', q', r' exist that satisfy the last condition. For example, setting $r = r' = 0$, it is easy to see that the resulting equation $qp' - pq' = 1$ can be satisfied for an arbitrary relatively prime pair of numbers p and q once suitable q' and r' are found (q' and r' are consequently relatively prime).

Example 4.9 For the remaining examples from Theorem 4.4, the proofs proceed in a similar way. In Table 2 we list the choices for translations that make the orientable double cover a complement in S^4 . Types of boundary components are identified in the order we have used here, note that this order is different from tables in [9]. The translations are listed in the order one sets them equal to 1. The letters RS, if applicable, stand in the place where we were forced to pass to a presentation of $H / \langle \langle \text{translations} \rangle \rangle$ via the Reidemeister-Schreier method. Examples that do not employ it are typically straightforward. Appearances of translations labeled t' indicate that the corresponding boundary component lifted to two components in the orientable double cover.

This completes the proof of Theorem 4.4. □

number	side-pairing	boundary	translation choice
1011	14FF28	GGGGG	$t_1 = c, t_2 = a, t_3 = k, t_4 = i,$ $t_5 = e^{-1}g$ (see [6])
71	13EB34	GGHGA	$t_1 = a^{-1}h, t_2 = a, t_4 = k,$ $t_5 = c^{-1}i, t'_5 = ct_5c^{-1}, t_3 = e^{-1}l$
23	1569A4	GAGAH	$t_1 = c^{-1}h, t_2 = a, t'_2 = ct_2c^{-1},$ $t_3 = e^{-1}d^{-1}, t_4 = k,$ $t'_4 = ct_4c^{-1}, t_5 = c^{-1}i^{-1}eg$
1092	53FFCA	AGGAGG	$t_1 = c^{-1}h, t'_1 = ct_1c^{-1}, t_2 = a^{-1}i^{-1},$ $t_3 = c^{-1}k^{-1}, t_4 = i^{-1}h^{-1}, t'_4 = ct_4c^{-1}$ $t_5 = e^{-1}g, t_6 = e^{-1}h$
231	1569F4	GBGBH	$t_1 = c^{-1}h, t_2 = a, t'_2 = ct_2c^{-1},$ $t_3 = e^{-1}d^{-1}, t_4 = k, t'_4 = ct_4c^{-1},$ $t_5 = a^{-1}k^{-1}ci^{-1}eg$
112	13C874	GGHBG	$t_1 = g, t_2 = a, t_4 = k, t'_4 = ct_4c^{-1},$ $t_5 = c^{-1}ieg, t_3 = e^{-1}d^{-1}l$
92	1348EC	GAHIH	$t_1 = g, t_4 = k^{-1}l^{-1},$ $t_2 = e, t_3 = e^{-1}d^{-1}l,$ RS, $t'_2 = cac^{-1}, t_5 = c^{-1}jak$
51	156A9C	HGABG	$t_2 = a, t_3 = c^{-1}l, t'_3 = ct_3c^{-1},$ $t_4 = i^{-1}h^{-1}, t'_4 = ct_4c^{-1},$ $t_5 = c^{-1}i^{-1}f^{-1}g, t_1 = c^{-1}ag^{-1}$
40	143CF9	GAGAJ	$t_1 = c, t_3 = e^{-1}k^{-1}, t_5 = e^{-1}h^{-1}fg,$ $t_4 = g^{-1}l, t_2 = a,$ RS, $t'_2 = i(e^{-1}j)i^{-1}, t'_4 = i(i^{-1}h)i^{-1}$

Table 2

We conclude with some related examples.

Example 4.10 (Examples of complements in simply-connected manifolds with higher Euler characteristic) It was shown in [6] that index- n cyclic covers of \widetilde{M}_{1011} are complements of $4n + 1$ tori in a simply-connected closed manifold N with Euler characteristic $2n$. We show that many other manifolds from Theorem 4.4 have an analogous property. Two propositions from [6] are used for this purpose.

Let M be a hyperbolic manifold with boundary components E_1, \dots, E_m and let

$H = \pi_1 M$, $K_i = \pi_1 E_i$. If we take the cover corresponding to a normal subgroup H_0 of H , [6] Proposition 2.3 asserts that the number of path-components of $p^{-1}(E_i)$ is equal to the index of the image of K_i in H/H_0 under the quotient map. Now let $t_i \in \pi_1 E_i$ be a translation normal in E_i , $i = 1, \dots, m$. Setting $H_0 = \langle\langle t_1, \dots, t_m \rangle\rangle_H$, suppose that H/H_0 is a finite group of order l , and that t_i is primitive $K_i \cap H_0$. Then [6] Proposition 2.4 states that the index- l cover of M corresponding to subgroup H_0 is a complement inside a simply-connected closed manifold.

From example 4.6 it is easily seen that $H/\langle\langle t_1, t_2, t'_2, t_3, t_4, t_5 \rangle\rangle_H = \mathbf{Z} = \langle gc \rangle = \langle [t'_5] \rangle$. It follows that $H/\langle\langle t_1, t_2, t'_2, t_3, t_4, t_5, (t'_5)^m \rangle\rangle_H = \mathbf{Z}_m$, so the above discussion shows that an m -fold cover of \widetilde{M}_{56} is a complement in a simply-connected 4-manifold N with $\chi(N) = 2m$. To find the link type, consider the generators of boundary components of \widetilde{M}_{56} and their images under the quotient map $H \rightarrow H/\langle\langle t_1, t_2, t'_2, t_3, t_4, t_5 \rangle\rangle_H$:

component	type	generators	images in $\mathbf{Z} = \langle gc \rangle$
\widetilde{E}_1	B	$a^{-1}b, g^{-1}h, c^{-1}h$	$1, g^{-2} = (gc)^{-2}, gc$
\widetilde{E}_2	B	$a, e, i^{-1}j$	$1, 1, 1$
\widetilde{E}'_2	B	$cac^{-1}, cec^{-1}, c(i^{-1}j)c^{-1}$	$1, 1, 1$
\widetilde{E}_3	A	$c^{-2}, e^{-1}dl, e^{-1}f$	$1, 1, 1$
\widetilde{E}_4	A	$g^{-1}h^{-1}, i^{-2}, i^{-1}k^{-1}$	$1, 1, 1$
\widetilde{E}_5	A	$a^{-1}k^{-1}a^{-1}k, a^{-1}k^{-1}eg, c^{-1}i$	$1, gc, 1$
\widetilde{E}'_5	A	$c(a^{-1}k^{-1}a^{-1}k)c^{-1},$ $c(a^{-1}k^{-1}eg)c^{-1}, c(c^{-1}i)c^{-1}$	$1, gc, 1$

The results quoted above imply that the number of components in the lift of a boundary component \widetilde{E} is simply the order of the group $\mathbf{Z}/\langle A, [t'_5]^m \rangle$, where A is the set of images of the generators of $\pi_1 \widetilde{E}$. Those orders are, respectively, 1, m , m , m , m , 1, 1.

Let $H_0 = \langle\langle t_1, t_2, t'_2, t_3, t_4, t_5, (t'_5)^m \rangle\rangle_H$. Note that $(t'_5)^m$ is primitive in $H_0 \cap \pi_1 \widetilde{E}'_5$, since powers smaller than m of t'_5 are not in H_0 because they map to nontrivial elements of $H/H_0 = \mathbf{Z}_m$.

The only generator of $\pi_1 \widetilde{E}_1$ with nontrivial holonomy is $c^{-1}h$, mapping under the restriction $\psi : \pi_1 \widetilde{E}_1 \rightarrow \mathbf{Z}_m$ to the generator of \mathbf{Z}_m . If m is even, elements of $\ker \psi$ will all have trivial holonomy, meaning that \widetilde{E}_1 lifts to a collection of manifolds of type A; if m is odd, some elements of $\ker \psi$ will have nontrivial holonomy, so \widetilde{E}_1 will lift to a collection of manifolds of type B.

Putting everything together, we see that an m -fold cyclic cover of \widetilde{M}_{56} is a complement inside a simply-connected closed manifold N with $\chi(N) = 2m$ of

$2m + 3$ tori and $2m$ Klein bottles, if m is even;
 $2m + 2$ tori and $2m + 1$ Klein bottles, if m is odd.

At the end of Example 4.8 we saw that $H / \langle \langle t_1, t'_1, t_2, t'_2, t_3, t_4 \rangle \rangle_H = \mathbf{Z} \oplus \mathbf{Z}$, where $\pi_1 \widetilde{E}_5$ and $\pi_1 \widetilde{E}'_5$ map surjectively to $\mathbf{Z} \oplus \mathbf{Z}$ under the quotient map. This now allows for many choices of t_5 and t'_5 that make $H / \langle \langle t_1, t'_1, t_2, t'_2, t_3, t_4, t_5, t'_5 \rangle \rangle_H$ finite. With notation from Example 4.8 it is clear that

$$H / \langle \langle t_1, t'_1, t_2, t'_2, t_3, t_4, t_5, t'_5 \rangle \rangle_H \text{ is finite} \iff \begin{vmatrix} q+r & q'+r' \\ p+r & p'+r' \end{vmatrix} \neq 0.$$

Note that t_5 and t'_5 do not have to be primitive in $\pi_1 \widetilde{E}_5$ or $\pi_1 \widetilde{E}'_5$, so there is no condition on $\gcd(p, q, r)$ or $\gcd(p', q', r')$. It turns out t_5 and t'_5 are always primitive in $H_0 \cap \pi_1 \widetilde{E}_5$ and $H_0 \cap \pi_1 \widetilde{E}'_5$, where $H_0 = \langle \langle t_1, t'_1, t_2, t'_2, t_3, t_4, t_5, t'_5 \rangle \rangle_H$.

The number of boundary components of the cover of \widetilde{M}_{36} corresponding to H_0 is determined by considering the order of the groups

$$\mathbf{Z} \oplus \mathbf{Z} / \langle (q+r, p+r), (q'+r', p'+r'), A \rangle,$$

where A is the set of images of generators of boundary components of \widetilde{M}_{56} . Care must be taken to ascertain whether boundary components of type B lift to type B or A.

Among the many possibilities here, we just consider the case $q = p' = r = r' = 0$, with p and q' arbitrary. Under those conditions, one can see that \widetilde{M}_{36} has a finite cover with deck group $\mathbf{Z}_p \oplus \mathbf{Z}_{q'}$ that is the complement in a simply-connected manifold N with $\chi(N) = 2pq'$ of

$$\begin{aligned} &(1 + 2p) \gcd(2, q') + \gcd(2p, q') + 2 + 2p \text{ tori, if } q' \text{ is even;} \\ &(1 + 2p) \gcd(2, q') + \gcd(2p, q') + 2 \text{ tori and } 2p \text{ Klein bottles, if } q' \text{ is odd.} \end{aligned}$$

The table below lists other examples of complements in a simply-connected manifold N that we found using the same method. The last column indicates which translations different from the ones in Example 4.9 are set equal to 1 to cause $H / \langle \langle \text{translations} \rangle \rangle_H$ to be a finite group. The cover is then the one corresponding to the kernel of $H \rightarrow H / \langle \langle \text{translations} \rangle \rangle_H$. The Euler characteristic of N is always twice the cardinality of the group of deck transformations.

A cover of mfd.	with deck group	is a complement in a simply-connected closed manifold of:	Altered choice of translations
\widetilde{M}_{1091}	$\mathbf{Z}_m \oplus \mathbf{Z}_n$	$4 \gcd(m, n) + 5$ tori	t_2^m and t_3^n
\widetilde{M}_{71}	\mathbf{Z}_m	$3m + 3$ tori	t_3^m
\widetilde{M}_{112}	\mathbf{Z}_m	$3m + 3$ tori, m even $3m + 1$ tori, 2 K. bottles, m odd	t_3^m
\widetilde{M}_{92}	\mathbf{Z}_m	$4m + 2$ tori, m even $4m + 1$ tori, 1 K. bottle, m odd	t_5^m
\widetilde{M}_{92}	\mathbf{Z}_m	$3m + 3$ tori, m even $3m + 2$ tori, 1 K. bottle, m odd	$(t'_2)^m$
\widetilde{M}_{40}	\mathbf{Z}_m	$2m + 4$ tori, m K. bottles	$(t'_2)^m$ or $(t'_4)^m$

Example 4.11 (More examples of aspherical homology spheres.) Ratcliffe and Tschantz produced the first examples (infinitely many) of aspherical 4-manifolds that are homology spheres. The construction in [10] used \widetilde{M}_{1011} : by choosing fibers t_i of boundary components E_i as in Example 4.9 and filling them in with a disc, one gets a manifold N with $\pi_1 N = 1$, thus $H_1 N = 0$. By Proposition 4.1, $H_1 \widetilde{M}_{1011} = \mathbf{Z}^5$, generated by the fibers t_1, \dots, t_5 . If the fibers are altered, $\pi_1 N$ will likely fail to be trivial. However, one can retain $H_1 N = 0$ if t_i is replaced with $t_i + s_i$, where s_i is a carefully chosen element of $H_1 E_i = \mathbf{Z}^3$ that can have arbitrarily large length. When fibers are chosen to be sufficiently long, the manifold N supports a metric of nonpositive curvature and is therefore aspherical (N is a homology sphere because $H_1 N = 0$ and $\chi(N) = 2$). We note that this construction goes through for any of the orientable double covers from 4.4 whose boundary components are all 3-tori (i.e. which are complements of a collection of tori in S^4), namely \widetilde{M}_{71} , \widetilde{M}_{23} , \widetilde{M}_{1092} and \widetilde{M}_{1091} .

References

- [1] **B N Apanasov**, *Conformal geometry of discrete groups and manifolds*, de Gruyter Expositions in Mathematics 32, Walter de Gruyter & Co., Berlin (2000)
- [2] **M H Freedman, F Quinn**, *Topology of 4-manifolds*, Princeton Mathematical Series 39, Princeton University Press, Princeton, NJ (1990)
- [3] **R E Gompf, A I Stipsicz**, *4-manifolds and Kirby calculus*, Graduate Studies in Mathematics 20, American Mathematical Society, Providence, RI (1999)

- [4] **W Hantzsche, H Wendt**, *Dreidimensionale euklidische Raumformen*, Math. Ann. 110 (1935) 593–611
- [5] **D Ivanšić**, *Embeddability of noncompact hyperbolic manifolds as complements of codimension-1 and -2 submanifolds*, Topology Appl. 120 (2002) 211–236
- [6] **D Ivanšić**, *Hyperbolic structure on a complement of tori in the 4-sphere*, Adv. Geom. 4 (2004) 119–139
- [7] **R C Lyndon, P E Schupp**, *Combinatorial group theory*, Classics in Mathematics, Springer-Verlag, Berlin (2001)
- [8] **J G Ratcliffe**, *Foundations of hyperbolic manifolds*, Graduate Texts in Mathematics 149, Springer-Verlag, New York (1994)
- [9] **J G Ratcliffe, S T Tschantz**, *The volume spectrum of hyperbolic 4-manifolds*, Experiment. Math. 9 (2000) 101–125
- [10] **J G Ratcliffe, S T Tschantz**, *Some examples of aspherical 4-manifolds that are homology 4-spheres*, Topology 44 (2005) 341–350
- [11] **J A Wolf**, *Spaces of constant curvature*, Publish or Perish Inc., Houston, TX (1984)

DI: *Department of Mathematics and Statistics, Murray State University
Murray, KY 42071, USA*

JR and ST: *Department of Mathematics, Vanderbilt University
Nashville, TN 37240, USA*

Email: dubravko.ivansic@murraystate.edu, ratclifj@math.vanderbilt.edu,
tschantz@math.vanderbilt.edu

Received: 1 March 2005 Revised: 28 June 2005