

C^1 actions on the mapping class groups on the circle

KAMLESH PARWANI

Let S be a connected orientable surface with finitely many punctures, finitely many boundary components, and genus at least 6. Then any C^1 action of the mapping class group of S on the circle is trivial.

The techniques used in the proof of this result permit us to show that products of Kazhdan groups and certain lattices cannot have C^1 faithful actions on the circle. We also prove that for $n \geq 6$, any C^1 action of $\text{Aut}(F_n)$ or $\text{Out}(F_n)$ on the circle factors through an action of $\mathbb{Z}/2\mathbb{Z}$.

[37E10](#); [57M60](#)

1 Introduction

Theorem 1.1 *Let S be a connected orientable surface with finitely many punctures, finitely many boundary components, and genus at least 6. Then any C^1 action of the mapping class group of S on the circle is trivial.*

Thurston has established that there are faithful (effective) C^0 actions of the mapping class group of S on the circle when S is a surface of negative Euler characteristic with nonempty boundary (see Short and Wiest [15] or Ghys [10]). Dehornoy in [5], using different techniques, had previously shown that braid groups (mapping class groups of the punctured sphere) act faithfully on the circle. The fact that mapping class groups act on the circle had been observed earlier by Nielsen in his classical works. [Theorem 1.1](#) above asserts that nontrivial actions do not exist under the smoothness assumption when the genus is sufficiently large.

In fact, any C^0 action of a finitely generated group on the circle is (topologically) conjugate to a Lipschitz action (see Deroin, Kleptsyn and Navas [6]). So the known actions may be considered to be Lipschitz, and therefore, can be assumed to be differentiable almost everywhere. The result above shows that when the genus is at least 6, the actions of mapping class groups are not smoothable, that is, they are not conjugate to C^1 actions.

[Theorem 1.1](#) provides an infinite family of examples that answer the following question posed by John Franks.

Question 1.2 (Franks [8]) Does there exist a finitely generated group which acts faithfully on the circle via homeomorphisms but for which there is no faithful C^1 action?

Note that another example that answers this question has been constructed by Calegari [4]. [Theorem 1.1](#) also generalizes the following result of Farb and Franks [7].

Theorem 1.3 (Farb and Franks) *Let S be compact surface of genus at least 3 and at most one puncture. Then any C^2 action of the mapping class group of S on the circle must be trivial.*

The main results of this paper are the following two theorems.

Theorem 1.4 *Let H and G be two finitely generated groups such that $H_1(G, \mathbb{Z}) = H_1(H, \mathbb{Z}) = 0$. Then for any C^1 action of $H \times G$ on the circle, either $H \times 1$ acts trivially or $1 \times G$ acts trivially.*

Theorem 1.5 *Let H and G be two finitely generated groups such that $H^1(G_0, \mathbb{R}) = H^1(H_0, \mathbb{R}) = 0$ for all finite index normal subgroups H_0 and G_0 of H and G respectively. Then for any C^1 action of $H \times G$ on the circle, the induced action of either $H \times 1$ or $1 \times G$ factors through an action of a finite group.*

[Theorem 1.1](#) follows directly from [Theorem 1.4](#). [Theorem 1.5](#) has implications for C^1 actions of finite index subgroups of mapping class groups. Even if the mapping class groups don't have property (T) (see Taherkhani [16]), it is still conceivable (and conjectured) that all finite index subgroups have trivial first cohomology. If this is true, [Theorem 1.5](#) would imply that the following conjecture is true when the genus of surface is at least 4.

Conjecture 1.6 *Let G be a finite index subgroup of the mapping class group of S , where S is a connected orientable surface with finitely many punctures, finitely many boundary components, and genus at least 2. Then any C^1 action of G on the circle cannot be faithful.*

[Theorem 1.4](#) also implies that for $n \geq 6$, any C^1 action of $\text{Aut}(F_n)$ or $\text{Out}(F_n)$ on the circle factors through an action of $\mathbb{Z}/2\mathbb{Z}$. [Theorem 1.5](#) can be applied to show that products of Kazhdan groups and certain lattices cannot have C^1 faithful actions on the circle. The proofs of the main results are given in [Section 3](#) and the other corollaries are discussed in greater detail in [Section 4](#).

Conversations with Benson Farb and John Franks were valuable in the preparation of this article. The author would also like to thank the referee for several useful comments and suggestions.

2 Tools

We now present several results that will be used in the proof of [Theorem 1.1](#) and [Theorem 1.4](#).

2.1 Algebraic properties of $\text{Mod}(S)$

Throughout this paper, unless otherwise stated, by a surface we mean a connected orientable surface with finitely many punctures and boundary components. The mapping class group of S , denoted by $\text{Mod}(S)$, is the group of isotopy classes of orientation preserving diffeomorphisms of S ; the diffeomorphisms and the isotopies fix all the punctures and fix the boundary pointwise.

When S is a surface of genus at least 6, there exists a simple closed separating curve that splits S into two subsurfaces M_1 and M_2 such that the genus of each of these subsurfaces is at least 3. So $\text{Mod}(S)$ contains the subgroup $\text{Mod}(M_1) \times \text{Mod}(M_2)$. We will show that either $\text{Mod}(M_1)$ acts trivially or $\text{Mod}(M_2)$ acts trivially in order to establish [Theorem 1.1](#).

It is known that if M is a surface of genus 2, $H_1(\text{Mod}(M), \mathbb{Z}) = \mathbb{Z}_{10}$, and if M has genus greater than 2, then $H_1(\text{Mod}(M), \mathbb{Z}) = 0$. We will also use the fact that if the genus of M is at least 2, then $\text{Mod}(M)$ is generated by Dehn twists about finitely many nonseparating simple closed curves. The survey article by Korkmaz [\[12\]](#) is a good reference for these results.

2.2 Thurston's Stability Lemma

Theorem 2.1 (Thurston [\[17\]](#)) *Let G be a finitely generated group acting on \mathbb{R}^n with a global fixed point x . If the action is C^1 and $Dg(x)$ is the identity for all $g \in G$, then either there is a nontrivial homomorphism of G into \mathbb{R} or G acts trivially.*

We will need the following result which is a direct consequence of Thurston's Stability Lemma ([Theorem 2.1](#)).

Lemma 2.2 *Let G be a finitely generated group with an orientation preserving action on the circle. If G acts with a global fixed point and $H^1(G, \mathbb{R}) = 0$, then G acts trivially.*

Proof Let x be the global fixed point. It suffices to show that $g'(x) = 1$ for all $g \in G$. So consider the homomorphism $L: G \rightarrow \mathbb{R}$ defined by $L(g) = \log(g'(x))$. Since $H^1(G, \mathbb{R}) = 0$, this must be the trivial homomorphism, which implies that $g'(x) = 1$ for all $g \in G$. \square

2.3 Rotation numbers

The subject matter of this subsection is well known. The interested reader may refer to Ghys [10] or even Farb and Franks [7] for more details.

Definition 2.3 Let G be a finitely generated subgroup of the orientation preserving homeomorphisms of the circle and let μ be a G -invariant probability measure. The *mean rotation number homomorphism* is the map $\rho: G \rightarrow \mathbb{R}/\mathbb{Z}$ defined by

$$g \rightarrow \int_{S^1} (\tilde{g} - \text{Id}) d\tilde{\mu} \pmod{1},$$

where \tilde{g} and $\tilde{\mu}$ are lifts of g and μ to the real line and the integral is over a single fundamental domain.

The fact that this map is a homomorphism follows easily from the assumption that G preserves μ . Note that the mean rotation number of a homeomorphism is the same as the translation number of a circle homeomorphism, which was originally defined by Poincaré.

Proposition 2.4 *Let G be a finitely generated group with an orientation preserving action on the circle and let μ be a G -invariant probability measure. The element $g \in G$ acts with a fixed point if $\rho(g) = 0$.*

Corollary 2.5 *Let G be a finitely generated group with an orientation preserving action on the circle and let μ be a G -invariant probability measure. If $H^1(G, \mathbb{Z}) = 0$, then G acts with a global fixed point.*

Proof Since G is perfect and the target of ρ , the rotation number homomorphism, is an abelian group, ρ must be trivial. The above proposition implies that every element acts with a fixed point. The support of μ is contained in the fixed point set of each element in G , and therefore, G must have a global fixed point. \square

Corollary 2.6 *Let G be a finitely generated group with an orientation preserving action on the circle and let μ be a G -invariant probability measure. If $H^1(G, \mathbb{R}) = 0$, then G has a periodic orbit. In particular, a finite index normal subgroup of G acts with a global fixed point.*

Proof The assumption $H^1(G, \mathbb{R}) = 0$ implies that the image of the mean rotation number homomorphism is a finite group, and therefore, the kernel K is a finite index normal subgroup. The proposition above implies that every element in K acts with a fixed point. Recall that the support of the invariant measure is contained in the fixed point sets of every element in K . So K , a finite index normal subgroup of G , acts with a global fixed point. \square

2.4 Hyperbolic fixed points

The proof of [Theorem 1.4](#) relies heavily on the following result, which guarantees the existence of an element with a finite fixed point set in the absence of an invariant probability measure.

Theorem 2.7 (Deroin, Kleptsyn and Navas [\[6\]](#)) *Let G be a countable group with an orientation preserving C^1 action on the circle. If there is no G -invariant probability measure for the action, then there exists an element $g \in G$ that only has hyperbolic fixed points. In particular, g has a nonempty finite set of fixed points.*

3 C^1 actions on the circle

Theorem 1.4 *Let H and G be two finitely generated groups such that $H_1(G, \mathbb{Z}) = H_1(H, \mathbb{Z}) = 0$. Then for any C^1 action of $H \times G$ on the circle, either $H \times 1$ acts trivially or $1 \times G$ acts trivially.*

Proof Suppose that there exists a C^1 action of $H \times G$ on the circle. Since H and G are both perfect, this action must be orientation preserving. We now consider the induced action of G on the circle. By the result of Deroin, Kleptsyn, and Navas, either there exists a probability measure μ that is G -invariant or there is an element $g \in G$ that has a finite number of fixed points. We treat these two cases separately.

Case 1 There exists a probability measure μ that is G -invariant.

Now consider the mean rotation number homomorphism. This has to be trivial, and so, every element acts with a fixed point. The support of the measure μ is contained in the intersection of the fixed point sets of all the elements. So G has a global fixed point. Now apply the corollary to Thurston's Stability Lemma ([Lemma 2.2](#)) to conclude that G acts trivially.

Case 2 There is an element g in G that has a finite number of fixed points.

In this case, there is a finite set—the set of hyperbolic fixed points—left invariant by the induced action of the group H . This implies that the action of the group H has an invariant measure, and now we may argue as above, with G replaced by H , to conclude that H acts trivially. \square

The proof of [Theorem 1.5](#) is almost identical to the one given above. The assumption of trivial cohomologies rather than trivial homologies guarantees the existence of periodic

points and not necessarily fixed points. These periodic points are fixed points for some finite index normal subgroup, which must act trivially for the same reasons presented above. The proof can be easily obtained by making minor adjustments to the arguments in the proof of [Theorem 1.4](#) above.

Theorem 1.5 *Let H and G be two finitely generated groups such that $H^1(G_0, \mathbb{R}) = H^1(H_0, \mathbb{R}) = 0$ for all finite index normal subgroups H_0 and G_0 of H and G respectively. Then for any C^1 action of $H \times G$ on the circle, the induced action of either $H \times 1$ or $1 \times G$ factors through an action of a finite group.*

Proof Suppose that there exists a C^1 action of $H \times G$ on the circle. Let H_0 and G_0 be the index 2 subgroups of H and G respectively that have induced orientation preserving actions. We now focus on the action of the group G_0 . By the result of Deroin, Kleptsyn, and Navas, either there exists a probability measure μ that is G_0 -invariant or there is an element $g \in G_0$ that has a finite number of fixed points. We treat these two cases separately.

Case 1 There exists a probability measure μ that is G_0 -invariant.

Apply [Corollary 2.6](#) to obtain a finite index normal subgroup G_1 that acts with a global fixed point. Now Thurston's Stability Lemma ([Lemma 2.2](#)) implies that G_1 acts trivially, and so, the induced action of G factors through an action of a finite group.

Case 2 There is an element g in G_0 that has a finite number of fixed points.

In this case, there is a finite set—the set of hyperbolic fixed points—left invariant by the induced action of the group H_0 . This implies that the action of the group H_0 has an invariant measure, and now we may argue as above, with G_0 replaced by H_0 , to conclude that the induced action of H factors through an action of a finite group. \square

We are now ready to prove [Theorem 1.1](#), which follows immediately from [Theorem 1.4](#) above.

Theorem 1.1 *Let S be a connected orientable surface with finitely many punctures, finitely many boundary components, and genus at least 6. Then any C^1 action of the mapping class group of S on the circle is trivial.*

Proof We may split the surface S up into two subsurfaces M_1 and M_2 as in [Section 2.1](#) to obtain the action of $\text{Mod}(M_1) \times \text{Mod}(M_2)$ on the circle. Note that $\text{Mod}(M_1)$ and $\text{Mod}(M_2)$ are both finitely generated and perfect (see Korkmaz [[12](#)]). [Theorem 1.4](#)

implies that either $\text{Mod}(M_1)$ acts trivially or $\text{Mod}(M_2)$ acts trivially. In both cases, a Dehn twist about a nonseparating simple closed curve acts trivially. Since all such Dehn twists are conjugate to each other (see Ivanov [11] for instance), all of them must act trivially. Now recall the $\text{Mod}(S)$ is generated by these Dehn twists. \square

Remark 3.1 Suppose that every finite index normal subgroup of $\text{Mod}(S)$ has trivial first cohomology when S is a surface of genus at least 2. Now consider a C^1 action of a finite index subgroup K of $\text{Mod}(S)$, where S has genus at least 4. As in the proof of the theorem above, we may split S into two subsurfaces M_1 and M_2 , each with genus at least 2. The action of K induces an action of $H \times G$, where H and G are finite index subgroups of $\text{Mod}(M_1)$ and $\text{Mod}(M_2)$ respectively. Under our assumptions, [Theorem 1.5](#) implies that the action of K cannot be faithful. This elaborates on the comments just before [Conjecture 1.6](#).

The following corollary is a direct consequence of [Theorem 1.4](#).

Corollary 3.2 For $1 \leq i \leq n$, let G_i be a finitely generated group with $H_1(G_i, \mathbb{Z}) = 0$. Then for any C^1 action of $G_1 \times G_2 \times \cdots \times G_n$ on the circle, there exists at most one G_i with a nontrivial induced action on the circle.

An analogous statement follows from [Theorem 1.5](#).

4 Other corollaries

We first apply [Theorem 1.4](#) to $\text{Aut}(F_n)$ and $\text{Out}(F_n)$ and then apply [Theorem 1.5](#) to show that products of Kazhdan groups cannot act smoothly on the circle.

4.1 $\text{Aut}(F_n)$ and $\text{Out}(F_n)$

We now discuss the actions of $\text{Aut}(F_n)$, the automorphism group of F_n , and $\text{Out}(F_n)$, the outer automorphism group of F_n , where F_n is the free group of rank n . The proof that the C^1 actions of $\text{Aut}(F_n)$ and $\text{Out}(F_n)$ on the circle are trivial is similar to the proof of [Theorem 1.1](#).

The lemma below follows directly from the proof of Lemma 4.1 in [7].

Lemma 4.1 For $n \geq 3$, $\text{Aut}(F_n)$ contains a finitely generated subgroup T of index 2 which has a set of generators $\{A_{ij}, B_{ij}\}$ with $i \neq j$, $1 \leq i \leq n$ and $1 \leq j \leq n$. The subgroup T has a following properties.

- $H_1(T, \mathbb{Z}) = 0$.
- A_{ij} is conjugate to A_{kl} and B_{ij} is conjugate to B_{kl} , for all positive integers i, j, k, l between 1 and n .

Theorem 4.2 For $n \geq 6$, any C^1 action of $\text{Aut}(F_n)$ or $\text{Out}(F_n)$ on the circle factors through an action of $\mathbb{Z}/2\mathbb{Z}$.

Proof Suppose $\text{Aut}(F_n)$, where $n \geq 6$, has a C^1 action on the circle. We may consider the induced action of $\text{Aut}(F_3) \times \text{Aut}(F_{n-3})$, which is a subgroup of $\text{Aut}(F_n)$, on the circle. Apply the lemma above to obtain an action of $H \times G$ on the circle, where H and G are the index 2 subgroups of $\text{Aut}(F_3)$ and $\text{Aut}(F_{n-3})$ respectively that play the role of the subgroup T in the lemma above.

Since H and G are finitely generated and perfect, [Theorem 1.4](#) implies that either H acts trivially or G acts trivially. In either case, some A_{ij} and some B_{kl} must act trivially. Now recall that $\text{Aut}(F_n)$ has an index two subgroup that is generated by elements conjugate to A_{ij} and B_{kl} . So an index two subgroup of $\text{Aut}(F_n)$ acts trivially.

Since there exists a natural homomorphism from $\text{Aut}(F_n)$ onto $\text{Out}(F_n)$, the same result holds for $\text{Out}(F_n)$ also. \square

Note that the result of Bridson and Vogtmann in [\[2\]](#) about C^0 actions implies this theorem. However, their proof strongly relies on the existence of finite order elements. We give our proof here because it is short, easy, and it has implications for torsion free finite index subgroups of $\text{Aut}(F_n)$ and $\text{Out}(F_n)$. It should also be noted that Farb and Franks prove the same result for C^2 actions in [\[7\]](#).

4.2 Kazhdan groups

It is well known that discrete Kazhdan groups are finitely generated and any finite index normal subgroup of a discrete Kazhdan group has trivial first cohomology. So [Theorem 1.5](#) implies that products of discrete Kazhdan cannot have faithful C^1 actions on the circle.

Corollary 4.3 Let H and G be two discrete Kazhdan groups. Then for any C^1 action of $H \times G$ on the circle, the induced action of either $H \times 1$ or $1 \times G$ factors through an action of a finite group.

This is related to a result of Navas about actions of groups with property (T) on the circle; in [13] he proves that for $\alpha > 1/2$, any $C^{1+\alpha}$ action of a discrete Kazhdan group on the circle factors through an action of a finite group.

Certain lattices like $SL(n, \mathbb{Z})$ for $n > 2$ are known to have Kazhdan's property (T). Also when $n > 5$, $SL(n, \mathbb{Z})$ contains two commuting sublattices isomorphic to $SL(3, \mathbb{Z})$. These facts yield the following corollary.

Corollary 4.4 *Any C^1 action of a finite index subgroup of $SL(n, \mathbb{Z})$, for $n > 5$, on the circle factors through an action of a finite group.*

This result is well known. It follows from the work of Morris (Witte) in [18] (also see Ghys [9] and Burger and Monod [3]). The corollary above holds for a large family of lattices that contain two commuting sublattices that satisfy the hypothesis of Theorem 1.5.

The interested reader should also see Bader, Furman and Shaker [1] and Navas [14] for results about actions of products of lattices and Kazhdan groups on the circle.

References

- [1] U Bader, A Furman, A Shaker, *Superrigidity, Weyl groups, and actions on the circle* [arXiv:math/0605276](#)
- [2] MR Bridson, K Vogtmann, *Homomorphisms from automorphism groups of free groups*, Bull. London Math. Soc. 35 (2003) 785–792 [MR2000025](#)
- [3] M Burger, N Monod, *Bounded cohomology of lattices in higher rank Lie groups*, J. Eur. Math. Soc. (JEMS) 1 (1999) 199–235 [MR1694584](#)
- [4] D Calegari, *Dynamical forcing of circular groups*, Trans. Amer. Math. Soc. 358 (2006) 3473–3491 [MR2218985](#)
- [5] P Dehornoy, *A fast method for comparing braids*, Adv. Math. 125 (1997) 200–235 [MR1434111](#)
- [6] B Deroin, V Kleptsyn, A Navas, *Sur la dynamique unidimensionnelle en régularité intermédiaire*, Acta Math. 199 (2007) 199–262 [MR2358052](#)
- [7] B Farb, J Franks, *Groups of homeomorphisms of one-manifolds I: actions of nonlinear groups* [arXiv:math/0107085](#)
- [8] J Franks, personal communication (2001)
- [9] É Ghys, *Actions de réseaux sur le cercle*, Invent. Math. 137 (1999) 199–231 [MR1703323](#)

- [10] **É Ghys**, *Groups acting on the circle*, Enseign. Math. (2) 47 (2001) 329–407 [MR1876932](#)
- [11] **N V Ivanov**, *Mapping class groups*, from: “Handbook of geometric topology”, North-Holland, Amsterdam (2002) 523–633 [MR1886678](#)
- [12] **M Korkmaz**, *Low-dimensional homology groups of mapping class groups: a survey*, Turkish J. Math. 26 (2002) 101–114 [MR1892804](#)
- [13] **A Navas**, *Actions de groupes de Kazhdan sur le cercle*, Ann. Sci. École Norm. Sup. (4) 35 (2002) 749–758 [MR1951442](#)
- [14] **A Navas**, *Quelques nouveaux phénomènes de rang 1 pour les groupes de difféomorphismes du cercle*, Comment. Math. Helv. 80 (2005) 355–375 [MR2142246](#)
- [15] **H Short, B Wiest**, *Orderings of mapping class groups after Thurston*, Enseign. Math. (2) 46 (2000) 279–312 [MR1805402](#)
- [16] **F Taherkhani**, *The Kazhdan property of the mapping class group of closed surfaces and the first cohomology group of its cofinite subgroups*, Experiment. Math. 9 (2000) 261–274 [MR1780211](#)
- [17] **W P Thurston**, *A generalization of the Reeb stability theorem*, Topology 13 (1974) 347–352 [MR0356087](#)
- [18] **D Witte**, *Arithmetic groups of higher \mathbf{Q} -rank cannot act on 1-manifolds*, Proc. Amer. Math. Soc. 122 (1994) 333–340 [MR1198459](#)

Department of Mathematics, Eastern Illinois University
600 Lincoln Avenue, Charleston, IL 61920, USA

forty2@math.northwestern.edu

Received: 22 February 2008 Revised: 19 March 2008